Person with disabilities experiencing problems accessing NCHS birth and death tables should contact **nchsed@cdc.gov**, or call 301-458-4688

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
United States												
All races, both sexes	226,661	692	289	136	133	200	273	372	501	756	1,626	2,883
Male	108,560	415	163	75	87	113	166	214	248	401	835	1,479
Female	118,101	277	126	61	46	87	107	158	253	355	791	1,404
White, both sexes	205,112	401	190	90	67	125	174	260	335	539	1,192	2,223
Male	97,448	236	107	52	45	66	105	145	164	284	626	1,145
Female	107,664	165	83	38	22	59	69	115	171	255	566	1,078
Black, both sexes	17,008	252	76	38	61	64	93	102	146	193	383	593
Male	8,615	154	46	20	40	39	57	62	71	102	184	292
Female	8,393	98	30	18	21	25	36	40	75	91	199	301
American Indian 1/, both sexes	1,076	14	8	3	3	3	3	3	7	11	27	29
Male	527	9	5	1	1	1	1	1	4	7	14	20
Female	549	5	3	2	2	2	2	2	3	4	13	9
Asian or Pacific Islander 2/, both sexes	3,465	25	15	5	2	8	3	7	13	13	24	38
Male	1,970	16	5	2	1	7	3	6	9	8	11	22
Female	1,495	9	10	3	1	1	-	1	4	5	13	16
Alabama												
All races, both sexes	4,111	4	2	1	3	3	5	8	9	17	31	67
Male	2,024	2	2	-	3	1	4	5	2	12	16	44
Female	2,087	2	-	1	-	2	1	3	7	5	15	23
White, both sexes	3,530	3	1	1	2	1	1	4	6	13	21	49
Male	1,704	1	1	-	2	-	1	2	2	9	10	32
Female	1,826	2	-	1	-	1	-	2	4	4	11	17
Black, both sexes	572	1	1	-	1	2	4	3	3	4	10	18
Male	317	1	1	-	1	1	3	2	-	3	6	12
Female	255	-	-	-	-	1	1	1	3	1	4	6
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Alaska												
All races, both sexes	250	3	1	-	-	1	1	-	3	2	3	4
Male	118	1	-	-	-	-	-	-	2	-	-	3
Female	132	2	1	-	-	1	1	-	1	2	3	1
White, both sexes	171	-	-	-	-	-	1	-	1	2	3	2
Male	83	-	-	-	-	-	-	-	1	-	-	1
Female	88	-	-	-	-	-	1	-	-	2	3	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	65	3	1	-	-	1	-	-	2	-	-	2
Male	30	1	-	-	-	-	-	-	1	-	-	2
Female	35	2	1	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	4,733	24	10	4	6	4	4	5	14	23	34	55
Male	2,374	14	7	2	2	2	2	3	4	15	13	27
Female	2,359	10	3	2	4	2	2	2	10	8	21	28
White, both sexes	4,515	18	5	2	1	2	2	4	13	15	26	49
Male	2,257	9	3	2	1	1	1	3	4	9	11	23
Female	2,258	9	2	-	-	1	1	1	9	6	15	26
Black, both sexes	81	5	3	-	2	1	2	1	-	3	3	3
Male	42	4	2	-	-	-	1	-	-	2	1	1
Female	39	1	1	-	2	1	1	1	-	1	2	2
American Indian 1/, both sexes	111	1	2	2	3	1	-	-	1	5	4	3
Male	60	1	2	-	1	1	-	-	-	4	-	3
Female	51	-	-	2	2	-	-	-	1	1	4	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Arkansas												
All races, both sexes	2,755	4	4	1	2	5	6	3	11	11	24	31
Male	1,383	2	3	1	1	4	1	1	2	8	14	18
Female	1,372	2	1	-	1	1	5	2	9	3	10	13
White, both sexes	2,534	2	2	1	1	4	3	2	8	11	22	21
Male	1,270	-	1	1	1	3	1	-	-	8	13	13
Female	1,264	2	1	-	-	1	2	2	8	3	9	8
Black, both sexes	209	2	2	-	1	1	3	1	2	-	2	10
Male	108	2	2	-	-	1	-	1	1	-	1	5
Female	101	-	-	-	1	-	3	-	1	-	1	5
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	23,297	76	39	15	15	23	24	37	42	69	147	250
Male	11,043	42	20	6	14	16	14	24	21	38	76	138
Female	12,254	34	19	9	1	7	10	13	21	31	71	112
White, both sexes	20,257	50	32	9	7	14	16	21	27	52	111	201
Male	9,401	26	16	3	6	11	8	13	13	30	62	111
Female	10,856	24	16	6	1	3	8	8	14	22	49	90
Black, both sexes	1,336	15	2	4	7	4	8	11	11	14	25	35
Male	669	10	2	2	7	-	6	7	6	6	10	16
Female	667	5	-	2	-	4	2	4	5	8	15	19
American Indian 1/, both sexes	104	1	-	-	-	-	-	2	-	1	3	-
Male	55	1	-	-	-	-	-	1	-	1	2	-
Female	49	-	-	-	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1,600	10	5	2	1	5	-	3	4	2	8	14
Male	918	5	2	1	1	5	-	3	2	1	2	11
Female	682	5	3	1	-	-	-	-	2	1	6	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Colorado												
All races, both sexes	3,211	7	7	2	-	2	6	8	11	8	17	40
Male	1,642	4	4	1	-	2	4	5	6	2	8	20
Female	1,569	3	3	1	-	-	2	3	5	6	9	20
White, both sexes	3,081	6	4	1	-	2	6	7	9	7	13	36
Male	1,579	3	2	1	-	2	4	4	5	2	6	17
Female	1,502	3	2	-	-	-	2	3	4	5	7	19
Black, both sexes	92	1	1	1	-	-	-	1	2	1	1	3
Male	45	1	1	-	-	-	-	1	1	-	-	2
Female	47	-	-	1	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	14	-	1	-	-	-	-	-	-	-	3	1
Male	7	-	1	-	-	-	-	-	-	-	2	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	24	-	1	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	1	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2,977	6	1	2	1	1	-	2	5	6	14	27
Male	1,313	2	1	1	1	-	-	-	3	3	10	12
Female	1,664	4	-	1	-	1	-	2	2	3	4	15
White, both sexes	2,841	2	1	2	-	-	-	2	3	4	9	22
Male	1,244	-	1	1	-	-	-	-	2	1	6	11
Female	1,597	2	-	1	-	-	-	2	1	3	3	11
Black, both sexes	127	4	-	-	1	1	-	-	1	2	5	5
Male	62	2	-	-	1	-	-	-	-	2	4	1
Female	65	2	-	-	-	1	-	-	1	-	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Delaware												
All races, both sexes	633	-	-	1	-	-	1	2	2	3	5	2
Male	305	-	-	1	-	-	1	-	2	2	3	2
Female	328	-	-	-	-	-	-	2	-	1	2	-
White, both sexes	557	-	-	1	-	-	-	1	1	1	2	1
Male	263	-	-	1	-	-	-	-	1	1	2	1
Female	294	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	71	-	-	-	-	-	1	1	1	2	3	1
Male	40	-	-	-	-	-	1	-	1	1	1	1
Female	31	-	-	-	-	-	-	1	-	1	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	265	3	-	-	1	-	-	1	-	4	4	9
Male	125	3	-	-	1	-	-	1	-	2	1	7
Female	140	-	-	-	-	-	-	-	-	2	3	2
White, both sexes	82	-	-	-	-	-	-	-	-	1	1	2
Male	41	-	-	-	-	-	-	-	-	-	-	2
Female	41	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	180	3	-	-	1	-	-	1	-	3	3	7
Male	82	3	-	-	1	-	-	1	-	2	1	5
Female	98	-	-	-	-	-	-	-	-	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Florida												
All races, both sexes	14,053	54	16	8	7	11	21	27	22	41	86	179
Male	6,873	31	9	5	4	6	13	16	10	21	43	83
Female	7,180	23	7	3	3	5	8	11	12	20	43	96
White, both sexes	13,086	32	7	6	4	10	12	19	14	31	68	137
Male	6,362	20	4	4	2	6	9	11	8	18	35	62
Female	6,724	12	3	2	2	4	3	8	6	13	33	75
Black, both sexes	896	21	9	2	3	1	9	7	8	9	17	40
Male	472	10	5	1	2	-	4	5	2	2	8	20
Female	424	11	4	1	1	1	5	2	6	7	9	20
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	55	1	-	-	-	-	-	1	-	1	1	2
Male	31	1	-	-	-	-	-	-	-	1	-	1
Female	24	-	-	-	-	-	-	1	-	-	1	1
Georgia												
All races, both sexes	6,400	32	11	6	8	8	12	18	25	28	75	101
Male	3,025	18	5	4	4	4	8	10	11	17	33	49
Female	3,375	14	6	2	4	4	4	8	14	11	42	52
White, both sexes	5,346	8	4	2	4	4	7	11	13	16	42	61
Male	2,507	5	-	2	1	3	5	6	7	11	22	29
Female	2,839	3	4	-	3	1	2	5	6	5	20	32
Black, both sexes	1,009	24	5	4	4	4	5	7	12	12	32	39
Male	495	13	3	2	3	1	3	4	4	6	10	20
Female	514	11	2	2	1	3	2	3	8	6	22	19
American Indian 1/, both sexes	7	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	38	-	1	-	-	-	-	-	-	-	1	1
Male	19	-	1	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Hawaii												
All races, both sexes	812	3	2	-	-	1	-	1	3	5	3	12
Male	449	-	1	-	-	-	-	1	3	4	1	6
Female	363	3	1	-	-	1	-	-	-	1	2	6
White, both sexes	221	1	1	-	-	-	-	-	-	1	1	5
Male	116	-	-	-	-	-	-	-	-	1	-	4
Female	105	1	1	-	-	-	-	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	588	2	1	-	-	1	-	1	3	4	2	7
Male	331	-	1	-	-	-	-	1	3	3	1	2
Female	257	2	-	-	-	1	-	-	-	1	1	5
Idaho												
All races, both sexes	1,064	1	1	-	-	2	2	2	-	3	3	17
Male	565	1	1	-	-	1	1	2	-	2	-	10
Female	499	-	-	-	-	1	1	-	-	1	3	7
White, both sexes	1,056	1	1	-	-	2	2	2	-	2	3	16
Male	561	1	1	-	-	1	1	2	-	2	-	9
Female	495	-	-	-	-	1	1	-	-	-	3	7
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Illinois												
All races, both sexes	9,537	34	9	9	5	10	13	11	39	39	68	116
Male	4,581	23	7	4	4	9	7	7	25	21	39	66
Female	4,956	11	2	5	1	1	6	4	14	18	29	50
White, both sexes	8,447	15	4	6	1	4	7	6	17	24	46	77
Male	4,011	10	3	4	1	3	4	4	10	13	28	45
Female	4,436	5	1	2	-	1	3	2	7	11	18	32
Black, both sexes	996	18	5	2	4	6	6	5	22	15	21	38
Male	517	12	4	-	3	6	3	3	15	8	11	21
Female	479	6	1	2	1	-	3	2	7	7	10	17
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	90	1	-	1	-	-	-	-	-	-	1	1
Male	52	1	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	1	-	-	-	-	-	-	1	1
Indiana												
All races, both sexes	5,640	25	6	1	4	5	7	5	11	25	36	88
Male	2,758	13	2	1	2	2	5	1	7	17	17	48
Female	2,882	12	4	-	2	3	2	4	4	8	19	40
White, both sexes	5,367	19	5	1	2	5	3	5	9	18	31	74
Male	2,612	12	1	1	2	2	3	1	6	12	15	38
Female	2,755	7	4	-	-	3	-	4	3	6	16	36
Black, both sexes	266	6	1	-	2	-	4	-	2	7	5	14
Male	143	1	1	-	-	-	2	-	1	5	2	10
Female	123	5	-	-	2	-	2	-	1	2	3	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Iowa												
All races, both sexes	3,016	5	5	1	1	2	3	2	8	9	18	24
Male	1,497	3	3	1	-	2	3	-	4	5	10	11
Female	1,519	2	2	-	1	-	-	2	4	4	8	13
White, both sexes	2,982	5	4	1	1	1	3	2	7	9	16	23
Male	1,477	3	3	1	-	1	3	-	3	5	9	10
Female	1,505	2	1	-	1	-	-	2	4	4	7	13
Black, both sexes	24	-	1	-	-	1	-	-	1	-	1	1
Male	17	-	-	-	-	1	-	-	1	-	1	1
Female	7	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2,618	7	4	3	-	-	1	8	8	9	16	32
Male	1,248	6	2	1	-	-	1	7	3	6	6	12
Female	1,370	1	2	2	-	-	-	1	5	3	10	20
White, both sexes	2,487	2	1	3	-	-	-	6	6	7	13	30
Male	1,183	1	-	1	-	-	-	5	1	4	5	12
Female	1,304	1	1	2	-	-	-	1	5	3	8	18
Black, both sexes	92	4	3	-	-	-	1	2	2	2	2	2
Male	46	4	2	-	-	-	1	2	2	2	-	-
Female	46	-	1	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	29	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	1	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Kentucky												
All races, both sexes	4,196	10	3	1	-	1	8	6	11	15	35	70
Male	2,019	5	3	1	-	1	5	3	9	13	19	35
Female	2,177	5	-	-	-	-	3	3	2	2	16	35
White, both sexes	3,982	8	3	1	-	-	8	5	11	13	30	63
Male	1,918	4	3	1	-	-	5	3	9	11	17	33
Female	2,064	4	-	-	-	-	3	2	2	2	13	30
Black, both sexes	205	2	-	-	-	1	-	1	-	2	5	7
Male	97	1	-	-	-	1	-	-	-	2	2	2
Female	108	1	-	-	-	-	-	1	-	-	3	5
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3,243	21	8	2	4	2	9	5	16	11	41	60
Male	1,618	10	7	1	4	-	5	3	5	7	18	29
Female	1,625	11	1	1	-	2	4	2	11	4	23	31
White, both sexes	2,553	4	3	1	3	1	4	3	8	7	22	42
Male	1,240	2	3	1	3	-	2	3	2	3	8	20
Female	1,313	2	-	-	-	1	2	-	6	4	14	22
Black, both sexes	679	17	5	1	1	1	5	2	8	4	18	18
Male	372	8	4	-	1	-	3	-	3	4	10	9
Female	307	9	1	1	-	1	2	2	5	-	8	9
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Maine												
All races, both sexes	1,345	2	1	-	1	-	1	1	2	6	11	16
Male	653	1	1	-	1	-	-	-	2	2	5	13
Female	692	1	-	-	-	-	1	1	-	4	6	3
White, both sexes	1,336	2	-	-	1	-	1	1	1	6	11	16
Male	648	1	-	-	1	-	-	-	1	2	5	13
Female	688	1	-	-	-	-	1	1	-	4	6	3
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3,743	10	3	2	3	1	5	6	7	12	42	56
Male	1,727	5	1	-	1	1	4	4	4	7	21	22
Female	2,016	5	2	2	2	-	1	2	3	5	21	34
White, both sexes	3,042	5	-	1	-	-	3	2	4	8	18	31
Male	1,390	1	-	-	-	-	2	1	3	4	5	12
Female	1,652	4	-	1	-	-	1	1	1	4	13	19
Black, both sexes	640	5	1	1	3	1	2	4	3	4	22	24
Male	309	4	-	-	1	1	2	3	1	3	15	10
Female	331	1	1	1	2	-	-	1	2	1	7	14
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	54	-	2	-	-	-	-	-	-	-	2	-
Male	26	-	1	-	-	-	-	-	-	-	1	-
Female	28	-	1	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Massachusetts												
All races, both sexes	5,605	8	8	3	1	1	3	3	7	11	25	41
Male	2,453	6	4	1	1	-	1	2	2	4	14	20
Female	3,152	2	4	2	-	1	2	1	5	7	11	21
White, both sexes	5,422	6	7	3	1	1	2	3	6	11	20	37
Male	2,361	4	3	1	1	-	1	2	1	4	13	18
Female	3,061	2	4	2	-	1	1	1	5	7	7	19
Black, both sexes	132	1	1	-	-	-	1	-	-	-	5	4
Male	58	1	1	-	-	-	-	-	-	-	1	2
Female	74	-	-	-	-	-	1	-	-	-	4	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	1	-	-	-	-	-	-	1	-	-	-
Male	32	1	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	7,645	19	7	5	5	6	9	13	15	26	58	84
Male	3,673	9	5	2	3	5	6	9	6	10	33	41
Female	3,972	10	2	3	2	1	3	4	9	16	25	43
White, both sexes	6,923	11	3	1	3	1	6	8	12	17	42	62
Male	3,286	5	3	-	2	1	4	4	6	6	26	28
Female	3,637	6	-	1	1	-	2	4	6	11	16	34
Black, both sexes	655	8	3	4	2	5	3	5	2	8	13	20
Male	354	4	2	2	1	4	2	5	-	3	6	11
Female	301	4	1	2	1	1	1	-	2	5	7	9
American Indian 1/, both sexes	45	-	1	-	-	-	-	-	-	1	2	2
Male	24	-	-	-	-	-	-	-	-	1	1	2
Female	21	-	1	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	1	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Minnesota												
All races, both sexes	3,076	6	3	1	-	1	1	3	3	6	15	26
Male	1,479	4	3	1	-	1	1	1	1	3	7	15
Female	1,597	2	-	-	-	-	-	2	2	3	8	11
White, both sexes	2,993	2	2	-	-	-	1	2	2	5	14	23
Male	1,430	1	2	-	-	-	1	1	-	3	6	14
Female	1,563	1	-	-	-	-	-	1	2	2	8	9
Black, both sexes	40	2	1	1	-	1	-	1	-	1	1	3
Male	23	2	1	1	-	1	-	-	-	-	1	1
Female	17	-	-	-	-	-	-	1	-	1	-	2
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	2	-	-	-	-	-	-	1	-	-	-
Male	16	1	-	-	-	-	-	-	1	-	-	-
Female	10	1	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2,583	19	3	2	3	5	2	13	9	11	29	41
Male	1,301	14	3	2	2	3	-	6	6	2	14	17
Female	1,282	5	-	-	1	2	2	7	3	9	15	24
White, both sexes	1,970	2	2	1	-	2	-	6	4	3	14	19
Male	976	2	2	1	-	-	-	2	4	-	6	7
Female	994	-	-	-	-	2	-	4	-	3	8	12
Black, both sexes	602	16	1	1	3	3	2	7	5	8	15	22
Male	319	11	1	1	2	3	-	4	2	2	8	10
Female	283	5	-	-	1	-	2	3	3	6	7	12
American Indian 1/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Missouri												
All races, both sexes	5,312	19	5	1	7	3	4	12	13	11	50	78
Male	2,520	11	1	-	5	1	4	7	5	7	26	43
Female	2,792	8	4	1	2	2	-	5	8	4	24	35
White, both sexes	4,934	8	4	1	3	2	3	10	8	6	36	65
Male	2,341	4	1	-	2	1	3	6	3	3	17	36
Female	2,593	4	3	1	1	1	-	4	5	3	19	29
Black, both sexes	351	11	1	-	3	1	1	2	5	5	12	13
Male	172	7	-	-	3	-	1	1	2	4	7	7
Female	179	4	1	-	-	1	-	1	3	1	5	6
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	1	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	1	-	-	-	-	-	-	-
Montana												
All races, both sexes	948	1	2	2	-	-	1	1	2	2	3	6
Male	440	1	1	1	-	-	1	-	2	2	1	4
Female	508	-	1	1	-	-	-	1	-	-	2	2
White, both sexes	904	-	2	2	-	-	1	-	1	1	3	6
Male	415	-	1	1	-	-	1	-	1	1	1	4
Female	489	-	1	1	-	-	-	-	-	-	2	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	41	1	-	-	-	-	-	1	1	1	-	-
Male	23	1	-	-	-	-	-	-	1	1	-	-
Female	18	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Nebraska												
All races, both sexes	1,540	2	-	1	-	5	1	2	5	6	9	12
Male	752	1	-	-	-	1	-	1	2	3	3	6
Female	788	1	-	1	-	4	1	1	3	3	6	6
White, both sexes	1,497	2	-	1	-	4	1	2	3	6	8	8
Male	730	1	-	-	-	1	-	1	-	3	3	4
Female	767	1	-	1	-	3	1	1	3	3	5	4
Black, both sexes	23	-	-	-	-	1	-	-	2	-	1	3
Male	11	-	-	-	-	-	-	-	2	-	-	1
Female	12	-	-	-	-	1	-	-	-	-	1	2
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,826	8	5	2	-	-	5	4	4	12	14	33
Male	897	6	3	1	-	-	4	3	1	3	10	15
Female	929	2	2	1	-	-	1	1	3	9	4	18
White, both sexes	1,699	5	5	2	-	-	5	2	4	7	10	28
Male	830	4	3	1	-	-	4	1	1	3	8	13
Female	869	1	2	1	-	-	1	1	3	4	2	15
Black, both sexes	79	3	-	-	-	-	-	2	-	5	2	3
Male	44	2	-	-	-	-	-	2	-	-	1	1
Female	35	1	-	-	-	-	-	-	-	5	1	2
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	40	-	-	-	-	-	-	-	-	-	2	2
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	21	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
New Hampshire												
All races, both sexes	1,012	1	1	-	-	-	1	1	1	3	2	6
Male	489	1	-	-	-	-	-	1	-	2	-	3
Female	523	-	1	-	-	-	1	-	1	1	2	3
White, both sexes	1,006	1	1	-	-	-	1	1	1	3	2	6
Male	485	1	-	-	-	-	-	1	-	2	-	3
Female	521	-	1	-	-	-	1	-	1	1	2	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	5,499	11	4	3	6	6	5	13	8	24	40	64
Male	2,507	6	3	2	5	2	4	7	6	8	23	31
Female	2,992	5	1	1	1	4	1	6	2	16	17	33
White, both sexes	4,910	4	2	1	3	2	4	10	7	16	28	42
Male	2,226	2	2	1	2	-	3	6	5	4	17	23
Female	2,684	2	-	-	1	2	1	4	2	12	11	19
Black, both sexes	514	7	2	2	3	3	1	3	1	7	11	22
Male	230	4	1	1	3	1	1	1	1	3	6	8
Female	284	3	1	1	-	2	-	2	-	4	5	14
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	70	-	-	-	-	1	-	-	-	1	1	-
Male	48	-	-	-	-	1	-	-	-	1	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
New Mexico												
All races, both sexes	1,508	4	2	2	2	1	1	3	4	4	12	27
Male	747	2	1	-	2	1	-	2	4	3	5	16
Female	761	2	1	2	-	-	1	1	-	1	7	11
White, both sexes	1,420	4	1	2	2	1	1	2	3	4	10	23
Male	704	2	1	-	2	1	-	1	3	3	4	13
Female	716	2	-	2	-	-	1	1	-	1	6	10
Black, both sexes	14	-	-	-	-	-	-	1	-	-	1	1
Male	9	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	72	-	1	-	-	-	-	-	1	-	1	3
Male	33	-	-	-	-	-	-	-	1	-	1	3
Female	39	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	13,299	37	19	7	7	16	17	22	24	37	103	150
Male	6,121	23	10	6	3	10	9	9	13	25	58	80
Female	7,178	14	9	1	4	6	8	13	11	12	45	70
White, both sexes	11,610	18	10	4	3	12	14	14	13	24	68	112
Male	5,301	11	7	3	1	6	7	7	7	19	43	61
Female	6,309	7	3	1	2	6	7	7	6	5	25	51
Black, both sexes	1,392	17	7	2	4	3	3	8	10	11	34	35
Male	632	11	3	2	2	3	2	2	6	5	14	17
Female	760	6	4	-	2	-	1	6	4	6	20	18
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	276	2	2	1	-	1	-	-	1	2	1	3
Male	177	1	-	1	-	1	-	-	-	1	1	2
Female	99	1	2	-	-	-	-	-	1	1	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
North Carolina												
All races, both sexes	7,329	22	9	5	3	15	12	9	22	28	62	129
Male	3,428	18	5	3	2	7	4	5	9	13	32	62
Female	3,901	4	4	2	1	8	8	4	13	15	30	67
White, both sexes	6,355	11	6	4	1	12	4	5	11	15	37	83
Male	2,921	10	2	2	1	5	2	3	6	5	19	40
Female	3,434	1	4	2	-	7	2	2	5	10	18	43
Black, both sexes	920	11	3	1	2	3	7	4	11	13	25	41
Male	478	8	3	1	1	2	2	2	3	8	13	18
Female	442	3	-	-	1	1	5	2	8	5	12	23
American Indian 1/, both sexes	42	-	-	-	-	-	1	-	-	-	-	5
Male	24	-	-	-	-	-	-	-	-	-	-	4
Female	18	-	-	-	-	-	1	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	523	1	1	-	-	-	1	-	-	-	2	3
Male	276	1	1	-	-	-	1	-	-	-	1	2
Female	247	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	506	1	1	-	-	-	1	-	-	-	2	2
Male	269	1	1	-	-	-	1	-	-	-	1	1
Female	237	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Ohio												
All races, both sexes	10,095	25	12	4	2	6	7	13	16	41	63	151
Male	4,871	19	5	3	2	2	4	9	9	25	33	74
Female	5,224	6	7	1	-	4	3	4	7	16	30	77
White, both sexes	9,275	14	9	2	1	4	6	7	10	28	52	120
Male	4,425	12	3	2	1	1	3	5	5	15	29	59
Female	4,850	2	6	-	-	3	3	2	5	13	23	61
Black, both sexes	796	11	2	2	1	2	1	6	6	13	11	30
Male	434	7	2	1	1	1	1	4	4	10	4	15
Female	362	4	-	1	-	1	-	2	2	3	7	15
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	17	-	1	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	1	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3,614	10	4	2	4	4	6	5	4	23	33	46
Male	1,762	6	3	1	2	2	3	1	3	8	13	26
Female	1,852	4	1	1	2	2	3	4	1	15	20	20
White, both sexes	3,313	6	4	2	1	2	3	5	2	21	27	37
Male	1,620	4	3	1	1	-	1	1	1	8	9	22
Female	1,693	2	1	1	-	2	2	4	1	13	18	15
Black, both sexes	130	2	-	-	3	2	1	-	2	-	-	5
Male	61	1	-	-	1	2	1	-	2	-	-	3
Female	69	1	-	-	2	-	-	-	-	-	-	2
American Indian 1/, both sexes	169	2	-	-	-	-	1	-	-	2	6	4
Male	80	1	-	-	-	-	-	-	-	-	4	1
Female	89	1	-	-	-	-	1	-	-	2	2	3
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Oregon												
All races, both sexes	2,915	7	3	-	-	1	5	1	8	4	11	38
Male	1,415	3	1	-	-	1	3	1	4	3	7	21
Female	1,500	4	2	-	-	-	2	-	4	1	4	17
White, both sexes	2,844	4	3	-	-	1	5	1	8	4	11	36
Male	1,378	-	1	-	-	1	3	1	4	3	7	20
Female	1,466	4	2	-	-	-	2	-	4	1	4	16
Black, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	27	1	-	-	-	-	-	-	-	-	-	-
Male	10	1	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	2	-	-	-	-	-	-	-	-	-	1
Male	13	2	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	11,248	21	6	6	8	7	10	16	16	30	60	125
Male	5,344	10	3	2	3	6	6	10	5	11	38	64
Female	5,904	11	3	4	5	1	4	6	11	19	22	61
White, both sexes	10,431	15	6	5	7	2	7	12	13	24	44	94
Male	4,961	5	3	2	3	2	4	7	5	10	29	48
Female	5,470	10	3	3	4	-	3	5	8	14	15	46
Black, both sexes	777	6	-	1	1	5	3	4	3	6	16	31
Male	361	5	-	-	-	4	2	3	-	1	9	16
Female	416	1	-	1	1	1	1	1	3	5	7	15
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Rhode Island												
All races, both sexes	961	2	1	1	-	-	-	3	-	3	6	6
Male	425	2	-	1	-	-	-	2	-	1	3	1
Female	536	-	1	-	-	-	-	1	-	2	3	5
White, both sexes	930	2	1	1	-	-	-	3	-	2	6	6
Male	408	2	-	1	-	-	-	2	-	1	3	1
Female	522	-	1	-	-	-	-	1	-	1	3	5
Black, both sexes	25	-	-	-	-	-	-	-	-	1	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3,443	10	6	2	2	1	6	9	9	16	23	47
Male	1,700	6	3	2	1	1	4	4	3	10	12	24
Female	1,743	4	3	-	1	-	2	5	6	6	11	23
White, both sexes	2,854	4	2	1	-	1	2	8	4	10	12	30
Male	1,378	3	2	1	-	1	1	3	1	6	7	15
Female	1,476	1	-	-	-	-	1	5	3	4	5	15
Black, both sexes	577	6	4	1	2	-	4	1	5	6	11	16
Male	315	3	1	1	1	-	3	1	2	4	5	9
Female	262	3	3	-	1	-	1	-	3	2	6	7
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
South Dakota												
All races, both sexes	671	2	-	1	-	-	1	-	3	1	4	4
Male	341	1	-	1	-	-	1	-	2	-	3	1
Female	330	1	-	-	-	-	-	-	1	1	1	3
White, both sexes	641	2	-	-	-	-	1	-	2	1	3	4
Male	322	1	-	-	-	-	1	-	2	-	2	1
Female	319	1	-	-	-	-	-	-	-	1	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	30	-	-	1	-	-	-	-	1	-	1	-
Male	19	-	-	1	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5,549	20	5	5	4	9	13	10	12	13	52	93
Male	2,653	11	2	4	4	5	7	7	8	8	29	42
Female	2,896	9	3	1	-	4	6	3	4	5	23	51
White, both sexes	5,057	16	2	3	2	5	8	7	8	10	42	72
Male	2,395	9	1	2	2	4	4	5	6	5	24	33
Female	2,662	7	1	1	-	1	4	2	2	5	18	39
Black, both sexes	484	4	3	2	2	4	5	3	4	3	10	21
Male	253	2	1	2	2	1	3	2	2	3	5	9
Female	231	2	2	-	-	3	2	1	2	-	5	12
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Texas												
All races, both sexes	13,948	49	26	14	10	15	19	29	31	48	114	199
Male	6,644	29	12	9	5	9	14	12	15	25	58	99
Female	7,304	20	14	5	5	6	5	17	16	23	56	100
White, both sexes	12,737	37	19	12	5	12	11	24	25	36	90	162
Male	6,053	25	11	8	2	7	7	10	12	18	45	82
Female	6,684	12	8	4	3	5	4	14	13	18	45	80
Black, both sexes	1,068	11	5	2	5	3	5	5	5	12	23	32
Male	523	3	1	1	3	2	4	2	2	7	13	15
Female	545	8	4	1	2	1	1	3	3	5	10	17
American Indian 1/, both sexes	26	-	-	-	-	-	1	-	-	-	1	3
Male	13	-	-	-	-	-	1	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	117	1	2	-	-	-	2	-	1	-	-	2
Male	55	1	-	-	-	-	2	-	1	-	-	1
Female	62	-	2	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	1,204	4	3	1	3	4	1	4	3	3	9	15
Male	587	3	1	-	2	-	1	2	1	1	1	5
Female	617	1	2	1	1	4	-	2	2	2	8	10
White, both sexes	1,177	4	3	-	3	4	1	4	3	2	9	15
Male	573	3	1	-	2	-	1	2	1	-	1	5
Female	604	1	2	-	1	4	-	2	2	2	8	10
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	1	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Vermont												
All races, both sexes	491	3	-	-	-	1	-	-	-	1	-	2
Male	221	2	-	-	-	-	-	-	-	-	-	-
Female	270	1	-	-	-	1	-	-	-	1	-	2
White, both sexes	491	3	-	-	-	1	-	-	-	1	-	2
Male	221	2	-	-	-	-	-	-	-	-	-	-
Female	270	1	-	-	-	1	-	-	-	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5,466	22	8	3	-	4	7	8	10	16	39	83
Male	2,612	13	7	2	-	3	7	7	7	6	22	43
Female	2,854	9	1	1	-	1	-	1	3	10	17	40
White, both sexes	4,717	14	6	-	-	2	3	6	6	8	26	65
Male	2,221	7	5	-	-	2	3	5	4	3	14	35
Female	2,496	7	1	-	-	-	-	1	2	5	12	30
Black, both sexes	686	6	2	3	-	2	4	1	4	8	12	17
Male	352	5	2	2	-	1	4	1	3	3	7	8
Female	334	1	-	1	-	1	-	-	1	5	5	9
American Indian 1/, both sexes	6	1	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	57	1	-	-	-	-	-	1	-	-	1	-
Male	35	1	-	-	-	-	-	1	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Washington												
All races, both sexes	4,456	8	6	3	3	2	2	8	7	18	27	46
Male	2,088	6	2	1	2	1	1	6	3	8	10	26
Female	2,368	2	4	2	1	1	1	2	4	10	17	20
White, both sexes	4,214	4	5	2	3	-	2	8	7	15	23	40
Male	1,972	3	2	1	2	-	1	6	3	7	9	21
Female	2,242	1	3	1	1	-	1	2	4	8	14	19
Black, both sexes	78	1	1	1	-	1	-	-	-	1	2	4
Male	41	1	-	-	-	1	-	-	-	1	-	3
Female	37	-	1	1	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	55	2	-	-	-	1	-	-	-	-	1	-
Male	18	1	-	-	-	-	-	-	-	-	-	-
Female	37	1	-	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	109	1	-	-	-	-	-	-	-	2	1	2
Male	57	1	-	-	-	-	-	-	-	-	1	2
Female	52	-	-	-	-	-	-	-	-	2	-	-
West Virginia												
All races, both sexes	2,230	3	4	-	-	3	2	4	5	4	22	28
Male	1,123	3	3	-	-	-	-	4	1	-	11	15
Female	1,107	-	1	-	-	3	2	-	4	4	11	13
White, both sexes	2,192	3	3	-	-	3	2	3	5	3	20	28
Male	1,103	3	3	-	-	-	-	3	1	-	11	15
Female	1,089	-	-	-	-	3	2	-	4	3	9	13
Black, both sexes	37	-	1	-	-	-	-	1	-	1	2	-
Male	20	-	-	-	-	-	-	1	-	-	-	-
Female	17	-	1	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Wisconsin												
All races, both sexes	4,208	16	1	-	2	2	2	2	8	7	25	31
Male	2,073	12	1	-	1	1	2	2	4	5	14	18
Female	2,135	4	-	-	1	1	-	-	4	2	11	13
White, both sexes	4,068	14	1	-	2	1	-	1	6	7	23	26
Male	2,004	10	1	-	1	-	-	1	3	5	12	16
Female	2,064	4	-	-	1	1	-	-	3	2	11	10
Black, both sexes	103	1	-	-	-	1	2	1	2	-	1	4
Male	53	1	-	-	-	1	2	1	1	-	1	1
Female	50	-	-	-	-	-	-	-	1	-	-	3
American Indian 1/, both sexes	24	1	-	-	-	-	-	-	-	-	1	1
Male	12	1	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	558	1	2	1	-	-	-	3	3	1	1	13
Male	278	-	1	-	-	-	-	1	1	1	1	10
Female	280	1	1	1	-	-	-	2	2	-	-	3
White, both sexes	549	1	2	1	-	-	-	3	3	1	1	12
Male	272	-	1	-	-	-	-	1	1	1	1	9
Female	277	1	1	1	-	-	-	2	2	-	-	3
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
United States												
All races, both sexes	4,869	7,689	11,563	16,942	24,656	35,643	43,740	38,762	24,186	8,899	1,844	7
Male	2,659	4,184	6,241	8,894	12,936	18,440	21,411	17,289	9,282	2,601	422	5
Female	2,210	3,505	5,322	8,048	11,720	17,203	22,329	21,473	14,904	6,298	1,422	2
White, both sexes	3,895	6,486	10,142	15,075	22,426	32,686	40,432	36,124	22,446	8,173	1,625	6
Male	2,117	3,546	5,397	7,840	11,678	16,844	19,704	16,053	8,575	2,356	359	4
Female	1,778	2,940	4,745	7,235	10,748	15,842	20,728	20,071	13,871	5,817	1,266	2
Black, both sexes	869	1,058	1,206	1,534	1,809	2,255	2,448	1,915	1,225	512	176	-
Male	484	561	720	861	1,015	1,214	1,220	844	439	146	44	-
Female	385	497	486	673	794	1,041	1,228	1,071	786	366	132	-
American Indian 1/, both sexes	52	63	91	121	126	166	146	113	57	21	9	-
Male	26	26	47	63	62	77	73	61	20	5	3	-
Female	26	37	44	58	64	89	73	52	37	16	6	-
Asian or Pacific Islander 2/, both sexes	53	82	124	212	295	536	714	610	458	193	34	1
Male	32	51	77	130	181	305	414	331	248	94	16	1
Female	21	31	47	82	114	231	300	279	210	99	18	-
Alabama												
All races, both sexes	105	185	244	375	541	676	758	605	332	116	24	-
Male	63	110	135	216	282	348	350	284	112	29	4	-
Female	42	75	109	159	259	328	408	321	220	87	20	-
White, both sexes	75	142	210	321	483	597	679	529	273	101	18	-
Male	47	85	110	180	244	298	314	251	88	23	4	-
Female	28	57	100	141	239	299	365	278	185	78	14	-
Black, both sexes	29	42	34	53	58	77	77	76	59	14	6	-
Male	15	25	25	36	38	49	36	33	24	6	-	-
Female	14	17	9	17	20	28	41	43	35	8	6	-
American Indian 1/, both sexes	-	1	-	1	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	1	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Alaska												
All races, both sexes	10	14	22	22	32	38	38	29	18	9	-	-
Male	6	7	15	11	12	21	16	13	6	5	-	-
Female	4	7	7	11	20	17	22	16	12	4	-	-
White, both sexes	5	10	15	12	25	32	26	21	10	6	-	-
Male	2	5	12	6	10	19	11	9	3	4	-	-
Female	3	5	3	6	15	13	15	12	7	2	-	-
Black, both sexes	2	1	-	-	1	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	3	5	8	6	5	11	4	8	3	-	-
Male	3	2	2	4	2	2	5	2	3	1	-	-
Female	-	1	3	4	4	3	6	2	5	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	-	1	1	3	-	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	1	1	-	1	1	2	-	-	-	-
Arizona												
All races, both sexes	111	164	246	416	574	828	899	771	393	120	28	-
Male	61	89	136	230	302	436	461	363	155	43	7	-
Female	50	75	110	186	272	392	438	408	238	77	21	-
White, both sexes	105	152	233	394	557	797	878	745	382	111	24	-
Male	58	84	130	218	294	418	444	349	151	39	5	-
Female	47	68	103	176	263	379	434	396	231	72	19	-
Black, both sexes	4	6	7	9	4	11	6	4	5	2	-	-
Male	2	3	5	5	3	6	6	1	-	-	-	-
Female	2	3	2	4	1	5	-	3	5	2	-	-
American Indian 1/, both sexes	2	6	6	11	10	16	13	15	4	2	4	-
Male	1	2	1	6	3	9	10	10	2	2	2	-
Female	1	4	5	5	7	7	3	5	2	-	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	3	4	2	7	2	5	-	-
Male	-	-	-	1	2	3	1	3	2	2	-	-
Female	-	-	-	1	1	1	1	4	-	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Arkansas												
All races, both sexes	82	107	183	222	302	399	515	428	282	108	25	-
Male	32	68	102	107	181	218	272	210	97	38	3	-
Female	50	39	81	115	121	181	243	218	185	70	22	-
White, both sexes	68	94	161	208	282	376	481	407	266	95	19	-
Male	26	62	90	99	169	203	255	198	91	34	2	-
Female	42	32	71	109	113	173	226	209	175	61	17	-
Black, both sexes	13	13	21	13	20	19	32	21	15	12	6	-
Male	6	6	12	8	12	13	16	12	6	3	1	-
Female	7	7	9	5	8	6	16	9	9	9	5	-
American Indian 1/, both sexes	1	-	-	1	-	4	2	-	-	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	1	-	-	1	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
California												
All races, both sexes	441	661	1,082	1,585	2,278	3,458	4,749	4,296	2,741	1,060	209	-
Male	240	367	609	823	1,161	1,744	2,252	1,949	1,090	339	60	-
Female	201	294	473	762	1,117	1,714	2,497	2,347	1,651	721	149	-
White, both sexes	358	551	913	1,368	1,985	3,026	4,182	3,807	2,423	917	187	-
Male	199	302	500	695	981	1,506	1,946	1,702	939	281	51	-
Female	159	249	413	673	1,004	1,520	2,236	2,105	1,484	636	136	-
Black, both sexes	63	77	108	121	151	171	195	163	96	46	9	-
Male	31	42	67	71	84	89	85	75	33	18	2	-
Female	32	35	41	50	67	82	110	88	63	28	7	-
American Indian 1/, both sexes	7	3	9	11	14	17	16	12	5	3	-	-
Male	2	3	7	5	12	7	8	4	2	-	-	-
Female	5	-	2	6	2	10	8	8	3	3	-	-
Asian or Pacific Islander 2/, both sexes	13	30	52	85	128	244	356	314	217	94	13	-
Male	8	20	35	52	84	142	213	168	116	40	7	-
Female	5	10	17	33	44	102	143	146	101	54	6	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Colorado												
All races, both sexes	80	117	153	237	364	512	631	537	331	117	24	-
Male	46	77	94	144	187	258	328	265	149	34	4	-
Female	34	40	59	93	177	254	303	272	182	83	20	-
White, both sexes	74	109	147	222	343	494	612	523	328	115	23	-
Male	43	73	90	137	175	252	319	258	148	34	4	-
Female	31	36	57	85	168	242	293	265	180	81	19	-
Black, both sexes	5	7	6	13	18	10	8	9	2	2	1	-
Male	2	4	4	6	10	4	4	4	1	-	-	-
Female	3	3	2	7	8	6	4	5	1	2	1	-
American Indian 1/, both sexes	1	1	-	1	-	3	-	3	-	-	-	-
Male	1	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	1	-	2	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	3	5	11	2	1	-	-	-
Male	-	-	-	1	2	1	5	2	-	-	-	-
Female	-	-	-	-	1	4	6	-	1	-	-	-
Connecticut												
All races, both sexes	50	73	118	153	219	408	596	630	452	185	27	1
Male	27	32	59	86	115	211	273	253	170	48	6	-
Female	23	41	59	67	104	197	323	377	282	137	21	1
White, both sexes	45	67	108	137	209	392	574	620	440	178	25	1
Male	25	28	53	77	109	201	260	250	166	47	6	-
Female	20	39	55	60	100	191	314	370	274	131	19	1
Black, both sexes	4	5	10	16	10	16	19	9	10	7	2	-
Male	1	3	6	9	6	10	12	2	2	1	-	-
Female	3	2	4	7	4	6	7	7	8	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	3	1	2	-	-	-
Male	1	1	-	-	-	-	1	1	2	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Delaware												
All races, both sexes	14	18	20	53	75	118	134	100	60	22	3	-
Male	8	7	15	28	38	64	59	43	24	8	-	-
Female	6	11	5	25	37	54	75	57	36	14	3	-
White, both sexes	12	15	15	41	65	105	124	91	58	21	3	-
Male	7	6	10	23	33	57	54	37	23	7	-	-
Female	5	9	5	18	32	48	70	54	35	14	3	-
Black, both sexes	2	3	5	11	9	12	9	8	2	1	-	-
Male	1	1	5	4	4	7	5	6	1	1	-	-
Female	1	2	-	7	5	5	4	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Dist. of Columbia												
All races, both sexes	6	13	12	19	18	39	53	40	29	11	3	-
Male	4	9	4	11	7	19	20	16	16	3	1	-
Female	2	4	8	8	11	20	33	24	13	8	2	-
White, both sexes	-	-	3	3	3	18	17	18	8	6	2	-
Male	-	-	2	3	3	11	6	8	5	1	-	-
Female	-	-	1	-	-	7	11	10	3	5	2	-
Black, both sexes	6	13	9	16	15	20	35	22	21	4	1	-
Male	4	9	2	8	4	7	13	8	11	2	1	-
Female	2	4	7	8	11	13	22	14	10	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Florida												
All races, both sexes	284	459	683	1,101	1,558	2,260	2,823	2,472	1,375	467	98	1
Male	155	269	378	594	816	1,202	1,397	1,106	549	145	20	1
Female	129	190	305	507	742	1,058	1,426	1,366	826	322	78	-
White, both sexes	242	395	607	1,008	1,457	2,133	2,694	2,373	1,304	440	92	1
Male	134	234	334	540	762	1,125	1,321	1,058	519	135	20	1
Female	108	161	273	468	695	1,008	1,373	1,315	785	305	72	-
Black, both sexes	40	62	74	88	94	113	114	91	62	26	6	-
Male	21	34	42	52	50	69	68	42	25	10	-	-
Female	19	28	32	36	44	44	46	49	37	16	6	-
American Indian 1/, both sexes	1	-	1	2	2	4	2	3	1	-	-	-
Male	-	-	1	1	1	2	1	2	-	-	-	-
Female	1	-	-	1	1	2	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	3	5	10	13	5	8	1	-	-
Male	-	1	1	1	3	6	7	4	5	-	-	-
Female	1	1	-	2	2	4	6	1	3	1	-	-
Georgia												
All races, both sexes	191	281	410	526	771	1,017	1,126	978	539	198	39	-
Male	91	148	225	285	408	530	524	414	176	55	6	-
Female	100	133	185	241	363	487	602	564	363	143	33	-
White, both sexes	133	208	343	450	659	883	973	868	471	159	27	-
Male	65	107	182	244	342	460	449	371	150	42	4	-
Female	68	101	161	206	317	423	524	497	321	117	23	-
Black, both sexes	56	73	65	73	107	131	140	104	63	37	12	-
Male	24	41	41	40	63	70	69	40	24	12	2	-
Female	32	32	24	33	44	61	71	64	39	25	10	-
American Indian 1/, both sexes	1	-	-	-	2	1	1	-	-	1	-	-
Male	1	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	3	3	2	12	6	5	1	-	-
Male	1	-	2	1	2	-	6	3	2	-	-	-
Female	-	-	-	2	1	2	6	3	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Hawaii												
All races, both sexes	18	24	27	46	64	108	155	148	125	56	11	-
Male	10	16	15	27	31	60	95	80	60	34	5	-
Female	8	8	12	19	33	48	60	68	65	22	6	-
White, both sexes	5	5	8	11	23	33	48	34	34	9	2	-
Male	3	2	7	6	12	18	26	18	16	3	-	-
Female	2	3	1	5	11	15	22	16	18	6	2	-
Black, both sexes	-	1	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	18	19	35	41	74	107	114	91	46	9	-
Male	7	14	8	21	19	41	69	62	44	30	5	-
Female	6	4	11	14	22	33	38	52	47	16	4	-
Idaho												
All races, both sexes	20	35	48	71	115	187	209	201	105	30	12	-
Male	11	18	29	44	58	105	117	103	46	12	4	-
Female	9	17	19	27	57	82	92	98	59	18	8	-
White, both sexes	20	34	47	71	115	186	209	199	104	30	12	-
Male	11	18	28	44	58	104	117	102	46	12	4	-
Female	9	16	19	27	57	82	92	97	58	18	8	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	1	-	1	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Illinois												
All races, both sexes	204	299	463	657	973	1,422	1,880	1,683	1,114	395	94	-
Male	106	150	259	342	517	740	934	765	421	119	16	-
Female	98	149	204	315	456	682	946	918	693	276	78	-
White, both sexes	143	231	394	574	846	1,269	1,703	1,572	1,050	375	83	-
Male	78	123	217	297	445	660	840	710	392	114	10	-
Female	65	108	177	277	401	609	863	862	658	261	73	-
Black, both sexes	57	67	68	77	119	136	157	93	54	18	8	-
Male	26	27	42	42	67	69	82	43	23	5	5	-
Female	31	40	26	35	52	67	75	50	31	13	3	-
American Indian 1/, both sexes	-	1	1	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	6	7	17	20	17	10	2	3	-
Male	2	-	-	3	5	11	12	11	6	-	1	-
Female	2	-	-	3	2	6	8	6	4	2	2	-
Indiana												
All races, both sexes	124	210	300	473	668	896	1,058	907	575	180	36	-
Male	74	117	160	256	358	461	508	423	223	55	8	-
Female	50	93	140	217	310	435	550	484	352	125	28	-
White, both sexes	109	196	285	453	639	857	1,009	878	562	175	32	-
Male	64	109	152	249	343	440	483	405	216	52	6	-
Female	45	87	133	204	296	417	526	473	346	123	26	-
Black, both sexes	14	13	15	19	28	38	49	28	12	5	4	-
Male	10	8	8	6	14	21	25	17	7	3	2	-
Female	4	5	7	13	14	17	24	11	5	2	2	-
American Indian 1/, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Iowa												
All races, both sexes	39	79	109	204	318	402	609	569	371	202	36	-
Male	18	43	60	111	180	217	345	262	141	73	5	-
Female	21	36	49	93	138	185	264	307	230	129	31	-
White, both sexes	37	79	108	201	311	399	603	567	367	202	36	-
Male	17	43	59	109	175	215	342	262	139	73	5	-
Female	20	36	49	92	136	184	261	305	228	129	31	-
Black, both sexes	1	-	1	2	4	2	5	2	2	-	-	-
Male	1	-	1	1	4	2	3	-	1	-	-	-
Female	-	-	-	1	-	-	2	2	1	-	-	-
American Indian 1/, both sexes	1	-	-	1	2	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	1	-	1	-	-	-
Kansas												
All races, both sexes	48	76	107	181	259	416	479	472	319	141	32	-
Male	28	41	43	101	138	207	241	217	141	41	6	-
Female	20	35	64	80	121	209	238	255	178	100	26	-
White, both sexes	44	71	102	169	244	400	454	460	310	135	30	-
Male	26	41	40	95	130	197	228	214	138	40	5	-
Female	18	30	62	74	114	203	226	246	172	95	25	-
Black, both sexes	1	4	3	6	8	13	18	7	6	6	2	-
Male	1	-	1	3	5	9	10	-	2	1	1	-
Female	-	4	2	3	3	4	8	7	4	5	1	-
American Indian 1/, both sexes	2	-	1	5	6	2	7	5	-	-	-	-
Male	1	-	1	2	3	-	3	3	-	-	-	-
Female	1	-	-	3	3	2	4	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	1	1	1	-	-	3	-	-	-
Male	-	-	1	1	-	1	-	-	1	-	-	-
Female	1	1	-	-	1	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Kentucky												
All races, both sexes	120	154	270	407	518	643	749	615	404	133	23	-
Male	56	90	141	228	277	325	366	247	152	36	7	-
Female	64	64	129	179	241	318	383	368	252	97	16	-
White, both sexes	108	143	256	386	495	614	716	598	379	122	23	-
Male	52	83	133	215	264	310	349	240	144	35	7	-
Female	56	60	123	171	231	304	367	358	235	87	16	-
Black, both sexes	12	11	12	20	23	27	31	17	23	11	-	-
Male	4	7	7	12	13	14	16	7	8	1	-	-
Female	8	4	5	8	10	13	15	10	15	10	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	1	1	-	1	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	1	-	-	-
Louisiana												
All races, both sexes	94	137	205	304	399	490	579	464	271	96	24	1
Male	59	74	107	158	228	253	299	205	107	33	5	1
Female	35	63	98	146	171	237	280	259	164	63	19	-
White, both sexes	58	102	148	238	316	392	494	394	221	76	15	1
Male	32	51	76	116	175	205	254	169	89	22	3	1
Female	26	51	72	122	141	187	240	225	132	54	12	-
Black, both sexes	36	35	55	66	82	96	83	67	50	20	9	-
Male	27	23	29	42	53	48	43	34	18	11	2	-
Female	9	12	26	24	29	48	40	33	32	9	7	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	2	1	2	-	-	-	-
Male	-	-	2	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Maine												
All races, both sexes	24	43	58	101	159	196	265	219	165	63	11	-
Male	15	24	34	47	84	108	133	106	60	15	2	-
Female	9	19	24	54	75	88	132	113	105	48	9	-
White, both sexes	24	43	57	100	159	193	264	219	164	63	11	-
Male	15	24	34	46	84	106	133	106	60	15	2	-
Female	9	19	23	54	75	87	131	113	104	48	9	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Maryland												
All races, both sexes	98	131	172	253	407	535	713	692	419	147	29	-
Male	55	72	89	123	202	284	330	287	177	33	5	-
Female	43	59	83	130	205	251	383	405	242	114	24	-
White, both sexes	51	103	128	190	325	453	609	590	360	137	24	-
Male	26	57	64	96	156	245	277	248	157	33	3	-
Female	25	46	64	94	169	208	332	342	203	104	21	-
Black, both sexes	47	28	44	56	73	74	90	96	49	10	3	-
Male	29	15	25	25	41	35	45	36	17	-	1	-
Female	18	13	19	31	32	39	45	60	32	10	2	-
American Indian 1/, both sexes	-	-	-	2	1	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	2	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	5	8	7	14	5	9	-	2	-
Male	-	-	-	2	5	3	8	2	3	-	1	-
Female	-	-	-	3	3	4	6	3	6	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Massachusetts												
All races, both sexes	68	150	232	307	520	854	1,085	1,109	773	325	71	-
Male	36	83	117	151	286	419	479	434	281	94	18	-
Female	32	67	115	156	234	435	606	675	492	231	53	-
White, both sexes	64	143	224	296	501	828	1,051	1,080	755	315	68	-
Male	33	79	113	144	272	403	465	423	273	91	17	-
Female	31	64	111	152	229	425	586	657	482	224	51	-
Black, both sexes	4	6	7	5	12	14	26	22	12	9	3	-
Male	3	3	4	2	7	9	11	7	3	3	1	-
Female	1	3	3	3	5	5	15	15	9	6	2	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	6	7	11	8	7	6	1	-	-
Male	-	-	-	5	7	6	3	4	5	-	-	-
Female	-	-	1	1	-	5	5	3	1	1	-	-
Michigan												
All races, both sexes	167	259	385	578	853	1,332	1,500	1,263	769	235	57	-
Male	96	151	205	292	426	677	742	572	302	62	19	-
Female	71	108	180	286	427	655	758	691	467	173	38	-
White, both sexes	126	211	330	531	783	1,221	1,393	1,159	728	226	49	-
Male	67	123	177	262	394	616	683	521	283	59	16	-
Female	59	88	153	269	389	605	710	638	445	167	33	-
Black, both sexes	35	46	51	36	62	98	102	98	38	9	7	-
Male	24	28	26	21	28	57	57	49	19	3	2	-
Female	11	18	25	15	34	41	45	49	19	6	5	-
American Indian 1/, both sexes	4	2	1	9	6	9	2	5	1	-	-	-
Male	3	-	-	7	4	3	1	2	-	-	-	-
Female	1	2	1	2	2	6	1	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	3	2	2	4	3	1	2	-	1	-
Male	2	-	2	2	-	1	1	-	-	-	1	-
Female	-	-	1	-	2	3	2	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Minnesota												
All races, both sexes	46	68	132	186	296	459	640	585	412	163	23	1
Male	24	43	69	99	166	251	311	254	178	42	4	1
Female	22	25	63	87	130	208	329	331	234	121	19	-
White, both sexes	42	65	124	178	288	448	627	581	405	161	23	-
Male	23	40	64	95	159	245	305	252	174	41	4	-
Female	19	25	60	83	129	203	322	329	231	120	19	-
Black, both sexes	3	2	3	5	2	4	6	2	2	-	-	-
Male	-	2	2	3	2	2	3	1	1	-	-	-
Female	3	-	1	2	-	2	3	1	1	-	-	-
American Indian 1/, both sexes	1	-	4	2	3	3	4	-	-	-	-	-
Male	1	-	2	-	2	2	3	-	-	-	-	-
Female	-	-	2	2	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	3	4	3	2	5	2	-	1
Male	-	1	1	1	3	2	-	1	3	1	-	1
Female	-	-	-	-	-	2	3	1	2	1	-	-
Mississippi												
All races, both sexes	74	121	161	216	308	431	429	377	206	101	21	1
Male	36	67	94	122	176	249	222	168	69	27	1	1
Female	38	54	67	94	132	182	207	209	137	74	20	-
White, both sexes	44	80	123	159	250	344	364	314	161	66	11	1
Male	18	41	68	92	135	201	193	135	49	18	1	1
Female	26	39	55	67	115	143	171	179	112	48	10	-
Black, both sexes	29	40	38	55	58	85	63	62	44	35	10	-
Male	17	25	26	29	41	47	29	32	20	9	-	-
Female	12	15	12	26	17	38	34	30	24	26	10	-
American Indian 1/, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	2	-	1	-	1	1	-	-	-
Male	1	1	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Missouri												
All races, both sexes	133	196	267	409	584	884	1,012	832	539	201	51	1
Male	75	102	131	201	319	446	505	362	210	46	13	-
Female	58	94	136	208	265	438	507	470	329	155	38	1
White, both sexes	118	181	241	374	545	836	950	789	513	192	48	1
Male	67	92	120	185	302	425	475	345	198	44	12	-
Female	51	89	121	189	243	411	475	444	315	148	36	1
Black, both sexes	13	14	23	33	38	40	59	42	23	9	3	-
Male	8	10	10	15	17	18	30	17	12	2	1	-
Female	5	4	13	18	21	22	29	25	11	7	2	-
American Indian 1/, both sexes	2	-	2	2	1	4	1	-	2	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	2	-	1	1	1	3	1	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	4	2	1	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	1	1	-	-	2	2	1	1	-	-	-
Montana												
All races, both sexes	18	24	51	70	117	152	186	163	105	37	5	-
Male	10	13	28	35	61	81	78	76	36	9	-	-
Female	8	11	23	35	56	71	108	87	69	28	5	-
White, both sexes	13	19	46	67	110	150	181	156	105	36	5	-
Male	7	11	24	33	58	80	76	71	36	9	-	-
Female	6	8	22	34	52	70	105	85	69	27	5	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	5	3	3	7	2	5	6	-	1	-	-
Male	3	2	3	2	3	1	2	4	-	-	-	-
Female	2	3	-	1	4	1	3	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Nebraska												
All races, both sexes	25	29	55	86	163	253	293	273	215	83	22	-
Male	12	17	30	43	90	143	151	132	85	25	7	-
Female	13	12	25	43	73	110	142	141	130	58	15	-
White, both sexes	22	29	52	81	157	246	288	271	212	83	21	-
Male	10	17	28	40	87	138	150	131	85	25	6	-
Female	12	12	24	41	70	108	138	140	127	58	15	-
Black, both sexes	2	-	-	2	3	2	2	1	3	-	1	-
Male	2	-	-	1	2	1	1	-	-	-	1	-
Female	-	-	-	1	1	1	1	1	3	-	-	-
American Indian 1/, both sexes	1	-	3	2	2	2	3	1	-	-	-	-
Male	-	-	2	1	1	1	-	1	-	-	-	-
Female	1	-	1	1	1	1	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	3	-	-	-	-	-	-
Male	-	-	-	1	-	3	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Nevada												
All races, both sexes	35	76	127	174	284	298	341	229	125	44	6	-
Male	19	50	58	87	144	156	172	101	48	14	2	-
Female	16	26	69	87	140	142	169	128	77	30	4	-
White, both sexes	30	65	117	166	265	277	324	220	119	42	6	-
Male	15	44	53	81	134	141	164	98	46	14	2	-
Female	15	21	64	85	131	136	160	122	73	28	4	-
Black, both sexes	5	8	8	4	13	9	10	4	2	1	-	-
Male	4	4	5	3	8	7	5	2	-	-	-	-
Female	1	4	3	1	5	2	5	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	2	1	2	-	1	2	-	-	-
Male	-	-	-	2	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	2	2	5	10	7	4	2	1	-	-
Male	-	2	-	1	2	7	3	1	1	-	-	-
Female	-	1	2	1	3	3	4	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
New Hampshire												
All races, both sexes	23	33	49	79	116	144	201	194	110	43	4	-
Male	13	18	24	35	56	85	110	87	42	11	1	-
Female	10	15	25	44	60	59	91	107	68	32	3	-
White, both sexes	23	33	49	79	115	142	201	193	109	42	4	-
Male	13	18	24	35	55	83	110	87	41	11	1	-
Female	10	15	25	44	60	59	91	106	68	31	3	-
Black, both sexes	-	-	-	-	-	2	-	-	1	1	-	-
Male	-	-	-	-	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
New Jersey												
All races, both sexes	104	150	205	304	549	874	1,117	1,079	657	235	41	-
Male	58	83	115	165	270	407	527	462	248	65	10	-
Female	46	67	90	139	279	467	590	617	409	170	31	-
White, both sexes	84	118	167	252	489	782	1,032	1,013	603	216	35	-
Male	45	70	92	135	238	360	487	435	231	59	9	-
Female	39	48	75	117	251	422	545	578	372	157	26	-
Black, both sexes	20	32	33	47	50	78	73	56	41	16	6	-
Male	13	13	19	25	24	38	34	22	8	3	1	-
Female	7	19	14	22	26	40	39	34	33	13	5	-
American Indian 1/, both sexes	-	-	2	-	1	2	-	-	-	-	-	-
Male	-	-	2	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	5	9	12	12	10	13	3	-	-
Male	-	-	2	5	8	8	6	5	9	3	-	-
Female	-	-	1	-	1	4	6	5	4	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
New Mexico												
All races, both sexes	29	44	91	98	159	249	333	210	161	52	20	-
Male	15	22	49	58	74	125	165	111	69	17	6	-
Female	14	22	42	40	85	124	168	99	92	35	14	-
White, both sexes	29	39	84	94	154	241	313	196	148	50	19	-
Male	15	19	46	55	72	123	154	103	64	17	6	-
Female	14	20	38	39	82	118	159	93	84	33	13	-
Black, both sexes	-	1	-	-	2	2	1	2	3	-	-	-
Male	-	1	-	-	1	2	1	1	2	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	4	7	4	3	6	18	11	10	2	1	-
Male	-	2	3	3	1	-	10	6	3	-	-	-
Female	-	2	4	1	2	6	8	5	7	2	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
New York												
All races, both sexes	272	358	644	862	1,217	1,969	2,525	2,527	1,660	696	130	-
Male	150	188	363	445	601	1,013	1,211	1,082	592	196	34	-
Female	122	170	281	417	616	956	1,314	1,445	1,068	500	96	-
White, both sexes	205	273	538	712	1,055	1,766	2,250	2,278	1,499	627	115	-
Male	114	148	293	363	520	897	1,082	975	528	179	30	-
Female	91	125	245	349	535	869	1,168	1,303	971	448	85	-
Black, both sexes	62	77	93	131	132	164	220	195	122	50	12	-
Male	34	34	58	68	63	91	96	74	36	8	3	-
Female	28	43	35	63	69	73	124	121	86	42	9	-
American Indian 1/, both sexes	-	1	1	2	4	8	1	3	1	-	-	-
Male	-	1	1	1	2	3	1	2	-	-	-	-
Female	-	-	-	1	2	5	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	5	7	12	17	26	31	54	51	38	19	3	-
Male	2	5	11	13	16	22	32	31	28	9	1	-
Female	3	2	1	4	10	9	22	20	10	10	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
North Carolina												
All races, both sexes	149	310	456	643	826	1,165	1,361	1,112	686	240	65	-
Male	89	163	247	336	409	578	648	475	243	69	11	-
Female	60	147	209	307	417	587	713	637	443	171	54	-
White, both sexes	105	247	370	560	729	1,049	1,229	1,001	614	208	54	-
Male	60	122	196	290	357	524	580	419	213	57	8	-
Female	45	125	174	270	372	525	649	582	401	151	46	-
Black, both sexes	40	61	80	76	93	111	125	102	70	31	10	-
Male	25	40	49	42	50	52	66	52	28	11	2	-
Female	15	21	31	34	43	59	59	50	42	20	8	-
American Indian 1/, both sexes	3	2	3	6	3	3	4	9	2	-	1	-
Male	3	1	2	4	1	1	1	4	2	-	1	-
Female	-	1	1	2	2	2	3	5	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	1	1	2	3	-	-	1	-	-
Male	1	-	-	-	1	1	1	-	-	1	-	-
Female	-	-	3	1	-	1	2	-	-	-	-	-
North Dakota												
All races, both sexes	4	10	17	24	50	88	90	106	88	32	6	-
Male	2	5	9	15	28	52	57	54	41	6	1	-
Female	2	5	8	9	22	36	33	52	47	26	5	-
White, both sexes	2	10	15	21	48	85	87	105	88	32	6	-
Male	1	5	8	15	27	51	55	54	41	6	1	-
Female	1	5	7	6	21	34	32	51	47	26	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	2	3	2	3	3	1	-	-	-	-
Male	1	-	1	-	1	1	2	-	-	-	-	-
Female	1	-	1	3	1	2	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Ohio												
All races, both sexes	235	375	539	790	1,150	1,704	1,891	1,661	998	347	65	-
Male	138	201	298	403	606	894	926	739	379	87	15	-
Female	97	174	241	387	544	810	965	922	619	260	50	-
White, both sexes	198	313	487	710	1,073	1,558	1,775	1,571	947	332	58	-
Male	115	170	260	354	561	813	865	694	360	86	12	-
Female	83	143	227	356	512	745	910	877	587	246	46	-
Black, both sexes	36	61	49	77	75	145	111	86	50	15	6	-
Male	22	31	37	46	45	81	57	43	18	1	3	-
Female	14	30	12	31	30	64	54	43	32	14	3	-
American Indian 1/, both sexes	-	-	2	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	2	-	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	3	1	-	4	3	1	-	1	-
Male	1	-	1	3	-	-	4	1	1	-	-	-
Female	-	1	-	-	1	-	-	2	-	-	1	-
Oklahoma												
All races, both sexes	99	169	230	357	460	618	611	513	291	102	23	-
Male	48	93	115	180	239	334	304	226	117	30	8	-
Female	51	76	115	177	221	284	307	287	174	72	15	-
White, both sexes	83	148	206	320	427	573	570	486	276	95	19	-
Male	40	82	102	164	225	310	286	212	112	28	8	-
Female	43	66	104	156	202	263	284	274	164	67	11	-
Black, both sexes	11	9	6	17	11	16	17	15	7	3	3	-
Male	5	5	2	7	6	7	8	7	3	1	-	-
Female	6	4	4	10	5	9	9	8	4	2	3	-
American Indian 1/, both sexes	5	12	18	20	22	29	23	12	8	4	1	-
Male	3	6	11	9	8	17	10	7	2	1	-	-
Female	2	6	7	11	14	12	13	5	6	3	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Oregon												
All races, both sexes	58	115	147	207	360	480	540	482	340	89	19	-
Male	36	62	69	103	180	236	278	231	139	33	4	-
Female	22	53	78	104	180	244	262	251	201	56	15	-
White, both sexes	56	110	139	203	353	465	531	477	333	86	18	-
Male	34	60	66	101	176	227	273	229	136	32	4	-
Female	22	50	73	102	177	238	258	248	197	54	14	-
Black, both sexes	1	3	4	2	3	3	5	2	1	1	-	-
Male	1	1	2	1	2	2	3	-	-	1	-	-
Female	-	2	2	1	1	1	2	2	1	-	-	-
American Indian 1/, both sexes	-	2	4	2	4	7	2	1	2	1	1	-
Male	-	1	1	1	2	2	1	-	1	-	-	-
Female	-	1	3	1	2	5	1	1	1	1	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	5	2	2	4	1	-	-
Male	1	-	-	-	-	5	1	2	2	-	-	-
Female	-	-	-	-	-	-	1	-	2	1	-	-
Pennsylvania												
All races, both sexes	187	316	488	667	1,093	1,765	2,331	2,201	1,335	456	104	-
Male	101	173	274	363	560	927	1,152	1,001	506	109	20	-
Female	86	143	214	304	533	838	1,179	1,200	829	347	84	-
White, both sexes	144	263	439	594	1,005	1,640	2,211	2,107	1,266	435	98	-
Male	75	148	248	322	514	866	1,094	967	486	104	19	-
Female	69	115	191	272	491	774	1,117	1,140	780	331	79	-
Black, both sexes	39	52	48	66	86	119	109	91	66	19	6	-
Male	23	25	26	36	44	58	51	34	18	5	1	-
Female	16	27	22	30	42	61	58	57	48	14	5	-
American Indian 1/, both sexes	2	-	1	1	-	1	1	-	-	1	-	-
Male	1	-	-	1	-	1	1	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	6	2	5	10	3	3	1	-	-
Male	2	-	-	4	2	2	6	-	2	-	-	-
Female	-	1	-	2	-	3	4	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Rhode Island												
All races, both sexes	15	25	42	61	92	140	203	193	114	40	14	-
Male	8	16	15	29	47	74	88	84	45	5	4	-
Female	7	9	27	32	45	66	115	109	69	35	10	-
White, both sexes	15	24	40	57	88	135	199	187	112	39	13	-
Male	8	15	14	26	44	71	87	82	43	4	4	-
Female	7	9	26	31	44	64	112	105	69	35	9	-
Black, both sexes	-	1	2	4	3	4	3	5	1	-	1	-
Male	-	1	1	3	2	2	-	2	1	-	-	-
Female	-	-	1	1	1	2	3	3	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
South Carolina												
All races, both sexes	96	159	235	327	439	552	631	494	259	94	26	-
Male	48	92	128	188	245	302	312	187	98	23	7	-
Female	48	67	107	139	194	250	319	307	161	71	19	-
White, both sexes	66	115	182	258	369	485	554	431	218	85	17	-
Male	33	62	94	145	202	262	273	158	84	21	4	-
Female	33	53	88	113	167	223	281	273	134	64	13	-
Black, both sexes	30	44	51	68	70	64	75	61	40	9	9	-
Male	15	30	32	42	43	40	38	27	13	2	3	-
Female	15	14	19	26	27	24	37	34	27	7	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	3	-	1	1	-	-	-
Male	-	-	2	1	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	3	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
South Dakota												
All races, both sexes	10	21	23	34	72	89	127	133	94	43	9	-
Male	4	6	10	17	49	49	77	69	37	12	2	-
Female	6	15	13	17	23	40	50	64	57	31	7	-
White, both sexes	8	16	20	33	66	84	125	132	92	43	9	-
Male	3	5	8	16	43	45	75	69	37	12	2	-
Female	5	11	12	17	23	39	50	63	55	31	7	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	5	3	1	6	5	2	1	2	-	-	-
Male	1	1	2	1	6	4	2	-	-	-	-	-
Female	1	4	1	-	-	1	-	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	150	244	353	508	647	875	994	855	489	165	33	-
Male	79	118	182	257	340	459	502	358	174	52	5	-
Female	71	126	171	251	307	416	492	497	315	113	28	-
White, both sexes	111	211	312	471	592	814	924	807	462	152	26	-
Male	56	106	154	235	310	423	464	338	161	49	4	-
Female	55	105	158	236	282	391	460	469	301	103	22	-
Black, both sexes	39	33	39	37	53	61	69	47	26	12	7	-
Male	23	12	27	22	29	36	37	20	12	2	1	-
Female	16	21	12	15	24	25	32	27	14	10	6	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	1	1	1	1	-	-
Male	-	-	1	-	1	-	1	-	1	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Texas												
All races, both sexes	321	530	793	1,175	1,625	2,204	2,659	2,158	1,339	501	89	-
Male	182	263	421	601	870	1,106	1,273	970	492	155	24	-
Female	139	267	372	574	755	1,098	1,386	1,188	847	346	65	-
White, both sexes	257	453	718	1,043	1,491	2,042	2,478	2,027	1,258	460	77	-
Male	149	233	381	527	795	1,024	1,185	911	462	140	19	-
Female	108	220	337	516	696	1,018	1,293	1,116	796	320	58	-
Black, both sexes	58	66	72	120	116	135	158	113	71	40	11	-
Male	31	26	37	71	67	70	77	49	23	14	5	-
Female	27	40	35	49	49	65	81	64	48	26	6	-
American Indian 1/, both sexes	2	3	-	5	1	2	6	1	1	-	-	-
Male	-	1	-	2	1	1	4	1	1	-	-	-
Female	2	2	-	3	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	8	3	7	17	25	17	17	9	1	1	-
Male	2	3	3	1	7	11	7	9	6	1	-	-
Female	2	5	-	6	10	14	10	8	3	-	1	-
Utah												
All races, both sexes	28	60	59	86	141	192	221	205	119	37	6	-
Male	20	20	27	43	84	101	108	104	52	10	1	-
Female	8	40	32	43	57	91	113	101	67	27	5	-
White, both sexes	27	59	56	85	141	186	213	203	118	37	4	-
Male	19	19	26	42	84	98	104	103	51	10	1	-
Female	8	40	30	43	57	88	109	100	67	27	3	-
Black, both sexes	-	-	1	-	-	2	2	-	-	-	1	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	1	-
American Indian 1/, both sexes	1	-	2	-	-	3	1	2	-	-	1	-
Male	1	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	2	1	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	1	5	-	1	-	-	-
Male	-	1	-	1	-	-	3	-	1	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Vermont												
All races, both sexes	5	13	34	25	50	80	117	78	49	27	6	-
Male	2	8	19	15	25	44	50	33	16	6	1	-
Female	3	5	15	10	25	36	67	45	33	21	5	-
White, both sexes	5	13	34	25	50	80	117	78	49	27	6	-
Male	2	8	19	15	25	44	50	33	16	6	1	-
Female	3	5	15	10	25	36	67	45	33	21	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	106	184	261	394	664	866	1,062	903	562	215	49	-
Male	58	96	148	201	340	437	544	404	200	56	11	-
Female	48	88	113	193	324	429	518	499	362	159	38	-
White, both sexes	76	152	219	322	575	753	933	823	498	190	40	-
Male	41	78	124	159	284	373	475	365	183	52	9	-
Female	35	74	95	163	291	380	458	458	315	138	31	-
Black, both sexes	30	28	40	69	81	101	118	71	59	22	8	-
Male	17	16	22	41	50	56	62	34	14	2	2	-
Female	13	12	18	28	31	45	56	37	45	20	6	-
American Indian 1/, both sexes	-	1	-	-	1	1	-	-	1	-	-	-
Male	-	1	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	2	3	7	11	11	9	4	3	1	-
Male	-	1	2	1	5	7	7	5	2	2	-	-
Female	-	2	-	2	2	4	4	4	2	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Washington												
All races, both sexes	115	157	232	346	453	687	860	792	481	171	31	1
Male	61	80	110	178	241	335	402	367	182	58	7	1
Female	54	77	122	168	212	352	458	425	299	113	24	-
White, both sexes	106	142	214	327	430	643	819	764	464	164	31	1
Male	57	76	104	167	230	316	375	353	176	55	7	1
Female	49	66	110	160	200	327	444	411	288	109	24	-
Black, both sexes	7	6	8	3	5	10	14	9	3	2	-	-
Male	4	2	4	2	4	3	10	4	1	1	-	-
Female	3	4	4	1	1	7	4	5	2	1	-	-
American Indian 1/, both sexes	1	7	5	7	8	10	5	5	3	-	-	-
Male	-	1	1	4	2	2	3	3	1	-	-	-
Female	1	6	4	3	6	8	2	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	5	9	10	24	22	14	11	5	-	-
Male	-	1	1	5	5	14	14	7	4	2	-	-
Female	1	1	4	4	5	10	8	7	7	3	-	-
West Virginia												
All races, both sexes	50	85	113	214	293	404	390	319	205	64	18	-
Male	28	52	59	106	163	220	200	148	81	25	4	-
Female	22	33	54	108	130	184	190	171	124	39	14	-
White, both sexes	49	85	111	209	289	402	385	314	204	59	15	-
Male	28	52	57	104	159	218	198	145	81	22	3	-
Female	21	33	54	105	130	184	187	169	123	37	12	-
Black, both sexes	1	-	2	5	4	2	5	5	-	5	3	-
Male	-	-	2	2	4	2	2	3	-	3	1	-
Female	1	-	-	3	-	-	3	2	-	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Wisconsin												
All races, both sexes	73	108	186	274	396	708	849	770	522	191	35	-
Male	41	61	105	138	221	387	427	349	219	57	8	-
Female	32	47	81	136	175	321	422	421	303	134	27	-
White, both sexes	61	100	173	258	382	683	837	758	514	186	35	-
Male	34	57	98	129	216	375	423	342	216	57	8	-
Female	27	43	75	129	166	308	414	416	298	129	27	-
Black, both sexes	11	7	10	12	13	18	7	5	7	1	-	-
Male	7	4	6	8	5	7	3	2	3	-	-	-
Female	4	3	4	4	8	11	4	3	4	1	-	-
American Indian 1/, both sexes	1	1	2	3	1	3	2	6	-	2	-	-
Male	-	-	1	1	-	3	-	4	-	-	-	-
Female	1	1	1	2	1	-	2	2	-	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	4	3	1	1	2	-	-
Male	-	-	-	-	-	2	1	1	-	-	-	-
Female	-	-	1	1	-	2	2	-	1	2	-	-
Wyoming												
All races, both sexes	11	20	24	35	67	79	123	90	63	14	7	-
Male	5	10	13	16	38	42	70	38	29	2	-	-
Female	6	10	11	19	29	37	53	52	34	12	7	-
White, both sexes	10	19	24	33	65	79	122	90	62	14	7	-
Male	5	9	13	14	36	42	70	38	29	2	-	-
Female	5	10	11	19	29	37	52	52	33	12	7	-
Black, both sexes	1	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	1	2	-	1	-	-	-	-	-
Male	-	1	-	1	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
United States												
All races, both sexes	173	11	11	6	-	4	3	1	4	-	3	9
Male	83	9	8	4	-	2	2	1	2	-	2	4
Female	90	2	3	2	-	2	1	-	2	-	1	5
White, both sexes	152	7	8	6	-	2	3	1	4	-	2	6
Male	70	5	6	4	-	2	2	1	2	-	2	3
Female	82	2	2	2	-	-	1	-	2	-	-	3
Black, both sexes	18	3	2	-	-	2	-	-	-	-	1	3
Male	11	3	2	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	2	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	1	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Arkansas												
All races, both sexes	4	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	1	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	1	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	13	1	1	1	-	-	-	-	1	-	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	6	-	1	1	-	-	-	-	1	-	-	-
White, both sexes	9	-	-	1	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	1	-	-	-	-	1	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Colorado												
All races, both sexes	6	-	1	-	-	-	-	-	-	-	-	1
Male	4	-	1	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	1	-	-	-	-	-	-	-	-	1
Male	4	-	1	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Delaware												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Florida												
All races, both sexes	13	1	1	-	-	-	1	-	-	-	-	-
Male	10	-	1	-	-	-	1	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	1	-	-	-	-	1	-	-	-	-	-
Male	8	-	-	-	-	-	1	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	4	-	2	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Illinois												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	3	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Iowa												
All races, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Kentucky												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	1	1	-	-	1	-	-	1	-	-	-
Male	3	1	1	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	3	1	1	-	-	-	-	-	1	-	-	-
Male	3	1	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Maine												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Massachusetts												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Minnesota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Missouri												
All races, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Nebraska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	-	-	2	-	-	-	-	-	-	-	1
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	2	-	-	1	-	-	-	-	-	-	-	1
White, both sexes	2	-	-	2	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	2	-	-	-	-	-	-	-	-	-	-
Male	3	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
New Mexico												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	14	-	-	-	-	-	1	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	1	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
North Carolina												
All races, both sexes	5	-	-	-	-	1	1	-	-	-	-	2
Male	3	-	-	-	-	1	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	4	-	-	-	-	1	1	-	-	-	-	1
Male	3	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Ohio												
All races, both sexes	10	1	1	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Oregon												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Texas												
All races, both sexes	11	-	-	1	-	1	-	-	-	-	-	1
Male	4	-	-	1	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	10	-	-	1	-	-	-	-	-	-	-	1
Male	4	-	-	1	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Vermont												
All races, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	7	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Washington												
All races, both sexes	5	-	-	-	-	-	-	1	-	-	-	1
Male	4	-	-	-	-	-	-	1	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	3	-	2	-	-	-	-	-	-	-	-	-
Male	2	-	2	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	2	-	-	-	-	-	-	-	-	-
Male	2	-	2	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
Wisconsin												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
United States												
All races, both sexes	8	9	4	5	5	15	12	17	27	14	5	-
Male	5	4	1	4	2	9	6	7	7	3	1	-
Female	3	5	3	1	3	6	6	10	20	11	4	-
White, both sexes	7	8	2	4	5	14	12	17	26	13	5	-
Male	4	3	-	3	2	9	6	7	6	2	1	-
Female	3	5	2	1	3	5	6	10	20	11	4	-
Black, both sexes	1	-	2	1	-	1	-	-	1	1	-	-
Male	1	-	1	1	-	-	-	-	1	1	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	1	1	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	2	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Arkansas												
All races, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	1	1	2	1	-	-	1	1	1	-	-
Male	-	1	1	2	1	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	1	1	-	1	1	-	-	1	1	1	-	-
Male	-	1	-	1	1	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Colorado												
All races, both sexes	1	1	-	-	1	-	-	1	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	1	1	-	-	1	-	-	1	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	3	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	2	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	3	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Florida												
All races, both sexes	1	1	-	1	-	1	1	2	2	-	1	-
Male	-	1	-	1	-	1	1	2	1	-	1	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	1	-	1	1	2	2	-	1	-
Male	-	-	-	1	-	1	1	2	1	-	1	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Illinois												
All races, both sexes	1	1	-	-	-	-	1	-	-	-	1	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	1	-	-	-	-	1	-	-	-	1	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Iowa												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Massachusetts												
All races, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	1	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Minnesota												
All races, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Missouri												
All races, both sexes	-	-	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Nebraska												
All races, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
New Mexico												
All races, both sexes	-	-	1	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	1	-
White, both sexes	-	-	1	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	1	-	-	-	3	2	3	2	1	-	-
Male	-	-	-	-	-	2	-	1	1	-	-	-
Female	-	1	-	-	-	1	2	2	1	1	-	-
White, both sexes	-	1	-	-	-	3	2	3	2	1	-	-
Male	-	-	-	-	-	2	-	1	1	-	-	-
Female	-	1	-	-	-	1	2	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
North Carolina												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Ohio												
All races, both sexes	-	-	-	-	-	2	1	-	4	1	-	-
Male	-	-	-	-	-	2	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	3	1	-	-
White, both sexes	-	-	-	-	-	2	1	-	4	1	-	-
Male	-	-	-	-	-	2	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Oregon												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	-	-	-	1	-	2	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	2	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Texas												
All races, both sexes	-	1	-	-	-	-	1	1	4	1	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	1	-	-	-	-	-	-	3	1	-	-
White, both sexes	-	1	-	-	-	-	1	1	4	1	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	1	-	-	-	-	-	-	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	1	1	-	2	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	2	1	-
White, both sexes	-	-	-	-	-	-	1	1	-	2	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Washington												
All races, both sexes	1	-	-	-	-	1	-	-	1	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	1	-	-	1	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
Wisconsin												
All races, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
United States												
All races, both sexes	15	-	1	1	-	1	1	-	1	-	-	-
Male	8	-	1	-	-	-	1	-	-	-	-	-
Female	7	-	-	1	-	1	-	-	1	-	-	-
White, both sexes	13	-	1	1	-	-	1	-	1	-	-	-
Male	7	-	1	-	-	-	1	-	-	-	-	-
Female	6	-	-	1	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Arkansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Colorado												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Florida												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Iowa												
All races, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Missouri												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Oregon												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
United States												
All races, both sexes	1	2	1	2	-	1	1	2	-	-	-	-
Male	1	2	-	2	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	2	-	-	-	-
White, both sexes	1	1	1	2	-	1	1	2	-	-	-	-
Male	1	1	-	2	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Arkansas												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Colorado												
All races, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Florida												
All races, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Oregon												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Washington												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
United States												
All races, both sexes	25	5	5	2	-	-	1	-	1	-	-	3
Male	15	5	3	1	-	-	-	-	1	-	-	-
Female	10	-	2	1	-	-	1	-	-	-	-	3
White, both sexes	20	4	3	2	-	-	1	-	1	-	-	1
Male	12	4	1	1	-	-	-	-	1	-	-	-
Female	8	-	2	1	-	-	1	-	-	-	-	1
Black, both sexes	5	1	2	-	-	-	-	-	-	-	-	2
Male	3	1	2	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Colorado												
All races, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Florida												
All races, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	-	2	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	1	-	-	-	-	-	-	1	-	-	-
Male	2	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	1	-	-	-
Male	2	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Minnesota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
New Mexico												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Ohio												
All races, both sexes	3	1	1	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Texas												
All races, both sexes	2	-	-	1	-	-	-	-	-	-	-	1
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	2	-	-	1	-	-	-	-	-	-	-	1
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Vermont												
All races, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
Wisconsin												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
United States												
All races, both sexes	-	1	1	1	1	3	-	-	1	-	-	-
Male	-	1	-	1	1	2	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	1	-	-	-
White, both sexes	-	1	1	1	1	3	-	-	1	-	-	-
Male	-	1	-	1	1	2	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Minnesota												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
New Mexico												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Ohio												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
United States												
All races, both sexes	133	6	5	3	-	3	1	1	2	-	3	6
Male	60	4	4	3	-	2	1	1	1	-	2	4
Female	73	2	1	-	-	1	-	-	1	-	1	2
White, both sexes	119	3	4	3	-	2	1	1	2	-	2	5
Male	51	1	4	3	-	2	1	1	1	-	2	3
Female	68	2	-	-	-	-	-	-	1	-	-	2
Black, both sexes	12	2	-	-	-	1	-	-	-	-	1	1
Male	8	2	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	1	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	4	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Arkansas												
All races, both sexes	3	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	1	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	1	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	9	1	1	-	-	-	-	-	1	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	1	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Colorado												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Delaware												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Florida												
All races, both sexes	10	1	-	-	-	-	1	-	-	-	-	-
Male	7	-	-	-	-	-	1	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	1	-	-	-	-	1	-	-	-	-	-
Male	7	-	-	-	-	-	1	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Illinois												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	3	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Iowa												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Kentucky												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Maine												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Massachusetts												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Minnesota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Missouri												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Nebraska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	2	-	-	-	-	-	-	-	-	-	-
Male	3	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
New Mexico												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
North Carolina												
All races, both sexes	3	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Ohio												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Oregon												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Texas												
All races, both sexes	9	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	7	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Washington												
All races, both sexes	4	-	-	-	-	-	-	1	-	-	-	1
Male	3	-	-	-	-	-	-	1	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	3	-	2	-	-	-	-	-	-	-	-	-
Male	2	-	2	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	2	-	-	-	-	-	-	-	-	-
Male	2	-	2	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Wisconsin												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
United States												
All races, both sexes	7	6	2	2	4	11	11	15	26	14	5	-
Male	4	1	1	1	1	6	6	7	7	3	1	-
Female	3	5	1	1	3	5	5	8	19	11	4	-
White, both sexes	6	6	-	1	4	10	11	15	25	13	5	-
Male	3	1	-	-	1	6	6	7	6	2	1	-
Female	3	5	-	1	3	4	5	8	19	11	4	-
Black, both sexes	1	-	2	1	-	1	-	-	1	1	-	-
Male	1	-	1	1	-	-	-	-	1	1	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Arkansas												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	-	1	1	-	-	-	1	1	1	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	1	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Colorado												
All races, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	3	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	2	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	3	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Florida												
All races, both sexes	1	-	-	-	-	1	1	2	2	-	1	-
Male	-	-	-	-	-	1	1	2	1	-	1	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	1	1	2	2	-	1	-
Male	-	-	-	-	-	1	1	2	1	-	1	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Illinois												
All races, both sexes	1	1	-	-	-	-	1	-	-	-	1	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	1	-	-	-	-	1	-	-	-	1	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Iowa												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Massachusetts												
All races, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Missouri												
All races, both sexes	-	-	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Nebraska												
All races, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	1	-	-	-	2	2	3	2	1	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	1	-	-	-	1	2	2	1	1	-	-
White, both sexes	-	1	-	-	-	2	2	3	2	1	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	1	-	-	-	1	2	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
North Carolina												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Ohio												
All races, both sexes	-	-	-	-	-	1	1	-	4	1	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	3	1	-	-
White, both sexes	-	-	-	-	-	1	1	-	4	1	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Texas												
All races, both sexes	-	1	-	-	-	-	1	1	4	1	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	1	-	-	-	-	-	-	3	1	-	-
White, both sexes	-	1	-	-	-	-	1	1	4	1	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	1	-	-	-	-	-	-	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	1	1	-	2	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	2	1	-
White, both sexes	-	-	-	-	-	-	1	1	-	2	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Wisconsin												
All races, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
United States												
All races, both sexes	226,488	681	278	130	133	196	270	371	497	756	1,623	2,874
Male	108,477	406	155	71	87	111	164	213	246	401	833	1,475
Female	118,011	275	123	59	46	85	106	158	251	355	790	1,399
White, both sexes	204,960	394	182	84	67	123	171	259	331	539	1,190	2,217
Male	97,378	231	101	48	45	64	103	144	162	284	624	1,142
Female	107,582	163	81	36	22	59	68	115	169	255	566	1,075
Black, both sexes	16,990	249	74	38	61	62	93	102	146	193	382	590
Male	8,604	151	44	20	40	39	57	62	71	102	184	291
Female	8,386	98	30	18	21	23	36	40	75	91	198	299
American Indian 1/, both sexes	1,076	14	8	3	3	3	3	3	7	11	27	29
Male	527	9	5	1	1	1	1	1	4	7	14	20
Female	549	5	3	2	2	2	2	2	3	4	13	9
Asian or Pacific Islander 2/, both sexes	3,462	24	14	5	2	8	3	7	13	13	24	38
Male	1,968	15	5	2	1	7	3	6	9	8	11	22
Female	1,494	9	9	3	1	1	-	1	4	5	13	16
Alabama												
All races, both sexes	4,106	4	1	1	3	3	5	8	9	17	31	67
Male	2,023	2	1	-	3	1	4	5	2	12	16	44
Female	2,083	2	-	1	-	2	1	3	7	5	15	23
White, both sexes	3,525	3	-	1	2	1	1	4	6	13	21	49
Male	1,703	1	-	-	2	-	1	2	2	9	10	32
Female	1,822	2	-	1	-	1	-	2	4	4	11	17
Black, both sexes	572	1	1	-	1	2	4	3	3	4	10	18
Male	317	1	1	-	1	1	3	2	-	3	6	12
Female	255	-	-	-	-	1	1	1	3	1	4	6
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Alaska												
All races, both sexes	250	3	1	-	-	1	1	-	3	2	3	4
Male	118	1	-	-	-	-	-	-	2	-	-	3
Female	132	2	1	-	-	1	1	-	1	2	3	1
White, both sexes	171	-	-	-	-	-	1	-	1	2	3	2
Male	83	-	-	-	-	-	-	-	1	-	-	1
Female	88	-	-	-	-	-	1	-	-	2	3	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	65	3	1	-	-	1	-	-	2	-	-	2
Male	30	1	-	-	-	-	-	-	1	-	-	2
Female	35	2	1	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	4,731	24	10	4	6	4	4	5	14	23	34	55
Male	2,372	14	7	2	2	2	2	3	4	15	13	27
Female	2,359	10	3	2	4	2	2	2	10	8	21	28
White, both sexes	4,513	18	5	2	1	2	2	4	13	15	26	49
Male	2,255	9	3	2	1	1	1	3	4	9	11	23
Female	2,258	9	2	-	-	1	1	1	9	6	15	26
Black, both sexes	81	5	3	-	2	1	2	1	-	3	3	3
Male	42	4	2	-	-	-	1	-	-	2	1	1
Female	39	1	1	-	2	1	1	1	-	1	2	2
American Indian 1/, both sexes	111	1	2	2	3	1	-	-	1	5	4	3
Male	60	1	2	-	1	1	-	-	-	4	-	3
Female	51	-	-	2	2	-	-	-	1	1	4	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Arkansas												
All races, both sexes	2,751	4	4	1	2	4	6	3	11	11	24	30
Male	1,381	2	3	1	1	3	1	1	2	8	14	17
Female	1,370	2	1	-	1	1	5	2	9	3	10	13
White, both sexes	2,530	2	2	1	1	3	3	2	8	11	22	20
Male	1,268	-	1	1	1	2	1	-	-	8	13	12
Female	1,262	2	1	-	-	1	2	2	8	3	9	8
Black, both sexes	209	2	2	-	1	1	3	1	2	-	2	10
Male	108	2	2	-	-	1	-	1	1	-	1	5
Female	101	-	-	-	1	-	3	-	1	-	1	5
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	23,284	75	38	14	15	23	24	37	41	69	147	250
Male	11,036	41	20	6	14	16	14	24	21	38	76	138
Female	12,248	34	18	8	1	7	10	13	20	31	71	112
White, both sexes	20,248	50	32	8	7	14	16	21	26	52	111	201
Male	9,397	26	16	3	6	11	8	13	13	30	62	111
Female	10,851	24	16	5	1	3	8	8	13	22	49	90
Black, both sexes	1,334	15	2	4	7	4	8	11	11	14	25	35
Male	667	10	2	2	7	-	6	7	6	6	10	16
Female	667	5	-	2	-	4	2	4	5	8	15	19
American Indian 1/, both sexes	104	1	-	-	-	-	-	2	-	1	3	-
Male	55	1	-	-	-	-	-	1	-	1	2	-
Female	49	-	-	-	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1,598	9	4	2	1	5	-	3	4	2	8	14
Male	917	4	2	1	1	5	-	3	2	1	2	11
Female	681	5	2	1	-	-	-	-	2	1	6	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Colorado												
All races, both sexes	3,205	7	6	2	-	2	6	8	11	8	17	39
Male	1,638	4	3	1	-	2	4	5	6	2	8	19
Female	1,567	3	3	1	-	-	2	3	5	6	9	20
White, both sexes	3,075	6	3	1	-	2	6	7	9	7	13	35
Male	1,575	3	1	1	-	2	4	4	5	2	6	16
Female	1,500	3	2	-	-	-	2	3	4	5	7	19
Black, both sexes	92	1	1	1	-	-	-	1	2	1	1	3
Male	45	1	1	-	-	-	-	1	1	-	-	2
Female	47	-	-	1	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	14	-	1	-	-	-	-	-	-	-	3	1
Male	7	-	1	-	-	-	-	-	-	-	2	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	24	-	1	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	1	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2,973	6	1	2	1	1	-	2	5	6	14	27
Male	1,312	2	1	1	1	-	-	-	3	3	10	12
Female	1,661	4	-	1	-	1	-	2	2	3	4	15
White, both sexes	2,837	2	1	2	-	-	-	2	3	4	9	22
Male	1,243	-	1	1	-	-	-	-	2	1	6	11
Female	1,594	2	-	1	-	-	-	2	1	3	3	11
Black, both sexes	127	4	-	-	1	1	-	-	1	2	5	5
Male	62	2	-	-	1	-	-	-	-	2	4	1
Female	65	2	-	-	-	1	-	-	1	-	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Delaware												
All races, both sexes	630	-	-	1	-	-	1	2	2	3	5	2
Male	305	-	-	1	-	-	1	-	2	2	3	2
Female	325	-	-	-	-	-	-	2	-	1	2	-
White, both sexes	554	-	-	1	-	-	-	1	1	1	2	1
Male	263	-	-	1	-	-	-	-	1	1	2	1
Female	291	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	71	-	-	-	-	-	1	1	1	2	3	1
Male	40	-	-	-	-	-	1	-	1	1	1	1
Female	31	-	-	-	-	-	-	1	-	1	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	264	3	-	-	1	-	-	1	-	4	4	9
Male	124	3	-	-	1	-	-	1	-	2	1	7
Female	140	-	-	-	-	-	-	-	-	2	3	2
White, both sexes	82	-	-	-	-	-	-	-	-	1	1	2
Male	41	-	-	-	-	-	-	-	-	-	-	2
Female	41	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	179	3	-	-	1	-	-	1	-	3	3	7
Male	81	3	-	-	1	-	-	1	-	2	1	5
Female	98	-	-	-	-	-	-	-	-	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Florida												
All races, both sexes	14,040	53	15	8	7	11	20	27	22	41	86	179
Male	6,863	31	8	5	4	6	12	16	10	21	43	83
Female	7,177	22	7	3	3	5	8	11	12	20	43	96
White, both sexes	13,075	31	7	6	4	10	11	19	14	31	68	137
Male	6,354	20	4	4	2	6	8	11	8	18	35	62
Female	6,721	11	3	2	2	4	3	8	6	13	33	75
Black, both sexes	895	21	8	2	3	1	9	7	8	9	17	40
Male	471	10	4	1	2	-	4	5	2	2	8	20
Female	424	11	4	1	1	1	5	2	6	7	9	20
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	54	1	-	-	-	-	-	1	-	1	1	2
Male	30	1	-	-	-	-	-	-	-	1	-	1
Female	24	-	-	-	-	-	-	1	-	-	1	1
Georgia												
All races, both sexes	6,396	32	9	6	8	8	12	18	25	28	75	101
Male	3,023	18	4	4	4	4	8	10	11	17	33	49
Female	3,373	14	5	2	4	4	4	8	14	11	42	52
White, both sexes	5,343	8	3	2	4	4	7	11	13	16	42	61
Male	2,506	5	-	2	1	3	5	6	7	11	22	29
Female	2,837	3	3	-	3	1	2	5	6	5	20	32
Black, both sexes	1,008	24	4	4	4	4	5	7	12	12	32	39
Male	494	13	2	2	3	1	3	4	4	6	10	20
Female	514	11	2	2	1	3	2	3	8	6	22	19
American Indian 1/, both sexes	7	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	38	-	1	-	-	-	-	-	-	-	1	1
Male	19	-	1	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Hawaii												
All races, both sexes	812	3	2	-	-	1	-	1	3	5	3	12
Male	449	-	1	-	-	-	-	1	3	4	1	6
Female	363	3	1	-	-	1	-	-	-	1	2	6
White, both sexes	221	1	1	-	-	-	-	-	-	1	1	5
Male	116	-	-	-	-	-	-	-	-	1	-	4
Female	105	1	1	-	-	-	-	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	588	2	1	-	-	1	-	1	3	4	2	7
Male	331	-	1	-	-	-	-	1	3	3	1	2
Female	257	2	-	-	-	1	-	-	-	1	1	5
Idaho												
All races, both sexes	1,063	1	1	-	-	2	2	2	-	3	3	16
Male	564	1	1	-	-	1	1	2	-	2	-	9
Female	499	-	-	-	-	1	1	-	-	1	3	7
White, both sexes	1,055	1	1	-	-	2	2	2	-	2	3	15
Male	560	1	1	-	-	1	1	2	-	2	-	8
Female	495	-	-	-	-	1	1	-	-	-	3	7
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Illinois												
All races, both sexes	9,533	34	9	9	5	10	13	11	39	39	68	116
Male	4,579	23	7	4	4	9	7	7	25	21	39	66
Female	4,954	11	2	5	1	1	6	4	14	18	29	50
White, both sexes	8,443	15	4	6	1	4	7	6	17	24	46	77
Male	4,009	10	3	4	1	3	4	4	10	13	28	45
Female	4,434	5	1	2	-	1	3	2	7	11	18	32
Black, both sexes	996	18	5	2	4	6	6	5	22	15	21	38
Male	517	12	4	-	3	6	3	3	15	8	11	21
Female	479	6	1	2	1	-	3	2	7	7	10	17
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	90	1	-	1	-	-	-	-	-	-	1	1
Male	52	1	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	1	-	-	-	-	-	-	1	1
Indiana												
All races, both sexes	5,637	25	6	-	4	5	7	5	11	25	36	88
Male	2,756	13	2	-	2	2	5	1	7	17	17	48
Female	2,881	12	4	-	2	3	2	4	4	8	19	40
White, both sexes	5,364	19	5	-	2	5	3	5	9	18	31	74
Male	2,610	12	1	-	2	2	3	1	6	12	15	38
Female	2,754	7	4	-	-	3	-	4	3	6	16	36
Black, both sexes	266	6	1	-	2	-	4	-	2	7	5	14
Male	143	1	1	-	-	-	2	-	1	5	2	10
Female	123	5	-	-	2	-	2	-	1	2	3	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Iowa												
All races, both sexes	3,014	5	4	1	1	2	3	2	8	9	18	24
Male	1,496	3	2	1	-	2	3	-	4	5	10	11
Female	1,518	2	2	-	1	-	-	2	4	4	8	13
White, both sexes	2,980	5	3	1	1	1	3	2	7	9	16	23
Male	1,476	3	2	1	-	1	3	-	3	5	9	10
Female	1,504	2	1	-	1	-	-	2	4	4	7	13
Black, both sexes	24	-	1	-	-	1	-	-	1	-	1	1
Male	17	-	-	-	-	1	-	-	1	-	1	1
Female	7	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2,617	7	4	3	-	-	1	8	8	9	16	32
Male	1,248	6	2	1	-	-	1	7	3	6	6	12
Female	1,369	1	2	2	-	-	-	1	5	3	10	20
White, both sexes	2,486	2	1	3	-	-	-	6	6	7	13	30
Male	1,183	1	-	1	-	-	-	5	1	4	5	12
Female	1,303	1	1	2	-	-	-	1	5	3	8	18
Black, both sexes	92	4	3	-	-	-	1	2	2	2	2	2
Male	46	4	2	-	-	-	1	2	2	2	-	-
Female	46	-	1	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	29	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	1	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Kentucky												
All races, both sexes	4,195	10	3	1	-	1	8	6	11	15	35	70
Male	2,019	5	3	1	-	1	5	3	9	13	19	35
Female	2,176	5	-	-	-	-	3	3	2	2	16	35
White, both sexes	3,981	8	3	1	-	-	8	5	11	13	30	63
Male	1,918	4	3	1	-	-	5	3	9	11	17	33
Female	2,063	4	-	-	-	-	3	2	2	2	13	30
Black, both sexes	205	2	-	-	-	1	-	1	-	2	5	7
Male	97	1	-	-	-	1	-	-	-	2	2	2
Female	108	1	-	-	-	-	-	1	-	-	3	5
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3,239	20	7	2	4	1	9	5	15	11	41	60
Male	1,615	9	6	1	4	-	5	3	4	7	18	29
Female	1,624	11	1	1	-	1	4	2	11	4	23	31
White, both sexes	2,550	3	2	1	3	1	4	3	7	7	22	42
Male	1,237	1	2	1	3	-	2	3	1	3	8	20
Female	1,313	2	-	-	-	1	2	-	6	4	14	22
Black, both sexes	678	17	5	1	1	-	5	2	8	4	18	18
Male	372	8	4	-	1	-	3	-	3	4	10	9
Female	306	9	1	1	-	-	2	2	5	-	8	9
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Maine												
All races, both sexes	1,344	2	1	-	1	-	1	1	2	6	11	16
Male	653	1	1	-	1	-	-	-	2	2	5	13
Female	691	1	-	-	-	-	1	1	-	4	6	3
White, both sexes	1,335	2	-	-	1	-	1	1	1	6	11	16
Male	648	1	-	-	1	-	-	-	1	2	5	13
Female	687	1	-	-	-	-	1	1	-	4	6	3
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3,742	10	3	2	3	1	5	6	7	12	41	56
Male	1,726	5	1	-	1	1	4	4	4	7	20	22
Female	2,016	5	2	2	2	-	1	2	3	5	21	34
White, both sexes	3,041	5	-	1	-	-	3	2	4	8	17	31
Male	1,389	1	-	-	-	-	2	1	3	4	4	12
Female	1,652	4	-	1	-	-	1	1	1	4	13	19
Black, both sexes	640	5	1	1	3	1	2	4	3	4	22	24
Male	309	4	-	-	1	1	2	3	1	3	15	10
Female	331	1	1	1	2	-	-	1	2	1	7	14
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	54	-	2	-	-	-	-	-	-	-	2	-
Male	26	-	1	-	-	-	-	-	-	-	1	-
Female	28	-	1	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Massachusetts												
All races, both sexes	5,603	8	8	3	1	1	3	3	7	11	25	41
Male	2,453	6	4	1	1	-	1	2	2	4	14	20
Female	3,150	2	4	2	-	1	2	1	5	7	11	21
White, both sexes	5,420	6	7	3	1	1	2	3	6	11	20	37
Male	2,361	4	3	1	1	-	1	2	1	4	13	18
Female	3,059	2	4	2	-	1	1	1	5	7	7	19
Black, both sexes	132	1	1	-	-	-	1	-	-	-	5	4
Male	58	1	1	-	-	-	-	-	-	-	1	2
Female	74	-	-	-	-	-	1	-	-	-	4	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	1	-	-	-	-	-	-	1	-	-	-
Male	32	1	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	7,642	19	7	5	5	6	9	13	15	26	58	84
Male	3,673	9	5	2	3	5	6	9	6	10	33	41
Female	3,969	10	2	3	2	1	3	4	9	16	25	43
White, both sexes	6,920	11	3	1	3	1	6	8	12	17	42	62
Male	3,286	5	3	-	2	1	4	4	6	6	26	28
Female	3,634	6	-	1	1	-	2	4	6	11	16	34
Black, both sexes	655	8	3	4	2	5	3	5	2	8	13	20
Male	354	4	2	2	1	4	2	5	-	3	6	11
Female	301	4	1	2	1	1	1	-	2	5	7	9
American Indian 1/, both sexes	45	-	1	-	-	-	-	-	-	1	2	2
Male	24	-	-	-	-	-	-	-	-	1	1	2
Female	21	-	1	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	1	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Minnesota												
All races, both sexes	3,074	6	3	1	-	1	1	3	3	6	15	26
Male	1,479	4	3	1	-	1	1	1	1	3	7	15
Female	1,595	2	-	-	-	-	-	2	2	3	8	11
White, both sexes	2,991	2	2	-	-	-	1	2	2	5	14	23
Male	1,430	1	2	-	-	-	1	1	-	3	6	14
Female	1,561	1	-	-	-	-	-	1	2	2	8	9
Black, both sexes	40	2	1	1	-	1	-	1	-	1	1	3
Male	23	2	1	1	-	1	-	-	-	-	1	1
Female	17	-	-	-	-	-	-	1	-	1	-	2
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	2	-	-	-	-	-	-	1	-	-	-
Male	16	1	-	-	-	-	-	-	1	-	-	-
Female	10	1	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2,582	19	3	2	3	5	2	13	9	11	29	41
Male	1,300	14	3	2	2	3	-	6	6	2	14	17
Female	1,282	5	-	-	1	2	2	7	3	9	15	24
White, both sexes	1,970	2	2	1	-	2	-	6	4	3	14	19
Male	976	2	2	1	-	-	-	2	4	-	6	7
Female	994	-	-	-	-	2	-	4	-	3	8	12
Black, both sexes	601	16	1	1	3	3	2	7	5	8	15	22
Male	318	11	1	1	2	3	-	4	2	2	8	10
Female	283	5	-	-	1	-	2	3	3	6	7	12
American Indian 1/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Missouri												
All races, both sexes	5,308	19	5	1	7	3	4	12	12	11	50	78
Male	2,517	11	1	-	5	1	4	7	5	7	26	43
Female	2,791	8	4	1	2	2	-	5	7	4	24	35
White, both sexes	4,930	8	4	1	3	2	3	10	7	6	36	65
Male	2,338	4	1	-	2	1	3	6	3	3	17	36
Female	2,592	4	3	1	1	1	-	4	4	3	19	29
Black, both sexes	351	11	1	-	3	1	1	2	5	5	12	13
Male	172	7	-	-	3	-	1	1	2	4	7	7
Female	179	4	1	-	-	1	-	1	3	1	5	6
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	1	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	1	-	-	-	-	-	-	-
Montana												
All races, both sexes	947	1	2	1	-	-	1	1	2	2	3	6
Male	439	1	1	-	-	-	1	-	2	2	1	4
Female	508	-	1	1	-	-	-	1	-	-	2	2
White, both sexes	903	-	2	1	-	-	1	-	1	1	3	6
Male	414	-	1	-	-	-	1	-	1	1	1	4
Female	489	-	1	1	-	-	-	-	-	-	2	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	41	1	-	-	-	-	-	1	1	1	-	-
Male	23	1	-	-	-	-	-	-	1	1	-	-
Female	18	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Nebraska												
All races, both sexes	1,537	2	-	1	-	5	1	2	5	6	9	12
Male	751	1	-	-	-	1	-	1	2	3	3	6
Female	786	1	-	1	-	4	1	1	3	3	6	6
White, both sexes	1,494	2	-	1	-	4	1	2	3	6	8	8
Male	729	1	-	-	-	1	-	1	-	3	3	4
Female	765	1	-	1	-	3	1	1	3	3	5	4
Black, both sexes	23	-	-	-	-	1	-	-	2	-	1	3
Male	11	-	-	-	-	-	-	-	2	-	-	1
Female	12	-	-	-	-	1	-	-	-	-	1	2
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,823	8	5	-	-	-	5	4	4	12	14	32
Male	896	6	3	-	-	-	4	3	1	3	10	15
Female	927	2	2	-	-	-	1	1	3	9	4	17
White, both sexes	1,697	5	5	-	-	-	5	2	4	7	10	28
Male	829	4	3	-	-	-	4	1	1	3	8	13
Female	868	1	2	-	-	-	1	1	3	4	2	15
Black, both sexes	78	3	-	-	-	-	-	2	-	5	2	2
Male	44	2	-	-	-	-	-	2	-	-	1	1
Female	34	1	-	-	-	-	-	-	-	5	1	1
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	40	-	-	-	-	-	-	-	-	-	2	2
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	21	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
New Hampshire												
All races, both sexes	1,011	1	1	-	-	-	1	1	1	3	2	6
Male	489	1	-	-	-	-	-	1	-	2	-	3
Female	522	-	1	-	-	-	1	-	1	1	2	3
White, both sexes	1,005	1	1	-	-	-	1	1	1	3	2	6
Male	485	1	-	-	-	-	-	1	-	2	-	3
Female	520	-	1	-	-	-	1	-	1	1	2	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	5,496	9	4	3	6	6	5	13	8	24	40	64
Male	2,504	4	3	2	5	2	4	7	6	8	23	31
Female	2,992	5	1	1	1	4	1	6	2	16	17	33
White, both sexes	4,909	4	2	1	3	2	4	10	7	16	28	42
Male	2,225	2	2	1	2	-	3	6	5	4	17	23
Female	2,684	2	-	-	1	2	1	4	2	12	11	19
Black, both sexes	512	5	2	2	3	3	1	3	1	7	11	22
Male	228	2	1	1	3	1	1	1	1	3	6	8
Female	284	3	1	1	-	2	-	2	-	4	5	14
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	70	-	-	-	-	1	-	-	-	1	1	-
Male	48	-	-	-	-	1	-	-	-	1	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
New Mexico												
All races, both sexes	1,505	4	2	2	2	1	1	3	4	4	12	27
Male	747	2	1	-	2	1	-	2	4	3	5	16
Female	758	2	1	2	-	-	1	1	-	1	7	11
White, both sexes	1,417	4	1	2	2	1	1	2	3	4	10	23
Male	704	2	1	-	2	1	-	1	3	3	4	13
Female	713	2	-	2	-	-	1	1	-	1	6	10
Black, both sexes	14	-	-	-	-	-	-	1	-	-	1	1
Male	9	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	72	-	1	-	-	-	-	-	1	-	1	3
Male	33	-	-	-	-	-	-	-	1	-	1	3
Female	39	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	13,285	37	19	7	7	16	16	22	24	37	102	150
Male	6,116	23	10	6	3	10	9	9	13	25	57	80
Female	7,169	14	9	1	4	6	7	13	11	12	45	70
White, both sexes	11,596	18	10	4	3	12	13	14	13	24	67	112
Male	5,296	11	7	3	1	6	7	7	7	19	42	61
Female	6,300	7	3	1	2	6	6	7	6	5	25	51
Black, both sexes	1,392	17	7	2	4	3	3	8	10	11	34	35
Male	632	11	3	2	2	3	2	2	6	5	14	17
Female	760	6	4	-	2	-	1	6	4	6	20	18
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	276	2	2	1	-	1	-	-	1	2	1	3
Male	177	1	-	1	-	1	-	-	-	1	1	2
Female	99	1	2	-	-	-	-	-	1	1	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
North Carolina												
All races, both sexes	7,324	22	9	5	3	14	11	9	22	28	62	127
Male	3,425	18	5	3	2	6	3	5	9	13	32	62
Female	3,899	4	4	2	1	8	8	4	13	15	30	65
White, both sexes	6,351	11	6	4	1	11	3	5	11	15	37	82
Male	2,918	10	2	2	1	4	1	3	6	5	19	40
Female	3,433	1	4	2	-	7	2	2	5	10	18	42
Black, both sexes	919	11	3	1	2	3	7	4	11	13	25	40
Male	478	8	3	1	1	2	2	2	3	8	13	18
Female	441	3	-	-	1	1	5	2	8	5	12	22
American Indian 1/, both sexes	42	-	-	-	-	-	1	-	-	-	-	5
Male	24	-	-	-	-	-	-	-	-	-	-	4
Female	18	-	-	-	-	-	1	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	523	1	1	-	-	-	1	-	-	-	2	3
Male	276	1	1	-	-	-	1	-	-	-	1	2
Female	247	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	506	1	1	-	-	-	1	-	-	-	2	2
Male	269	1	1	-	-	-	1	-	-	-	1	1
Female	237	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Ohio												
All races, both sexes	10,085	24	11	4	2	6	7	13	16	41	63	151
Male	4,866	18	5	3	2	2	4	9	9	25	33	74
Female	5,219	6	6	1	-	4	3	4	7	16	30	77
White, both sexes	9,266	14	8	2	1	4	6	7	10	28	52	120
Male	4,421	12	3	2	1	1	3	5	5	15	29	59
Female	4,845	2	5	-	-	3	3	2	5	13	23	61
Black, both sexes	795	10	2	2	1	2	1	6	6	13	11	30
Male	433	6	2	1	1	1	1	4	4	10	4	15
Female	362	4	-	1	-	1	-	2	2	3	7	15
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	17	-	1	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	1	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3,612	10	4	2	4	4	6	5	4	23	33	45
Male	1,761	6	3	1	2	2	3	1	3	8	13	26
Female	1,851	4	1	1	2	2	3	4	1	15	20	19
White, both sexes	3,312	6	4	2	1	2	3	5	2	21	27	36
Male	1,620	4	3	1	1	-	1	1	1	8	9	22
Female	1,692	2	1	1	-	2	2	4	1	13	18	14
Black, both sexes	129	2	-	-	3	2	1	-	2	-	-	5
Male	60	1	-	-	1	2	1	-	2	-	-	3
Female	69	1	-	-	2	-	-	-	-	-	-	2
American Indian 1/, both sexes	169	2	-	-	-	-	1	-	-	2	6	4
Male	80	1	-	-	-	-	-	-	-	-	4	1
Female	89	1	-	-	-	-	1	-	-	2	2	3
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Oregon												
All races, both sexes	2,913	7	3	-	-	1	5	1	8	4	11	38
Male	1,415	3	1	-	-	1	3	1	4	3	7	21
Female	1,498	4	2	-	-	-	2	-	4	1	4	17
White, both sexes	2,842	4	3	-	-	1	5	1	8	4	11	36
Male	1,378	-	1	-	-	1	3	1	4	3	7	20
Female	1,464	4	2	-	-	-	2	-	4	1	4	16
Black, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	27	1	-	-	-	-	-	-	-	-	-	-
Male	10	1	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	2	-	-	-	-	-	-	-	-	-	1
Male	13	2	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	11,244	21	6	6	8	7	10	16	16	30	60	125
Male	5,342	10	3	2	3	6	6	10	5	11	38	64
Female	5,902	11	3	4	5	1	4	6	11	19	22	61
White, both sexes	10,428	15	6	5	7	2	7	12	13	24	44	94
Male	4,959	5	3	2	3	2	4	7	5	10	29	48
Female	5,469	10	3	3	4	-	3	5	8	14	15	46
Black, both sexes	776	6	-	1	1	5	3	4	3	6	16	31
Male	361	5	-	-	-	4	2	3	-	1	9	16
Female	415	1	-	1	1	1	1	1	3	5	7	15
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Rhode Island												
All races, both sexes	961	2	1	1	-	-	-	3	-	3	6	6
Male	425	2	-	1	-	-	-	2	-	1	3	1
Female	536	-	1	-	-	-	-	1	-	2	3	5
White, both sexes	930	2	1	1	-	-	-	3	-	2	6	6
Male	408	2	-	1	-	-	-	2	-	1	3	1
Female	522	-	1	-	-	-	-	1	-	1	3	5
Black, both sexes	25	-	-	-	-	-	-	-	-	1	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3,441	10	6	2	2	1	6	9	9	16	22	47
Male	1,700	6	3	2	1	1	4	4	3	10	12	24
Female	1,741	4	3	-	1	-	2	5	6	6	10	23
White, both sexes	2,854	4	2	1	-	1	2	8	4	10	12	30
Male	1,378	3	2	1	-	1	1	3	1	6	7	15
Female	1,476	1	-	-	-	-	1	5	3	4	5	15
Black, both sexes	575	6	4	1	2	-	4	1	5	6	10	16
Male	315	3	1	1	1	-	3	1	2	4	5	9
Female	260	3	3	-	1	-	1	-	3	2	5	7
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
South Dakota												
All races, both sexes	670	2	-	1	-	-	1	-	3	1	4	4
Male	340	1	-	1	-	-	1	-	2	-	3	1
Female	330	1	-	-	-	-	-	-	1	1	1	3
White, both sexes	640	2	-	-	-	-	1	-	2	1	3	4
Male	321	1	-	-	-	-	1	-	2	-	2	1
Female	319	1	-	-	-	-	-	-	-	1	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	30	-	-	1	-	-	-	-	1	-	1	-
Male	19	-	-	1	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5,547	20	5	5	4	9	13	10	11	13	52	93
Male	2,651	11	2	4	4	5	7	7	7	8	29	42
Female	2,896	9	3	1	-	4	6	3	4	5	23	51
White, both sexes	5,055	16	2	3	2	5	8	7	7	10	42	72
Male	2,393	9	1	2	2	4	4	5	5	5	24	33
Female	2,662	7	1	1	-	1	4	2	2	5	18	39
Black, both sexes	484	4	3	2	2	4	5	3	4	3	10	21
Male	253	2	1	2	2	1	3	2	2	3	5	9
Female	231	2	2	-	-	3	2	1	2	-	5	12
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Texas												
All races, both sexes	13,937	49	26	13	10	14	19	29	31	48	114	198
Male	6,640	29	12	8	5	9	14	12	15	25	58	99
Female	7,297	20	14	5	5	5	5	17	16	23	56	99
White, both sexes	12,727	37	19	11	5	12	11	24	25	36	90	161
Male	6,049	25	11	7	2	7	7	10	12	18	45	82
Female	6,678	12	8	4	3	5	4	14	13	18	45	79
Black, both sexes	1,067	11	5	2	5	2	5	5	5	12	23	32
Male	523	3	1	1	3	2	4	2	2	7	13	15
Female	544	8	4	1	2	-	1	3	3	5	10	17
American Indian 1/, both sexes	26	-	-	-	-	-	1	-	-	-	1	3
Male	13	-	-	-	-	-	1	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	117	1	2	-	-	-	2	-	1	-	-	2
Male	55	1	-	-	-	-	2	-	1	-	-	1
Female	62	-	2	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	1,203	4	3	1	3	4	1	4	3	3	9	15
Male	587	3	1	-	2	-	1	2	1	1	1	5
Female	616	1	2	1	1	4	-	2	2	2	8	10
White, both sexes	1,176	4	3	-	3	4	1	4	3	2	9	15
Male	573	3	1	-	2	-	1	2	1	-	1	5
Female	603	1	2	-	1	4	-	2	2	2	8	10
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	1	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Vermont												
All races, both sexes	489	1	-	-	-	1	-	-	-	1	-	2
Male	219	-	-	-	-	-	-	-	-	-	-	-
Female	270	1	-	-	-	1	-	-	-	1	-	2
White, both sexes	489	1	-	-	-	1	-	-	-	1	-	2
Male	219	-	-	-	-	-	-	-	-	-	-	-
Female	270	1	-	-	-	1	-	-	-	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5,459	20	8	3	-	4	7	8	10	16	39	83
Male	2,611	12	7	2	-	3	7	7	7	6	22	43
Female	2,848	8	1	1	-	1	-	1	3	10	17	40
White, both sexes	4,710	12	6	-	-	2	3	6	6	8	26	65
Male	2,220	6	5	-	-	2	3	5	4	3	14	35
Female	2,490	6	1	-	-	-	-	1	2	5	12	30
Black, both sexes	686	6	2	3	-	2	4	1	4	8	12	17
Male	352	5	2	2	-	1	4	1	3	3	7	8
Female	334	1	-	1	-	1	-	-	1	5	5	9
American Indian 1/, both sexes	6	1	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	57	1	-	-	-	-	-	1	-	-	1	-
Male	35	1	-	-	-	-	-	1	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Washington												
All races, both sexes	4,451	8	6	3	3	2	2	7	7	18	27	45
Male	2,084	6	2	1	2	1	1	5	3	8	10	25
Female	2,367	2	4	2	1	1	1	2	4	10	17	20
White, both sexes	4,210	4	5	2	3	-	2	7	7	15	23	40
Male	1,969	3	2	1	2	-	1	5	3	7	9	21
Female	2,241	1	3	1	1	-	1	2	4	8	14	19
Black, both sexes	77	1	1	1	-	1	-	-	-	1	2	3
Male	40	1	-	-	-	1	-	-	-	1	-	2
Female	37	-	1	1	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	55	2	-	-	-	1	-	-	-	-	1	-
Male	18	1	-	-	-	-	-	-	-	-	-	-
Female	37	1	-	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	109	1	-	-	-	-	-	-	-	2	1	2
Male	57	1	-	-	-	-	-	-	-	-	1	2
Female	52	-	-	-	-	-	-	-	-	2	-	-
West Virginia												
All races, both sexes	2,227	3	2	-	-	3	2	4	5	4	22	28
Male	1,121	3	1	-	-	-	-	4	1	-	11	15
Female	1,106	-	1	-	-	3	2	-	4	4	11	13
White, both sexes	2,189	3	1	-	-	3	2	3	5	3	20	28
Male	1,101	3	1	-	-	-	-	3	1	-	11	15
Female	1,088	-	-	-	-	3	2	-	4	3	9	13
Black, both sexes	37	-	1	-	-	-	-	1	-	1	2	-
Male	20	-	-	-	-	-	-	1	-	-	-	-
Female	17	-	1	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
Wisconsin												
All races, both sexes	4,205	15	1	-	2	2	2	2	8	7	25	31
Male	2,072	11	1	-	1	1	2	2	4	5	14	18
Female	2,133	4	-	-	1	1	-	-	4	2	11	13
White, both sexes	4,065	13	1	-	2	1	-	1	6	7	23	26
Male	2,003	9	1	-	1	-	-	1	3	5	12	16
Female	2,062	4	-	-	1	1	-	-	3	2	11	10
Black, both sexes	103	1	-	-	-	1	2	1	2	-	1	4
Male	53	1	-	-	-	1	2	1	1	-	1	1
Female	50	-	-	-	-	-	-	-	1	-	-	3
American Indian 1/, both sexes	24	1	-	-	-	-	-	-	-	-	1	1
Male	12	1	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	558	1	2	1	-	-	-	3	3	1	1	13
Male	278	-	1	-	-	-	-	1	1	1	1	10
Female	280	1	1	1	-	-	-	2	2	-	-	3
White, both sexes	549	1	2	1	-	-	-	3	3	1	1	12
Male	272	-	1	-	-	-	-	1	1	1	1	9
Female	277	1	1	1	-	-	-	2	2	-	-	3
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
United States												
All races, both sexes	4,861	7,680	11,559	16,937	24,651	35,628	43,728	38,745	24,159	8,885	1,839	7
Male	2,654	4,180	6,240	8,890	12,934	18,431	21,405	17,282	9,275	2,598	421	5
Female	2,207	3,500	5,319	8,047	11,717	17,197	22,323	21,463	14,884	6,287	1,418	2
White, both sexes	3,888	6,478	10,140	15,071	22,421	32,672	40,420	36,107	22,420	8,160	1,620	6
Male	2,113	3,543	5,397	7,837	11,676	16,835	19,698	16,046	8,569	2,354	358	4
Female	1,775	2,935	4,743	7,234	10,745	15,837	20,722	20,061	13,851	5,806	1,262	2
Black, both sexes	868	1,058	1,204	1,533	1,809	2,254	2,448	1,915	1,224	511	176	-
Male	483	561	719	860	1,015	1,214	1,220	844	438	145	44	-
Female	385	497	485	673	794	1,040	1,228	1,071	786	366	132	-
American Indian 1/, both sexes	52	63	91	121	126	166	146	113	57	21	9	-
Male	26	26	47	63	62	77	73	61	20	5	3	-
Female	26	37	44	58	64	89	73	52	37	16	6	-
Asian or Pacific Islander 2/, both sexes	53	81	124	212	295	536	714	610	458	193	34	1
Male	32	50	77	130	181	305	414	331	248	94	16	1
Female	21	31	47	82	114	231	300	279	210	99	18	-
Alabama												
All races, both sexes	105	185	244	375	540	675	758	605	330	116	24	-
Male	63	110	135	216	282	348	350	284	112	29	4	-
Female	42	75	109	159	258	327	408	321	218	87	20	-
White, both sexes	75	142	210	321	482	596	679	529	271	101	18	-
Male	47	85	110	180	244	298	314	251	88	23	4	-
Female	28	57	100	141	238	298	365	278	183	78	14	-
Black, both sexes	29	42	34	53	58	77	77	76	59	14	6	-
Male	15	25	25	36	38	49	36	33	24	6	-	-
Female	14	17	9	17	20	28	41	43	35	8	6	-
American Indian 1/, both sexes	-	1	-	1	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	1	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Alaska												
All races, both sexes	10	14	22	22	32	38	38	29	18	9	-	-
Male	6	7	15	11	12	21	16	13	6	5	-	-
Female	4	7	7	11	20	17	22	16	12	4	-	-
White, both sexes	5	10	15	12	25	32	26	21	10	6	-	-
Male	2	5	12	6	10	19	11	9	3	4	-	-
Female	3	5	3	6	15	13	15	12	7	2	-	-
Black, both sexes	2	1	-	-	1	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	3	5	8	6	5	11	4	8	3	-	-
Male	3	2	2	4	2	2	5	2	3	1	-	-
Female	-	1	3	4	4	3	6	2	5	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	-	1	1	3	-	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	1	1	-	1	1	2	-	-	-	-
Arizona												
All races, both sexes	111	164	246	415	574	828	898	771	393	120	28	-
Male	61	89	136	229	302	436	460	363	155	43	7	-
Female	50	75	110	186	272	392	438	408	238	77	21	-
White, both sexes	105	152	233	393	557	797	877	745	382	111	24	-
Male	58	84	130	217	294	418	443	349	151	39	5	-
Female	47	68	103	176	263	379	434	396	231	72	19	-
Black, both sexes	4	6	7	9	4	11	6	4	5	2	-	-
Male	2	3	5	5	3	6	6	1	-	-	-	-
Female	2	3	2	4	1	5	-	3	5	2	-	-
American Indian 1/, both sexes	2	6	6	11	10	16	13	15	4	2	4	-
Male	1	2	1	6	3	9	10	10	2	2	2	-
Female	1	4	5	5	7	7	3	5	2	-	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	3	4	2	7	2	5	-	-
Male	-	-	-	1	2	3	1	3	2	2	-	-
Female	-	-	-	1	1	1	1	4	-	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Arkansas												
All races, both sexes	82	107	182	221	302	399	515	428	282	108	25	-
Male	32	68	102	107	181	218	272	210	97	38	3	-
Female	50	39	80	114	121	181	243	218	185	70	22	-
White, both sexes	68	94	160	207	282	376	481	407	266	95	19	-
Male	26	62	90	99	169	203	255	198	91	34	2	-
Female	42	32	70	108	113	173	226	209	175	61	17	-
Black, both sexes	13	13	21	13	20	19	32	21	15	12	6	-
Male	6	6	12	8	12	13	16	12	6	3	1	-
Female	7	7	9	5	8	6	16	9	9	9	5	-
American Indian 1/, both sexes	1	-	-	1	-	4	2	-	-	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	1	-	-	1	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
California												
All races, both sexes	440	660	1,081	1,583	2,277	3,458	4,749	4,295	2,740	1,059	209	-
Male	240	366	608	821	1,160	1,744	2,252	1,948	1,090	339	60	-
Female	200	294	473	762	1,117	1,714	2,497	2,347	1,650	720	149	-
White, both sexes	357	550	913	1,367	1,984	3,026	4,182	3,806	2,422	916	187	-
Male	199	301	500	694	980	1,506	1,946	1,701	939	281	51	-
Female	158	249	413	673	1,004	1,520	2,236	2,105	1,483	635	136	-
Black, both sexes	63	77	107	120	151	171	195	163	96	46	9	-
Male	31	42	66	70	84	89	85	75	33	18	2	-
Female	32	35	41	50	67	82	110	88	63	28	7	-
American Indian 1/, both sexes	7	3	9	11	14	17	16	12	5	3	-	-
Male	2	3	7	5	12	7	8	4	2	-	-	-
Female	5	-	2	6	2	10	8	8	3	3	-	-
Asian or Pacific Islander 2/, both sexes	13	30	52	85	128	244	356	314	217	94	13	-
Male	8	20	35	52	84	142	213	168	116	40	7	-
Female	5	10	17	33	44	102	143	146	101	54	6	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Colorado												
All races, both sexes	79	116	153	237	363	512	631	536	331	117	24	-
Male	45	76	94	144	187	258	328	265	149	34	4	-
Female	34	40	59	93	176	254	303	271	182	83	20	-
White, both sexes	73	108	147	222	342	494	612	522	328	115	23	-
Male	42	72	90	137	175	252	319	258	148	34	4	-
Female	31	36	57	85	167	242	293	264	180	81	19	-
Black, both sexes	5	7	6	13	18	10	8	9	2	2	1	-
Male	2	4	4	6	10	4	4	4	1	-	-	-
Female	3	3	2	7	8	6	4	5	1	2	1	-
American Indian 1/, both sexes	1	1	-	1	-	3	-	3	-	-	-	-
Male	1	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	1	-	2	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	3	5	11	2	1	-	-	-
Male	-	-	-	1	2	1	5	2	-	-	-	-
Female	-	-	-	-	1	4	6	-	1	-	-	-
Connecticut												
All races, both sexes	50	73	118	153	219	408	596	630	449	184	27	1
Male	27	32	59	86	115	211	273	253	169	48	6	-
Female	23	41	59	67	104	197	323	377	280	136	21	1
White, both sexes	45	67	108	137	209	392	574	620	437	177	25	1
Male	25	28	53	77	109	201	260	250	165	47	6	-
Female	20	39	55	60	100	191	314	370	272	130	19	1
Black, both sexes	4	5	10	16	10	16	19	9	10	7	2	-
Male	1	3	6	9	6	10	12	2	2	1	-	-
Female	3	2	4	7	4	6	7	7	8	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	3	1	2	-	-	-
Male	1	1	-	-	-	-	1	1	2	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Delaware												
All races, both sexes	14	18	20	53	75	118	134	99	59	21	3	-
Male	8	7	15	28	38	64	59	43	24	8	-	-
Female	6	11	5	25	37	54	75	56	35	13	3	-
White, both sexes	12	15	15	41	65	105	124	90	57	20	3	-
Male	7	6	10	23	33	57	54	37	23	7	-	-
Female	5	9	5	18	32	48	70	53	34	13	3	-
Black, both sexes	2	3	5	11	9	12	9	8	2	1	-	-
Male	1	1	5	4	4	7	5	6	1	1	-	-
Female	1	2	-	7	5	5	4	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Dist. of Columbia												
All races, both sexes	6	13	12	19	18	39	53	40	29	10	3	-
Male	4	9	4	11	7	19	20	16	16	2	1	-
Female	2	4	8	8	11	20	33	24	13	8	2	-
White, both sexes	-	-	3	3	3	18	17	18	8	6	2	-
Male	-	-	2	3	3	11	6	8	5	1	-	-
Female	-	-	1	-	-	7	11	10	3	5	2	-
Black, both sexes	6	13	9	16	15	20	35	22	21	3	1	-
Male	4	9	2	8	4	7	13	8	11	1	1	-
Female	2	4	7	8	11	13	22	14	10	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Florida												
All races, both sexes	283	458	683	1,100	1,558	2,259	2,822	2,470	1,373	467	97	1
Male	155	268	378	593	816	1,201	1,396	1,104	548	145	19	1
Female	128	190	305	507	742	1,058	1,426	1,366	825	322	78	-
White, both sexes	241	395	607	1,007	1,457	2,132	2,693	2,371	1,302	440	91	1
Male	134	234	334	539	762	1,124	1,320	1,056	518	135	19	1
Female	107	161	273	468	695	1,008	1,373	1,315	784	305	72	-
Black, both sexes	40	62	74	88	94	113	114	91	62	26	6	-
Male	21	34	42	52	50	69	68	42	25	10	-	-
Female	19	28	32	36	44	44	46	49	37	16	6	-
American Indian 1/, both sexes	1	-	1	2	2	4	2	3	1	-	-	-
Male	-	-	1	1	1	2	1	2	-	-	-	-
Female	1	-	-	1	1	2	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	3	5	10	13	5	8	1	-	-
Male	-	-	1	1	3	6	7	4	5	-	-	-
Female	1	1	-	2	2	4	6	1	3	1	-	-
Georgia												
All races, both sexes	191	281	410	526	770	1,017	1,126	978	538	198	39	-
Male	91	148	225	285	407	530	524	414	176	55	6	-
Female	100	133	185	241	363	487	602	564	362	143	33	-
White, both sexes	133	208	343	450	658	883	973	868	470	159	27	-
Male	65	107	182	244	341	460	449	371	150	42	4	-
Female	68	101	161	206	317	423	524	497	320	117	23	-
Black, both sexes	56	73	65	73	107	131	140	104	63	37	12	-
Male	24	41	41	40	63	70	69	40	24	12	2	-
Female	32	32	24	33	44	61	71	64	39	25	10	-
American Indian 1/, both sexes	1	-	-	-	2	1	1	-	-	1	-	-
Male	1	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	3	3	2	12	6	5	1	-	-
Male	1	-	2	1	2	-	6	3	2	-	-	-
Female	-	-	-	2	1	2	6	3	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Hawaii												
All races, both sexes	18	24	27	46	64	108	155	148	125	56	11	-
Male	10	16	15	27	31	60	95	80	60	34	5	-
Female	8	8	12	19	33	48	60	68	65	22	6	-
White, both sexes	5	5	8	11	23	33	48	34	34	9	2	-
Male	3	2	7	6	12	18	26	18	16	3	-	-
Female	2	3	1	5	11	15	22	16	18	6	2	-
Black, both sexes	-	1	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	18	19	35	41	74	107	114	91	46	9	-
Male	7	14	8	21	19	41	69	62	44	30	5	-
Female	6	4	11	14	22	33	38	52	47	16	4	-
Idaho												
All races, both sexes	20	35	48	71	115	187	209	201	105	30	12	-
Male	11	18	29	44	58	105	117	103	46	12	4	-
Female	9	17	19	27	57	82	92	98	59	18	8	-
White, both sexes	20	34	47	71	115	186	209	199	104	30	12	-
Male	11	18	28	44	58	104	117	102	46	12	4	-
Female	9	16	19	27	57	82	92	97	58	18	8	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	1	-	1	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Illinois												
All races, both sexes	203	298	463	657	973	1,422	1,879	1,683	1,114	395	93	-
Male	105	150	259	342	517	740	933	765	421	119	16	-
Female	98	148	204	315	456	682	946	918	693	276	77	-
White, both sexes	142	230	394	574	846	1,269	1,702	1,572	1,050	375	82	-
Male	77	123	217	297	445	660	839	710	392	114	10	-
Female	65	107	177	277	401	609	863	862	658	261	72	-
Black, both sexes	57	67	68	77	119	136	157	93	54	18	8	-
Male	26	27	42	42	67	69	82	43	23	5	5	-
Female	31	40	26	35	52	67	75	50	31	13	3	-
American Indian 1/, both sexes	-	1	1	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	6	7	17	20	17	10	2	3	-
Male	2	-	-	3	5	11	12	11	6	-	1	-
Female	2	-	-	3	2	6	8	6	4	2	2	-
Indiana												
All races, both sexes	123	210	300	473	668	896	1,058	907	574	180	36	-
Male	73	117	160	256	358	461	508	423	223	55	8	-
Female	50	93	140	217	310	435	550	484	351	125	28	-
White, both sexes	108	196	285	453	639	857	1,009	878	561	175	32	-
Male	63	109	152	249	343	440	483	405	216	52	6	-
Female	45	87	133	204	296	417	526	473	345	123	26	-
Black, both sexes	14	13	15	19	28	38	49	28	12	5	4	-
Male	10	8	8	6	14	21	25	17	7	3	2	-
Female	4	5	7	13	14	17	24	11	5	2	2	-
American Indian 1/, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Iowa												
All races, both sexes	39	78	109	204	318	402	609	569	371	202	36	-
Male	18	43	60	111	180	217	345	262	141	73	5	-
Female	21	35	49	93	138	185	264	307	230	129	31	-
White, both sexes	37	78	108	201	311	399	603	567	367	202	36	-
Male	17	43	59	109	175	215	342	262	139	73	5	-
Female	20	35	49	92	136	184	261	305	228	129	31	-
Black, both sexes	1	-	1	2	4	2	5	2	2	-	-	-
Male	1	-	1	1	4	2	3	-	1	-	-	-
Female	-	-	-	1	-	-	2	2	1	-	-	-
American Indian 1/, both sexes	1	-	-	1	2	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	1	-	1	-	-	-
Kansas												
All races, both sexes	48	76	107	181	259	416	479	472	319	140	32	-
Male	28	41	43	101	138	207	241	217	141	41	6	-
Female	20	35	64	80	121	209	238	255	178	99	26	-
White, both sexes	44	71	102	169	244	400	454	460	310	134	30	-
Male	26	41	40	95	130	197	228	214	138	40	5	-
Female	18	30	62	74	114	203	226	246	172	94	25	-
Black, both sexes	1	4	3	6	8	13	18	7	6	6	2	-
Male	1	-	1	3	5	9	10	-	2	1	1	-
Female	-	4	2	3	3	4	8	7	4	5	1	-
American Indian 1/, both sexes	2	-	1	5	6	2	7	5	-	-	-	-
Male	1	-	1	2	3	-	3	3	-	-	-	-
Female	1	-	-	3	3	2	4	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	1	1	1	-	-	3	-	-	-
Male	-	-	1	1	-	1	-	-	1	-	-	-
Female	1	1	-	-	1	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Kentucky												
All races, both sexes	120	154	270	407	518	643	748	615	404	133	23	-
Male	56	90	141	228	277	325	366	247	152	36	7	-
Female	64	64	129	179	241	318	382	368	252	97	16	-
White, both sexes	108	143	256	386	495	614	715	598	379	122	23	-
Male	52	83	133	215	264	310	349	240	144	35	7	-
Female	56	60	123	171	231	304	366	358	235	87	16	-
Black, both sexes	12	11	12	20	23	27	31	17	23	11	-	-
Male	4	7	7	12	13	14	16	7	8	1	-	-
Female	8	4	5	8	10	13	15	10	15	10	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	1	1	-	1	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	1	-	-	-
Louisiana												
All races, both sexes	94	137	205	304	399	490	579	464	271	96	24	1
Male	59	74	107	158	228	253	299	205	107	33	5	1
Female	35	63	98	146	171	237	280	259	164	63	19	-
White, both sexes	58	102	148	238	316	392	494	394	221	76	15	1
Male	32	51	76	116	175	205	254	169	89	22	3	1
Female	26	51	72	122	141	187	240	225	132	54	12	-
Black, both sexes	36	35	55	66	82	96	83	67	50	20	9	-
Male	27	23	29	42	53	48	43	34	18	11	2	-
Female	9	12	26	24	29	48	40	33	32	9	7	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	2	1	2	-	-	-	-
Male	-	-	2	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Maine												
All races, both sexes	24	43	58	101	159	196	265	219	165	63	10	-
Male	15	24	34	47	84	108	133	106	60	15	2	-
Female	9	19	24	54	75	88	132	113	105	48	8	-
White, both sexes	24	43	57	100	159	193	264	219	164	63	10	-
Male	15	24	34	46	84	106	133	106	60	15	2	-
Female	9	19	23	54	75	87	131	113	104	48	8	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Maryland												
All races, both sexes	98	131	172	253	407	535	713	692	419	147	29	-
Male	55	72	89	123	202	284	330	287	177	33	5	-
Female	43	59	83	130	205	251	383	405	242	114	24	-
White, both sexes	51	103	128	190	325	453	609	590	360	137	24	-
Male	26	57	64	96	156	245	277	248	157	33	3	-
Female	25	46	64	94	169	208	332	342	203	104	21	-
Black, both sexes	47	28	44	56	73	74	90	96	49	10	3	-
Male	29	15	25	25	41	35	45	36	17	-	1	-
Female	18	13	19	31	32	39	45	60	32	10	2	-
American Indian 1/, both sexes	-	-	-	2	1	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	2	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	5	8	7	14	5	9	-	2	-
Male	-	-	-	2	5	3	8	2	3	-	1	-
Female	-	-	-	3	3	4	6	3	6	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Massachusetts												
All races, both sexes	68	150	232	307	519	853	1,085	1,109	773	325	71	-
Male	36	83	117	151	286	419	479	434	281	94	18	-
Female	32	67	115	156	233	434	606	675	492	231	53	-
White, both sexes	64	143	224	296	500	827	1,051	1,080	755	315	68	-
Male	33	79	113	144	272	403	465	423	273	91	17	-
Female	31	64	111	152	228	424	586	657	482	224	51	-
Black, both sexes	4	6	7	5	12	14	26	22	12	9	3	-
Male	3	3	4	2	7	9	11	7	3	3	1	-
Female	1	3	3	3	5	5	15	15	9	6	2	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	6	7	11	8	7	6	1	-	-
Male	-	-	-	5	7	6	3	4	5	-	-	-
Female	-	-	1	1	-	5	5	3	1	1	-	-
Michigan												
All races, both sexes	167	259	385	578	853	1,331	1,500	1,262	769	234	57	-
Male	96	151	205	292	426	677	742	572	302	62	19	-
Female	71	108	180	286	427	654	758	690	467	172	38	-
White, both sexes	126	211	330	531	783	1,220	1,393	1,158	728	225	49	-
Male	67	123	177	262	394	616	683	521	283	59	16	-
Female	59	88	153	269	389	604	710	637	445	166	33	-
Black, both sexes	35	46	51	36	62	98	102	98	38	9	7	-
Male	24	28	26	21	28	57	57	49	19	3	2	-
Female	11	18	25	15	34	41	45	49	19	6	5	-
American Indian 1/, both sexes	4	2	1	9	6	9	2	5	1	-	-	-
Male	3	-	-	7	4	3	1	2	-	-	-	-
Female	1	2	1	2	2	6	1	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	3	2	2	4	3	1	2	-	1	-
Male	2	-	2	2	-	1	1	-	-	-	1	-
Female	-	-	1	-	2	3	2	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Minnesota												
All races, both sexes	46	68	132	186	296	458	640	585	411	163	23	1
Male	24	43	69	99	166	251	311	254	178	42	4	1
Female	22	25	63	87	130	207	329	331	233	121	19	-
White, both sexes	42	65	124	178	288	447	627	581	404	161	23	-
Male	23	40	64	95	159	245	305	252	174	41	4	-
Female	19	25	60	83	129	202	322	329	230	120	19	-
Black, both sexes	3	2	3	5	2	4	6	2	2	-	-	-
Male	-	2	2	3	2	2	3	1	1	-	-	-
Female	3	-	1	2	-	2	3	1	1	-	-	-
American Indian 1/, both sexes	1	-	4	2	3	3	4	-	-	-	-	-
Male	1	-	2	-	2	2	3	-	-	-	-	-
Female	-	-	2	2	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	3	4	3	2	5	2	-	1
Male	-	1	1	1	3	2	-	1	3	1	-	1
Female	-	-	-	-	-	2	3	1	2	1	-	-
Mississippi												
All races, both sexes	74	121	161	216	308	431	429	377	205	101	21	1
Male	36	67	94	122	176	249	222	168	68	27	1	1
Female	38	54	67	94	132	182	207	209	137	74	20	-
White, both sexes	44	80	123	159	250	344	364	314	161	66	11	1
Male	18	41	68	92	135	201	193	135	49	18	1	1
Female	26	39	55	67	115	143	171	179	112	48	10	-
Black, both sexes	29	40	38	55	58	85	63	62	43	35	10	-
Male	17	25	26	29	41	47	29	32	19	9	-	-
Female	12	15	12	26	17	38	34	30	24	26	10	-
American Indian 1/, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	2	-	1	-	1	1	-	-	-
Male	1	1	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Missouri												
All races, both sexes	133	196	267	409	584	883	1,011	832	539	200	51	1
Male	75	102	131	201	319	445	504	362	210	45	13	-
Female	58	94	136	208	265	438	507	470	329	155	38	1
White, both sexes	118	181	241	374	545	835	949	789	513	191	48	1
Male	67	92	120	185	302	424	474	345	198	43	12	-
Female	51	89	121	189	243	411	475	444	315	148	36	1
Black, both sexes	13	14	23	33	38	40	59	42	23	9	3	-
Male	8	10	10	15	17	18	30	17	12	2	1	-
Female	5	4	13	18	21	22	29	25	11	7	2	-
American Indian 1/, both sexes	2	-	2	2	1	4	1	-	2	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	2	-	1	1	1	3	1	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	4	2	1	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	1	1	-	-	2	2	1	1	-	-	-
Montana												
All races, both sexes	18	24	51	70	117	152	186	163	105	37	5	-
Male	10	13	28	35	61	81	78	76	36	9	-	-
Female	8	11	23	35	56	71	108	87	69	28	5	-
White, both sexes	13	19	46	67	110	150	181	156	105	36	5	-
Male	7	11	24	33	58	80	76	71	36	9	-	-
Female	6	8	22	34	52	70	105	85	69	27	5	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	5	3	3	7	2	5	6	-	1	-	-
Male	3	2	3	2	3	1	2	4	-	-	-	-
Female	2	3	-	1	4	1	3	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Nebraska												
All races, both sexes	24	29	55	86	163	252	292	273	215	83	22	-
Male	12	17	30	43	90	142	151	132	85	25	7	-
Female	12	12	25	43	73	110	141	141	130	58	15	-
White, both sexes	21	29	52	81	157	245	287	271	212	83	21	-
Male	10	17	28	40	87	137	150	131	85	25	6	-
Female	11	12	24	41	70	108	137	140	127	58	15	-
Black, both sexes	2	-	-	2	3	2	2	1	3	-	1	-
Male	2	-	-	1	2	1	1	-	-	-	1	-
Female	-	-	-	1	1	1	1	1	3	-	-	-
American Indian 1/, both sexes	1	-	3	2	2	2	3	1	-	-	-	-
Male	-	-	2	1	1	1	-	1	-	-	-	-
Female	1	-	1	1	1	1	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	3	-	-	-	-	-	-
Male	-	-	-	1	-	3	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Nevada												
All races, both sexes	35	76	127	174	284	298	341	229	125	44	6	-
Male	19	50	58	87	144	156	172	101	48	14	2	-
Female	16	26	69	87	140	142	169	128	77	30	4	-
White, both sexes	30	65	117	166	265	277	324	220	119	42	6	-
Male	15	44	53	81	134	141	164	98	46	14	2	-
Female	15	21	64	85	131	136	160	122	73	28	4	-
Black, both sexes	5	8	8	4	13	9	10	4	2	1	-	-
Male	4	4	5	3	8	7	5	2	-	-	-	-
Female	1	4	3	1	5	2	5	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	2	1	2	-	1	2	-	-	-
Male	-	-	-	2	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	2	2	5	10	7	4	2	1	-	-
Male	-	2	-	1	2	7	3	1	1	-	-	-
Female	-	1	2	1	3	3	4	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
New Hampshire												
All races, both sexes	23	33	49	79	116	144	201	194	109	43	4	-
Male	13	18	24	35	56	85	110	87	42	11	1	-
Female	10	15	25	44	60	59	91	107	67	32	3	-
White, both sexes	23	33	49	79	115	142	201	193	108	42	4	-
Male	13	18	24	35	55	83	110	87	41	11	1	-
Female	10	15	25	44	60	59	91	106	67	31	3	-
Black, both sexes	-	-	-	-	-	2	-	-	1	1	-	-
Male	-	-	-	-	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
New Jersey												
All races, both sexes	104	150	205	304	549	874	1,117	1,078	657	235	41	-
Male	58	83	115	165	270	407	527	461	248	65	10	-
Female	46	67	90	139	279	467	590	617	409	170	31	-
White, both sexes	84	118	167	252	489	782	1,032	1,012	603	216	35	-
Male	45	70	92	135	238	360	487	434	231	59	9	-
Female	39	48	75	117	251	422	545	578	372	157	26	-
Black, both sexes	20	32	33	47	50	78	73	56	41	16	6	-
Male	13	13	19	25	24	38	34	22	8	3	1	-
Female	7	19	14	22	26	40	39	34	33	13	5	-
American Indian 1/, both sexes	-	-	2	-	1	2	-	-	-	-	-	-
Male	-	-	2	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	5	9	12	12	10	13	3	-	-
Male	-	-	2	5	8	8	6	5	9	3	-	-
Female	-	-	1	-	1	4	6	5	4	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
New Mexico												
All races, both sexes	29	44	90	98	159	249	333	210	161	51	19	-
Male	15	22	49	58	74	125	165	111	69	17	6	-
Female	14	22	41	40	85	124	168	99	92	34	13	-
White, both sexes	29	39	83	94	154	241	313	196	148	49	18	-
Male	15	19	46	55	72	123	154	103	64	17	6	-
Female	14	20	37	39	82	118	159	93	84	32	12	-
Black, both sexes	-	1	-	-	2	2	1	2	3	-	-	-
Male	-	1	-	-	1	2	1	1	2	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	4	7	4	3	6	18	11	10	2	1	-
Male	-	2	3	3	1	-	10	6	3	-	-	-
Female	-	2	4	1	2	6	8	5	7	2	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
New York												
All races, both sexes	272	357	644	862	1,217	1,966	2,523	2,524	1,658	695	130	-
Male	150	188	363	445	601	1,011	1,211	1,081	591	196	34	-
Female	122	169	281	417	616	955	1,312	1,443	1,067	499	96	-
White, both sexes	205	272	538	712	1,055	1,763	2,248	2,275	1,497	626	115	-
Male	114	148	293	363	520	895	1,082	974	527	179	30	-
Female	91	124	245	349	535	868	1,166	1,301	970	447	85	-
Black, both sexes	62	77	93	131	132	164	220	195	122	50	12	-
Male	34	34	58	68	63	91	96	74	36	8	3	-
Female	28	43	35	63	69	73	124	121	86	42	9	-
American Indian 1/, both sexes	-	1	1	2	4	8	1	3	1	-	-	-
Male	-	1	1	1	2	3	1	2	-	-	-	-
Female	-	-	-	1	2	5	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	5	7	12	17	26	31	54	51	38	19	3	-
Male	2	5	11	13	16	22	32	31	28	9	1	-
Female	3	2	1	4	10	9	22	20	10	10	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
North Carolina												
All races, both sexes	149	310	456	643	826	1,164	1,361	1,112	686	240	65	-
Male	89	163	247	336	409	577	648	475	243	69	11	-
Female	60	147	209	307	417	587	713	637	443	171	54	-
White, both sexes	105	247	370	560	729	1,048	1,229	1,001	614	208	54	-
Male	60	122	196	290	357	523	580	419	213	57	8	-
Female	45	125	174	270	372	525	649	582	401	151	46	-
Black, both sexes	40	61	80	76	93	111	125	102	70	31	10	-
Male	25	40	49	42	50	52	66	52	28	11	2	-
Female	15	21	31	34	43	59	59	50	42	20	8	-
American Indian 1/, both sexes	3	2	3	6	3	3	4	9	2	-	1	-
Male	3	1	2	4	1	1	1	4	2	-	1	-
Female	-	1	1	2	2	2	3	5	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	1	1	2	3	-	-	1	-	-
Male	1	-	-	-	1	1	1	-	-	1	-	-
Female	-	-	3	1	-	1	2	-	-	-	-	-
North Dakota												
All races, both sexes	4	10	17	24	50	88	90	106	88	32	6	-
Male	2	5	9	15	28	52	57	54	41	6	1	-
Female	2	5	8	9	22	36	33	52	47	26	5	-
White, both sexes	2	10	15	21	48	85	87	105	88	32	6	-
Male	1	5	8	15	27	51	55	54	41	6	1	-
Female	1	5	7	6	21	34	32	51	47	26	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	2	3	2	3	3	1	-	-	-	-
Male	1	-	1	-	1	1	2	-	-	-	-	-
Female	1	-	1	3	1	2	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Ohio												
All races, both sexes	235	375	539	790	1,150	1,702	1,890	1,661	994	346	65	-
Male	138	201	298	403	606	892	925	739	378	87	15	-
Female	97	174	241	387	544	810	965	922	616	259	50	-
White, both sexes	198	313	487	710	1,073	1,556	1,774	1,571	943	331	58	-
Male	115	170	260	354	561	811	864	694	359	86	12	-
Female	83	143	227	356	512	745	910	877	584	245	46	-
Black, both sexes	36	61	49	77	75	145	111	86	50	15	6	-
Male	22	31	37	46	45	81	57	43	18	1	3	-
Female	14	30	12	31	30	64	54	43	32	14	3	-
American Indian 1/, both sexes	-	-	2	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	2	-	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	3	1	-	4	3	1	-	1	-
Male	1	-	1	3	-	-	4	1	1	-	-	-
Female	-	1	-	-	1	-	-	2	-	-	1	-
Oklahoma												
All races, both sexes	98	169	230	357	460	618	611	513	291	102	23	-
Male	47	93	115	180	239	334	304	226	117	30	8	-
Female	51	76	115	177	221	284	307	287	174	72	15	-
White, both sexes	83	148	206	320	427	573	570	486	276	95	19	-
Male	40	82	102	164	225	310	286	212	112	28	8	-
Female	43	66	104	156	202	263	284	274	164	67	11	-
Black, both sexes	10	9	6	17	11	16	17	15	7	3	3	-
Male	4	5	2	7	6	7	8	7	3	1	-	-
Female	6	4	4	10	5	9	9	8	4	2	3	-
American Indian 1/, both sexes	5	12	18	20	22	29	23	12	8	4	1	-
Male	3	6	11	9	8	17	10	7	2	1	-	-
Female	2	6	7	11	14	12	13	5	6	3	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Oregon												
All races, both sexes	58	115	147	207	360	480	539	482	339	89	19	-
Male	36	62	69	103	180	236	278	231	139	33	4	-
Female	22	53	78	104	180	244	261	251	200	56	15	-
White, both sexes	56	110	139	203	353	465	530	477	332	86	18	-
Male	34	60	66	101	176	227	273	229	136	32	4	-
Female	22	50	73	102	177	238	257	248	196	54	14	-
Black, both sexes	1	3	4	2	3	3	5	2	1	1	-	-
Male	1	1	2	1	2	2	3	-	-	1	-	-
Female	-	2	2	1	1	1	2	2	1	-	-	-
American Indian 1/, both sexes	-	2	4	2	4	7	2	1	2	1	1	-
Male	-	1	1	1	2	2	1	-	1	-	-	-
Female	-	1	3	1	2	5	1	1	1	1	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	5	2	2	4	1	-	-
Male	1	-	-	-	-	5	1	2	2	-	-	-
Female	-	-	-	-	-	-	1	-	2	1	-	-
Pennsylvania												
All races, both sexes	187	315	488	667	1,093	1,764	2,331	2,199	1,335	456	104	-
Male	101	172	274	363	560	927	1,152	1,000	506	109	20	-
Female	86	143	214	304	533	837	1,179	1,199	829	347	84	-
White, both sexes	144	262	439	594	1,005	1,640	2,211	2,105	1,266	435	98	-
Male	75	147	248	322	514	866	1,094	966	486	104	19	-
Female	69	115	191	272	491	774	1,117	1,139	780	331	79	-
Black, both sexes	39	52	48	66	86	118	109	91	66	19	6	-
Male	23	25	26	36	44	58	51	34	18	5	1	-
Female	16	27	22	30	42	60	58	57	48	14	5	-
American Indian 1/, both sexes	2	-	1	1	-	1	1	-	-	1	-	-
Male	1	-	-	1	-	1	1	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	6	2	5	10	3	3	1	-	-
Male	2	-	-	4	2	2	6	-	2	-	-	-
Female	-	1	-	2	-	3	4	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Rhode Island												
All races, both sexes	15	25	42	61	92	140	203	193	114	40	14	-
Male	8	16	15	29	47	74	88	84	45	5	4	-
Female	7	9	27	32	45	66	115	109	69	35	10	-
White, both sexes	15	24	40	57	88	135	199	187	112	39	13	-
Male	8	15	14	26	44	71	87	82	43	4	4	-
Female	7	9	26	31	44	64	112	105	69	35	9	-
Black, both sexes	-	1	2	4	3	4	3	5	1	-	1	-
Male	-	1	1	3	2	2	-	2	1	-	-	-
Female	-	-	1	1	1	2	3	3	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
South Carolina												
All races, both sexes	96	159	234	327	439	552	631	494	259	94	26	-
Male	48	92	128	188	245	302	312	187	98	23	7	-
Female	48	67	106	139	194	250	319	307	161	71	19	-
White, both sexes	66	115	182	258	369	485	554	431	218	85	17	-
Male	33	62	94	145	202	262	273	158	84	21	4	-
Female	33	53	88	113	167	223	281	273	134	64	13	-
Black, both sexes	30	44	50	68	70	64	75	61	40	9	9	-
Male	15	30	32	42	43	40	38	27	13	2	3	-
Female	15	14	18	26	27	24	37	34	27	7	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	3	-	1	1	-	-	-
Male	-	-	2	1	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	3	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
South Dakota												
All races, both sexes	10	21	23	34	72	89	127	133	93	43	9	-
Male	4	6	10	17	49	49	77	69	36	12	2	-
Female	6	15	13	17	23	40	50	64	57	31	7	-
White, both sexes	8	16	20	33	66	84	125	132	91	43	9	-
Male	3	5	8	16	43	45	75	69	36	12	2	-
Female	5	11	12	17	23	39	50	63	55	31	7	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	5	3	1	6	5	2	1	2	-	-	-
Male	1	1	2	1	6	4	2	-	-	-	-	-
Female	1	4	1	-	-	1	-	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	150	244	353	508	647	875	994	855	489	164	33	-
Male	79	118	182	257	340	459	502	358	174	51	5	-
Female	71	126	171	251	307	416	492	497	315	113	28	-
White, both sexes	111	211	312	471	592	814	924	807	462	151	26	-
Male	56	106	154	235	310	423	464	338	161	48	4	-
Female	55	105	158	236	282	391	460	469	301	103	22	-
Black, both sexes	39	33	39	37	53	61	69	47	26	12	7	-
Male	23	12	27	22	29	36	37	20	12	2	1	-
Female	16	21	12	15	24	25	32	27	14	10	6	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	1	1	1	1	-	-
Male	-	-	1	-	1	-	1	-	1	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Texas												
All races, both sexes	321	529	793	1,175	1,625	2,204	2,658	2,157	1,335	500	89	-
Male	182	263	421	601	870	1,106	1,272	969	491	155	24	-
Female	139	266	372	574	755	1,098	1,386	1,188	844	345	65	-
White, both sexes	257	452	718	1,043	1,491	2,042	2,477	2,026	1,254	459	77	-
Male	149	233	381	527	795	1,024	1,184	910	461	140	19	-
Female	108	219	337	516	696	1,018	1,293	1,116	793	319	58	-
Black, both sexes	58	66	72	120	116	135	158	113	71	40	11	-
Male	31	26	37	71	67	70	77	49	23	14	5	-
Female	27	40	35	49	49	65	81	64	48	26	6	-
American Indian 1/, both sexes	2	3	-	5	1	2	6	1	1	-	-	-
Male	-	1	-	2	1	1	4	1	1	-	-	-
Female	2	2	-	3	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	8	3	7	17	25	17	17	9	1	1	-
Male	2	3	3	1	7	11	7	9	6	1	-	-
Female	2	5	-	6	10	14	10	8	3	-	1	-
Utah												
All races, both sexes	28	60	59	86	141	192	221	204	119	37	6	-
Male	20	20	27	43	84	101	108	104	52	10	1	-
Female	8	40	32	43	57	91	113	100	67	27	5	-
White, both sexes	27	59	56	85	141	186	213	202	118	37	4	-
Male	19	19	26	42	84	98	104	103	51	10	1	-
Female	8	40	30	43	57	88	109	99	67	27	3	-
Black, both sexes	-	-	1	-	-	2	2	-	-	-	1	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	1	-
American Indian 1/, both sexes	1	-	2	-	-	3	1	2	-	-	1	-
Male	1	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	2	1	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	1	5	-	1	-	-	-
Male	-	1	-	1	-	-	3	-	1	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Vermont												
All races, both sexes	5	13	34	25	50	80	117	78	49	27	6	-
Male	2	8	19	15	25	44	50	33	16	6	1	-
Female	3	5	15	10	25	36	67	45	33	21	5	-
White, both sexes	5	13	34	25	50	80	117	78	49	27	6	-
Male	2	8	19	15	25	44	50	33	16	6	1	-
Female	3	5	15	10	25	36	67	45	33	21	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	106	184	261	394	664	866	1,061	902	562	213	48	-
Male	58	96	148	201	340	437	544	404	200	56	11	-
Female	48	88	113	193	324	429	517	498	362	157	37	-
White, both sexes	76	152	219	322	575	753	932	822	498	188	39	-
Male	41	78	124	159	284	373	475	365	183	52	9	-
Female	35	74	95	163	291	380	457	457	315	136	30	-
Black, both sexes	30	28	40	69	81	101	118	71	59	22	8	-
Male	17	16	22	41	50	56	62	34	14	2	2	-
Female	13	12	18	28	31	45	56	37	45	20	6	-
American Indian 1/, both sexes	-	1	-	-	1	1	-	-	1	-	-	-
Male	-	1	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	2	3	7	11	11	9	4	3	1	-
Male	-	1	2	1	5	7	7	5	2	2	-	-
Female	-	2	-	2	2	4	4	4	2	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Washington												
All races, both sexes	114	157	232	346	453	686	860	792	480	171	31	1
Male	60	80	110	178	241	334	402	367	182	58	7	1
Female	54	77	122	168	212	352	458	425	298	113	24	-
White, both sexes	105	142	214	327	430	642	819	764	463	164	31	1
Male	56	76	104	167	230	315	375	353	176	55	7	1
Female	49	66	110	160	200	327	444	411	287	109	24	-
Black, both sexes	7	6	8	3	5	10	14	9	3	2	-	-
Male	4	2	4	2	4	3	10	4	1	1	-	-
Female	3	4	4	1	1	7	4	5	2	1	-	-
American Indian 1/, both sexes	1	7	5	7	8	10	5	5	3	-	-	-
Male	-	1	1	4	2	2	3	3	1	-	-	-
Female	1	6	4	3	6	8	2	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	5	9	10	24	22	14	11	5	-	-
Male	-	1	1	5	5	14	14	7	4	2	-	-
Female	1	1	4	4	5	10	8	7	7	3	-	-
West Virginia												
All races, both sexes	50	85	113	214	293	404	390	318	205	64	18	-
Male	28	52	59	106	163	220	200	148	81	25	4	-
Female	22	33	54	108	130	184	190	170	124	39	14	-
White, both sexes	49	85	111	209	289	402	385	313	204	59	15	-
Male	28	52	57	104	159	218	198	145	81	22	3	-
Female	21	33	54	105	130	184	187	168	123	37	12	-
Black, both sexes	1	-	2	5	4	2	5	5	-	5	3	-
Male	-	-	2	2	4	2	2	3	-	3	1	-
Female	1	-	-	3	-	-	3	2	-	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J09-J98,U04)												
Wisconsin												
All races, both sexes	73	107	186	274	396	708	849	769	522	191	35	-
Male	41	61	105	138	221	387	427	349	219	57	8	-
Female	32	46	81	136	175	321	422	420	303	134	27	-
White, both sexes	61	99	173	258	382	683	837	757	514	186	35	-
Male	34	57	98	129	216	375	423	342	216	57	8	-
Female	27	42	75	129	166	308	414	415	298	129	27	-
Black, both sexes	11	7	10	12	13	18	7	5	7	1	-	-
Male	7	4	6	8	5	7	3	2	3	-	-	-
Female	4	3	4	4	8	11	4	3	4	1	-	-
American Indian 1/, both sexes	1	1	2	3	1	3	2	6	-	2	-	-
Male	-	-	1	1	-	3	-	4	-	-	-	-
Female	1	1	1	2	1	-	2	2	-	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	4	3	1	1	2	-	-
Male	-	-	-	-	-	2	1	1	-	-	-	-
Female	-	-	1	1	-	2	2	-	1	2	-	-
Wyoming												
All races, both sexes	11	20	24	35	67	79	123	90	63	14	7	-
Male	5	10	13	16	38	42	70	38	29	2	-	-
Female	6	10	11	19	29	37	53	52	34	12	7	-
White, both sexes	10	19	24	33	65	79	122	90	62	14	7	-
Male	5	9	13	14	36	42	70	38	29	2	-	-
Female	5	10	11	19	29	37	52	52	33	12	7	-
Black, both sexes	1	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	1	2	-	1	-	-	-	-	-
Male	-	1	-	1	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
United States												
All races, both sexes	849	18	19	16	9	6	7	3	3	3	7	7
Male	362	11	12	5	6	2	3	2	2	1	3	4
Female	487	7	7	11	3	4	4	1	1	2	4	3
White, both sexes	788	13	11	12	8	4	5	3	1	2	6	5
Male	332	9	8	4	5	1	1	2	1	-	3	4
Female	456	4	3	8	3	3	4	1	-	2	3	1
Black, both sexes	41	5	4	-	1	1	2	-	2	-	1	2
Male	21	2	3	-	1	-	2	-	1	-	-	-
Female	20	3	1	-	-	1	-	-	1	-	1	2
American Indian 1/, both sexes	3	-	-	1	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	4	3	-	1	-	-	-	-	-	-
Male	8	-	1	1	-	1	-	-	-	-	-	-
Female	9	-	3	2	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	15	-	-	-	-	-	1	-	-	-	-	-
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	18	-	-	1	-	-	-	-	-	1	-	-
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	1	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Arkansas												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	46	2	2	3	-	2	-	1	-	1	1	1
Male	22	2	1	1	-	2	-	1	-	-	-	1
Female	24	-	1	2	-	-	-	-	-	1	1	-
White, both sexes	39	1	2	1	-	1	-	1	-	1	-	1
Male	17	1	1	-	-	1	-	1	-	-	-	1
Female	22	-	1	1	-	-	-	-	-	1	-	-
Black, both sexes	3	1	-	-	-	-	-	-	-	-	1	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	2	-	1	-	-	-	-	-	-
Male	3	-	-	1	-	1	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Colorado												
All races, both sexes	19	1	-	-	-	-	-	-	-	-	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	1	-	-	-	-	-	-	-	-	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	19	1	-	-	-	1	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	1	-	-	-	1	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Delaware												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Florida												
All races, both sexes	28	2	2	2	-	1	-	-	-	-	-	-
Male	14	-	2	1	-	-	-	-	-	-	-	-
Female	14	2	-	1	-	1	-	-	-	-	-	-
White, both sexes	25	2	1	2	-	1	-	-	-	-	-	-
Male	12	-	1	1	-	-	-	-	-	-	-	-
Female	13	2	-	1	-	1	-	-	-	-	-	-
Black, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	16	-	-	-	1	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	1	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Hawaii												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	6	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Illinois												
All races, both sexes	39	-	-	2	-	-	-	-	2	-	1	-
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	25	-	-	2	-	-	-	-	1	-	1	-
White, both sexes	35	-	-	1	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	1	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	2	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	17	-	-	-	1	-	-	-	-	-	-	-
Male	12	-	-	-	1	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	1	-	-	-	-	-	-	-
Male	12	-	-	-	1	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Iowa												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	19	-	1	1	-	-	-	-	-	-	-	-
Male	5	-	1	-	-	-	-	-	-	-	-	-
Female	14	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	1	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Kentucky												
All races, both sexes	18	1	1	-	-	-	-	-	1	-	1	-
Male	11	-	1	-	-	-	-	-	1	-	1	-
Female	7	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	1	1	-	-	-	-	-	1	-	1	-
Male	11	-	1	-	-	-	-	-	1	-	1	-
Female	7	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	6	1	-	-	1	-	-	-	-	1	1	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	1	1	-
White, both sexes	5	-	-	-	1	-	-	-	-	1	1	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Maine												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	15	-	2	1	-	-	-	-	-	-	-	2
Male	6	-	1	-	-	-	-	-	-	-	-	1
Female	9	-	1	1	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	1	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	2	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Massachusetts												
All races, both sexes	22	-	1	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	1	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	23	2	-	-	-	-	-	-	-	-	-	-
Male	10	1	-	-	-	-	-	-	-	-	-	-
Female	13	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Minnesota												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Missouri												
All races, both sexes	28	-	-	-	1	-	-	-	-	-	2	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	1	-	-	-	-	-	1	-
White, both sexes	28	-	-	-	1	-	-	-	-	-	2	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	1	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Nebraska												
All races, both sexes	14	1	-	-	-	-	-	-	-	-	-	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	1	-	-	-	-	-	-	-	-	-	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
New Hampshire												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
New Mexico												
All races, both sexes	11	1	1	1	-	-	-	-	-	-	-	-
Male	5	1	1	-	-	-	-	-	-	-	-	-
Female	6	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	11	1	1	1	-	-	-	-	-	-	-	-
Male	5	1	1	-	-	-	-	-	-	-	-	-
Female	6	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	41	1	2	-	1	1	1	-	-	-	-	-
Male	14	1	1	-	1	-	1	-	-	-	-	-
Female	27	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	38	1	1	-	1	1	-	-	-	-	-	-
Male	12	1	1	-	1	-	-	-	-	-	-	-
Female	26	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
North Carolina												
All races, both sexes	21	1	-	1	1	1	-	-	-	-	-	-
Male	8	1	-	1	1	-	-	-	-	-	-	-
Female	13	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	21	1	-	1	1	1	-	-	-	-	-	-
Male	8	1	-	1	1	-	-	-	-	-	-	-
Female	13	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Ohio												
All races, both sexes	37	-	1	-	-	-	-	-	-	-	-	-
Male	12	-	1	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	1	-	-	-	-	-	-	-	-	-
Male	11	-	1	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Oregon												
All races, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	21	-	-	1	-	-	1	-	-	-	-	-
Male	8	-	-	-	-	-	1	-	-	-	-	-
Female	13	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	1	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Rhode Island												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	15	-	1	1	-	-	-	-	-	-	-	-
Male	9	-	-	1	-	-	-	-	-	-	-	-
Female	6	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	1	-	-	-	-	-	-	-	-
Male	5	-	-	1	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	1	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
South Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	17	1	-	1	-	-	1	-	-	-	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	17	1	-	1	-	-	1	-	-	-	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Texas												
All races, both sexes	59	1	3	1	1	-	-	1	-	-	-	3
Male	19	1	1	1	1	-	-	1	-	-	-	1
Female	40	-	2	-	-	-	-	-	-	-	-	2
White, both sexes	51	1	2	1	-	-	-	1	-	-	-	2
Male	18	1	1	1	-	-	-	1	-	-	-	1
Female	33	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	1	-	-	-	-	-	-	1
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	11	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Vermont												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	34	1	1	-	-	-	-	-	-	-	-	-
Male	12	-	1	-	-	-	-	-	-	-	-	-
Female	22	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	1	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	21	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Washington												
All races, both sexes	23	-	-	-	1	-	-	-	-	-	-	1
Male	10	-	-	-	1	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	1	-	-	-	-	-	-	1
Male	9	-	-	-	1	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	11	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J09-J11)												
Wisconsin												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	1	-
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
United States												
All races, both sexes	19	21	22	31	50	93	143	156	133	67	16	-
Male	9	10	10	15	27	39	65	62	49	22	3	-
Female	10	11	12	16	23	54	78	94	84	45	13	-
White, both sexes	18	21	20	27	48	90	135	153	125	66	15	-
Male	8	10	9	14	26	37	60	62	44	22	2	-
Female	10	11	11	13	22	53	75	91	81	44	13	-
Black, both sexes	1	-	2	2	1	2	8	1	5	1	-	-
Male	1	-	1	1	-	1	5	-	3	-	-	-
Female	-	-	1	1	1	1	3	1	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	1	1	-	1	3	-	1	-
Male	-	-	-	-	1	1	-	-	2	-	1	-
Female	-	-	-	2	-	-	-	1	1	-	-	-
Alabama												
All races, both sexes	-	-	1	-	1	2	4	3	1	2	-	-
Male	-	-	1	-	-	-	2	1	-	1	-	-
Female	-	-	-	-	1	2	2	2	1	1	-	-
White, both sexes	-	-	-	-	1	2	4	3	1	2	-	-
Male	-	-	-	-	-	-	2	1	-	1	-	-
Female	-	-	-	-	1	2	2	2	1	1	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Alaska												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	3	-	-	1	2	2	3	4	-	-	-
Male	-	2	-	-	1	-	2	3	1	-	-	-
Female	1	1	-	-	-	2	-	-	3	-	-	-
White, both sexes	1	3	-	-	-	2	2	3	4	-	-	-
Male	-	2	-	-	-	-	2	3	1	-	-	-
Female	1	1	-	-	-	2	-	-	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Arkansas												
All races, both sexes	-	-	-	1	2	-	2	2	2	2	-	-
Male	-	-	-	-	1	-	2	-	1	-	-	-
Female	-	-	-	1	1	-	-	2	1	2	-	-
White, both sexes	-	-	-	1	2	-	1	2	2	2	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	1	1	-	-	2	1	2	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4	-	-	3	2	3	5	8	5	2	1	-
Male	1	-	-	2	1	1	3	4	1	-	1	-
Female	3	-	-	1	1	2	2	4	4	2	-	-
White, both sexes	4	-	-	3	2	3	5	8	4	2	-	-
Male	1	-	-	2	1	1	3	4	-	-	-	-
Female	3	-	-	1	1	2	2	4	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Colorado												
All races, both sexes	-	-	-	-	1	-	4	6	3	3	1	-
Male	-	-	-	-	1	-	1	2	2	-	-	-
Female	-	-	-	-	-	-	3	4	1	3	1	-
White, both sexes	-	-	-	-	1	-	4	6	3	3	1	-
Male	-	-	-	-	1	-	1	2	2	-	-	-
Female	-	-	-	-	-	-	3	4	1	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	2	4	2	4	5	-	-
Male	-	-	-	-	-	2	1	-	2	2	-	-
Female	-	-	-	-	-	-	3	2	2	3	-	-
White, both sexes	-	-	-	-	-	2	4	2	4	5	-	-
Male	-	-	-	-	-	2	1	-	2	2	-	-
Female	-	-	-	-	-	-	3	2	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Delaware												
All races, both sexes	-	-	-	1	-	-	-	1	-	2	-	-
Male	-	-	-	1	-	-	-	-	-	2	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	2	-	-
Male	-	-	-	1	-	-	-	-	-	2	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Florida												
All races, both sexes	-	1	3	3	2	3	3	4	2	-	-	-
Male	-	1	-	2	2	2	1	3	-	-	-	-
Female	-	-	3	1	-	1	2	1	2	-	-	-
White, both sexes	-	1	3	2	2	3	2	4	2	-	-	-
Male	-	1	-	2	2	2	-	3	-	-	-	-
Female	-	-	3	-	-	1	2	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	1	-	1	2	-	5	2	3	1	-	-
Male	-	-	-	-	-	-	2	2	-	-	-	-
Female	-	1	-	1	2	-	3	-	3	1	-	-
White, both sexes	-	1	-	1	2	-	4	2	3	-	-	-
Male	-	-	-	-	-	-	2	2	-	-	-	-
Female	-	1	-	1	2	-	2	-	3	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	2	2	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	2	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Illinois												
All races, both sexes	2	-	2	-	4	6	7	6	5	1	1	-
Male	1	-	1	-	2	-	5	2	2	-	-	-
Female	1	-	1	-	2	6	2	4	3	1	1	-
White, both sexes	2	-	2	-	3	6	7	6	5	1	1	-
Male	1	-	1	-	2	-	5	2	2	-	-	-
Female	1	-	1	-	1	6	2	4	3	1	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	1	1	-	2	2	5	3	2	-	-
Male	-	-	-	1	-	2	1	3	2	2	-	-
Female	-	-	1	-	-	-	1	2	1	-	-	-
White, both sexes	-	-	1	1	-	2	2	5	3	2	-	-
Male	-	-	-	1	-	2	1	3	2	2	-	-
Female	-	-	1	-	-	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Iowa												
All races, both sexes	-	-	-	2	-	4	3	2	4	1	-	-
Male	-	-	-	-	-	2	-	2	2	-	-	-
Female	-	-	-	2	-	2	3	-	2	1	-	-
White, both sexes	-	-	-	2	-	4	3	2	4	1	-	-
Male	-	-	-	-	-	2	-	2	2	-	-	-
Female	-	-	-	2	-	2	3	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	1	1	1	2	7	3	1	-	-
Male	-	-	-	1	-	-	-	2	-	1	-	-
Female	1	-	-	-	1	1	2	5	3	-	-	-
White, both sexes	1	-	-	1	1	1	2	7	3	1	-	-
Male	-	-	-	1	-	-	-	2	-	1	-	-
Female	1	-	-	-	1	1	2	5	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Kentucky												
All races, both sexes	-	1	1	-	1	5	2	2	1	1	-	-
Male	-	1	-	-	-	3	1	2	-	1	-	-
Female	-	-	1	-	1	2	1	-	1	-	-	-
White, both sexes	-	1	1	-	1	5	2	2	1	1	-	-
Male	-	1	-	-	-	3	1	2	-	1	-	-
Female	-	-	1	-	1	2	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Maine												
All races, both sexes	-	-	-	1	-	5	2	3	2	3	-	-
Male	-	-	-	-	-	3	-	2	1	2	-	-
Female	-	-	-	1	-	2	2	1	1	1	-	-
White, both sexes	-	-	-	1	-	5	2	3	2	3	-	-
Male	-	-	-	-	-	3	-	2	1	2	-	-
Female	-	-	-	1	-	2	2	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	2	1	-	4	-	1	1	-	-
Male	-	-	-	-	1	-	2	-	1	-	-	-
Female	1	-	-	2	-	-	2	-	-	1	-	-
White, both sexes	1	-	-	1	1	-	3	-	1	1	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	1	-	-	1	-	-	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Massachusetts												
All races, both sexes	-	1	1	1	1	5	5	3	2	2	-	-
Male	-	-	-	1	-	2	2	1	-	-	-	-
Female	-	1	1	-	1	3	3	2	2	2	-	-
White, both sexes	-	1	1	1	1	5	5	3	2	2	-	-
Male	-	-	-	1	-	2	2	1	-	-	-	-
Female	-	1	1	-	1	3	3	2	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	2	1	3	4	1	3	4	1	2	-
Male	-	-	2	1	2	1	-	2	1	-	-	-
Female	-	-	-	-	1	3	1	1	3	1	2	-
White, both sexes	-	-	2	1	3	4	1	3	4	1	2	-
Male	-	-	2	1	2	1	-	2	1	-	-	-
Female	-	-	-	-	1	3	1	1	3	1	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Minnesota												
All races, both sexes	-	2	-	-	2	2	4	7	8	4	1	-
Male	-	1	-	-	2	1	3	5	3	1	1	-
Female	-	1	-	-	-	1	1	2	5	3	-	-
White, both sexes	-	2	-	-	2	2	4	7	8	4	1	-
Male	-	1	-	-	2	1	3	5	3	1	1	-
Female	-	1	-	-	-	1	1	2	5	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	1	1	1	1	-	1	-	-	-
Male	1	-	-	1	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	1	-	-	1	1	1	1	-	1	-	-	-
Male	1	-	-	1	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Missouri												
All races, both sexes	2	1	-	-	-	4	6	5	4	2	1	-
Male	2	-	-	-	-	4	5	2	-	-	-	-
Female	-	1	-	-	-	-	1	3	4	2	1	-
White, both sexes	2	1	-	-	-	4	6	5	4	2	1	-
Male	2	-	-	-	-	4	5	2	-	-	-	-
Female	-	1	-	-	-	-	1	3	4	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	-	-	-	1	2	2	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	-	-
White, both sexes	-	-	1	-	-	-	1	2	2	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Nebraska												
All races, both sexes	-	-	-	-	-	1	3	4	4	1	-	-
Male	-	-	-	-	-	-	2	2	3	1	-	-
Female	-	-	-	-	-	1	1	2	1	-	-	-
White, both sexes	-	-	-	-	-	1	3	4	4	1	-	-
Male	-	-	-	-	-	-	2	2	3	1	-	-
Female	-	-	-	-	-	1	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
New Hampshire												
All races, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
New Mexico												
All races, both sexes	1	-	-	-	-	1	2	2	2	-	-	-
Male	1	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	1	2	1	-	-	-
White, both sexes	1	-	-	-	-	1	2	2	2	-	-	-
Male	1	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	1	-	-	1	3	11	5	8	2	2	-
Male	-	1	-	-	1	1	4	-	2	1	-	-
Female	2	-	-	-	-	2	7	5	6	1	2	-
White, both sexes	2	1	-	-	1	3	10	5	8	2	2	-
Male	-	1	-	-	1	1	3	-	2	1	-	-
Female	2	-	-	-	-	2	7	5	6	1	2	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
North Carolina												
All races, both sexes	-	1	1	2	1	-	3	3	5	-	1	-
Male	-	-	-	1	1	-	1	1	1	-	-	-
Female	-	1	1	1	-	-	2	2	4	-	1	-
White, both sexes	-	1	1	2	1	-	3	3	5	-	1	-
Male	-	-	-	1	1	-	1	1	1	-	-	-
Female	-	1	1	1	-	-	2	2	4	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	2	-	1	-	1	-	3	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	1	-	1	-	2	-	-	-
White, both sexes	-	-	2	-	1	-	1	-	3	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	1	-	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Ohio												
All races, both sexes	1	-	-	3	2	4	5	7	11	2	1	-
Male	1	-	-	1	1	1	2	-	5	-	-	-
Female	-	-	-	2	1	3	3	7	6	2	1	-
White, both sexes	1	-	-	3	2	3	5	7	10	2	1	-
Male	1	-	-	1	1	-	2	-	5	-	-	-
Female	-	-	-	2	1	3	3	7	5	2	1	-
Black, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	1	2	1	1	4	-	1	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	4	-	1	-	-
White, both sexes	-	-	-	1	2	1	1	3	-	1	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Oregon												
All races, both sexes	-	-	-	-	-	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	1	2	4	7	4	1	-	-
Male	-	-	-	-	1	-	2	3	1	-	-	-
Female	-	-	-	-	-	2	2	4	3	1	-	-
White, both sexes	-	-	-	-	1	2	4	7	4	1	-	-
Male	-	-	-	-	1	-	2	3	1	-	-	-
Female	-	-	-	-	-	2	2	4	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	1	1	1	-	2	3	4	-	1	-
Male	-	-	-	1	-	-	2	2	3	-	-	-
Female	-	-	1	-	1	-	-	1	1	-	1	-
White, both sexes	-	-	1	-	1	-	2	3	1	-	1	-
Male	-	-	-	-	-	-	2	2	-	-	-	-
Female	-	-	1	-	1	-	-	1	1	-	1	-
Black, both sexes	-	-	-	1	-	-	-	-	2	-	-	-
Male	-	-	-	1	-	-	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
South Dakota												
All races, both sexes	-	-	1	-	1	2	3	1	-	1	1	-
Male	-	-	-	-	1	1	3	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	1	1	-
White, both sexes	-	-	1	-	1	2	3	1	-	1	1	-
Male	-	-	-	-	1	1	3	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	1	-	1	2	2	4	1	1	2	-	-
Male	-	-	-	1	-	1	2	-	1	1	-	-
Female	-	1	-	-	2	1	2	1	-	1	-	-
White, both sexes	-	1	-	1	2	2	4	1	1	2	-	-
Male	-	-	-	1	-	1	2	-	1	1	-	-
Female	-	1	-	-	2	1	2	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Texas												
All races, both sexes	-	3	2	1	3	9	6	12	9	4	-	-
Male	-	2	1	-	-	2	2	1	3	2	-	-
Female	-	1	1	1	3	7	4	11	6	2	-	-
White, both sexes	-	3	1	-	3	8	5	12	8	4	-	-
Male	-	2	1	-	-	2	2	1	3	2	-	-
Female	-	1	-	-	3	6	3	11	5	2	-	-
Black, both sexes	-	-	1	1	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	1	-	1	-	1	3	3	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	1	-	1	-	1	3	2	-	-	-	-
White, both sexes	-	1	-	1	-	1	3	3	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	1	-	1	-	1	3	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Vermont												
All races, both sexes	-	-	1	-	1	2	2	2	1	1	-	-
Male	-	-	1	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	1	1	-	-
White, both sexes	-	-	1	-	1	2	2	2	1	1	-	-
Male	-	-	1	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	-	-	3	4	9	5	4	4	1	-
Male	-	-	-	-	2	1	3	3	1	-	1	-
Female	1	1	-	-	1	3	6	2	3	4	-	-
White, both sexes	1	1	-	-	3	4	8	4	4	4	1	-
Male	-	-	-	-	2	1	2	3	1	-	1	-
Female	1	1	-	-	1	3	6	1	3	4	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Washington												
All races, both sexes	-	1	1	-	2	2	2	5	6	1	1	-
Male	-	1	1	-	2	1	-	-	3	-	-	-
Female	-	-	-	-	-	1	2	5	3	1	1	-
White, both sexes	-	1	1	-	2	1	2	5	5	1	1	-
Male	-	1	1	-	2	-	-	-	3	-	-	-
Female	-	-	-	-	-	1	2	5	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
West Virginia												
All races, both sexes	-	1	-	-	-	1	3	3	-	2	-	-
Male	-	1	-	-	-	-	1	1	-	1	-	-
Female	-	-	-	-	-	1	2	2	-	1	-	-
White, both sexes	-	1	-	-	-	1	2	3	-	2	-	-
Male	-	1	-	-	-	-	1	1	-	1	-	-
Female	-	-	-	-	-	1	1	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J09-J11)												
Wisconsin												
All races, both sexes	1	-	1	-	2	7	2	10	3	6	1	-
Male	1	-	1	-	2	5	2	6	-	2	-	-
Female	-	-	-	-	-	2	-	4	3	4	1	-
White, both sexes	-	-	1	-	2	7	2	10	3	6	1	-
Male	-	-	1	-	2	5	2	6	-	2	-	-
Female	-	-	-	-	-	2	-	4	3	4	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
United States												
All races, both sexes	55,477	245	106	24	19	67	104	138	191	288	543	824
Male	25,288	152	51	12	12	34	62	81	90	155	303	468
Female	30,189	93	55	12	7	33	42	57	101	133	240	356
White, both sexes	48,613	148	72	19	14	55	74	102	132	212	408	601
Male	21,978	94	35	12	10	27	46	58	67	113	233	343
Female	26,635	54	37	7	4	28	28	44	65	99	175	258
Black, both sexes	5,270	85	30	3	5	10	28	31	44	66	119	200
Male	2,465	53	13	-	2	5	16	21	15	37	61	109
Female	2,805	32	17	3	3	5	12	10	29	29	58	91
American Indian 1/, both sexes	264	3	4	1	-	1	2	2	5	5	10	9
Male	136	-	3	-	-	1	-	-	2	3	6	8
Female	128	3	1	1	-	-	2	2	3	2	4	1
Asian or Pacific Islander 2/, both sexes	1,330	9	-	1	-	1	-	3	10	5	6	14
Male	709	5	-	-	-	1	-	2	6	2	3	8
Female	621	4	-	1	-	-	-	1	4	3	3	6
Alabama												
All races, both sexes	903	1	-	-	1	1	1	5	4	5	9	22
Male	417	-	-	-	1	-	-	3	-	5	7	15
Female	486	1	-	-	-	1	1	2	4	-	2	7
White, both sexes	738	1	-	-	1	1	-	3	3	4	5	15
Male	340	-	-	-	1	-	-	1	-	4	3	9
Female	398	1	-	-	-	1	-	2	3	-	2	6
Black, both sexes	162	-	-	-	-	-	1	1	1	1	4	7
Male	75	-	-	-	-	-	-	1	-	1	4	6
Female	87	-	-	-	-	-	1	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Alaska												
All races, both sexes	48	2	1	-	-	-	-	-	2	1	-	2
Male	28	-	-	-	-	-	-	-	1	-	-	1
Female	20	2	1	-	-	-	-	-	1	1	-	1
White, both sexes	30	-	-	-	-	-	-	-	-	1	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	17	2	1	-	-	-	-	-	2	-	-	1
Male	10	-	-	-	-	-	-	-	1	-	-	1
Female	7	2	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1,148	12	5	1	1	3	3	2	8	10	16	22
Male	578	8	2	-	-	2	1	2	1	5	7	11
Female	570	4	3	1	1	1	2	-	7	5	9	11
White, both sexes	1,047	9	3	-	-	2	2	2	7	6	12	17
Male	521	5	1	-	-	1	1	2	1	2	5	7
Female	526	4	2	-	-	1	1	-	6	4	7	10
Black, both sexes	30	3	1	-	1	-	1	-	-	2	1	2
Male	18	3	-	-	-	-	-	-	-	1	1	1
Female	12	-	1	-	1	-	1	-	-	1	-	1
American Indian 1/, both sexes	58	-	1	1	-	1	-	-	1	2	2	3
Male	33	-	1	-	-	1	-	-	-	2	-	3
Female	25	-	-	1	-	-	-	-	1	-	2	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Arkansas												
All races, both sexes	761	2	1	-	1	3	3	-	3	3	10	14
Male	335	1	1	-	-	2	1	-	-	2	5	9
Female	426	1	-	-	1	1	2	-	3	1	5	5
White, both sexes	683	1	1	-	-	3	1	-	3	3	8	9
Male	297	-	1	-	-	2	1	-	-	2	4	6
Female	386	1	-	-	-	1	-	-	3	1	4	3
Black, both sexes	75	1	-	-	1	-	2	-	-	-	2	5
Male	36	1	-	-	-	-	-	-	-	-	1	3
Female	39	-	-	-	1	-	2	-	-	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7,292	24	15	2	1	7	10	13	15	27	58	79
Male	3,438	13	9	1	1	4	4	9	8	14	36	50
Female	3,854	11	6	1	-	3	6	4	7	13	22	29
White, both sexes	6,110	18	14	2	1	5	8	10	9	24	49	63
Male	2,840	11	8	1	1	3	3	8	7	13	31	37
Female	3,270	7	6	1	-	2	5	2	2	11	18	26
Black, both sexes	474	3	1	-	-	1	2	2	4	2	7	10
Male	219	2	1	-	-	-	1	1	1	1	4	8
Female	255	1	-	-	-	1	1	1	3	1	3	2
American Indian 1/, both sexes	26	-	-	-	-	-	-	1	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	682	3	-	-	-	1	-	-	2	1	1	6
Male	366	-	-	-	-	1	-	-	-	-	-	5
Female	316	3	-	-	-	-	-	-	2	1	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Colorado												
All races, both sexes	595	3	3	2	-	1	4	4	5	5	6	14
Male	295	2	1	1	-	1	2	3	1	1	4	8
Female	300	1	2	1	-	-	2	1	4	4	2	6
White, both sexes	569	3	2	1	-	1	4	4	4	5	5	12
Male	283	2	-	1	-	1	2	3	1	1	3	7
Female	286	1	2	-	-	-	2	1	3	4	2	5
Black, both sexes	16	-	-	1	-	-	-	-	1	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	1	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	4	-	1	-	-	-	-	-	-	-	1	-
Male	4	-	1	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	781	3	-	-	-	-	-	1	2	1	3	6
Male	327	2	-	-	-	-	-	-	1	1	2	2
Female	454	1	-	-	-	-	-	1	1	-	1	4
White, both sexes	737	1	-	-	-	-	-	1	-	-	1	5
Male	308	-	-	-	-	-	-	-	-	-	1	2
Female	429	1	-	-	-	-	-	1	-	-	-	3
Black, both sexes	38	2	-	-	-	-	-	-	1	1	2	1
Male	15	2	-	-	-	-	-	-	-	1	1	-
Female	23	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Delaware												
All races, both sexes	134	-	-	-	-	-	1	-	2	1	2	2
Male	61	-	-	-	-	-	1	-	2	-	2	2
Female	73	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	117	-	-	-	-	-	-	-	1	-	2	1
Male	52	-	-	-	-	-	-	-	1	-	2	1
Female	65	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	16	-	-	-	-	-	1	-	1	1	-	1
Male	9	-	-	-	-	-	1	-	1	-	-	1
Female	7	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	76	1	-	-	-	-	-	-	-	1	1	3
Male	37	1	-	-	-	-	-	-	-	1	-	2
Female	39	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	54	1	-	-	-	-	-	-	-	1	1	3
Male	24	1	-	-	-	-	-	-	-	1	-	2
Female	30	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Florida												
All races, both sexes	2,426	20	4	2	-	1	9	8	7	11	24	43
Male	1,232	13	2	1	-	1	5	3	3	7	14	22
Female	1,194	7	2	1	-	-	4	5	4	4	10	21
White, both sexes	2,169	11	3	2	-	1	4	4	4	10	20	28
Male	1,113	8	2	1	-	1	3	1	3	7	12	15
Female	1,056	3	1	1	-	-	1	3	1	3	8	13
Black, both sexes	244	9	1	-	-	-	5	3	3	1	4	14
Male	115	5	-	-	-	-	2	2	-	-	2	7
Female	129	4	1	-	-	-	3	1	3	1	2	7
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	1	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	1
Georgia												
All races, both sexes	1,447	9	6	-	1	2	7	4	10	12	28	24
Male	636	6	2	-	1	-	5	3	4	10	14	17
Female	811	3	4	-	-	2	2	1	6	2	14	7
White, both sexes	1,141	2	2	-	1	1	4	2	6	6	15	12
Male	496	1	-	-	1	-	3	1	3	6	9	9
Female	645	1	2	-	-	1	1	1	3	-	6	3
Black, both sexes	299	7	3	-	-	1	3	2	4	6	13	12
Male	134	5	1	-	-	-	2	2	1	4	5	8
Female	165	2	2	-	-	1	1	-	3	2	8	4
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Hawaii												
All races, both sexes	255	1	1	-	-	-	-	1	3	1	1	5
Male	137	-	-	-	-	-	-	1	3	-	-	3
Female	118	1	1	-	-	-	-	-	-	1	1	2
White, both sexes	64	-	1	-	-	-	-	-	-	-	-	3
Male	33	-	-	-	-	-	-	-	-	-	-	3
Female	31	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	191	1	-	-	-	-	-	1	3	1	1	2
Male	104	-	-	-	-	-	-	1	3	-	-	-
Female	87	1	-	-	-	-	-	-	-	1	1	2
Idaho												
All races, both sexes	219	-	1	-	-	1	-	2	-	1	1	9
Male	114	-	1	-	-	1	-	2	-	1	-	5
Female	105	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	217	-	1	-	-	1	-	2	-	1	1	8
Male	113	-	1	-	-	1	-	2	-	1	-	4
Female	104	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Illinois												
All races, both sexes	2,629	14	3	3	1	4	2	4	14	17	22	37
Male	1,265	8	2	2	1	4	2	2	7	7	15	28
Female	1,364	6	1	1	-	-	-	2	7	10	7	9
White, both sexes	2,279	7	1	2	1	3	1	2	8	11	15	25
Male	1,081	5	1	2	1	3	1	1	4	4	11	19
Female	1,198	2	-	-	-	-	-	1	4	7	4	6
Black, both sexes	323	6	2	1	-	1	1	2	6	6	7	12
Male	167	2	1	-	-	1	1	1	3	3	4	9
Female	156	4	1	1	-	-	-	1	3	3	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	1	-	-	-	-	-	-	-	-	-	-
Male	17	1	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,112	10	4	-	-	3	3	3	5	8	10	21
Male	549	4	2	-	-	1	3	1	4	7	5	15
Female	563	6	2	-	-	2	-	2	1	1	5	6
White, both sexes	1,041	7	3	-	-	3	2	3	4	5	10	16
Male	509	4	1	-	-	1	2	1	3	5	5	11
Female	532	3	2	-	-	2	-	2	1	-	5	5
Black, both sexes	68	3	1	-	-	-	1	-	1	3	-	5
Male	38	-	1	-	-	-	1	-	1	2	-	4
Female	30	3	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Iowa												
All races, both sexes	754	3	-	-	-	-	2	-	5	2	5	8
Male	318	2	-	-	-	-	2	-	3	2	3	6
Female	436	1	-	-	-	-	-	-	2	-	2	2
White, both sexes	747	3	-	-	-	-	2	-	5	2	5	7
Male	312	2	-	-	-	-	2	-	3	2	3	5
Female	435	1	-	-	-	-	-	-	2	-	2	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	628	2	2	-	-	-	-	2	3	1	4	8
Male	259	1	1	-	-	-	-	2	1	1	-	2
Female	369	1	1	-	-	-	-	-	2	-	4	6
White, both sexes	599	1	-	-	-	-	-	1	3	1	3	8
Male	243	-	-	-	-	-	-	1	1	1	-	2
Female	356	1	-	-	-	-	-	-	2	-	3	6
Black, both sexes	23	1	2	-	-	-	-	1	-	-	1	-
Male	11	1	1	-	-	-	-	1	-	-	-	-
Female	12	-	1	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Kentucky												
All races, both sexes	909	4	-	-	-	-	6	1	5	6	8	13
Male	420	2	-	-	-	-	5	1	4	6	3	7
Female	489	2	-	-	-	-	1	-	1	-	5	6
White, both sexes	855	3	-	-	-	-	6	1	5	6	7	12
Male	394	2	-	-	-	-	5	1	4	6	2	6
Female	461	1	-	-	-	-	1	-	1	-	5	6
Black, both sexes	53	1	-	-	-	-	-	-	-	-	1	1
Male	26	-	-	-	-	-	-	-	-	-	1	1
Female	27	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	827	10	3	1	1	1	3	2	7	3	15	25
Male	385	2	2	1	1	-	2	1	1	3	6	11
Female	442	8	1	-	-	1	1	1	6	-	9	14
White, both sexes	594	2	1	1	1	1	2	1	3	2	11	17
Male	272	-	1	1	1	-	2	1	-	2	4	9
Female	322	2	-	-	-	1	-	-	3	-	7	8
Black, both sexes	232	8	2	-	-	-	1	1	4	1	4	8
Male	112	2	1	-	-	-	-	-	1	1	2	2
Female	120	6	1	-	-	-	1	1	3	-	2	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Maine												
All races, both sexes	253	1	-	-	-	-	1	-	2	1	2	3
Male	101	-	-	-	-	-	-	-	2	1	1	3
Female	152	1	-	-	-	-	1	-	-	-	1	-
White, both sexes	248	1	-	-	-	-	1	-	1	1	2	3
Male	98	-	-	-	-	-	-	-	1	1	1	3
Female	150	1	-	-	-	-	1	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,067	5	-	1	-	-	3	1	1	5	19	29
Male	481	2	-	-	-	-	3	1	1	4	10	12
Female	586	3	-	1	-	-	-	-	-	1	9	17
White, both sexes	829	4	-	-	-	-	2	-	1	2	9	12
Male	374	1	-	-	-	-	2	-	1	1	3	5
Female	455	3	-	-	-	-	-	-	-	1	6	7
Black, both sexes	211	1	-	1	-	-	1	1	-	3	8	16
Male	94	1	-	-	-	-	1	1	-	3	6	7
Female	117	-	-	1	-	-	-	-	-	-	2	9
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	2	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Massachusetts												
All races, both sexes	1,728	2	2	1	-	1	-	-	2	4	7	16
Male	734	2	-	-	-	-	-	-	1	2	6	11
Female	994	-	2	1	-	1	-	-	1	2	1	5
White, both sexes	1,669	1	2	1	-	1	-	-	1	4	7	14
Male	704	1	-	-	-	-	-	-	-	2	6	9
Female	965	-	2	1	-	1	-	-	1	2	1	5
Black, both sexes	44	1	-	-	-	-	-	-	-	-	-	2
Male	24	1	-	-	-	-	-	-	-	-	-	2
Female	20	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,656	6	3	1	-	1	-	4	7	9	18	16
Male	772	3	3	-	-	1	-	3	2	2	12	9
Female	884	3	-	1	-	-	-	1	5	7	6	7
White, both sexes	1,447	4	2	1	-	-	-	2	6	8	12	10
Male	663	1	2	-	-	-	-	1	2	2	10	6
Female	784	3	-	1	-	-	-	1	4	6	2	4
Black, both sexes	197	2	1	-	-	1	-	2	-	1	6	6
Male	105	2	1	-	-	1	-	2	-	-	2	3
Female	92	-	-	-	-	-	-	-	-	1	4	3
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Minnesota												
All races, both sexes	600	3	2	-	-	-	1	-	-	2	4	7
Male	247	3	2	-	-	-	1	-	-	1	1	5
Female	353	-	-	-	-	-	-	-	-	1	3	2
White, both sexes	589	1	2	-	-	-	1	-	-	2	4	6
Male	241	1	2	-	-	-	1	-	-	1	1	5
Female	348	-	-	-	-	-	-	-	-	1	3	1
Black, both sexes	5	1	-	-	-	-	-	-	-	-	-	1
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	615	4	1	1	-	2	1	5	2	4	4	12
Male	262	4	1	1	-	-	-	2	2	-	2	6
Female	353	-	-	-	-	2	1	3	-	4	2	6
White, both sexes	414	2	1	1	-	2	-	1	2	2	2	5
Male	170	2	1	1	-	-	-	-	2	-	1	1
Female	244	-	-	-	-	2	-	1	-	2	1	4
Black, both sexes	200	2	-	-	-	-	1	4	-	2	2	7
Male	91	2	-	-	-	-	-	2	-	-	1	5
Female	109	-	-	-	-	-	1	2	-	2	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Missouri												
All races, both sexes	1,257	6	1	-	1	2	1	5	6	2	12	19
Male	535	4	-	-	1	1	1	2	2	1	4	13
Female	722	2	1	-	-	1	-	3	4	1	8	6
White, both sexes	1,165	2	-	-	1	2	1	5	3	1	9	14
Male	498	2	-	-	1	1	1	2	1	-	3	10
Female	667	-	-	-	-	1	-	3	2	1	6	4
Black, both sexes	87	4	1	-	-	-	-	-	3	1	3	5
Male	37	2	-	-	-	-	-	-	1	1	1	3
Female	50	2	1	-	-	-	-	-	2	-	2	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	163	-	-	-	-	-	1	1	-	1	1	3
Male	77	-	-	-	-	-	1	-	-	1	1	3
Female	86	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	151	-	-	-	-	-	1	-	-	-	1	3
Male	70	-	-	-	-	-	1	-	-	-	1	3
Female	81	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	1	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Nebraska												
All races, both sexes	331	1	-	-	-	2	1	2	1	3	5	3
Male	139	-	-	-	-	-	-	1	-	-	2	1
Female	192	1	-	-	-	2	1	1	1	3	3	2
White, both sexes	325	1	-	-	-	2	1	2	1	3	5	3
Male	137	-	-	-	-	-	-	1	-	-	2	1
Female	188	1	-	-	-	2	1	1	1	3	3	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	426	3	4	-	-	-	2	3	3	5	3	11
Male	223	1	3	-	-	-	1	3	1	2	3	6
Female	203	2	1	-	-	-	1	-	2	3	-	5
White, both sexes	376	2	4	-	-	-	2	1	3	3	2	9
Male	195	1	3	-	-	-	1	1	1	2	2	5
Female	181	1	1	-	-	-	1	-	2	1	-	4
Black, both sexes	33	1	-	-	-	-	-	2	-	2	1	-
Male	22	-	-	-	-	-	-	2	-	-	1	-
Female	11	1	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
New Hampshire												
All races, both sexes	210	-	1	-	-	-	1	-	-	-	1	1
Male	109	-	-	-	-	-	-	-	-	-	-	-
Female	101	-	1	-	-	-	1	-	-	-	1	1
White, both sexes	205	-	1	-	-	-	1	-	-	-	1	1
Male	106	-	-	-	-	-	-	-	-	-	-	-
Female	99	-	1	-	-	-	1	-	-	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,312	2	1	-	2	1	2	7	6	9	9	14
Male	603	1	-	-	2	-	1	5	5	3	7	9
Female	709	1	1	-	-	1	1	2	1	6	2	5
White, both sexes	1,175	2	-	-	-	1	2	5	6	6	6	10
Male	538	1	-	-	-	-	1	4	5	-	5	8
Female	637	1	-	-	-	1	1	1	1	6	1	2
Black, both sexes	119	-	1	-	2	-	-	2	-	3	3	4
Male	50	-	-	-	2	-	-	1	-	3	2	1
Female	69	-	1	-	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
New Mexico												
All races, both sexes	342	2	-	-	-	1	1	1	2	2	5	15
Male	166	1	-	-	-	1	-	-	2	2	4	9
Female	176	1	-	-	-	-	1	1	-	-	1	6
White, both sexes	316	2	-	-	-	1	1	1	1	2	4	12
Male	153	1	-	-	-	1	-	-	1	2	3	7
Female	163	1	-	-	-	-	1	1	-	-	1	5
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	1	-	1	2
Male	12	-	-	-	-	-	-	-	1	-	1	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4,851	12	10	-	1	3	4	7	7	18	42	51
Male	2,159	8	5	-	-	2	1	3	4	11	25	27
Female	2,692	4	5	-	1	1	3	4	3	7	17	24
White, both sexes	3,995	7	5	-	1	2	4	6	4	12	29	35
Male	1,771	5	3	-	-	1	1	2	3	10	19	20
Female	2,224	2	2	-	1	1	3	4	1	2	10	15
Black, both sexes	694	5	5	-	-	1	-	1	2	5	12	14
Male	299	3	2	-	-	1	-	1	1	-	5	6
Female	395	2	3	-	-	-	-	-	1	5	7	8
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	159	-	-	-	-	-	-	-	1	1	1	2
Male	89	-	-	-	-	-	-	-	-	1	1	1
Female	70	-	-	-	-	-	-	-	1	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
North Carolina												
All races, both sexes	1,684	10	5	2	-	5	6	3	6	13	28	36
Male	702	7	3	1	-	3	3	2	3	6	15	19
Female	982	3	2	1	-	2	3	1	3	7	13	17
White, both sexes	1,418	5	4	2	-	4	2	1	5	7	18	22
Male	583	4	2	1	-	2	1	1	3	2	10	12
Female	835	1	2	1	-	2	1	-	2	5	8	10
Black, both sexes	256	5	1	-	-	1	3	2	1	6	10	13
Male	114	3	1	-	-	1	2	1	-	4	5	6
Female	142	2	-	-	-	-	1	1	1	2	5	7
American Indian 1/, both sexes	7	-	-	-	-	-	1	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	131	-	-	-	-	-	-	-	-	-	-	1
Male	58	-	-	-	-	-	-	-	-	-	-	-
Female	73	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	129	-	-	-	-	-	-	-	-	-	-	1
Male	56	-	-	-	-	-	-	-	-	-	-	-
Female	73	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Ohio												
All races, both sexes	1,910	11	6	1	1	1	2	6	3	14	20	28
Male	833	10	3	1	1	-	1	5	2	8	13	13
Female	1,077	1	3	-	-	1	1	1	1	6	7	15
White, both sexes	1,739	5	4	1	1	1	2	4	2	9	17	24
Male	749	5	1	1	1	-	1	3	2	5	11	11
Female	990	-	3	-	-	1	1	1	-	4	6	13
Black, both sexes	168	6	2	-	-	-	-	2	1	5	3	4
Male	82	5	2	-	-	-	-	2	-	3	2	2
Female	86	1	-	-	-	-	-	-	1	2	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	869	3	2	1	-	2	2	3	-	12	12	13
Male	392	1	1	-	-	-	1	-	-	4	6	8
Female	477	2	1	1	-	2	1	3	-	8	6	5
White, both sexes	789	2	2	1	-	2	1	3	-	10	9	10
Male	358	1	1	-	-	-	1	-	-	4	4	6
Female	431	1	1	1	-	2	-	3	-	6	5	4
Black, both sexes	44	-	-	-	-	-	-	-	-	-	-	3
Male	17	-	-	-	-	-	-	-	-	-	-	2
Female	27	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	36	1	-	-	-	-	1	-	-	2	3	-
Male	17	-	-	-	-	-	-	-	-	-	2	-
Female	19	1	-	-	-	-	1	-	-	2	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Oregon												
All races, both sexes	517	2	-	-	-	-	1	1	3	1	6	5
Male	255	1	-	-	-	-	1	1	1	1	3	4
Female	262	1	-	-	-	-	-	-	2	-	3	1
White, both sexes	510	1	-	-	-	-	1	1	3	1	6	5
Male	253	-	-	-	-	-	1	1	1	1	3	4
Female	257	1	-	-	-	-	-	-	2	-	3	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2,695	8	2	-	2	1	5	5	4	10	17	35
Male	1,191	5	-	-	1	1	3	2	1	3	11	20
Female	1,504	3	2	-	1	-	2	3	3	7	6	15
White, both sexes	2,489	6	2	-	2	1	4	5	3	9	13	24
Male	1,092	3	-	-	1	1	2	2	1	2	11	14
Female	1,397	3	2	-	1	-	2	3	2	7	2	10
Black, both sexes	196	2	-	-	-	-	1	-	1	1	4	11
Male	96	2	-	-	-	-	1	-	-	1	-	6
Female	100	-	-	-	-	-	-	-	1	-	4	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Rhode Island												
All races, both sexes	265	-	-	-	-	-	-	1	-	2	4	-
Male	109	-	-	-	-	-	-	1	-	1	2	-
Female	156	-	-	-	-	-	-	-	-	1	2	-
White, both sexes	254	-	-	-	-	-	-	1	-	1	4	-
Male	103	-	-	-	-	-	-	1	-	1	2	-
Female	151	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	709	1	-	-	-	-	2	2	6	6	6	13
Male	331	-	-	-	-	-	2	1	3	3	3	7
Female	378	1	-	-	-	-	-	1	3	3	3	6
White, both sexes	542	-	-	-	-	-	1	2	2	6	3	10
Male	244	-	-	-	-	-	1	1	1	3	1	5
Female	298	-	-	-	-	-	-	1	1	3	2	5
Black, both sexes	165	1	-	-	-	-	1	-	4	-	3	3
Male	87	-	-	-	-	-	1	-	2	-	2	2
Female	78	1	-	-	-	-	-	-	2	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
South Dakota												
All races, both sexes	163	-	-	-	-	-	-	-	1	-	4	2
Male	68	-	-	-	-	-	-	-	-	-	3	-
Female	95	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	154	-	-	-	-	-	-	-	-	-	3	2
Male	62	-	-	-	-	-	-	-	-	-	2	-
Female	92	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	1	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,518	8	1	-	1	7	4	4	2	8	15	26
Male	645	6	1	-	1	4	2	3	1	5	8	13
Female	873	2	-	-	-	3	2	1	1	3	7	13
White, both sexes	1,345	5	-	-	1	5	4	4	1	6	10	18
Male	560	4	-	-	1	4	2	3	1	3	6	10
Female	785	1	-	-	-	1	2	1	-	3	4	8
Black, both sexes	170	3	1	-	-	2	-	-	1	2	5	8
Male	82	2	1	-	-	-	-	-	-	2	2	3
Female	88	1	-	-	-	2	-	-	1	-	3	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Texas												
All races, both sexes	3,283	14	8	3	2	4	5	15	13	20	40	63
Male	1,436	9	2	3	-	3	4	7	6	12	22	29
Female	1,847	5	6	-	2	1	1	8	7	8	18	34
White, both sexes	2,903	10	5	3	1	4	3	13	10	12	31	50
Male	1,269	6	2	3	-	3	2	7	4	7	17	24
Female	1,634	4	3	-	1	1	1	6	6	5	14	26
Black, both sexes	330	3	3	-	1	-	2	2	2	8	9	13
Male	146	2	-	-	-	-	2	-	1	5	5	5
Female	184	1	3	-	1	-	-	2	1	3	4	8
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	46	1	-	-	-	-	-	-	1	-	-	-
Male	20	1	-	-	-	-	-	-	1	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	332	2	2	1	-	2	-	2	-	3	2	5
Male	142	2	-	-	-	-	-	1	-	1	-	2
Female	190	-	2	1	-	2	-	1	-	2	2	3
White, both sexes	320	2	2	-	-	2	-	2	-	2	2	5
Male	137	2	-	-	-	-	-	1	-	-	-	2
Female	183	-	2	-	-	2	-	1	-	2	2	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	1	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Vermont												
All races, both sexes	67	-	-	-	-	-	-	-	-	-	-	-
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	67	-	-	-	-	-	-	-	-	-	-	-
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,247	5	1	-	-	2	3	3	4	5	13	14
Male	582	5	1	-	-	1	3	2	2	2	6	9
Female	665	-	-	-	-	1	-	1	2	3	7	5
White, both sexes	1,060	3	1	-	-	1	2	2	2	4	10	10
Male	497	3	1	-	-	1	2	1	1	2	4	7
Female	563	-	-	-	-	-	-	1	1	2	6	3
Black, both sexes	167	2	-	-	-	1	1	1	2	1	3	4
Male	74	2	-	-	-	-	1	1	1	-	2	2
Female	93	-	-	-	-	1	-	-	1	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Washington												
All races, both sexes	791	4	4	-	2	1	1	3	3	10	10	7
Male	364	4	1	-	1	1	-	1	2	5	5	2
Female	427	-	3	-	1	-	1	2	1	5	5	5
White, both sexes	737	3	3	-	2	-	1	3	3	9	9	7
Male	341	3	1	-	1	-	-	1	2	5	5	2
Female	396	-	2	-	1	-	1	2	1	4	4	5
Black, both sexes	15	-	1	-	-	1	-	-	-	-	-	-
Male	6	-	-	-	-	1	-	-	-	-	-	-
Female	9	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	26	1	-	-	-	-	-	-	-	1	-	-
Male	12	1	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	1	-	-
West Virginia												
All races, both sexes	448	3	1	-	-	1	-	2	2	2	4	7
Male	213	3	-	-	-	-	-	2	-	-	2	4
Female	235	-	1	-	-	1	-	-	2	2	2	3
White, both sexes	438	3	-	-	-	1	-	2	2	2	4	7
Male	208	3	-	-	-	-	-	2	-	-	2	4
Female	230	-	-	-	-	1	-	-	2	2	2	3
Black, both sexes	9	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Wisconsin												
All races, both sexes	980	6	-	-	-	1	-	-	4	1	7	8
Male	452	5	-	-	-	-	-	-	3	1	3	5
Female	528	1	-	-	-	1	-	-	1	-	4	3
White, both sexes	946	5	-	-	-	1	-	-	4	1	7	7
Male	436	4	-	-	-	-	-	-	3	1	3	5
Female	510	1	-	-	-	1	-	-	1	-	4	2
Black, both sexes	26	1	-	-	-	-	-	-	-	-	-	1
Male	13	1	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	113	-	-	1	-	-	-	-	1	-	-	4
Male	47	-	-	-	-	-	-	-	-	-	-	3
Female	66	-	-	1	-	-	-	-	1	-	-	1
White, both sexes	111	-	-	1	-	-	-	-	1	-	-	3
Male	45	-	-	-	-	-	-	-	-	-	-	2
Female	66	-	-	1	-	-	-	-	1	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
United States												
All races, both sexes	1,157	1,370	1,741	2,368	3,612	6,272	10,160	11,519	9,384	4,274	1,068	3
Male	705	820	1,037	1,344	1,985	3,333	4,878	5,002	3,368	1,176	218	2
Female	452	550	704	1,024	1,627	2,939	5,282	6,517	6,016	3,098	850	1
White, both sexes	868	1,087	1,429	1,897	3,089	5,480	9,045	10,486	8,568	3,879	945	3
Male	519	660	843	1,081	1,689	2,926	4,359	4,562	3,060	1,051	188	2
Female	349	427	586	816	1,400	2,554	4,686	5,924	5,508	2,828	757	1
Black, both sexes	254	249	260	397	406	591	824	716	577	277	98	-
Male	166	140	165	213	224	310	353	276	187	76	23	-
Female	88	109	95	184	182	281	471	440	390	201	75	-
American Indian 1/, both sexes	15	11	15	17	23	28	39	37	18	13	6	-
Male	6	5	6	12	15	10	26	22	7	4	-	-
Female	9	6	9	5	8	18	13	15	11	9	6	-
Asian or Pacific Islander 2/, both sexes	20	23	37	57	94	173	252	280	221	105	19	-
Male	14	15	23	38	57	87	140	142	114	45	7	-
Female	6	8	14	19	37	86	112	138	107	60	12	-
Alabama												
All races, both sexes	25	34	31	62	66	127	170	161	116	49	13	-
Male	22	19	19	39	33	65	81	64	33	9	2	-
Female	3	15	12	23	33	62	89	97	83	40	11	-
White, both sexes	14	27	25	50	55	106	143	140	93	42	10	-
Male	13	16	16	30	27	54	72	60	25	7	2	-
Female	1	11	9	20	28	52	71	80	68	35	8	-
Black, both sexes	10	7	6	12	11	21	26	21	23	7	3	-
Male	8	3	3	9	6	11	9	4	8	2	-	-
Female	2	4	3	3	5	10	17	17	15	5	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Alaska												
All races, both sexes	4	1	2	3	1	3	8	6	4	8	-	-
Male	2	1	1	2	1	3	5	4	2	5	-	-
Female	2	-	1	1	-	-	3	2	2	3	-	-
White, both sexes	2	1	1	2	-	2	5	6	3	6	-	-
Male	1	1	1	2	-	2	2	4	1	4	-	-
Female	1	-	-	-	-	-	3	2	2	2	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	1	1	3	-	1	2	-	-
Male	1	-	-	-	1	1	3	-	1	1	-	-
Female	-	-	1	1	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	39	37	61	75	95	138	182	216	147	56	19	-
Male	25	21	38	50	53	70	100	108	52	20	2	-
Female	14	16	23	25	42	68	82	108	95	36	17	-
White, both sexes	36	31	53	70	87	126	170	204	144	49	17	-
Male	24	19	33	46	48	65	91	100	51	17	2	-
Female	12	12	20	24	39	61	79	104	93	32	15	-
Black, both sexes	2	2	4	2	2	3	2	-	1	1	-	-
Male	1	-	4	1	2	2	2	-	-	-	-	-
Female	1	2	-	1	-	1	-	-	1	1	-	-
American Indian 1/, both sexes	1	4	4	3	4	8	8	9	2	2	2	-
Male	-	2	1	3	2	3	6	6	1	2	-	-
Female	1	2	3	-	2	5	2	3	1	-	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	2	3	-	4	-	-
Male	-	-	-	-	1	-	1	2	-	1	-	-
Female	-	-	-	-	1	1	1	1	-	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Arkansas												
All races, both sexes	21	19	30	31	46	77	147	147	132	55	16	-
Male	9	13	17	13	21	40	72	68	39	20	2	-
Female	12	6	13	18	25	37	75	79	93	35	14	-
White, both sexes	17	16	27	28	43	71	134	138	121	48	11	-
Male	7	11	15	12	20	36	66	61	35	17	1	-
Female	10	5	12	16	23	35	68	77	86	31	10	-
Black, both sexes	4	3	3	3	3	5	12	9	10	7	5	-
Male	2	2	2	1	1	3	5	7	4	3	1	-
Female	2	1	1	2	2	2	7	2	6	4	4	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
California												
All races, both sexes	92	119	223	303	459	783	1,420	1,611	1,294	607	130	-
Male	60	68	132	180	243	429	686	771	512	179	29	-
Female	32	51	91	123	216	354	734	840	782	428	101	-
White, both sexes	75	100	186	234	363	644	1,186	1,353	1,124	523	119	-
Male	48	58	112	140	181	358	564	645	436	148	27	-
Female	27	42	74	94	182	286	622	708	688	375	92	-
Black, both sexes	10	10	22	37	50	49	83	94	52	30	5	-
Male	7	4	11	19	31	26	32	42	16	12	-	-
Female	3	6	11	18	19	23	51	52	36	18	5	-
American Indian 1/, both sexes	1	1	-	2	2	5	5	5	1	2	-	-
Male	-	1	-	1	2	-	4	3	1	-	-	-
Female	1	-	-	1	-	5	1	2	-	2	-	-
Asian or Pacific Islander 2/, both sexes	6	8	15	30	44	85	146	159	117	52	6	-
Male	5	5	9	20	29	45	86	81	59	19	2	-
Female	1	3	6	10	15	40	60	78	58	33	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Colorado												
All races, both sexes	16	20	14	17	36	60	109	127	88	50	11	-
Male	8	14	7	11	17	35	68	61	33	15	2	-
Female	8	6	7	6	19	25	41	66	55	35	9	-
White, both sexes	15	20	14	13	35	55	104	125	88	48	11	-
Male	8	14	7	9	17	33	65	59	33	15	2	-
Female	7	6	7	4	18	22	39	66	55	33	9	-
Black, both sexes	1	-	-	4	1	2	2	-	-	2	-	-
Male	-	-	-	2	-	1	1	-	-	-	-	-
Female	1	-	-	2	1	1	1	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	3	1	-	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
Connecticut												
All races, both sexes	6	10	24	23	30	84	150	192	146	82	18	-
Male	4	7	17	13	17	43	64	78	50	23	3	-
Female	2	3	7	10	13	41	86	114	96	59	15	-
White, both sexes	6	9	20	21	29	81	141	188	140	77	17	-
Male	4	6	14	12	16	42	61	76	49	22	3	-
Female	2	3	6	9	13	39	80	112	91	55	14	-
Black, both sexes	-	1	4	2	1	3	6	3	5	5	1	-
Male	-	1	3	1	1	1	2	1	-	1	-	-
Female	-	-	1	1	-	2	4	2	5	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	1	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Delaware												
All races, both sexes	3	3	1	7	12	21	27	25	19	7	1	-
Male	2	2	1	3	4	11	10	14	6	1	-	-
Female	1	1	-	4	8	10	17	11	13	6	1	-
White, both sexes	2	3	1	4	11	17	26	23	18	7	1	-
Male	1	2	1	3	4	9	9	12	6	1	-	-
Female	1	1	-	1	7	8	17	11	12	6	1	-
Black, both sexes	1	-	-	3	1	3	1	2	1	-	-	-
Male	1	-	-	-	-	2	1	2	-	-	-	-
Female	-	-	-	3	1	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	4	1	6	6	5	16	13	9	5	2	-
Male	2	3	1	3	4	2	4	8	5	-	1	-
Female	1	1	-	3	2	3	12	5	4	5	1	-
White, both sexes	-	-	1	1	2	2	5	6	-	3	1	-
Male	-	-	1	1	2	1	2	6	-	-	-	-
Female	-	-	-	-	-	1	3	-	-	3	1	-
Black, both sexes	3	4	-	5	4	3	11	7	9	1	1	-
Male	2	3	-	2	2	1	2	2	5	-	1	-
Female	1	1	-	3	2	2	9	5	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Florida												
All races, both sexes	56	59	88	148	169	295	454	487	360	145	35	1
Male	39	40	54	92	105	174	230	231	146	42	7	1
Female	17	19	34	56	64	121	224	256	214	103	28	-
White, both sexes	48	49	69	113	146	268	433	458	332	133	32	1
Male	33	34	43	74	92	161	219	222	136	38	7	1
Female	15	15	26	39	54	107	214	236	196	95	25	-
Black, both sexes	7	10	19	35	22	25	20	27	25	11	3	-
Male	6	6	11	18	13	12	11	8	8	4	-	-
Female	1	4	8	17	9	13	9	19	17	7	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	2	2	1	-	-
Male	-	-	-	-	-	1	-	1	2	-	-	-
Female	-	-	-	-	1	1	1	1	-	1	-	-
Georgia												
All races, both sexes	40	50	53	69	96	177	279	276	196	87	21	-
Male	23	29	34	45	47	88	131	104	57	14	2	-
Female	17	21	19	24	49	89	148	172	139	73	19	-
White, both sexes	25	36	32	49	76	140	239	242	165	72	14	-
Male	14	22	19	36	38	72	112	91	47	11	1	-
Female	11	14	13	13	38	68	127	151	118	61	13	-
Black, both sexes	14	14	19	20	20	37	39	33	30	15	7	-
Male	8	7	13	9	9	16	18	12	10	3	1	-
Female	6	7	6	11	11	21	21	21	20	12	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	-	-	-	1	1	1	-	-	-
Male	1	-	2	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Hawaii												
All races, both sexes	9	5	4	6	15	27	33	60	50	26	7	-
Male	5	2	1	4	7	14	20	37	23	14	3	-
Female	4	3	3	2	8	13	13	23	27	12	4	-
White, both sexes	3	1	1	-	6	6	12	11	14	5	1	-
Male	2	-	1	-	4	4	5	5	7	2	-	-
Female	1	1	-	-	2	2	7	6	7	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	4	3	6	9	21	21	49	36	21	6	-
Male	3	2	-	4	3	10	15	32	16	12	3	-
Female	3	2	3	2	6	11	6	17	20	9	3	-
Idaho												
All races, both sexes	6	4	6	5	11	23	39	45	35	19	11	-
Male	3	3	3	2	5	12	23	23	18	9	3	-
Female	3	1	3	3	6	11	16	22	17	10	8	-
White, both sexes	6	4	6	5	11	23	39	45	34	19	11	-
Male	3	3	3	2	5	12	23	23	18	9	3	-
Female	3	1	3	3	6	11	16	22	16	10	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Illinois												
All races, both sexes	45	62	88	114	167	283	506	530	452	202	59	-
Male	31	31	58	73	91	162	260	247	156	66	12	-
Female	14	31	30	41	76	121	246	283	296	136	47	-
White, both sexes	34	43	73	82	135	246	434	490	422	193	51	-
Male	27	24	45	52	76	145	218	229	141	65	7	-
Female	7	19	28	30	59	101	216	261	281	128	44	-
Black, both sexes	11	19	15	30	30	37	64	33	26	8	6	-
Male	4	7	13	19	14	17	36	15	12	1	4	-
Female	7	12	2	11	16	20	28	18	14	7	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	2	-	8	7	4	1	2	-
Male	-	-	-	2	1	-	6	3	3	-	1	-
Female	-	-	-	-	1	-	2	4	1	1	1	-
Indiana												
All races, both sexes	31	28	34	48	73	132	211	225	183	62	18	-
Male	22	20	20	28	41	69	98	116	68	20	5	-
Female	9	8	14	20	32	63	113	109	115	42	13	-
White, both sexes	27	25	33	46	69	125	195	215	178	58	17	-
Male	19	17	19	26	39	66	92	110	65	18	5	-
Female	8	8	14	20	30	59	103	105	113	40	12	-
Black, both sexes	3	3	1	2	3	7	16	9	5	4	1	-
Male	3	3	1	2	1	3	6	5	3	2	-	-
Female	-	-	-	-	2	4	10	4	2	2	1	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Iowa												
All races, both sexes	3	14	9	26	39	54	137	152	160	111	24	-
Male	3	9	7	15	24	29	75	59	45	31	3	-
Female	-	5	2	11	15	25	62	93	115	80	21	-
White, both sexes	3	14	8	25	38	54	135	152	159	111	24	-
Male	3	9	6	14	23	29	73	59	45	31	3	-
Female	-	5	2	11	15	25	62	93	114	80	21	-
Black, both sexes	-	-	1	1	1	-	2	-	-	-	-	-
Male	-	-	1	1	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Kansas												
All races, both sexes	12	13	9	16	25	58	108	134	134	78	19	-
Male	7	9	5	9	17	25	51	54	48	21	5	-
Female	5	4	4	7	8	33	57	80	86	57	14	-
White, both sexes	10	12	9	15	24	57	103	129	131	74	18	-
Male	6	9	5	8	16	24	48	52	46	20	4	-
Female	4	3	4	7	8	33	55	77	85	54	14	-
Black, both sexes	-	1	-	-	1	1	5	3	2	4	1	-
Male	-	-	-	-	1	1	3	-	1	1	1	-
Female	-	1	-	-	-	-	2	3	1	3	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	2	-	-	-	-
Male	1	-	-	-	-	-	-	2	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Kentucky												
All races, both sexes	22	28	30	47	73	111	168	164	142	66	15	-
Male	9	14	21	30	44	53	83	69	45	20	4	-
Female	13	14	9	17	29	58	85	95	97	46	11	-
White, both sexes	21	28	26	44	68	106	159	160	125	63	15	-
Male	9	14	18	28	40	50	78	67	40	20	4	-
Female	12	14	8	16	28	56	81	93	85	43	11	-
Black, both sexes	1	-	4	3	5	5	9	4	16	3	-	-
Male	-	-	3	2	4	3	5	2	5	-	-	-
Female	1	-	1	1	1	2	4	2	11	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Louisiana												
All races, both sexes	20	28	32	49	60	104	170	124	114	43	11	1
Male	14	16	17	24	37	50	81	55	45	12	3	1
Female	6	12	15	25	23	54	89	69	69	31	8	-
White, both sexes	12	18	20	36	40	67	137	96	89	30	6	1
Male	7	8	11	15	24	30	65	42	39	7	2	1
Female	5	10	9	21	16	37	72	54	50	23	4	-
Black, both sexes	8	10	12	13	20	37	32	28	25	13	5	-
Male	7	8	6	9	13	20	15	13	6	5	1	-
Female	1	2	6	4	7	17	17	15	19	8	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Maine												
All races, both sexes	3	4	4	5	24	20	52	38	62	26	5	-
Male	1	2	1	1	9	12	27	16	21	4	-	-
Female	2	2	3	4	15	8	25	22	41	22	5	-
White, both sexes	3	4	4	5	24	17	52	38	61	26	5	-
Male	1	2	1	1	9	10	27	16	21	4	-	-
Female	2	2	3	4	15	7	25	22	40	22	5	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Maryland												
All races, both sexes	36	35	28	38	69	107	181	227	189	73	20	-
Male	24	26	13	22	42	59	79	86	76	18	3	-
Female	12	9	15	16	27	48	102	141	113	55	17	-
White, both sexes	18	26	17	25	50	85	151	187	158	66	16	-
Male	12	19	8	16	31	49	69	72	66	18	1	-
Female	6	7	9	9	19	36	82	115	92	48	15	-
Black, both sexes	18	9	11	11	15	20	25	37	25	7	2	-
Male	12	7	5	4	8	8	8	13	9	-	1	-
Female	6	2	6	7	7	12	17	24	16	7	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	3	2	5	3	5	-	2	-
Male	-	-	-	2	3	2	2	1	1	-	1	-
Female	-	-	-	-	-	-	3	2	4	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Massachusetts												
All races, both sexes	20	28	49	48	103	183	333	396	326	162	45	-
Male	11	17	31	20	65	94	151	144	123	46	10	-
Female	9	11	18	28	38	89	182	252	203	116	35	-
White, both sexes	19	26	47	45	99	174	322	384	319	159	44	-
Male	10	15	30	20	61	90	146	141	120	44	9	-
Female	9	11	17	25	38	84	176	243	199	115	35	-
Black, both sexes	1	2	2	2	2	6	9	9	4	3	1	-
Male	1	2	1	-	2	3	5	3	1	2	1	-
Female	-	-	1	2	-	3	4	6	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	3	2	3	3	-	-	-
Male	-	-	-	-	2	1	-	-	2	-	-	-
Female	-	-	-	1	-	2	2	3	1	-	-	-
Michigan												
All races, both sexes	42	44	59	63	106	214	306	337	276	115	29	-
Male	28	27	34	38	59	120	142	145	110	28	6	-
Female	14	17	25	25	47	94	164	192	166	87	23	-
White, both sexes	25	33	49	53	93	190	264	301	257	111	26	-
Male	15	20	29	29	53	107	122	129	103	26	6	-
Female	10	13	20	24	40	83	142	172	154	85	20	-
Black, both sexes	15	11	9	8	13	22	41	34	18	4	3	-
Male	11	7	5	7	6	13	20	16	7	2	-	-
Female	4	4	4	1	7	9	21	18	11	2	3	-
American Indian 1/, both sexes	1	-	-	2	-	1	-	1	1	-	-	-
Male	1	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	1	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Minnesota												
All races, both sexes	8	9	12	15	27	48	96	137	144	75	10	-
Male	3	7	7	8	18	21	46	51	59	13	1	-
Female	5	2	5	7	9	27	50	86	85	62	9	-
White, both sexes	7	9	11	14	27	47	94	136	143	75	10	-
Male	3	7	6	7	18	21	45	51	58	13	1	-
Female	4	2	5	7	9	26	49	85	85	62	9	-
Black, both sexes	1	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Mississippi												
All races, both sexes	22	20	19	31	55	85	107	97	80	50	13	-
Male	12	13	11	12	35	40	51	32	21	16	1	-
Female	10	7	8	19	20	45	56	65	59	34	12	-
White, both sexes	12	9	13	11	42	56	84	75	58	30	6	-
Male	5	5	6	5	25	27	42	23	13	10	1	-
Female	7	4	7	6	17	29	42	52	45	20	5	-
Black, both sexes	10	10	6	20	13	29	23	22	22	20	7	-
Male	7	7	5	7	10	13	9	9	8	6	-	-
Female	3	3	1	13	3	16	14	13	14	14	7	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Missouri												
All races, both sexes	25	28	31	44	75	154	216	271	220	104	33	1
Male	18	17	22	20	43	76	108	101	70	24	7	-
Female	7	11	9	24	32	78	108	170	150	80	26	1
White, both sexes	22	25	26	38	67	145	203	260	210	99	31	1
Male	16	15	19	18	41	72	103	100	64	23	6	-
Female	6	10	7	20	26	73	100	160	146	76	25	1
Black, both sexes	3	2	4	6	7	8	13	11	9	5	2	-
Male	2	2	3	2	2	4	5	1	6	1	1	-
Female	1	-	1	4	5	4	8	10	3	4	1	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	5	2	6	5	11	13	25	37	34	15	3	-
Male	3	2	4	3	4	10	11	19	11	4	-	-
Female	2	-	2	2	7	3	14	18	23	11	3	-
White, both sexes	4	1	5	5	11	13	22	34	34	14	3	-
Male	2	1	3	3	4	10	9	18	11	4	-	-
Female	2	-	2	2	7	3	13	16	23	10	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	1	-	-	-	3	2	-	1	-	-
Male	1	1	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Nebraska												
All races, both sexes	5	5	8	10	13	35	40	68	80	36	13	-
Male	2	4	6	5	8	18	17	30	31	12	2	-
Female	3	1	2	5	5	17	23	38	49	24	11	-
White, both sexes	4	5	7	10	13	34	38	68	79	36	13	-
Male	1	4	6	5	8	17	17	30	31	12	2	-
Female	3	1	1	5	5	17	21	38	48	24	11	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	13	18	24	34	52	69	71	60	36	13	2	-
Male	8	14	13	19	26	42	32	27	15	6	1	-
Female	5	4	11	15	26	27	39	33	21	7	1	-
White, both sexes	9	15	20	34	44	59	65	55	34	13	2	-
Male	4	11	11	19	22	34	30	26	15	6	1	-
Female	5	4	9	15	22	25	35	29	19	7	1	-
Black, both sexes	4	2	3	-	5	6	4	2	1	-	-	-
Male	4	2	2	-	3	6	1	1	-	-	-	-
Female	-	-	1	-	2	-	3	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	2	4	2	3	-	-	-	-
Male	-	1	-	-	1	2	1	-	-	-	-	-
Female	-	-	1	-	1	2	1	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
New Hampshire												
All races, both sexes	4	2	3	6	9	22	32	56	50	20	2	-
Male	3	1	1	4	4	17	20	32	20	6	1	-
Female	1	1	2	2	5	5	12	24	30	14	1	-
White, both sexes	4	2	3	6	8	21	32	55	49	19	2	-
Male	3	1	1	4	3	16	20	32	19	6	1	-
Female	1	1	2	2	5	5	12	23	30	13	1	-
Black, both sexes	-	-	-	-	-	1	-	-	1	1	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
New Jersey												
All races, both sexes	16	30	31	43	98	146	232	282	248	108	25	-
Male	10	21	22	24	64	64	109	121	99	28	8	-
Female	6	9	9	19	34	82	123	161	149	80	17	-
White, both sexes	12	23	25	36	91	128	208	266	227	98	23	-
Male	7	16	18	21	60	56	100	113	92	23	8	-
Female	5	7	7	15	31	72	108	153	135	75	15	-
Black, both sexes	4	7	4	7	7	14	21	15	15	8	2	-
Male	3	5	2	3	4	5	7	7	2	3	-	-
Female	1	2	2	4	3	9	14	8	13	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	3	3	1	6	2	-	-
Male	-	-	2	-	-	2	2	1	5	2	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
New Mexico												
All races, both sexes	9	7	14	10	24	37	64	53	60	24	11	-
Male	6	5	7	6	13	16	33	25	28	6	2	-
Female	3	2	7	4	11	21	31	28	32	18	9	-
White, both sexes	9	6	12	9	23	35	61	48	56	23	10	-
Male	6	5	7	5	12	16	31	22	26	6	2	-
Female	3	1	5	4	11	19	30	26	30	17	8	-
Black, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	2	1	1	2	3	4	3	1	1	-
Male	-	-	-	1	1	-	2	3	1	-	-	-
Female	-	1	2	-	-	2	1	1	2	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	109	111	155	188	313	565	874	1,042	857	390	92	-
Male	63	64	91	108	156	320	400	429	313	107	22	-
Female	46	47	64	80	157	245	474	613	544	283	70	-
White, both sexes	75	73	119	130	235	462	731	891	749	343	82	-
Male	43	45	64	77	118	257	342	370	275	96	20	-
Female	32	28	55	53	117	205	389	521	474	247	62	-
Black, both sexes	31	33	30	51	61	83	112	125	81	33	9	-
Male	19	15	21	26	28	51	44	48	21	5	2	-
Female	12	18	9	25	33	32	68	77	60	28	7	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	6	7	17	19	31	25	26	14	1	-
Male	1	4	6	5	10	12	14	11	17	6	-	-
Female	2	1	-	2	7	7	17	14	9	8	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
North Carolina												
All races, both sexes	30	55	53	72	106	202	281	329	281	116	45	-
Male	17	31	38	35	56	94	130	126	78	30	5	-
Female	13	24	15	37	50	108	151	203	203	86	40	-
White, both sexes	18	43	43	54	87	175	250	297	245	98	38	-
Male	7	24	32	25	48	83	116	114	67	25	4	-
Female	11	19	11	29	39	92	134	183	178	73	34	-
Black, both sexes	11	12	10	17	19	26	30	29	35	18	7	-
Male	9	7	6	10	8	11	14	10	10	5	1	-
Female	2	5	4	7	11	15	16	19	25	13	6	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	3	1	-	-	-
Male	1	-	-	-	-	-	-	2	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
North Dakota												
All races, both sexes	1	1	2	2	8	14	17	34	34	15	2	-
Male	1	1	1	1	6	8	9	18	13	-	-	-
Female	-	-	1	1	2	6	8	16	21	15	2	-
White, both sexes	-	1	2	2	7	14	17	34	34	15	2	-
Male	-	1	1	1	5	8	9	18	13	-	-	-
Female	-	-	1	1	2	6	8	16	21	15	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Ohio												
All races, both sexes	45	72	69	89	139	221	340	393	286	129	34	-
Male	24	36	34	50	78	106	154	159	103	25	7	-
Female	21	36	35	39	61	115	186	234	183	104	27	-
White, both sexes	37	59	58	74	125	203	314	374	273	122	30	-
Male	19	33	26	43	68	95	146	150	97	25	6	-
Female	18	26	32	31	57	108	168	224	176	97	24	-
Black, both sexes	7	13	11	15	14	18	26	18	13	7	3	-
Male	4	3	8	7	10	11	8	8	6	-	1	-
Female	3	10	3	8	4	7	18	10	7	7	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Oklahoma												
All races, both sexes	29	30	29	39	67	119	158	176	113	49	10	-
Male	13	16	18	21	32	65	74	76	40	13	3	-
Female	16	14	11	18	35	54	84	100	73	36	7	-
White, both sexes	23	23	24	33	60	111	148	165	109	45	8	-
Male	10	13	15	19	27	62	72	71	37	12	3	-
Female	13	10	9	14	33	49	76	94	72	33	5	-
Black, both sexes	6	4	2	2	3	6	5	8	2	2	1	-
Male	3	2	-	-	2	3	-	3	2	-	-	-
Female	3	2	2	2	1	3	5	5	-	2	1	-
American Indian 1/, both sexes	-	3	3	4	4	2	5	3	2	2	1	-
Male	-	1	3	2	3	-	2	2	1	1	-	-
Female	-	2	-	2	1	2	3	1	1	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Oregon												
All races, both sexes	7	12	14	19	35	44	91	112	109	44	11	-
Male	3	9	7	11	19	21	51	58	47	16	1	-
Female	4	3	7	8	16	23	40	54	62	28	10	-
White, both sexes	7	12	13	19	35	44	89	112	108	43	10	-
Male	3	9	7	11	19	21	50	58	47	16	1	-
Female	4	3	6	8	16	23	39	54	61	27	9	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	1	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Pennsylvania												
All races, both sexes	44	55	71	81	149	280	503	658	489	226	50	-
Male	28	35	42	51	77	150	258	285	162	49	7	-
Female	16	20	29	30	72	130	245	373	327	177	43	-
White, both sexes	33	42	59	64	133	258	469	627	470	216	49	-
Male	18	27	36	38	68	141	243	273	157	47	7	-
Female	15	15	23	26	65	117	226	354	313	169	42	-
Black, both sexes	10	13	12	16	15	20	31	30	19	9	1	-
Male	9	8	6	12	8	9	15	12	5	2	-	-
Female	1	5	6	4	7	11	16	18	14	7	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	2	3	1	-	-	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	3	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Rhode Island												
All races, both sexes	3	5	7	10	14	31	45	55	55	23	10	-
Male	2	4	4	6	5	16	21	23	18	4	2	-
Female	1	1	3	4	9	15	24	32	37	19	8	-
White, both sexes	3	5	7	8	14	29	45	53	53	22	9	-
Male	2	4	4	4	5	15	21	23	16	3	2	-
Female	1	1	3	4	9	14	24	30	37	19	7	-
Black, both sexes	-	-	-	2	-	1	-	1	1	-	1	-
Male	-	-	-	2	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
South Carolina												
All races, both sexes	21	33	26	48	43	86	129	138	97	39	13	-
Male	12	21	18	28	26	43	64	48	35	12	5	-
Female	9	12	8	20	17	43	65	90	62	27	8	-
White, both sexes	11	22	17	26	29	73	105	117	76	35	7	-
Male	7	13	9	17	18	35	53	40	28	10	2	-
Female	4	9	8	9	11	38	52	77	48	25	5	-
Black, both sexes	10	11	9	22	14	11	24	21	21	4	6	-
Male	5	8	9	11	8	8	11	8	7	2	3	-
Female	5	3	-	11	6	3	13	13	14	2	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
South Dakota												
All races, both sexes	4	-	1	4	6	7	27	44	37	19	7	-
Male	2	-	-	4	4	4	14	21	13	1	2	-
Female	2	-	1	-	2	3	13	23	24	18	5	-
White, both sexes	3	-	1	3	4	5	26	44	37	19	7	-
Male	2	-	-	3	2	3	13	21	13	1	2	-
Female	1	-	1	-	2	2	13	23	24	18	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	2	2	1	-	-	-	-	-
Male	-	-	-	1	2	1	1	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	38	38	54	73	115	184	291	329	218	84	18	-
Male	23	18	27	34	56	99	127	129	65	20	3	-
Female	15	20	27	39	59	85	164	200	153	64	15	-
White, both sexes	24	31	42	66	102	158	259	311	205	78	15	-
Male	14	14	17	30	48	86	111	124	60	19	3	-
Female	10	17	25	36	54	72	148	187	145	59	12	-
Black, both sexes	14	7	12	7	12	26	31	18	13	5	3	-
Male	9	4	10	4	7	13	15	5	5	-	-	-
Female	5	3	2	3	5	13	16	13	8	5	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	1	-	-
Male	-	-	-	-	1	-	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Texas												
All races, both sexes	88	105	117	168	235	364	609	622	517	228	43	-
Male	51	56	66	90	142	187	253	252	159	70	13	-
Female	37	49	51	78	93	177	356	370	358	158	30	-
White, both sexes	69	83	96	143	214	323	547	576	473	202	35	-
Male	41	43	55	76	129	164	232	240	145	60	9	-
Female	28	40	41	67	85	159	315	336	328	142	26	-
Black, both sexes	17	20	20	21	17	29	53	39	39	25	7	-
Male	10	12	10	13	11	18	17	11	11	9	4	-
Female	7	8	10	8	6	11	36	28	28	16	3	-
American Indian 1/, both sexes	2	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	2	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	2	4	12	9	7	5	1	1	-
Male	-	1	1	-	2	5	4	1	3	1	-	-
Female	-	1	-	2	2	7	5	6	2	-	1	-
Utah												
All races, both sexes	8	12	9	15	21	37	45	72	66	23	5	-
Male	4	4	4	5	9	17	24	37	24	7	1	-
Female	4	8	5	10	12	20	21	35	42	16	4	-
White, both sexes	8	11	8	14	21	36	43	71	65	23	3	-
Male	4	3	4	4	9	17	23	37	23	7	1	-
Female	4	8	4	10	12	19	20	34	42	16	2	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	1	1	-	1	-	-	-
Male	-	1	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Vermont												
All races, both sexes	-	1	2	2	5	8	9	18	13	6	3	-
Male	-	1	1	2	3	4	5	9	3	1	-	-
Female	-	-	1	-	2	4	4	9	10	5	3	-
White, both sexes	-	1	2	2	5	8	9	18	13	6	3	-
Male	-	1	1	2	3	4	5	9	3	1	-	-
Female	-	-	1	-	2	4	4	9	10	5	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	26	23	35	59	86	151	248	252	205	89	23	-
Male	14	14	21	32	45	88	138	108	70	18	3	-
Female	12	9	14	27	41	63	110	144	135	71	20	-
White, both sexes	14	18	30	47	72	127	210	229	180	78	20	-
Male	8	11	18	26	36	76	117	100	65	15	3	-
Female	6	7	12	21	36	51	93	129	115	63	17	-
Black, both sexes	12	5	3	12	11	20	37	18	22	9	3	-
Male	6	3	1	6	7	10	20	6	4	2	-	-
Female	6	2	2	6	4	10	17	12	18	7	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	3	4	1	5	3	2	-	-
Male	-	-	2	-	2	2	1	2	1	1	-	-
Female	-	-	-	-	1	2	-	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Washington												
All races, both sexes	21	21	30	33	37	82	128	164	135	79	16	-
Male	13	13	17	18	17	46	55	86	46	27	4	-
Female	8	8	13	15	20	36	73	78	89	52	12	-
White, both sexes	18	18	27	29	31	77	117	159	131	74	16	-
Male	12	12	16	15	16	43	50	84	44	25	4	-
Female	6	6	11	14	15	34	67	75	87	49	12	-
Black, both sexes	2	2	-	1	1	2	2	1	-	2	-	-
Male	1	1	-	1	-	-	1	-	-	1	-	-
Female	1	1	-	-	1	2	1	1	-	1	-	-
American Indian 1/, both sexes	1	1	-	1	3	1	4	1	-	-	-	-
Male	-	-	-	1	1	1	2	-	-	-	-	-
Female	1	1	-	-	2	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	2	2	2	5	3	4	3	-	-
Male	-	-	1	1	-	2	2	2	2	1	-	-
Female	-	-	2	1	2	-	3	1	2	2	-	-
West Virginia												
All races, both sexes	6	11	12	24	34	59	85	87	65	31	12	-
Male	5	9	7	12	21	32	39	40	21	14	2	-
Female	1	2	5	12	13	27	46	47	44	17	10	-
White, both sexes	6	11	12	24	33	58	85	85	64	28	11	-
Male	5	9	7	12	20	31	39	39	21	12	2	-
Female	1	2	5	12	13	27	46	46	43	16	9	-
Black, both sexes	-	-	-	-	1	1	-	2	-	3	1	-
Male	-	-	-	-	1	1	-	1	-	2	-	-
Female	-	-	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Wisconsin												
All races, both sexes	16	15	25	32	49	111	168	222	192	97	26	-
Male	11	11	15	16	29	66	79	92	79	29	8	-
Female	5	4	10	16	20	45	89	130	113	68	18	-
White, both sexes	15	13	23	27	48	107	162	217	189	94	26	-
Male	10	10	15	13	29	63	76	89	78	29	8	-
Female	5	3	8	14	19	44	86	128	111	65	18	-
Black, both sexes	1	2	2	5	1	4	4	2	2	1	-	-
Male	1	1	-	3	-	3	2	1	1	-	-	-
Female	-	1	2	2	1	1	2	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	3	-	1	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
Wyoming												
All races, both sexes	-	-	2	3	5	12	23	18	29	8	7	-
Male	-	-	1	2	3	4	15	6	12	1	-	-
Female	-	-	1	1	2	8	8	12	17	7	7	-
White, both sexes	-	-	2	3	4	12	23	18	29	8	7	-
Male	-	-	1	2	2	4	15	6	12	1	-	-
Female	-	-	1	1	2	8	8	12	17	7	7	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
United States												
All races, both sexes	242	58	19	4	2	3	3	1	1	2	10	5
Male	116	36	9	2	1	2	-	-	-	2	6	2
Female	126	22	10	2	1	1	3	1	1	-	4	3
White, both sexes	198	34	13	4	1	3	3	1	1	2	8	5
Male	90	22	6	2	1	2	-	-	-	2	5	2
Female	108	12	7	2	-	1	3	1	1	-	3	3
Black, both sexes	33	20	4	-	1	-	-	-	-	-	2	-
Male	19	13	1	-	-	-	-	-	-	-	1	-
Female	14	7	3	-	1	-	-	-	-	-	1	-
American Indian 1/, both sexes	4	1	2	-	-	-	-	-	-	-	-	-
Male	2	-	2	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	3	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	10	1	2	-	1	1	-	-	-	-	1	-
Male	5	1	-	-	-	1	-	-	-	-	1	-
Female	5	-	2	-	1	-	-	-	-	-	-	-
White, both sexes	6	1	1	-	-	1	-	-	-	-	-	-
Male	4	1	-	-	-	1	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	1	-	1	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Arkansas												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	38	5	2	-	-	-	2	-	-	-	1	1
Male	19	2	1	-	-	-	-	-	-	-	1	1
Female	19	3	1	-	-	-	2	-	-	-	-	-
White, both sexes	34	2	2	-	-	-	2	-	-	-	1	1
Male	17	1	1	-	-	-	-	-	-	-	1	1
Female	17	1	1	-	-	-	2	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Colorado												
All races, both sexes	7	1	1	1	-	-	-	-	-	-	1	-
Male	3	-	1	1	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	6	1	-	1	-	-	-	-	-	-	1	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Florida												
All races, both sexes	13	6	-	-	-	-	-	-	-	-	1	-
Male	4	2	-	-	-	-	-	-	-	-	-	-
Female	9	4	-	-	-	-	-	-	-	-	1	-
White, both sexes	7	3	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	5	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	3	-	-	-	-	-	-	-	-	1	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	4	2	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	8	2	1	-	1	-	-	-	-	-	2	-
Male	6	2	1	-	1	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	4	-	-	-	1	-	-	-	-	-	2	-
Male	2	-	-	-	1	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	2	-	-	-	-	-	-	-	-	-	-
Male	3	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Illinois												
All races, both sexes	9	1	1	-	-	-	-	-	1	-	-	-
Male	5	-	1	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	2	1	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	6	5	1	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	4	3	1	-	-	-	-	-	-	-	-	-
White, both sexes	4	3	1	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	2	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Iowa												
All races, both sexes	3	1	-	-	-	-	-	-	-	1	-	-
Male	2	1	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	-	-	-	1	-	-
Male	2	1	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Kentucky												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Maine												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Massachusetts												
All races, both sexes	4	2	-	-	-	-	-	-	-	-	-	1
Male	3	2	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	-	-	-	-	-	1
Male	2	1	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	11	1	1	-	-	-	-	-	-	-	-	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	8	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	1	1	-	-	-	-	-	-	-	-	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	8	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Minnesota												
All races, both sexes	4	2	-	-	-	-	-	-	-	1	-	-
Male	3	2	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	-	-	-	1	-	-
Male	2	1	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	2	-	-	-	-	-	1	-	-	-	-
Male	3	2	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Missouri												
All races, both sexes	4	1	-	-	-	-	-	-	-	-	1	-
Male	3	1	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Nebraska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	1	3	-	-	-	1	-	-	-	-	-
Male	3	-	3	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	1	-	-	-	-	-
White, both sexes	5	1	3	-	-	-	1	-	-	-	-	-
Male	3	-	3	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
New Mexico												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	15	4	2	-	-	-	-	-	-	-	-	1
Male	6	3	-	-	-	-	-	-	-	-	-	-
Female	9	1	2	-	-	-	-	-	-	-	-	1
White, both sexes	11	2	-	-	-	-	-	-	-	-	-	1
Male	5	2	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	2	2	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	3	1	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
North Carolina												
All races, both sexes	10	4	1	1	-	2	-	-	-	-	-	-
Male	5	3	-	1	-	1	-	-	-	-	-	-
Female	5	1	1	-	-	1	-	-	-	-	-	-
White, both sexes	8	2	1	1	-	2	-	-	-	-	-	-
Male	4	2	-	1	-	1	-	-	-	-	-	-
Female	4	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Ohio												
All races, both sexes	9	2	-	-	-	-	-	-	-	-	1	1
Male	5	2	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	8	1	-	-	-	-	-	-	-	-	1	1
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	8	1	1	1	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	1	1	-	-	-	-	-	-	-	-
White, both sexes	7	-	1	1	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	1	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Oregon												
All races, both sexes	7	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	9	2	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	5	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	2	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	5	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Texas												
All races, both sexes	12	2	2	-	-	-	-	-	-	-	-	-
Male	4	1	1	-	-	-	-	-	-	-	-	-
Female	8	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	11	2	2	-	-	-	-	-	-	-	-	-
Male	3	1	1	-	-	-	-	-	-	-	-	-
Female	8	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	4	2	1	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	4	2	1	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Washington												
All races, both sexes	7	3	-	-	-	-	-	-	-	-	-	-
Male	5	3	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	2	-	-	-	-	-	-	-	-	-	-
Male	4	2	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	6	2	-	-	-	-	-	-	-	-	-	1
Male	3	2	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	6	2	-	-	-	-	-	-	-	-	-	1
Male	3	2	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Wisconsin												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
United States												
All races, both sexes	6	4	4	8	13	19	27	25	21	6	1	-
Male	2	3	2	4	8	9	12	9	6	1	-	-
Female	4	1	2	4	5	10	15	16	15	5	1	-
White, both sexes	6	3	4	7	9	18	24	24	21	6	1	-
Male	2	2	2	3	5	9	10	8	6	1	-	-
Female	4	1	2	4	4	9	14	16	15	5	1	-
Black, both sexes	-	1	-	1	2	-	1	1	-	-	-	-
Male	-	1	-	1	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	2	-	-	-	-	-
Male	-	-	-	-	2	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	1	1	-	2	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Arkansas												
All races, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2	1	1	1	5	3	2	8	3	1	-	-
Male	-	1	-	1	4	1	1	4	2	-	-	-
Female	2	-	1	-	1	2	1	4	1	1	-	-
White, both sexes	2	1	1	1	4	3	2	8	3	1	-	-
Male	-	1	-	1	3	1	1	4	2	-	-	-
Female	2	-	1	-	1	2	1	4	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Colorado												
All races, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Florida												
All races, both sexes	-	1	-	1	1	-	-	1	2	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	2	-	-	-
White, both sexes	-	1	-	-	-	-	-	1	2	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
Black, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Illinois												
All races, both sexes	1	-	-	-	-	2	2	1	-	-	-	-
Male	1	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	1	-	-	-	-	2	1	1	-	-	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Iowa												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Kentucky												
All races, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	5	1	-	2	-	1	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	3	1	-	2	-	1	-
White, both sexes	-	-	-	-	-	5	1	-	2	-	1	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	3	1	-	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Missouri												
All races, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
New Hampshire												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	1	1	1	1	1	3	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	3	-	-	-
White, both sexes	-	-	-	1	1	1	1	1	3	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
North Carolina												
All races, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Ohio												
All races, both sexes	-	-	-	-	-	-	3	1	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	1	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	3	1	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	1	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	1	-	-	1	1	2	-	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	1	-	-	1	1	2	-	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Oregon												
All races, both sexes	1	-	-	1	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	-	-	1	1	1	-	-	-
White, both sexes	1	-	-	1	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	1	-	2	3	-	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	2	3	-	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Texas												
All races, both sexes	2	-	-	1	1	-	2	1	-	1	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	2	1	-	1	-	-
White, both sexes	2	-	-	1	-	-	2	1	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	2	1	-	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Washington												
All races, both sexes	-	-	1	-	-	-	2	1	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	1	-	-	-	2	1	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	2	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	2	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
United States												
All races, both sexes	4,566	43	29	4	3	11	20	17	26	38	81	124
Male	2,395	23	12	3	1	3	13	11	9	18	41	63
Female	2,171	20	17	1	2	8	7	6	17	20	40	61
White, both sexes	3,939	28	18	3	3	10	17	14	17	33	65	82
Male	2,065	16	7	3	1	2	11	9	7	17	35	38
Female	1,874	12	11	-	2	8	6	5	10	16	30	44
Black, both sexes	483	11	11	-	-	-	3	3	6	4	11	40
Male	246	6	5	-	-	-	2	2	1	1	4	23
Female	237	5	6	-	-	-	1	1	5	3	7	17
American Indian 1/, both sexes	30	1	-	-	-	1	-	-	1	-	4	2
Male	18	-	-	-	-	1	-	-	-	-	2	2
Female	12	1	-	-	-	-	-	-	1	-	2	-
Asian or Pacific Islander 2/, both sexes	114	3	-	1	-	-	-	-	2	1	1	-
Male	66	1	-	-	-	-	-	-	1	-	-	-
Female	48	2	-	1	-	-	-	-	1	1	1	-
Alabama												
All races, both sexes	57	-	-	-	1	-	-	-	-	-	-	3
Male	29	-	-	-	1	-	-	-	-	-	-	2
Female	28	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	47	-	-	-	1	-	-	-	-	-	-	1
Male	21	-	-	-	1	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Alaska												
All races, both sexes	6	1	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	396	2	2	-	-	2	-	-	3	4	6	7
Male	208	2	1	-	-	1	-	-	-	1	2	2
Female	188	-	1	-	-	1	-	-	3	3	4	5
White, both sexes	370	1	2	-	-	1	-	-	2	3	4	6
Male	195	1	1	-	-	-	-	-	-	1	2	1
Female	175	-	1	-	-	1	-	-	2	2	2	5
Black, both sexes	9	1	-	-	-	-	-	-	-	1	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	14	-	-	-	-	1	-	-	1	-	2	1
Male	6	-	-	-	-	1	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	1	-	2	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Arkansas												
All races, both sexes	35	-	-	-	-	-	1	-	-	1	-	-
Male	18	-	-	-	-	-	1	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	1	-	-	1	-	-
Male	15	-	-	-	-	-	1	-	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	738	5	4	-	-	1	2	2	3	6	18	19
Male	384	1	2	-	-	-	1	1	-	5	10	11
Female	354	4	2	-	-	1	1	1	3	1	8	8
White, both sexes	628	3	3	-	-	1	2	2	1	6	15	14
Male	319	1	1	-	-	-	1	1	-	5	9	8
Female	309	2	2	-	-	1	1	1	1	1	6	6
Black, both sexes	46	1	1	-	-	-	-	-	1	-	1	5
Male	26	-	1	-	-	-	-	-	-	-	-	3
Female	20	1	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	61	1	-	-	-	-	-	-	1	-	1	-
Male	37	-	-	-	-	-	-	-	-	-	-	-
Female	24	1	-	-	-	-	-	-	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Colorado												
All races, both sexes	55	1	1	-	-	-	-	2	-	-	-	3
Male	30	1	-	-	-	-	-	2	-	-	-	1
Female	25	-	1	-	-	-	-	-	-	-	-	2
White, both sexes	51	1	1	-	-	-	-	2	-	-	-	3
Male	29	1	-	-	-	-	-	2	-	-	-	1
Female	22	-	1	-	-	-	-	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	36	-	-	-	-	-	-	-	1	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Delaware												
All races, both sexes	13	-	-	-	-	-	1	-	-	-	-	1
Male	7	-	-	-	-	-	1	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	1	-	-	-	-	1
Male	3	-	-	-	-	-	1	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Florida												
All races, both sexes	235	3	1	-	-	-	3	1	-	-	3	3
Male	128	2	-	-	-	-	1	1	-	-	2	1
Female	107	1	1	-	-	-	2	-	-	-	1	2
White, both sexes	216	2	-	-	-	-	2	1	-	-	3	2
Male	118	1	-	-	-	-	1	1	-	-	2	1
Female	98	1	-	-	-	-	1	-	-	-	1	1
Black, both sexes	19	1	1	-	-	-	1	-	-	-	-	1
Male	10	1	-	-	-	-	-	-	-	-	-	-
Female	9	-	1	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	79	1	2	-	-	1	1	-	1	-	3	3
Male	48	1	1	-	-	-	1	-	-	-	2	2
Female	31	-	1	-	-	1	-	-	1	-	1	1
White, both sexes	64	-	-	-	-	1	1	-	-	-	2	2
Male	38	-	-	-	-	-	1	-	-	-	2	1
Female	26	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	15	1	2	-	-	-	-	-	1	-	1	1
Male	10	1	1	-	-	-	-	-	-	-	-	1
Female	5	-	1	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Hawaii												
All races, both sexes	13	1	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	1	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Illinois												
All races, both sexes	103	1	1	2	-	-	-	-	2	1	1	2
Male	56	-	1	2	-	-	-	-	2	-	1	1
Female	47	1	-	-	-	-	-	-	-	1	-	1
White, both sexes	93	1	1	2	-	-	-	-	1	1	1	1
Male	47	-	1	2	-	-	-	-	1	-	1	-
Female	46	1	-	-	-	-	-	-	-	1	-	1
Black, both sexes	10	-	-	-	-	-	-	-	1	-	-	1
Male	9	-	-	-	-	-	-	-	1	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	54	-	-	-	-	-	2	-	-	-	1	2
Male	36	-	-	-	-	-	2	-	-	-	-	2
Female	18	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	47	-	-	-	-	-	1	-	-	-	1	1
Male	31	-	-	-	-	-	1	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	-	-	1	-	-	-	-	1
Male	5	-	-	-	-	-	1	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Iowa												
All races, both sexes	27	1	-	-	-	-	1	-	-	-	-	1
Male	15	1	-	-	-	-	1	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	1	-	-	-	-	1	-	-	-	-	1
Male	15	1	-	-	-	-	1	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	27	1	1	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	13	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	26	1	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	12	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Kentucky												
All races, both sexes	67	1	-	-	-	-	-	-	-	1	-	1
Male	38	-	-	-	-	-	-	-	-	1	-	-
Female	29	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	63	1	-	-	-	-	-	-	-	1	-	1
Male	34	-	-	-	-	-	-	-	-	1	-	-
Female	29	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	31	-	1	-	-	-	1	-	-	-	-	4
Male	20	-	1	-	-	-	1	-	-	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	21	-	-	-	-	-	1	-	-	-	-	2
Male	12	-	-	-	-	-	1	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	10	-	1	-	-	-	-	-	-	-	-	2
Male	8	-	1	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Maine												
All races, both sexes	11	1	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	1	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	75	2	-	-	-	-	1	-	-	-	2	5
Male	43	-	-	-	-	-	1	-	-	-	1	4
Female	32	2	-	-	-	-	-	-	-	-	1	1
White, both sexes	61	2	-	-	-	-	1	-	-	-	1	2
Male	32	-	-	-	-	-	1	-	-	-	-	2
Female	29	2	-	-	-	-	-	-	-	-	1	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	3
Male	11	-	-	-	-	-	-	-	-	-	1	2
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Massachusetts												
All races, both sexes	74	-	2	-	-	-	-	-	-	1	1	-
Male	37	-	-	-	-	-	-	-	-	-	1	-
Female	37	-	2	-	-	-	-	-	-	1	-	-
White, both sexes	71	-	2	-	-	-	-	-	-	1	1	-
Male	36	-	-	-	-	-	-	-	-	-	1	-
Female	35	-	2	-	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	97	-	-	-	-	-	-	-	-	-	2	2
Male	53	-	-	-	-	-	-	-	-	-	1	1
Female	44	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	86	-	-	-	-	-	-	-	-	-	2	1
Male	43	-	-	-	-	-	-	-	-	-	1	-
Female	43	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Minnesota												
All races, both sexes	32	1	1	-	-	-	-	-	-	-	1	2
Male	18	1	1	-	-	-	-	-	-	-	-	2
Female	14	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	29	-	1	-	-	-	-	-	-	-	1	2
Male	17	-	1	-	-	-	-	-	-	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	40	1	-	-	-	-	-	-	-	1	-	3
Male	25	1	-	-	-	-	-	-	-	-	-	3
Female	15	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	24	1	-	-	-	-	-	-	-	1	-	-
Male	14	1	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	3
Male	11	-	-	-	-	-	-	-	-	-	-	3
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Missouri												
All races, both sexes	58	1	-	-	-	-	-	1	1	-	-	2
Male	31	-	-	-	-	-	-	-	-	-	-	2
Female	27	1	-	-	-	-	-	1	1	-	-	-
White, both sexes	52	-	-	-	-	-	-	1	1	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	6	1	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	3	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Nebraska												
All races, both sexes	15	-	-	-	-	-	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	15	-	-	-	-	-	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	18	1	-	-	-	-	-	-	1	2	-	-
Male	14	-	-	-	-	-	-	-	-	2	-	-
Female	4	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	1	2	-	-
Male	13	-	-	-	-	-	-	-	-	2	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
New Hampshire												
All races, both sexes	29	-	1	-	-	-	1	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	1	-	-	-	1	-	-	-	-	1
White, both sexes	28	-	1	-	-	-	1	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	1	-	-	-	1	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	35	-	-	-	-	1	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	1	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
New Mexico												
All races, both sexes	24	2	-	-	-	-	1	-	-	-	-	2
Male	12	1	-	-	-	-	-	-	-	-	-	1
Female	12	1	-	-	-	-	1	-	-	-	-	1
White, both sexes	24	2	-	-	-	-	1	-	-	-	-	2
Male	12	1	-	-	-	-	-	-	-	-	-	1
Female	12	1	-	-	-	-	1	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	621	1	2	-	-	2	-	3	-	3	8	10
Male	301	-	1	-	-	1	-	1	-	2	5	5
Female	320	1	1	-	-	1	-	2	-	1	3	5
White, both sexes	449	1	1	-	-	2	-	2	-	3	5	6
Male	224	-	-	-	-	1	-	-	-	2	3	4
Female	225	1	1	-	-	1	-	2	-	1	2	2
Black, both sexes	151	-	1	-	-	-	-	1	-	-	3	4
Male	61	-	1	-	-	-	-	1	-	-	2	1
Female	90	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
North Carolina												
All races, both sexes	95	2	1	-	-	1	1	-	-	2	5	4
Male	54	-	1	-	-	1	-	-	-	-	3	1
Female	41	2	-	-	-	-	1	-	-	2	2	3
White, both sexes	79	1	1	-	-	1	1	-	-	1	3	2
Male	50	-	1	-	-	1	-	-	-	-	2	1
Female	29	1	-	-	-	-	1	-	-	1	1	1
Black, both sexes	16	1	-	-	-	-	-	-	-	1	2	2
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	12	1	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Ohio												
All races, both sexes	246	2	1	-	-	-	-	1	1	3	3	8
Male	114	2	1	-	-	-	-	1	1	2	1	2
Female	132	-	-	-	-	-	-	-	-	1	2	6
White, both sexes	232	1	-	-	-	-	-	-	1	3	3	6
Male	105	1	-	-	-	-	-	-	1	2	1	1
Female	127	-	-	-	-	-	-	-	-	1	2	5
Black, both sexes	14	1	1	-	-	-	-	1	-	-	-	2
Male	9	1	1	-	-	-	-	1	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	67	-	1	-	-	1	-	1	-	1	4	-
Male	31	-	1	-	-	-	-	-	-	-	3	-
Female	36	-	-	-	-	1	-	1	-	1	1	-
White, both sexes	58	-	1	-	-	1	-	1	-	1	4	-
Male	26	-	1	-	-	-	-	-	-	-	3	-
Female	32	-	-	-	-	1	-	1	-	1	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Oregon												
All races, both sexes	61	-	-	-	-	-	1	-	-	-	2	1
Male	31	-	-	-	-	-	1	-	-	-	1	1
Female	30	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	61	-	-	-	-	-	1	-	-	-	2	1
Male	31	-	-	-	-	-	1	-	-	-	1	1
Female	30	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	96	-	1	-	-	-	-	-	1	1	2	3
Male	52	-	-	-	-	-	-	-	-	-	1	2
Female	44	-	1	-	-	-	-	-	1	1	1	1
White, both sexes	90	-	1	-	-	-	-	-	-	1	2	3
Male	49	-	-	-	-	-	-	-	-	-	1	2
Female	41	-	1	-	-	-	-	-	-	1	1	1
Black, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Rhode Island												
All races, both sexes	15	-	-	-	-	-	-	1	-	1	-	-
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	14	-	-	-	-	-	-	1	-	-	-	-
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	68	-	-	-	-	-	-	1	2	3	1	1
Male	35	-	-	-	-	-	-	1	-	1	-	1
Female	33	-	-	-	-	-	-	-	2	2	1	-
White, both sexes	51	-	-	-	-	-	-	1	1	3	1	-
Male	29	-	-	-	-	-	-	1	-	1	-	-
Female	22	-	-	-	-	-	-	-	1	2	1	-
Black, both sexes	17	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
South Dakota												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	151	1	-	-	-	-	1	-	-	2	2	5
Male	73	-	-	-	-	-	1	-	-	1	-	3
Female	78	1	-	-	-	-	-	-	-	1	2	2
White, both sexes	138	-	-	-	-	-	1	-	-	1	2	3
Male	67	-	-	-	-	-	1	-	-	-	-	3
Female	71	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	12	1	-	-	-	-	-	-	-	1	-	2
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	7	1	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Texas												
All races, both sexes	280	4	2	1	1	1	-	2	5	2	11	10
Male	136	4	-	1	-	-	-	1	3	1	4	3
Female	144	-	2	-	1	1	-	1	2	1	7	7
White, both sexes	238	2	1	1	1	1	-	1	4	2	8	6
Male	122	2	-	1	-	-	-	1	2	1	4	1
Female	116	-	1	-	1	1	-	-	2	1	4	5
Black, both sexes	34	2	1	-	-	-	-	1	-	-	3	4
Male	10	2	-	-	-	-	-	-	-	-	-	2
Female	24	-	1	-	-	-	-	1	-	-	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	21	-	-	1	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	1	-	-	-	-	-	-	-	1
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Vermont												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	63	2	-	-	-	-	1	-	-	-	2	1
Male	44	2	-	-	-	-	1	-	-	-	1	1
Female	19	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	54	2	-	-	-	-	1	-	-	-	2	1
Male	38	2	-	-	-	-	1	-	-	-	1	1
Female	16	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Washington												
All races, both sexes	156	1	2	-	1	-	1	1	2	3	2	4
Male	84	1	1	-	-	-	-	1	2	1	1	2
Female	72	-	1	-	1	-	1	-	-	2	1	2
White, both sexes	141	1	1	-	1	-	1	1	2	2	2	4
Male	77	1	1	-	-	-	-	1	2	1	1	2
Female	64	-	-	-	1	-	1	-	-	1	1	2
Black, both sexes	7	-	1	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
West Virginia												
All races, both sexes	24	-	1	-	-	1	-	1	1	-	-	1
Male	14	-	-	-	-	-	-	1	-	-	-	1
Female	10	-	1	-	-	1	-	-	1	-	-	-
White, both sexes	22	-	-	-	-	1	-	1	1	-	-	1
Male	14	-	-	-	-	-	-	1	-	-	-	1
Female	8	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Wisconsin												
All races, both sexes	56	3	-	-	-	-	-	-	1	-	-	1
Male	28	3	-	-	-	-	-	-	1	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	51	3	-	-	-	-	-	-	1	-	-	1
Male	24	3	-	-	-	-	-	-	1	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
United States												
All races, both sexes	196	196	241	287	377	609	797	738	492	196	41	-
Male	120	133	146	169	224	355	423	364	200	57	7	-
Female	76	63	95	118	153	254	374	374	292	139	34	-
White, both sexes	151	165	207	232	333	535	701	666	444	178	37	-
Male	89	110	124	144	193	315	371	336	181	51	5	-
Female	62	55	83	88	140	220	330	330	263	127	32	-
Black, both sexes	40	22	26	46	29	53	66	59	37	13	3	-
Male	28	16	18	20	19	31	29	23	12	4	2	-
Female	12	6	8	26	10	22	37	36	25	9	1	-
American Indian 1/, both sexes	3	4	1	2	1	2	4	3	-	1	-	-
Male	1	2	-	2	1	-	4	2	-	1	-	-
Female	2	2	1	-	-	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	7	7	14	19	26	10	11	4	1	-
Male	2	5	4	3	11	9	19	3	7	1	-	-
Female	-	-	3	4	3	10	7	7	4	3	1	-
Alabama												
All races, both sexes	3	3	2	3	6	9	14	6	7	-	-	-
Male	3	2	2	2	2	4	6	3	2	-	-	-
Female	-	1	-	1	4	5	8	3	5	-	-	-
White, both sexes	2	3	2	2	6	9	13	4	4	-	-	-
Male	2	2	2	1	2	4	5	2	-	-	-	-
Female	-	1	-	1	4	5	8	2	4	-	-	-
Black, both sexes	-	-	-	1	-	-	1	2	3	-	-	-
Male	-	-	-	1	-	-	1	1	2	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Alaska												
All races, both sexes	-	-	-	2	-	-	-	1	-	-	-	-
Male	-	-	-	2	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	2	-	-	-	1	-	-	-	-
Male	-	-	-	2	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	18	17	33	32	32	45	62	76	36	15	4	-
Male	10	8	20	21	18	24	35	43	15	5	-	-
Female	8	9	13	11	14	21	27	33	21	10	4	-
White, both sexes	17	14	30	31	31	42	61	73	36	12	4	-
Male	9	7	18	20	17	23	34	42	15	4	-	-
Female	8	7	12	11	14	19	27	31	21	8	4	-
Black, both sexes	1	1	2	1	1	1	-	-	-	-	-	-
Male	1	-	2	1	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	1	-	-	2	1	2	-	1	-	-
Male	-	1	-	-	-	-	1	1	-	1	-	-
Female	-	1	1	-	-	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Arkansas												
All races, both sexes	2	2	1	1	4	2	9	8	4	-	-	-
Male	2	2	1	-	2	1	3	3	2	-	-	-
Female	-	-	-	1	2	1	6	5	2	-	-	-
White, both sexes	2	2	-	1	4	2	7	7	3	-	-	-
Male	2	2	-	-	2	1	3	2	1	-	-	-
Female	-	-	-	1	2	1	4	5	2	-	-	-
Black, both sexes	-	-	1	-	-	-	2	1	1	-	-	-
Male	-	-	1	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	19	21	32	40	55	91	150	138	82	42	8	-
Male	15	13	19	25	30	52	77	74	34	12	2	-
Female	4	8	13	15	25	39	73	64	48	30	6	-
White, both sexes	15	17	30	31	46	73	122	124	76	40	7	-
Male	12	9	18	20	23	41	59	67	31	11	2	-
Female	3	8	12	11	23	32	63	57	45	29	5	-
Black, both sexes	2	1	-	4	3	6	8	9	3	1	-	-
Male	2	1	-	3	3	4	3	4	1	1	-	-
Female	-	-	-	1	-	2	5	5	2	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	2	5	6	12	19	5	3	1	1	-
Male	1	3	1	2	4	7	14	3	2	-	-	-
Female	-	-	1	3	2	5	5	2	1	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Colorado												
All races, both sexes	3	6	2	2	4	6	6	14	3	1	1	-
Male	1	5	2	-	1	4	3	8	1	1	-	-
Female	2	1	-	2	3	2	3	6	2	-	1	-
White, both sexes	3	6	2	1	4	4	6	13	3	1	1	-
Male	1	5	2	-	1	4	3	7	1	1	-	-
Female	2	1	-	1	3	-	3	6	2	-	1	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	1	-	2	2	4	9	6	8	1	-	-
Male	1	1	-	1	-	3	5	4	8	-	-	-
Female	-	-	-	1	2	1	4	2	-	1	-	-
White, both sexes	1	1	-	2	2	4	9	6	8	1	-	-
Male	1	1	-	1	-	3	5	4	8	-	-	-
Female	-	-	-	1	2	1	4	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Delaware												
All races, both sexes	1	1	1	1	2	2	2	-	1	-	-	-
Male	1	-	1	-	1	2	-	-	-	-	-	-
Female	-	1	-	1	1	-	2	-	1	-	-	-
White, both sexes	-	1	1	-	2	2	2	-	1	-	-	-
Male	-	-	1	-	1	2	-	-	-	-	-	-
Female	-	1	-	-	1	-	2	-	1	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	1	-	1	2	-	-	-	-	-	-	-
Male	1	1	-	-	2	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	1	1	-	-	-	-	-	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Florida												
All races, both sexes	8	7	7	23	20	37	37	44	21	16	1	-
Male	5	5	4	14	16	25	18	20	9	4	1	-
Female	3	2	3	9	4	12	19	24	12	12	-	-
White, both sexes	7	7	5	19	17	35	36	42	21	16	1	-
Male	4	5	3	11	14	24	17	20	9	4	1	-
Female	3	2	2	8	3	11	19	22	12	12	-	-
Black, both sexes	1	-	2	4	3	2	1	2	-	-	-	-
Male	1	-	1	3	2	1	1	-	-	-	-	-
Female	-	-	1	1	1	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	5	4	2	8	8	6	15	10	8	1	-	-
Male	3	3	2	4	4	2	11	7	5	-	-	-
Female	2	1	-	4	4	4	4	3	3	1	-	-
White, both sexes	4	3	-	6	8	5	14	10	7	1	-	-
Male	2	2	-	3	4	2	10	7	4	-	-	-
Female	2	1	-	3	4	3	4	3	3	1	-	-
Black, both sexes	1	1	2	2	-	1	1	-	1	-	-	-
Male	1	1	2	1	-	-	1	-	1	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Hawaii												
All races, both sexes	-	-	1	1	2	1	2	2	2	-	-	-
Male	-	-	1	-	1	-	1	1	1	-	-	-
Female	-	-	-	1	1	1	1	1	1	-	-	-
White, both sexes	-	-	1	-	-	-	1	2	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	1	-	2	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	1	1	1	-	-	1	-	-	-
Idaho												
All races, both sexes	-	2	1	-	1	3	3	1	1	-	-	-
Male	-	2	1	-	1	2	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
White, both sexes	-	2	1	-	1	3	3	1	1	-	-	-
Male	-	2	1	-	1	2	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Illinois												
All races, both sexes	3	4	4	5	7	13	23	22	9	3	-	-
Male	1	2	2	4	7	7	15	7	3	1	-	-
Female	2	2	2	1	-	6	8	15	6	2	-	-
White, both sexes	2	4	4	5	6	12	20	21	8	3	-	-
Male	-	2	2	4	6	7	12	6	2	1	-	-
Female	2	2	2	1	-	5	8	15	6	2	-	-
Black, both sexes	1	-	-	-	1	1	3	1	1	-	-	-
Male	1	-	-	-	1	-	3	1	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	7	4	4	1	3	11	9	5	4	1	-	-
Male	5	4	2	1	1	8	5	3	3	-	-	-
Female	2	-	2	-	2	3	4	2	1	1	-	-
White, both sexes	5	4	4	1	3	11	8	3	4	1	-	-
Male	4	4	2	1	1	8	5	1	3	-	-	-
Female	1	-	2	-	2	3	3	2	1	1	-	-
Black, both sexes	1	-	-	-	-	-	1	2	-	-	-	-
Male	1	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Iowa												
All races, both sexes	-	2	-	1	4	5	2	6	-	3	1	-
Male	-	1	-	-	3	2	1	4	-	1	-	-
Female	-	1	-	1	1	3	1	2	-	2	1	-
White, both sexes	-	2	-	1	4	5	2	6	-	3	1	-
Male	-	1	-	-	3	2	1	4	-	1	-	-
Female	-	1	-	1	1	3	1	2	-	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	1	1	3	-	3	6	4	4	1	1	-
Male	1	1	-	1	-	1	5	3	2	-	-	-
Female	-	-	1	2	-	2	1	1	2	1	1	-
White, both sexes	1	1	1	3	-	3	6	4	4	1	1	-
Male	1	1	-	1	-	1	5	3	2	-	-	-
Female	-	-	1	2	-	2	1	1	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Kentucky												
All races, both sexes	2	2	5	7	9	11	11	9	4	3	1	-
Male	1	1	3	6	6	6	6	5	2	1	-	-
Female	1	1	2	1	3	5	5	4	2	2	1	-
White, both sexes	2	2	4	7	8	11	10	8	4	3	1	-
Male	1	1	2	6	5	6	5	4	2	1	-	-
Female	1	1	2	1	3	5	5	4	2	2	1	-
Black, both sexes	-	-	1	-	1	-	1	1	-	-	-	-
Male	-	-	1	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	-	4	3	3	2	4	5	1	1	-	-
Male	2	-	2	2	3	1	-	4	1	1	-	-
Female	-	-	2	1	-	1	4	1	-	-	-	-
White, both sexes	1	-	3	2	1	2	4	4	-	1	-	-
Male	1	-	2	1	1	1	-	3	-	1	-	-
Female	-	-	1	1	-	1	4	1	-	-	-	-
Black, both sexes	1	-	1	1	2	-	-	1	1	-	-	-
Male	1	-	-	1	2	-	-	1	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Maine												
All races, both sexes	-	-	2	-	1	1	3	1	2	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	2	-	-	1	1	1	2	-	-	-
White, both sexes	-	-	2	-	1	1	3	1	2	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	2	-	-	1	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	8	5	3	3	8	12	8	8	8	2	-	-
Male	6	3	3	3	2	9	5	3	2	1	-	-
Female	2	2	-	-	6	3	3	5	6	1	-	-
White, both sexes	5	3	1	3	7	11	8	8	7	2	-	-
Male	3	1	1	3	2	8	5	3	2	1	-	-
Female	2	2	-	-	5	3	3	5	5	1	-	-
Black, both sexes	3	2	2	-	1	1	-	-	1	-	-	-
Male	3	2	2	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Massachusetts												
All races, both sexes	2	1	4	5	10	8	13	6	14	4	3	-
Male	1	-	4	2	8	3	8	2	6	1	1	-
Female	1	1	-	3	2	5	5	4	8	3	2	-
White, both sexes	2	1	4	4	10	7	13	6	14	4	2	-
Male	1	-	4	2	8	3	8	2	6	1	-	-
Female	1	1	-	2	2	4	5	4	8	3	2	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	8	6	4	7	11	14	19	7	13	3	1	-
Male	6	3	2	3	9	6	13	4	5	-	-	-
Female	2	3	2	4	2	8	6	3	8	3	1	-
White, both sexes	5	5	4	7	11	11	17	6	13	3	1	-
Male	3	2	2	3	9	4	11	3	5	-	-	-
Female	2	3	2	4	2	7	6	3	8	3	1	-
Black, both sexes	3	1	-	-	-	3	2	1	-	-	-	-
Male	3	1	-	-	-	2	2	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Minnesota												
All races, both sexes	2	1	2	4	2	5	1	4	5	1	-	-
Male	-	1	1	3	2	4	1	-	2	-	-	-
Female	2	-	1	1	-	1	-	4	3	1	-	-
White, both sexes	1	1	2	4	2	5	1	3	5	1	-	-
Male	-	1	1	3	2	4	1	-	2	-	-	-
Female	1	-	1	1	-	1	-	3	3	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Mississippi												
All races, both sexes	3	2	-	6	4	7	5	1	4	3	-	-
Male	1	2	-	3	3	5	2	1	2	2	-	-
Female	2	-	-	3	1	2	3	-	2	1	-	-
White, both sexes	2	-	-	2	3	6	5	1	1	2	-	-
Male	1	-	-	2	2	4	2	1	-	1	-	-
Female	1	-	-	-	1	2	3	-	1	1	-	-
Black, both sexes	1	2	-	4	1	1	-	-	3	1	-	-
Male	-	2	-	1	1	1	-	-	2	1	-	-
Female	1	-	-	3	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Missouri												
All races, both sexes	3	5	3	2	9	7	7	9	6	1	1	-
Male	2	3	2	1	5	4	3	5	3	1	-	-
Female	1	2	1	1	4	3	4	4	3	-	1	-
White, both sexes	3	5	3	2	6	7	7	9	6	1	1	-
Male	2	3	2	1	4	4	3	5	3	1	-	-
Female	1	2	1	1	2	3	4	4	3	-	1	-
Black, both sexes	-	-	-	-	3	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	1	1	2	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	-	2	-	1	-	1	-	-	-
White, both sexes	1	-	1	1	2	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	-	2	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Nebraska												
All races, both sexes	-	1	-	-	1	1	2	4	2	2	-	-
Male	-	1	-	-	1	-	-	4	2	2	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
White, both sexes	-	1	-	-	1	1	2	4	2	2	-	-
Male	-	1	-	-	1	-	-	4	2	2	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	3	1	1	1	1	2	2	1	1	-	-
Male	1	3	1	-	1	1	2	1	1	1	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	1	2	1	1	1	1	2	2	1	1	-	-
Male	1	2	1	-	1	1	2	1	1	1	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
New Hampshire												
All races, both sexes	2	-	-	2	1	9	4	4	3	1	-	-
Male	1	-	-	1	1	7	3	3	1	1	-	-
Female	1	-	-	1	-	2	1	1	2	-	-	-
White, both sexes	2	-	-	2	1	8	4	4	3	1	-	-
Male	1	-	-	1	1	6	3	3	1	1	-	-
Female	1	-	-	1	-	2	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	-	2	1	4	2	7	9	6	1	-	-
Male	1	-	2	1	3	2	3	4	3	-	-	-
Female	1	-	-	-	1	-	4	5	3	1	-	-
White, both sexes	2	-	1	1	4	2	6	7	6	1	-	-
Male	1	-	1	1	3	2	3	3	3	-	-	-
Female	1	-	-	-	1	-	3	4	3	1	-	-
Black, both sexes	-	-	1	-	-	-	1	2	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
New Mexico												
All races, both sexes	2	1	3	1	1	1	6	-	3	-	1	-
Male	1	1	3	1	-	-	4	-	-	-	-	-
Female	1	-	-	-	1	1	2	-	3	-	1	-
White, both sexes	2	1	3	1	1	1	6	-	3	-	1	-
Male	1	1	3	1	-	-	4	-	-	-	-	-
Female	1	-	-	-	1	1	2	-	3	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	20	26	27	28	39	94	111	119	84	35	9	-
Male	14	17	21	17	20	57	52	48	32	7	1	-
Female	6	9	6	11	19	37	59	71	52	28	8	-
White, both sexes	12	18	18	12	29	71	77	90	65	29	8	-
Male	10	13	13	10	13	43	38	40	27	6	1	-
Female	2	5	5	2	16	28	39	50	38	23	7	-
Black, both sexes	8	8	8	15	5	20	28	28	15	6	1	-
Male	4	4	7	6	2	12	10	8	2	1	-	-
Female	4	4	1	9	3	8	18	20	13	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	5	3	6	1	4	-	-	-
Male	-	-	1	1	5	2	4	-	3	-	-	-
Female	-	-	-	-	-	1	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
North Carolina												
All races, both sexes	1	9	10	3	8	13	13	12	8	2	-	-
Male	1	7	7	3	4	6	8	8	2	2	-	-
Female	-	2	3	-	4	7	5	4	6	-	-	-
White, both sexes	-	9	8	3	6	9	13	12	7	2	-	-
Male	-	7	7	3	4	4	8	8	2	2	-	-
Female	-	2	1	-	2	5	5	4	5	-	-	-
Black, both sexes	1	-	2	-	2	4	-	-	1	-	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	2	-	2	2	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	2	2	3	2	1	-	-	-	-
Male	1	-	-	1	2	1	2	1	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
White, both sexes	-	-	-	2	2	3	2	1	-	-	-	-
Male	-	-	-	1	2	1	2	1	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Ohio												
All races, both sexes	9	6	12	12	26	30	50	35	26	18	3	-
Male	3	4	6	8	15	19	24	13	8	4	-	-
Female	6	2	6	4	11	11	26	22	18	14	3	-
White, both sexes	9	6	12	10	25	28	47	35	25	18	3	-
Male	3	4	6	6	14	18	24	13	7	4	-	-
Female	6	2	6	4	11	10	23	22	18	14	3	-
Black, both sexes	-	-	-	2	1	2	3	-	1	-	-	-
Male	-	-	-	2	1	1	-	-	1	-	-	-
Female	-	-	-	-	-	1	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	5	3	2	3	5	11	11	12	6	1	-	-
Male	3	2	1	2	-	7	5	4	3	-	-	-
Female	2	1	1	1	5	4	6	8	3	1	-	-
White, both sexes	3	1	2	2	5	11	9	10	6	1	-	-
Male	2	-	1	1	-	7	4	4	3	-	-	-
Female	1	1	1	1	5	4	5	6	3	1	-	-
Black, both sexes	2	1	-	-	-	-	1	2	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Oregon												
All races, both sexes	5	3	4	4	3	5	12	8	8	5	-	-
Male	3	2	2	1	2	2	6	7	2	1	-	-
Female	2	1	2	3	1	3	6	1	6	4	-	-
White, both sexes	5	3	4	4	3	5	12	8	8	5	-	-
Male	3	2	2	1	2	2	6	7	2	1	-	-
Female	2	1	2	3	1	3	6	1	6	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	9	3	7	7	10	13	16	12	9	2	-	-
Male	4	3	3	5	6	9	7	6	6	-	-	-
Female	5	-	4	2	4	4	9	6	3	2	-	-
White, both sexes	8	3	6	6	10	12	15	12	9	2	-	-
Male	3	3	3	4	6	9	6	6	6	-	-	-
Female	5	-	3	2	4	3	9	6	3	2	-	-
Black, both sexes	1	-	1	1	-	1	1	-	-	-	-	-
Male	1	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Rhode Island												
All races, both sexes	1	1	-	1	3	2	2	-	3	-	-	-
Male	-	1	-	1	1	2	-	-	2	-	-	-
Female	1	-	-	-	2	-	2	-	1	-	-	-
White, both sexes	1	1	-	1	3	2	2	-	3	-	-	-
Male	-	1	-	1	1	2	-	-	2	-	-	-
Female	1	-	-	-	2	-	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	3	2	7	8	8	12	8	5	3	2	-
Male	1	1	2	2	7	4	8	2	2	1	2	-
Female	1	2	-	5	1	4	4	6	3	2	-	-
White, both sexes	2	2	2	2	7	7	9	8	2	3	1	-
Male	1	-	2	2	6	3	7	2	2	1	1	-
Female	1	2	-	-	1	4	2	6	-	2	-	-
Black, both sexes	-	1	-	5	1	1	3	-	3	-	1	-
Male	-	1	-	-	1	1	1	-	-	-	1	-
Female	-	-	-	5	-	-	2	-	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
South Dakota												
All races, both sexes	1	-	-	-	1	-	1	6	-	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	1	-	-	-	1	-	-	4	-	-	-	-
White, both sexes	1	-	-	-	1	-	1	6	-	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	1	-	-	-	1	-	-	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	8	12	14	12	22	29	22	13	4	2	-
Male	1	6	3	9	6	14	13	11	3	2	-	-
Female	1	2	9	5	6	8	16	11	10	2	2	-
White, both sexes	1	8	10	14	12	20	27	21	13	3	2	-
Male	1	6	2	9	6	13	11	11	3	1	-	-
Female	-	2	8	5	6	7	16	10	10	2	2	-
Black, both sexes	1	-	2	-	-	2	2	1	-	-	-	-
Male	-	-	1	-	-	1	2	-	-	-	-	-
Female	1	-	1	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Texas												
All races, both sexes	12	15	17	18	21	28	48	41	31	9	1	-
Male	7	10	10	8	16	12	25	20	9	2	-	-
Female	5	5	7	10	5	16	23	21	22	7	1	-
White, both sexes	8	13	16	16	19	26	45	33	29	5	1	-
Male	4	9	9	8	15	12	25	18	9	1	-	-
Female	4	4	7	8	4	14	20	15	20	4	1	-
Black, both sexes	4	1	-	2	1	-	3	6	2	4	-	-
Male	3	-	-	-	-	-	-	2	-	1	-	-
Female	1	1	-	2	1	-	3	4	2	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	2	-	2	-	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	2	-	-	-	-
Utah												
All races, both sexes	2	1	3	-	2	5	-	2	4	-	-	-
Male	1	1	1	-	-	4	-	1	1	-	-	-
Female	1	-	2	-	2	1	-	1	3	-	-	-
White, both sexes	2	1	3	-	2	5	-	2	4	-	-	-
Male	1	1	1	-	-	4	-	1	1	-	-	-
Female	1	-	2	-	2	1	-	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Vermont												
All races, both sexes	-	-	1	1	4	1	3	-	1	-	-	-
Male	-	-	1	1	2	-	2	-	-	-	-	-
Female	-	-	-	-	2	1	1	-	1	-	-	-
White, both sexes	-	-	1	1	4	1	3	-	1	-	-	-
Male	-	-	1	1	2	-	2	-	-	-	-	-
Female	-	-	-	-	2	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	1	3	3	4	15	12	9	5	2	-	-
Male	2	1	2	2	3	11	11	4	3	-	-	-
Female	1	-	1	1	1	4	1	5	2	2	-	-
White, both sexes	2	1	1	3	2	14	10	9	4	2	-	-
Male	2	1	1	2	1	10	9	4	3	-	-	-
Female	-	-	-	1	1	4	1	5	1	2	-	-
Black, both sexes	1	-	1	-	2	1	2	-	1	-	-	-
Male	-	-	-	-	2	1	2	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Washington												
All races, both sexes	9	9	11	8	8	23	22	26	16	6	1	-
Male	4	6	4	5	5	12	13	16	7	3	-	-
Female	5	3	7	3	3	11	9	10	9	3	1	-
White, both sexes	7	7	9	7	7	22	20	26	15	5	1	-
Male	3	5	4	4	4	12	11	16	6	3	-	-
Female	4	2	5	3	3	10	9	10	9	2	1	-
Black, both sexes	2	1	-	-	-	1	1	-	-	1	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	1	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	1	1	3	1	4	3	3	2	-	-	-
Male	1	1	1	-	1	3	3	1	1	-	-	-
Female	-	-	-	3	-	1	-	2	1	-	-	-
White, both sexes	1	1	1	3	1	4	3	3	1	-	-	-
Male	1	1	1	-	1	3	3	1	1	-	-	-
Female	-	-	-	3	-	1	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Wisconsin												
All races, both sexes	2	4	3	4	1	12	6	10	7	2	-	-
Male	2	3	2	2	-	7	2	3	3	-	-	-
Female	-	1	1	2	1	5	4	7	4	2	-	-
White, both sexes	1	3	3	4	1	10	5	10	7	2	-	-
Male	1	2	2	2	-	5	2	3	3	-	-	-
Female	-	1	1	2	1	5	3	7	4	2	-	-
Black, both sexes	1	1	-	-	-	2	1	-	-	-	-	-
Male	1	1	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	-	1	2	-	1	-	-	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	2	-	1	-	-	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
United States												
All races, both sexes	50,669	144	58	16	14	53	81	120	164	248	452	695
Male	22,777	93	30	7	10	29	49	70	81	135	256	403
Female	27,892	51	28	9	4	24	32	50	83	113	196	292
White, both sexes	44,476	86	41	12	10	42	54	87	114	177	335	514
Male	19,823	56	22	7	8	23	35	49	60	94	193	303
Female	24,653	30	19	5	2	19	19	38	54	83	142	211
Black, both sexes	4,754	54	15	3	4	10	25	28	38	62	106	160
Male	2,200	34	7	-	2	5	14	19	14	36	56	86
Female	2,554	20	8	3	2	5	11	9	24	26	50	74
American Indian 1/, both sexes	230	1	2	1	-	-	2	2	4	5	6	7
Male	116	-	1	-	-	-	-	-	2	3	4	6
Female	114	1	1	1	-	-	2	2	2	2	2	1
Asian or Pacific Islander 2/, both sexes	1,209	3	-	-	-	1	-	3	8	4	5	14
Male	638	3	-	-	-	1	-	2	5	2	3	8
Female	571	-	-	-	-	-	-	1	3	2	2	6
Alabama												
All races, both sexes	844	1	-	-	-	1	1	5	4	5	9	19
Male	388	-	-	-	-	-	-	3	-	5	7	13
Female	456	1	-	-	-	1	1	2	4	-	2	6
White, both sexes	689	1	-	-	-	1	-	3	3	4	5	14
Male	319	-	-	-	-	-	-	1	-	4	3	9
Female	370	1	-	-	-	1	-	2	3	-	2	5
Black, both sexes	153	-	-	-	-	-	1	1	1	1	4	5
Male	68	-	-	-	-	-	-	1	-	1	4	4
Female	85	-	-	-	-	-	1	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Alaska												
All races, both sexes	42	1	1	-	-	-	-	-	2	1	-	-
Male	24	-	-	-	-	-	-	-	1	-	-	-
Female	18	1	1	-	-	-	-	-	1	1	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	1	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	1	1	-	-	-	-	-	2	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	6	1	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	742	9	1	1	-	-	3	2	5	6	9	15
Male	365	5	1	-	-	-	1	2	1	4	4	9
Female	377	4	-	1	-	-	2	-	4	2	5	6
White, both sexes	671	7	-	-	-	-	2	2	5	3	8	11
Male	322	3	-	-	-	-	1	2	1	1	3	6
Female	349	4	-	-	-	-	1	-	4	2	5	5
Black, both sexes	18	2	-	-	-	-	1	-	-	1	-	2
Male	10	2	-	-	-	-	-	-	-	1	-	1
Female	8	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	43	-	1	1	-	-	-	-	-	2	-	2
Male	27	-	1	-	-	-	-	-	-	2	-	2
Female	16	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Arkansas												
All races, both sexes	723	2	1	-	1	3	2	-	3	2	10	14
Male	317	1	1	-	-	2	-	-	-	1	5	9
Female	406	1	-	-	1	1	2	-	3	1	5	5
White, both sexes	650	1	1	-	-	3	-	-	3	2	8	9
Male	282	-	1	-	-	2	-	-	-	1	4	6
Female	368	1	-	-	-	1	-	-	3	1	4	3
Black, both sexes	70	1	-	-	1	-	2	-	-	-	2	5
Male	33	1	-	-	-	-	-	-	-	-	1	3
Female	37	-	-	-	1	-	2	-	-	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	6,516	14	9	2	1	6	6	11	12	21	39	59
Male	3,035	10	6	1	1	4	3	8	8	9	25	38
Female	3,481	4	3	1	-	2	3	3	4	12	14	21
White, both sexes	5,448	13	9	2	1	4	4	8	8	18	33	48
Male	2,504	9	6	1	1	3	2	7	7	8	21	28
Female	2,944	4	3	1	-	1	2	1	1	10	12	20
Black, both sexes	427	1	-	-	-	1	2	2	3	2	6	5
Male	192	1	-	-	-	-	1	1	1	1	4	5
Female	235	-	-	-	-	1	1	1	2	1	2	-
American Indian 1/, both sexes	23	-	-	-	-	-	-	1	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	618	-	-	-	-	1	-	-	1	1	-	6
Male	328	-	-	-	-	1	-	-	-	-	-	5
Female	290	-	-	-	-	-	-	-	1	1	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Colorado												
All races, both sexes	533	1	1	1	-	1	4	2	5	5	5	11
Male	262	1	-	-	-	1	2	1	1	1	4	7
Female	271	-	1	1	-	-	2	1	4	4	1	4
White, both sexes	512	1	1	-	-	1	4	2	4	5	4	9
Male	252	1	-	-	-	1	2	1	1	1	3	6
Female	260	-	1	-	-	-	2	1	3	4	1	3
Black, both sexes	14	-	-	1	-	-	-	-	1	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	1	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	744	2	-	-	-	-	-	1	1	1	3	5
Male	303	1	-	-	-	-	-	-	1	1	2	2
Female	441	1	-	-	-	-	-	1	-	-	1	3
White, both sexes	703	1	-	-	-	-	-	1	-	-	1	5
Male	285	-	-	-	-	-	-	-	-	-	1	2
Female	418	1	-	-	-	-	-	1	-	-	-	3
Black, both sexes	35	1	-	-	-	-	-	-	-	1	2	-
Male	14	1	-	-	-	-	-	-	-	1	1	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Delaware												
All races, both sexes	121	-	-	-	-	-	-	-	2	1	2	1
Male	54	-	-	-	-	-	-	-	2	-	2	1
Female	67	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	108	-	-	-	-	-	-	-	1	-	2	1
Male	48	-	-	-	-	-	-	-	1	-	2	1
Female	60	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	1	1	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	69	1	-	-	-	-	-	-	-	1	1	2
Male	33	1	-	-	-	-	-	-	-	1	-	2
Female	36	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	48	1	-	-	-	-	-	-	-	1	1	2
Male	21	1	-	-	-	-	-	-	-	1	-	2
Female	27	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Florida												
All races, both sexes	2,178	11	3	2	-	1	6	7	7	11	20	40
Male	1,100	9	2	1	-	1	4	2	3	7	12	21
Female	1,078	2	1	1	-	-	2	5	4	4	8	19
White, both sexes	1,946	6	3	2	-	1	2	3	4	10	17	26
Male	993	6	2	1	-	1	2	-	3	7	10	14
Female	953	-	1	1	-	-	-	3	1	3	7	12
Black, both sexes	219	5	-	-	-	-	4	3	3	1	3	13
Male	103	3	-	-	-	-	2	2	-	-	2	7
Female	116	2	-	-	-	-	2	1	3	1	1	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	1	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	1
Georgia												
All races, both sexes	1,360	6	3	-	-	1	6	4	9	12	23	21
Male	582	3	-	-	-	-	4	3	4	10	11	15
Female	778	3	3	-	-	1	2	1	5	2	12	6
White, both sexes	1,073	2	2	-	-	-	3	2	6	6	11	10
Male	456	1	-	-	-	-	2	1	3	6	6	8
Female	617	1	2	-	-	-	1	1	3	-	5	2
Black, both sexes	281	4	1	-	-	1	3	2	3	6	12	11
Male	121	2	-	-	-	-	2	2	1	4	5	7
Female	160	2	1	-	-	1	1	-	2	2	7	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Hawaii												
All races, both sexes	241	-	-	-	-	-	-	1	3	1	1	5
Male	131	-	-	-	-	-	-	1	3	-	-	3
Female	110	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	59	-	-	-	-	-	-	-	-	-	-	3
Male	31	-	-	-	-	-	-	-	-	-	-	3
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	182	-	-	-	-	-	-	1	3	1	1	2
Male	100	-	-	-	-	-	-	1	3	-	-	-
Female	82	-	-	-	-	-	-	-	-	1	1	2
Idaho												
All races, both sexes	207	-	1	-	-	1	-	2	-	1	1	9
Male	106	-	1	-	-	1	-	2	-	1	-	5
Female	101	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	205	-	1	-	-	1	-	2	-	1	1	8
Male	105	-	1	-	-	1	-	2	-	1	-	4
Female	100	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Illinois												
All races, both sexes	2,517	12	1	1	1	4	2	4	11	16	21	35
Male	1,204	8	-	-	1	4	2	2	5	7	14	27
Female	1,313	4	1	1	-	-	-	2	6	9	7	8
White, both sexes	2,180	6	-	-	1	3	1	2	6	10	14	24
Male	1,031	5	-	-	1	3	1	1	3	4	10	19
Female	1,149	1	-	-	-	-	-	1	3	6	4	5
Black, both sexes	311	5	1	1	-	1	1	2	5	6	7	11
Male	157	2	-	-	-	1	1	1	2	3	4	8
Female	154	3	1	1	-	-	-	1	3	3	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	1	-	-	-	-	-	-	-	-	-	-
Male	16	1	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,052	5	3	-	-	3	1	3	5	8	9	19
Male	511	2	2	-	-	1	1	1	4	7	5	13
Female	541	3	1	-	-	2	-	2	1	1	4	6
White, both sexes	990	4	2	-	-	3	1	3	4	5	9	15
Male	476	2	1	-	-	1	1	1	3	5	5	10
Female	514	2	1	-	-	2	-	2	1	-	4	5
Black, both sexes	60	1	1	-	-	-	-	-	1	3	-	4
Male	33	-	1	-	-	-	-	-	1	2	-	3
Female	27	1	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Iowa												
All races, both sexes	724	1	-	-	-	-	1	-	5	1	5	7
Male	301	-	-	-	-	-	1	-	3	1	3	5
Female	423	1	-	-	-	-	-	-	2	-	2	2
White, both sexes	717	1	-	-	-	-	1	-	5	1	5	6
Male	295	-	-	-	-	-	1	-	3	1	3	4
Female	422	1	-	-	-	-	-	-	2	-	2	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	599	1	1	-	-	-	-	2	3	1	4	8
Male	243	1	1	-	-	-	-	2	1	1	-	2
Female	356	-	-	-	-	-	-	-	2	-	4	6
White, both sexes	571	-	-	-	-	-	-	1	3	1	3	8
Male	227	-	-	-	-	-	-	1	1	1	-	2
Female	344	-	-	-	-	-	-	-	2	-	3	6
Black, both sexes	22	1	1	-	-	-	-	1	-	-	1	-
Male	11	1	1	-	-	-	-	1	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Kentucky												
All races, both sexes	839	2	-	-	-	-	6	1	5	5	8	12
Male	379	1	-	-	-	-	5	1	4	5	3	7
Female	460	1	-	-	-	-	1	-	1	-	5	5
White, both sexes	789	1	-	-	-	-	6	1	5	5	7	11
Male	357	1	-	-	-	-	5	1	4	5	2	6
Female	432	-	-	-	-	-	1	-	1	-	5	5
Black, both sexes	49	1	-	-	-	-	-	-	-	-	1	1
Male	22	-	-	-	-	-	-	-	-	-	1	1
Female	27	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	794	9	2	1	1	1	2	2	7	3	15	21
Male	364	2	1	1	1	-	1	1	1	3	6	9
Female	430	7	1	-	-	1	1	1	6	-	9	12
White, both sexes	572	1	1	1	1	1	1	1	3	2	11	15
Male	260	-	1	1	1	-	1	1	-	2	4	8
Female	312	1	-	-	-	1	-	-	3	-	7	7
Black, both sexes	221	8	1	-	-	-	1	1	4	1	4	6
Male	103	2	-	-	-	-	-	-	1	1	2	1
Female	118	6	1	-	-	-	1	1	3	-	2	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Maine												
All races, both sexes	241	-	-	-	-	-	1	-	2	1	2	3
Male	98	-	-	-	-	-	-	-	2	1	1	3
Female	143	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	236	-	-	-	-	-	1	-	1	1	2	3
Male	95	-	-	-	-	-	-	-	1	1	1	3
Female	141	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	992	3	-	1	-	-	2	1	1	5	17	24
Male	438	2	-	-	-	-	2	1	1	4	9	8
Female	554	1	-	1	-	-	-	-	-	1	8	16
White, both sexes	768	2	-	-	-	-	1	-	1	2	8	10
Male	342	1	-	-	-	-	1	-	1	1	3	3
Female	426	1	-	-	-	-	-	-	-	1	5	7
Black, both sexes	197	1	-	1	-	-	1	1	-	3	7	13
Male	83	1	-	-	-	-	1	1	-	3	5	5
Female	114	-	-	1	-	-	-	-	-	-	2	8
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	2	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Massachusetts												
All races, both sexes	1,650	-	-	1	-	1	-	-	2	3	6	15
Male	694	-	-	-	-	-	-	-	1	2	5	10
Female	956	-	-	1	-	1	-	-	1	1	1	5
White, both sexes	1,595	-	-	1	-	1	-	-	1	3	6	13
Male	666	-	-	-	-	-	-	-	-	2	5	8
Female	929	-	-	1	-	1	-	-	1	1	1	5
Black, both sexes	40	-	-	-	-	-	-	-	-	-	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	2
Female	18	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,548	5	2	1	-	1	-	4	7	9	16	14
Male	716	3	2	-	-	1	-	3	2	2	11	8
Female	832	2	-	1	-	-	-	1	5	7	5	6
White, both sexes	1,350	3	1	1	-	-	-	2	6	8	10	9
Male	617	1	1	-	-	-	-	1	2	2	9	6
Female	733	2	-	1	-	-	-	1	4	6	1	3
Black, both sexes	186	2	1	-	-	1	-	2	-	1	6	5
Male	95	2	1	-	-	1	-	2	-	-	2	2
Female	91	-	-	-	-	-	-	-	-	1	4	3
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Minnesota												
All races, both sexes	564	-	1	-	-	-	1	-	-	1	3	5
Male	226	-	1	-	-	-	1	-	-	-	1	3
Female	338	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	557	-	1	-	-	-	1	-	-	1	3	4
Male	222	-	1	-	-	-	1	-	-	-	1	3
Female	335	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	571	1	1	1	-	2	1	4	2	3	4	9
Male	234	1	1	1	-	-	-	2	2	-	2	3
Female	337	-	-	-	-	2	1	2	-	3	2	6
White, both sexes	388	1	1	1	-	2	-	-	2	1	2	5
Male	155	1	1	1	-	-	-	-	2	-	1	1
Female	233	-	-	-	-	2	-	-	-	1	1	4
Black, both sexes	182	-	-	-	-	-	1	4	-	2	2	4
Male	78	-	-	-	-	-	-	2	-	-	1	2
Female	104	-	-	-	-	-	1	2	-	2	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Missouri												
All races, both sexes	1,195	4	1	-	1	2	1	4	5	2	11	17
Male	501	3	-	-	1	1	1	2	2	1	3	11
Female	694	1	1	-	-	1	-	2	3	1	8	6
White, both sexes	1,110	2	-	-	1	2	1	4	2	1	8	14
Male	468	2	-	-	1	1	1	2	1	-	2	10
Female	642	-	-	-	-	1	-	2	1	1	6	4
Black, both sexes	80	2	1	-	-	-	-	-	3	1	3	3
Male	33	1	-	-	-	-	-	-	1	1	1	1
Female	47	1	1	-	-	-	-	-	2	-	2	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	155	-	-	-	-	-	1	1	-	1	1	2
Male	75	-	-	-	-	-	1	-	-	1	1	2
Female	80	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	143	-	-	-	-	-	1	-	-	-	1	2
Male	68	-	-	-	-	-	1	-	-	-	1	2
Female	75	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	1	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Nebraska												
All races, both sexes	315	1	-	-	-	2	1	2	-	3	5	2
Male	129	-	-	-	-	-	-	1	-	-	2	1
Female	186	1	-	-	-	2	1	1	-	3	3	1
White, both sexes	309	1	-	-	-	2	1	2	-	3	5	2
Male	127	-	-	-	-	-	-	1	-	-	2	1
Female	182	1	-	-	-	2	1	1	-	3	3	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	403	1	1	-	-	-	1	3	2	3	3	11
Male	206	1	-	-	-	-	1	3	1	-	3	6
Female	197	-	1	-	-	-	-	-	1	3	-	5
White, both sexes	355	1	1	-	-	-	1	1	2	1	2	9
Male	179	1	-	-	-	-	1	1	1	-	2	5
Female	176	-	1	-	-	-	-	-	1	1	-	4
Black, both sexes	32	-	-	-	-	-	-	2	-	2	1	-
Male	22	-	-	-	-	-	-	2	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
New Hampshire												
All races, both sexes	180	-	-	-	-	-	-	-	-	-	1	-
Male	90	-	-	-	-	-	-	-	-	-	-	-
Female	90	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	177	-	-	-	-	-	-	-	-	-	1	-
Male	89	-	-	-	-	-	-	-	-	-	-	-
Female	88	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,276	2	1	-	2	-	2	7	6	9	9	14
Male	584	1	-	-	2	-	1	5	5	3	7	9
Female	692	1	1	-	-	-	1	2	1	6	2	5
White, both sexes	1,143	2	-	-	-	-	2	5	6	6	6	10
Male	521	1	-	-	-	-	1	4	5	-	5	8
Female	622	1	-	-	-	-	1	1	1	6	1	2
Black, both sexes	115	-	1	-	2	-	-	2	-	3	3	4
Male	48	-	-	-	2	-	-	1	-	3	2	1
Female	67	-	1	-	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
New Mexico												
All races, both sexes	317	-	-	-	-	1	-	1	2	2	5	13
Male	153	-	-	-	-	1	-	-	2	2	4	8
Female	164	-	-	-	-	-	-	1	-	-	1	5
White, both sexes	291	-	-	-	-	1	-	1	1	2	4	10
Male	140	-	-	-	-	1	-	-	1	2	3	6
Female	151	-	-	-	-	-	-	1	-	-	1	4
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	1	-	1	2
Male	12	-	-	-	-	-	-	-	1	-	1	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4,215	7	6	-	1	1	4	4	7	15	34	40
Male	1,852	5	4	-	-	1	1	2	4	9	20	22
Female	2,363	2	2	-	1	-	3	2	3	6	14	18
White, both sexes	3,535	4	4	-	1	-	4	4	4	9	24	28
Male	1,542	3	3	-	-	-	1	2	3	8	16	16
Female	1,993	1	1	-	1	-	3	2	1	1	8	12
Black, both sexes	539	3	2	-	-	1	-	-	2	5	9	10
Male	237	2	1	-	-	1	-	-	1	-	3	5
Female	302	1	1	-	-	-	-	-	1	5	6	5
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	138	-	-	-	-	-	-	-	1	1	1	2
Male	73	-	-	-	-	-	-	-	-	1	1	1
Female	65	-	-	-	-	-	-	-	1	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
North Carolina												
All races, both sexes	1,579	4	3	1	-	2	5	3	6	11	23	32
Male	643	4	2	-	-	1	3	2	3	6	12	18
Female	936	-	1	1	-	1	2	1	3	5	11	14
White, both sexes	1,331	2	2	1	-	1	1	1	5	6	15	20
Male	529	2	1	-	-	-	1	1	3	2	8	11
Female	802	-	1	1	-	1	-	-	2	4	7	9
Black, both sexes	238	2	1	-	-	1	3	2	1	5	8	11
Male	109	2	1	-	-	1	2	1	-	4	4	6
Female	129	-	-	-	-	-	1	1	1	1	4	5
American Indian 1/, both sexes	7	-	-	-	-	-	1	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	119	-	-	-	-	-	-	-	-	-	-	-
Male	50	-	-	-	-	-	-	-	-	-	-	-
Female	69	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	118	-	-	-	-	-	-	-	-	-	-	-
Male	49	-	-	-	-	-	-	-	-	-	-	-
Female	69	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Ohio												
All races, both sexes	1,655	7	5	1	1	1	2	5	2	11	16	19
Male	714	6	2	1	1	-	1	4	1	6	12	11
Female	941	1	3	-	-	1	1	1	1	5	4	8
White, both sexes	1,499	3	4	1	1	1	2	4	1	6	13	17
Male	640	3	1	1	1	-	1	3	1	3	10	10
Female	859	-	3	-	-	1	1	1	-	3	3	7
Black, both sexes	153	4	1	-	-	-	-	1	1	5	3	2
Male	72	3	1	-	-	-	-	1	-	3	2	1
Female	81	1	-	-	-	-	-	-	1	2	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	794	2	-	-	-	1	2	2	-	11	8	13
Male	358	1	-	-	-	-	1	-	-	4	3	8
Female	436	1	-	-	-	1	1	2	-	7	5	5
White, both sexes	724	2	-	-	-	1	1	2	-	9	5	10
Male	329	1	-	-	-	-	1	-	-	4	1	6
Female	395	1	-	-	-	1	-	2	-	5	4	4
Black, both sexes	38	-	-	-	-	-	-	-	-	-	-	3
Male	15	-	-	-	-	-	-	-	-	-	-	2
Female	23	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	32	-	-	-	-	-	1	-	-	2	3	-
Male	14	-	-	-	-	-	-	-	-	-	2	-
Female	18	-	-	-	-	-	1	-	-	2	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Oregon												
All races, both sexes	449	1	-	-	-	-	-	1	3	1	4	4
Male	223	1	-	-	-	-	-	1	1	1	2	3
Female	226	-	-	-	-	-	-	-	2	-	2	1
White, both sexes	443	-	-	-	-	-	-	1	3	1	4	4
Male	221	-	-	-	-	-	-	1	1	1	2	3
Female	222	-	-	-	-	-	-	-	2	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2,590	6	1	-	2	1	5	5	3	9	14	32
Male	1,135	5	-	-	1	1	3	2	1	3	9	18
Female	1,455	1	1	-	1	-	2	3	2	6	5	14
White, both sexes	2,390	4	1	-	2	1	4	5	3	8	10	21
Male	1,039	3	-	-	1	1	2	2	1	2	9	12
Female	1,351	1	1	-	1	-	2	3	2	6	1	9
Black, both sexes	190	2	-	-	-	-	1	-	-	1	4	11
Male	93	2	-	-	-	-	1	-	-	1	-	6
Female	97	-	-	-	-	-	-	-	-	-	4	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Rhode Island												
All races, both sexes	249	-	-	-	-	-	-	-	-	1	4	-
Male	101	-	-	-	-	-	-	-	-	1	2	-
Female	148	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	239	-	-	-	-	-	-	-	-	1	4	-
Male	95	-	-	-	-	-	-	-	-	1	2	-
Female	144	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	640	1	-	-	-	-	2	1	4	3	5	12
Male	296	-	-	-	-	-	2	-	3	2	3	6
Female	344	1	-	-	-	-	-	1	1	1	2	6
White, both sexes	490	-	-	-	-	-	1	1	1	3	2	10
Male	215	-	-	-	-	-	1	-	1	2	1	5
Female	275	-	-	-	-	-	-	1	-	1	1	5
Black, both sexes	148	1	-	-	-	-	1	-	3	-	3	2
Male	81	-	-	-	-	-	1	-	2	-	2	1
Female	67	1	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
South Dakota												
All races, both sexes	151	-	-	-	-	-	-	-	1	-	2	1
Male	63	-	-	-	-	-	-	-	-	-	1	-
Female	88	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	143	-	-	-	-	-	-	-	-	-	2	1
Male	58	-	-	-	-	-	-	-	-	-	1	-
Female	85	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,365	6	1	-	1	7	3	4	2	6	13	21
Male	571	5	1	-	1	4	1	3	1	4	8	10
Female	794	1	-	-	-	3	2	1	1	2	5	11
White, both sexes	1,205	4	-	-	1	5	3	4	1	5	8	15
Male	492	3	-	-	1	4	1	3	1	3	6	7
Female	713	1	-	-	-	1	2	1	-	2	2	8
Black, both sexes	158	2	1	-	-	2	-	-	1	1	5	6
Male	77	2	1	-	-	-	-	-	-	1	2	3
Female	81	-	-	-	-	2	-	-	1	-	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Texas												
All races, both sexes	2,991	8	4	2	1	3	5	13	8	18	29	53
Male	1,296	4	1	2	-	3	4	6	3	11	18	26
Female	1,695	4	3	-	1	-	1	7	5	7	11	27
White, both sexes	2,654	6	2	2	-	3	3	12	6	10	23	44
Male	1,144	3	1	2	-	3	2	6	2	6	13	23
Female	1,510	3	1	-	-	-	1	6	4	4	10	21
Black, both sexes	295	1	2	-	1	-	2	1	2	8	6	9
Male	135	-	-	-	-	-	2	-	1	5	5	3
Female	160	1	2	-	1	-	-	1	1	3	1	6
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	38	1	-	-	-	-	-	-	-	-	-	-
Male	16	1	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	307	-	1	-	-	2	-	2	-	3	2	4
Male	131	-	-	-	-	-	-	1	-	1	-	2
Female	176	-	1	-	-	2	-	1	-	2	2	2
White, both sexes	296	-	1	-	-	2	-	2	-	2	2	4
Male	126	-	-	-	-	-	-	1	-	-	-	2
Female	170	-	1	-	-	2	-	1	-	2	2	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Vermont												
All races, both sexes	56	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	56	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,182	3	1	-	-	2	2	3	4	5	11	13
Male	537	3	1	-	-	1	2	2	2	2	5	8
Female	645	-	-	-	-	1	-	1	2	3	6	5
White, both sexes	1,004	1	1	-	-	1	1	2	2	4	8	9
Male	458	1	1	-	-	1	1	1	1	2	3	6
Female	546	-	-	-	-	-	-	1	1	2	5	3
Black, both sexes	159	2	-	-	-	1	1	1	2	1	3	4
Male	69	2	-	-	-	-	1	1	1	-	2	2
Female	90	-	-	-	-	1	-	-	1	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Washington												
All races, both sexes	628	-	2	-	1	1	-	2	1	7	8	3
Male	275	-	-	-	1	1	-	-	-	4	4	-
Female	353	-	2	-	-	-	-	2	1	3	4	3
White, both sexes	590	-	2	-	1	-	-	2	1	7	7	3
Male	260	-	-	-	1	-	-	-	-	4	4	-
Female	330	-	2	-	-	-	-	2	1	3	3	3
Black, both sexes	8	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	1	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	418	1	-	-	-	-	-	1	1	2	4	5
Male	196	1	-	-	-	-	-	1	-	-	2	3
Female	222	-	-	-	-	-	-	-	1	2	2	2
White, both sexes	410	1	-	-	-	-	-	1	1	2	4	5
Male	191	1	-	-	-	-	-	1	-	-	2	3
Female	219	-	-	-	-	-	-	-	1	2	2	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Wisconsin												
All races, both sexes	922	3	-	-	-	1	-	-	3	1	7	7
Male	423	2	-	-	-	-	-	-	2	1	3	5
Female	499	1	-	-	-	1	-	-	1	-	4	2
White, both sexes	893	2	-	-	-	1	-	-	3	1	7	6
Male	411	1	-	-	-	-	-	-	2	1	3	5
Female	482	1	-	-	-	1	-	-	1	-	4	1
Black, both sexes	21	1	-	-	-	-	-	-	-	-	-	1
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	107	-	-	-	-	-	-	-	1	-	-	4
Male	44	-	-	-	-	-	-	-	-	-	-	3
Female	63	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	105	-	-	-	-	-	-	-	1	-	-	3
Male	42	-	-	-	-	-	-	-	-	-	-	2
Female	63	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
United States												
All races, both sexes	955	1,170	1,496	2,073	3,222	5,644	9,336	10,756	8,871	4,072	1,026	3
Male	583	684	889	1,171	1,753	2,969	4,443	4,629	3,162	1,118	211	2
Female	372	486	607	902	1,469	2,675	4,893	6,127	5,709	2,954	815	1
White, both sexes	711	919	1,218	1,658	2,747	4,927	8,320	9,796	8,103	3,695	907	3
Male	428	548	717	934	1,491	2,602	3,978	4,218	2,873	999	183	2
Female	283	371	501	724	1,256	2,325	4,342	5,578	5,230	2,696	724	1
Black, both sexes	214	226	234	350	375	538	757	656	540	264	95	-
Male	138	123	147	192	204	279	324	252	175	72	21	-
Female	76	103	87	158	171	259	433	404	365	192	74	-
American Indian 1/, both sexes	12	7	14	15	22	25	35	34	18	12	6	-
Male	5	3	6	10	14	10	22	20	7	3	-	-
Female	7	4	8	5	8	15	13	14	11	9	6	-
Asian or Pacific Islander 2/, both sexes	18	18	30	50	78	154	224	270	210	101	18	-
Male	12	10	19	35	44	78	119	139	107	44	7	-
Female	6	8	11	15	34	76	105	131	103	57	11	-
Alabama												
All races, both sexes	22	31	28	59	60	118	156	155	109	48	13	-
Male	19	17	17	37	31	61	75	61	31	9	2	-
Female	3	14	11	22	29	57	81	94	78	39	11	-
White, both sexes	12	24	22	48	49	97	130	136	89	41	10	-
Male	11	14	14	29	25	50	67	58	25	7	2	-
Female	1	10	8	19	24	47	63	78	64	34	8	-
Black, both sexes	10	7	6	11	11	21	25	19	20	7	3	-
Male	8	3	3	8	6	11	8	3	6	2	-	-
Female	2	4	3	3	5	10	17	16	14	5	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Alaska												
All races, both sexes	4	1	2	1	1	3	8	5	4	8	-	-
Male	2	1	1	-	1	3	5	3	2	5	-	-
Female	2	-	1	1	-	-	3	2	2	3	-	-
White, both sexes	2	1	1	-	-	2	5	5	3	6	-	-
Male	1	1	1	-	-	2	2	3	1	4	-	-
Female	1	-	-	-	-	-	3	2	2	2	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	1	1	3	-	1	2	-	-
Male	1	-	-	-	1	1	3	-	1	1	-	-
Female	-	-	1	1	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	21	20	28	43	63	92	119	140	109	41	15	-
Male	15	13	18	29	35	46	64	65	36	15	2	-
Female	6	7	10	14	28	46	55	75	73	26	13	-
White, both sexes	19	17	23	39	56	84	108	131	106	37	13	-
Male	15	12	15	26	31	42	56	58	35	13	2	-
Female	4	5	8	13	25	42	52	73	71	24	11	-
Black, both sexes	1	1	2	1	1	2	2	-	1	1	-	-
Male	-	-	2	-	1	1	2	-	-	-	-	-
Female	1	1	-	1	-	1	-	-	1	1	-	-
American Indian 1/, both sexes	1	2	3	3	4	5	7	7	2	1	2	-
Male	-	1	1	3	2	3	5	5	1	1	-	-
Female	1	1	2	-	2	2	2	2	1	-	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	2	2	-	2	-	-
Male	-	-	-	-	1	-	1	2	-	1	-	-
Female	-	-	-	-	1	1	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Arkansas												
All races, both sexes	19	17	29	29	42	74	138	138	128	55	16	-
Male	7	11	16	13	19	39	69	65	37	20	2	-
Female	12	6	13	16	23	35	69	73	91	35	14	-
White, both sexes	15	14	27	26	39	68	127	130	118	48	11	-
Male	5	9	15	12	18	35	63	59	34	17	1	-
Female	10	5	12	14	21	33	64	71	84	31	10	-
Black, both sexes	4	3	2	3	3	5	10	8	9	7	5	-
Male	2	2	1	1	1	3	5	6	3	3	1	-
Female	2	1	1	2	2	2	5	2	6	4	4	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
California												
All races, both sexes	71	97	190	262	399	689	1,268	1,465	1,209	564	122	-
Male	45	54	113	154	209	376	608	693	476	167	27	-
Female	26	43	77	108	190	313	660	772	733	397	95	-
White, both sexes	58	82	155	202	313	568	1,062	1,221	1,045	482	112	-
Male	36	48	94	119	155	316	504	574	403	137	25	-
Female	22	34	61	83	158	252	558	647	642	345	87	-
Black, both sexes	8	9	22	33	47	43	75	85	49	29	5	-
Male	5	3	11	16	28	22	29	38	15	11	-	-
Female	3	6	11	17	19	21	46	47	34	18	5	-
American Indian 1/, both sexes	-	1	-	2	2	5	4	5	1	2	-	-
Male	-	1	-	1	2	-	3	3	1	-	-	-
Female	-	-	-	1	-	5	1	2	-	2	-	-
Asian or Pacific Islander 2/, both sexes	5	5	13	25	37	73	127	154	114	51	5	-
Male	4	2	8	18	24	38	72	78	57	19	2	-
Female	1	3	5	7	13	35	55	76	57	32	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Colorado												
All races, both sexes	13	14	12	15	32	53	102	112	85	49	10	-
Male	7	9	5	11	16	30	65	53	32	14	2	-
Female	6	5	7	4	16	23	37	59	53	35	8	-
White, both sexes	12	14	12	12	31	50	97	111	85	47	10	-
Male	7	9	5	9	16	28	62	52	32	14	2	-
Female	5	5	7	3	15	22	35	59	53	33	8	-
Black, both sexes	1	-	-	3	1	1	2	-	-	2	-	-
Male	-	-	-	2	-	1	1	-	-	-	-	-
Female	1	-	-	1	1	-	1	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	3	1	-	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Connecticut												
All races, both sexes	5	9	24	21	28	80	141	186	138	81	18	-
Male	3	6	17	12	17	40	59	74	42	23	3	-
Female	2	3	7	9	11	40	82	112	96	58	15	-
White, both sexes	5	8	20	19	27	77	132	182	132	76	17	-
Male	3	5	14	11	16	39	56	72	41	22	3	-
Female	2	3	6	8	11	38	76	110	91	54	14	-
Black, both sexes	-	1	4	2	1	3	6	3	5	5	1	-
Male	-	1	3	1	1	1	2	1	-	1	-	-
Female	-	-	1	1	-	2	4	2	5	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	1	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Delaware												
All races, both sexes	2	2	-	6	10	19	25	25	18	7	1	-
Male	1	2	-	3	3	9	10	14	6	1	-	-
Female	1	-	-	3	7	10	15	11	12	6	1	-
White, both sexes	2	2	-	4	9	15	24	23	17	7	1	-
Male	1	2	-	3	3	7	9	12	6	1	-	-
Female	1	-	-	1	6	8	15	11	11	6	1	-
Black, both sexes	-	-	-	2	1	3	1	2	1	-	-	-
Male	-	-	-	-	-	2	1	2	-	-	-	-
Female	-	-	-	2	1	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	3	1	5	4	5	16	13	9	5	2	-
Male	1	2	1	3	2	2	4	8	5	-	1	-
Female	-	1	-	2	2	3	12	5	4	5	1	-
White, both sexes	-	-	1	1	1	2	5	6	-	3	1	-
Male	-	-	1	1	1	1	2	6	-	-	-	-
Female	-	-	-	-	-	1	3	-	-	3	1	-
Black, both sexes	1	3	-	4	3	3	11	7	9	1	1	-
Male	1	2	-	2	1	1	2	2	5	-	1	-
Female	-	1	-	2	2	2	9	5	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Florida												
All races, both sexes	48	51	81	124	148	258	417	442	337	129	34	1
Male	34	34	50	77	89	149	212	211	137	38	6	1
Female	14	17	31	47	59	109	205	231	200	91	28	-
White, both sexes	41	41	64	94	129	233	397	415	309	117	31	1
Male	29	28	40	63	78	137	202	202	127	34	6	1
Female	12	13	24	31	51	96	195	213	182	83	25	-
Black, both sexes	6	10	17	30	18	23	19	25	25	11	3	-
Male	5	6	10	14	11	11	10	8	8	4	-	-
Female	1	4	7	16	7	12	9	17	17	7	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	2	2	1	-	-
Male	-	-	-	-	-	1	-	1	2	-	-	-
Female	-	-	-	-	1	1	1	1	-	1	-	-
Georgia												
All races, both sexes	35	46	51	61	88	171	264	265	188	85	21	-
Male	20	26	32	41	43	86	120	96	52	14	2	-
Female	15	20	19	20	45	85	144	169	136	71	19	-
White, both sexes	21	33	32	43	68	135	225	232	158	70	14	-
Male	12	20	19	33	34	70	102	84	43	11	1	-
Female	9	13	13	10	34	65	123	148	115	59	13	-
Black, both sexes	13	13	17	18	20	36	38	32	29	15	7	-
Male	7	6	11	8	9	16	17	11	9	3	1	-
Female	6	7	6	10	11	20	21	21	20	12	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	-	-	-	1	1	1	-	-	-
Male	1	-	2	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Hawaii												
All races, both sexes	9	5	3	5	13	26	30	58	48	26	7	-
Male	5	2	-	4	6	14	18	36	22	14	3	-
Female	4	3	3	1	7	12	12	22	26	12	4	-
White, both sexes	3	1	-	-	6	6	11	9	14	5	1	-
Male	2	-	-	-	4	4	5	4	7	2	-	-
Female	1	1	-	-	2	2	6	5	7	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	4	3	5	7	20	19	49	34	21	6	-
Male	3	2	-	4	2	10	13	32	15	12	3	-
Female	3	2	3	1	5	10	6	17	19	9	3	-
Idaho												
All races, both sexes	6	2	5	5	10	20	36	44	34	19	11	-
Male	3	1	2	2	4	10	21	23	18	9	3	-
Female	3	1	3	3	6	10	15	21	16	10	8	-
White, both sexes	6	2	5	5	10	20	36	44	33	19	11	-
Male	3	1	2	2	4	10	21	23	18	9	3	-
Female	3	1	3	3	6	10	15	21	15	10	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Illinois												
All races, both sexes	41	58	84	109	160	268	481	507	443	199	59	-
Male	29	29	56	69	84	153	244	240	153	65	12	-
Female	12	29	28	40	76	115	237	267	290	134	47	-
White, both sexes	31	39	69	77	129	232	413	468	414	190	51	-
Male	26	22	43	48	70	136	206	223	139	64	7	-
Female	5	17	26	29	59	96	207	245	275	126	44	-
Black, both sexes	10	19	15	30	29	36	61	32	25	8	6	-
Male	3	7	13	19	13	17	33	14	11	1	4	-
Female	7	12	2	11	16	19	28	18	14	7	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	2	-	7	7	4	1	2	-
Male	-	-	-	2	1	-	5	3	3	-	1	-
Female	-	-	-	-	1	-	2	4	1	1	1	-
Indiana												
All races, both sexes	24	24	30	47	70	121	202	220	179	61	18	-
Male	17	16	18	27	40	61	93	113	65	20	5	-
Female	7	8	12	20	30	60	109	107	114	41	13	-
White, both sexes	22	21	29	45	66	114	187	212	174	57	17	-
Male	15	13	17	25	38	58	87	109	62	18	5	-
Female	7	8	12	20	28	56	100	103	112	39	12	-
Black, both sexes	2	3	1	2	3	7	15	7	5	4	1	-
Male	2	3	1	2	1	3	6	3	3	2	-	-
Female	-	-	-	-	2	4	9	4	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Iowa												
All races, both sexes	3	12	9	25	35	48	135	146	160	108	23	-
Male	3	8	7	15	21	27	74	55	45	30	3	-
Female	-	4	2	10	14	21	61	91	115	78	20	-
White, both sexes	3	12	8	24	34	48	133	146	159	108	23	-
Male	3	8	6	14	20	27	72	55	45	30	3	-
Female	-	4	2	10	14	21	61	91	114	78	20	-
Black, both sexes	-	-	1	1	1	-	2	-	-	-	-	-
Male	-	-	1	1	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Kansas												
All races, both sexes	11	12	8	13	25	55	101	130	129	77	18	-
Male	6	8	5	8	17	24	45	51	45	21	5	-
Female	5	4	3	5	8	31	56	79	84	56	13	-
White, both sexes	9	11	8	12	24	54	96	125	126	73	17	-
Male	5	8	5	7	16	23	42	49	43	20	4	-
Female	4	3	3	5	8	31	54	76	83	53	13	-
Black, both sexes	-	1	-	-	1	1	5	3	2	4	1	-
Male	-	-	-	-	1	1	3	-	1	1	1	-
Female	-	1	-	-	-	-	2	3	1	3	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	2	-	-	-	-
Male	1	-	-	-	-	-	-	2	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Kentucky												
All races, both sexes	20	26	25	40	63	100	157	155	137	63	14	-
Male	8	13	18	24	37	47	77	64	42	19	4	-
Female	12	13	7	16	26	53	80	91	95	44	10	-
White, both sexes	19	26	22	37	59	95	149	152	120	60	14	-
Male	8	13	16	22	34	44	73	63	37	19	4	-
Female	11	13	6	15	25	51	76	89	83	41	10	-
Black, both sexes	1	-	3	3	4	5	8	3	16	3	-	-
Male	-	-	2	2	3	3	4	1	5	-	-	-
Female	1	-	1	1	1	2	4	2	11	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Louisiana												
All races, both sexes	18	27	28	46	57	102	166	119	113	42	11	1
Male	12	15	15	22	34	49	81	51	44	11	3	1
Female	6	12	13	24	23	53	85	68	69	31	8	-
White, both sexes	11	18	17	34	39	65	133	92	89	29	6	1
Male	6	8	9	14	23	29	65	39	39	6	2	1
Female	5	10	8	20	16	36	68	53	50	23	4	-
Black, both sexes	7	9	11	12	18	37	32	27	24	13	5	-
Male	6	7	6	8	11	20	15	12	5	5	1	-
Female	1	2	5	4	7	17	17	15	19	8	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Maine												
All races, both sexes	3	4	2	5	23	19	49	37	59	26	5	-
Male	1	2	1	1	8	12	25	16	21	4	-	-
Female	2	2	1	4	15	7	24	21	38	22	5	-
White, both sexes	3	4	2	5	23	16	49	37	58	26	5	-
Male	1	2	1	1	8	10	25	16	21	4	-	-
Female	2	2	1	4	15	6	24	21	37	22	5	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Maryland												
All races, both sexes	28	30	25	35	61	95	173	219	181	71	20	-
Male	18	23	10	19	40	50	74	83	74	17	3	-
Female	10	7	15	16	21	45	99	136	107	54	17	-
White, both sexes	13	23	16	22	43	74	143	179	151	64	16	-
Male	9	18	7	13	29	41	64	69	64	17	1	-
Female	4	5	9	9	14	33	79	110	87	47	15	-
Black, both sexes	15	7	9	11	14	19	25	37	24	7	2	-
Male	9	5	3	4	8	7	8	13	9	-	1	-
Female	6	2	6	7	6	12	17	24	15	7	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	3	2	5	3	5	-	2	-
Male	-	-	-	2	3	2	2	1	1	-	1	-
Female	-	-	-	-	-	-	3	2	4	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Massachusetts												
All races, both sexes	18	27	45	43	93	175	320	390	311	158	42	-
Male	10	17	27	18	57	91	143	142	117	45	9	-
Female	8	10	18	25	36	84	177	248	194	113	33	-
White, both sexes	17	25	43	41	89	167	309	378	304	155	42	-
Male	9	15	26	18	53	87	138	139	114	43	9	-
Female	8	10	17	23	36	80	171	239	190	112	33	-
Black, both sexes	1	2	2	1	2	5	9	9	4	3	-	-
Male	1	2	1	-	2	3	5	3	1	2	-	-
Female	-	-	1	1	-	2	4	6	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	3	2	3	3	-	-	-
Male	-	-	-	-	2	1	-	-	2	-	-	-
Female	-	-	-	1	-	2	2	3	1	-	-	-
Michigan												
All races, both sexes	34	38	55	56	95	195	286	330	261	112	27	-
Male	22	24	32	35	50	112	129	141	105	28	6	-
Female	12	14	23	21	45	83	157	189	156	84	21	-
White, both sexes	20	28	45	46	82	174	246	295	242	108	24	-
Male	12	18	27	26	44	101	111	126	98	26	6	-
Female	8	10	18	20	38	73	135	169	144	82	18	-
Black, both sexes	12	10	9	8	13	19	39	33	18	4	3	-
Male	8	6	5	7	6	11	18	15	7	2	-	-
Female	4	4	4	1	7	8	21	18	11	2	3	-
American Indian 1/, both sexes	1	-	-	2	-	1	-	1	1	-	-	-
Male	1	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	1	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Minnesota												
All races, both sexes	6	8	10	11	25	43	94	133	139	74	10	-
Male	3	6	6	5	16	17	45	51	57	13	1	-
Female	3	2	4	6	9	26	49	82	82	61	9	-
White, both sexes	6	8	9	10	25	42	92	133	138	74	10	-
Male	3	6	5	4	16	17	44	51	56	13	1	-
Female	3	2	4	6	9	25	48	82	82	61	9	-
Black, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	19	18	19	25	51	78	102	95	76	47	13	-
Male	11	11	11	9	32	35	49	30	19	14	1	-
Female	8	7	8	16	19	43	53	65	57	33	12	-
White, both sexes	10	9	13	9	39	50	79	73	57	28	6	-
Male	4	5	6	3	23	23	40	21	13	9	1	-
Female	6	4	7	6	16	27	39	52	44	19	5	-
Black, both sexes	9	8	6	16	12	28	23	22	19	19	7	-
Male	7	5	5	6	9	12	9	9	6	5	-	-
Female	2	3	1	10	3	16	14	13	13	14	7	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Missouri												
All races, both sexes	22	23	28	42	66	147	207	262	214	103	32	1
Male	16	14	20	19	38	72	104	96	67	23	7	-
Female	6	9	8	23	28	75	103	166	147	80	25	1
White, both sexes	19	20	23	36	61	138	194	251	204	98	30	1
Male	14	12	17	17	37	68	99	95	61	22	6	-
Female	5	8	6	19	24	70	95	156	143	76	24	1
Black, both sexes	3	2	4	6	4	8	13	11	9	5	2	-
Male	2	2	3	2	1	4	5	1	6	1	1	-
Female	1	-	1	4	3	4	8	10	3	4	1	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	2	5	4	9	13	24	37	33	15	3	-
Male	3	2	4	2	4	10	11	19	11	4	-	-
Female	1	-	1	2	5	3	13	18	22	11	3	-
White, both sexes	3	1	4	4	9	13	21	34	33	14	3	-
Male	2	1	3	2	4	10	9	18	11	4	-	-
Female	1	-	1	2	5	3	12	16	22	10	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	1	-	-	-	3	2	-	1	-	-
Male	1	1	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Nebraska												
All races, both sexes	5	4	8	10	12	34	38	64	78	33	13	-
Male	2	3	6	5	7	18	17	26	29	10	2	-
Female	3	1	2	5	5	16	21	38	49	23	11	-
White, both sexes	4	4	7	10	12	33	36	64	77	33	13	-
Male	1	3	6	5	7	17	17	26	29	10	2	-
Female	3	1	1	5	5	16	19	38	48	23	11	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	12	15	23	33	51	68	69	58	35	12	2	-
Male	7	11	12	19	25	41	30	26	14	5	1	-
Female	5	4	11	14	26	27	39	32	21	7	1	-
White, both sexes	8	13	19	33	43	58	63	53	33	12	2	-
Male	3	9	10	19	21	33	28	25	14	5	1	-
Female	5	4	9	14	22	25	35	28	19	7	1	-
Black, both sexes	4	2	3	-	5	6	4	2	1	-	-	-
Male	4	2	2	-	3	6	1	1	-	-	-	-
Female	-	-	1	-	2	-	3	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	4	2	3	-	-	-	-
Male	-	-	-	-	1	2	1	-	-	-	-	-
Female	-	-	1	-	1	2	1	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
New Hampshire												
All races, both sexes	2	2	3	4	7	13	28	52	47	19	2	-
Male	2	1	1	3	2	10	17	29	19	5	1	-
Female	-	1	2	1	5	3	11	23	28	14	1	-
White, both sexes	2	2	3	4	7	13	28	51	46	18	2	-
Male	2	1	1	3	2	10	17	29	18	5	1	-
Female	-	1	2	1	5	3	11	22	28	13	1	-
Black, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
New Jersey												
All races, both sexes	14	30	29	42	94	144	224	273	242	107	25	-
Male	9	21	20	23	61	62	106	117	96	28	8	-
Female	5	9	9	19	33	82	118	156	146	79	17	-
White, both sexes	10	23	24	35	87	126	201	259	221	97	23	-
Male	6	16	17	20	57	54	97	110	89	23	8	-
Female	4	7	7	15	30	72	104	149	132	74	15	-
Black, both sexes	4	7	3	7	7	14	20	13	15	8	2	-
Male	3	5	1	3	4	5	7	6	2	3	-	-
Female	1	2	2	4	3	9	13	7	13	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	3	3	1	6	2	-	-
Male	-	-	2	-	-	2	2	1	5	2	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
New Mexico												
All races, both sexes	7	6	11	9	23	36	58	53	56	24	10	-
Male	5	4	4	5	13	16	29	25	27	6	2	-
Female	2	2	7	4	10	20	29	28	29	18	8	-
White, both sexes	7	5	9	8	22	34	55	48	52	23	9	-
Male	5	4	4	4	12	16	27	22	25	6	2	-
Female	2	1	5	4	10	18	28	26	27	17	7	-
Black, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	2	1	1	2	3	4	3	1	1	-
Male	-	-	-	1	1	-	2	3	1	-	-	-
Female	-	1	2	-	-	2	1	1	2	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	89	85	128	159	273	470	762	922	770	355	83	-
Male	49	47	70	90	136	262	347	381	281	100	21	-
Female	40	38	58	69	137	208	415	541	489	255	62	-
White, both sexes	63	55	101	117	205	390	653	800	681	314	74	-
Male	33	32	51	66	105	213	303	330	248	90	19	-
Female	30	23	50	51	100	177	350	470	433	224	55	-
Black, both sexes	23	25	22	36	56	63	84	97	66	27	8	-
Male	15	11	14	20	26	39	34	40	19	4	2	-
Female	8	14	8	16	30	24	50	57	47	23	6	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	5	6	12	16	25	24	22	14	1	-
Male	1	4	5	4	5	10	10	11	14	6	-	-
Female	2	1	-	2	7	6	15	13	8	8	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
North Carolina												
All races, both sexes	29	45	43	69	98	189	267	317	273	114	45	-
Male	16	24	31	32	52	88	122	118	76	28	5	-
Female	13	21	12	37	46	101	145	199	197	86	40	-
White, both sexes	18	33	35	51	81	166	236	285	238	96	38	-
Male	7	17	25	22	44	79	108	106	65	23	4	-
Female	11	16	10	29	37	87	128	179	173	73	34	-
Black, both sexes	10	12	8	17	17	22	30	29	34	18	7	-
Male	8	7	6	10	8	9	14	10	10	5	1	-
Female	2	5	2	7	9	13	16	19	24	13	6	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	3	1	-	-	-
Male	1	-	-	-	-	-	-	2	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	2	-	6	11	15	33	34	15	2	-
Male	-	1	1	-	4	7	7	17	13	-	-	-
Female	-	-	1	-	2	4	8	16	21	15	2	-
White, both sexes	-	1	2	-	5	11	15	33	34	15	2	-
Male	-	1	1	-	3	7	7	17	13	-	-	-
Female	-	-	1	-	2	4	8	16	21	15	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Ohio												
All races, both sexes	36	66	57	77	113	191	287	357	260	110	31	-
Male	21	32	28	42	63	87	128	146	95	20	7	-
Female	15	34	29	35	50	104	159	211	165	90	24	-
White, both sexes	28	53	46	64	100	175	264	338	248	103	27	-
Male	16	29	20	37	54	77	120	137	90	20	6	-
Female	12	24	26	27	46	98	144	201	158	83	21	-
Black, both sexes	7	13	11	13	13	16	23	18	12	7	3	-
Male	4	3	8	5	9	10	8	8	5	-	1	-
Female	3	10	3	8	4	6	15	10	7	7	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Oklahoma												
All races, both sexes	24	27	26	36	62	107	146	162	107	48	10	-
Male	10	14	16	19	32	57	69	71	37	13	3	-
Female	14	13	10	17	30	50	77	91	70	35	7	-
White, both sexes	20	22	21	31	55	99	138	153	103	44	8	-
Male	8	13	13	18	27	54	68	66	34	12	3	-
Female	12	9	8	13	28	45	70	87	69	32	5	-
Black, both sexes	4	3	2	2	3	6	4	6	2	2	1	-
Male	2	1	-	-	2	3	-	3	2	-	-	-
Female	2	2	2	2	1	3	4	3	-	2	1	-
American Indian 1/, both sexes	-	2	3	3	4	2	4	3	2	2	1	-
Male	-	-	3	1	3	-	1	2	1	1	-	-
Female	-	2	-	2	1	2	3	1	1	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Oregon												
All races, both sexes	1	9	10	14	32	38	78	103	100	39	11	-
Male	-	7	5	10	17	18	45	51	45	15	1	-
Female	1	2	5	4	15	20	33	52	55	24	10	-
White, both sexes	1	9	9	14	32	38	77	103	99	38	10	-
Male	-	7	5	10	17	18	44	51	45	15	1	-
Female	1	2	4	4	15	20	33	52	54	23	9	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	1	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Pennsylvania												
All races, both sexes	35	52	64	74	138	267	485	643	480	224	50	-
Male	24	32	39	46	71	141	250	277	156	49	7	-
Female	11	20	25	28	67	126	235	366	324	175	43	-
White, both sexes	25	39	53	58	122	246	452	612	461	214	49	-
Male	15	24	33	34	62	132	236	265	151	47	7	-
Female	10	15	20	24	60	114	216	347	310	167	42	-
Black, both sexes	9	13	11	15	15	19	30	30	19	9	1	-
Male	8	8	6	11	8	9	14	12	5	2	-	-
Female	1	5	5	4	7	10	16	18	14	7	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	2	3	1	-	-	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	3	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Rhode Island												
All races, both sexes	2	4	7	9	11	29	43	55	51	23	10	-
Male	2	3	4	5	4	14	21	23	16	4	2	-
Female	-	1	3	4	7	15	22	32	35	19	8	-
White, both sexes	2	4	7	7	11	27	43	53	49	22	9	-
Male	2	3	4	3	4	13	21	23	14	3	2	-
Female	-	1	3	4	7	14	22	30	35	19	7	-
Black, both sexes	-	-	-	2	-	1	-	1	1	-	1	-
Male	-	-	-	2	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
South Carolina												
All races, both sexes	19	30	24	41	35	78	117	129	92	36	11	-
Male	11	20	16	26	19	39	56	46	33	11	3	-
Female	8	10	8	15	16	39	61	83	59	25	8	-
White, both sexes	9	20	15	24	22	66	96	108	74	32	6	-
Male	6	13	7	15	12	32	46	38	26	9	1	-
Female	3	7	8	9	10	34	50	70	48	23	5	-
Black, both sexes	10	10	9	17	13	10	21	21	18	4	5	-
Male	5	7	9	11	7	7	10	8	7	2	2	-
Female	5	3	-	6	6	3	11	13	11	2	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
South Dakota												
All races, both sexes	3	-	1	4	5	7	26	38	37	19	7	-
Male	2	-	-	4	4	4	13	19	13	1	2	-
Female	1	-	1	-	1	3	13	19	24	18	5	-
White, both sexes	2	-	1	3	3	5	25	38	37	19	7	-
Male	2	-	-	3	2	3	12	19	13	1	2	-
Female	-	-	1	-	1	2	13	19	24	18	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	2	2	1	-	-	-	-	-
Male	-	-	-	1	2	1	1	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	36	30	42	59	103	161	262	307	205	80	16	-
Male	22	12	24	25	50	85	114	118	62	18	3	-
Female	14	18	18	34	53	76	148	189	143	62	13	-
White, both sexes	23	23	32	52	90	137	232	290	192	75	13	-
Male	13	8	15	21	42	73	100	113	57	18	3	-
Female	10	15	17	31	48	64	132	177	135	57	10	-
Black, both sexes	13	7	10	7	12	24	29	17	13	5	3	-
Male	9	4	9	4	7	12	13	5	5	-	-	-
Female	4	3	1	3	5	12	16	12	8	5	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Texas												
All races, both sexes	74	90	100	149	213	336	559	580	486	218	42	-
Male	43	46	56	82	125	175	228	232	150	68	13	-
Female	31	44	44	67	88	161	331	348	336	150	29	-
White, both sexes	59	70	80	126	195	297	500	542	444	196	34	-
Male	36	34	46	68	114	152	207	222	136	59	9	-
Female	23	36	34	58	81	145	293	320	308	137	25	-
Black, both sexes	13	19	20	19	15	29	50	33	37	21	7	-
Male	7	12	10	13	10	18	17	9	11	8	4	-
Female	6	7	10	6	5	11	33	24	26	13	3	-
American Indian 1/, both sexes	2	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	2	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	3	10	9	5	5	1	1	-
Male	-	-	-	-	1	5	4	1	3	1	-	-
Female	-	1	-	2	2	5	5	4	2	-	1	-
Utah												
All races, both sexes	6	11	6	15	19	32	45	70	61	23	5	-
Male	3	3	3	5	9	13	24	36	23	7	1	-
Female	3	8	3	10	10	19	21	34	38	16	4	-
White, both sexes	6	10	5	14	19	31	43	69	60	23	3	-
Male	3	2	3	4	9	13	23	36	22	7	1	-
Female	3	8	2	10	10	18	20	33	38	16	2	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	1	1	-	1	-	-	-
Male	-	1	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Vermont												
All races, both sexes	-	1	1	1	1	7	6	18	12	6	3	-
Male	-	1	-	1	1	4	3	9	3	1	-	-
Female	-	-	1	-	-	3	3	9	9	5	3	-
White, both sexes	-	1	1	1	1	7	6	18	12	6	3	-
Male	-	1	-	1	1	4	3	9	3	1	-	-
Female	-	-	1	-	-	3	3	9	9	5	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	23	22	32	56	81	135	236	243	200	87	23	-
Male	12	13	19	30	41	77	127	104	67	18	3	-
Female	11	9	13	26	40	58	109	139	133	69	20	-
White, both sexes	12	17	29	44	69	112	200	220	176	76	20	-
Male	6	10	17	24	34	66	108	96	62	15	3	-
Female	6	7	12	20	35	46	92	124	114	61	17	-
Black, both sexes	11	5	2	12	9	19	35	18	21	9	3	-
Male	6	3	1	6	5	9	18	6	4	2	-	-
Female	5	2	1	6	4	10	17	12	17	7	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	3	4	1	5	3	2	-	-
Male	-	-	1	-	2	2	1	2	1	1	-	-
Female	-	-	-	-	1	2	-	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Washington												
All races, both sexes	12	12	18	25	29	59	104	137	119	73	15	-
Male	9	7	12	13	12	34	41	70	39	24	4	-
Female	3	5	6	12	17	25	63	67	80	49	11	-
White, both sexes	11	11	17	22	24	55	95	132	116	69	15	-
Male	9	7	11	11	12	31	38	68	38	22	4	-
Female	2	4	6	11	12	24	57	64	78	47	11	-
Black, both sexes	-	1	-	1	1	1	1	1	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	1	-	-
Female	-	1	-	-	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	2	1	3	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	2	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	2	2	5	3	3	3	-	-
Male	-	-	1	1	-	2	2	2	1	1	-	-
Female	-	-	-	1	2	-	3	1	2	2	-	-
West Virginia												
All races, both sexes	5	10	11	19	32	55	82	84	63	31	12	-
Male	4	8	6	11	20	29	36	39	20	14	2	-
Female	1	2	5	8	12	26	46	45	43	17	10	-
White, both sexes	5	10	11	19	31	54	82	82	63	28	11	-
Male	4	8	6	11	19	28	36	38	20	12	2	-
Female	1	2	5	8	12	26	46	44	43	16	9	-
Black, both sexes	-	-	-	-	1	1	-	2	-	3	1	-
Male	-	-	-	-	1	1	-	1	-	2	-	-
Female	-	-	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Wisconsin												
All races, both sexes	14	11	22	28	48	99	161	212	184	95	26	-
Male	9	8	13	14	29	59	76	89	76	29	8	-
Female	5	3	9	14	19	40	85	123	108	66	18	-
White, both sexes	14	10	20	23	47	97	156	207	181	92	26	-
Male	9	8	13	11	29	58	73	86	75	29	8	-
Female	5	2	7	12	18	39	83	121	106	63	18	-
Black, both sexes	-	1	2	5	1	2	3	2	2	1	-	-
Male	-	-	-	3	-	1	2	1	1	-	-	-
Female	-	1	2	2	1	1	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	3	-	1	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
Wyoming												
All races, both sexes	-	-	1	3	5	11	21	18	28	8	7	-
Male	-	-	1	2	3	4	13	6	11	1	-	-
Female	-	-	-	1	2	7	8	12	17	7	7	-
White, both sexes	-	-	1	3	4	11	21	18	28	8	7	-
Male	-	-	1	2	2	4	13	6	11	1	-	-
Female	-	-	-	1	2	7	8	12	17	7	7	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
United States												
All races, both sexes	297	52	14	1	2	1	3	-	-	2	6	5
Male	119	30	6	-	-	1	3	-	-	-	4	5
Female	178	22	8	1	2	-	-	-	-	2	2	-
White, both sexes	251	35	8	1	1	1	2	-	-	1	5	3
Male	93	21	3	-	-	1	2	-	-	-	3	3
Female	158	14	5	1	1	-	-	-	-	1	2	-
Black, both sexes	37	14	5	-	1	-	1	-	-	1	1	2
Male	22	7	3	-	-	-	1	-	-	-	1	2
Female	15	7	2	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	3	1	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	5	1	1	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	2	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	1	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Arkansas												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	27	5	1	-	-	-	-	-	-	-	1	-
Male	9	4	-	-	-	-	-	-	-	-	-	-
Female	18	1	1	-	-	-	-	-	-	-	1	-
White, both sexes	21	4	1	-	-	-	-	-	-	-	1	-
Male	7	3	-	-	-	-	-	-	-	-	-	-
Female	14	1	1	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Colorado												
All races, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Florida												
All races, both sexes	13	-	2	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	2	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	12	6	-	-	-	-	-	-	-	-	1	-
Male	3	3	-	-	-	-	-	-	-	-	-	-
Female	9	3	-	-	-	-	-	-	-	-	1	-
White, both sexes	9	3	-	-	-	-	-	-	-	-	1	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	8	2	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	3	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Illinois												
All races, both sexes	14	2	-	-	-	-	-	-	-	-	-	1
Male	7	2	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	1	-	-	-	-	-	-	-	-	-	1
Male	4	1	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	7	1	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Iowa												
All races, both sexes	9	-	2	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	2	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	4	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Kentucky												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	2	-	-	1	-	-	-	-	1	1	-
Male	3	2	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	1	-	-	-	-	1	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	4	2	-	-	1	-	-	-	-	-	1	-
Male	3	2	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Massachusetts												
All races, both sexes	7	-	3	-	-	-	-	-	-	-	-	-
Male	3	-	2	-	-	-	-	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	2	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	13	2	1	-	-	-	-	-	-	-	-	-
Male	4	1	1	-	-	-	-	-	-	-	-	-
Female	9	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	1	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Minnesota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Missouri												
All races, both sexes	5	2	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	2	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Nebraska												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	17	3	2	-	-	-	-	-	-	-	-	1
Male	8	1	1	-	-	-	-	-	-	-	-	1
Female	9	2	1	-	-	-	-	-	-	-	-	-
White, both sexes	15	1	2	-	-	-	-	-	-	-	-	1
Male	8	1	1	-	-	-	-	-	-	-	-	1
Female	7	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
North Carolina												
All races, both sexes	8	3	-	-	-	1	-	-	-	-	-	-
Male	6	3	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	3	-	-	-	1	-	-	-	-	-	-
Male	5	3	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Ohio												
All races, both sexes	32	4	-	-	-	-	-	-	-	-	-	1
Male	15	3	-	-	-	-	-	-	-	-	-	1
Female	17	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	4	-	-	-	-	-	-	-	-	-	1
Male	15	3	-	-	-	-	-	-	-	-	-	1
Female	17	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4	1	-	-	-	-	-	-	-	-	2	-
Male	3	1	-	-	-	-	-	-	-	-	2	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	2	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Oregon												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	22	2	-	-	1	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	15	2	-	-	1	-	-	-	-	-	-	-
White, both sexes	21	1	-	-	1	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	14	1	-	-	1	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Rhode Island												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	4	1	1	-	-	-	-	-	-	-	-	-
Male	4	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
South Dakota												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4	3	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	2	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Texas												
All races, both sexes	17	4	1	1	-	-	1	-	-	-	-	-
Male	8	4	1	-	-	-	1	-	-	-	-	-
Female	9	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	15	4	1	1	-	-	1	-	-	-	-	-
Male	7	4	1	-	-	-	1	-	-	-	-	-
Female	8	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	7	1	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	1	-	-
White, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Washington												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute lower respiratory infections (J20-J22,U04)												
Wisconsin												
All races, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
United States												
All races, both sexes	5	6	7	8	16	25	36	39	48	15	6	-
Male	2	2	4	1	7	12	11	12	17	2	-	-
Female	3	4	3	7	9	13	25	27	31	13	6	-
White, both sexes	3	3	5	8	14	21	36	38	46	14	6	-
Male	1	-	3	1	6	9	11	12	15	2	-	-
Female	2	3	2	7	8	12	25	26	31	12	6	-
Black, both sexes	1	3	2	-	2	2	-	-	2	-	-	-
Male	-	2	1	-	1	2	-	-	2	-	-	-
Female	1	1	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	1	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	2	-	-	2	4	5	1	4	2	-	-
Male	-	-	-	-	-	2	1	-	1	1	-	-
Female	-	2	-	-	2	2	4	1	3	1	-	-
White, both sexes	-	1	-	-	2	2	5	-	4	1	-	-
Male	-	-	-	-	-	1	1	-	1	1	-	-
Female	-	1	-	-	2	1	4	-	3	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Colorado												
All races, both sexes	-	-	-	-	1	2	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	2	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Florida												
All races, both sexes	-	-	-	-	1	3	2	3	2	-	-	-
Male	-	-	-	-	-	1	1	2	1	-	-	-
Female	-	-	-	-	1	2	1	1	1	-	-	-
White, both sexes	-	-	-	-	1	3	2	3	2	-	-	-
Male	-	-	-	-	-	1	1	2	1	-	-	-
Female	-	-	-	-	1	2	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	2	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	2	-	-	-
White, both sexes	-	-	-	-	-	-	2	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Illinois												
All races, both sexes	-	1	1	-	1	-	2	1	3	1	1	-
Male	-	1	-	-	-	-	-	1	1	1	-	-
Female	-	-	1	-	1	-	2	-	2	-	1	-
White, both sexes	-	-	1	-	1	-	2	1	2	1	1	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	1	-	1	-	2	-	2	-	1	-
Black, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	1	1	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	-	-	-	-
White, both sexes	-	-	-	-	1	1	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Iowa												
All races, both sexes	1	-	-	1	1	1	1	2	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	1	-	-	1	-	1	-	2	-	-	-	-
White, both sexes	1	-	-	1	1	1	1	2	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	1	-	-	1	-	1	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	1	2	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
White, both sexes	-	-	-	-	-	-	1	2	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	1	-	-	-	2	-	2	2	1	2	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	2	1	2	-
White, both sexes	-	-	-	-	-	2	-	2	2	1	2	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	2	1	2	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Missouri												
All races, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Nebraska												
All races, both sexes	-	-	-	1	1	1	-	-	3	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	1	1	-	-	2	1	-	-
White, both sexes	-	-	-	1	1	1	-	-	3	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	1	1	-	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	2	-	2	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	1	1	-	-	2	3	2	2	-	-
Male	-	-	1	-	-	-	1	2	1	-	-	-
Female	-	-	-	1	-	-	1	1	1	2	-	-
White, both sexes	-	-	1	1	-	-	2	3	2	2	-	-
Male	-	-	1	-	-	-	1	2	1	-	-	-
Female	-	-	-	1	-	-	1	1	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	2	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Ohio												
All races, both sexes	2	1	-	3	1	2	3	7	5	2	1	-
Male	1	-	-	1	1	2	2	3	1	-	-	-
Female	1	1	-	2	-	-	1	4	4	2	1	-
White, both sexes	2	1	-	3	1	2	3	7	5	2	1	-
Male	1	-	-	1	1	2	2	3	1	-	-	-
Female	1	1	-	2	-	-	1	4	4	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	1	2	2	-	6	5	1	1	-
Male	-	-	-	-	1	1	-	1	3	-	-	-
Female	-	-	-	1	1	1	-	5	2	1	1	-
White, both sexes	-	-	-	1	2	2	-	6	5	1	1	-
Male	-	-	-	-	1	1	-	1	3	-	-	-
Female	-	-	-	1	1	1	-	5	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Rhode Island												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
South Dakota												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Texas												
All races, both sexes	-	-	1	-	1	1	2	-	4	1	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	4	1	-	-
White, both sexes	-	-	-	-	-	1	2	-	4	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	4	1	-	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Vermont												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	1	-	3	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	3	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22,U04)												
Wisconsin												
All races, both sexes	-	-	-	1	-	-	-	-	3	1	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	1	-	-	-	-	1	1	-	-
White, both sexes	-	-	-	1	-	-	-	-	3	1	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	1	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
United States												
All races, both sexes	214	52	14	1	2	1	3	-	-	1	6	5
Male	96	30	6	-	-	1	3	-	-	-	4	5
Female	118	22	8	1	2	-	-	-	-	1	2	-
White, both sexes	173	35	8	1	1	1	2	-	-	1	5	3
Male	73	21	3	-	-	1	2	-	-	-	3	3
Female	100	14	5	1	1	-	-	-	-	1	2	-
Black, both sexes	34	14	5	-	1	-	1	-	-	-	1	2
Male	21	7	3	-	-	-	1	-	-	-	1	2
Female	13	7	2	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	3	1	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	5	1	1	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	2	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	1	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Arkansas												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	24	5	1	-	-	-	-	-	-	-	1	-
Male	6	4	-	-	-	-	-	-	-	-	-	-
Female	18	1	1	-	-	-	-	-	-	-	1	-
White, both sexes	19	4	1	-	-	-	-	-	-	-	1	-
Male	5	3	-	-	-	-	-	-	-	-	-	-
Female	14	1	1	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Colorado												
All races, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Florida												
All races, both sexes	12	-	2	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	2	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	10	6	-	-	-	-	-	-	-	-	1	-
Male	3	3	-	-	-	-	-	-	-	-	-	-
Female	7	3	-	-	-	-	-	-	-	-	1	-
White, both sexes	7	3	-	-	-	-	-	-	-	-	1	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	6	2	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	3	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Illinois												
All races, both sexes	11	2	-	-	-	-	-	-	-	-	-	1
Male	6	2	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	1	-	-	-	-	-	-	-	-	-	1
Male	3	1	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	7	1	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Iowa												
All races, both sexes	5	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	2	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Kentucky												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	2	-	-	1	-	-	-	-	1	1	-
Male	3	2	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	1	-	-	-	-	1	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	4	2	-	-	1	-	-	-	-	-	1	-
Male	3	2	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Massachusetts												
All races, both sexes	5	-	3	-	-	-	-	-	-	-	-	-
Male	3	-	2	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	2	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	7	2	1	-	-	-	-	-	-	-	-	-
Male	3	1	1	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	1	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Minnesota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Missouri												
All races, both sexes	4	2	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	13	3	2	-	-	-	-	-	-	-	-	1
Male	7	1	1	-	-	-	-	-	-	-	-	1
Female	6	2	1	-	-	-	-	-	-	-	-	-
White, both sexes	11	1	2	-	-	-	-	-	-	-	-	1
Male	7	1	1	-	-	-	-	-	-	-	-	1
Female	4	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
North Carolina												
All races, both sexes	6	3	-	-	-	1	-	-	-	-	-	-
Male	5	3	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	3	-	-	-	1	-	-	-	-	-	-
Male	5	3	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Ohio												
All races, both sexes	22	4	-	-	-	-	-	-	-	-	-	1
Male	11	3	-	-	-	-	-	-	-	-	-	1
Female	11	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	4	-	-	-	-	-	-	-	-	-	1
Male	11	3	-	-	-	-	-	-	-	-	-	1
Female	11	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	2	-
Male	3	1	-	-	-	-	-	-	-	-	2	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	2	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	14	2	-	-	1	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	9	2	-	-	1	-	-	-	-	-	-	-
White, both sexes	13	1	-	-	1	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	1	-	-	1	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	4	1	1	-	-	-	-	-	-	-	-	-
Male	4	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
South Dakota												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	3	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Texas												
All races, both sexes	14	4	1	1	-	-	1	-	-	-	-	-
Male	8	4	1	-	-	-	1	-	-	-	-	-
Female	6	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	13	4	1	1	-	-	1	-	-	-	-	-
Male	7	4	1	-	-	-	1	-	-	-	-	-
Female	6	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Washington												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Wisconsin												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
United States												
All races, both sexes	4	4	5	6	12	15	25	27	22	6	3	-
Male	1	2	3	1	6	6	8	11	8	1	-	-
Female	3	2	2	5	6	9	17	16	14	5	3	-
White, both sexes	3	1	4	6	10	12	25	26	21	5	3	-
Male	1	-	2	1	5	4	8	11	7	1	-	-
Female	2	1	2	5	5	8	17	15	14	4	3	-
Black, both sexes	1	3	1	-	2	2	-	-	1	-	-	-
Male	-	2	1	-	1	2	-	-	1	-	-	-
Female	1	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	2	-	-	2	2	5	1	4	1	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	2	-	-	2	2	4	1	3	1	-	-
White, both sexes	-	1	-	-	2	1	5	-	4	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	1	-	-	2	1	4	-	3	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Colorado												
All races, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Florida												
All races, both sexes	-	-	-	-	1	2	2	3	2	-	-	-
Male	-	-	-	-	-	-	1	2	1	-	-	-
Female	-	-	-	-	1	2	1	1	1	-	-	-
White, both sexes	-	-	-	-	1	2	2	3	2	-	-	-
Male	-	-	-	-	-	-	1	2	1	-	-	-
Female	-	-	-	-	1	2	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Illinois												
All races, both sexes	-	1	1	-	1	-	1	1	1	1	1	-
Male	-	1	-	-	-	-	-	1	-	1	-	-
Female	-	-	1	-	1	-	1	-	1	-	1	-
White, both sexes	-	-	1	-	1	-	1	1	1	1	1	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	1	-	1	-	1	-	1	-	1	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	1	1	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	-	-	-	-
White, both sexes	-	-	-	-	1	1	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Iowa												
All races, both sexes	1	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	1	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	1	-	-	-	-	-	2	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Missouri												
All races, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	1	1	-	-	2	3	-	-	-	-
Male	-	-	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
White, both sexes	-	-	1	1	-	-	2	3	-	-	-	-
Male	-	-	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Ohio												
All races, both sexes	2	-	-	3	1	-	3	5	2	1	-	-
Male	1	-	-	1	1	-	2	2	-	-	-	-
Female	1	-	-	2	-	-	1	3	2	1	-	-
White, both sexes	2	-	-	3	1	-	3	5	2	1	-	-
Male	1	-	-	1	1	-	2	2	-	-	-	-
Female	1	-	-	2	-	-	1	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	2	2	-	4	1	-	1	-
Male	-	-	-	-	1	1	-	1	1	-	-	-
Female	-	-	-	-	1	1	-	3	-	-	1	-
White, both sexes	-	-	-	-	2	2	-	4	1	-	1	-
Male	-	-	-	-	1	1	-	1	1	-	-	-
Female	-	-	-	-	1	1	-	3	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Texas												
All races, both sexes	-	-	-	-	1	1	2	-	2	1	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	2	1	-	-
White, both sexes	-	-	-	-	-	1	2	-	2	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	2	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Vermont												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
Wisconsin												
All races, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
United States												
All races, both sexes	83	-	-	-	-	-	-	-	-	1	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	60	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	78	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Colorado												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Florida												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Illinois												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Iowa												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Massachusetts												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Minnesota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Missouri												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Nebraska												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
North Carolina												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Ohio												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Oregon												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Texas												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified acute lower respiratory infections (J22,U04)												
Wisconsin												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
United States												
All races, both sexes	1	2	2	2	4	10	11	12	26	9	3	-
Male	1	-	1	-	1	6	3	1	9	1	-	-
Female	-	2	1	2	3	4	8	11	17	8	3	-
White, both sexes	-	2	1	2	4	9	11	12	25	9	3	-
Male	-	-	1	-	1	5	3	1	8	1	-	-
Female	-	2	-	2	3	4	8	11	17	8	3	-
Black, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	1	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	2	-	-	-	1	-	-
Male	-	-	-	-	-	2	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Colorado												
All races, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Florida												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Illinois												
All races, both sexes	-	-	-	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Iowa												
All races, both sexes	-	-	-	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	2	-	-	1	1	2	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	1	2	-
White, both sexes	-	-	-	-	-	2	-	-	1	1	2	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	1	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Missouri												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Nebraska												
All races, both sexes	-	-	-	1	1	1	-	-	3	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	1	1	-	-	2	1	-	-
White, both sexes	-	-	-	1	1	1	-	-	3	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	1	1	-	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	1	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	2	2	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	2	-	-
White, both sexes	-	-	-	-	-	-	-	-	2	2	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Ohio												
All races, both sexes	-	1	-	-	-	2	-	2	3	1	1	-
Male	-	-	-	-	-	2	-	1	1	-	-	-
Female	-	1	-	-	-	-	-	1	2	1	1	-
White, both sexes	-	1	-	-	-	2	-	2	3	1	1	-
Male	-	-	-	-	-	2	-	1	1	-	-	-
Female	-	1	-	-	-	-	-	1	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	1	-	-	-	2	4	1	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	1	-	-	-	2	2	1	-	-
White, both sexes	-	-	-	1	-	-	-	2	4	1	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	1	-	-	-	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Rhode Island												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
South Dakota												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Texas												
All races, both sexes	-	-	1	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	2	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute lower respiratory infections (J22,U04)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	2	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	2	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
United States												
All races, both sexes	478	18	7	2	3	6	1	6	10	13	15	15
Male	248	11	3	2	2	5	-	5	8	9	9	8
Female	230	7	4	-	1	1	1	1	2	4	6	7
White, both sexes	372	11	5	-	2	2	1	6	5	11	11	10
Male	191	5	2	-	1	1	-	5	4	8	8	5
Female	181	6	3	-	1	1	1	1	1	3	3	5
Black, both sexes	86	6	1	1	1	3	-	-	5	2	4	5
Male	46	5	1	1	1	3	-	-	4	1	1	3
Female	40	1	-	-	-	-	-	-	1	1	3	2
American Indian 1/, both sexes	3	1	1	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	1	-	1	-	-	-	-	-	-
Male	9	-	-	1	-	1	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Arkansas												
All races, both sexes	3	1	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	39	4	1	-	-	1	-	2	1	1	2	-
Male	24	4	-	-	-	1	-	2	1	1	1	-
Female	15	-	1	-	-	-	-	-	-	-	1	-
White, both sexes	27	2	1	-	-	-	-	2	-	1	2	-
Male	16	2	-	-	-	-	-	2	-	1	1	-
Female	11	-	1	-	-	-	-	-	-	-	1	-
Black, both sexes	5	2	-	-	-	-	-	-	1	-	-	-
Male	4	2	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	1	-	-	-	-	-	-
Male	4	-	-	-	-	1	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Colorado												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Florida												
All races, both sexes	25	-	-	-	1	1	-	-	-	2	3	2
Male	14	-	-	-	-	1	-	-	-	2	3	1
Female	11	-	-	-	1	-	-	-	-	-	-	1
White, both sexes	21	-	-	-	1	1	-	-	-	2	2	1
Male	12	-	-	-	-	1	-	-	-	2	2	-
Female	9	-	-	-	1	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	15	1	-	-	-	-	-	-	1	-	2	1
Male	7	-	-	-	-	-	-	-	1	-	1	1
Female	8	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	1	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Hawaii												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Illinois												
All races, both sexes	25	2	1	-	-	1	-	1	1	1	-	-
Male	14	2	-	-	-	1	-	1	1	1	-	-
Female	11	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	18	1	1	-	-	-	-	1	1	-	-	-
Male	8	1	-	-	-	-	-	1	1	-	-	-
Female	10	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	7	1	-	-	-	1	-	-	-	1	-	-
Male	6	1	-	-	-	1	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	9	1	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	3	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	1	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Iowa												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	5	1	-	-	-	-	-	-	-	-	-	1
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	-	-	-	-	-	-	-	-	1
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Kentucky												
All races, both sexes	11	-	-	-	-	-	-	1	-	1	1	-
Male	7	-	-	-	-	-	-	1	-	1	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	1	-	1	1	-
Male	7	-	-	-	-	-	-	1	-	1	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Maine												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Massachusetts												
All races, both sexes	9	2	-	-	-	-	-	-	-	1	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	1	-	-
White, both sexes	9	2	-	-	-	-	-	-	-	1	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	24	-	-	-	-	-	-	-	1	2	-	1
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	18	-	-	-	-	-	-	-	-	2	-	1
White, both sexes	16	-	-	-	-	-	-	-	1	1	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Minnesota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	7	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	1	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Missouri												
All races, both sexes	13	-	1	-	1	-	-	-	-	-	1	-
Male	9	-	1	-	1	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	9	-	1	-	-	-	-	-	-	-	1	-
Male	6	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	1	-	-	-	-	-	-	-
Male	3	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Nebraska												
All races, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
New Mexico												
All races, both sexes	4	-	1	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	19	2	-	1	-	1	-	-	-	-	-	-
Male	8	1	-	1	-	-	-	-	-	-	-	-
Female	11	1	-	-	-	1	-	-	-	-	-	-
White, both sexes	15	2	-	-	-	1	-	-	-	-	-	-
Male	6	1	-	-	-	-	-	-	-	-	-	-
Female	9	1	-	-	-	1	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
North Carolina												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Ohio												
All races, both sexes	31	-	-	-	-	1	-	1	1	3	1	1
Male	14	-	-	-	-	1	-	-	1	2	-	-
Female	17	-	-	-	-	-	-	1	-	1	1	1
White, both sexes	27	-	-	-	-	-	-	1	-	3	-	1
Male	12	-	-	-	-	-	-	-	-	2	-	-
Female	15	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	3	-	-	-	-	1	-	-	1	-	1	-
Male	2	-	-	-	-	1	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Oregon												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	15	1	-	-	1	-	-	1	-	-	-	-
Male	6	-	-	-	1	-	-	1	-	-	-	-
Female	9	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	1	-	-	1	-	-	1	-	-	-	-
Male	6	-	-	-	1	-	-	1	-	-	-	-
Female	7	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
South Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	15	1	-	-	-	-	-	-	1	-	-	1
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	8	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	11	1	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Texas												
All races, both sexes	38	-	-	-	-	-	-	-	-	1	-	1
Male	19	-	-	-	-	-	-	-	-	1	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	33	-	-	-	-	-	-	-	-	1	-	1
Male	16	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	13	-	1	1	-	-	-	-	-	-	1	-
Male	5	-	1	1	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	1	1	-	-	-	-	-	-	1	-
Male	4	-	1	1	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Washington												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	6	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
Wisconsin												
All races, both sexes	9	1	-	-	-	-	-	-	-	-	1	-
Male	7	1	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
United States												
All races, both sexes	35	30	36	28	38	60	50	61	34	8	2	-
Male	16	15	19	13	21	35	22	28	16	1	-	-
Female	19	15	17	15	17	25	28	33	18	7	2	-
White, both sexes	23	20	24	21	29	54	45	54	30	6	2	-
Male	11	10	14	9	16	30	21	26	14	1	-	-
Female	12	10	10	12	13	24	24	28	16	5	2	-
Black, both sexes	9	8	10	7	8	3	4	6	2	1	-	-
Male	3	4	5	4	4	3	1	2	-	-	-	-
Female	6	4	5	3	4	-	3	4	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	2	-	1	2	1	1	2	1	-	-
Male	2	1	-	-	1	1	-	-	2	-	-	-
Female	1	1	2	-	-	1	1	1	-	1	-	-
Alabama												
All races, both sexes	2	1	-	-	1	-	-	2	-	-	-	-
Male	2	-	-	-	1	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Alaska												
All races, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	1	-	2	2	2	-	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	-	-	-	-
White, both sexes	-	-	-	1	-	2	2	1	-	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Arkansas												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	6	5	1	1	2	5	2	1	3	1	-	-
Male	3	4	-	-	1	5	-	-	1	-	-	-
Female	3	1	1	1	1	-	2	1	2	1	-	-
White, both sexes	5	3	-	-	2	4	2	1	2	-	-	-
Male	3	2	-	-	1	4	-	-	-	-	-	-
Female	2	1	-	-	1	-	2	1	2	-	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	-	-	1	-	-	1	1	-	-
Male	-	1	-	-	-	1	-	-	1	-	-	-
Female	1	-	1	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Colorado												
All races, both sexes	-	1	2	-	2	-	1	1	-	-	-	-
Male	-	1	2	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	-	1	2	-	2	-	1	1	-	-	-	-
Male	-	1	2	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	-	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	1	-	1	1	-	-	-	-
White, both sexes	-	1	-	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Florida												
All races, both sexes	1	-	3	1	1	2	2	5	1	-	-	-
Male	-	-	2	-	-	2	2	1	-	-	-	-
Female	1	-	1	1	1	-	-	4	1	-	-	-
White, both sexes	-	-	2	1	1	2	2	5	1	-	-	-
Male	-	-	2	-	-	2	2	1	-	-	-	-
Female	-	-	-	1	1	-	-	4	1	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	1	4	-	1	-	-	1	-	-	-	-
Male	1	1	-	-	1	-	-	1	-	-	-	-
Female	2	-	4	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	2	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	2	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	2	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	1	-	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Hawaii												
All races, both sexes	1	1	-	-	-	1	-	-	1	-	-	-
Male	1	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Illinois												
All races, both sexes	2	-	1	2	-	6	2	-	4	-	1	-
Male	1	-	-	1	-	3	-	-	3	-	-	-
Female	1	-	1	1	-	3	2	-	1	-	1	-
White, both sexes	1	-	-	1	-	5	2	-	4	-	1	-
Male	-	-	-	-	-	2	-	-	3	-	-	-
Female	1	-	-	1	-	3	2	-	1	-	1	-
Black, both sexes	1	-	1	1	-	1	-	-	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	2	-	1	-	2	-	1	1	-	-	-
Male	-	1	-	1	-	2	-	-	1	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	2	-	1	1	-	-	-
Male	-	-	-	1	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Iowa												
All races, both sexes	2	1	-	-	-	2	1	2	-	-	-	-
Male	1	1	-	-	-	-	-	2	-	-	-	-
Female	1	-	-	-	-	2	1	-	-	-	-	-
White, both sexes	2	1	-	-	-	2	1	2	-	-	-	-
Male	1	1	-	-	-	-	-	2	-	-	-	-
Female	1	-	-	-	-	2	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Kentucky												
All races, both sexes	-	-	-	1	1	1	3	2	-	-	-	-
Male	-	-	-	1	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	-	-	-	-
White, both sexes	-	-	-	1	1	1	3	1	-	-	-	-
Male	-	-	-	1	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	1	1	-	2	2	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	2	1	-	-	-	-
White, both sexes	-	-	1	1	-	1	2	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Maine												
All races, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	2	-	1	1	1	2	-	1	-	-
Male	1	-	2	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	1	-	-
White, both sexes	1	-	-	-	1	1	1	2	-	-	-	-
Male	1	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	2	-	-	-	-	-	-	1	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Massachusetts												
All races, both sexes	1	-	-	-	1	1	1	-	2	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	1	-	-	-	-	1	1	-	1	-	-	-
White, both sexes	1	-	-	-	1	1	1	-	2	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	1	-	-	-	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	2	1	1	2	3	2	4	3	-	-	-
Male	1	1	-	-	-	-	2	1	-	-	-	-
Female	1	1	1	1	2	3	-	3	3	-	-	-
White, both sexes	1	1	-	-	-	3	2	4	3	-	-	-
Male	1	1	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	3	-	3	3	-	-	-
Black, both sexes	1	1	1	1	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	1	1	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Minnesota												
All races, both sexes	-	-	-	-	-	2	-	2	-	-	-	-
Male	-	-	-	-	-	2	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	-	1	-	-	1	1	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Missouri												
All races, both sexes	-	-	2	1	1	1	2	3	-	-	-	-
Male	-	-	2	-	-	-	2	3	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	1	1	3	-	-	-	-
Male	-	-	1	-	-	-	1	3	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	1	1	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Nebraska												
All races, both sexes	-	-	-	-	-	2	-	2	-	-	-	-
Male	-	-	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	2	-	2	-	-	-	-
Male	-	-	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
New Hampshire												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	1	2	2	-	2	2	1	1	-	-	-
Male	-	1	1	2	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	2	1	1	-	-	-
White, both sexes	-	1	2	1	-	2	2	1	1	-	-	-
Male	-	1	1	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	2	1	1	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
New Mexico												
All races, both sexes	-	-	-	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	2	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	2	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	1	1	-	4	3	3	2	-	-	-	-
Male	-	-	-	-	2	2	1	1	-	-	-	-
Female	1	1	1	-	2	1	2	1	-	-	-	-
White, both sexes	1	-	1	-	3	3	2	2	-	-	-	-
Male	-	-	-	-	1	2	1	1	-	-	-	-
Female	1	-	1	-	2	1	1	1	-	-	-	-
Black, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
North Carolina												
All races, both sexes	1	-	-	2	2	1	-	1	-	-	-	-
Male	1	-	-	2	-	1	-	1	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Ohio												
All races, both sexes	-	3	4	2	3	2	2	1	3	2	1	-
Male	-	-	2	-	2	2	1	1	2	-	-	-
Female	-	3	2	2	1	-	1	-	1	2	1	-
White, both sexes	-	2	4	2	3	2	2	1	3	2	1	-
Male	-	-	2	-	2	2	1	1	2	-	-	-
Female	-	2	2	2	1	-	1	-	1	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	1	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Oregon												
All races, both sexes	1	1	-	2	1	1	1	1	1	-	-	-
Male	-	-	-	1	1	1	1	1	1	-	-	-
Female	1	1	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	2	1	1	1	1	1	-	-	-
Male	-	-	-	1	1	1	1	1	1	-	-	-
Female	1	1	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	1	-	-	1	2	3	4	-	-	-	-
Male	-	1	-	-	-	1	1	1	-	-	-	-
Female	1	-	-	-	1	1	2	3	-	-	-	-
White, both sexes	1	1	-	-	1	2	3	2	-	-	-	-
Male	-	1	-	-	-	1	1	1	-	-	-	-
Female	1	-	-	-	1	1	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	1	1	1	-	1	2	1	-	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	-	-	2	1	-	-	-	-
White, both sexes	1	-	1	1	-	-	2	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	2	1	-	-	-	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
South Dakota												
All races, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	2	1	-	2	3	1	-	1	-	-	-
Male	1	-	1	-	1	2	1	-	-	-	-	-
Female	1	2	-	-	1	1	-	-	1	-	-	-
White, both sexes	-	2	1	-	1	3	1	-	1	-	-	-
Male	-	-	1	-	1	2	1	-	-	-	-	-
Female	-	2	-	-	-	1	-	-	1	-	-	-
Black, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Texas												
All races, both sexes	1	-	3	4	3	5	7	9	4	-	-	-
Male	-	-	2	1	2	3	4	5	1	-	-	-
Female	1	-	1	3	1	2	3	4	3	-	-	-
White, both sexes	1	-	2	4	2	5	5	8	4	-	-	-
Male	-	-	1	1	1	3	4	4	1	-	-	-
Female	1	-	1	3	1	2	1	4	3	-	-	-
Black, both sexes	-	-	1	-	1	-	1	1	-	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	1	1	2	-	-	1	3	1	1	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	3	1	1	-	-
White, both sexes	-	1	-	1	-	-	-	3	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	3	-	1	-	-
Black, both sexes	-	-	1	1	-	-	1	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Washington												
All races, both sexes	1	1	-	-	1	2	-	-	2	2	-	-
Male	1	1	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	2	-	-	2	1	-	-
White, both sexes	1	1	-	-	-	2	-	-	1	2	-	-
Male	1	1	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	2	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	1	2	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	2	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Wisconsin												
All races, both sexes	-	-	2	-	1	1	-	-	3	-	-	-
Male	-	-	1	-	1	-	-	-	3	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	2	-	1	1	-	-	3	-	-	-
Male	-	-	1	-	1	-	-	-	3	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
United States												
All races, both sexes	124,583	30	43	52	63	61	90	113	142	222	619	1,320
Male	59,260	20	29	33	45	39	54	66	71	105	291	617
Female	65,323	10	14	19	18	22	36	47	71	117	328	703
White, both sexes	114,993	15	26	24	18	21	48	64	77	146	445	1,076
Male	54,043	10	17	18	14	12	28	38	32	69	213	498
Female	60,950	5	9	6	4	9	20	26	45	77	232	578
Black, both sexes	7,730	11	15	28	41	37	38	46	62	74	152	216
Male	4,136	6	12	15	29	25	22	25	36	34	69	103
Female	3,594	5	3	13	12	12	16	21	26	40	83	113
American Indian 1/, both sexes	508	2	-	-	2	-	1	-	1	1	11	13
Male	234	2	-	-	1	-	1	-	1	1	6	7
Female	274	-	-	-	1	-	-	-	-	-	5	6
Asian or Pacific Islander 2/, both sexes	1,352	2	2	-	2	3	3	3	2	1	11	15
Male	847	2	-	-	1	2	3	3	2	1	3	9
Female	505	-	2	-	1	1	-	-	-	-	8	6
Alabama												
All races, both sexes	2,309	-	1	-	2	-	1	1	3	5	14	26
Male	1,155	-	1	-	2	-	1	1	1	2	5	15
Female	1,154	-	-	-	-	-	-	-	2	3	9	11
White, both sexes	2,068	-	-	-	1	-	-	-	1	4	9	22
Male	999	-	-	-	1	-	-	-	1	2	4	14
Female	1,069	-	-	-	-	-	-	-	-	2	5	8
Black, both sexes	238	-	1	-	1	-	1	1	2	1	5	4
Male	155	-	1	-	1	-	1	1	-	-	1	1
Female	83	-	-	-	-	-	-	-	2	1	4	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Alaska												
All races, both sexes	140	-	-	-	-	-	-	-	-	1	2	-
Male	58	-	-	-	-	-	-	-	-	-	-	-
Female	82	-	-	-	-	-	-	-	-	1	2	-
White, both sexes	100	-	-	-	-	-	-	-	-	1	2	-
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	59	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2,799	1	2	1	4	-	-	1	3	4	6	19
Male	1,358	1	2	1	2	-	-	1	1	4	2	7
Female	1,441	-	-	-	2	-	-	-	2	-	4	12
White, both sexes	2,744	-	2	1	1	-	-	1	3	3	5	18
Male	1,332	-	2	1	1	-	-	1	1	3	2	7
Female	1,412	-	-	-	-	-	-	-	2	-	3	11
Black, both sexes	33	-	-	-	1	-	-	-	-	1	1	1
Male	15	-	-	-	-	-	-	-	-	1	-	-
Female	18	-	-	-	1	-	-	-	-	-	1	1
American Indian 1/, both sexes	16	1	-	-	2	-	-	-	-	-	-	-
Male	5	1	-	-	1	-	-	-	-	-	-	-
Female	11	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Arkansas												
All races, both sexes	1,498	-	1	-	-	1	2	1	3	3	11	7
Male	786	-	1	-	-	1	-	-	1	2	7	3
Female	712	-	-	-	-	-	2	1	2	1	4	4
White, both sexes	1,414	-	-	-	-	-	1	1	1	3	11	6
Male	742	-	-	-	-	-	-	-	-	2	7	3
Female	672	-	-	-	-	-	1	1	1	1	4	3
Black, both sexes	79	-	1	-	-	1	1	-	1	-	-	1
Male	43	-	1	-	-	1	-	-	-	-	-	-
Female	36	-	-	-	-	-	1	-	1	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	12,829	3	8	7	7	10	11	15	14	23	55	117
Male	5,948	2	5	4	7	6	8	9	7	11	21	58
Female	6,881	1	3	3	-	4	3	6	7	12	34	59
White, both sexes	11,360	2	7	3	-	6	6	5	8	11	35	96
Male	5,090	1	4	2	-	5	4	2	1	5	16	47
Female	6,270	1	3	1	-	1	2	3	7	6	19	49
Black, both sexes	712	-	1	4	6	3	5	7	4	11	13	17
Male	382	-	1	2	6	-	4	4	4	5	4	7
Female	330	-	-	2	-	3	1	3	-	6	9	10
American Indian 1/, both sexes	59	-	-	-	-	-	-	-	-	1	1	-
Male	30	-	-	-	-	-	-	-	-	1	-	-
Female	29	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	698	1	-	-	1	1	-	3	2	-	6	4
Male	446	1	-	-	1	1	-	3	2	-	1	4
Female	252	-	-	-	-	-	-	-	-	-	5	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Colorado												
All races, both sexes	1,932	-	-	-	-	1	1	2	4	1	7	15
Male	971	-	-	-	-	1	1	1	3	-	3	5
Female	961	-	-	-	-	-	-	1	1	1	4	10
White, both sexes	1,861	-	-	-	-	1	1	2	3	1	4	14
Male	940	-	-	-	-	1	1	1	2	-	2	4
Female	921	-	-	-	-	-	-	1	1	1	2	10
Black, both sexes	54	-	-	-	-	-	-	-	1	-	1	1
Male	26	-	-	-	-	-	-	-	1	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1,458	-	-	2	1	-	-	-	-	4	4	12
Male	648	-	-	1	1	-	-	-	-	1	2	5
Female	810	-	-	1	-	-	-	-	-	3	2	7
White, both sexes	1,404	-	-	2	-	-	-	-	-	3	3	9
Male	617	-	-	1	-	-	-	-	-	-	1	4
Female	787	-	-	1	-	-	-	-	-	3	2	5
Black, both sexes	52	-	-	-	1	-	-	-	-	1	1	3
Male	29	-	-	-	1	-	-	-	-	1	1	1
Female	23	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Delaware												
All races, both sexes	344	-	-	-	-	-	-	2	-	-	1	-
Male	158	-	-	-	-	-	-	-	-	-	1	-
Female	186	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	303	-	-	-	-	-	-	1	-	-	-	-
Male	134	-	-	-	-	-	-	-	-	-	-	-
Female	169	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	37	-	-	-	-	-	-	1	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	126	1	-	-	-	-	-	1	-	2	3	4
Male	60	1	-	-	-	-	-	1	-	-	1	3
Female	66	-	-	-	-	-	-	-	-	2	2	1
White, both sexes	45	-	-	-	-	-	-	-	-	1	1	2
Male	19	-	-	-	-	-	-	-	-	-	-	2
Female	26	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	80	1	-	-	-	-	-	1	-	1	2	2
Male	40	1	-	-	-	-	-	1	-	-	1	1
Female	40	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Florida												
All races, both sexes	8,912	3	2	4	4	3	7	9	10	12	32	94
Male	4,189	2	1	3	3	-	5	7	4	2	15	40
Female	4,723	1	1	1	1	3	2	2	6	10	17	54
White, both sexes	8,434	1	1	2	2	2	4	6	5	8	28	77
Male	3,917	1	-	2	2	-	3	5	2	1	13	32
Female	4,517	-	1	-	-	2	1	1	3	7	15	45
Black, both sexes	444	2	1	2	2	1	3	3	5	4	3	16
Male	251	1	1	1	1	-	2	2	2	1	2	7
Female	193	1	-	1	1	1	1	1	3	3	1	9
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
Georgia												
All races, both sexes	3,387	-	1	5	4	2	2	7	4	11	23	45
Male	1,631	-	1	3	2	1	1	4	2	3	9	17
Female	1,756	-	-	2	2	1	1	3	2	8	14	28
White, both sexes	2,953	-	-	2	1	-	-	4	-	5	12	27
Male	1,383	-	-	2	-	-	-	2	-	1	5	8
Female	1,570	-	-	-	1	-	-	2	-	4	7	19
Black, both sexes	409	-	1	3	3	2	2	3	4	6	10	17
Male	236	-	1	1	2	1	1	2	2	2	3	9
Female	173	-	-	2	1	1	1	1	2	4	7	8
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Hawaii												
All races, both sexes	293	-	-	-	-	1	-	-	-	1	-	5
Male	161	-	-	-	-	-	-	-	-	1	-	2
Female	132	-	-	-	-	1	-	-	-	-	-	3
White, both sexes	105	-	-	-	-	-	-	-	-	-	-	2
Male	53	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	186	-	-	-	-	1	-	-	-	1	-	3
Male	106	-	-	-	-	-	-	-	-	1	-	1
Female	80	-	-	-	-	1	-	-	-	-	-	2
Idaho												
All races, both sexes	644	-	-	-	-	-	-	-	-	-	1	6
Male	336	-	-	-	-	-	-	-	-	-	-	3
Female	308	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	640	-	-	-	-	-	-	-	-	-	1	6
Male	333	-	-	-	-	-	-	-	-	-	-	3
Female	307	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Illinois												
All races, both sexes	4,731	-	2	4	2	4	7	5	18	13	28	51
Male	2,238	-	2	2	2	4	3	4	14	8	15	21
Female	2,493	-	-	2	-	-	4	1	4	5	13	30
White, both sexes	4,253	-	-	3	-	-	3	2	6	9	18	32
Male	1,981	-	-	2	-	-	1	2	4	6	10	14
Female	2,272	-	-	1	-	-	2	-	2	3	8	18
Black, both sexes	446	-	2	1	2	4	4	3	12	4	10	18
Male	239	-	2	-	2	4	2	2	10	2	5	7
Female	207	-	-	1	-	-	2	1	2	2	5	11
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Indiana												
All races, both sexes	3,291	2	1	-	3	1	3	1	2	10	17	47
Male	1,591	2	-	-	1	1	1	-	1	5	7	23
Female	1,700	-	1	-	2	-	2	1	1	5	10	24
White, both sexes	3,151	2	1	-	1	1	1	1	1	8	14	42
Male	1,524	2	-	-	1	1	1	-	1	3	6	20
Female	1,627	-	1	-	-	-	-	1	-	5	8	22
Black, both sexes	136	-	-	-	2	-	2	-	1	2	3	5
Male	66	-	-	-	-	-	-	-	-	2	1	3
Female	70	-	-	-	2	-	2	-	1	-	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Iowa												
All races, both sexes	1,655	-	-	-	-	1	1	-	3	3	11	8
Male	868	-	-	-	-	1	1	-	1	1	6	3
Female	787	-	-	-	-	-	-	-	2	2	5	5
White, both sexes	1,641	-	-	-	-	-	1	-	2	3	9	8
Male	860	-	-	-	-	-	1	-	-	1	5	3
Female	781	-	-	-	-	-	-	-	2	2	4	5
Black, both sexes	8	-	-	-	-	1	-	-	1	-	1	-
Male	7	-	-	-	-	1	-	-	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1,501	-	-	-	-	-	1	4	4	5	9	20
Male	756	-	-	-	-	-	1	3	2	2	4	8
Female	745	-	-	-	-	-	-	1	2	3	5	12
White, both sexes	1,430	-	-	-	-	-	-	4	2	5	7	19
Male	725	-	-	-	-	-	-	3	-	2	3	8
Female	705	-	-	-	-	-	-	1	2	3	4	11
Black, both sexes	46	-	-	-	-	-	1	-	2	-	1	1
Male	21	-	-	-	-	-	1	-	2	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Kentucky												
All races, both sexes	2,407	-	1	-	-	1	2	2	1	5	16	49
Male	1,160	-	1	-	-	1	-	1	1	4	9	23
Female	1,247	-	-	-	-	-	2	1	-	1	7	26
White, both sexes	2,295	-	1	-	-	-	2	1	1	4	14	44
Male	1,108	-	1	-	-	-	-	1	1	3	9	22
Female	1,187	-	-	-	-	-	2	-	-	1	5	22
Black, both sexes	106	-	-	-	-	1	-	1	-	1	2	5
Male	48	-	-	-	-	1	-	-	-	1	-	1
Female	58	-	-	-	-	-	-	1	-	-	2	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,687	2	2	1	1	-	1	1	2	3	12	21
Male	863	2	2	-	1	-	1	1	1	1	5	11
Female	824	-	-	1	-	-	-	-	1	2	7	10
White, both sexes	1,395	-	-	-	-	-	-	1	1	2	6	15
Male	683	-	-	-	-	-	-	1	-	-	2	6
Female	712	-	-	-	-	-	-	-	1	2	4	9
Black, both sexes	287	2	2	1	1	-	1	-	1	1	6	6
Male	176	2	2	-	1	-	1	-	1	1	3	5
Female	111	-	-	1	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Maine												
All races, both sexes	783	-	-	-	1	-	-	-	-	1	4	7
Male	390	-	-	-	1	-	-	-	-	-	2	6
Female	393	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	781	-	-	-	1	-	-	-	-	1	4	7
Male	390	-	-	-	1	-	-	-	-	-	2	6
Female	391	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,832	-	1	-	2	1	1	3	3	2	12	16
Male	792	-	-	-	1	1	1	1	1	-	5	3
Female	1,040	-	1	-	1	-	-	2	2	2	7	13
White, both sexes	1,559	-	-	-	-	-	-	1	1	2	4	13
Male	667	-	-	-	-	-	-	-	1	-	-	2
Female	892	-	-	-	-	-	-	1	-	2	4	11
Black, both sexes	255	-	1	-	2	1	1	2	2	-	8	3
Male	116	-	-	-	1	1	1	1	-	-	5	1
Female	139	-	1	-	1	-	-	1	2	-	3	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Massachusetts												
All races, both sexes	2,535	-	-	-	-	-	2	2	2	1	13	16
Male	1,065	-	-	-	-	-	-	2	1	-	5	5
Female	1,470	-	-	-	-	-	2	-	1	1	8	11
White, both sexes	2,462	-	-	-	-	-	1	2	2	1	8	15
Male	1,027	-	-	-	-	-	-	2	1	-	4	5
Female	1,435	-	-	-	-	-	1	-	1	1	4	10
Black, both sexes	54	-	-	-	-	-	1	-	-	-	5	1
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	33	-	-	-	-	-	1	-	-	-	4	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4,474	-	1	4	3	5	4	8	3	6	26	51
Male	2,139	-	-	2	2	4	3	5	1	5	14	23
Female	2,335	-	1	2	1	1	1	3	2	1	12	28
White, both sexes	4,138	-	-	-	1	1	4	5	1	3	18	41
Male	1,959	-	-	-	1	1	3	2	1	3	10	15
Female	2,179	-	-	-	-	-	1	3	-	-	8	26
Black, both sexes	303	-	1	4	2	4	-	3	2	3	6	8
Male	162	-	-	2	1	3	-	3	-	2	3	6
Female	141	-	1	2	1	1	-	-	2	1	3	2
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	-	-	2	2
Male	14	-	-	-	-	-	-	-	-	-	1	2
Female	12	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Minnesota												
All races, both sexes	1,770	-	-	1	-	1	-	1	1	1	4	13
Male	872	-	-	1	-	1	-	-	-	-	3	7
Female	898	-	-	-	-	-	-	1	1	1	1	6
White, both sexes	1,728	-	-	-	-	-	-	-	1	-	3	12
Male	847	-	-	-	-	-	-	-	-	-	2	6
Female	881	-	-	-	-	-	-	-	1	-	1	6
Black, both sexes	22	-	-	1	-	1	-	1	-	1	1	1
Male	14	-	-	1	-	1	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,378	-	-	-	2	1	-	3	4	5	14	19
Male	735	-	-	-	2	1	-	2	1	1	7	7
Female	643	-	-	-	-	-	-	1	3	4	7	12
White, both sexes	1,122	-	-	-	-	-	-	1	-	1	7	11
Male	581	-	-	-	-	-	-	1	-	-	3	4
Female	541	-	-	-	-	-	-	-	-	1	4	7
Black, both sexes	253	-	-	-	2	1	-	2	4	4	7	8
Male	153	-	-	-	2	1	-	1	1	1	4	3
Female	100	-	-	-	-	-	-	1	3	3	3	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Missouri												
All races, both sexes	3,017	1	-	-	3	-	3	5	4	7	24	45
Male	1,441	-	-	-	2	-	3	3	3	6	14	25
Female	1,576	1	-	-	1	-	-	2	1	1	10	20
White, both sexes	2,834	1	-	-	1	-	2	3	3	4	18	40
Male	1,345	-	-	-	1	-	2	2	2	3	10	22
Female	1,489	1	-	-	-	-	-	1	1	1	8	18
Black, both sexes	171	-	-	-	1	-	1	2	1	3	5	5
Male	92	-	-	-	1	-	1	1	1	3	3	3
Female	79	-	-	-	-	-	-	1	-	-	2	2
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	1	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
Montana												
All races, both sexes	579	-	-	1	-	-	-	-	-	-	-	3
Male	260	-	-	-	-	-	-	-	-	-	-	1
Female	319	-	-	1	-	-	-	-	-	-	-	2
White, both sexes	558	-	-	1	-	-	-	-	-	-	-	3
Male	248	-	-	-	-	-	-	-	-	-	-	1
Female	310	-	-	1	-	-	-	-	-	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Nebraska												
All races, both sexes	878	-	-	-	-	1	-	-	2	1	1	5
Male	437	-	-	-	-	-	-	-	1	1	-	3
Female	441	-	-	-	-	1	-	-	1	-	1	2
White, both sexes	853	-	-	-	-	-	-	-	1	1	-	3
Male	423	-	-	-	-	-	-	-	-	1	-	2
Female	430	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	16	-	-	-	-	1	-	-	1	-	1	1
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	8	-	-	-	-	1	-	-	-	-	1	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,066	-	-	-	-	-	1	-	1	4	5	13
Male	486	-	-	-	-	-	1	-	-	1	3	4
Female	580	-	-	-	-	-	-	-	1	3	2	9
White, both sexes	1,020	-	-	-	-	-	1	-	1	3	4	12
Male	461	-	-	-	-	-	1	-	-	1	3	3
Female	559	-	-	-	-	-	-	-	1	2	1	9
Black, both sexes	28	-	-	-	-	-	-	-	-	1	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
New Hampshire												
All races, both sexes	603	-	-	-	-	-	-	-	1	2	1	3
Male	277	-	-	-	-	-	-	-	-	2	-	2
Female	326	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	602	-	-	-	-	-	-	-	1	2	1	3
Male	276	-	-	-	-	-	-	-	-	2	-	2
Female	326	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2,862	-	1	-	2	4	2	1	2	10	18	27
Male	1,231	-	1	-	2	2	2	-	1	3	6	10
Female	1,631	-	-	-	-	2	-	1	1	7	12	17
White, both sexes	2,569	-	-	-	1	-	1	1	1	6	12	20
Male	1,093	-	-	-	1	-	1	-	-	3	4	8
Female	1,476	-	-	-	-	-	-	1	1	3	8	12
Black, both sexes	261	-	1	-	1	3	1	-	1	4	5	7
Male	119	-	1	-	1	1	1	-	1	-	2	2
Female	142	-	-	-	-	2	-	-	-	4	3	5
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	1	-	-	-	-	1	-
Male	17	-	-	-	-	1	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
New Mexico												
All races, both sexes	888	-	-	-	2	-	-	1	-	1	5	9
Male	434	-	-	-	2	-	-	1	-	1	-	5
Female	454	-	-	-	-	-	-	-	-	-	5	4
White, both sexes	866	-	-	-	2	-	-	1	-	1	4	8
Male	422	-	-	-	2	-	-	1	-	1	-	4
Female	444	-	-	-	-	-	-	-	-	-	4	4
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6,329	1	3	3	4	3	6	11	9	10	37	63
Male	2,886	1	1	3	2	3	2	3	3	6	22	33
Female	3,443	-	2	-	2	-	4	8	6	4	15	30
White, both sexes	5,706	1	-	2	-	1	5	4	5	7	24	47
Male	2,568	1	-	2	-	1	2	2	1	4	16	26
Female	3,138	-	-	-	-	-	3	2	4	3	8	21
Black, both sexes	528	-	2	1	4	2	1	7	4	3	13	16
Male	247	-	1	1	2	2	-	1	2	2	6	7
Female	281	-	1	-	2	-	1	6	2	1	7	9
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	79	-	1	-	-	-	-	-	-	-	-	-
Male	62	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
North Carolina												
All races, both sexes	4,017	1	-	1	1	4	3	5	7	9	18	48
Male	1,912	1	-	1	1	1	-	2	3	5	7	21
Female	2,105	-	-	-	-	3	3	3	4	4	11	27
White, both sexes	3,581	-	-	-	-	2	1	3	4	5	8	36
Male	1,667	-	-	-	-	-	-	1	2	3	2	15
Female	1,914	-	-	-	-	2	1	2	2	2	6	21
Black, both sexes	412	1	-	1	1	2	2	2	3	4	10	10
Male	231	1	-	1	1	1	-	1	1	2	5	4
Female	181	-	-	-	-	1	2	1	2	2	5	6
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	283	-	-	-	-	-	1	-	-	-	-	1
Male	158	-	-	-	-	-	1	-	-	-	-	1
Female	125	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	274	-	-	-	-	-	1	-	-	-	-	1
Male	156	-	-	-	-	-	1	-	-	-	-	1
Female	118	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Ohio												
All races, both sexes	6,054	1	2	2	1	1	3	4	6	13	31	80
Male	2,903	1	1	1	1	-	2	3	4	8	14	38
Female	3,151	-	1	1	-	1	1	1	2	5	17	42
White, both sexes	5,595	1	2	-	-	1	2	2	3	9	26	61
Male	2,650	1	1	-	-	-	1	2	2	5	13	29
Female	2,945	-	1	-	-	1	1	-	1	4	13	32
Black, both sexes	448	-	-	2	1	-	1	2	3	4	5	18
Male	248	-	-	1	1	-	1	1	2	3	1	9
Female	200	-	-	1	-	-	-	1	1	1	4	9
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2,192	2	2	1	4	2	3	1	1	6	12	23
Male	1,100	2	2	1	2	2	2	1	1	2	4	11
Female	1,092	-	-	-	2	-	1	-	-	4	8	12
White, both sexes	2,028	1	2	1	1	-	1	1	-	6	9	19
Male	1,017	1	2	1	1	-	-	1	-	2	2	10
Female	1,011	-	-	-	-	-	1	-	-	4	7	9
Black, both sexes	66	-	-	-	3	2	1	-	1	-	-	1
Male	34	-	-	-	1	2	1	-	1	-	-	-
Female	32	-	-	-	2	-	-	-	-	-	-	1
American Indian 1/, both sexes	96	1	-	-	-	-	-	-	-	-	3	3
Male	48	1	-	-	-	-	-	-	-	-	2	1
Female	48	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Oregon												
All races, both sexes	1,828	1	1	-	-	1	1	-	2	2	5	24
Male	882	-	-	-	-	1	1	-	2	2	4	15
Female	946	1	1	-	-	-	-	-	-	-	1	9
White, both sexes	1,785	1	1	-	-	1	1	-	2	2	5	23
Male	859	-	-	-	-	1	1	-	2	2	4	15
Female	926	1	1	-	-	-	-	-	-	-	1	8
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	5,621	1	1	2	2	1	1	4	3	8	21	53
Male	2,616	-	1	1	-	-	-	4	1	2	14	26
Female	3,005	1	-	1	2	1	1	-	2	6	7	27
White, both sexes	5,244	1	1	2	2	-	-	1	1	3	15	43
Male	2,435	-	1	1	-	-	-	1	1	2	9	20
Female	2,809	1	-	1	2	-	-	-	-	1	6	23
Black, both sexes	359	-	-	-	-	1	1	3	2	5	6	10
Male	168	-	-	-	-	-	-	3	-	-	5	6
Female	191	-	-	-	-	1	1	-	2	5	1	4
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Rhode Island												
All races, both sexes	485	-	-	1	-	-	-	-	-	-	2	3
Male	202	-	-	1	-	-	-	-	-	-	1	-
Female	283	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	472	-	-	1	-	-	-	-	-	-	2	3
Male	195	-	-	1	-	-	-	-	-	-	1	-
Female	277	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1,935	-	-	1	2	-	2	2	-	3	9	22
Male	968	-	-	1	1	-	-	1	-	2	6	12
Female	967	-	-	-	1	-	2	1	-	1	3	10
White, both sexes	1,677	-	-	-	-	-	1	2	-	1	5	14
Male	821	-	-	-	-	-	-	1	-	1	4	7
Female	856	-	-	-	-	-	1	1	-	-	1	7
Black, both sexes	252	-	-	1	2	-	1	-	-	2	4	8
Male	142	-	-	1	1	-	-	-	-	1	2	5
Female	110	-	-	-	1	-	1	-	-	1	2	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
South Dakota												
All races, both sexes	376	-	-	-	-	-	1	-	1	-	-	2
Male	204	-	-	-	-	-	1	-	1	-	-	1
Female	172	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	362	-	-	-	-	-	1	-	1	-	-	2
Male	194	-	-	-	-	-	1	-	1	-	-	1
Female	168	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2,980	-	1	3	3	2	6	2	2	2	25	46
Male	1,455	-	-	3	3	1	4	1	2	2	14	22
Female	1,525	-	1	-	-	1	2	1	-	-	11	24
White, both sexes	2,762	-	1	1	1	-	3	2	1	1	22	41
Male	1,332	-	-	1	1	-	2	1	1	1	12	18
Female	1,430	-	1	-	-	-	1	1	-	-	10	23
Black, both sexes	216	-	-	2	2	2	3	-	1	1	3	5
Male	123	-	-	2	2	1	2	-	1	1	2	4
Female	93	-	-	-	-	1	1	-	-	-	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Texas												
All races, both sexes	7,639	6	5	5	1	5	8	3	10	11	38	81
Male	3,667	2	4	3	1	4	6	-	4	4	16	37
Female	3,972	4	1	2	-	1	2	3	6	7	22	44
White, both sexes	7,088	2	3	3	-	3	2	2	8	10	30	65
Male	3,384	2	3	2	-	2	1	-	3	4	12	31
Female	3,704	-	-	1	-	1	1	2	5	6	18	34
Black, both sexes	498	4	1	2	1	2	3	1	2	1	8	12
Male	257	-	1	1	1	2	2	-	1	-	4	5
Female	241	4	-	1	-	-	1	1	1	1	4	7
American Indian 1/, both sexes	13	-	-	-	-	-	1	-	-	-	-	2
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	40	-	1	-	-	-	2	-	-	-	-	2
Male	20	-	-	-	-	-	2	-	-	-	-	1
Female	20	-	1	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	591	2	1	-	-	1	-	-	1	-	3	8
Male	310	1	1	-	-	-	-	-	-	-	1	3
Female	281	1	-	-	-	1	-	-	1	-	2	5
White, both sexes	581	2	1	-	-	1	-	-	1	-	3	8
Male	304	1	1	-	-	-	-	-	-	-	1	3
Female	277	1	-	-	-	1	-	-	1	-	2	5
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Vermont												
All races, both sexes	320	-	-	-	-	-	-	-	-	1	-	2
Male	142	-	-	-	-	-	-	-	-	-	-	-
Female	178	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	320	-	-	-	-	-	-	-	-	1	-	2
Male	142	-	-	-	-	-	-	-	-	-	-	-
Female	178	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2,697	1	2	2	-	1	1	1	-	5	7	42
Male	1,245	1	1	1	-	1	1	1	-	2	4	19
Female	1,452	-	1	1	-	-	-	-	-	3	3	23
White, both sexes	2,415	-	2	-	-	-	1	1	-	2	5	37
Male	1,089	-	1	-	-	-	1	1	-	-	3	16
Female	1,326	-	1	-	-	-	-	-	-	2	2	21
Black, both sexes	266	-	-	2	-	1	-	-	-	3	2	4
Male	145	-	-	1	-	1	-	-	-	2	1	3
Female	121	-	-	1	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	14	1	-	-	-	-	-	-	-	-	-	-
Male	10	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Washington												
All races, both sexes	2,662	1	-	1	-	-	-	1	-	2	9	19
Male	1,205	1	-	-	-	-	-	1	-	1	1	11
Female	1,457	-	-	1	-	-	-	-	-	1	8	8
White, both sexes	2,550	-	-	-	-	-	-	1	-	1	9	18
Male	1,151	-	-	-	-	-	-	1	-	-	1	10
Female	1,399	-	-	-	-	-	-	-	-	1	8	8
Black, both sexes	35	1	-	1	-	-	-	-	-	1	-	-
Male	19	1	-	-	-	-	-	-	-	1	-	-
Female	16	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	-	-	-	-	-	-	-	-	-	-	1
Male	29	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1,267	-	-	-	-	1	1	1	3	1	11	14
Male	594	-	-	-	-	-	-	1	1	-	6	7
Female	673	-	-	-	-	1	1	-	2	1	5	7
White, both sexes	1,249	-	-	-	-	1	1	1	3	-	10	14
Male	585	-	-	-	-	-	-	1	1	-	6	7
Female	664	-	-	-	-	1	1	-	2	-	4	7
Black, both sexes	18	-	-	-	-	-	-	-	-	1	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Wisconsin												
All races, both sexes	2,354	-	1	-	2	1	1	1	3	1	12	11
Male	1,139	-	1	-	1	1	1	1	1	1	6	9
Female	1,215	-	-	-	1	-	-	-	2	-	6	2
White, both sexes	2,277	-	1	-	2	-	-	-	2	1	10	10
Male	1,099	-	1	-	1	-	-	-	-	1	4	8
Female	1,178	-	-	-	1	-	-	-	2	-	6	2
Black, both sexes	55	-	-	-	-	1	1	1	1	-	1	-
Male	30	-	-	-	-	1	1	1	1	-	1	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	345	-	-	-	-	-	-	1	-	1	-	5
Male	178	-	-	-	-	-	-	-	-	1	-	3
Female	167	-	-	-	-	-	-	1	-	-	-	2
White, both sexes	339	-	-	-	-	-	-	1	-	1	-	5
Male	175	-	-	-	-	-	-	-	-	1	-	3
Female	164	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
United States												
All races, both sexes	2,604	4,699	7,676	11,601	16,635	22,770	24,636	18,703	9,363	2,738	399	4
Male	1,341	2,427	3,955	5,799	8,353	11,371	11,796	8,231	3,658	843	113	3
Female	1,263	2,272	3,721	5,802	8,282	11,399	12,840	10,472	5,705	1,895	286	1
White, both sexes	2,160	4,100	6,900	10,609	15,446	21,241	23,149	17,692	8,841	2,534	358	3
Male	1,106	2,103	3,492	5,231	7,685	10,527	10,959	7,708	3,423	760	98	2
Female	1,054	1,997	3,408	5,378	7,761	10,714	12,190	9,984	5,418	1,774	260	1
Black, both sexes	405	534	671	839	979	1,193	1,110	745	360	138	36	-
Male	216	293	397	478	550	643	610	370	142	50	11	-
Female	189	241	274	361	429	550	500	375	218	88	25	-
American Indian 1/, both sexes	20	31	55	61	76	94	72	44	16	7	1	-
Male	8	12	32	28	34	42	31	21	5	1	1	-
Female	12	19	23	33	42	52	41	23	11	6	-	-
Asian or Pacific Islander 2/, both sexes	19	34	50	92	134	242	305	222	146	59	4	1
Male	11	19	34	62	84	159	196	132	88	32	3	1
Female	8	15	16	30	50	83	109	90	58	27	1	-
Alabama												
All races, both sexes	53	102	170	258	366	421	435	288	114	45	4	-
Male	25	60	85	140	190	219	208	140	48	11	1	-
Female	28	42	85	118	176	202	227	148	66	34	3	-
White, both sexes	44	80	154	227	336	387	402	259	97	41	4	-
Male	21	45	72	119	168	195	186	124	38	8	1	-
Female	23	35	82	108	168	192	216	135	59	33	3	-
Black, both sexes	9	22	16	31	30	33	32	29	17	3	-	-
Male	4	15	13	21	22	23	22	16	10	3	-	-
Female	5	7	3	10	8	10	10	13	7	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Alaska												
All races, both sexes	4	11	14	15	24	22	21	17	8	1	-	-
Male	2	5	10	6	8	10	6	7	4	-	-	-
Female	2	6	4	9	16	12	15	10	4	1	-	-
White, both sexes	3	8	9	8	20	18	14	13	4	-	-	-
Male	1	3	7	3	8	9	4	4	2	-	-	-
Female	2	5	2	5	12	9	10	9	2	-	-	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	3	4	5	3	3	6	1	4	1	-	-
Male	1	2	2	2	-	1	2	1	2	-	-	-
Female	-	1	2	3	3	2	4	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	1	1	2	-	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	1	1	1	-	-	-	-
Arizona												
All races, both sexes	50	105	152	271	388	563	566	430	183	45	5	-
Male	24	58	79	143	191	297	271	186	71	15	2	-
Female	26	47	73	128	197	266	295	244	112	30	3	-
White, both sexes	49	102	148	260	384	553	562	424	179	44	5	-
Male	24	56	78	137	190	292	268	183	70	14	2	-
Female	25	46	70	123	194	261	294	241	109	30	3	-
Black, both sexes	1	2	2	7	2	6	3	3	3	-	-	-
Male	-	2	1	4	1	2	3	1	-	-	-	-
Female	1	-	1	3	1	4	-	2	3	-	-	-
American Indian 1/, both sexes	-	1	2	3	2	2	1	2	-	-	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	1	2	2	2	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	-	1	1	1	-	-
Male	-	-	-	1	-	2	-	1	1	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Arkansas												
All races, both sexes	40	65	129	163	212	267	271	197	96	26	3	-
Male	12	41	70	81	129	146	145	99	38	9	1	-
Female	28	24	59	82	83	121	126	98	58	17	2	-
White, both sexes	34	58	117	157	205	255	256	190	93	23	3	-
Male	9	38	63	77	125	138	137	97	36	9	1	-
Female	25	20	54	80	80	117	119	93	57	14	2	-
Black, both sexes	5	7	12	6	7	10	14	7	3	3	-	-
Male	3	3	7	4	4	8	8	2	2	-	-	-
Female	2	4	5	2	3	2	6	5	1	3	-	-
American Indian 1/, both sexes	1	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	256	422	714	1,053	1,537	2,242	2,671	2,146	1,134	329	55	-
Male	134	233	393	513	756	1,075	1,231	899	436	119	21	-
Female	122	189	321	540	781	1,167	1,440	1,247	698	210	34	-
White, both sexes	205	351	600	938	1,377	2,002	2,405	1,953	1,024	276	50	-
Male	109	188	313	441	659	932	1,075	796	379	94	17	-
Female	96	163	287	497	718	1,070	1,330	1,157	645	182	33	-
Black, both sexes	43	54	79	71	84	106	97	58	31	15	3	-
Male	21	34	51	45	43	57	47	29	11	5	2	-
Female	22	20	28	26	41	49	50	29	20	10	1	-
American Indian 1/, both sexes	4	1	7	6	8	11	10	7	2	1	-	-
Male	1	1	6	3	7	6	4	1	-	-	-	-
Female	3	-	1	3	1	5	6	6	2	1	-	-
Asian or Pacific Islander 2/, both sexes	4	16	28	38	68	123	159	128	77	37	2	-
Male	3	10	23	24	47	80	105	73	46	20	2	-
Female	1	6	5	14	21	43	54	55	31	17	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Colorado												
All races, both sexes	47	75	111	190	262	352	393	269	155	40	7	-
Male	25	49	69	112	131	166	187	133	73	12	-	-
Female	22	26	42	78	131	186	206	136	82	28	7	-
White, both sexes	43	67	106	180	251	342	384	263	153	40	6	-
Male	23	45	66	108	125	163	183	131	73	12	-	-
Female	20	22	40	72	126	179	201	132	80	28	6	-
Black, both sexes	4	7	5	8	10	7	3	5	1	-	1	-
Male	2	4	3	3	6	3	1	2	-	-	-	-
Female	2	3	2	5	4	4	2	3	1	-	1	-
American Indian 1/, both sexes	-	1	-	1	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	6	-	1	-	-	-
Male	-	-	-	1	-	-	3	-	-	-	-	-
Female	-	-	-	-	1	1	3	-	1	-	-	-
Connecticut												
All races, both sexes	32	41	71	94	143	243	289	284	177	55	5	1
Male	18	16	32	53	75	118	135	111	66	12	2	-
Female	14	25	39	41	68	125	154	173	111	43	3	1
White, both sexes	29	37	68	86	139	234	281	281	173	54	4	1
Male	16	14	31	47	73	113	129	111	63	12	2	-
Female	13	23	37	39	66	121	152	170	110	42	2	1
Black, both sexes	2	4	3	8	4	9	8	3	3	1	1	-
Male	1	2	1	6	2	5	6	-	2	-	-	-
Female	1	2	2	2	2	4	2	3	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Delaware												
All races, both sexes	7	10	13	35	55	75	73	43	19	9	2	-
Male	3	3	8	19	28	37	31	14	9	5	-	-
Female	4	7	5	16	27	38	42	29	10	4	2	-
White, both sexes	6	9	11	28	48	68	65	38	19	8	2	-
Male	3	2	6	15	24	33	27	11	9	4	-	-
Female	3	7	5	13	24	35	38	27	10	4	2	-
Black, both sexes	1	1	2	6	6	7	7	4	-	1	-	-
Male	-	1	2	3	3	4	4	3	-	1	-	-
Female	1	-	-	3	3	3	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	6	5	10	9	25	29	15	12	1	1	-
Male	2	3	1	7	2	13	13	5	8	-	-	-
Female	-	3	4	3	7	12	16	10	4	1	1	-
White, both sexes	-	-	1	2	-	10	9	11	6	1	1	-
Male	-	-	1	2	-	6	2	2	4	-	-	-
Female	-	-	-	-	-	4	7	9	2	1	1	-
Black, both sexes	2	6	4	8	9	14	20	4	6	-	-	-
Male	2	3	-	5	2	6	11	3	4	-	-	-
Female	-	3	4	3	7	8	9	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Florida												
All races, both sexes	185	302	486	779	1,140	1,575	1,820	1,478	712	220	35	-
Male	96	163	265	408	568	790	858	614	272	65	8	-
Female	89	139	221	371	572	785	962	864	440	155	27	-
White, both sexes	162	271	450	740	1,077	1,503	1,737	1,427	686	211	34	-
Male	85	147	241	384	534	743	807	585	260	62	8	-
Female	77	124	209	356	543	760	930	842	426	149	26	-
Black, both sexes	22	31	35	36	59	67	73	46	23	9	1	-
Male	11	16	23	22	32	44	44	25	11	3	-	-
Female	11	15	12	14	27	23	29	21	12	6	1	-
American Indian 1/, both sexes	-	-	-	2	2	3	2	3	-	-	-	-
Male	-	-	-	1	1	1	1	2	-	-	-	-
Female	-	-	-	1	1	2	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	2	2	8	2	3	-	-	-
Male	-	-	1	1	1	2	6	2	1	-	-	-
Female	1	-	-	-	1	-	2	-	2	-	-	-
Georgia												
All races, both sexes	96	175	257	361	517	590	591	439	190	61	6	-
Male	46	91	137	180	267	307	264	212	61	23	-	-
Female	50	84	120	181	250	283	327	227	129	38	6	-
White, both sexes	72	138	232	322	461	530	520	399	174	50	4	-
Male	34	67	118	159	231	272	227	190	50	17	-	-
Female	38	71	114	163	230	258	293	209	124	33	4	-
Black, both sexes	24	37	25	38	53	58	62	36	14	9	2	-
Male	12	24	19	20	34	35	32	21	10	5	-	-
Female	12	13	6	18	19	23	30	15	4	4	2	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	9	4	2	1	-	-
Male	-	-	-	1	1	-	5	1	1	-	-	-
Female	-	-	-	-	1	1	4	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Hawaii												
All races, both sexes	7	12	10	27	34	49	62	44	29	10	2	-
Male	4	8	4	14	17	28	38	25	13	7	-	-
Female	3	4	6	13	17	21	24	19	16	3	2	-
White, both sexes	2	3	3	9	15	21	24	13	10	2	1	-
Male	1	1	2	4	8	10	14	7	5	-	-	-
Female	1	2	1	5	7	11	10	6	5	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	9	7	18	19	27	38	31	19	7	1	-
Male	3	7	2	10	9	17	24	18	8	6	-	-
Female	2	2	5	8	10	10	14	13	11	1	1	-
Idaho												
All races, both sexes	9	22	33	54	83	142	123	116	45	9	1	-
Male	6	9	21	36	39	76	72	54	16	3	1	-
Female	3	13	12	18	44	66	51	62	29	6	-	-
White, both sexes	9	22	32	54	83	141	123	114	45	9	1	-
Male	6	9	20	36	39	75	72	53	16	3	1	-
Female	3	13	12	18	44	66	51	61	29	6	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Illinois												
All races, both sexes	112	165	276	421	604	821	949	740	391	104	14	-
Male	50	84	141	203	308	409	460	332	148	27	1	-
Female	62	81	135	218	296	412	489	408	243	77	13	-
White, both sexes	74	133	238	384	547	742	887	700	366	96	13	-
Male	32	69	119	186	272	367	423	312	138	24	-	-
Female	42	64	119	198	275	375	464	388	228	72	13	-
Black, both sexes	37	31	38	36	54	69	59	35	19	7	1	-
Male	18	15	22	17	34	36	34	16	7	3	1	-
Female	19	16	16	19	20	33	25	19	12	4	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	2	10	3	5	6	1	-	-
Male	-	-	-	-	2	6	3	4	3	-	-	-
Female	1	-	-	1	-	4	-	1	3	1	-	-
Indiana												
All races, both sexes	63	132	212	335	473	587	615	464	254	62	7	-
Male	35	67	107	171	248	296	287	207	110	21	1	-
Female	28	65	105	164	225	291	328	257	144	41	6	-
White, both sexes	55	125	201	321	453	564	594	448	250	62	6	-
Male	30	64	102	167	239	285	276	196	109	21	-	-
Female	25	61	99	154	214	279	318	252	141	41	6	-
Black, both sexes	8	6	11	13	20	22	21	16	3	-	1	-
Male	5	3	5	3	9	11	11	11	1	-	1	-
Female	3	3	6	10	11	11	10	5	2	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Iowa												
All races, both sexes	30	56	81	150	225	267	360	281	118	52	8	-
Male	12	29	43	77	126	147	201	140	54	24	2	-
Female	18	27	38	73	99	120	159	141	64	28	6	-
White, both sexes	28	56	81	150	222	265	359	280	117	52	8	-
Male	11	29	43	77	124	146	201	140	53	24	2	-
Female	17	27	38	73	98	119	158	140	64	28	6	-
Black, both sexes	1	-	-	-	2	1	-	1	-	-	-	-
Male	1	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Kansas												
All races, both sexes	24	51	81	128	201	295	282	232	121	35	8	-
Male	15	26	30	67	108	153	142	114	67	13	1	-
Female	9	25	51	61	93	142	140	118	54	22	7	-
White, both sexes	23	49	78	120	191	281	265	228	118	33	7	-
Male	15	26	29	63	102	145	134	114	67	13	1	-
Female	8	23	49	57	89	136	131	114	51	20	6	-
Black, both sexes	-	1	2	4	5	11	11	2	2	2	1	-
Male	-	-	-	2	3	7	6	-	-	-	-	-
Female	-	1	2	2	2	4	5	2	2	2	1	-
American Indian 1/, both sexes	-	-	1	4	4	2	6	2	-	-	-	-
Male	-	-	1	2	3	-	2	-	-	-	-	-
Female	-	-	-	2	1	2	4	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	1	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Kentucky												
All races, both sexes	71	101	200	292	358	405	419	291	152	37	4	-
Male	36	56	101	149	180	198	209	116	61	12	2	-
Female	35	45	99	143	178	207	210	175	91	25	2	-
White, both sexes	64	92	192	280	341	387	401	284	149	34	4	-
Male	32	51	97	143	171	190	200	113	60	12	2	-
Female	32	41	95	137	170	197	201	171	89	22	2	-
Black, both sexes	7	9	6	11	17	16	17	7	3	3	-	-
Male	4	5	3	5	9	7	8	3	1	-	-	-
Female	3	4	3	6	8	9	9	4	2	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	1	-	-	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	47	78	118	207	253	287	281	227	98	37	8	-
Male	27	44	59	111	141	147	145	107	40	15	2	-
Female	20	34	59	96	112	140	136	120	58	22	6	-
White, both sexes	27	61	91	166	210	244	249	202	84	31	5	-
Male	13	33	48	84	111	126	125	90	33	10	1	-
Female	14	28	43	82	99	118	124	112	51	21	4	-
Black, both sexes	20	17	26	41	43	42	31	23	14	6	3	-
Male	14	11	10	27	30	21	19	15	7	5	1	-
Female	6	6	16	14	13	21	12	8	7	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	1	2	-	-	-	-
Male	-	-	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Maine												
All races, both sexes	18	32	42	78	110	137	153	120	61	17	2	-
Male	12	17	27	32	58	73	75	59	24	3	1	-
Female	6	15	15	46	52	64	78	61	37	14	1	-
White, both sexes	18	32	41	78	110	137	152	120	61	17	2	-
Male	12	17	27	32	58	73	75	59	24	3	1	-
Female	6	15	14	46	52	64	77	61	37	14	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	35	59	106	165	247	317	372	298	141	46	5	-
Male	16	26	50	72	106	158	161	119	61	8	2	-
Female	19	33	56	93	141	159	211	179	80	38	3	-
White, both sexes	20	51	84	133	215	277	323	261	124	45	5	-
Male	8	24	39	60	86	139	137	107	54	8	2	-
Female	12	27	45	73	129	138	186	154	70	37	3	-
Black, both sexes	15	8	22	29	30	38	42	36	14	1	-	-
Male	8	2	11	12	19	18	20	11	5	-	-	-
Female	7	6	11	17	11	20	22	25	9	1	-	-
American Indian 1/, both sexes	-	-	-	2	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	7	-	3	-	-	-
Male	-	-	-	-	1	-	4	-	2	-	-	-
Female	-	-	-	1	1	1	3	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Massachusetts												
All races, both sexes	32	83	139	197	296	472	481	454	249	84	12	-
Male	17	39	61	93	153	217	201	176	73	19	3	-
Female	15	44	78	104	143	255	280	278	176	65	9	-
White, both sexes	30	80	137	191	286	460	471	445	243	79	11	-
Male	15	37	60	88	147	207	196	173	70	19	3	-
Female	15	43	77	103	139	253	275	272	173	60	8	-
Black, both sexes	2	2	2	2	8	7	8	7	4	4	1	-
Male	2	1	1	1	4	6	3	1	1	-	-	-
Female	-	1	1	1	4	1	5	6	3	4	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	4	2	5	2	2	2	1	-	-
Male	-	-	-	4	2	4	2	2	2	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Michigan												
All races, both sexes	88	164	255	412	604	870	909	663	317	66	15	-
Male	46	95	126	201	280	419	449	311	126	18	9	-
Female	42	69	129	211	324	451	460	352	191	48	6	-
White, both sexes	79	141	225	384	565	805	865	621	305	63	11	-
Male	40	80	112	183	264	386	424	289	121	17	7	-
Female	39	61	113	201	301	419	441	332	184	46	4	-
Black, both sexes	9	23	30	20	34	55	41	39	12	3	4	-
Male	6	15	14	12	13	30	23	21	5	1	2	-
Female	3	8	16	8	21	25	18	18	7	2	2	-
American Indian 1/, both sexes	-	-	-	6	4	7	2	3	-	-	-	-
Male	-	-	-	4	3	2	1	1	-	-	-	-
Female	-	-	-	2	1	5	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	1	3	1	-	-	-	-	-
Male	-	-	-	2	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Minnesota												
All races, both sexes	22	39	90	139	217	303	405	314	165	47	6	1
Male	10	22	49	71	110	173	198	132	77	16	1	1
Female	12	17	41	68	107	130	207	182	88	31	5	-
White, both sexes	21	38	86	136	213	298	397	311	161	45	6	-
Male	9	21	47	70	107	170	195	130	74	15	1	-
Female	12	17	39	66	106	128	202	181	87	30	5	-
Black, both sexes	-	-	3	2	2	3	4	2	-	-	-	-
Male	-	-	2	1	2	2	2	1	-	-	-	-
Female	-	-	1	1	-	1	2	1	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	2	2	2	-	-	-	-	-
Male	1	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	1	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	2	1	4	2	-	1
Male	-	1	-	-	-	-	-	1	3	1	-	1
Female	-	-	-	-	-	-	2	-	1	1	-	-
Mississippi												
All races, both sexes	34	69	107	143	189	259	233	185	73	33	4	1
Male	14	40	58	84	106	156	127	92	27	9	-	1
Female	20	29	49	59	83	103	106	93	46	24	4	-
White, both sexes	24	50	86	118	157	217	206	156	62	23	2	1
Male	8	28	44	68	83	130	109	76	20	6	-	1
Female	16	22	42	50	74	87	97	80	42	17	2	-
Black, both sexes	10	19	21	24	32	42	27	28	10	10	2	-
Male	6	12	14	16	23	26	18	15	7	3	-	-
Female	4	7	7	8	9	16	9	13	3	7	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Missouri												
All races, both sexes	76	129	189	288	406	579	594	389	203	64	8	-
Male	39	62	81	134	210	283	293	179	86	15	3	-
Female	37	67	108	154	196	296	301	210	117	49	5	-
White, both sexes	69	122	175	270	381	555	557	368	198	60	7	-
Male	35	57	76	126	197	272	272	167	84	14	3	-
Female	34	65	99	144	184	283	285	201	114	46	4	-
Black, both sexes	6	7	13	17	25	20	35	21	4	4	1	-
Male	4	5	4	8	13	9	21	12	2	1	-	-
Female	2	2	9	9	12	11	14	9	2	3	1	-
American Indian 1/, both sexes	1	-	1	1	-	3	1	-	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	-	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	12	16	38	46	87	113	116	98	37	11	1	-
Male	6	9	19	20	45	58	42	45	13	2	-	-
Female	6	7	19	26	42	55	74	53	24	9	1	-
White, both sexes	9	12	34	45	83	112	116	94	37	11	1	-
Male	5	8	16	19	44	57	42	41	13	2	-	-
Female	4	4	18	26	39	55	74	53	24	9	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	4	2	1	4	1	-	4	-	-	-	-
Male	1	1	2	1	1	1	-	4	-	-	-	-
Female	2	3	-	-	3	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Nebraska												
All races, both sexes	12	19	38	62	123	158	195	136	90	28	7	-
Male	7	9	17	29	68	92	98	71	31	6	4	-
Female	5	10	21	33	55	66	97	65	59	22	3	-
White, both sexes	11	19	38	59	118	155	192	134	88	28	6	-
Male	6	9	17	27	65	89	97	70	31	6	3	-
Female	5	10	21	32	53	66	95	64	57	22	3	-
Black, both sexes	1	-	-	2	2	1	2	1	2	-	1	-
Male	1	-	-	1	2	1	1	-	-	-	1	-
Female	-	-	-	1	-	-	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	2	-	-	-	-	-	-
Male	-	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Nevada												
All races, both sexes	12	44	85	102	198	188	198	132	61	19	3	-
Male	4	24	35	46	100	84	97	57	24	5	1	-
Female	8	20	50	56	98	104	101	75	37	14	2	-
White, both sexes	11	38	81	98	190	181	192	128	59	18	3	-
Male	4	22	32	43	95	80	94	55	22	5	1	-
Female	7	16	49	55	95	101	98	73	37	13	2	-
Black, both sexes	1	4	3	4	6	2	4	2	-	-	-	-
Male	-	1	3	3	4	-	2	1	-	-	-	-
Female	1	3	-	1	2	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	-	2	5	2	1	1	1	-	-
Male	-	1	-	-	1	4	1	1	1	-	-	-
Female	-	1	1	-	1	1	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
New Hampshire												
All races, both sexes	18	27	30	57	86	104	122	104	36	10	2	-
Male	9	16	16	24	37	56	65	36	11	3	-	-
Female	9	11	14	33	49	48	57	68	25	7	2	-
White, both sexes	18	27	30	57	86	103	122	104	36	10	2	-
Male	9	16	16	24	37	55	65	36	11	3	-	-
Female	9	11	14	33	49	48	57	68	25	7	2	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	65	77	131	196	329	538	613	507	245	83	11	-
Male	34	39	64	102	139	240	276	202	82	24	2	-
Female	31	38	67	94	190	298	337	305	163	59	9	-
White, both sexes	57	60	107	164	292	488	572	480	222	78	7	-
Male	30	32	48	82	120	217	254	194	74	24	1	-
Female	27	28	59	82	172	271	318	286	148	54	6	-
Black, both sexes	8	17	22	30	32	43	35	24	18	5	4	-
Male	4	7	14	18	15	20	19	8	4	-	1	-
Female	4	10	8	12	17	23	16	16	14	5	3	-
American Indian 1/, both sexes	-	-	2	-	1	1	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	4	6	6	3	5	-	-	-
Male	-	-	-	2	4	3	3	-	4	-	-	-
Female	-	-	-	-	-	3	3	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
New Mexico												
All races, both sexes	13	24	63	67	107	174	211	117	70	18	6	-
Male	4	9	30	40	48	90	101	65	27	8	3	-
Female	9	15	33	27	59	84	110	52	43	10	3	-
White, both sexes	13	23	60	66	106	172	204	115	67	18	6	-
Male	4	8	29	39	48	88	98	63	26	8	3	-
Female	9	15	31	27	58	84	106	52	41	10	3	-
Black, both sexes	-	1	-	-	-	2	1	1	2	-	-	-
Male	-	1	-	-	-	2	1	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	3	1	1	-	5	1	1	-	-	-
Male	-	-	1	1	-	-	2	1	-	-	-	-
Female	-	-	2	-	1	-	3	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
New York												
All races, both sexes	115	188	368	535	700	1,122	1,276	1,061	583	207	24	-
Male	59	96	203	257	321	541	620	449	196	57	8	-
Female	56	92	165	278	379	581	656	612	387	150	16	-
White, both sexes	88	156	321	467	635	1,032	1,172	986	538	193	22	-
Male	45	81	174	218	291	494	562	414	176	51	7	-
Female	43	75	147	249	344	538	610	572	362	142	15	-
Black, both sexes	26	30	44	62	54	74	84	55	35	10	1	-
Male	13	14	26	34	24	36	42	20	11	3	-	-
Female	13	16	18	28	30	38	42	35	24	7	1	-
American Indian 1/, both sexes	-	1	1	1	3	7	1	2	-	-	-	-
Male	-	1	1	-	1	3	1	2	-	-	-	-
Female	-	-	-	1	2	4	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	5	8	9	19	18	10	4	1	-
Male	1	-	2	5	5	8	15	13	9	3	1	-
Female	-	1	-	-	3	1	4	5	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
North Carolina												
All races, both sexes	79	191	304	445	559	749	782	499	239	66	7	-
Male	46	93	160	230	270	370	360	217	95	26	3	-
Female	33	98	144	215	289	379	422	282	144	40	4	-
White, both sexes	64	164	248	400	510	687	712	451	222	58	6	-
Male	39	74	125	204	244	339	320	191	86	20	2	-
Female	25	90	123	196	266	348	392	260	136	38	4	-
Black, both sexes	14	26	52	42	46	60	67	45	16	8	-	-
Male	6	19	34	24	24	30	38	25	8	6	-	-
Female	8	7	18	18	22	30	29	20	8	2	-	-
American Indian 1/, both sexes	1	1	2	3	2	1	2	3	1	-	1	-
Male	1	-	1	2	1	-	1	1	1	-	1	-
Female	-	1	1	1	1	1	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	6	10	15	32	60	51	53	39	11	2	-
Male	1	2	5	10	17	34	35	27	21	3	1	-
Female	1	4	5	5	15	26	16	26	18	8	1	-
White, both sexes	1	6	8	14	31	58	50	52	39	11	2	-
Male	1	2	4	10	17	34	34	27	21	3	1	-
Female	-	4	4	4	14	24	16	25	18	8	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	2	1	1	2	1	1	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	1	1	1	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Ohio												
All races, both sexes	139	234	385	572	809	1,167	1,163	874	427	120	20	-
Male	81	118	207	276	397	581	565	401	165	33	6	-
Female	58	116	178	296	412	586	598	473	262	87	14	-
White, both sexes	116	196	354	521	760	1,072	1,097	829	408	116	19	-
Male	68	96	186	245	369	531	528	374	160	33	6	-
Female	48	100	168	276	391	541	569	455	248	83	13	-
Black, both sexes	23	38	30	51	48	95	62	42	18	4	1	-
Male	13	22	21	31	28	50	34	26	4	-	-	-
Female	10	16	9	20	20	45	28	16	14	4	1	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	3	1	-	-	-
Male	-	-	-	-	-	-	3	1	1	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Oklahoma												
All races, both sexes	59	114	178	267	352	415	357	238	115	36	4	-
Male	28	64	86	127	190	217	186	106	51	14	1	-
Female	31	50	92	140	162	198	171	132	64	22	3	-
White, both sexes	54	106	162	245	327	388	331	228	110	33	3	-
Male	26	59	76	118	182	199	172	101	50	13	1	-
Female	28	47	86	127	145	189	159	127	60	20	2	-
Black, both sexes	3	4	4	11	8	7	10	6	3	1	1	-
Male	1	2	2	6	4	3	6	3	1	1	-	-
Female	2	2	2	5	4	4	4	3	2	-	1	-
American Indian 1/, both sexes	2	4	12	11	17	20	15	4	2	2	-	-
Male	1	3	8	3	4	15	8	2	-	-	-	-
Female	1	1	4	8	13	5	7	2	2	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Oregon												
All races, both sexes	32	84	113	156	271	353	343	276	133	27	3	-
Male	21	42	55	75	128	173	166	129	56	10	2	-
Female	11	42	58	81	143	180	177	147	77	17	1	-
White, both sexes	31	79	107	153	265	343	338	274	131	25	3	-
Male	20	40	52	73	125	167	163	128	55	9	2	-
Female	11	39	55	80	140	176	175	146	76	16	1	-
Black, both sexes	-	3	3	1	3	1	3	1	1	1	-	-
Male	-	1	2	1	2	-	2	-	-	1	-	-
Female	-	2	1	-	1	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	2	3	2	3	4	1	-	1	-	-	-
Male	-	1	1	1	1	1	-	-	1	-	-	-
Female	-	1	2	1	2	3	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	5	1	1	-	1	-	-
Male	1	-	-	-	-	5	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Pennsylvania												
All races, both sexes	93	177	309	437	688	1,059	1,211	946	457	128	19	-
Male	42	90	167	229	329	535	549	412	173	37	4	-
Female	51	87	142	208	359	524	662	534	284	91	15	-
White, both sexes	72	150	280	394	640	991	1,163	912	434	120	19	-
Male	32	78	152	207	306	498	524	397	168	34	4	-
Female	40	72	128	187	334	493	639	515	266	86	15	-
Black, both sexes	19	27	28	38	48	65	43	33	23	7	-	-
Male	9	12	15	18	23	34	20	15	5	3	-	-
Female	10	15	13	20	25	31	23	18	18	4	-	-
American Indian 1/, both sexes	1	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	4	-	2	5	1	-	1	-	-
Male	1	-	-	3	-	2	5	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Rhode Island												
All races, both sexes	7	13	26	35	66	78	114	94	34	10	2	-
Male	4	7	6	13	34	40	40	39	15	1	1	-
Female	3	6	20	22	32	38	74	55	19	9	1	-
White, both sexes	7	13	26	33	62	78	110	91	34	10	2	-
Male	4	7	6	12	31	40	39	37	15	1	1	-
Female	3	6	20	21	31	38	71	54	19	9	1	-
Black, both sexes	-	-	-	2	3	-	3	3	-	-	-	-
Male	-	-	-	1	2	-	-	2	-	-	-	-
Female	-	-	-	1	1	-	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	55	99	165	220	298	347	355	218	96	36	5	-
Male	23	57	85	119	159	188	178	88	40	8	-	-
Female	32	42	80	101	139	159	177	130	56	28	5	-
White, both sexes	38	80	133	186	262	315	322	194	88	32	4	-
Male	15	43	66	97	136	169	159	77	38	8	-	-
Female	23	37	67	89	126	146	163	117	50	24	4	-
Black, both sexes	17	19	30	33	36	32	31	23	8	4	1	-
Male	8	14	17	21	23	19	18	10	2	-	-	-
Female	9	5	13	12	13	13	13	13	6	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	-	-	1	-	-	-	-
Male	-	-	2	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
South Dakota												
All races, both sexes	5	13	18	25	55	64	78	58	41	14	1	-
Male	2	4	9	10	37	33	48	33	18	7	-	-
Female	3	9	9	15	18	31	30	25	23	7	1	-
White, both sexes	4	10	16	25	51	63	77	57	40	14	1	-
Male	1	3	7	10	33	32	47	33	18	7	-	-
Female	3	7	9	15	18	31	30	24	22	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	3	2	-	4	1	1	1	1	-	-	-
Male	1	1	2	-	4	1	1	-	-	-	-	-
Female	-	2	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	86	154	241	362	447	547	504	326	168	47	6	-
Male	42	71	119	175	230	275	263	142	67	19	-	-
Female	44	83	122	187	217	272	241	184	101	28	6	-
White, both sexes	71	136	218	338	414	520	475	310	157	45	5	-
Male	35	65	105	161	213	257	248	133	61	17	-	-
Female	36	71	113	177	201	263	227	177	96	28	5	-
Black, both sexes	15	18	22	24	32	27	29	16	11	2	1	-
Male	7	6	14	14	17	18	15	9	6	2	-	-
Female	8	12	8	10	15	9	14	7	5	-	1	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Texas												
All races, both sexes	154	313	516	807	1,080	1,435	1,468	1,033	505	131	24	-
Male	86	140	265	397	554	705	713	480	200	40	6	-
Female	68	173	251	410	526	730	755	553	305	91	18	-
White, both sexes	123	276	479	726	997	1,348	1,396	980	488	125	22	-
Male	68	129	243	354	509	662	674	450	192	37	6	-
Female	55	147	236	372	488	686	722	530	296	88	16	-
Black, both sexes	29	34	36	76	74	78	63	48	15	6	2	-
Male	17	11	21	41	41	39	35	26	6	3	-	-
Female	12	23	15	35	33	39	28	22	9	3	2	-
American Indian 1/, both sexes	-	1	-	2	1	2	4	-	-	-	-	-
Male	-	-	-	1	1	1	2	-	-	-	-	-
Female	-	1	-	1	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	3	8	7	5	5	2	-	-	-
Male	1	-	1	1	3	3	2	4	2	-	-	-
Female	1	2	-	2	5	4	3	1	-	-	-	-
Utah												
All races, both sexes	12	32	33	63	87	118	111	83	28	7	1	-
Male	11	9	14	36	56	66	53	41	17	1	-	-
Female	1	23	19	27	31	52	58	42	11	6	1	-
White, both sexes	11	32	33	63	87	114	106	83	28	7	1	-
Male	10	9	14	36	56	64	50	41	17	1	-	-
Female	1	23	19	27	31	50	56	42	11	6	1	-
Black, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	2	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	4	-	-	-	-	-
Male	-	-	-	-	-	-	3	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Vermont												
All races, both sexes	3	11	25	18	39	58	85	41	24	12	1	-
Male	1	6	15	9	18	30	36	15	9	3	-	-
Female	2	5	10	9	21	28	49	26	15	9	1	-
White, both sexes	3	11	25	18	39	58	85	41	24	12	1	-
Male	1	6	15	9	18	30	36	15	9	3	-	-
Female	2	5	10	9	21	28	49	26	15	9	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	43	108	163	237	411	480	538	397	186	59	13	-
Male	26	55	81	114	195	219	260	169	71	19	5	-
Female	17	53	82	123	216	261	278	228	115	40	8	-
White, both sexes	34	95	140	204	367	431	491	372	169	52	12	-
Male	20	48	71	93	167	193	233	156	63	19	4	-
Female	14	47	69	111	200	238	258	216	106	33	8	-
Black, both sexes	9	13	23	33	40	45	42	25	16	7	1	-
Male	6	7	10	21	25	22	25	13	7	-	1	-
Female	3	6	13	12	15	23	17	12	9	7	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	4	5	-	1	-	-	-
Male	-	-	-	-	2	4	2	-	1	-	-	-
Female	-	-	-	-	1	-	3	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Washington												
All races, both sexes	67	101	159	245	335	487	547	425	210	47	5	1
Male	37	46	72	121	168	229	247	176	75	17	1	1
Female	30	55	87	124	167	258	300	249	135	30	4	-
White, both sexes	63	94	148	237	324	460	528	411	204	46	5	1
Male	35	46	67	116	162	217	234	170	74	16	1	1
Female	28	48	81	121	162	243	294	241	130	30	4	-
Black, both sexes	3	1	7	2	1	6	8	3	1	-	-	-
Male	2	-	4	1	1	3	6	-	-	-	-	-
Female	1	1	3	1	-	3	2	3	1	-	-	-
American Indian 1/, both sexes	-	5	3	2	5	7	1	3	2	-	-	-
Male	-	-	1	1	1	-	1	2	-	-	-	-
Female	-	5	2	1	4	7	-	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	4	5	14	10	8	3	1	-	-
Male	-	-	-	3	4	9	6	4	1	1	-	-
Female	1	1	1	1	1	5	4	4	2	-	-	-
West Virginia												
All races, both sexes	32	55	79	139	200	261	226	134	89	18	2	-
Male	15	27	39	56	102	133	110	58	33	5	1	-
Female	17	28	40	83	98	128	116	76	56	13	1	-
White, both sexes	31	55	77	134	199	261	222	133	89	17	1	-
Male	15	27	37	54	101	133	108	57	33	5	-	-
Female	16	28	40	80	98	128	114	76	56	12	1	-
Black, both sexes	1	-	2	5	1	-	4	1	-	1	1	-
Male	-	-	2	2	1	-	2	1	-	-	1	-
Female	1	-	-	3	-	-	2	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Wisconsin												
All races, both sexes	41	77	118	201	274	468	499	374	212	54	3	-
Male	23	39	68	96	147	237	240	165	88	14	-	-
Female	18	38	50	105	127	231	259	209	124	40	3	-
White, both sexes	33	71	109	194	266	453	493	369	208	52	3	-
Male	18	36	62	91	143	231	239	163	87	14	-	-
Female	15	35	47	103	123	222	254	206	121	38	3	-
Black, both sexes	7	5	7	5	8	9	3	2	4	-	-	-
Male	5	3	5	4	4	2	1	-	1	-	-	-
Female	2	2	2	1	4	7	2	2	3	-	-	-
American Indian 1/, both sexes	1	1	2	2	-	2	2	2	-	1	-	-
Male	-	-	1	1	-	2	-	1	-	-	-	-
Female	1	1	1	1	-	-	2	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	4	1	1	-	1	-	-
Male	-	-	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	1	-	-
Wyoming												
All races, both sexes	8	16	20	27	49	62	76	55	21	4	-	-
Male	3	9	11	11	29	34	41	25	11	-	-	-
Female	5	7	9	16	20	28	35	30	10	4	-	-
White, both sexes	7	15	20	26	48	62	75	55	20	4	-	-
Male	3	8	11	10	28	34	41	25	11	-	-	-
Female	4	7	9	16	20	28	34	30	9	4	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	1	1	-	1	-	-	-	-	-
Male	-	1	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
United States												
All races, both sexes	123,610	30	43	52	63	61	87	112	140	221	613	1,309
Male	58,976	20	29	33	45	39	51	66	70	105	288	610
Female	64,634	10	14	19	18	22	36	46	70	116	325	699
White, both sexes	114,102	15	26	24	18	21	45	63	75	145	442	1,068
Male	53,799	10	17	18	14	12	25	38	31	69	212	492
Female	60,303	5	9	6	4	9	20	25	44	76	230	576
Black, both sexes	7,695	11	15	28	41	37	38	46	62	74	149	216
Male	4,114	6	12	15	29	25	22	25	36	34	67	103
Female	3,581	5	3	13	12	12	16	21	26	40	82	113
American Indian 1/, both sexes	503	2	-	-	2	-	1	-	1	1	11	13
Male	231	2	-	-	1	-	1	-	1	1	6	7
Female	272	-	-	-	1	-	-	-	-	-	5	6
Asian or Pacific Islander 2/, both sexes	1,310	2	2	-	2	3	3	3	2	1	11	12
Male	832	2	-	-	1	2	3	3	2	1	3	8
Female	478	-	2	-	1	1	-	-	-	-	8	4
Alabama												
All races, both sexes	2,297	-	1	-	2	-	1	1	3	5	14	26
Male	1,153	-	1	-	2	-	1	1	1	2	5	15
Female	1,144	-	-	-	-	-	-	-	2	3	9	11
White, both sexes	2,056	-	-	-	1	-	-	-	1	4	9	22
Male	997	-	-	-	1	-	-	-	1	2	4	14
Female	1,059	-	-	-	-	-	-	-	-	2	5	8
Black, both sexes	238	-	1	-	1	-	1	1	2	1	5	4
Male	155	-	1	-	1	-	1	1	-	-	1	1
Female	83	-	-	-	-	-	-	-	2	1	4	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Alaska												
All races, both sexes	140	-	-	-	-	-	-	-	-	1	2	-
Male	58	-	-	-	-	-	-	-	-	-	-	-
Female	82	-	-	-	-	-	-	-	-	1	2	-
White, both sexes	100	-	-	-	-	-	-	-	-	1	2	-
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	59	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2,771	1	2	1	4	-	-	1	3	4	6	19
Male	1,344	1	2	1	2	-	-	1	1	4	2	7
Female	1,427	-	-	-	2	-	-	-	2	-	4	12
White, both sexes	2,717	-	2	1	1	-	-	1	3	3	5	18
Male	1,319	-	2	1	1	-	-	1	1	3	2	7
Female	1,398	-	-	-	-	-	-	-	2	-	3	11
Black, both sexes	33	-	-	-	1	-	-	-	-	1	1	1
Male	15	-	-	-	-	-	-	-	-	1	-	-
Female	18	-	-	-	1	-	-	-	-	-	1	1
American Indian 1/, both sexes	15	1	-	-	2	-	-	-	-	-	-	-
Male	4	1	-	-	1	-	-	-	-	-	-	-
Female	11	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Arkansas												
All races, both sexes	1,489	-	1	-	-	1	2	1	3	3	11	6
Male	784	-	1	-	-	1	-	-	1	2	7	2
Female	705	-	-	-	-	-	2	1	2	1	4	4
White, both sexes	1,405	-	-	-	-	-	1	1	1	3	11	5
Male	740	-	-	-	-	-	-	-	-	2	7	2
Female	665	-	-	-	-	-	1	1	1	1	4	3
Black, both sexes	79	-	1	-	-	1	1	-	1	-	-	1
Male	43	-	1	-	-	1	-	-	-	-	-	-
Female	36	-	-	-	-	-	1	-	1	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	12,709	3	8	7	7	10	10	15	14	23	55	115
Male	5,916	2	5	4	7	6	7	9	7	11	21	56
Female	6,793	1	3	3	-	4	3	6	7	12	34	59
White, both sexes	11,258	2	7	3	-	6	5	5	8	11	35	94
Male	5,065	1	4	2	-	5	3	2	1	5	16	45
Female	6,193	1	3	1	-	1	2	3	7	6	19	49
Black, both sexes	712	-	1	4	6	3	5	7	4	11	13	17
Male	382	-	1	2	6	-	4	4	4	5	4	7
Female	330	-	-	2	-	3	1	3	-	6	9	10
American Indian 1/, both sexes	59	-	-	-	-	-	-	-	-	1	1	-
Male	30	-	-	-	-	-	-	-	-	1	-	-
Female	29	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	680	1	-	-	1	1	-	3	2	-	6	4
Male	439	1	-	-	1	1	-	3	2	-	1	4
Female	241	-	-	-	-	-	-	-	-	-	5	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Colorado												
All races, both sexes	1,920	-	-	-	-	1	1	2	4	1	7	15
Male	969	-	-	-	-	1	1	1	3	-	3	5
Female	951	-	-	-	-	-	-	1	1	1	4	10
White, both sexes	1,850	-	-	-	-	1	1	2	3	1	4	14
Male	938	-	-	-	-	1	1	1	2	-	2	4
Female	912	-	-	-	-	-	-	1	1	1	2	10
Black, both sexes	54	-	-	-	-	-	-	-	1	-	1	1
Male	26	-	-	-	-	-	-	-	1	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1,448	-	-	2	1	-	-	-	-	3	4	12
Male	647	-	-	1	1	-	-	-	-	1	2	5
Female	801	-	-	1	-	-	-	-	-	2	2	7
White, both sexes	1,395	-	-	2	-	-	-	-	-	2	3	9
Male	616	-	-	1	-	-	-	-	-	-	1	4
Female	779	-	-	1	-	-	-	-	-	2	2	5
Black, both sexes	51	-	-	-	1	-	-	-	-	1	1	3
Male	29	-	-	-	1	-	-	-	-	1	1	1
Female	22	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Delaware												
All races, both sexes	344	-	-	-	-	-	-	2	-	-	1	-
Male	158	-	-	-	-	-	-	-	-	-	1	-
Female	186	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	303	-	-	-	-	-	-	1	-	-	-	-
Male	134	-	-	-	-	-	-	-	-	-	-	-
Female	169	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	37	-	-	-	-	-	-	1	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	126	1	-	-	-	-	-	1	-	2	3	4
Male	60	1	-	-	-	-	-	1	-	-	1	3
Female	66	-	-	-	-	-	-	-	-	2	2	1
White, both sexes	45	-	-	-	-	-	-	-	-	1	1	2
Male	19	-	-	-	-	-	-	-	-	-	-	2
Female	26	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	80	1	-	-	-	-	-	1	-	1	2	2
Male	40	1	-	-	-	-	-	1	-	-	1	1
Female	40	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Florida												
All races, both sexes	8,841	3	2	4	4	3	7	9	10	12	32	94
Male	4,167	2	1	3	3	-	5	7	4	2	15	40
Female	4,674	1	1	1	1	3	2	2	6	10	17	54
White, both sexes	8,364	1	1	2	2	2	4	6	5	8	28	77
Male	3,896	1	-	2	2	-	3	5	2	1	13	32
Female	4,468	-	1	-	-	2	1	1	3	7	15	45
Black, both sexes	443	2	1	2	2	1	3	3	5	4	3	16
Male	250	1	1	1	1	-	2	2	2	1	2	7
Female	193	1	-	1	1	1	1	1	3	3	1	9
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
Georgia												
All races, both sexes	3,362	-	1	5	4	2	2	7	4	11	23	45
Male	1,625	-	1	3	2	1	1	4	2	3	9	17
Female	1,737	-	-	2	2	1	1	3	2	8	14	28
White, both sexes	2,928	-	-	2	1	-	-	4	-	5	12	27
Male	1,377	-	-	2	-	-	-	2	-	1	5	8
Female	1,551	-	-	-	1	-	-	2	-	4	7	19
Black, both sexes	409	-	1	3	3	2	2	3	4	6	10	17
Male	236	-	1	1	2	1	1	2	2	2	3	9
Female	173	-	-	2	1	1	1	1	2	4	7	8
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Hawaii												
All races, both sexes	279	-	-	-	-	1	-	-	-	1	-	3
Male	158	-	-	-	-	-	-	-	-	1	-	1
Female	121	-	-	-	-	1	-	-	-	-	-	2
White, both sexes	101	-	-	-	-	-	-	-	-	-	-	1
Male	51	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	176	-	-	-	-	1	-	-	-	1	-	2
Male	105	-	-	-	-	-	-	-	-	1	-	1
Female	71	-	-	-	-	1	-	-	-	-	-	1
Idaho												
All races, both sexes	639	-	-	-	-	-	-	-	-	-	1	6
Male	334	-	-	-	-	-	-	-	-	-	-	3
Female	305	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	635	-	-	-	-	-	-	-	-	-	1	6
Male	331	-	-	-	-	-	-	-	-	-	-	3
Female	304	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Illinois												
All races, both sexes	4,691	-	2	4	2	4	7	5	17	13	28	50
Male	2,221	-	2	2	2	4	3	4	14	8	15	21
Female	2,470	-	-	2	-	-	4	1	3	5	13	29
White, both sexes	4,218	-	-	3	-	-	3	2	5	9	18	32
Male	1,966	-	-	2	-	-	1	2	4	6	10	14
Female	2,252	-	-	1	-	-	2	-	1	3	8	18
Black, both sexes	443	-	2	1	2	4	4	3	12	4	10	18
Male	237	-	2	-	2	4	2	2	10	2	5	7
Female	206	-	-	1	-	-	2	1	2	2	5	11
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	3,268	2	1	-	3	1	3	1	2	10	16	47
Male	1,585	2	-	-	1	1	1	-	1	5	7	23
Female	1,683	-	1	-	2	-	2	1	1	5	9	24
White, both sexes	3,129	2	1	-	1	1	1	1	1	8	13	42
Male	1,519	2	-	-	1	1	1	-	1	3	6	20
Female	1,610	-	1	-	-	-	-	1	-	5	7	22
Black, both sexes	135	-	-	-	2	-	2	-	1	2	3	5
Male	65	-	-	-	-	-	-	-	-	2	1	3
Female	70	-	-	-	2	-	2	-	1	-	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Iowa												
All races, both sexes	1,640	-	-	-	-	1	1	-	3	3	11	8
Male	865	-	-	-	-	1	1	-	1	1	6	3
Female	775	-	-	-	-	-	-	-	2	2	5	5
White, both sexes	1,626	-	-	-	-	-	1	-	2	3	9	8
Male	857	-	-	-	-	-	1	-	-	1	5	3
Female	769	-	-	-	-	-	-	-	2	2	4	5
Black, both sexes	8	-	-	-	-	1	-	-	1	-	1	-
Male	7	-	-	-	-	1	-	-	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1,481	-	-	-	-	-	1	4	4	5	9	20
Male	754	-	-	-	-	-	1	3	2	2	4	8
Female	727	-	-	-	-	-	-	1	2	3	5	12
White, both sexes	1,411	-	-	-	-	-	-	4	2	5	7	19
Male	723	-	-	-	-	-	-	3	-	2	3	8
Female	688	-	-	-	-	-	-	1	2	3	4	11
Black, both sexes	46	-	-	-	-	-	1	-	2	-	1	1
Male	21	-	-	-	-	-	1	-	2	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Kentucky												
All races, both sexes	2,391	-	1	-	-	1	2	2	1	5	16	48
Male	1,156	-	1	-	-	1	-	1	1	4	9	22
Female	1,235	-	-	-	-	-	2	1	-	1	7	26
White, both sexes	2,279	-	1	-	-	-	2	1	1	4	14	43
Male	1,104	-	1	-	-	-	-	1	1	3	9	21
Female	1,175	-	-	-	-	-	2	-	-	1	5	22
Black, both sexes	106	-	-	-	-	1	-	1	-	1	2	5
Male	48	-	-	-	-	1	-	-	-	1	-	1
Female	58	-	-	-	-	-	-	1	-	-	2	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,681	2	2	1	1	-	1	1	2	3	12	21
Male	862	2	2	-	1	-	1	1	1	1	5	11
Female	819	-	-	1	-	-	-	-	1	2	7	10
White, both sexes	1,391	-	-	-	-	-	-	1	1	2	6	15
Male	683	-	-	-	-	-	-	1	-	-	2	6
Female	708	-	-	-	-	-	-	-	1	2	4	9
Black, both sexes	286	2	2	1	1	-	1	-	1	1	6	6
Male	175	2	2	-	1	-	1	-	1	1	3	5
Female	111	-	-	1	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Maine												
All races, both sexes	776	-	-	-	1	-	-	-	-	1	4	7
Male	389	-	-	-	1	-	-	-	-	-	2	6
Female	387	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	774	-	-	-	1	-	-	-	-	1	4	7
Male	389	-	-	-	1	-	-	-	-	-	2	6
Female	385	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,817	-	1	-	2	1	1	3	3	2	12	16
Male	787	-	-	-	1	1	1	1	1	-	5	3
Female	1,030	-	1	-	1	-	-	2	2	2	7	13
White, both sexes	1,545	-	-	-	-	-	-	1	1	2	4	13
Male	663	-	-	-	-	-	-	-	1	-	-	2
Female	882	-	-	-	-	-	-	1	-	2	4	11
Black, both sexes	254	-	1	-	2	1	1	2	2	-	8	3
Male	115	-	-	-	1	1	1	1	-	-	5	1
Female	139	-	1	-	1	-	-	1	2	-	3	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Massachusetts												
All races, both sexes	2,509	-	-	-	-	-	2	2	2	1	13	16
Male	1,055	-	-	-	-	-	-	2	1	-	5	5
Female	1,454	-	-	-	-	-	2	-	1	1	8	11
White, both sexes	2,439	-	-	-	-	-	1	2	2	1	8	15
Male	1,018	-	-	-	-	-	-	2	1	-	4	5
Female	1,421	-	-	-	-	-	1	-	1	1	4	10
Black, both sexes	51	-	-	-	-	-	1	-	-	-	5	1
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	31	-	-	-	-	-	1	-	-	-	4	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4,455	-	1	4	3	5	3	8	3	6	25	51
Male	2,130	-	-	2	2	4	2	5	1	5	13	23
Female	2,325	-	1	2	1	1	1	3	2	1	12	28
White, both sexes	4,121	-	-	-	1	1	3	5	1	3	18	41
Male	1,951	-	-	-	1	1	2	2	1	3	10	15
Female	2,170	-	-	-	-	-	1	3	-	-	8	26
Black, both sexes	301	-	1	4	2	4	-	3	2	3	5	8
Male	161	-	-	2	1	3	-	3	-	2	2	6
Female	140	-	1	2	1	1	-	-	2	1	3	2
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	-	-	2	2
Male	14	-	-	-	-	-	-	-	-	-	1	2
Female	12	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Minnesota												
All races, both sexes	1,754	-	-	1	-	1	-	1	1	1	4	13
Male	866	-	-	1	-	1	-	-	-	-	3	7
Female	888	-	-	-	-	-	-	1	1	1	1	6
White, both sexes	1,712	-	-	-	-	-	-	-	1	-	3	12
Male	841	-	-	-	-	-	-	-	-	-	2	6
Female	871	-	-	-	-	-	-	-	1	-	1	6
Black, both sexes	22	-	-	1	-	1	-	1	-	1	1	1
Male	14	-	-	1	-	1	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,372	-	-	-	2	1	-	3	4	5	14	19
Male	735	-	-	-	2	1	-	2	1	1	7	7
Female	637	-	-	-	-	-	-	1	3	4	7	12
White, both sexes	1,116	-	-	-	-	-	-	1	-	1	7	11
Male	581	-	-	-	-	-	-	1	-	-	3	4
Female	535	-	-	-	-	-	-	-	-	1	4	7
Black, both sexes	253	-	-	-	2	1	-	2	4	4	7	8
Male	153	-	-	-	2	1	-	1	1	1	4	3
Female	100	-	-	-	-	-	-	1	3	3	3	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Missouri												
All races, both sexes	2,993	1	-	-	3	-	3	5	4	7	24	45
Male	1,428	-	-	-	2	-	3	3	3	6	14	25
Female	1,565	1	-	-	1	-	-	2	1	1	10	20
White, both sexes	2,810	1	-	-	1	-	2	3	3	4	18	40
Male	1,332	-	-	-	1	-	2	2	2	3	10	22
Female	1,478	1	-	-	-	-	-	1	1	1	8	18
Black, both sexes	171	-	-	-	1	-	1	2	1	3	5	5
Male	92	-	-	-	1	-	1	1	1	3	3	3
Female	79	-	-	-	-	-	-	1	-	-	2	2
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	1	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
Montana												
All races, both sexes	577	-	-	1	-	-	-	-	-	-	-	3
Male	259	-	-	-	-	-	-	-	-	-	-	1
Female	318	-	-	1	-	-	-	-	-	-	-	2
White, both sexes	557	-	-	1	-	-	-	-	-	-	-	3
Male	248	-	-	-	-	-	-	-	-	-	-	1
Female	309	-	-	1	-	-	-	-	-	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Nebraska												
All races, both sexes	869	-	-	-	-	1	-	-	2	1	1	4
Male	435	-	-	-	-	-	-	-	1	1	-	2
Female	434	-	-	-	-	1	-	-	1	-	1	2
White, both sexes	845	-	-	-	-	-	-	-	1	1	-	3
Male	422	-	-	-	-	-	-	-	-	1	-	2
Female	423	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	16	-	-	-	-	1	-	-	1	-	1	1
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	8	-	-	-	-	1	-	-	-	-	1	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,064	-	-	-	-	-	1	-	1	4	5	13
Male	485	-	-	-	-	-	1	-	-	1	3	4
Female	579	-	-	-	-	-	-	-	1	3	2	9
White, both sexes	1,018	-	-	-	-	-	1	-	1	3	4	12
Male	460	-	-	-	-	-	1	-	-	1	3	3
Female	558	-	-	-	-	-	-	-	1	2	1	9
Black, both sexes	28	-	-	-	-	-	-	-	-	1	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
New Hampshire												
All races, both sexes	598	-	-	-	-	-	-	-	1	2	1	3
Male	277	-	-	-	-	-	-	-	-	2	-	2
Female	321	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	597	-	-	-	-	-	-	-	1	2	1	3
Male	276	-	-	-	-	-	-	-	-	2	-	2
Female	321	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2,840	-	1	-	2	4	2	1	2	10	18	27
Male	1,224	-	1	-	2	2	2	-	1	3	6	10
Female	1,616	-	-	-	-	2	-	1	1	7	12	17
White, both sexes	2,549	-	-	-	1	-	1	1	1	6	12	20
Male	1,088	-	-	-	1	-	1	-	-	3	4	8
Female	1,461	-	-	-	-	-	-	1	1	3	8	12
Black, both sexes	259	-	1	-	1	3	1	-	1	4	5	7
Male	117	-	1	-	1	1	1	-	1	-	2	2
Female	142	-	-	-	-	2	-	-	-	4	3	5
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	1	-	-	-	-	1	-
Male	17	-	-	-	-	1	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
New Mexico												
All races, both sexes	881	-	-	-	2	-	-	1	-	1	5	9
Male	433	-	-	-	2	-	-	1	-	1	-	5
Female	448	-	-	-	-	-	-	-	-	-	5	4
White, both sexes	861	-	-	-	2	-	-	1	-	1	4	8
Male	421	-	-	-	2	-	-	1	-	1	-	4
Female	440	-	-	-	-	-	-	-	-	-	4	4
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6,290	1	3	3	4	3	6	11	9	10	36	63
Male	2,870	1	1	3	2	3	2	3	3	6	21	33
Female	3,420	-	2	-	2	-	4	8	6	4	15	30
White, both sexes	5,672	1	-	2	-	1	5	4	5	7	23	47
Male	2,556	1	-	2	-	1	2	2	1	4	15	26
Female	3,116	-	-	-	-	-	3	2	4	3	8	21
Black, both sexes	526	-	2	1	4	2	1	7	4	3	13	16
Male	246	-	1	1	2	2	-	1	2	2	6	7
Female	280	-	1	-	2	-	1	6	2	1	7	9
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	76	-	1	-	-	-	-	-	-	-	-	-
Male	59	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
North Carolina												
All races, both sexes	3,983	1	-	1	1	4	3	5	7	9	16	47
Male	1,906	1	-	1	1	1	-	2	3	5	6	21
Female	2,077	-	-	-	-	3	3	3	4	4	10	26
White, both sexes	3,550	-	-	-	-	2	1	3	4	5	7	35
Male	1,663	-	-	-	-	-	-	1	2	3	2	15
Female	1,887	-	-	-	-	2	1	2	2	2	5	20
Black, both sexes	409	1	-	1	1	2	2	2	3	4	9	10
Male	229	1	-	1	1	1	-	1	1	2	4	4
Female	180	-	-	-	-	1	2	1	2	2	5	6
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	281	-	-	-	-	-	1	-	-	-	-	1
Male	157	-	-	-	-	-	1	-	-	-	-	1
Female	124	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	272	-	-	-	-	-	1	-	-	-	-	1
Male	155	-	-	-	-	-	1	-	-	-	-	1
Female	117	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Ohio												
All races, both sexes	6,025	1	2	2	1	1	3	4	6	13	30	80
Male	2,897	1	1	1	1	-	2	3	4	8	14	38
Female	3,128	-	1	1	-	1	1	1	2	5	16	42
White, both sexes	5,569	1	2	-	-	1	2	2	3	9	26	61
Male	2,645	1	1	-	-	-	1	2	2	5	13	29
Female	2,924	-	1	-	-	1	1	-	1	4	13	32
Black, both sexes	445	-	-	2	1	-	1	2	3	4	4	18
Male	247	-	-	1	1	-	1	1	2	3	1	9
Female	198	-	-	1	-	-	-	1	1	1	3	9
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2,183	2	2	1	4	2	3	1	1	6	12	23
Male	1,097	2	2	1	2	2	2	1	1	2	4	11
Female	1,086	-	-	-	2	-	1	-	-	4	8	12
White, both sexes	2,020	1	2	1	1	-	1	1	-	6	9	19
Male	1,015	1	2	1	1	-	-	1	-	2	2	10
Female	1,005	-	-	-	-	-	1	-	-	4	7	9
Black, both sexes	66	-	-	-	3	2	1	-	1	-	-	1
Male	34	-	-	-	1	2	1	-	1	-	-	-
Female	32	-	-	-	2	-	-	-	-	-	-	1
American Indian 1/, both sexes	95	1	-	-	-	-	-	-	-	-	3	3
Male	47	1	-	-	-	-	-	-	-	-	2	1
Female	48	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Oregon												
All races, both sexes	1,810	1	1	-	-	1	1	-	2	2	5	24
Male	875	-	-	-	-	1	1	-	2	2	4	15
Female	935	1	1	-	-	-	-	-	-	-	1	9
White, both sexes	1,767	1	1	-	-	1	1	-	2	2	5	23
Male	852	-	-	-	-	1	1	-	2	2	4	15
Female	915	1	1	-	-	-	-	-	-	-	1	8
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	5,572	1	1	2	2	1	1	4	2	8	21	53
Male	2,598	-	1	1	-	-	-	4	-	2	14	26
Female	2,974	1	-	1	2	1	1	-	2	6	7	27
White, both sexes	5,198	1	1	2	2	-	-	1	-	3	15	43
Male	2,419	-	1	1	-	-	-	1	-	2	9	20
Female	2,779	1	-	1	2	-	-	-	-	1	6	23
Black, both sexes	356	-	-	-	-	1	1	3	2	5	6	10
Male	166	-	-	-	-	-	-	3	-	-	5	6
Female	190	-	-	-	-	1	1	-	2	5	1	4
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Rhode Island												
All races, both sexes	475	-	-	1	-	-	-	-	-	-	2	3
Male	200	-	-	1	-	-	-	-	-	-	1	-
Female	275	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	462	-	-	1	-	-	-	-	-	-	2	3
Male	193	-	-	1	-	-	-	-	-	-	1	-
Female	269	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1,919	-	-	1	2	-	2	1	-	3	9	22
Male	964	-	-	1	1	-	-	1	-	2	6	12
Female	955	-	-	-	1	-	2	-	-	1	3	10
White, both sexes	1,663	-	-	-	-	-	1	1	-	1	5	14
Male	819	-	-	-	-	-	-	1	-	1	4	7
Female	844	-	-	-	-	-	1	-	-	-	1	7
Black, both sexes	250	-	-	1	2	-	1	-	-	2	4	8
Male	140	-	-	1	1	-	-	-	-	1	2	5
Female	110	-	-	-	1	-	1	-	-	1	2	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
South Dakota												
All races, both sexes	373	-	-	-	-	-	1	-	1	-	-	2
Male	203	-	-	-	-	-	1	-	1	-	-	1
Female	170	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	359	-	-	-	-	-	1	-	1	-	-	2
Male	193	-	-	-	-	-	1	-	1	-	-	1
Female	166	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2,956	-	1	3	3	2	6	2	2	2	25	46
Male	1,450	-	-	3	3	1	4	1	2	2	14	22
Female	1,506	-	1	-	-	1	2	1	-	-	11	24
White, both sexes	2,738	-	1	1	1	-	3	2	1	1	22	41
Male	1,327	-	-	1	1	-	2	1	1	1	12	18
Female	1,411	-	1	-	-	-	1	1	-	-	10	23
Black, both sexes	216	-	-	2	2	2	3	-	1	1	3	5
Male	123	-	-	2	2	1	2	-	1	1	2	4
Female	93	-	-	-	-	1	1	-	-	-	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Texas												
All races, both sexes	7,570	6	5	5	1	5	7	3	10	11	38	80
Male	3,646	2	4	3	1	4	5	-	4	4	16	36
Female	3,924	4	1	2	-	1	2	3	6	7	22	44
White, both sexes	7,025	2	3	3	-	3	1	2	8	10	30	64
Male	3,366	2	3	2	-	2	-	-	3	4	12	30
Female	3,659	-	-	1	-	1	1	2	5	6	18	34
Black, both sexes	494	4	1	2	1	2	3	1	2	1	8	12
Male	255	-	1	1	1	2	2	-	1	-	4	5
Female	239	4	-	1	-	-	1	1	1	1	4	7
American Indian 1/, both sexes	13	-	-	-	-	-	1	-	-	-	-	2
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	38	-	1	-	-	-	2	-	-	-	-	2
Male	19	-	-	-	-	-	2	-	-	-	-	1
Female	19	-	1	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	587	2	1	-	-	1	-	-	1	-	3	8
Male	308	1	1	-	-	-	-	-	-	-	1	3
Female	279	1	-	-	-	1	-	-	1	-	2	5
White, both sexes	578	2	1	-	-	1	-	-	1	-	3	8
Male	303	1	1	-	-	-	-	-	-	-	1	3
Female	275	1	-	-	-	1	-	-	1	-	2	5
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Vermont												
All races, both sexes	316	-	-	-	-	-	-	-	-	1	-	2
Male	142	-	-	-	-	-	-	-	-	-	-	-
Female	174	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	316	-	-	-	-	-	-	-	-	1	-	2
Male	142	-	-	-	-	-	-	-	-	-	-	-
Female	174	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2,673	1	2	2	-	1	1	1	-	5	7	41
Male	1,239	1	1	1	-	1	1	1	-	2	4	19
Female	1,434	-	1	1	-	-	-	-	-	3	3	22
White, both sexes	2,393	-	2	-	-	-	1	1	-	2	5	36
Male	1,085	-	1	-	-	-	1	1	-	-	3	16
Female	1,308	-	1	-	-	-	-	-	-	2	2	20
Black, both sexes	264	-	-	2	-	1	-	-	-	3	2	4
Male	143	-	-	1	-	1	-	-	-	2	1	3
Female	121	-	-	1	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	14	1	-	-	-	-	-	-	-	-	-	-
Male	10	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Washington												
All races, both sexes	2,634	1	-	1	-	-	-	1	-	2	9	19
Male	1,199	1	-	-	-	-	-	1	-	1	1	11
Female	1,435	-	-	1	-	-	-	-	-	1	8	8
White, both sexes	2,524	-	-	-	-	-	-	1	-	1	9	18
Male	1,146	-	-	-	-	-	-	1	-	-	1	10
Female	1,378	-	-	-	-	-	-	-	-	1	8	8
Black, both sexes	35	1	-	1	-	-	-	-	-	1	-	-
Male	19	1	-	-	-	-	-	-	-	1	-	-
Female	16	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	47	-	-	-	-	-	-	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1,261	-	-	-	-	1	1	1	3	1	11	14
Male	594	-	-	-	-	-	-	1	1	-	6	7
Female	667	-	-	-	-	1	1	-	2	1	5	7
White, both sexes	1,244	-	-	-	-	1	1	1	3	-	10	14
Male	585	-	-	-	-	-	-	1	1	-	6	7
Female	659	-	-	-	-	1	1	-	2	-	4	7
Black, both sexes	17	-	-	-	-	-	-	-	-	1	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Wisconsin												
All races, both sexes	2,340	-	1	-	2	1	1	1	3	1	12	11
Male	1,136	-	1	-	1	1	1	1	1	1	6	9
Female	1,204	-	-	-	1	-	-	-	2	-	6	2
White, both sexes	2,263	-	1	-	2	-	-	-	2	1	10	10
Male	1,096	-	1	-	1	-	-	-	-	1	4	8
Female	1,167	-	-	-	1	-	-	-	2	-	6	2
Black, both sexes	55	-	-	-	-	1	1	1	1	-	1	-
Male	30	-	-	-	-	1	1	1	1	-	1	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	340	-	-	-	-	-	-	1	-	1	-	5
Male	176	-	-	-	-	-	-	-	-	1	-	3
Female	164	-	-	-	-	-	-	1	-	-	-	2
White, both sexes	334	-	-	-	-	-	-	1	-	1	-	5
Male	173	-	-	-	-	-	-	-	-	1	-	3
Female	161	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
United States												
All races, both sexes	2,583	4,676	7,639	11,546	16,537	22,630	24,404	18,501	9,250	2,714	395	4
Male	1,327	2,416	3,939	5,782	8,323	11,321	11,730	8,182	3,647	838	112	3
Female	1,256	2,260	3,700	5,764	8,214	11,309	12,674	10,319	5,603	1,876	283	1
White, both sexes	2,144	4,081	6,867	10,558	15,360	21,113	22,931	17,501	8,736	2,511	355	3
Male	1,095	2,095	3,479	5,217	7,659	10,484	10,898	7,665	3,414	755	98	2
Female	1,049	1,986	3,388	5,341	7,701	10,629	12,033	9,836	5,322	1,756	257	1
Black, both sexes	403	531	668	838	973	1,188	1,105	742	358	137	35	-
Male	215	290	394	477	547	639	608	368	142	50	10	-
Female	188	241	274	361	426	549	497	374	216	87	25	-
American Indian 1/, both sexes	18	31	54	60	75	94	72	44	16	7	1	-
Male	6	12	32	27	34	42	31	21	5	1	1	-
Female	12	19	22	33	41	52	41	23	11	6	-	-
Asian or Pacific Islander 2/, both sexes	18	33	50	90	129	235	296	214	140	59	4	1
Male	11	19	34	61	83	156	193	128	86	32	3	1
Female	7	14	16	29	46	79	103	86	54	27	1	-
Alabama												
All races, both sexes	53	101	168	258	365	421	432	284	114	44	4	-
Male	25	60	84	140	190	219	207	140	48	11	1	-
Female	28	41	84	118	175	202	225	144	66	33	3	-
White, both sexes	44	79	152	227	335	387	399	255	97	40	4	-
Male	21	45	71	119	168	195	185	124	38	8	1	-
Female	23	34	81	108	167	192	214	131	59	32	3	-
Black, both sexes	9	22	16	31	30	33	32	29	17	3	-	-
Male	4	15	13	21	22	23	22	16	10	3	-	-
Female	5	7	3	10	8	10	10	13	7	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Alaska												
All races, both sexes	4	11	14	15	24	22	21	17	8	1	-	-
Male	2	5	10	6	8	10	6	7	4	-	-	-
Female	2	6	4	9	16	12	15	10	4	1	-	-
White, both sexes	3	8	9	8	20	18	14	13	4	-	-	-
Male	1	3	7	3	8	9	4	4	2	-	-	-
Female	2	5	2	5	12	9	10	9	2	-	-	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	3	4	5	3	3	6	1	4	1	-	-
Male	1	2	2	2	-	1	2	1	2	-	-	-
Female	-	1	2	3	3	2	4	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	1	1	2	-	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	1	1	1	-	-	-	-
Arizona												
All races, both sexes	49	105	151	267	383	560	560	425	180	45	5	-
Male	24	58	78	141	188	296	267	184	70	15	2	-
Female	25	47	73	126	195	264	293	241	110	30	3	-
White, both sexes	48	102	147	257	379	550	556	419	176	44	5	-
Male	24	56	77	136	187	291	264	181	69	14	2	-
Female	24	46	70	121	192	259	292	238	107	30	3	-
Black, both sexes	1	2	2	7	2	6	3	3	3	-	-	-
Male	-	2	1	4	1	2	3	1	-	-	-	-
Female	1	-	1	3	1	4	-	2	3	-	-	-
American Indian 1/, both sexes	-	1	2	2	2	2	1	2	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	2	2	2	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	-	1	1	1	-	-
Male	-	-	-	1	-	2	-	1	1	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Arkansas												
All races, both sexes	40	65	129	162	210	265	270	195	96	26	3	-
Male	12	41	70	81	129	146	145	98	38	9	1	-
Female	28	24	59	81	81	119	125	97	58	17	2	-
White, both sexes	34	58	117	156	203	253	255	188	93	23	3	-
Male	9	38	63	77	125	138	137	96	36	9	1	-
Female	25	20	54	79	78	115	118	92	57	14	2	-
Black, both sexes	5	7	12	6	7	10	14	7	3	3	-	-
Male	3	3	7	4	4	8	8	2	2	-	-	-
Female	2	4	5	2	3	2	6	5	1	3	-	-
American Indian 1/, both sexes	1	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	255	420	707	1,046	1,527	2,227	2,638	2,121	1,121	327	53	-
Male	133	233	392	512	752	1,069	1,221	897	433	118	21	-
Female	122	187	315	534	775	1,158	1,417	1,224	688	209	32	-
White, both sexes	204	350	593	932	1,369	1,990	2,377	1,931	1,014	274	48	-
Male	108	188	312	440	656	928	1,066	795	378	93	17	-
Female	96	162	281	492	713	1,062	1,311	1,136	636	181	31	-
Black, both sexes	43	54	79	71	84	106	97	58	31	15	3	-
Male	21	34	51	45	43	57	47	29	11	5	2	-
Female	22	20	28	26	41	49	50	29	20	10	1	-
American Indian 1/, both sexes	4	1	7	6	8	11	10	7	2	1	-	-
Male	1	1	6	3	7	6	4	1	-	-	-	-
Female	3	-	1	3	1	5	6	6	2	1	-	-
Asian or Pacific Islander 2/, both sexes	4	15	28	37	66	120	154	125	74	37	2	-
Male	3	10	23	24	46	78	104	72	44	20	2	-
Female	1	5	5	13	20	42	50	53	30	17	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Colorado												
All races, both sexes	47	75	111	189	261	351	390	267	151	40	7	-
Male	25	49	69	112	131	166	186	133	72	12	-	-
Female	22	26	42	77	130	185	204	134	79	28	7	-
White, both sexes	43	67	106	179	250	341	382	261	149	40	6	-
Male	23	45	66	108	125	163	182	131	72	12	-	-
Female	20	22	40	71	125	178	200	130	77	28	6	-
Black, both sexes	4	7	5	8	10	7	3	5	1	-	1	-
Male	2	4	3	3	6	3	1	2	-	-	-	-
Female	2	3	2	5	4	4	2	3	1	-	1	-
American Indian 1/, both sexes	-	1	-	1	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	5	-	1	-	-	-
Male	-	-	-	1	-	-	3	-	-	-	-	-
Female	-	-	-	-	1	1	2	-	1	-	-	-
Connecticut												
All races, both sexes	32	41	71	94	142	243	289	279	175	54	5	1
Male	18	16	32	53	75	118	135	110	66	12	2	-
Female	14	25	39	41	67	125	154	169	109	42	3	1
White, both sexes	29	37	68	86	139	234	281	276	171	53	4	1
Male	16	14	31	47	73	113	129	110	63	12	2	-
Female	13	23	37	39	66	121	152	166	108	41	2	1
Black, both sexes	2	4	3	8	3	9	8	3	3	1	1	-
Male	1	2	1	6	2	5	6	-	2	-	-	-
Female	1	2	2	2	1	4	2	3	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Delaware												
All races, both sexes	7	10	13	35	55	75	73	43	19	9	2	-
Male	3	3	8	19	28	37	31	14	9	5	-	-
Female	4	7	5	16	27	38	42	29	10	4	2	-
White, both sexes	6	9	11	28	48	68	65	38	19	8	2	-
Male	3	2	6	15	24	33	27	11	9	4	-	-
Female	3	7	5	13	24	35	38	27	10	4	2	-
Black, both sexes	1	1	2	6	6	7	7	4	-	1	-	-
Male	-	1	2	3	3	4	4	3	-	1	-	-
Female	1	-	-	3	3	3	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	6	5	10	9	25	29	15	12	1	1	-
Male	2	3	1	7	2	13	13	5	8	-	-	-
Female	-	3	4	3	7	12	16	10	4	1	1	-
White, both sexes	-	-	1	2	-	10	9	11	6	1	1	-
Male	-	-	1	2	-	6	2	2	4	-	-	-
Female	-	-	-	-	-	4	7	9	2	1	1	-
Black, both sexes	2	6	4	8	9	14	20	4	6	-	-	-
Male	2	3	-	5	2	6	11	3	4	-	-	-
Female	-	3	4	3	7	8	9	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Florida												
All races, both sexes	184	299	484	776	1,135	1,569	1,805	1,456	700	218	35	-
Male	96	161	264	407	568	786	856	604	270	65	8	-
Female	88	138	220	369	567	783	949	852	430	153	27	-
White, both sexes	161	268	448	737	1,072	1,497	1,722	1,406	674	209	34	-
Male	85	145	240	383	534	739	805	576	258	62	8	-
Female	76	123	208	354	538	758	917	830	416	147	26	-
Black, both sexes	22	31	35	36	59	67	73	45	23	9	1	-
Male	11	16	23	22	32	44	44	24	11	3	-	-
Female	11	15	12	14	27	23	29	21	12	6	1	-
American Indian 1/, both sexes	-	-	-	2	2	3	2	3	-	-	-	-
Male	-	-	-	1	1	1	1	2	-	-	-	-
Female	-	-	-	1	1	2	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	2	2	8	2	3	-	-	-
Male	-	-	1	1	1	2	6	2	1	-	-	-
Female	1	-	-	-	1	-	2	-	2	-	-	-
Georgia												
All races, both sexes	96	174	256	360	514	586	584	432	189	61	6	-
Male	46	90	137	179	267	304	264	211	61	23	-	-
Female	50	84	119	181	247	282	320	221	128	38	6	-
White, both sexes	72	137	231	321	458	526	513	392	173	50	4	-
Male	34	66	118	158	231	269	227	189	50	17	-	-
Female	38	71	113	163	227	257	286	203	123	33	4	-
Black, both sexes	24	37	25	38	53	58	62	36	14	9	2	-
Male	12	24	19	20	34	35	32	21	10	5	-	-
Female	12	13	6	18	19	23	30	15	4	4	2	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	9	4	2	1	-	-
Male	-	-	-	1	1	-	5	1	1	-	-	-
Female	-	-	-	-	1	1	4	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Hawaii												
All races, both sexes	7	12	10	27	32	46	61	41	26	10	2	-
Male	4	8	4	14	17	27	38	24	13	7	-	-
Female	3	4	6	13	15	19	23	17	13	3	2	-
White, both sexes	2	3	3	9	15	19	24	13	9	2	1	-
Male	1	1	2	4	8	9	14	7	5	-	-	-
Female	1	2	1	5	7	10	10	6	4	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	9	7	18	17	26	37	28	17	7	1	-
Male	3	7	2	10	9	17	24	17	8	6	-	-
Female	2	2	5	8	8	9	13	11	9	1	1	-
Idaho												
All races, both sexes	9	22	33	53	82	141	123	114	45	9	1	-
Male	6	9	21	35	38	76	72	54	16	3	1	-
Female	3	13	12	18	44	65	51	60	29	6	-	-
White, both sexes	9	22	32	53	82	140	123	112	45	9	1	-
Male	6	9	20	35	38	75	72	53	16	3	1	-
Female	3	13	12	18	44	65	51	59	29	6	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Illinois												
All races, both sexes	111	164	275	419	601	816	939	731	386	103	14	-
Male	49	83	141	203	307	407	454	328	147	26	1	-
Female	62	81	134	216	294	409	485	403	239	77	13	-
White, both sexes	73	132	237	382	544	738	879	691	362	95	13	-
Male	31	68	119	186	271	366	418	308	137	23	-	-
Female	42	64	118	196	273	372	461	383	225	72	13	-
Black, both sexes	37	31	38	36	54	68	57	35	19	7	1	-
Male	18	15	22	17	34	35	33	16	7	3	1	-
Female	19	16	16	19	20	33	24	19	12	4	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	2	10	3	5	5	1	-	-
Male	-	-	-	-	2	6	3	4	3	-	-	-
Female	1	-	-	1	-	4	-	1	2	1	-	-
Indiana												
All races, both sexes	62	131	212	335	470	585	609	457	254	60	7	-
Male	35	67	107	171	247	295	284	206	110	21	1	-
Female	27	64	105	164	223	290	325	251	144	39	6	-
White, both sexes	54	124	201	321	451	562	588	441	250	60	6	-
Male	30	64	102	167	239	284	273	195	109	21	-	-
Female	24	60	99	154	212	278	315	246	141	39	6	-
Black, both sexes	8	6	11	13	19	22	21	16	3	-	1	-
Male	5	3	5	3	8	11	11	11	1	-	1	-
Female	3	3	6	10	11	11	10	5	2	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Iowa												
All races, both sexes	30	55	81	149	222	265	357	279	115	52	8	-
Male	12	28	43	77	125	147	200	140	54	24	2	-
Female	18	27	38	72	97	118	157	139	61	28	6	-
White, both sexes	28	55	81	149	219	263	356	278	114	52	8	-
Male	11	28	43	77	123	146	200	140	53	24	2	-
Female	17	27	38	72	96	117	156	138	61	28	6	-
Black, both sexes	1	-	-	-	2	1	-	1	-	-	-	-
Male	1	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Kansas												
All races, both sexes	23	50	81	128	199	292	278	227	117	35	8	-
Male	15	26	30	67	108	152	141	114	67	13	1	-
Female	8	24	51	61	91	140	137	113	50	22	7	-
White, both sexes	23	48	78	120	189	278	261	223	114	33	7	-
Male	15	26	29	63	102	144	133	114	67	13	1	-
Female	8	22	49	57	87	134	128	109	47	20	6	-
Black, both sexes	-	1	2	4	5	11	11	2	2	2	1	-
Male	-	-	-	2	3	7	6	-	-	-	-	-
Female	-	1	2	2	2	4	5	2	2	2	1	-
American Indian 1/, both sexes	-	-	1	4	4	2	6	2	-	-	-	-
Male	-	-	1	2	3	-	2	-	-	-	-	-
Female	-	-	-	2	1	2	4	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Kentucky												
All races, both sexes	71	101	199	292	352	403	417	287	152	37	4	-
Male	36	56	100	149	179	198	208	116	61	12	2	-
Female	35	45	99	143	173	205	209	171	91	25	2	-
White, both sexes	64	92	191	280	335	385	399	280	149	34	4	-
Male	32	51	96	143	170	190	199	113	60	12	2	-
Female	32	41	95	137	165	195	200	167	89	22	2	-
Black, both sexes	7	9	6	11	17	16	17	7	3	3	-	-
Male	4	5	3	5	9	7	8	3	1	-	-	-
Female	3	4	3	6	8	9	9	4	2	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	1	-	-	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	47	78	118	206	251	285	280	227	98	37	8	-
Male	27	44	59	111	140	147	145	107	40	15	2	-
Female	20	34	59	95	111	138	135	120	58	22	6	-
White, both sexes	27	61	91	165	209	243	248	202	84	31	5	-
Male	13	33	48	84	111	126	125	90	33	10	1	-
Female	14	28	43	81	98	117	123	112	51	21	4	-
Black, both sexes	20	17	26	41	42	42	31	23	14	6	3	-
Male	14	11	10	27	29	21	19	15	7	5	1	-
Female	6	6	16	14	13	21	12	8	7	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	2	-	-	-	-
Male	-	-	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Maine												
All races, both sexes	18	32	42	77	109	136	150	120	60	17	2	-
Male	12	17	27	32	57	73	75	59	24	3	1	-
Female	6	15	15	45	52	63	75	61	36	14	1	-
White, both sexes	18	32	41	77	109	136	149	120	60	17	2	-
Male	12	17	27	32	57	73	75	59	24	3	1	-
Female	6	15	14	45	52	63	74	61	36	14	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	34	58	105	163	247	313	369	296	141	45	5	-
Male	15	26	50	72	106	157	160	118	61	7	2	-
Female	19	32	55	91	141	156	209	178	80	38	3	-
White, both sexes	19	50	83	131	215	274	320	259	124	44	5	-
Male	7	24	39	60	86	139	136	106	54	7	2	-
Female	12	26	44	71	129	135	184	153	70	37	3	-
Black, both sexes	15	8	22	29	30	37	42	36	14	1	-	-
Male	8	2	11	12	19	17	20	11	5	-	-	-
Female	7	6	11	17	11	20	22	25	9	1	-	-
American Indian 1/, both sexes	-	-	-	2	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	7	-	3	-	-	-
Male	-	-	-	-	1	-	4	-	2	-	-	-
Female	-	-	-	1	1	1	3	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Massachusetts												
All races, both sexes	32	82	137	194	296	470	477	444	247	82	12	-
Male	17	38	59	91	153	216	199	174	73	19	3	-
Female	15	44	78	103	143	254	278	270	174	63	9	-
White, both sexes	30	80	135	188	286	458	467	435	242	78	11	-
Male	15	37	58	86	147	206	194	171	70	19	3	-
Female	15	43	77	102	139	252	273	264	172	59	8	-
Black, both sexes	2	1	2	2	8	7	8	7	3	3	1	-
Male	2	-	1	1	4	6	3	1	1	-	-	-
Female	-	1	1	1	4	1	5	6	2	3	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	4	2	5	2	2	2	1	-	-
Male	-	-	-	4	2	4	2	2	2	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Michigan												
All races, both sexes	86	164	255	412	603	865	906	659	315	66	15	-
Male	44	95	126	201	280	417	448	309	126	18	9	-
Female	42	69	129	211	323	448	458	350	189	48	6	-
White, both sexes	77	141	225	384	565	800	862	617	303	63	11	-
Male	38	80	112	183	264	384	423	287	121	17	7	-
Female	39	61	113	201	301	416	439	330	182	46	4	-
Black, both sexes	9	23	30	20	33	55	41	39	12	3	4	-
Male	6	15	14	12	13	30	23	21	5	1	2	-
Female	3	8	16	8	20	25	18	18	7	2	2	-
American Indian 1/, both sexes	-	-	-	6	4	7	2	3	-	-	-	-
Male	-	-	-	4	3	2	1	1	-	-	-	-
Female	-	-	-	2	1	5	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	1	3	1	-	-	-	-	-
Male	-	-	-	2	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Minnesota												
All races, both sexes	21	39	90	139	217	298	401	310	164	46	6	1
Male	9	22	49	71	110	169	197	132	77	16	1	1
Female	12	17	41	68	107	129	204	178	87	30	5	-
White, both sexes	20	38	86	136	213	293	393	307	160	44	6	-
Male	8	21	47	70	107	166	194	130	74	15	1	-
Female	12	17	39	66	106	127	199	177	86	29	5	-
Black, both sexes	-	-	3	2	2	3	4	2	-	-	-	-
Male	-	-	2	1	2	2	2	1	-	-	-	-
Female	-	-	1	1	-	1	2	1	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	2	2	2	-	-	-	-	-
Male	1	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	1	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	2	1	4	2	-	1
Male	-	1	-	-	-	-	-	1	3	1	-	1
Female	-	-	-	-	-	-	2	-	1	1	-	-
Mississippi												
All races, both sexes	34	69	106	142	188	259	230	185	73	33	4	1
Male	14	40	58	84	106	156	127	92	27	9	-	1
Female	20	29	48	58	82	103	103	93	46	24	4	-
White, both sexes	24	50	85	117	156	217	203	156	62	23	2	1
Male	8	28	44	68	83	130	109	76	20	6	-	1
Female	16	22	41	49	73	87	94	80	42	17	2	-
Black, both sexes	10	19	21	24	32	42	27	28	10	10	2	-
Male	6	12	14	16	23	26	18	15	7	3	-	-
Female	4	7	7	8	9	16	9	13	3	7	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Missouri												
All races, both sexes	76	128	189	287	402	576	586	386	200	63	8	-
Male	39	61	81	134	206	280	291	177	86	14	3	-
Female	37	67	108	153	196	296	295	209	114	49	5	-
White, both sexes	69	121	175	269	377	552	549	365	195	59	7	-
Male	35	56	76	126	193	269	270	165	84	13	3	-
Female	34	65	99	143	184	283	279	200	111	46	4	-
Black, both sexes	6	7	13	17	25	20	35	21	4	4	1	-
Male	4	5	4	8	13	9	21	12	2	1	-	-
Female	2	2	9	9	12	11	14	9	2	3	1	-
American Indian 1/, both sexes	1	-	1	1	-	3	1	-	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	-	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	11	16	38	46	87	112	116	98	37	11	1	-
Male	5	9	19	20	45	58	42	45	13	2	-	-
Female	6	7	19	26	42	54	74	53	24	9	1	-
White, both sexes	9	12	34	45	83	111	116	94	37	11	1	-
Male	5	8	16	19	44	57	42	41	13	2	-	-
Female	4	4	18	26	39	54	74	53	24	9	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	4	2	1	4	1	-	4	-	-	-	-
Male	-	1	2	1	1	1	-	4	-	-	-	-
Female	2	3	-	-	3	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Nebraska												
All races, both sexes	12	19	37	60	122	158	193	135	89	28	7	-
Male	7	9	17	28	68	92	98	71	31	6	4	-
Female	5	10	20	32	54	66	95	64	58	22	3	-
White, both sexes	11	19	37	57	117	155	190	133	87	28	6	-
Male	6	9	17	26	65	89	97	70	31	6	3	-
Female	5	10	20	31	52	66	93	63	56	22	3	-
Black, both sexes	1	-	-	2	2	1	2	1	2	-	1	-
Male	1	-	-	1	2	1	1	-	-	-	1	-
Female	-	-	-	1	-	-	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	2	-	-	-	-	-	-
Male	-	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Nevada												
All races, both sexes	12	44	85	101	198	188	197	132	61	19	3	-
Male	4	24	35	45	100	84	97	57	24	5	1	-
Female	8	20	50	56	98	104	100	75	37	14	2	-
White, both sexes	11	38	81	97	190	181	191	128	59	18	3	-
Male	4	22	32	42	95	80	94	55	22	5	1	-
Female	7	16	49	55	95	101	97	73	37	13	2	-
Black, both sexes	1	4	3	4	6	2	4	2	-	-	-	-
Male	-	1	3	3	4	-	2	1	-	-	-	-
Female	1	3	-	1	2	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	-	2	5	2	1	1	1	-	-
Male	-	1	-	-	1	4	1	1	1	-	-	-
Female	-	1	1	-	1	1	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
New Hampshire												
All races, both sexes	18	27	30	57	85	103	119	104	36	10	2	-
Male	9	16	16	24	37	56	65	36	11	3	-	-
Female	9	11	14	33	48	47	54	68	25	7	2	-
White, both sexes	18	27	30	57	85	102	119	104	36	10	2	-
Male	9	16	16	24	37	55	65	36	11	3	-	-
Female	9	11	14	33	48	47	54	68	25	7	2	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	64	76	130	195	326	536	608	503	242	83	10	-
Male	33	39	63	102	138	240	274	201	82	24	1	-
Female	31	37	67	93	188	296	334	302	160	59	9	-
White, both sexes	56	59	107	163	289	486	567	476	219	78	7	-
Male	29	32	48	82	119	217	252	193	74	24	1	-
Female	27	27	59	81	170	269	315	283	145	54	6	-
Black, both sexes	8	17	21	30	32	43	35	24	18	5	3	-
Male	4	7	13	18	15	20	19	8	4	-	-	-
Female	4	10	8	12	17	23	16	16	14	5	3	-
American Indian 1/, both sexes	-	-	2	-	1	1	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	4	6	6	3	5	-	-	-
Male	-	-	-	2	4	3	3	-	4	-	-	-
Female	-	-	-	-	-	3	3	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
New Mexico												
All races, both sexes	13	24	62	66	106	174	209	116	69	18	6	-
Male	4	9	30	40	48	90	101	64	27	8	3	-
Female	9	15	32	26	58	84	108	52	42	10	3	-
White, both sexes	13	23	60	65	106	172	202	114	66	18	6	-
Male	4	8	29	39	48	88	98	62	26	8	3	-
Female	9	15	31	26	58	84	104	52	40	10	3	-
Black, both sexes	-	1	-	-	-	2	1	1	2	-	-	-
Male	-	1	-	-	-	2	1	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	2	1	-	-	5	1	1	-	-	-
Male	-	-	1	1	-	-	2	1	-	-	-	-
Female	-	-	1	-	-	-	3	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
New York												
All races, both sexes	112	187	368	531	698	1,117	1,266	1,052	579	207	24	-
Male	58	95	203	254	320	540	616	445	196	57	8	-
Female	54	92	165	277	378	577	650	607	383	150	16	-
White, both sexes	86	156	321	464	633	1,028	1,162	978	534	193	22	-
Male	44	81	174	216	290	494	558	411	176	51	7	-
Female	42	75	147	248	343	534	604	567	358	142	15	-
Black, both sexes	25	29	44	62	54	74	84	55	35	10	1	-
Male	13	13	26	34	24	36	42	20	11	3	-	-
Female	12	16	18	28	30	38	42	35	24	7	1	-
American Indian 1/, both sexes	-	1	1	1	3	7	1	2	-	-	-	-
Male	-	1	1	-	1	3	1	2	-	-	-	-
Female	-	-	-	1	2	4	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	4	8	8	19	17	10	4	1	-
Male	1	-	2	4	5	7	15	12	9	3	1	-
Female	-	1	-	-	3	1	4	5	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
North Carolina												
All races, both sexes	78	189	302	443	558	744	775	493	235	65	7	-
Male	45	93	159	230	269	370	360	215	95	26	3	-
Female	33	96	143	213	289	374	415	278	140	39	4	-
White, both sexes	63	162	247	398	509	682	706	445	218	57	6	-
Male	38	74	125	204	243	339	320	189	86	20	2	-
Female	25	88	122	194	266	343	386	256	132	37	4	-
Black, both sexes	14	26	51	42	46	60	66	45	16	8	-	-
Male	6	19	33	24	24	30	38	25	8	6	-	-
Female	8	7	18	18	22	30	28	20	8	2	-	-
American Indian 1/, both sexes	1	1	2	3	2	1	2	3	1	-	1	-
Male	1	-	1	2	1	-	1	1	1	-	1	-
Female	-	1	1	1	1	1	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	6	10	15	32	60	50	53	38	11	2	-
Male	1	2	5	10	17	34	34	27	21	3	1	-
Female	1	4	5	5	15	26	16	26	17	8	1	-
White, both sexes	1	6	8	14	31	58	49	52	38	11	2	-
Male	1	2	4	10	17	34	33	27	21	3	1	-
Female	-	4	4	4	14	24	16	25	17	8	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	2	1	1	2	1	1	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	1	1	1	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Ohio												
All races, both sexes	138	234	383	571	806	1,160	1,157	869	424	120	20	-
Male	80	118	205	276	397	580	564	400	165	33	6	-
Female	58	116	178	295	409	580	593	469	259	87	14	-
White, both sexes	115	196	352	520	757	1,067	1,091	824	405	116	19	-
Male	67	96	184	245	369	531	527	373	160	33	6	-
Female	48	100	168	275	388	536	564	451	245	83	13	-
Black, both sexes	23	38	30	51	48	93	62	42	18	4	1	-
Male	13	22	21	31	28	49	34	26	4	-	-	-
Female	10	16	9	20	20	44	28	16	14	4	1	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	3	1	-	-	-
Male	-	-	-	-	-	-	3	1	1	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Oklahoma												
All races, both sexes	58	114	178	267	350	413	355	238	115	34	4	-
Male	27	64	86	127	189	217	185	106	51	14	1	-
Female	31	50	92	140	161	196	170	132	64	20	3	-
White, both sexes	54	106	162	245	325	386	329	228	110	31	3	-
Male	26	59	76	118	181	199	171	101	50	13	1	-
Female	28	47	86	127	144	187	158	127	60	18	2	-
Black, both sexes	3	4	4	11	8	7	10	6	3	1	1	-
Male	1	2	2	6	4	3	6	3	1	1	-	-
Female	2	2	2	5	4	4	4	3	2	-	1	-
American Indian 1/, both sexes	1	4	12	11	17	20	15	4	2	2	-	-
Male	-	3	8	3	4	15	8	2	-	-	-	-
Female	1	1	4	8	13	5	7	2	2	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Oregon												
All races, both sexes	31	84	113	156	268	350	342	270	130	26	3	-
Male	21	42	55	75	127	170	165	127	56	10	2	-
Female	10	42	58	81	141	180	177	143	74	16	1	-
White, both sexes	30	79	107	153	262	340	337	268	128	24	3	-
Male	20	40	52	73	124	164	162	126	55	9	2	-
Female	10	39	55	80	138	176	175	142	73	15	1	-
Black, both sexes	-	3	3	1	3	1	3	1	1	1	-	-
Male	-	1	2	1	2	-	2	-	-	1	-	-
Female	-	2	1	-	1	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	2	3	2	3	4	1	-	1	-	-	-
Male	-	1	1	1	1	1	-	-	1	-	-	-
Female	-	1	2	1	2	3	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	5	1	1	-	1	-	-
Male	1	-	-	-	-	5	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Pennsylvania												
All races, both sexes	91	177	306	435	687	1,050	1,202	937	446	126	19	-
Male	40	90	166	227	329	531	545	410	172	36	4	-
Female	51	87	140	208	358	519	657	527	274	90	15	-
White, both sexes	71	150	277	392	639	982	1,154	904	424	118	19	-
Male	31	78	151	205	306	494	520	396	167	33	4	-
Female	40	72	126	187	333	488	634	508	257	85	15	-
Black, both sexes	18	27	28	38	48	65	43	32	22	7	-	-
Male	8	12	15	18	23	34	20	14	5	3	-	-
Female	10	15	13	20	25	31	23	18	17	4	-	-
American Indian 1/, both sexes	1	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	4	-	2	5	1	-	1	-	-
Male	1	-	-	3	-	2	5	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Rhode Island												
All races, both sexes	7	12	26	34	65	76	112	92	33	10	2	-
Male	4	7	6	13	34	39	39	39	15	1	1	-
Female	3	5	20	21	31	37	73	53	18	9	1	-
White, both sexes	7	12	26	32	61	76	108	89	33	10	2	-
Male	4	7	6	12	31	39	38	37	15	1	1	-
Female	3	5	20	20	30	37	70	52	18	9	1	-
Black, both sexes	-	-	-	2	3	-	3	3	-	-	-	-
Male	-	-	-	1	2	-	-	2	-	-	-	-
Female	-	-	-	1	1	-	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	55	98	164	220	296	344	351	216	95	35	5	-
Male	23	56	84	119	158	188	178	87	40	8	-	-
Female	32	42	80	101	138	156	173	129	55	27	5	-
White, both sexes	38	80	133	186	260	312	318	192	87	31	4	-
Male	15	43	66	97	135	169	159	76	38	8	-	-
Female	23	37	67	89	125	143	159	116	49	23	4	-
Black, both sexes	17	18	29	33	36	32	31	23	8	4	1	-
Male	8	13	16	21	23	19	18	10	2	-	-	-
Female	9	5	13	12	13	13	13	13	6	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	-	-	1	-	-	-	-
Male	-	-	2	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
South Dakota												
All races, both sexes	5	13	18	25	54	63	77	58	41	14	1	-
Male	2	4	9	10	37	32	48	33	18	7	-	-
Female	3	9	9	15	17	31	29	25	23	7	1	-
White, both sexes	4	10	16	25	50	62	76	57	40	14	1	-
Male	1	3	7	10	33	31	47	33	18	7	-	-
Female	3	7	9	15	17	31	29	24	22	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	3	2	-	4	1	1	1	1	-	-	-
Male	1	1	2	-	4	1	1	-	-	-	-	-
Female	-	2	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	86	154	239	358	446	543	496	325	166	45	6	-
Male	42	71	119	175	229	273	262	142	66	19	-	-
Female	44	83	120	183	217	270	234	183	100	26	6	-
White, both sexes	71	136	216	334	413	516	467	309	155	43	5	-
Male	35	65	105	161	212	255	247	133	60	17	-	-
Female	36	71	111	173	201	261	220	176	95	26	5	-
Black, both sexes	15	18	22	24	32	27	29	16	11	2	1	-
Male	7	6	14	14	17	18	15	9	6	2	-	-
Female	8	12	8	10	15	9	14	7	5	-	1	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Texas												
All races, both sexes	154	312	514	802	1,071	1,425	1,450	1,017	500	130	24	-
Male	86	140	263	395	551	702	709	475	200	40	6	-
Female	68	172	251	407	520	723	741	542	300	90	18	-
White, both sexes	123	275	477	722	990	1,338	1,379	966	483	124	22	-
Male	68	129	241	353	506	659	671	446	192	37	6	-
Female	55	146	236	369	484	679	708	520	291	87	16	-
Black, both sexes	29	34	36	75	73	78	62	47	15	6	2	-
Male	17	11	21	40	41	39	34	26	6	3	-	-
Female	12	23	15	35	32	39	28	21	9	3	2	-
American Indian 1/, both sexes	-	1	-	2	1	2	4	-	-	-	-	-
Male	-	-	-	1	1	1	2	-	-	-	-	-
Female	-	1	-	1	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	3	7	7	5	4	2	-	-	-
Male	1	-	1	1	3	3	2	3	2	-	-	-
Female	1	2	-	2	4	4	3	1	-	-	-	-
Utah												
All races, both sexes	12	32	33	63	87	117	109	82	28	7	1	-
Male	11	9	14	36	56	66	51	41	17	1	-	-
Female	1	23	19	27	31	51	58	41	11	6	1	-
White, both sexes	11	32	33	63	87	113	105	82	28	7	1	-
Male	10	9	14	36	56	64	49	41	17	1	-	-
Female	1	23	19	27	31	49	56	41	11	6	1	-
Black, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	2	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Vermont												
All races, both sexes	3	11	24	18	39	58	84	40	23	12	1	-
Male	1	6	15	9	18	30	36	15	9	3	-	-
Female	2	5	9	9	21	28	48	25	14	9	1	-
White, both sexes	3	11	24	18	39	58	84	40	23	12	1	-
Male	1	6	15	9	18	30	36	15	9	3	-	-
Female	2	5	9	9	21	28	48	25	14	9	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	43	107	162	236	408	477	531	395	182	59	12	-
Male	26	54	80	114	194	218	258	169	71	19	5	-
Female	17	53	82	122	214	259	273	226	111	40	7	-
White, both sexes	34	94	139	203	365	429	484	370	165	52	11	-
Male	20	47	70	93	167	193	231	156	63	19	4	-
Female	14	47	69	110	198	236	253	214	102	33	7	-
Black, both sexes	9	13	23	33	39	44	42	25	16	7	1	-
Male	6	7	10	21	24	21	25	13	7	-	1	-
Female	3	6	13	12	15	23	17	12	9	7	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	4	5	-	1	-	-	-
Male	-	-	-	-	2	4	2	-	1	-	-	-
Female	-	-	-	-	1	-	3	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Washington												
All races, both sexes	67	100	159	245	333	482	538	421	203	47	5	1
Male	37	45	72	121	168	228	244	176	74	17	1	1
Female	30	55	87	124	165	254	294	245	129	30	4	-
White, both sexes	63	93	148	237	322	456	520	407	197	46	5	1
Male	35	45	67	116	162	216	232	170	73	16	1	1
Female	28	48	81	121	160	240	288	237	124	30	4	-
Black, both sexes	3	1	7	2	1	6	8	3	1	-	-	-
Male	2	-	4	1	1	3	6	-	-	-	-	-
Female	1	1	3	1	-	3	2	3	1	-	-	-
American Indian 1/, both sexes	-	5	3	2	5	7	1	3	2	-	-	-
Male	-	-	1	1	1	-	1	2	-	-	-	-
Female	-	5	2	1	4	7	-	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	4	5	13	9	8	3	1	-	-
Male	-	-	-	3	4	9	5	4	1	1	-	-
Female	1	1	1	1	1	4	4	4	2	-	-	-
West Virginia												
All races, both sexes	32	55	79	139	199	259	224	133	89	18	2	-
Male	15	27	39	56	102	133	110	58	33	5	1	-
Female	17	28	40	83	97	126	114	75	56	13	1	-
White, both sexes	31	55	77	134	198	259	221	132	89	17	1	-
Male	15	27	37	54	101	133	108	57	33	5	-	-
Female	16	28	40	80	97	126	113	75	56	12	1	-
Black, both sexes	1	-	2	5	1	-	3	1	-	1	1	-
Male	-	-	2	2	1	-	2	1	-	-	1	-
Female	1	-	-	3	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Wisconsin												
All races, both sexes	41	77	118	201	273	467	493	370	210	54	3	-
Male	23	39	68	96	147	236	238	165	88	14	-	-
Female	18	38	50	105	126	231	255	205	122	40	3	-
White, both sexes	33	71	109	194	265	452	487	365	206	52	3	-
Male	18	36	62	91	143	230	237	163	87	14	-	-
Female	15	35	47	103	122	222	250	202	119	38	3	-
Black, both sexes	7	5	7	5	8	9	3	2	4	-	-	-
Male	5	3	5	4	4	2	1	-	1	-	-	-
Female	2	2	2	1	4	7	2	2	3	-	-	-
American Indian 1/, both sexes	1	1	2	2	-	2	2	2	-	1	-	-
Male	-	-	1	1	-	2	-	1	-	-	-	-
Female	1	1	1	1	-	-	2	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	4	1	1	-	1	-	-
Male	-	-	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	1	-	-
Wyoming												
All races, both sexes	8	16	19	27	47	60	76	55	21	4	-	-
Male	3	9	11	11	28	33	41	25	11	-	-	-
Female	5	7	8	16	19	27	35	30	10	4	-	-
White, both sexes	7	15	19	26	46	60	75	55	20	4	-	-
Male	3	8	11	10	27	33	41	25	11	-	-	-
Female	4	7	8	16	19	27	34	30	9	4	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	1	1	-	1	-	-	-	-	-
Male	-	1	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
United States												
All races, both sexes	400	14	6	1	1	2	4	1	2	2	1	4
Male	160	9	6	-	1	2	2	1	2	1	1	3
Female	240	5	-	1	-	-	2	-	-	1	-	1
White, both sexes	354	6	5	1	-	2	4	1	2	2	1	2
Male	142	4	5	-	-	2	2	1	2	1	1	1
Female	212	2	-	1	-	-	2	-	-	1	-	1
Black, both sexes	38	7	1	-	1	-	-	-	-	-	-	1
Male	15	4	1	-	1	-	-	-	-	-	-	1
Female	23	3	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	1	-	-	-	-	-	-	-	-	-	1
Male	3	1	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Arkansas												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	35	-	1	-	-	1	-	-	-	-	-	-
Male	13	-	1	-	-	1	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	1	-	-	1	-	-	-	-	-	-
Male	13	-	1	-	-	1	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Colorado												
All races, both sexes	7	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Florida												
All races, both sexes	23	2	-	-	-	-	-	-	2	-	-	-
Male	9	2	-	-	-	-	-	-	2	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	1	-	-	-	-	-	-	2	-	-	-
Male	7	1	-	-	-	-	-	-	2	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	9	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Hawaii												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Illinois												
All races, both sexes	14	-	-	1	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	1	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	10	1	-	-	-	-	-	-	-	-	1	-
Male	4	1	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	1	-	-	-	-	-	-	-	-	1	-
Male	4	1	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Iowa												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Kentucky												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	7	2	-	-	-	-	-	-	-	-	-	-
Male	4	2	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Maine												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Massachusetts												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Minnesota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Missouri												
All races, both sexes	12	1	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	8	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	1	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	7	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Nebraska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
New Hampshire												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
New Mexico												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	11	-	-	-	1	-	-	-	-	-	-	-
Male	7	-	-	-	1	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
North Carolina												
All races, both sexes	13	1	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Ohio												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Oregon												
All races, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	33	-	1	-	-	-	-	-	-	1	-	-
Male	16	-	1	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	30	-	1	-	-	-	-	-	-	1	-	-
Male	14	-	1	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Rhode Island												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
South Dakota												
All races, both sexes	6	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	13	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Texas												
All races, both sexes	27	4	2	-	-	1	-	-	-	-	-	1
Male	12	1	2	-	-	1	-	-	-	-	-	1
Female	15	3	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	1	1	-	-	1	-	-	-	-	-	-
Male	10	1	1	-	-	1	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	3	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	3	3	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	8	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
Wisconsin												
All races, both sexes	7	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
United States												
All races, both sexes	10	13	14	17	23	41	67	79	56	31	11	-
Male	4	6	5	7	9	16	27	27	18	11	2	-
Female	6	7	9	10	14	25	40	52	38	20	9	-
White, both sexes	7	12	13	15	20	36	60	73	52	31	9	-
Male	3	6	4	7	8	14	25	26	17	11	2	-
Female	4	6	9	8	12	22	35	47	35	20	7	-
Black, both sexes	3	1	1	2	3	5	4	5	3	-	1	-
Male	1	-	1	-	1	2	1	1	1	-	-	-
Female	2	1	-	2	2	3	3	4	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	-	1	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	1	-
Alabama												
All races, both sexes	-	1	-	-	-	2	1	1	2	1	1	-
Male	-	1	-	-	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	1	1	1	-
White, both sexes	-	1	-	-	-	2	1	1	2	1	1	-
Male	-	1	-	-	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Arkansas												
All races, both sexes	-	1	2	1	1	2	-	-	1	-	-	-
Male	-	1	1	1	-	1	-	-	1	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
White, both sexes	-	1	2	1	1	2	-	-	1	-	-	-
Male	-	1	1	1	-	1	-	-	1	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	3	-	2	1	3	3	12	4	5	-	-
Male	-	-	-	-	-	1	-	8	1	1	-	-
Female	-	3	-	2	1	2	3	4	3	4	-	-
White, both sexes	-	3	-	2	1	3	3	12	4	5	-	-
Male	-	-	-	-	-	1	-	8	1	1	-	-
Female	-	3	-	2	1	2	3	4	3	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Colorado												
All races, both sexes	-	-	-	-	-	1	1	2	-	1	1	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	1	-
White, both sexes	-	-	-	-	-	1	1	2	-	1	1	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	1	1	1	1	4	4	1	-	-
Male	-	-	-	-	1	1	-	1	2	1	-	-
Female	-	-	-	1	-	-	1	3	2	-	-	-
White, both sexes	-	-	-	1	-	1	1	4	4	1	-	-
Male	-	-	-	-	-	1	-	1	2	1	-	-
Female	-	-	-	1	-	-	1	3	2	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Florida												
All races, both sexes	1	1	1	1	1	1	7	5	1	-	-	-
Male	1	-	-	-	1	-	2	-	1	-	-	-
Female	-	1	1	1	-	1	5	5	-	-	-	-
White, both sexes	1	1	1	1	1	1	5	5	1	-	-	-
Male	1	-	-	-	1	-	1	-	1	-	-	-
Female	-	1	1	1	-	1	4	5	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	2	5	1	-	-	-
Male	-	-	-	-	-	-	2	2	-	-	-	-
Female	-	-	-	-	-	-	-	3	1	-	-	-
White, both sexes	-	-	-	-	-	-	2	4	1	-	-	-
Male	-	-	-	-	-	-	2	2	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	1	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Illinois												
All races, both sexes	-	1	-	1	2	2	2	3	1	1	-	-
Male	-	1	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	1	2	2	1	2	-	1	-	-
White, both sexes	-	1	-	1	2	2	2	3	-	1	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	2	2	1	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	1	-	1	-	1	1	2	2	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	1	-	1	-	-	-	2	2	-	-	-
White, both sexes	-	1	-	1	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	1	-	1	-	-	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Iowa												
All races, both sexes	-	-	1	-	-	1	1	2	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	-	2	-	1	-	-
White, both sexes	-	-	1	-	-	1	1	2	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	-	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	1	-	-	1	-	2	1	-
Male	-	-	-	-	-	-	-	-	-	2	-	-
Female	-	-	-	-	1	-	-	1	-	-	1	-
White, both sexes	-	-	-	-	1	-	-	1	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	2	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Kentucky												
All races, both sexes	2	-	1	-	-	-	1	2	4	1	-	-
Male	-	-	1	-	-	-	-	1	-	1	-	-
Female	2	-	-	-	-	-	1	1	4	-	-	-
White, both sexes	2	-	1	-	-	-	1	2	4	1	-	-
Male	-	-	1	-	-	-	-	1	-	1	-	-
Female	2	-	-	-	-	-	1	1	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	2	-	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	1	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	2	-	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	2	-	2	2	6	-	-	-
Male	-	-	-	-	1	-	1	1	4	-	-	-
Female	-	-	-	-	1	-	1	1	2	-	-	-
White, both sexes	-	-	-	-	2	-	1	1	5	-	-	-
Male	-	-	-	-	1	-	1	-	4	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Massachusetts												
All races, both sexes	-	-	-	1	-	2	2	1	1	1	-	-
Male	-	-	-	1	-	1	-	1	-	1	-	-
Female	-	-	-	-	-	1	2	-	1	-	-	-
White, both sexes	-	-	-	1	-	2	2	1	1	1	-	-
Male	-	-	-	1	-	1	-	1	-	1	-	-
Female	-	-	-	-	-	1	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	2	1	-	-	-	4	4	1	-	1	-
Male	-	2	-	-	-	-	2	-	-	-	1	-
Female	1	-	1	-	-	-	2	4	1	-	-	-
White, both sexes	1	2	1	-	-	-	4	4	1	-	1	-
Male	-	2	-	-	-	-	2	-	-	-	1	-
Female	1	-	1	-	-	-	2	4	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Minnesota												
All races, both sexes	-	-	-	-	1	1	1	-	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	1	-	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	-	1	2	1	-	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
Black, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Missouri												
All races, both sexes	-	-	1	1	1	1	4	1	-	1	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	3	1	-	1	-	-
White, both sexes	-	-	1	1	1	1	3	1	-	1	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	2	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
New Hampshire												
All races, both sexes	-	-	-	-	-	1	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	1	1	1	-	1	2	-	2	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	1	1	-	-	1	1	-	2	-	-	-
White, both sexes	-	-	1	1	-	1	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	1	-	-	-
Black, both sexes	-	1	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
New Mexico												
All races, both sexes	1	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	1	1	1	-	-	-
White, both sexes	1	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	1	-	-	2	1	3	3	-	-	-
Male	-	-	1	-	-	1	1	1	2	-	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-
White, both sexes	-	-	1	-	-	1	1	3	3	-	-	-
Male	-	-	1	-	-	1	1	1	2	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
North Carolina												
All races, both sexes	-	-	-	2	1	2	3	1	2	1	-	-
Male	-	-	-	1	-	-	2	-	1	-	-	-
Female	-	-	-	1	1	2	1	1	1	1	-	-
White, both sexes	-	-	-	1	1	1	3	-	2	1	-	-
Male	-	-	-	1	-	-	2	-	1	-	-	-
Female	-	-	-	-	1	1	1	-	1	1	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Ohio												
All races, both sexes	-	-	-	-	-	2	1	4	3	4	-	-
Male	-	-	-	-	-	1	1	2	1	1	-	-
Female	-	-	-	-	-	1	-	2	2	3	-	-
White, both sexes	-	-	-	-	-	1	1	4	3	4	-	-
Male	-	-	-	-	-	1	1	2	1	1	-	-
Female	-	-	-	-	-	-	-	2	2	3	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	1	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Oregon												
All races, both sexes	-	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	-	1	-	2	5	9	4	5	2	-	-
Male	3	-	-	-	1	3	5	1	2	-	-	-
Female	-	-	1	-	1	2	4	3	3	2	-	-
White, both sexes	2	-	1	-	2	4	9	3	5	2	-	-
Male	2	-	-	-	1	2	5	1	2	-	-	-
Female	-	-	1	-	1	2	4	2	3	2	-	-
Black, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Rhode Island												
All races, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	1	-	1	-	-	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
South Dakota												
All races, both sexes	-	-	-	-	-	1	1	1	-	2	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	1	1	-	-	1	-	-
White, both sexes	-	-	-	-	-	1	1	1	-	2	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	1	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	-	-	1	1	-	1	3	4	1	-	-
Male	-	-	-	-	1	-	-	2	-	-	-	-
Female	1	-	-	1	-	-	1	1	4	1	-	-
White, both sexes	-	-	-	1	1	-	1	3	4	1	-	-
Male	-	-	-	-	1	-	-	2	-	-	-	-
Female	-	-	-	1	-	-	1	1	4	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Texas												
All races, both sexes	-	-	1	1	1	2	4	4	2	1	3	-
Male	-	-	1	1	-	-	3	1	-	-	1	-
Female	-	-	-	-	1	2	1	3	2	1	2	-
White, both sexes	-	-	1	1	1	2	4	4	2	1	3	-
Male	-	-	1	1	-	-	3	1	-	-	1	-
Female	-	-	-	-	1	2	1	3	2	1	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Vermont												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	3	2	2	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	2	1	2	-	-	-	-	-
White, both sexes	-	-	-	-	2	2	2	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Wisconsin												
All races, both sexes	-	-	1	-	1	-	1	1	1	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	-	1	-	1	-	1	-
White, both sexes	-	-	1	-	1	-	1	1	1	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	-	1	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
United States												
All races, both sexes	340	5	6	-	-	2	1	2	-	4	3	6
Male	139	3	3	-	-	1	1	1	-	4	3	2
Female	201	2	3	-	-	1	-	1	-	-	-	4
White, both sexes	303	3	4	-	-	2	-	1	-	3	2	5
Male	117	2	2	-	-	1	-	1	-	3	2	1
Female	186	1	2	-	-	1	-	-	-	-	-	4
Black, both sexes	28	1	2	-	-	-	-	1	-	1	1	1
Male	16	-	1	-	-	-	-	-	-	1	1	1
Female	12	1	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	1	-	-	-	-	1	-	-	-	-	-
Male	6	1	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Arkansas												
All races, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	43	1	2	-	-	-	-	1	-	-	-	-
Male	14	1	1	-	-	-	-	-	-	-	-	-
Female	29	-	1	-	-	-	-	1	-	-	-	-
White, both sexes	37	-	2	-	-	-	-	-	-	-	-	-
Male	11	-	1	-	-	-	-	-	-	-	-	-
Female	26	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Colorado												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Florida												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Hawaii												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Illinois												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Iowa												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Kentucky												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Maine												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Massachusetts												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	22	-	-	-	-	-	-	-	-	1	1	1
Male	16	-	-	-	-	-	-	-	-	1	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	1	1	1
Male	16	-	-	-	-	-	-	-	-	1	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Minnesota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Missouri												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Nebraska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
New Hampshire												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	25	1	-	-	-	-	-	-	-	1	2	1
Male	9	1	-	-	-	-	-	-	-	1	2	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	1	-	-	-	-	-	-	-	1	1	-
Male	6	1	-	-	-	-	-	-	-	1	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
North Carolina												
All races, both sexes	10	-	-	-	-	1	-	1	-	1	-	-
Male	5	-	-	-	-	-	-	1	-	1	-	-
Female	5	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	1	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Ohio												
All races, both sexes	23	1	-	-	-	-	-	-	-	1	-	-
Male	9	1	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	1	-	-	-	-	-	-	-	1	-	-
Male	7	1	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Oregon												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	15	1	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	13	1	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Texas												
All races, both sexes	22	1	1	-	-	1	-	-	-	-	-	-
Male	13	-	1	-	-	1	-	-	-	-	-	-
Female	9	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	1	-	-	1	-	-	-	-	-	-
Male	12	-	1	-	-	1	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	10	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Washington												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic bronchitis (J41-J42)												
Wisconsin												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
United States												
All races, both sexes	7	10	12	18	31	41	51	66	43	25	7	-
Male	4	7	5	9	15	17	22	22	15	3	2	-
Female	3	3	7	9	16	24	29	44	28	22	5	-
White, both sexes	7	5	12	17	28	37	45	63	43	20	6	-
Male	4	3	5	8	13	13	19	21	15	3	1	-
Female	3	2	7	9	15	24	26	42	28	17	5	-
Black, both sexes	-	5	-	1	3	3	4	2	-	3	-	-
Male	-	4	-	1	2	3	2	-	-	-	-	-
Female	-	1	-	-	1	-	2	2	-	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	2	1	-
Male	-	-	-	-	-	1	1	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	2	-	-
Alabama												
All races, both sexes	1	2	-	-	-	-	2	-	-	-	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Arkansas												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	1	1	1	7	6	12	3	5	3	-
Male	-	-	-	1	1	1	3	4	-	-	2	-
Female	-	-	1	-	-	6	3	8	3	5	1	-
White, both sexes	-	-	1	1	1	6	6	12	3	3	2	-
Male	-	-	-	1	1	-	3	4	-	-	1	-
Female	-	-	1	-	-	6	3	8	3	3	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Florida												
All races, both sexes	1	-	1	3	1	6	5	4	5	1	-	-
Male	-	-	-	1	1	-	2	1	1	1	-	-
Female	1	-	1	2	-	6	3	3	4	-	-	-
White, both sexes	1	-	1	2	1	6	5	4	5	1	-	-
Male	-	-	-	-	1	-	2	1	1	1	-	-
Female	1	-	1	2	-	6	3	3	4	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	1	1	1	1	4	3	-	1	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	1	1	-	1	3	2	-	1	-
White, both sexes	-	-	-	1	1	1	1	4	3	-	1	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	1	1	-	1	3	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Illinois												
All races, both sexes	-	-	-	-	2	2	-	3	2	1	-	-
Male	-	-	-	-	-	2	-	1	1	-	-	-
Female	-	-	-	-	2	-	-	2	1	1	-	-
White, both sexes	-	-	-	-	2	2	-	2	2	1	-	-
Male	-	-	-	-	-	2	-	-	1	-	-	-
Female	-	-	-	-	2	-	-	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	1	1	2	2	2	3	1	-	-
Male	-	-	-	1	-	1	1	1	1	-	-	-
Female	-	-	-	-	1	1	1	1	2	1	-	-
White, both sexes	-	-	-	1	1	1	2	2	3	1	-	-
Male	-	-	-	1	-	-	1	1	1	-	-	-
Female	-	-	-	-	1	1	1	1	2	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Iowa												
All races, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Kentucky												
All races, both sexes	-	-	-	2	-	-	1	1	1	-	-	-
Male	-	-	-	1	-	-	1	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	2	-	-	1	1	1	-	-	-
Male	-	-	-	1	-	-	1	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Massachusetts												
All races, both sexes	-	-	1	-	2	2	-	4	-	-	-	-
Male	-	-	-	-	1	1	-	2	-	-	-	-
Female	-	-	1	-	1	1	-	2	-	-	-	-
White, both sexes	-	-	1	-	2	2	-	4	-	-	-	-
Male	-	-	-	-	1	1	-	2	-	-	-	-
Female	-	-	1	-	1	1	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	2	-	1	4	1	4	4	1	-	-	-
Male	1	2	-	-	3	1	3	2	1	-	-	-
Female	1	-	-	1	1	-	1	2	-	-	-	-
White, both sexes	2	2	-	1	4	1	3	4	1	-	-	-
Male	1	2	-	-	3	1	3	2	1	-	-	-
Female	1	-	-	1	1	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	1	2	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	1	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Missouri												
All races, both sexes	-	-	-	-	-	-	3	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	3	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
New Hampshire												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	1	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	1	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	1	1	2	3	3	-	-	1	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	1	2	1	3	-	-	1	-
White, both sexes	-	-	-	1	1	2	1	3	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	2	-	3	-	-	1	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	1	1	1	1	1	4	4	4	3	-	-
Male	-	1	-	1	-	-	-	1	1	-	-	-
Female	-	-	1	-	1	1	4	3	3	3	-	-
White, both sexes	-	-	1	1	1	1	4	3	4	3	-	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	-	-	1	-	1	1	4	2	3	3	-	-
Black, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
North Carolina												
All races, both sexes	-	-	1	1	-	1	1	1	1	1	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	1	-	-
White, both sexes	-	-	1	1	-	-	1	1	1	1	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Ohio												
All races, both sexes	1	1	1	-	2	3	2	4	4	3	-	-
Male	-	1	-	-	1	2	1	-	2	-	-	-
Female	1	-	1	-	1	1	1	4	2	3	-	-
White, both sexes	1	-	1	-	1	3	2	4	4	3	-	-
Male	-	-	-	-	-	2	1	-	2	-	-	-
Female	1	-	1	-	1	1	1	4	2	3	-	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	1	-	-	1	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Oregon												
All races, both sexes	-	-	-	-	1	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	2	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	3	2	4	1	3	-	-	-
Male	-	-	-	-	2	-	2	-	2	-	-	-
Female	-	-	-	-	1	2	2	1	1	-	-	-
White, both sexes	-	-	-	-	2	2	3	1	3	-	-	-
Male	-	-	-	-	1	-	1	-	2	-	-	-
Female	-	-	-	-	1	2	2	1	1	-	-	-
Black, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	2	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	1	1	-	1	-	-	1	-	1	-
Male	-	-	-	1	-	1	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	1	1	-	1	-	-	1	-	1	-
Male	-	-	-	1	-	1	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Texas												
All races, both sexes	-	1	3	1	2	5	-	4	-	3	-	-
Male	-	-	3	1	1	3	-	2	-	1	-	-
Female	-	1	-	-	1	2	-	2	-	2	-	-
White, both sexes	-	-	3	1	2	4	-	3	-	3	-	-
Male	-	-	3	1	1	2	-	2	-	1	-	-
Female	-	-	-	-	1	2	-	1	-	2	-	-
Black, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	1	-	2	2	-	1	3	-	-	-	-
Male	-	-	-	1	1	-	-	3	-	-	-	-
Female	-	1	-	1	1	-	1	-	-	-	-	-
White, both sexes	-	1	-	2	2	-	1	3	-	-	-	-
Male	-	-	-	1	1	-	-	3	-	-	-	-
Female	-	1	-	1	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Washington												
All races, both sexes	2	-	-	-	2	-	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	2	-	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Wisconsin												
All races, both sexes	-	-	-	-	1	2	-	1	2	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	1	1	-	-	-
White, both sexes	-	-	-	-	1	2	-	1	2	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
United States												
All races, both sexes	12,551	2	-	1	1	-	1	2	7	10	76	158
Male	6,318	2	-	1	-	-	-	1	5	6	43	80
Female	6,233	-	-	-	1	-	1	1	2	4	33	78
White, both sexes	11,689	1	-	1	-	-	1	2	5	7	59	134
Male	5,784	1	-	1	-	-	-	1	3	4	34	64
Female	5,905	-	-	-	-	-	1	1	2	3	25	70
Black, both sexes	707	1	-	-	-	-	-	-	1	3	17	20
Male	430	1	-	-	-	-	-	-	1	2	9	13
Female	277	-	-	-	-	-	-	-	-	1	8	7
American Indian 1/, both sexes	32	-	-	-	-	-	-	-	1	-	-	1
Male	16	-	-	-	-	-	-	-	1	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	123	-	-	-	1	-	-	-	-	-	-	3
Male	88	-	-	-	-	-	-	-	-	-	-	2
Female	35	-	-	-	1	-	-	-	-	-	-	1
Alabama												
All races, both sexes	188	-	-	-	-	-	-	-	-	1	3	3
Male	101	-	-	-	-	-	-	-	-	-	1	1
Female	87	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	167	-	-	-	-	-	-	-	-	1	3	3
Male	85	-	-	-	-	-	-	-	-	-	1	1
Female	82	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Alaska												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	352	-	-	-	-	-	-	-	-	-	1	4
Male	173	-	-	-	-	-	-	-	-	-	-	2
Female	179	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	348	-	-	-	-	-	-	-	-	-	1	4
Male	170	-	-	-	-	-	-	-	-	-	-	2
Female	178	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Arkansas												
All races, both sexes	182	-	-	-	-	-	-	-	1	-	3	2
Male	106	-	-	-	-	-	-	-	1	-	2	1
Female	76	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	173	-	-	-	-	-	-	-	-	-	3	2
Male	101	-	-	-	-	-	-	-	-	-	2	1
Female	72	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,213	-	-	-	-	-	-	-	-	-	5	13
Male	590	-	-	-	-	-	-	-	-	-	3	6
Female	623	-	-	-	-	-	-	-	-	-	2	7
White, both sexes	1,099	-	-	-	-	-	-	-	-	-	3	9
Male	510	-	-	-	-	-	-	-	-	-	2	3
Female	589	-	-	-	-	-	-	-	-	-	1	6
Black, both sexes	69	-	-	-	-	-	-	-	-	-	2	3
Male	44	-	-	-	-	-	-	-	-	-	1	2
Female	25	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	41	-	-	-	-	-	-	-	-	-	-	1
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Colorado												
All races, both sexes	154	-	-	-	-	-	-	-	-	-	1	1
Male	79	-	-	-	-	-	-	-	-	-	-	-
Female	75	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	145	-	-	-	-	-	-	-	-	-	-	1
Male	77	-	-	-	-	-	-	-	-	-	-	-
Female	68	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	155	-	-	-	-	-	-	-	-	-	-	1
Male	84	-	-	-	-	-	-	-	-	-	-	1
Female	71	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	144	-	-	-	-	-	-	-	-	-	-	1
Male	78	-	-	-	-	-	-	-	-	-	-	1
Female	66	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Delaware												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Florida												
All races, both sexes	846	-	-	-	-	-	-	-	-	1	1	10
Male	399	-	-	-	-	-	-	-	-	-	-	4
Female	447	-	-	-	-	-	-	-	-	1	1	6
White, both sexes	804	-	-	-	-	-	-	-	-	-	1	9
Male	369	-	-	-	-	-	-	-	-	-	-	3
Female	435	-	-	-	-	-	-	-	-	-	1	6
Black, both sexes	41	-	-	-	-	-	-	-	-	1	-	1
Male	29	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	362	-	-	1	-	-	-	1	-	-	2	6
Male	199	-	-	1	-	-	-	-	-	-	1	3
Female	163	-	-	-	-	-	-	1	-	-	1	3
White, both sexes	330	-	-	1	-	-	-	1	-	-	2	3
Male	173	-	-	1	-	-	-	-	-	-	1	1
Female	157	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	31	-	-	-	-	-	-	-	-	-	-	3
Male	25	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Hawaii												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	82	-	-	-	-	-	-	-	-	-	-	-
Male	43	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	82	-	-	-	-	-	-	-	-	-	-	-
Male	43	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Illinois												
All races, both sexes	566	-	-	-	-	-	-	-	-	1	5	6
Male	278	-	-	-	-	-	-	-	-	1	2	1
Female	288	-	-	-	-	-	-	-	-	-	3	5
White, both sexes	518	-	-	-	-	-	-	-	-	-	3	6
Male	246	-	-	-	-	-	-	-	-	-	1	1
Female	272	-	-	-	-	-	-	-	-	-	2	5
Black, both sexes	44	-	-	-	-	-	-	-	-	1	2	-
Male	29	-	-	-	-	-	-	-	-	1	1	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	364	-	-	-	-	-	-	-	-	2	2	9
Male	191	-	-	-	-	-	-	-	-	1	1	7
Female	173	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	350	-	-	-	-	-	-	-	-	2	2	7
Male	186	-	-	-	-	-	-	-	-	1	1	6
Female	164	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Iowa												
All races, both sexes	105	-	-	-	-	-	-	-	-	-	-	1
Male	60	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	104	-	-	-	-	-	-	-	-	-	-	1
Male	60	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	129	-	-	-	-	-	-	-	-	-	1	2
Male	62	-	-	-	-	-	-	-	-	-	1	1
Female	67	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	122	-	-	-	-	-	-	-	-	-	1	2
Male	60	-	-	-	-	-	-	-	-	-	1	1
Female	62	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Kentucky												
All races, both sexes	220	-	-	-	-	-	1	-	-	-	3	3
Male	104	-	-	-	-	-	-	-	-	-	2	1
Female	116	-	-	-	-	-	1	-	-	-	1	2
White, both sexes	213	-	-	-	-	-	1	-	-	-	3	3
Male	100	-	-	-	-	-	-	-	-	-	2	1
Female	113	-	-	-	-	-	1	-	-	-	1	2
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	178	-	-	-	-	-	-	-	-	-	2	3
Male	109	-	-	-	-	-	-	-	-	-	2	2
Female	69	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	155	-	-	-	-	-	-	-	-	-	2	2
Male	91	-	-	-	-	-	-	-	-	-	2	1
Female	64	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Maine												
All races, both sexes	73	-	-	-	-	-	-	-	-	-	1	1
Male	48	-	-	-	-	-	-	-	-	-	1	1
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	-	1	1
Male	48	-	-	-	-	-	-	-	-	-	1	1
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	209	-	-	-	-	-	-	-	1	-	2	2
Male	97	-	-	-	-	-	-	-	1	-	1	1
Female	112	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	176	-	-	-	-	-	-	-	1	-	-	2
Male	82	-	-	-	-	-	-	-	1	-	-	1
Female	94	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	30	-	-	-	-	-	-	-	-	-	2	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Massachusetts												
All races, both sexes	265	-	-	-	-	-	-	-	-	-	-	1
Male	118	-	-	-	-	-	-	-	-	-	-	1
Female	147	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	258	-	-	-	-	-	-	-	-	-	-	1
Male	115	-	-	-	-	-	-	-	-	-	-	1
Female	143	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	536	-	-	-	-	-	-	-	-	1	3	4
Male	249	-	-	-	-	-	-	-	-	1	2	3
Female	287	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	500	-	-	-	-	-	-	-	-	-	2	3
Male	230	-	-	-	-	-	-	-	-	-	1	2
Female	270	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	30	-	-	-	-	-	-	-	-	1	1	-
Male	16	-	-	-	-	-	-	-	-	1	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Minnesota												
All races, both sexes	108	-	-	-	-	-	-	-	-	-	1	-
Male	57	-	-	-	-	-	-	-	-	-	1	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	105	-	-	-	-	-	-	-	-	-	-	-
Male	55	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	88	-	-	-	-	-	-	1	-	-	3	2
Male	49	-	-	-	-	-	-	1	-	-	1	1
Female	39	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	79	-	-	-	-	-	-	1	-	-	3	2
Male	46	-	-	-	-	-	-	1	-	-	1	1
Female	33	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Missouri												
All races, both sexes	270	-	-	-	1	-	-	-	1	-	3	4
Male	142	-	-	-	-	-	-	-	1	-	2	3
Female	128	-	-	-	1	-	-	-	-	-	1	1
White, both sexes	253	-	-	-	-	-	-	-	1	-	2	3
Male	131	-	-	-	-	-	-	-	1	-	1	2
Female	122	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	16	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Montana												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Nebraska												
All races, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	34	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	62	-	-	-	-	-	-	-	-	-	-	-
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	62	-	-	-	-	-	-	-	1	-	1	1
Male	28	-	-	-	-	-	-	-	-	-	1	-
Female	34	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	61	-	-	-	-	-	-	-	1	-	1	1
Male	27	-	-	-	-	-	-	-	-	-	1	-
Female	34	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
New Hampshire												
All races, both sexes	72	-	-	-	-	-	-	-	-	-	1	1
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	72	-	-	-	-	-	-	-	-	-	1	1
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	411	-	-	-	-	-	-	-	-	-	-	3
Male	185	-	-	-	-	-	-	-	-	-	-	1
Female	226	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	385	-	-	-	-	-	-	-	-	-	-	3
Male	173	-	-	-	-	-	-	-	-	-	-	1
Female	212	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
New Mexico												
All races, both sexes	100	-	-	-	-	-	-	-	-	1	2	1
Male	52	-	-	-	-	-	-	-	-	1	-	-
Female	48	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	99	-	-	-	-	-	-	-	-	1	2	1
Male	51	-	-	-	-	-	-	-	-	1	-	-
Female	48	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	601	-	-	-	-	-	-	-	-	1	5	9
Male	287	-	-	-	-	-	-	-	-	1	4	7
Female	314	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	553	-	-	-	-	-	-	-	-	1	5	7
Male	265	-	-	-	-	-	-	-	-	1	4	5
Female	288	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	38	-	-	-	-	-	-	-	-	-	-	2
Male	16	-	-	-	-	-	-	-	-	-	-	2
Female	22	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
North Carolina												
All races, both sexes	449	-	-	-	-	-	-	-	-	-	2	10
Male	247	-	-	-	-	-	-	-	-	-	-	6
Female	202	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	408	-	-	-	-	-	-	-	-	-	-	10
Male	218	-	-	-	-	-	-	-	-	-	-	6
Female	190	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	39	-	-	-	-	-	-	-	-	-	2	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Ohio												
All races, both sexes	723	-	-	-	-	-	-	-	1	-	3	8
Male	350	-	-	-	-	-	-	-	1	-	2	5
Female	373	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	669	-	-	-	-	-	-	-	1	-	3	7
Male	318	-	-	-	-	-	-	-	1	-	2	4
Female	351	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	54	-	-	-	-	-	-	-	-	-	-	1
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	198	-	-	-	-	-	-	-	-	1	1	2
Male	114	-	-	-	-	-	-	-	-	1	-	1
Female	84	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	189	-	-	-	-	-	-	-	-	1	1	1
Male	110	-	-	-	-	-	-	-	-	1	-	1
Female	79	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Oregon												
All races, both sexes	214	-	-	-	-	-	-	-	-	-	-	4
Male	103	-	-	-	-	-	-	-	-	-	-	1
Female	111	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	211	-	-	-	-	-	-	-	-	-	-	3
Male	102	-	-	-	-	-	-	-	-	-	-	1
Female	109	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	560	-	-	-	-	-	-	-	-	-	2	3
Male	272	-	-	-	-	-	-	-	-	-	2	1
Female	288	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	522	-	-	-	-	-	-	-	-	-	1	2
Male	249	-	-	-	-	-	-	-	-	-	1	1
Female	273	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	36	-	-	-	-	-	-	-	-	-	1	1
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Rhode Island												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	180	-	-	-	-	-	-	-	-	-	3	4
Male	103	-	-	-	-	-	-	-	-	-	3	2
Female	77	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	159	-	-	-	-	-	-	-	-	-	3	2
Male	88	-	-	-	-	-	-	-	-	-	3	1
Female	71	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	21	-	-	-	-	-	-	-	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
South Dakota												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	289	-	-	-	-	-	-	-	-	-	2	8
Male	143	-	-	-	-	-	-	-	-	-	-	4
Female	146	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	270	-	-	-	-	-	-	-	-	-	2	7
Male	131	-	-	-	-	-	-	-	-	-	-	3
Female	139	-	-	-	-	-	-	-	-	-	2	4
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Texas												
All races, both sexes	824	-	-	-	-	-	-	-	1	-	5	13
Male	440	-	-	-	-	-	-	-	1	-	3	4
Female	384	-	-	-	-	-	-	-	-	-	2	9
White, both sexes	754	-	-	-	-	-	-	-	-	-	3	12
Male	397	-	-	-	-	-	-	-	-	-	2	4
Female	357	-	-	-	-	-	-	-	-	-	1	8
Black, both sexes	63	-	-	-	-	-	-	-	1	-	2	1
Male	38	-	-	-	-	-	-	-	1	-	1	-
Female	25	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	51	1	-	-	-	-	-	-	-	-	1	2
Male	31	1	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	51	1	-	-	-	-	-	-	-	-	1	2
Male	31	1	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Vermont												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	291	-	-	-	-	-	-	-	-	1	3	3
Male	137	-	-	-	-	-	-	-	-	-	2	1
Female	154	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	266	-	-	-	-	-	-	-	-	1	1	3
Male	121	-	-	-	-	-	-	-	-	-	1	1
Female	145	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	21	-	-	-	-	-	-	-	-	-	2	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Washington												
All races, both sexes	207	1	-	-	-	-	-	-	-	-	-	2
Male	110	1	-	-	-	-	-	-	-	-	-	1
Female	97	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	201	-	-	-	-	-	-	-	-	-	-	2
Male	104	-	-	-	-	-	-	-	-	-	-	1
Female	97	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	103	-	-	-	-	-	-	-	1	-	1	2
Male	49	-	-	-	-	-	-	-	-	-	1	2
Female	54	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	103	-	-	-	-	-	-	-	1	-	1	2
Male	49	-	-	-	-	-	-	-	-	-	1	2
Female	54	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
Wisconsin												
All races, both sexes	217	-	-	-	-	-	-	-	-	-	2	-
Male	118	-	-	-	-	-	-	-	-	-	2	-
Female	99	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	212	-	-	-	-	-	-	-	-	-	2	-
Male	114	-	-	-	-	-	-	-	-	-	2	-
Female	98	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
United States												
All races, both sexes	296	575	948	1,368	1,905	2,444	2,355	1,547	679	149	25	2
Male	187	308	503	730	967	1,258	1,150	739	288	38	10	2
Female	109	267	445	638	938	1,186	1,205	808	391	111	15	-
White, both sexes	262	521	858	1,258	1,791	2,290	2,221	1,474	643	136	23	2
Male	157	271	438	660	907	1,160	1,070	702	268	32	9	2
Female	105	250	420	598	884	1,130	1,151	772	375	104	14	-
Black, both sexes	31	50	86	89	92	122	103	58	24	8	2	-
Male	27	34	62	57	47	72	60	28	13	3	1	-
Female	4	16	24	32	45	50	43	30	11	5	1	-
American Indian 1/, both sexes	1	1	2	5	8	5	5	1	1	1	-	-
Male	1	1	1	3	2	3	3	-	-	-	-	-
Female	-	-	1	2	6	2	2	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	2	3	2	16	14	27	26	14	11	4	-	-
Male	2	2	2	10	11	23	17	9	7	3	-	-
Female	-	1	-	6	3	4	9	5	4	1	-	-
Alabama												
All races, both sexes	5	15	19	20	28	43	27	14	6	4	-	-
Male	2	8	12	8	14	25	16	8	4	2	-	-
Female	3	7	7	12	14	18	11	6	2	2	-	-
White, both sexes	5	13	17	16	26	40	24	11	6	2	-	-
Male	2	6	10	5	13	23	13	6	4	1	-	-
Female	3	7	7	11	13	17	11	5	2	1	-	-
Black, both sexes	-	2	2	4	2	3	3	3	-	1	-	-
Male	-	2	2	3	1	2	3	2	-	1	-	-
Female	-	-	-	1	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Alaska												
All races, both sexes	1	3	3	3	4	2	3	2	-	-	-	-
Male	-	2	3	1	3	-	1	1	-	-	-	-
Female	1	1	-	2	1	2	2	1	-	-	-	-
White, both sexes	1	2	3	3	4	2	2	2	-	-	-	-
Male	-	1	3	1	3	-	-	1	-	-	-	-
Female	1	1	-	2	1	2	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	6	14	14	27	51	85	74	41	28	6	1	-
Male	3	7	7	12	25	48	36	21	10	1	1	-
Female	3	7	7	15	26	37	38	20	18	5	-	-
White, both sexes	6	13	14	26	50	84	74	41	28	6	1	-
Male	3	6	7	11	25	47	36	21	10	1	1	-
Female	3	7	7	15	25	37	38	20	18	5	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Arkansas												
All races, both sexes	3	9	16	25	27	34	35	20	6	1	-	-
Male	2	6	10	12	20	22	18	8	3	1	-	-
Female	1	3	6	13	7	12	17	12	3	-	-	-
White, both sexes	2	9	15	25	27	32	32	19	6	1	-	-
Male	1	6	10	12	20	21	16	8	3	1	-	-
Female	1	3	5	13	7	11	16	11	3	-	-	-
Black, both sexes	1	-	1	-	-	2	3	1	-	-	-	-
Male	1	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	1	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	26	59	77	115	156	242	232	187	81	19	1	-
Male	16	37	45	58	74	120	109	82	34	6	-	-
Female	10	22	32	57	82	122	123	105	47	13	1	-
White, both sexes	23	53	63	103	147	220	209	177	75	16	1	-
Male	13	33	32	51	68	107	90	75	31	5	-	-
Female	10	20	31	52	79	113	119	102	44	11	1	-
Black, both sexes	2	6	11	6	5	13	12	4	4	1	-	-
Male	2	4	10	4	2	5	9	3	2	-	-	-
Female	-	2	1	2	3	8	3	1	2	1	-	-
American Indian 1/, both sexes	-	-	1	1	1	1	-	-	-	-	-	-
Male	-	-	1	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	5	3	8	11	6	2	2	-	-
Male	1	-	2	2	3	7	10	4	1	1	-	-
Female	-	-	-	3	-	1	1	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Colorado												
All races, both sexes	2	5	16	21	19	23	29	28	4	4	1	-
Male	1	3	8	16	8	10	20	11	1	1	-	-
Female	1	2	8	5	11	13	9	17	3	3	1	-
White, both sexes	2	5	14	20	17	22	27	28	4	4	1	-
Male	1	3	6	16	8	10	20	11	1	1	-	-
Female	1	2	8	4	9	12	7	17	3	3	1	-
Black, both sexes	-	-	2	1	2	1	1	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	2	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	1	5	4	19	14	35	30	21	19	6	-	-
Male	1	1	2	14	6	22	14	14	9	-	-	-
Female	-	4	2	5	8	13	16	7	10	6	-	-
White, both sexes	1	4	3	17	14	32	28	21	17	6	-	-
Male	1	-	2	12	6	21	13	14	8	-	-	-
Female	-	4	1	5	8	11	15	7	9	6	-	-
Black, both sexes	-	1	1	2	-	3	2	-	1	-	-	-
Male	-	1	-	2	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	2	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Delaware												
All races, both sexes	2	-	-	4	12	7	9	-	1	-	-	-
Male	1	-	-	3	6	4	3	-	1	-	-	-
Female	1	-	-	1	6	3	6	-	-	-	-	-
White, both sexes	2	-	-	1	10	7	8	-	1	-	-	-
Male	1	-	-	1	6	4	3	-	1	-	-	-
Female	1	-	-	-	4	3	5	-	-	-	-	-
Black, both sexes	-	-	-	3	2	-	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	1	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	1	-	2	-	2	2	-	-	-
Male	-	-	-	1	-	1	-	-	2	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	1	-	1	-	-	2	-	-	-
Male	-	-	-	1	-	-	-	-	2	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Florida												
All races, both sexes	20	31	50	79	123	169	182	110	56	11	3	-
Male	13	18	24	41	62	80	73	58	23	1	2	-
Female	7	13	26	38	61	89	109	52	33	10	1	-
White, both sexes	18	27	48	75	117	161	174	108	54	9	3	-
Male	11	14	22	39	58	73	70	56	21	-	2	-
Female	7	13	26	36	59	88	104	52	33	9	1	-
Black, both sexes	2	4	2	4	6	7	8	2	2	2	-	-
Male	2	4	2	2	4	6	3	2	2	1	-	-
Female	-	-	-	2	2	1	5	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	11	21	36	47	69	67	54	34	10	3	-	-
Male	7	10	25	28	38	36	26	24	-	-	-	-
Female	4	11	11	19	31	31	28	10	10	3	-	-
White, both sexes	9	16	33	46	66	63	48	30	10	2	-	-
Male	5	6	22	27	35	32	23	20	-	-	-	-
Female	4	10	11	19	31	31	25	10	10	2	-	-
Black, both sexes	2	5	3	1	3	4	6	3	-	1	-	-
Male	2	4	3	1	3	4	3	3	-	-	-	-
Female	-	1	-	-	-	-	3	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Hawaii												
All races, both sexes	-	2	-	5	5	12	12	5	4	1	1	-
Male	-	2	-	3	3	7	7	3	2	-	-	-
Female	-	-	-	2	2	5	5	2	2	1	1	-
White, both sexes	-	-	-	-	3	5	5	1	1	1	1	-
Male	-	-	-	-	1	2	3	-	1	-	-	-
Female	-	-	-	-	2	3	2	1	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	5	2	7	7	4	3	-	-	-
Male	-	2	-	3	2	5	4	3	1	-	-	-
Female	-	-	-	2	-	2	3	1	2	-	-	-
Idaho												
All races, both sexes	2	2	5	9	17	17	13	11	6	-	-	-
Male	1	1	3	8	6	6	9	6	3	-	-	-
Female	1	1	2	1	11	11	4	5	3	-	-	-
White, both sexes	2	2	5	9	17	17	13	11	6	-	-	-
Male	1	1	3	8	6	6	9	6	3	-	-	-
Female	1	1	2	1	11	11	4	5	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Illinois												
All races, both sexes	17	17	35	51	86	107	122	72	41	6	-	-
Male	10	11	16	27	47	62	53	34	13	1	-	-
Female	7	6	19	24	39	45	69	38	28	5	-	-
White, both sexes	14	14	30	49	78	97	118	66	37	6	-	-
Male	8	9	13	25	41	55	51	31	10	1	-	-
Female	6	5	17	24	37	42	67	35	27	5	-	-
Black, both sexes	3	3	5	2	8	9	4	5	2	-	-	-
Male	2	2	3	2	6	6	2	2	2	-	-	-
Female	1	1	2	-	2	3	2	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	2	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Indiana												
All races, both sexes	7	18	28	46	65	64	61	41	15	6	-	-
Male	4	8	17	29	33	29	31	20	9	2	-	-
Female	3	10	11	17	32	35	30	21	6	4	-	-
White, both sexes	7	16	27	45	63	62	59	40	14	6	-	-
Male	4	8	16	28	32	29	30	20	9	2	-	-
Female	3	8	11	17	31	33	29	20	5	4	-	-
Black, both sexes	-	2	1	1	2	2	2	1	1	-	-	-
Male	-	-	1	1	1	-	1	-	-	-	-	-
Female	-	2	-	-	1	2	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Iowa												
All races, both sexes	1	4	2	10	19	13	31	16	6	-	2	-
Male	-	4	1	4	12	10	19	8	1	-	1	-
Female	1	-	1	6	7	3	12	8	5	-	1	-
White, both sexes	1	4	2	10	18	13	31	16	6	-	2	-
Male	-	4	1	4	12	10	19	8	1	-	1	-
Female	1	-	1	6	6	3	12	8	5	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	3	11	15	15	26	18	21	12	2	1	-
Male	2	1	5	10	5	15	7	8	6	1	-	-
Female	-	2	6	5	10	11	11	13	6	1	1	-
White, both sexes	2	3	11	15	14	24	15	20	12	2	1	-
Male	2	1	5	10	5	14	6	8	6	1	-	-
Female	-	2	6	5	9	10	9	12	6	1	1	-
Black, both sexes	-	-	-	-	1	2	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Kentucky												
All races, both sexes	8	9	32	28	33	40	21	28	9	4	1	-
Male	6	5	16	16	19	15	6	14	3	1	-	-
Female	2	4	16	12	14	25	15	14	6	3	1	-
White, both sexes	8	8	31	26	33	39	21	27	8	4	1	-
Male	6	4	15	15	19	15	6	14	2	1	-	-
Female	2	4	16	11	14	24	15	13	6	3	1	-
Black, both sexes	-	1	1	2	-	1	-	1	1	-	-	-
Male	-	1	1	1	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	10	10	17	28	27	21	36	16	7	1	-	-
Male	9	3	8	14	21	14	24	8	4	-	-	-
Female	1	7	9	14	6	7	12	8	3	1	-	-
White, both sexes	5	9	13	25	24	20	34	13	7	1	-	-
Male	4	3	5	12	18	13	22	7	4	-	-	-
Female	1	6	8	13	6	7	12	6	3	1	-	-
Black, both sexes	5	1	4	3	3	1	2	3	-	-	-	-
Male	5	-	3	2	3	1	2	1	-	-	-	-
Female	-	1	1	1	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Maine												
All races, both sexes	4	3	5	13	6	13	15	11	1	-	-	-
Male	2	2	3	7	5	8	12	7	-	-	-	-
Female	2	1	2	6	1	5	3	4	1	-	-	-
White, both sexes	4	3	5	13	6	13	15	11	1	-	-	-
Male	2	2	3	7	5	8	12	7	-	-	-	-
Female	2	1	2	6	1	5	3	4	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	7	16	23	36	35	44	27	8	2	2	-
Male	1	3	8	8	16	16	23	13	5	-	1	-
Female	3	4	8	15	20	19	21	14	3	2	1	-
White, both sexes	4	5	12	18	34	30	37	22	7	2	2	-
Male	1	2	6	5	14	15	19	13	4	-	1	-
Female	3	3	6	13	20	15	18	9	3	2	1	-
Black, both sexes	-	2	4	5	2	5	5	5	-	-	-	-
Male	-	1	2	3	2	1	3	-	-	-	-	-
Female	-	1	2	2	-	4	2	5	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Massachusetts												
All races, both sexes	2	13	23	23	38	58	52	32	16	6	1	-
Male	1	6	12	12	19	26	20	14	6	1	-	-
Female	1	7	11	11	19	32	32	18	10	5	1	-
White, both sexes	1	13	23	22	37	57	51	31	16	5	1	-
Male	-	6	12	12	19	25	19	14	6	1	-	-
Female	1	7	11	10	18	32	32	17	10	4	1	-
Black, both sexes	1	-	-	1	1	1	1	1	-	1	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	13	29	37	53	86	113	108	62	22	4	1	-
Male	10	16	14	26	34	55	47	31	8	1	1	-
Female	3	13	23	27	52	58	61	31	14	3	-	-
White, both sexes	12	24	34	48	82	106	103	59	22	4	1	-
Male	9	14	12	24	32	51	45	30	8	1	1	-
Female	3	10	22	24	50	55	58	29	14	3	-	-
Black, both sexes	1	5	3	3	2	6	5	3	-	-	-	-
Male	1	2	2	1	1	4	2	1	-	-	-	-
Female	-	3	1	2	1	2	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	1	1	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Minnesota												
All races, both sexes	3	3	6	11	18	26	20	11	7	2	-	-
Male	2	2	3	6	8	18	11	4	2	-	-	-
Female	1	1	3	5	10	8	9	7	5	2	-	-
White, both sexes	2	3	6	11	18	26	19	11	7	2	-	-
Male	1	2	3	6	8	18	11	4	2	-	-	-
Female	1	1	3	5	10	8	8	7	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	3	11	9	4	20	16	12	3	1	1	1
Male	-	-	8	4	2	15	9	7	-	-	-	1
Female	1	3	3	5	2	5	7	5	3	1	1	-
White, both sexes	1	3	8	8	4	19	15	11	2	1	-	1
Male	-	-	6	4	2	14	9	7	-	-	-	1
Female	1	3	2	4	2	5	6	4	2	1	-	-
Black, both sexes	-	-	3	-	-	1	1	1	1	-	1	-
Male	-	-	2	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Missouri												
All races, both sexes	7	12	21	35	35	51	56	33	9	2	-	-
Male	3	7	10	16	23	23	30	15	7	2	-	-
Female	4	5	11	19	12	28	26	18	2	-	-	-
White, both sexes	6	10	17	35	35	51	52	31	9	1	-	-
Male	2	6	9	16	23	23	27	13	7	1	-	-
Female	4	4	8	19	12	28	25	18	2	-	-	-
Black, both sexes	1	2	4	-	-	-	4	2	-	1	-	-
Male	1	1	1	-	-	-	3	2	-	1	-	-
Female	-	1	3	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	6	3	9	10	10	6	2	-	-	-
Male	1	-	1	1	5	7	2	3	1	-	-	-
Female	1	-	5	2	4	3	8	3	1	-	-	-
White, both sexes	2	-	6	3	9	10	10	6	2	-	-	-
Male	1	-	1	1	5	7	2	3	1	-	-	-
Female	1	-	5	2	4	3	8	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Nebraska												
All races, both sexes	-	2	3	10	10	11	14	9	4	1	1	-
Male	-	1	3	6	6	5	5	4	3	-	1	-
Female	-	1	-	4	4	6	9	5	1	1	-	-
White, both sexes	-	2	3	10	10	10	13	9	4	1	-	-
Male	-	1	3	6	6	4	4	4	3	-	-	-
Female	-	1	-	4	4	6	9	5	1	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	3	10	7	19	5	10	1	3	1	-	-
Male	-	1	2	2	10	3	5	1	3	-	-	-
Female	-	2	8	5	9	2	5	-	-	1	-	-
White, both sexes	-	3	10	7	19	4	10	1	3	1	-	-
Male	-	1	2	2	10	2	5	1	3	-	-	-
Female	-	2	8	5	9	2	5	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
New Hampshire												
All races, both sexes	2	3	4	10	6	18	10	10	7	-	-	-
Male	-	2	2	6	2	11	5	1	3	-	-	-
Female	2	1	2	4	4	7	5	9	4	-	-	-
White, both sexes	2	3	4	10	6	18	10	10	7	-	-	-
Male	-	2	2	6	2	11	5	1	3	-	-	-
Female	2	1	2	4	4	7	5	9	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	9	13	18	36	65	87	72	68	33	6	1	-
Male	6	6	8	21	29	44	30	26	12	2	-	-
Female	3	7	10	15	36	43	42	42	21	4	1	-
White, both sexes	9	13	16	32	59	81	71	63	31	6	1	-
Male	6	6	6	19	26	41	29	26	11	2	-	-
Female	3	7	10	13	33	40	42	37	20	4	1	-
Black, both sexes	-	-	2	3	6	5	1	4	2	-	-	-
Male	-	-	2	1	3	2	1	-	1	-	-	-
Female	-	-	-	2	3	3	-	4	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
New Mexico												
All races, both sexes	1	3	11	10	15	19	21	8	4	3	1	-
Male	1	2	3	7	7	11	11	4	3	1	1	-
Female	-	1	8	3	8	8	10	4	1	2	-	-
White, both sexes	1	3	11	10	15	18	21	8	4	3	1	-
Male	1	2	3	7	7	10	11	4	3	1	1	-
Female	-	1	8	3	8	8	10	4	1	2	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	17	13	45	73	71	112	116	87	39	13	-	-
Male	12	9	24	33	32	51	61	39	10	4	-	-
Female	5	4	21	40	39	61	55	48	29	9	-	-
White, both sexes	13	12	39	64	62	106	109	86	37	12	-	-
Male	10	9	20	30	28	49	57	39	10	3	-	-
Female	3	3	19	34	34	57	52	47	27	9	-	-
Black, both sexes	4	-	6	7	6	6	5	-	2	-	-	-
Male	2	-	4	2	1	2	3	-	-	-	-	-
Female	2	-	2	5	5	4	2	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	3	-	2	1	-	1	-	-
Male	-	-	-	1	3	-	1	-	-	1	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
North Carolina												
All races, both sexes	10	28	36	54	79	77	86	49	18	-	-	-
Male	7	13	22	35	40	44	47	25	8	-	-	-
Female	3	15	14	19	39	33	39	24	10	-	-	-
White, both sexes	9	24	30	48	72	72	81	44	18	-	-	-
Male	6	9	18	29	35	41	44	22	8	-	-	-
Female	3	15	12	19	37	31	37	22	10	-	-	-
Black, both sexes	1	4	6	5	6	5	5	5	-	-	-	-
Male	1	4	4	5	5	3	3	3	-	-	-	-
Female	-	-	2	-	1	2	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	-	1	2	6	4	1	-	-	-	-
Male	-	1	-	1	-	2	3	-	-	-	-	-
Female	-	-	-	-	2	4	1	1	-	-	-	-
White, both sexes	-	1	-	1	1	6	4	1	-	-	-	-
Male	-	1	-	1	-	2	3	-	-	-	-	-
Female	-	-	-	-	1	4	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Ohio												
All races, both sexes	24	39	49	65	124	141	132	84	42	9	2	-
Male	17	23	27	34	59	64	63	38	16	1	-	-
Female	7	16	22	31	65	77	69	46	26	8	2	-
White, both sexes	21	34	43	57	117	127	126	83	39	9	2	-
Male	14	21	22	28	57	55	60	38	15	1	-	-
Female	7	13	21	29	60	72	66	45	24	8	2	-
Black, both sexes	3	5	6	8	7	14	6	1	3	-	-	-
Male	3	2	5	6	2	9	3	-	1	-	-	-
Female	-	3	1	2	5	5	3	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	9	9	21	24	29	43	34	14	8	3	-	-
Male	4	5	11	13	19	29	18	8	5	-	-	-
Female	5	4	10	11	10	14	16	6	3	3	-	-
White, both sexes	9	9	20	22	27	41	34	14	7	3	-	-
Male	4	5	11	11	19	27	18	8	5	-	-	-
Female	5	4	9	11	8	14	16	6	2	3	-	-
Black, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	2	2	-	-	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	1	-	2	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Oregon												
All races, both sexes	2	9	17	19	32	54	44	23	9	-	1	-
Male	2	5	8	8	16	28	21	11	2	-	1	-
Female	-	4	9	11	16	26	23	12	7	-	-	-
White, both sexes	2	9	17	19	32	52	44	23	9	-	1	-
Male	2	5	8	8	16	27	21	11	2	-	1	-
Female	-	4	9	11	16	25	23	12	7	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	8	16	42	62	93	111	106	82	29	5	1	-
Male	4	7	22	37	46	46	49	41	14	2	1	-
Female	4	9	20	25	47	65	57	41	15	3	-	-
White, both sexes	7	14	39	53	87	106	99	81	28	4	1	-
Male	3	5	21	31	43	42	45	40	14	2	1	-
Female	4	9	18	22	44	64	54	41	14	2	-	-
Black, both sexes	1	2	3	8	6	5	6	1	1	1	-	-
Male	1	2	1	5	3	4	3	1	-	-	-	-
Female	-	-	2	3	3	1	3	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Rhode Island												
All races, both sexes	-	4	4	1	6	3	8	2	-	-	-	-
Male	-	2	1	1	3	2	3	-	-	-	-	-
Female	-	2	3	-	3	1	5	2	-	-	-	-
White, both sexes	-	4	4	1	6	3	8	2	-	-	-	-
Male	-	2	1	1	3	2	3	-	-	-	-	-
Female	-	2	3	-	3	1	5	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	6	14	14	20	31	26	33	22	5	2	-	-
Male	4	9	9	11	15	17	20	8	4	1	-	-
Female	2	5	5	9	16	9	13	14	1	1	-	-
White, both sexes	6	13	11	18	24	24	31	20	5	2	-	-
Male	4	8	7	10	10	15	19	6	4	1	-	-
Female	2	5	4	8	14	9	12	14	1	1	-	-
Black, both sexes	-	1	3	2	7	2	2	2	-	-	-	-
Male	-	1	2	1	5	2	1	2	-	-	-	-
Female	-	-	1	1	2	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
South Dakota												
All races, both sexes	1	2	2	4	2	7	5	7	4	-	-	-
Male	-	-	2	1	1	3	3	3	-	-	-	-
Female	1	2	-	3	1	4	2	4	4	-	-	-
White, both sexes	1	2	2	4	2	7	5	6	4	-	-	-
Male	-	-	2	1	1	3	3	3	-	-	-	-
Female	1	2	-	3	1	4	2	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	17	27	43	51	53	47	27	5	3	-	-
Male	4	8	12	22	28	28	22	12	1	2	-	-
Female	2	9	15	21	23	25	25	15	4	1	-	-
White, both sexes	6	17	22	42	47	50	43	26	5	3	-	-
Male	4	8	8	22	26	26	19	12	1	2	-	-
Female	2	9	14	20	21	24	24	14	4	1	-	-
Black, both sexes	-	-	5	1	3	3	4	1	-	-	-	-
Male	-	-	4	-	2	2	3	-	-	-	-	-
Female	-	-	1	1	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Texas												
All races, both sexes	20	57	82	107	119	152	146	82	33	6	1	-
Male	16	28	51	61	62	73	80	41	19	1	-	-
Female	4	29	31	46	57	79	66	41	14	5	1	-
White, both sexes	16	55	77	94	107	140	136	76	31	6	1	-
Male	12	27	47	51	58	67	75	36	17	1	-	-
Female	4	28	30	43	49	73	61	40	14	5	1	-
Black, both sexes	3	2	5	13	10	10	9	6	1	-	-	-
Male	3	1	4	10	3	5	4	5	1	-	-	-
Female	-	1	1	3	7	5	5	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	2	2	-	-	1	-	-	-
Male	1	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Utah												
All races, both sexes	2	5	3	5	14	8	8	2	-	-	-	-
Male	2	1	2	3	10	5	6	-	-	-	-	-
Female	-	4	1	2	4	3	2	2	-	-	-	-
White, both sexes	2	5	3	5	14	8	8	2	-	-	-	-
Male	2	1	2	3	10	5	6	-	-	-	-	-
Female	-	4	1	2	4	3	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Vermont												
All races, both sexes	1	1	3	1	4	10	8	4	1	1	-	-
Male	1	1	2	1	3	5	5	2	-	-	-	-
Female	-	-	1	-	1	5	3	2	1	1	-	-
White, both sexes	1	1	3	1	4	10	8	4	1	1	-	-
Male	1	1	2	1	3	5	5	2	-	-	-	-
Female	-	-	1	-	1	5	3	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5	12	24	37	48	59	50	31	15	2	1	-
Male	3	6	10	17	24	29	23	13	8	1	-	-
Female	2	6	14	20	24	30	27	18	7	1	1	-
White, both sexes	4	12	22	36	44	51	46	29	14	2	1	-
Male	3	6	8	17	22	22	21	12	7	1	-	-
Female	1	6	14	19	22	29	25	17	7	1	1	-
Black, both sexes	1	-	2	1	2	8	3	2	-	-	-	-
Male	-	-	2	-	-	7	2	1	-	-	-	-
Female	1	-	-	1	2	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	1	-	1	-	-	-
Male	-	-	-	-	2	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Washington												
All races, both sexes	6	6	19	21	30	36	35	34	14	2	-	1
Male	4	1	8	11	16	21	16	20	8	2	-	1
Female	2	5	11	10	14	15	19	14	6	-	-	-
White, both sexes	6	6	19	20	30	34	34	34	14	1	-	1
Male	4	1	8	10	16	19	15	20	8	1	-	1
Female	2	5	11	10	14	15	19	14	6	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	-	-	-	1	-	-
Male	-	-	-	1	-	2	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	3	7	11	9	19	23	13	8	6	-	-	-
Male	-	4	5	3	8	13	6	4	3	-	-	-
Female	3	3	6	6	11	10	7	4	3	-	-	-
White, both sexes	3	7	11	9	19	23	13	8	6	-	-	-
Male	-	4	5	3	8	13	6	4	3	-	-	-
Female	3	3	6	6	11	10	7	4	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Wisconsin												
All races, both sexes	4	13	11	25	29	45	40	29	18	1	-	-
Male	2	9	6	11	16	28	19	16	9	-	-	-
Female	2	4	5	14	13	17	21	13	9	1	-	-
White, both sexes	4	12	10	25	29	43	40	29	17	1	-	-
Male	2	8	6	11	16	26	19	16	8	-	-	-
Female	2	4	4	14	13	17	21	13	9	1	-	-
Black, both sexes	-	1	1	-	-	1	-	-	1	-	-	-
Male	-	1	-	-	-	1	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	2	1	5	3	3	-	1	-	-	-
Male	-	1	2	-	1	2	3	-	-	-	-	-
Female	-	-	-	1	4	1	-	-	1	-	-	-
White, both sexes	-	1	2	1	5	3	3	-	1	-	-	-
Male	-	1	2	-	1	2	3	-	-	-	-	-
Female	-	-	-	1	4	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
United States												
All races, both sexes	106,706	3	5	5	7	1	2	15	29	64	301	864
Male	51,063	3	3	2	3	-	2	7	14	32	148	429
Female	55,643	-	2	3	4	1	-	8	15	32	153	435
White, both sexes	99,259	1	4	3	4	1	1	9	20	55	238	759
Male	46,951	1	3	2	2	-	1	4	9	27	119	371
Female	52,308	-	1	1	2	1	-	5	11	28	119	388
Black, both sexes	5,965	-	-	2	3	-	1	6	9	9	53	92
Male	3,230	-	-	-	1	-	1	3	5	5	21	50
Female	2,735	-	-	2	2	-	-	3	4	4	32	42
American Indian 1/, both sexes	449	2	-	-	-	-	-	-	-	-	7	10
Male	208	2	-	-	-	-	-	-	-	-	6	5
Female	241	-	-	-	-	-	-	-	-	-	1	5
Asian or Pacific Islander 2/, both sexes	1,033	-	1	-	-	-	-	-	-	-	3	3
Male	674	-	-	-	-	-	-	-	-	-	2	3
Female	359	-	1	-	-	-	-	-	-	-	1	-
Alabama												
All races, both sexes	2,037	-	-	-	-	-	-	-	1	-	5	22
Male	1,018	-	-	-	-	-	-	-	-	-	2	14
Female	1,019	-	-	-	-	-	-	-	1	-	3	8
White, both sexes	1,842	-	-	-	-	-	-	-	-	-	4	18
Male	893	-	-	-	-	-	-	-	-	-	2	13
Female	949	-	-	-	-	-	-	-	-	-	2	5
Black, both sexes	193	-	-	-	-	-	-	-	1	-	1	4
Male	124	-	-	-	-	-	-	-	-	-	-	1
Female	69	-	-	-	-	-	-	-	1	-	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Alaska												
All races, both sexes	108	-	-	-	-	-	-	-	-	-	1	-
Male	46	-	-	-	-	-	-	-	-	-	-	-
Female	62	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	73	-	-	-	-	-	-	-	-	-	1	-
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2,337	1	-	-	-	-	-	-	-	-	1	13
Male	1,136	1	-	-	-	-	-	-	-	-	1	5
Female	1,201	-	-	-	-	-	-	-	-	-	-	8
White, both sexes	2,295	-	-	-	-	-	-	-	-	-	1	13
Male	1,116	-	-	-	-	-	-	-	-	-	1	5
Female	1,179	-	-	-	-	-	-	-	-	-	-	8
Black, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Arkansas												
All races, both sexes	1,246	-	-	-	-	-	-	-	1	1	4	4
Male	657	-	-	-	-	-	-	-	-	1	4	1
Female	589	-	-	-	-	-	-	-	1	-	-	3
White, both sexes	1,184	-	-	-	-	-	-	-	1	1	4	3
Male	622	-	-	-	-	-	-	-	-	1	4	1
Female	562	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	58	-	-	-	-	-	-	-	-	-	-	1
Male	35	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	10,945	1	2	3	-	-	2	3	3	4	21	65
Male	5,126	1	1	1	-	-	2	2	1	1	11	34
Female	5,819	-	1	2	-	-	-	1	2	3	10	31
White, both sexes	9,777	1	2	1	-	-	1	1	3	2	17	58
Male	4,426	1	1	1	-	-	1	-	1	-	10	30
Female	5,351	-	1	-	-	-	-	1	2	2	7	28
Black, both sexes	549	-	-	2	-	-	1	2	-	2	4	5
Male	299	-	-	-	-	-	1	2	-	1	1	2
Female	250	-	-	2	-	-	-	-	-	1	3	3
American Indian 1/, both sexes	52	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	567	-	-	-	-	-	-	-	-	-	-	2
Male	376	-	-	-	-	-	-	-	-	-	-	2
Female	191	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Colorado												
All races, both sexes	1,712	-	-	-	-	-	-	1	-	-	3	10
Male	871	-	-	-	-	-	-	-	-	-	2	3
Female	841	-	-	-	-	-	-	1	-	-	1	7
White, both sexes	1,657	-	-	-	-	-	-	1	-	-	2	10
Male	845	-	-	-	-	-	-	-	-	-	1	3
Female	812	-	-	-	-	-	-	1	-	-	1	7
Black, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1,231	-	-	1	-	-	-	-	-	-	-	6
Male	543	-	-	-	-	-	-	-	-	-	-	2
Female	688	-	-	1	-	-	-	-	-	-	-	4
White, both sexes	1,201	-	-	1	-	-	-	-	-	-	-	6
Male	526	-	-	-	-	-	-	-	-	-	-	2
Female	675	-	-	1	-	-	-	-	-	-	-	4
Black, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Delaware												
All races, both sexes	303	-	-	-	-	-	-	1	-	-	-	-
Male	137	-	-	-	-	-	-	-	-	-	-	-
Female	166	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	271	-	-	-	-	-	-	1	-	-	-	-
Male	117	-	-	-	-	-	-	-	-	-	-	-
Female	154	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	105	-	-	-	-	-	-	-	-	-	-	2
Male	50	-	-	-	-	-	-	-	-	-	-	2
Female	55	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	65	-	-	-	-	-	-	-	-	-	-	1
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Florida												
All races, both sexes	7,775	-	-	-	-	1	-	4	1	8	24	67
Male	3,697	-	-	-	-	-	-	3	1	2	14	33
Female	4,078	-	-	-	-	1	-	1	-	6	10	34
White, both sexes	7,401	-	-	-	-	1	-	3	-	7	21	59
Male	3,479	-	-	-	-	-	-	3	-	1	12	28
Female	3,922	-	-	-	-	1	-	-	-	6	9	31
Black, both sexes	346	-	-	-	-	-	-	1	1	1	3	8
Male	201	-	-	-	-	-	-	-	1	1	2	5
Female	145	-	-	-	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,860	-	-	-	1	-	-	2	-	4	12	26
Male	1,378	-	-	-	-	-	-	1	-	1	5	10
Female	1,482	-	-	-	1	-	-	1	-	3	7	16
White, both sexes	2,527	-	-	-	1	-	-	1	-	2	7	21
Male	1,182	-	-	-	-	-	-	-	-	-	3	7
Female	1,345	-	-	-	1	-	-	1	-	2	4	14
Black, both sexes	314	-	-	-	-	-	-	1	-	2	4	5
Male	187	-	-	-	-	-	-	1	-	1	1	3
Female	127	-	-	-	-	-	-	-	-	1	3	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Hawaii												
All races, both sexes	206	-	-	-	-	-	-	-	-	-	-	1
Male	118	-	-	-	-	-	-	-	-	-	-	-
Female	88	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	80	-	-	-	-	-	-	-	-	-	-	1
Male	43	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	124	-	-	-	-	-	-	-	-	-	-	-
Male	73	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	541	-	-	-	-	-	-	-	-	-	1	5
Male	285	-	-	-	-	-	-	-	-	-	-	3
Female	256	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	537	-	-	-	-	-	-	-	-	-	1	5
Male	282	-	-	-	-	-	-	-	-	-	-	3
Female	255	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Illinois												
All races, both sexes	3,932	-	-	-	-	-	-	-	5	3	13	26
Male	1,856	-	-	-	-	-	-	-	5	2	5	12
Female	2,076	-	-	-	-	-	-	-	-	1	8	14
White, both sexes	3,593	-	-	-	-	-	-	-	3	3	10	19
Male	1,679	-	-	-	-	-	-	-	3	2	5	10
Female	1,914	-	-	-	-	-	-	-	-	1	5	9
Black, both sexes	316	-	-	-	-	-	-	-	2	-	3	7
Male	163	-	-	-	-	-	-	-	2	-	-	2
Female	153	-	-	-	-	-	-	-	-	-	3	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2,810	-	-	-	2	-	-	-	-	2	9	32
Male	1,366	-	-	-	1	-	-	-	-	1	3	13
Female	1,444	-	-	-	1	-	-	-	-	1	6	19
White, both sexes	2,703	-	-	-	1	-	-	-	-	2	8	30
Male	1,310	-	-	-	1	-	-	-	-	1	3	12
Female	1,393	-	-	-	-	-	-	-	-	1	5	18
Black, both sexes	105	-	-	-	1	-	-	-	-	-	1	2
Male	55	-	-	-	-	-	-	-	-	-	-	1
Female	50	-	-	-	1	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Iowa												
All races, both sexes	1,489	-	-	-	-	-	-	-	1	1	6	7
Male	789	-	-	-	-	-	-	-	-	-	4	3
Female	700	-	-	-	-	-	-	-	1	1	2	4
White, both sexes	1,481	-	-	-	-	-	-	-	1	1	6	7
Male	785	-	-	-	-	-	-	-	-	-	4	3
Female	696	-	-	-	-	-	-	-	1	1	2	4
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1,310	-	-	-	-	-	-	1	2	3	5	15
Male	678	-	-	-	-	-	-	1	1	2	2	6
Female	632	-	-	-	-	-	-	-	1	1	3	9
White, both sexes	1,253	-	-	-	-	-	-	1	1	3	4	14
Male	651	-	-	-	-	-	-	1	-	2	1	6
Female	602	-	-	-	-	-	-	-	1	1	3	8
Black, both sexes	35	-	-	-	-	-	-	-	1	-	-	1
Male	18	-	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Kentucky												
All races, both sexes	2,096	-	-	-	-	-	-	-	-	2	10	39
Male	1,025	-	-	-	-	-	-	-	-	1	5	19
Female	1,071	-	-	-	-	-	-	-	-	1	5	20
White, both sexes	2,001	-	-	-	-	-	-	-	-	2	9	36
Male	979	-	-	-	-	-	-	-	-	1	5	18
Female	1,022	-	-	-	-	-	-	-	-	1	4	18
Black, both sexes	89	-	-	-	-	-	-	-	-	-	1	3
Male	42	-	-	-	-	-	-	-	-	-	-	1
Female	47	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,445	-	-	-	-	-	-	-	-	-	6	13
Male	730	-	-	-	-	-	-	-	-	-	2	6
Female	715	-	-	-	-	-	-	-	-	-	4	7
White, both sexes	1,204	-	-	-	-	-	-	-	-	-	2	10
Male	584	-	-	-	-	-	-	-	-	-	-	4
Female	620	-	-	-	-	-	-	-	-	-	2	6
Black, both sexes	237	-	-	-	-	-	-	-	-	-	4	3
Male	142	-	-	-	-	-	-	-	-	-	2	2
Female	95	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Maine												
All races, both sexes	686	-	-	-	-	-	-	-	-	-	3	6
Male	336	-	-	-	-	-	-	-	-	-	1	5
Female	350	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	684	-	-	-	-	-	-	-	-	-	3	6
Male	336	-	-	-	-	-	-	-	-	-	1	5
Female	348	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,536	-	-	-	-	-	-	-	-	-	4	10
Male	659	-	-	-	-	-	-	-	-	-	2	2
Female	877	-	-	-	-	-	-	-	-	-	2	8
White, both sexes	1,329	-	-	-	-	-	-	-	-	-	2	8
Male	564	-	-	-	-	-	-	-	-	-	-	1
Female	765	-	-	-	-	-	-	-	-	-	2	7
Black, both sexes	192	-	-	-	-	-	-	-	-	-	2	2
Male	88	-	-	-	-	-	-	-	-	-	2	1
Female	104	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Massachusetts												
All races, both sexes	2,175	-	-	-	-	-	-	-	2	-	8	10
Male	918	-	-	-	-	-	-	-	1	-	3	3
Female	1,257	-	-	-	-	-	-	-	1	-	5	7
White, both sexes	2,122	-	-	-	-	-	-	-	2	-	6	10
Male	886	-	-	-	-	-	-	-	1	-	2	3
Female	1,236	-	-	-	-	-	-	-	1	-	4	7
Black, both sexes	35	-	-	-	-	-	-	-	-	-	2	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3,761	-	-	-	-	-	-	-	2	-	12	36
Male	1,814	-	-	-	-	-	-	-	1	-	6	15
Female	1,947	-	-	-	-	-	-	-	1	-	6	21
White, both sexes	3,503	-	-	-	-	-	-	-	1	-	10	30
Male	1,671	-	-	-	-	-	-	-	1	-	4	10
Female	1,832	-	-	-	-	-	-	-	-	-	6	20
Black, both sexes	233	-	-	-	-	-	-	-	1	-	1	5
Male	128	-	-	-	-	-	-	-	-	-	1	4
Female	105	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Minnesota												
All races, both sexes	1,593	-	-	-	-	-	-	1	-	-	1	7
Male	792	-	-	-	-	-	-	-	-	-	1	4
Female	801	-	-	-	-	-	-	1	-	-	-	3
White, both sexes	1,564	-	-	-	-	-	-	-	-	-	1	7
Male	773	-	-	-	-	-	-	-	-	-	1	4
Female	791	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	15	-	-	-	-	-	-	1	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,228	-	-	-	1	-	-	-	-	2	9	12
Male	669	-	-	-	1	-	-	-	-	1	6	5
Female	559	-	-	-	-	-	-	-	-	1	3	7
White, both sexes	1,016	-	-	-	-	-	-	-	-	1	4	7
Male	528	-	-	-	-	-	-	-	-	-	2	2
Female	488	-	-	-	-	-	-	-	-	1	2	5
Black, both sexes	210	-	-	-	1	-	-	-	-	1	5	5
Male	140	-	-	-	1	-	-	-	-	1	4	3
Female	70	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Missouri												
All races, both sexes	2,639	-	-	-	1	-	-	1	2	1	15	34
Male	1,253	-	-	-	1	-	-	-	1	1	9	19
Female	1,386	-	-	-	-	-	-	1	1	-	6	15
White, both sexes	2,492	-	-	-	1	-	-	1	2	1	12	33
Male	1,180	-	-	-	1	-	-	-	1	1	7	19
Female	1,312	-	-	-	-	-	-	1	1	-	5	14
Black, both sexes	137	-	-	-	-	-	-	-	-	-	2	1
Male	69	-	-	-	-	-	-	-	-	-	1	-
Female	68	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	511	-	-	-	-	-	-	-	-	-	-	2
Male	234	-	-	-	-	-	-	-	-	-	-	-
Female	277	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	492	-	-	-	-	-	-	-	-	-	-	2
Male	223	-	-	-	-	-	-	-	-	-	-	-
Female	269	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Nebraska												
All races, both sexes	775	-	-	-	-	-	-	-	-	-	1	3
Male	393	-	-	-	-	-	-	-	-	-	-	2
Female	382	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	757	-	-	-	-	-	-	-	-	-	-	2
Male	385	-	-	-	-	-	-	-	-	-	-	2
Female	372	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	980	-	-	-	-	-	-	-	-	3	2	9
Male	452	-	-	-	-	-	-	-	-	1	2	3
Female	528	-	-	-	-	-	-	-	-	2	-	6
White, both sexes	941	-	-	-	-	-	-	-	-	3	2	9
Male	429	-	-	-	-	-	-	-	-	1	2	3
Female	512	-	-	-	-	-	-	-	-	2	-	6
Black, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
New Hampshire												
All races, both sexes	512	-	-	-	-	-	-	-	-	2	-	2
Male	240	-	-	-	-	-	-	-	-	2	-	2
Female	272	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	511	-	-	-	-	-	-	-	-	2	-	2
Male	239	-	-	-	-	-	-	-	-	2	-	2
Female	272	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2,302	-	-	-	-	-	-	-	-	3	9	12
Male	995	-	-	-	-	-	-	-	-	1	2	5
Female	1,307	-	-	-	-	-	-	-	-	2	7	7
White, both sexes	2,084	-	-	-	-	-	-	-	-	2	7	11
Male	890	-	-	-	-	-	-	-	-	1	2	4
Female	1,194	-	-	-	-	-	-	-	-	1	5	7
Black, both sexes	193	-	-	-	-	-	-	-	-	1	1	1
Male	90	-	-	-	-	-	-	-	-	-	-	1
Female	103	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
New Mexico												
All races, both sexes	752	-	-	-	-	-	-	-	-	-	2	5
Male	369	-	-	-	-	-	-	-	-	-	-	4
Female	383	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	735	-	-	-	-	-	-	-	-	-	1	5
Male	360	-	-	-	-	-	-	-	-	-	-	4
Female	375	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	5,412	-	-	-	-	-	-	1	3	2	13	36
Male	2,484	-	-	-	-	-	-	-	1	1	7	19
Female	2,928	-	-	-	-	-	-	1	2	1	6	17
White, both sexes	4,938	-	-	-	-	-	-	-	2	2	9	32
Male	2,234	-	-	-	-	-	-	-	-	1	6	18
Female	2,704	-	-	-	-	-	-	-	2	1	3	14
Black, both sexes	397	-	-	-	-	-	-	1	1	-	4	4
Male	191	-	-	-	-	-	-	-	1	-	1	1
Female	206	-	-	-	-	-	-	1	-	-	3	3
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	62	-	-	-	-	-	-	-	-	-	-	-
Male	50	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
North Carolina												
All races, both sexes	3,388	-	-	-	-	-	-	-	-	-	5	29
Male	1,605	-	-	-	-	-	-	-	-	-	3	13
Female	1,783	-	-	-	-	-	-	-	-	-	2	16
White, both sexes	3,054	-	-	-	-	-	-	-	-	-	1	24
Male	1,414	-	-	-	-	-	-	-	-	-	-	9
Female	1,640	-	-	-	-	-	-	-	-	-	1	15
Black, both sexes	314	-	-	-	-	-	-	-	-	-	4	3
Male	179	-	-	-	-	-	-	-	-	-	3	2
Female	135	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	257	-	-	-	-	-	-	-	-	-	-	1
Male	147	-	-	-	-	-	-	-	-	-	-	1
Female	110	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	249	-	-	-	-	-	-	-	-	-	-	1
Male	145	-	-	-	-	-	-	-	-	-	-	1
Female	104	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Ohio												
All races, both sexes	5,112	-	-	-	-	-	-	-	1	6	19	61
Male	2,480	-	-	-	-	-	-	-	-	4	10	29
Female	2,632	-	-	-	-	-	-	-	1	2	9	32
White, both sexes	4,750	-	-	-	-	-	-	-	-	6	16	48
Male	2,281	-	-	-	-	-	-	-	-	4	9	23
Female	2,469	-	-	-	-	-	-	-	-	2	7	25
Black, both sexes	351	-	-	-	-	-	-	-	1	-	3	12
Male	194	-	-	-	-	-	-	-	-	-	1	6
Female	157	-	-	-	-	-	-	-	1	-	2	6
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,929	1	-	-	-	-	-	-	-	3	8	19
Male	959	1	-	-	-	-	-	-	-	1	4	10
Female	970	-	-	-	-	-	-	-	-	2	4	9
White, both sexes	1,788	-	-	-	-	-	-	-	-	3	5	16
Male	888	-	-	-	-	-	-	-	-	1	2	9
Female	900	-	-	-	-	-	-	-	-	2	3	7
Black, both sexes	52	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	88	1	-	-	-	-	-	-	-	-	3	3
Male	45	1	-	-	-	-	-	-	-	-	2	1
Female	43	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Oregon												
All races, both sexes	1,508	-	-	-	-	-	-	-	-	-	1	13
Male	737	-	-	-	-	-	-	-	-	-	-	10
Female	771	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	1,471	-	-	-	-	-	-	-	-	-	1	13
Male	717	-	-	-	-	-	-	-	-	-	-	10
Female	754	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4,814	-	-	-	-	-	-	-	-	2	6	36
Male	2,257	-	-	-	-	-	-	-	-	2	6	22
Female	2,557	-	-	-	-	-	-	-	-	-	-	14
White, both sexes	4,522	-	-	-	-	-	-	-	-	2	5	31
Male	2,119	-	-	-	-	-	-	-	-	2	5	17
Female	2,403	-	-	-	-	-	-	-	-	-	-	14
Black, both sexes	279	-	-	-	-	-	-	-	-	-	1	5
Male	129	-	-	-	-	-	-	-	-	-	1	5
Female	150	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Rhode Island												
All races, both sexes	438	-	-	1	-	-	-	-	-	-	1	3
Male	185	-	-	1	-	-	-	-	-	-	-	-
Female	253	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	425	-	-	1	-	-	-	-	-	-	1	3
Male	178	-	-	1	-	-	-	-	-	-	-	-
Female	247	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1,668	-	-	-	1	-	-	-	-	1	2	15
Male	832	-	-	-	-	-	-	-	-	1	1	8
Female	836	-	-	-	1	-	-	-	-	-	1	7
White, both sexes	1,468	-	-	-	-	-	-	-	-	1	1	11
Male	718	-	-	-	-	-	-	-	-	1	1	5
Female	750	-	-	-	-	-	-	-	-	-	-	6
Black, both sexes	194	-	-	-	1	-	-	-	-	-	1	4
Male	109	-	-	-	-	-	-	-	-	-	-	3
Female	85	-	-	-	1	-	-	-	-	-	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
South Dakota												
All races, both sexes	324	-	-	-	-	-	-	-	-	-	-	1
Male	185	-	-	-	-	-	-	-	-	-	-	1
Female	139	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	311	-	-	-	-	-	-	-	-	-	-	1
Male	175	-	-	-	-	-	-	-	-	-	-	1
Female	136	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2,569	-	-	-	-	-	-	-	1	1	18	28
Male	1,265	-	-	-	-	-	-	-	1	1	11	15
Female	1,304	-	-	-	-	-	-	-	-	-	7	13
White, both sexes	2,402	-	-	-	-	-	-	-	1	1	18	27
Male	1,173	-	-	-	-	-	-	-	1	1	11	14
Female	1,229	-	-	-	-	-	-	-	-	-	7	13
Black, both sexes	166	-	-	-	-	-	-	-	-	-	-	1
Male	92	-	-	-	-	-	-	-	-	-	-	1
Female	74	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Texas												
All races, both sexes	6,473	-	2	-	-	-	-	-	2	7	17	49
Male	3,104	-	1	-	-	-	-	-	-	3	6	27
Female	3,369	-	1	-	-	-	-	-	2	4	11	22
White, both sexes	6,063	-	1	-	-	-	-	-	1	6	14	41
Male	2,892	-	1	-	-	-	-	-	-	3	6	23
Female	3,171	-	-	-	-	-	-	-	1	3	8	18
Black, both sexes	375	-	-	-	-	-	-	-	1	1	3	7
Male	197	-	-	-	-	-	-	-	-	-	-	4
Female	178	-	-	-	-	-	-	-	1	1	3	3
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	25	-	1	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	1	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	494	-	-	-	-	-	-	-	1	-	-	5
Male	263	-	-	-	-	-	-	-	-	-	-	2
Female	231	-	-	-	-	-	-	-	1	-	-	3
White, both sexes	487	-	-	-	-	-	-	-	1	-	-	5
Male	259	-	-	-	-	-	-	-	-	-	-	2
Female	228	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Vermont												
All races, both sexes	271	-	-	-	-	-	-	-	-	-	-	1
Male	120	-	-	-	-	-	-	-	-	-	-	-
Female	151	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	271	-	-	-	-	-	-	-	-	-	-	1
Male	120	-	-	-	-	-	-	-	-	-	-	-
Female	151	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2,286	-	1	-	-	-	-	-	-	2	3	25
Male	1,057	-	1	-	-	-	-	-	-	1	1	11
Female	1,229	-	-	-	-	-	-	-	-	1	2	14
White, both sexes	2,063	-	1	-	-	-	-	-	-	1	3	23
Male	936	-	1	-	-	-	-	-	-	-	1	10
Female	1,127	-	-	-	-	-	-	-	-	1	2	13
Black, both sexes	214	-	-	-	-	-	-	-	-	1	-	1
Male	115	-	-	-	-	-	-	-	-	1	-	1
Female	99	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Washington												
All races, both sexes	2,343	-	-	-	-	-	-	-	-	-	7	17
Male	1,060	-	-	-	-	-	-	-	-	-	1	10
Female	1,283	-	-	-	-	-	-	-	-	-	6	7
White, both sexes	2,250	-	-	-	-	-	-	-	-	-	7	16
Male	1,019	-	-	-	-	-	-	-	-	-	1	9
Female	1,231	-	-	-	-	-	-	-	-	-	6	7
Black, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1,126	-	-	-	-	-	-	-	1	-	7	12
Male	536	-	-	-	-	-	-	-	1	-	3	5
Female	590	-	-	-	-	-	-	-	-	-	4	7
White, both sexes	1,111	-	-	-	-	-	-	-	1	-	6	12
Male	527	-	-	-	-	-	-	-	1	-	3	5
Female	584	-	-	-	-	-	-	-	-	-	3	7
Black, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
Wisconsin												
All races, both sexes	2,045	-	-	-	1	-	-	-	-	1	7	9
Male	993	-	-	-	-	-	-	-	-	1	3	9
Female	1,052	-	-	-	1	-	-	-	-	-	4	-
White, both sexes	1,982	-	-	-	1	-	-	-	-	1	6	8
Male	964	-	-	-	-	-	-	-	-	1	2	8
Female	1,018	-	-	-	1	-	-	-	-	-	4	-
Black, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	311	-	-	-	-	-	-	-	-	-	-	3
Male	164	-	-	-	-	-	-	-	-	-	-	2
Female	147	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	305	-	-	-	-	-	-	-	-	-	-	3
Male	161	-	-	-	-	-	-	-	-	-	-	2
Female	144	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
United States												
All races, both sexes	1,981	3,828	6,423	9,933	14,345	19,838	21,571	16,491	8,233	2,424	341	2
Male	1,017	2,006	3,347	4,970	7,252	9,941	10,435	7,313	3,274	768	96	1
Female	964	1,822	3,076	4,963	7,093	9,897	11,136	9,178	4,959	1,656	245	1
White, both sexes	1,695	3,387	5,829	9,118	13,351	18,530	20,300	15,612	7,780	2,252	309	1
Male	871	1,766	2,986	4,498	6,676	9,221	9,707	6,844	3,067	691	85	-
Female	824	1,621	2,843	4,620	6,675	9,309	10,593	8,768	4,713	1,561	224	1
Black, both sexes	267	390	502	697	825	1,028	963	650	321	117	30	-
Male	141	216	299	402	479	554	537	334	126	47	9	-
Female	126	174	203	295	346	474	426	316	195	70	21	-
American Indian 1/, both sexes	16	29	51	53	65	89	64	41	15	6	1	-
Male	5	11	31	23	32	39	26	21	5	1	1	-
Female	11	18	20	30	33	50	38	20	10	5	-	-
Asian or Pacific Islander 2/, both sexes	3	22	41	65	104	191	244	188	117	49	1	1
Male	-	13	31	47	65	127	165	114	76	29	1	1
Female	3	9	10	18	39	64	79	74	41	20	-	-
Alabama												
All races, both sexes	43	80	144	233	333	370	395	265	104	39	3	-
Male	20	47	70	130	175	189	189	130	42	9	1	-
Female	23	33	74	103	158	181	206	135	62	30	2	-
White, both sexes	35	64	134	208	306	340	366	240	87	37	3	-
Male	16	37	61	114	155	168	170	117	32	7	1	-
Female	19	27	73	94	151	172	196	123	55	30	2	-
Black, both sexes	8	16	10	25	27	29	28	25	17	2	-	-
Male	4	10	9	16	20	20	19	13	10	2	-	-
Female	4	6	1	9	7	9	9	12	7	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Alaska												
All races, both sexes	3	4	10	11	19	20	16	15	8	1	-	-
Male	2	2	7	5	5	10	5	6	4	-	-	-
Female	1	2	3	6	14	10	11	9	4	1	-	-
White, both sexes	2	3	6	4	15	16	11	11	4	-	-	-
Male	1	1	4	2	5	9	4	3	2	-	-	-
Female	1	2	2	2	10	7	7	8	2	-	-	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	3	5	3	3	5	1	4	1	-	-
Male	1	1	2	2	-	1	1	1	2	-	-	-
Female	-	-	1	3	3	2	4	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	1	-	2	-	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	-	-	-
Arizona												
All races, both sexes	38	85	132	231	325	471	477	372	150	37	4	-
Male	20	49	68	125	159	247	227	160	60	13	1	-
Female	18	36	64	106	166	224	250	212	90	24	3	-
White, both sexes	37	83	128	223	322	462	473	367	146	36	4	-
Male	20	48	67	121	158	243	224	157	59	12	1	-
Female	17	35	61	102	164	219	249	210	87	24	3	-
Black, both sexes	1	1	2	7	2	6	3	3	3	-	-	-
Male	-	1	1	4	1	2	3	1	-	-	-	-
Female	1	-	1	3	1	4	-	2	3	-	-	-
American Indian 1/, both sexes	-	1	2	1	1	2	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	2	1	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	1	1	-	-
Male	-	-	-	-	-	1	-	1	1	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Arkansas												
All races, both sexes	30	51	109	135	177	223	230	171	83	24	3	-
Male	9	31	59	67	108	119	127	89	33	8	1	-
Female	21	20	50	68	69	104	103	82	50	16	2	-
White, both sexes	26	46	99	129	170	213	220	165	82	22	3	-
Male	7	29	52	63	104	112	121	87	32	8	1	-
Female	19	17	47	66	66	101	99	78	50	14	2	-
Black, both sexes	3	5	10	6	7	8	9	6	1	2	-	-
Male	2	2	7	4	4	7	6	2	1	-	-	-
Female	1	3	3	2	3	1	3	4	-	2	-	-
American Indian 1/, both sexes	1	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	196	325	595	903	1,348	1,931	2,346	1,866	1,000	285	46	-
Male	104	184	340	445	669	926	1,096	792	391	107	18	-
Female	92	141	255	458	679	1,005	1,250	1,074	609	178	28	-
White, both sexes	161	272	511	806	1,206	1,729	2,123	1,696	905	239	43	-
Male	85	148	277	380	582	804	964	700	342	83	16	-
Female	76	124	234	426	624	925	1,159	996	563	156	27	-
Black, both sexes	32	41	56	63	76	91	80	52	27	12	3	-
Male	18	26	38	41	41	50	36	26	9	5	2	-
Female	14	15	18	22	35	41	44	26	18	7	1	-
American Indian 1/, both sexes	3	1	6	5	7	10	10	7	2	1	-	-
Male	1	1	5	2	6	5	4	1	-	-	-	-
Female	2	-	1	3	1	5	6	6	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	11	22	29	59	101	133	111	66	33	-	-
Male	-	9	20	22	40	67	92	65	40	19	-	-
Female	-	2	2	7	19	34	41	46	26	14	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Colorado												
All races, both sexes	40	67	94	166	240	325	352	233	142	34	5	-
Male	22	45	61	94	122	156	164	120	71	11	-	-
Female	18	22	33	72	118	169	188	113	71	23	5	-
White, both sexes	37	60	91	157	231	316	346	227	141	34	4	-
Male	21	41	60	90	116	153	160	118	71	11	-	-
Female	16	19	31	67	115	163	186	109	70	23	4	-
Black, both sexes	3	6	3	7	8	6	2	5	1	-	1	-
Male	1	4	1	3	6	3	1	2	-	-	-	-
Female	2	2	2	4	2	3	1	3	1	-	1	-
American Indian 1/, both sexes	-	1	-	1	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	4	-	-	-	-	-
Male	-	-	-	1	-	-	3	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Connecticut												
All races, both sexes	26	34	60	71	124	205	255	251	147	45	5	1
Male	14	15	29	38	68	95	120	94	55	11	2	-
Female	12	19	31	33	56	110	135	157	92	34	3	1
White, both sexes	25	31	58	66	123	199	249	249	145	44	4	1
Male	14	14	28	35	67	91	115	94	53	11	2	-
Female	11	17	30	31	56	108	134	155	92	33	2	1
Black, both sexes	1	3	2	5	1	6	6	2	2	1	1	-
Male	-	1	1	3	1	4	5	-	2	-	-	-
Female	1	2	1	2	-	2	1	2	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Delaware												
All races, both sexes	5	9	12	31	43	67	64	43	17	9	2	-
Male	2	2	7	16	22	33	28	14	8	5	-	-
Female	3	7	5	15	21	34	36	29	9	4	2	-
White, both sexes	4	8	11	27	38	60	57	38	17	8	2	-
Male	2	1	6	14	18	29	24	11	8	4	-	-
Female	2	7	5	13	20	31	33	27	9	4	2	-
Black, both sexes	1	1	1	3	4	7	7	4	-	1	-	-
Male	-	1	1	1	3	4	4	3	-	1	-	-
Female	1	-	-	2	1	3	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	4	4	9	9	23	29	12	10	1	1	-
Male	1	2	1	6	2	12	13	5	6	-	-	-
Female	-	2	3	3	7	11	16	7	4	1	1	-
White, both sexes	-	-	1	2	-	9	9	9	6	1	1	-
Male	-	-	1	2	-	5	2	2	4	-	-	-
Female	-	-	-	-	-	4	7	7	2	1	1	-
Black, both sexes	1	4	3	7	9	13	20	3	4	-	-	-
Male	1	2	-	4	2	6	11	3	2	-	-	-
Female	-	2	3	3	7	7	9	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Florida												
All races, both sexes	152	252	425	685	991	1,385	1,593	1,319	634	202	32	-
Male	78	140	237	364	496	704	773	539	244	63	6	-
Female	74	112	188	321	495	681	820	780	390	139	26	-
White, both sexes	135	231	394	654	940	1,322	1,523	1,275	610	195	31	-
Male	70	129	216	343	470	664	727	515	234	61	6	-
Female	65	102	178	311	470	658	796	760	376	134	25	-
Black, both sexes	17	21	30	28	47	59	62	39	21	7	1	-
Male	8	11	20	19	24	38	41	20	9	2	-	-
Female	9	10	10	9	23	21	21	19	12	5	1	-
American Indian 1/, both sexes	-	-	-	2	2	3	2	3	-	-	-	-
Male	-	-	-	1	1	1	1	2	-	-	-	-
Female	-	-	-	1	1	2	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	1	6	2	3	-	-	-
Male	-	-	1	1	1	1	4	2	1	-	-	-
Female	-	-	-	-	1	-	2	-	2	-	-	-
Georgia												
All races, both sexes	76	141	214	305	436	513	514	384	170	57	5	-
Male	37	74	112	147	227	265	234	183	60	22	-	-
Female	39	67	102	158	209	248	280	201	110	35	5	-
White, both sexes	58	116	194	270	389	459	453	351	155	47	3	-
Male	28	58	96	129	196	234	201	165	49	16	-	-
Female	30	58	98	141	193	225	252	186	106	31	3	-
Black, both sexes	18	25	20	35	44	52	53	31	14	8	2	-
Male	9	16	16	18	29	31	29	18	10	5	-	-
Female	9	9	4	17	15	21	24	13	4	3	2	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	8	2	1	1	-	-
Male	-	-	-	-	1	-	4	-	1	-	-	-
Female	-	-	-	-	1	1	4	2	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Hawaii												
All races, both sexes	2	8	9	17	25	34	45	36	21	8	-	-
Male	1	4	4	8	14	20	28	21	11	7	-	-
Female	1	4	5	9	11	14	17	15	10	1	-	-
White, both sexes	1	3	3	6	12	14	19	12	8	1	-	-
Male	1	1	2	3	7	7	11	7	4	-	-	-
Female	-	2	1	3	5	7	8	5	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	5	6	11	13	19	26	24	13	6	-	-
Male	-	3	2	5	7	12	17	14	7	6	-	-
Female	1	2	4	6	6	7	9	10	6	-	-	-
Idaho												
All races, both sexes	7	17	26	44	65	122	110	100	37	6	1	-
Male	5	7	16	27	32	68	63	48	13	2	1	-
Female	2	10	10	17	33	54	47	52	24	4	-	-
White, both sexes	7	17	25	44	65	121	110	98	37	6	1	-
Male	5	7	15	27	32	67	63	47	13	2	1	-
Female	2	10	10	17	33	54	47	51	24	4	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Illinois												
All races, both sexes	68	138	232	360	506	692	810	642	331	92	14	-
Male	30	68	124	174	257	340	398	287	130	23	1	-
Female	38	70	108	186	249	352	412	355	201	69	13	-
White, both sexes	49	115	202	328	460	628	754	611	314	84	13	-
Male	21	57	106	160	229	307	364	271	124	20	-	-
Female	28	58	96	168	231	321	390	340	190	64	13	-
Black, both sexes	19	22	30	31	43	56	53	28	14	7	1	-
Male	9	11	18	14	26	28	31	14	4	3	1	-
Female	10	11	12	17	17	28	22	14	10	4	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	8	3	3	3	1	-	-
Male	-	-	-	-	2	5	3	2	2	-	-	-
Female	-	-	-	1	-	3	-	1	1	1	-	-
Indiana												
All races, both sexes	52	105	181	282	402	511	539	407	229	50	7	-
Male	31	57	89	141	214	262	250	184	100	19	1	-
Female	21	48	92	141	188	249	289	223	129	31	6	-
White, both sexes	46	102	171	270	385	492	521	392	227	50	6	-
Male	26	54	85	138	207	252	240	173	99	19	-	-
Female	20	48	86	132	178	240	281	219	128	31	6	-
Black, both sexes	6	3	10	11	17	18	18	15	2	-	1	-
Male	5	3	4	2	7	10	10	11	1	-	1	-
Female	1	-	6	9	10	8	8	4	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Iowa												
All races, both sexes	26	50	77	139	202	248	315	256	104	51	6	-
Male	10	24	42	73	113	136	178	130	51	24	1	-
Female	16	26	35	66	89	112	137	126	53	27	5	-
White, both sexes	25	50	77	139	200	246	314	255	103	51	6	-
Male	10	24	42	73	111	135	178	130	50	24	1	-
Female	15	26	35	66	89	111	136	125	53	27	5	-
Black, both sexes	-	-	-	-	2	1	-	1	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Kansas												
All races, both sexes	18	47	68	110	181	263	260	200	103	28	6	-
Male	12	25	24	56	103	136	134	105	60	10	1	-
Female	6	22	44	54	78	127	126	95	43	18	5	-
White, both sexes	18	45	65	102	172	252	246	197	100	27	6	-
Male	12	25	23	52	97	129	127	105	60	10	1	-
Female	6	20	42	50	75	123	119	92	40	17	5	-
Black, both sexes	-	1	2	4	4	8	10	1	2	1	-	-
Male	-	-	-	2	3	6	6	-	-	-	-	-
Female	-	1	2	2	1	2	4	1	2	1	-	-
American Indian 1/, both sexes	-	-	1	4	4	2	4	2	-	-	-	-
Male	-	-	1	2	3	-	1	-	-	-	-	-
Female	-	-	-	2	1	2	3	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Kentucky												
All races, both sexes	54	90	164	256	314	360	389	250	135	30	3	-
Male	28	50	83	131	159	183	198	100	56	10	2	-
Female	26	40	81	125	155	177	191	150	79	20	1	-
White, both sexes	48	82	158	247	298	343	371	244	133	27	3	-
Male	24	46	80	126	150	175	189	97	56	10	2	-
Female	24	36	78	121	148	168	182	147	77	17	1	-
Black, both sexes	6	8	4	8	16	15	17	6	2	3	-	-
Male	4	4	2	4	9	7	8	3	-	-	-	-
Female	2	4	2	4	7	8	9	3	2	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	1	-	-	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	32	63	97	172	223	259	240	206	90	36	8	-
Male	18	38	51	95	119	130	120	98	36	15	2	-
Female	14	25	46	77	104	129	120	108	54	21	6	-
White, both sexes	19	51	76	136	184	218	211	186	76	30	5	-
Male	9	30	43	71	93	110	102	82	29	10	1	-
Female	10	21	33	65	91	108	109	104	47	20	4	-
Black, both sexes	13	12	20	36	39	41	28	18	14	6	3	-
Male	9	8	7	24	26	20	17	14	7	5	1	-
Female	4	4	13	12	13	21	11	4	7	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	2	-	-	-	-
Male	-	-	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Maine												
All races, both sexes	14	29	37	62	100	123	135	104	56	16	1	-
Male	10	15	24	25	52	65	63	51	22	2	1	-
Female	4	14	13	37	48	58	72	53	34	14	-	-
White, both sexes	14	29	36	62	100	123	134	104	56	16	1	-
Male	10	15	24	25	52	65	63	51	22	2	1	-
Female	4	14	12	37	48	58	71	53	34	14	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	25	48	86	135	205	277	319	260	122	42	3	-
Male	11	23	40	63	87	140	132	102	49	7	1	-
Female	14	25	46	72	118	137	187	158	73	35	2	-
White, both sexes	13	43	70	112	176	243	280	230	108	41	3	-
Male	6	22	32	55	69	123	114	91	43	7	1	-
Female	7	21	38	57	107	120	166	139	65	34	2	-
Black, both sexes	12	5	16	20	27	32	34	29	12	1	-	-
Male	5	1	8	8	17	16	15	10	5	-	-	-
Female	7	4	8	12	10	16	19	19	7	1	-	-
American Indian 1/, both sexes	-	-	-	2	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	5	-	2	-	-	-
Male	-	-	-	-	1	-	3	-	1	-	-	-
Female	-	-	-	1	1	1	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Massachusetts												
All races, both sexes	29	67	110	167	254	403	416	402	227	70	10	-
Male	15	32	47	77	131	188	178	157	66	17	3	-
Female	14	35	63	90	123	215	238	245	161	53	7	-
White, both sexes	29	65	108	162	246	392	408	395	222	68	9	-
Male	15	31	46	72	126	179	174	154	63	17	3	-
Female	14	34	62	90	120	213	234	241	159	51	6	-
Black, both sexes	-	1	2	1	6	6	6	5	3	2	1	-
Male	-	-	1	1	3	5	2	1	1	-	-	-
Female	-	1	1	-	3	1	4	4	2	2	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	4	2	5	2	2	2	-	-	-
Male	-	-	-	4	2	4	2	2	2	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Michigan												
All races, both sexes	65	129	204	353	504	737	777	580	288	61	13	-
Male	30	74	109	175	239	358	391	275	117	17	7	-
Female	35	55	95	178	265	379	386	305	171	44	6	-
White, both sexes	57	113	184	331	472	680	740	541	276	59	9	-
Male	25	62	98	159	227	329	369	254	112	16	5	-
Female	32	51	86	172	245	351	371	287	164	43	4	-
Black, both sexes	8	16	20	16	29	48	35	36	12	2	4	-
Male	5	12	11	11	10	26	20	20	5	1	2	-
Female	3	4	9	5	19	22	15	16	7	1	2	-
American Indian 1/, both sexes	-	-	-	4	3	6	1	3	-	-	-	-
Male	-	-	-	3	2	2	1	1	-	-	-	-
Female	-	-	-	1	1	4	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	3	1	-	-	-	-	-
Male	-	-	-	2	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Minnesota												
All races, both sexes	15	31	82	126	197	269	370	296	150	42	5	1
Male	7	18	45	65	101	149	185	127	73	15	1	1
Female	8	13	37	61	96	120	185	169	77	27	4	-
White, both sexes	15	31	78	124	193	264	365	293	147	41	5	-
Male	7	18	43	64	98	146	182	125	70	14	1	-
Female	8	13	35	60	95	118	183	168	77	27	4	-
Black, both sexes	-	-	3	1	2	3	3	2	-	-	-	-
Male	-	-	2	1	2	2	2	1	-	-	-	-
Female	-	-	1	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	2	2	1	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	1	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	3	1	-	1
Male	-	-	-	-	-	-	-	1	3	1	-	1
Female	-	-	-	-	-	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	30	63	90	132	182	232	207	169	67	29	3	-
Male	14	38	47	80	103	139	116	84	27	8	-	-
Female	16	25	43	52	79	93	91	85	40	21	3	-
White, both sexes	22	47	75	109	151	194	183	143	57	21	2	-
Male	8	28	37	64	80	115	98	69	20	5	-	-
Female	14	19	38	45	71	79	85	74	37	16	2	-
Black, both sexes	8	16	15	23	31	38	24	25	9	8	1	-
Male	6	10	10	16	23	24	18	14	7	3	-	-
Female	2	6	5	7	8	14	6	11	2	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Missouri												
All races, both sexes	65	113	159	246	361	518	521	349	187	58	8	-
Male	33	54	68	116	182	256	259	161	78	12	3	-
Female	32	59	91	130	179	262	262	188	109	46	5	-
White, both sexes	60	108	150	229	337	495	489	330	182	55	7	-
Male	31	50	65	108	169	245	241	151	76	12	3	-
Female	29	58	85	121	168	250	248	179	106	43	4	-
Black, both sexes	4	5	8	16	24	19	31	19	4	3	1	-
Male	2	4	2	8	13	9	18	10	2	-	-	-
Female	2	1	6	8	11	10	13	9	2	3	1	-
American Indian 1/, both sexes	1	-	1	1	-	3	1	-	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	-	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	9	15	30	42	77	101	104	88	32	10	1	-
Male	4	9	18	19	40	50	40	41	11	2	-	-
Female	5	6	12	23	37	51	64	47	21	8	1	-
White, both sexes	7	11	27	41	73	100	104	84	32	10	1	-
Male	4	8	15	18	39	49	40	37	11	2	-	-
Female	3	3	12	23	34	51	64	47	21	8	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	4	2	1	4	1	-	4	-	-	-	-
Male	-	1	2	1	1	1	-	4	-	-	-	-
Female	2	3	-	-	3	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Nebraska												
All races, both sexes	10	16	34	49	111	147	174	121	80	24	5	-
Male	6	8	14	21	61	87	92	66	27	6	3	-
Female	4	8	20	28	50	60	82	55	53	18	2	-
White, both sexes	10	16	34	46	106	145	172	119	78	24	5	-
Male	6	8	14	19	58	85	92	65	27	6	3	-
Female	4	8	20	27	48	60	80	54	51	18	2	-
Black, both sexes	-	-	-	2	2	1	1	1	2	-	-	-
Male	-	-	-	1	2	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Nevada												
All races, both sexes	12	37	74	92	179	183	184	129	55	18	3	-
Male	4	22	33	43	90	81	91	55	21	5	1	-
Female	8	15	41	49	89	102	93	74	34	13	2	-
White, both sexes	11	32	70	89	171	177	179	125	53	17	3	-
Male	4	20	30	40	85	78	88	53	19	5	1	-
Female	7	12	40	49	86	99	91	72	34	12	2	-
Black, both sexes	1	3	3	3	6	2	4	2	-	-	-	-
Male	-	1	3	3	4	-	2	1	-	-	-	-
Female	1	2	-	-	2	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	-	2	4	1	1	1	1	-	-
Male	-	1	-	-	1	3	1	1	1	-	-	-
Female	-	1	1	-	1	1	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
New Hampshire												
All races, both sexes	14	24	26	47	76	83	108	92	28	9	1	-
Male	8	14	14	18	33	44	59	35	8	3	-	-
Female	6	10	12	29	43	39	49	57	20	6	1	-
White, both sexes	14	24	26	47	76	82	108	92	28	9	1	-
Male	8	14	14	18	33	43	59	35	8	3	-	-
Female	6	10	12	29	43	39	49	57	20	6	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	42	57	105	150	250	439	524	424	204	75	8	-
Male	20	31	53	79	105	195	240	171	70	22	1	-
Female	22	26	52	71	145	244	284	253	134	53	7	-
White, both sexes	38	43	87	125	224	397	488	402	184	71	5	-
Male	18	25	42	62	91	175	221	163	63	22	1	-
Female	20	18	45	63	133	222	267	239	121	49	4	-
Black, both sexes	4	14	16	24	23	37	30	20	15	4	3	-
Male	2	6	9	16	11	18	16	8	3	-	-	-
Female	2	8	7	8	12	19	14	12	12	4	3	-
American Indian 1/, both sexes	-	-	2	-	-	1	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	3	4	6	2	5	-	-	-
Male	-	-	-	1	3	2	3	-	4	-	-	-
Female	-	-	-	-	-	2	3	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
New Mexico												
All races, both sexes	9	20	49	54	91	155	180	103	64	15	5	-
Male	2	7	27	32	41	79	87	57	24	7	2	-
Female	7	13	22	22	50	76	93	46	40	8	3	-
White, both sexes	9	19	47	53	91	154	174	101	61	15	5	-
Male	2	6	26	31	41	78	85	55	23	7	2	-
Female	7	13	21	22	50	76	89	46	38	8	3	-
Black, both sexes	-	1	-	-	-	1	1	1	2	-	-	-
Male	-	1	-	-	-	1	1	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	2	1	-	-	4	1	1	-	-	-
Male	-	-	1	1	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	3	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
New York												
All races, both sexes	68	154	300	437	616	988	1,127	936	522	185	24	-
Male	34	78	169	216	286	485	546	399	182	53	8	-
Female	34	76	131	221	330	503	581	537	340	132	16	-
White, both sexes	59	135	266	385	563	908	1,033	869	479	174	22	-
Male	29	69	148	183	260	442	494	367	162	48	7	-
Female	30	66	118	202	303	466	539	502	317	126	15	-
Black, both sexes	9	18	31	50	46	65	76	50	33	8	1	-
Male	5	8	18	31	23	33	37	19	11	3	-	-
Female	4	10	13	19	23	32	39	31	22	5	1	-
American Indian 1/, both sexes	-	1	1	-	3	7	1	2	-	-	-	-
Male	-	1	1	-	1	3	1	2	-	-	-	-
Female	-	-	-	-	2	4	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	4	8	17	15	10	3	1	-
Male	-	-	2	2	2	7	14	11	9	2	1	-
Female	-	-	-	-	2	1	3	4	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
North Carolina												
All races, both sexes	64	156	257	378	461	659	671	434	206	61	7	-
Male	37	77	133	190	221	324	307	188	84	25	3	-
Female	27	79	124	188	240	335	364	246	122	36	4	-
White, both sexes	54	138	213	342	423	606	611	393	190	53	6	-
Male	32	65	105	171	201	298	272	165	75	19	2	-
Female	22	73	108	171	222	308	339	228	115	34	4	-
Black, both sexes	9	17	40	34	36	51	58	39	15	8	-	-
Male	4	12	27	18	18	25	34	22	8	6	-	-
Female	5	5	13	16	18	26	24	17	7	2	-	-
American Indian 1/, both sexes	1	1	2	2	1	1	1	2	1	-	1	-
Male	1	-	1	1	1	-	-	1	1	-	1	-
Female	-	1	1	1	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	5	10	14	30	54	46	49	35	9	2	-
Male	1	1	5	9	17	32	31	26	20	3	1	-
Female	1	4	5	5	13	22	15	23	15	6	1	-
White, both sexes	1	5	8	13	30	52	45	48	35	9	2	-
Male	1	1	4	9	17	32	30	26	20	3	1	-
Female	-	4	4	4	13	20	15	22	15	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	2	1	-	2	1	1	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	1	1	-	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Ohio												
All races, both sexes	101	180	321	500	671	999	1,010	765	360	101	17	-
Male	56	91	174	240	334	508	498	356	144	30	6	-
Female	45	89	147	260	337	491	512	409	216	71	11	-
White, both sexes	84	152	301	458	631	922	950	723	346	97	16	-
Male	48	73	161	216	309	469	464	329	140	30	6	-
Female	36	79	140	242	322	453	486	394	206	67	10	-
Black, both sexes	17	28	19	42	39	77	56	39	13	4	1	-
Male	8	18	13	24	25	39	31	26	3	-	-	-
Female	9	10	6	18	14	38	25	13	10	4	1	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	3	1	-	-	-
Male	-	-	-	-	-	-	3	1	1	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Oklahoma												
All races, both sexes	46	102	155	238	317	366	316	220	105	29	4	-
Male	22	58	74	112	170	186	165	97	46	12	1	-
Female	24	44	81	126	147	180	151	123	59	17	3	-
White, both sexes	43	95	140	219	295	341	291	210	101	26	3	-
Male	22	53	64	105	162	170	151	92	45	11	1	-
Female	21	42	76	114	133	171	140	118	56	15	2	-
Black, both sexes	2	3	4	8	8	7	9	6	3	1	1	-
Male	-	2	2	4	4	3	6	3	1	1	-	-
Female	2	1	2	4	4	4	3	3	2	-	1	-
American Indian 1/, both sexes	1	4	11	11	14	18	15	4	1	2	-	-
Male	-	3	8	3	4	13	8	2	-	-	-	-
Female	1	1	3	8	10	5	7	2	1	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Oregon												
All races, both sexes	24	69	91	134	230	291	283	233	113	24	2	-
Male	18	36	45	65	108	141	139	113	51	10	1	-
Female	6	33	46	69	122	150	144	120	62	14	1	-
White, both sexes	24	64	86	132	224	283	278	231	111	22	2	-
Male	18	34	42	64	105	136	136	112	50	9	1	-
Female	6	30	44	68	119	147	142	119	61	13	1	-
Black, both sexes	-	3	2	1	3	1	3	1	1	1	-	-
Male	-	1	2	1	2	-	2	-	-	1	-	-
Female	-	2	-	-	1	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	2	3	1	3	3	1	-	1	-	-	-
Male	-	1	1	-	1	1	-	-	1	-	-	-
Female	-	1	2	1	2	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	4	1	1	-	1	-	-
Male	-	-	-	-	-	4	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Pennsylvania												
All races, both sexes	64	151	250	364	581	922	1,067	840	397	117	17	-
Male	29	77	137	188	278	477	487	367	151	33	3	-
Female	35	74	113	176	303	445	580	473	246	84	14	-
White, both sexes	51	129	228	332	542	861	1,028	809	377	110	17	-
Male	23	68	127	172	259	445	468	354	146	30	3	-
Female	28	61	101	160	283	416	560	455	231	80	14	-
Black, both sexes	12	22	21	29	39	58	36	30	20	6	-	-
Male	6	9	10	13	19	29	16	13	5	3	-	-
Female	6	13	11	16	20	29	20	17	15	3	-	-
American Indian 1/, both sexes	1	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	2	3	1	-	1	-	-
Male	-	-	-	2	-	2	3	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Rhode Island												
All races, both sexes	7	8	21	33	58	72	104	87	31	10	2	-
Male	4	5	5	12	31	36	36	39	14	1	1	-
Female	3	3	16	21	27	36	68	48	17	9	1	-
White, both sexes	7	8	21	31	54	72	100	84	31	10	2	-
Male	4	5	5	11	28	36	35	37	14	1	1	-
Female	3	3	16	20	26	36	65	47	17	9	1	-
Black, both sexes	-	-	-	2	3	-	3	3	-	-	-	-
Male	-	-	-	1	2	-	-	2	-	-	-	-
Female	-	-	-	1	1	-	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	37	81	144	195	254	314	311	190	88	31	4	-
Male	14	44	73	106	139	169	155	79	36	7	-	-
Female	23	37	71	89	115	145	156	111	52	24	4	-
White, both sexes	28	66	119	166	229	286	281	169	80	28	3	-
Male	10	34	58	87	122	152	137	70	34	7	-	-
Female	18	32	61	79	107	134	144	99	46	21	3	-
Black, both sexes	9	15	23	28	25	28	28	20	8	3	1	-
Male	4	10	13	18	17	17	17	8	2	-	-	-
Female	5	5	10	10	8	11	11	12	6	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	-	-	1	-	-	-	-
Male	-	-	2	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
South Dakota												
All races, both sexes	4	10	13	20	52	54	71	50	36	12	1	-
Male	2	4	6	9	36	29	45	29	18	6	-	-
Female	2	6	7	11	16	25	26	21	18	6	1	-
White, both sexes	3	7	11	20	48	53	70	50	35	12	1	-
Male	1	3	4	9	32	28	44	29	18	6	-	-
Female	2	4	7	11	16	25	26	21	17	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	3	2	-	4	1	1	-	1	-	-	-
Male	1	1	2	-	4	1	1	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	71	132	208	311	386	482	447	288	152	40	4	-
Male	36	61	106	151	197	242	240	126	62	16	-	-
Female	35	71	102	160	189	240	207	162	90	24	4	-
White, both sexes	61	117	192	289	359	459	422	273	141	38	4	-
Male	30	56	97	138	184	226	228	117	56	14	-	-
Female	31	61	95	151	175	233	194	156	85	24	4	-
Black, both sexes	10	15	15	22	27	23	25	15	11	2	-	-
Male	6	5	9	13	13	16	12	9	6	2	-	-
Female	4	10	6	9	14	7	13	6	5	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Texas												
All races, both sexes	124	234	413	676	931	1,248	1,279	905	451	116	19	-
Male	64	106	201	325	480	623	624	423	179	37	5	-
Female	60	128	212	351	451	625	655	482	272	79	14	-
White, both sexes	101	206	385	615	867	1,180	1,218	863	436	111	18	-
Male	52	98	184	295	440	588	591	399	173	34	5	-
Female	49	108	201	320	427	592	627	464	263	77	13	-
Black, both sexes	22	25	27	57	58	63	53	38	14	5	1	-
Male	12	8	16	28	37	33	30	21	5	3	-	-
Female	10	17	11	29	21	30	23	17	9	2	1	-
American Indian 1/, both sexes	-	1	-	2	1	2	3	-	-	-	-	-
Male	-	-	-	1	1	1	1	-	-	-	-	-
Female	-	1	-	1	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	2	5	3	5	4	1	-	-	-
Male	-	-	1	1	2	1	2	3	1	-	-	-
Female	1	2	-	1	3	2	3	1	-	-	-	-
Utah												
All races, both sexes	10	23	28	58	71	103	93	73	22	6	1	-
Male	9	7	11	33	45	58	43	40	14	1	-	-
Female	1	16	17	25	26	45	50	33	8	5	1	-
White, both sexes	9	23	28	58	71	99	91	73	22	6	1	-
Male	8	7	11	33	45	56	42	40	14	1	-	-
Female	1	16	17	25	26	43	49	33	8	5	1	-
Black, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	2	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Vermont												
All races, both sexes	-	10	21	17	34	47	74	36	20	10	1	-
Male	-	5	13	8	14	25	30	13	9	3	-	-
Female	-	5	8	9	20	22	44	23	11	7	1	-
White, both sexes	-	10	21	17	34	47	74	36	20	10	1	-
Male	-	5	13	8	14	25	30	13	9	3	-	-
Female	-	5	8	9	20	22	44	23	11	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	31	87	135	194	350	410	468	352	160	57	11	-
Male	18	45	69	94	168	185	232	150	59	18	5	-
Female	13	42	66	100	182	225	236	202	101	39	6	-
White, both sexes	30	78	116	163	313	372	428	330	145	50	10	-
Male	17	39	62	74	143	169	207	138	53	18	4	-
Female	13	39	54	89	170	203	221	192	92	32	6	-
Black, both sexes	1	9	19	31	35	35	37	22	15	7	1	-
Male	1	6	7	20	24	13	23	12	6	-	1	-
Female	-	3	12	11	11	22	14	10	9	7	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	3	3	-	-	-	-	-
Male	-	-	-	-	-	3	2	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Washington												
All races, both sexes	50	89	132	220	293	438	491	382	178	42	4	-
Male	29	43	61	109	148	203	224	155	64	13	-	-
Female	21	46	71	111	145	235	267	227	114	29	4	-
White, both sexes	47	82	122	213	285	415	475	369	173	42	4	-
Male	28	43	56	105	145	193	214	149	63	13	-	-
Female	19	39	66	108	140	222	261	220	110	29	4	-
Black, both sexes	2	1	7	2	1	6	7	3	1	-	-	-
Male	1	-	4	1	1	3	5	-	-	-	-	-
Female	1	1	3	1	-	3	2	3	1	-	-	-
American Indian 1/, both sexes	-	5	3	2	5	7	1	3	2	-	-	-
Male	-	-	1	1	1	-	1	2	-	-	-	-
Female	-	5	2	1	4	7	-	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	3	2	10	8	7	2	-	-	-
Male	-	-	-	2	1	7	4	4	1	-	-	-
Female	1	1	-	1	1	3	4	3	1	-	-	-
West Virginia												
All races, both sexes	27	46	65	129	179	235	205	122	81	15	2	-
Male	13	23	33	53	94	120	102	54	30	4	1	-
Female	14	23	32	76	85	115	103	68	51	11	1	-
White, both sexes	26	46	63	125	178	235	202	121	81	14	1	-
Male	13	23	31	51	93	120	100	53	30	4	-	-
Female	13	23	32	74	85	115	102	68	51	10	1	-
Black, both sexes	1	-	2	4	1	-	3	1	-	1	1	-
Male	-	-	2	2	1	-	2	1	-	-	1	-
Female	1	-	-	2	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Wisconsin												
All races, both sexes	35	59	103	174	239	411	441	331	181	51	2	-
Male	20	29	60	84	130	205	215	147	76	14	-	-
Female	15	30	43	90	109	206	226	184	105	37	2	-
White, both sexes	28	54	96	167	232	398	435	326	178	50	2	-
Male	16	27	55	79	126	201	214	145	76	14	-	-
Female	12	27	41	88	106	197	221	181	102	36	2	-
Black, both sexes	6	4	5	5	7	8	3	2	3	-	-	-
Male	4	2	4	4	4	1	1	-	-	-	-	-
Female	2	2	1	1	3	7	2	2	3	-	-	-
American Indian 1/, both sexes	1	1	2	2	-	2	2	2	-	1	-	-
Male	-	-	1	1	-	2	-	1	-	-	-	-
Female	1	1	1	1	-	-	2	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
Wyoming												
All races, both sexes	8	13	17	26	42	56	69	53	20	4	-	-
Male	3	7	9	11	27	31	38	25	11	-	-	-
Female	5	6	8	15	15	25	31	28	9	4	-	-
White, both sexes	7	12	17	25	41	56	68	53	19	4	-	-
Male	3	6	9	10	26	31	38	25	11	-	-	-
Female	4	6	8	15	15	25	30	28	8	4	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	1	1	-	1	-	-	-	-	-
Male	-	1	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
United States												
All races, both sexes	3,613	6	26	45	54	56	79	92	102	141	232	277
Male	1,296	3	17	30	41	36	46	56	49	62	93	96
Female	2,317	3	9	15	13	20	33	36	53	79	139	181
White, both sexes	2,497	4	13	19	14	16	39	50	48	78	142	168
Male	805	2	7	15	12	9	22	31	17	34	56	55
Female	1,692	2	6	4	2	7	17	19	31	44	86	113
Black, both sexes	957	2	12	26	37	37	37	39	52	61	78	102
Male	423	1	10	15	27	25	21	22	30	26	36	38
Female	534	1	2	11	10	12	16	17	22	35	42	64
American Indian 1/, both sexes	20	-	-	-	2	-	1	-	-	1	4	2
Male	7	-	-	-	1	-	1	-	-	1	-	1
Female	13	-	-	-	1	-	-	-	-	-	4	1
Asian or Pacific Islander 2/, both sexes	139	-	1	-	1	3	2	3	2	1	8	5
Male	61	-	-	-	1	2	2	3	2	1	1	2
Female	78	-	1	-	-	1	-	-	-	-	7	3
Alabama												
All races, both sexes	58	-	1	-	2	-	1	1	2	4	6	1
Male	27	-	1	-	2	-	1	1	1	2	2	-
Female	31	-	-	-	-	-	-	-	1	2	4	1
White, both sexes	35	-	-	-	1	-	-	-	1	3	2	1
Male	14	-	-	-	1	-	-	-	1	2	1	-
Female	21	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	23	-	1	-	1	-	1	1	1	1	4	-
Male	13	-	1	-	1	-	1	1	-	-	1	-
Female	10	-	-	-	-	-	-	-	1	1	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Alaska												
All races, both sexes	11	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	8	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	74	-	1	1	4	-	-	1	3	4	4	2
Male	32	-	1	1	2	-	-	1	1	4	1	-
Female	42	-	-	-	2	-	-	-	2	-	3	2
White, both sexes	66	-	1	1	1	-	-	1	3	3	3	1
Male	30	-	1	1	1	-	-	1	1	3	1	-
Female	36	-	-	-	-	-	-	-	2	-	2	1
Black, both sexes	4	-	-	-	1	-	-	-	-	1	1	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	1	-	-	-	-	-	1	1
American Indian 1/, both sexes	4	-	-	-	2	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	3	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Arkansas												
All races, both sexes	51	-	-	-	-	1	2	1	1	2	4	-
Male	14	-	-	-	-	1	-	-	-	1	1	-
Female	37	-	-	-	-	-	2	1	1	1	3	-
White, both sexes	39	-	-	-	-	-	1	1	-	2	4	-
Male	11	-	-	-	-	-	-	-	-	1	1	-
Female	28	-	-	-	-	-	1	1	-	1	3	-
Black, both sexes	12	-	-	-	-	1	1	-	1	-	-	-
Male	3	-	-	-	-	1	-	-	-	-	-	-
Female	9	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	473	1	3	4	7	9	8	11	11	19	29	37
Male	173	-	2	3	7	5	5	7	6	10	7	16
Female	300	1	1	1	-	4	3	4	5	9	22	21
White, both sexes	310	1	2	2	-	5	4	4	5	9	15	27
Male	105	-	1	1	-	4	2	2	-	5	4	12
Female	205	1	1	1	-	1	2	2	5	4	11	15
Black, both sexes	91	-	1	2	6	3	4	4	4	9	7	9
Male	38	-	1	2	6	-	3	2	4	4	2	3
Female	53	-	-	-	-	3	1	2	-	5	5	6
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	69	-	-	-	1	1	-	3	2	-	6	1
Male	29	-	-	-	1	1	-	3	2	-	1	1
Female	40	-	-	-	-	-	-	-	-	-	5	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Colorado												
All races, both sexes	46	-	-	-	-	1	-	1	4	1	3	4
Male	16	-	-	-	-	1	-	1	3	-	1	2
Female	30	-	-	-	-	-	-	-	1	1	2	2
White, both sexes	40	-	-	-	-	1	-	1	3	1	2	3
Male	13	-	-	-	-	1	-	1	2	-	1	1
Female	27	-	-	-	-	-	-	-	1	1	1	2
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	1	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	46	-	-	1	1	-	-	-	-	3	4	4
Male	13	-	-	1	1	-	-	-	-	1	2	1
Female	33	-	-	-	-	-	-	-	-	2	2	3
White, both sexes	35	-	-	1	-	-	-	-	-	2	3	1
Male	6	-	-	1	-	-	-	-	-	-	1	-
Female	29	-	-	-	-	-	-	-	-	2	2	1
Black, both sexes	10	-	-	-	1	-	-	-	-	1	1	3
Male	6	-	-	-	1	-	-	-	-	1	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Delaware												
All races, both sexes	6	-	-	-	-	-	-	1	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	1	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	12	1	-	-	-	-	-	1	-	2	3	1
Male	5	1	-	-	-	-	-	1	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	2	2	1
White, both sexes	2	-	-	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	10	1	-	-	-	-	-	1	-	1	2	1
Male	5	1	-	-	-	-	-	1	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Florida												
All races, both sexes	170	1	2	4	4	2	7	5	7	3	7	17
Male	55	-	1	3	3	-	5	4	1	-	1	3
Female	115	1	1	1	1	2	2	1	6	3	6	14
White, both sexes	113	-	1	2	2	1	4	3	3	1	6	9
Male	35	-	-	2	2	-	3	2	-	-	1	1
Female	78	-	1	-	-	1	1	1	3	1	5	8
Black, both sexes	53	1	1	2	2	1	3	2	4	2	-	7
Male	18	-	1	1	1	-	2	2	1	-	-	1
Female	35	1	-	1	1	1	1	-	3	2	-	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	1	-
Georgia												
All races, both sexes	119	-	1	4	3	2	2	3	4	7	9	13
Male	40	-	1	2	2	1	1	2	2	2	3	4
Female	79	-	-	2	1	1	1	1	2	5	6	9
White, both sexes	51	-	-	1	-	-	-	1	-	3	3	3
Male	14	-	-	1	-	-	-	1	-	1	1	-
Female	37	-	-	-	-	-	-	-	-	2	2	3
Black, both sexes	63	-	1	3	3	2	2	2	4	4	6	9
Male	24	-	1	1	2	1	1	1	2	1	2	4
Female	39	-	-	2	1	1	1	1	2	3	4	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Hawaii												
All races, both sexes	21	-	-	-	-	1	-	-	-	1	-	1
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	1	-	-	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	1	-	-	-	-	-	1
Idaho												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Illinois												
All races, both sexes	169	-	2	3	2	4	7	5	12	9	10	18
Male	79	-	2	2	2	4	3	4	9	5	8	8
Female	90	-	-	1	-	-	4	1	3	4	2	10
White, both sexes	85	-	-	2	-	-	3	2	2	6	5	7
Male	35	-	-	2	-	-	1	2	1	4	4	3
Female	50	-	-	-	-	-	2	-	1	2	1	4
Black, both sexes	82	-	2	1	2	4	4	3	10	3	5	11
Male	44	-	2	-	2	4	2	2	8	1	4	5
Female	38	-	-	1	-	-	2	1	2	2	1	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	72	1	1	-	1	1	3	1	2	6	4	6
Male	19	1	-	-	-	1	1	-	1	3	2	3
Female	53	-	1	-	1	-	2	1	1	3	2	3
White, both sexes	56	1	1	-	-	1	1	1	1	4	2	5
Male	15	1	-	-	-	1	1	-	1	1	1	2
Female	41	-	1	-	-	-	-	1	-	3	1	3
Black, both sexes	15	-	-	-	1	-	2	-	1	2	2	1
Male	4	-	-	-	-	-	-	-	-	2	1	1
Female	11	-	-	-	1	-	2	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Iowa												
All races, both sexes	37	-	-	-	-	1	1	-	2	2	5	-
Male	12	-	-	-	-	1	1	-	1	1	2	-
Female	25	-	-	-	-	-	-	-	1	1	3	-
White, both sexes	32	-	-	-	-	-	1	-	1	2	3	-
Male	8	-	-	-	-	-	1	-	-	1	1	-
Female	24	-	-	-	-	-	-	-	1	1	2	-
Black, both sexes	4	-	-	-	-	1	-	-	1	-	1	-
Male	4	-	-	-	-	1	-	-	1	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	34	-	-	-	-	-	1	3	2	2	3	2
Male	11	-	-	-	-	-	1	2	1	-	1	1
Female	23	-	-	-	-	-	-	1	1	2	2	1
White, both sexes	29	-	-	-	-	-	-	3	1	2	2	2
Male	9	-	-	-	-	-	-	2	-	-	1	1
Female	20	-	-	-	-	-	-	1	1	2	1	1
Black, both sexes	5	-	-	-	-	-	1	-	1	-	1	-
Male	2	-	-	-	-	-	1	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Kentucky												
All races, both sexes	59	-	1	-	-	1	1	2	1	3	3	6
Male	21	-	1	-	-	1	-	1	1	3	2	2
Female	38	-	-	-	-	-	1	1	-	-	1	4
White, both sexes	49	-	1	-	-	-	1	1	1	2	2	4
Male	19	-	1	-	-	-	-	1	1	2	2	2
Female	30	-	-	-	-	-	1	-	-	-	-	2
Black, both sexes	10	-	-	-	-	1	-	1	-	1	1	2
Male	2	-	-	-	-	1	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	1	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	49	-	2	1	1	-	1	1	2	3	4	5
Male	19	-	2	-	1	-	1	1	1	1	1	3
Female	30	-	-	1	-	-	-	-	1	2	3	2
White, both sexes	26	-	-	-	-	-	-	1	1	2	2	3
Male	6	-	-	-	-	-	-	1	-	-	-	1
Female	20	-	-	-	-	-	-	-	1	2	2	2
Black, both sexes	23	-	2	1	1	-	1	-	1	1	2	2
Male	13	-	2	-	1	-	1	-	1	1	1	2
Female	10	-	-	1	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Maine												
All races, both sexes	13	-	-	-	1	-	-	-	-	1	-	-
Male	4	-	-	-	1	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	13	-	-	-	1	-	-	-	-	1	-	-
Male	4	-	-	-	1	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	55	-	-	-	2	1	1	3	2	2	6	4
Male	21	-	-	-	1	1	1	1	-	-	2	-
Female	34	-	-	-	1	-	-	2	2	2	4	4
White, both sexes	27	-	-	-	-	-	-	1	-	2	2	3
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	1	-	2	2	3
Black, both sexes	28	-	-	-	2	1	1	2	2	-	4	1
Male	13	-	-	-	1	1	1	1	-	-	2	-
Female	15	-	-	-	1	-	-	1	2	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Massachusetts												
All races, both sexes	52	-	-	-	-	-	2	2	-	1	5	5
Male	11	-	-	-	-	-	-	2	-	-	2	1
Female	41	-	-	-	-	-	2	-	-	1	3	4
White, both sexes	42	-	-	-	-	-	1	2	-	1	2	4
Male	9	-	-	-	-	-	-	2	-	-	2	1
Female	33	-	-	-	-	-	1	-	-	1	-	3
Black, both sexes	9	-	-	-	-	-	1	-	-	-	3	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	-	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	121	-	1	4	3	5	3	8	1	4	9	9
Male	45	-	-	2	2	4	2	5	-	3	4	3
Female	76	-	1	2	1	1	1	3	1	1	5	6
White, both sexes	83	-	-	-	1	1	3	5	-	2	5	7
Male	29	-	-	-	1	1	2	2	-	2	4	2
Female	54	-	-	-	-	-	1	3	-	-	1	5
Black, both sexes	37	-	1	4	2	4	-	3	1	2	3	2
Male	16	-	-	2	1	3	-	3	-	1	-	1
Female	21	-	1	2	1	1	-	-	1	1	3	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Minnesota												
All races, both sexes	47	-	-	1	-	1	-	-	1	1	2	5
Male	14	-	-	1	-	1	-	-	-	-	1	3
Female	33	-	-	-	-	-	-	-	1	1	1	2
White, both sexes	37	-	-	-	-	-	-	-	1	-	2	4
Male	10	-	-	-	-	-	-	-	-	-	1	2
Female	27	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	6	-	-	1	-	1	-	-	-	1	-	1
Male	3	-	-	1	-	1	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	46	-	-	-	1	1	-	2	4	3	2	5
Male	14	-	-	-	1	1	-	1	1	-	-	1
Female	32	-	-	-	-	-	-	1	3	3	2	4
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	30	-	-	-	1	1	-	2	4	3	2	3
Male	8	-	-	-	1	1	-	1	1	-	-	-
Female	22	-	-	-	-	-	-	1	3	3	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Missouri												
All races, both sexes	68	-	-	-	1	-	3	4	1	5	6	7
Male	27	-	-	-	1	-	3	3	1	4	3	3
Female	41	-	-	-	-	-	-	1	-	1	3	4
White, both sexes	50	-	-	-	-	-	2	2	-	2	4	4
Male	15	-	-	-	-	-	2	2	-	1	2	1
Female	35	-	-	-	-	-	-	-	-	1	2	3
Black, both sexes	18	-	-	-	1	-	1	2	1	3	2	3
Male	12	-	-	-	1	-	1	1	1	3	1	2
Female	6	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	16	-	-	1	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	1	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Nebraska												
All races, both sexes	24	-	-	-	-	1	-	-	2	1	-	1
Male	7	-	-	-	-	-	-	-	1	1	-	-
Female	17	-	-	-	-	1	-	-	1	-	-	1
White, both sexes	21	-	-	-	-	-	-	-	1	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	16	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	3	-	-	-	-	1	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	18	-	-	-	-	-	1	-	-	1	2	3
Male	5	-	-	-	-	-	1	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	12	-	-	-	-	-	1	-	-	-	1	2
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
New Hampshire												
All races, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	108	-	1	-	2	4	2	1	2	7	9	12
Male	40	-	1	-	2	2	2	-	1	2	4	4
Female	68	-	-	-	-	2	-	1	1	5	5	8
White, both sexes	67	-	-	-	1	-	1	1	1	4	5	6
Male	23	-	-	-	1	-	1	-	-	2	2	3
Female	44	-	-	-	-	-	-	1	1	2	3	3
Black, both sexes	37	-	1	-	1	3	1	-	1	3	4	6
Male	15	-	1	-	1	1	1	-	1	-	2	1
Female	22	-	-	-	-	2	-	-	-	3	2	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
New Mexico												
All races, both sexes	24	-	-	-	2	-	-	1	-	-	1	3
Male	11	-	-	-	2	-	-	1	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	22	-	-	-	2	-	-	1	-	-	1	2
Male	9	-	-	-	2	-	-	1	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	241	-	3	3	3	3	6	10	6	6	16	17
Male	83	-	1	3	1	3	2	3	2	3	8	6
Female	158	-	2	-	2	-	4	7	4	3	8	11
White, both sexes	151	-	-	2	-	1	5	4	3	3	8	8
Male	45	-	-	2	-	1	2	2	1	1	4	3
Female	106	-	-	-	-	-	3	2	2	2	4	5
Black, both sexes	85	-	2	1	3	2	1	6	3	3	8	9
Male	35	-	1	1	1	2	-	1	1	2	4	3
Female	50	-	1	-	2	-	1	5	2	1	4	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
North Carolina												
All races, both sexes	123	-	-	1	1	3	3	4	7	8	9	8
Male	44	-	-	1	1	1	-	1	3	4	3	2
Female	79	-	-	-	-	2	3	3	4	4	6	6
White, both sexes	71	-	-	-	-	1	1	2	4	5	6	1
Male	24	-	-	-	-	-	-	-	2	3	2	-
Female	47	-	-	-	-	1	1	2	2	2	4	1
Black, both sexes	51	-	-	1	1	2	2	2	3	3	3	7
Male	19	-	-	1	1	1	-	1	1	1	1	2
Female	32	-	-	-	-	1	2	1	2	2	2	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	7	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Ohio												
All races, both sexes	153	-	2	2	1	1	3	4	4	6	8	11
Male	52	-	1	1	1	-	2	3	3	3	2	4
Female	101	-	1	1	-	1	1	1	1	3	6	7
White, both sexes	116	-	2	-	-	1	2	2	2	2	7	6
Male	33	-	1	-	-	-	1	2	1	-	2	2
Female	83	-	1	-	-	1	1	-	1	2	5	4
Black, both sexes	37	-	-	2	1	-	1	2	2	4	1	5
Male	19	-	-	1	1	-	1	1	2	3	-	2
Female	18	-	-	1	-	-	-	1	-	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	47	-	2	1	4	2	2	1	1	2	3	2
Male	19	-	2	1	2	2	1	1	1	-	-	-
Female	28	-	-	-	2	-	1	-	-	2	3	2
White, both sexes	35	-	2	1	1	-	1	1	-	2	3	2
Male	13	-	2	1	1	-	-	1	-	-	-	-
Female	22	-	-	-	-	-	1	-	-	2	3	2
Black, both sexes	11	-	-	-	3	2	1	-	1	-	-	-
Male	6	-	-	-	1	2	1	-	1	-	-	-
Female	5	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Oregon												
All races, both sexes	79	-	1	-	-	1	1	-	2	2	4	6
Male	35	-	-	-	-	1	1	-	2	2	4	4
Female	44	-	1	-	-	-	-	-	-	-	-	2
White, both sexes	76	-	1	-	-	1	1	-	2	2	4	6
Male	33	-	-	-	-	1	1	-	2	2	4	4
Female	43	-	1	-	-	-	-	-	-	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	150	-	-	2	2	1	1	4	2	5	13	13
Male	47	-	-	1	-	-	-	4	-	-	6	3
Female	103	-	-	1	2	1	1	-	2	5	7	10
White, both sexes	111	-	-	2	2	-	-	1	-	-	9	9
Male	33	-	-	1	-	-	-	1	-	-	3	2
Female	78	-	-	1	2	-	-	-	-	-	6	7
Black, both sexes	36	-	-	-	-	1	1	3	2	5	4	4
Male	12	-	-	-	-	-	-	3	-	-	3	1
Female	24	-	-	-	-	1	1	-	2	5	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Rhode Island												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	63	-	-	1	1	-	1	1	-	2	4	3
Male	27	-	-	1	1	-	-	1	-	1	2	2
Female	36	-	-	-	-	-	1	-	-	1	2	1
White, both sexes	32	-	-	-	-	-	-	1	-	-	1	1
Male	11	-	-	-	-	-	-	1	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	31	-	-	1	1	-	1	-	-	2	3	2
Male	16	-	-	1	1	-	-	-	-	1	2	1
Female	15	-	-	-	-	-	1	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
South Dakota												
All races, both sexes	9	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	80	-	1	3	3	2	5	2	1	1	5	10
Male	36	-	-	3	3	1	4	1	1	1	3	3
Female	44	-	1	-	-	1	1	1	-	-	2	7
White, both sexes	49	-	1	1	1	-	2	2	-	-	2	7
Male	17	-	-	1	1	-	2	1	-	-	1	1
Female	32	-	1	-	-	-	-	1	-	-	1	6
Black, both sexes	31	-	-	2	2	2	3	-	1	1	3	3
Male	19	-	-	2	2	1	2	-	1	1	2	2
Female	12	-	-	-	-	1	1	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Texas												
All races, both sexes	224	1	-	5	1	3	7	3	7	4	16	17
Male	77	1	-	3	1	2	5	-	3	1	7	4
Female	147	-	-	2	-	1	2	3	4	3	9	13
White, both sexes	168	1	-	3	-	1	1	2	7	4	13	11
Male	55	1	-	2	-	-	-	-	3	1	4	3
Female	113	-	-	1	-	1	1	2	4	3	9	8
Black, both sexes	49	-	-	2	1	2	3	1	-	-	3	4
Male	19	-	-	1	1	2	2	-	-	-	3	1
Female	30	-	-	1	-	-	1	1	-	-	-	3
American Indian 1/, both sexes	2	-	-	-	-	-	1	-	-	-	-	1
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	2	-	-	-	-	1
Male	2	-	-	-	-	-	2	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	38	1	1	-	-	1	-	-	-	-	2	1
Male	12	-	1	-	-	-	-	-	-	-	1	-
Female	26	1	-	-	-	1	-	-	-	-	1	1
White, both sexes	36	1	1	-	-	1	-	-	-	-	2	1
Male	11	-	1	-	-	-	-	-	-	-	1	-
Female	25	1	-	-	-	1	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Vermont												
All races, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	78	-	-	2	-	1	1	1	-	2	1	13
Male	37	-	-	1	-	1	1	1	-	1	1	7
Female	41	-	-	1	-	-	-	-	-	1	-	6
White, both sexes	48	-	-	-	-	-	1	1	-	-	1	10
Male	21	-	-	-	-	-	1	1	-	-	1	5
Female	27	-	-	-	-	-	-	-	-	-	-	5
Black, both sexes	28	-	-	2	-	1	-	-	-	2	-	3
Male	15	-	-	1	-	1	-	-	-	1	-	2
Female	13	-	-	1	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Washington												
All races, both sexes	79	-	-	1	-	-	-	1	-	2	2	-
Male	26	-	-	-	-	-	-	1	-	1	-	-
Female	53	-	-	1	-	-	-	-	-	1	2	-
White, both sexes	68	-	-	-	-	-	-	1	-	1	2	-
Male	20	-	-	-	-	-	-	1	-	-	-	-
Female	48	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	3	-	-	1	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	28	-	-	-	-	1	1	1	1	1	3	-
Male	7	-	-	-	-	-	-	1	-	-	2	-
Female	21	-	-	-	-	1	1	-	1	1	1	-
White, both sexes	26	-	-	-	-	1	1	1	1	-	3	-
Male	7	-	-	-	-	-	-	1	-	-	2	-
Female	19	-	-	-	-	1	1	-	1	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Wisconsin												
All races, both sexes	65	-	-	-	1	1	1	1	3	-	3	2
Male	21	-	-	-	1	1	1	1	1	-	1	-
Female	44	-	-	-	-	-	-	-	2	-	2	2
White, both sexes	56	-	-	-	1	-	-	-	2	-	2	2
Male	14	-	-	-	1	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	2	-	2	2
Black, both sexes	8	-	-	-	-	1	1	1	1	-	1	-
Male	7	-	-	-	-	1	1	1	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	11	-	-	-	-	-	-	1	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	11	-	-	-	-	-	-	1	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
United States												
All races, both sexes	289	250	242	210	233	266	360	318	239	85	11	-
Male	115	89	79	66	80	89	96	81	52	18	2	-
Female	174	161	163	144	153	177	264	237	187	67	9	-
White, both sexes	173	156	155	150	170	220	305	279	218	72	8	-
Male	60	49	46	44	55	76	77	72	47	18	1	-
Female	113	107	109	106	115	144	228	207	171	54	7	-
Black, both sexes	102	85	79	49	50	30	31	27	10	9	2	-
Male	46	36	32	17	18	8	8	5	2	-	-	-
Female	56	49	47	32	32	22	23	22	8	9	2	-
American Indian 1/, both sexes	1	1	1	2	2	-	2	1	-	-	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	1	1	1	1	2	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	8	7	9	11	16	22	11	11	4	1	-
Male	9	4	1	4	7	5	9	4	3	-	1	-
Female	4	4	6	5	4	11	13	7	8	4	-	-
Alabama												
All races, both sexes	4	3	5	5	4	6	7	4	2	-	-	-
Male	2	2	2	2	1	3	2	2	1	-	-	-
Female	2	1	3	3	3	3	5	2	1	-	-	-
White, both sexes	3	1	1	3	3	5	6	3	2	-	-	-
Male	2	1	-	-	-	2	2	1	1	-	-	-
Female	1	-	1	3	3	3	4	2	1	-	-	-
Black, both sexes	1	2	4	2	1	1	1	1	-	-	-	-
Male	-	1	2	2	1	1	-	1	-	-	-	-
Female	1	1	2	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Alaska												
All races, both sexes	-	4	1	1	1	-	2	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	3	1	1	1	-	2	-	-	-	-	-
White, both sexes	-	3	-	1	1	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	1	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Arizona												
All races, both sexes	5	5	5	9	6	4	7	9	2	2	-	-
Male	1	2	3	4	3	1	3	3	-	1	-	-
Female	4	3	2	5	3	3	4	6	2	1	-	-
White, both sexes	5	5	5	8	6	4	7	8	2	2	-	-
Male	1	2	3	4	3	1	3	3	-	1	-	-
Female	4	3	2	4	3	3	4	5	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Arkansas												
All races, both sexes	7	3	2	1	5	6	5	4	6	1	-	-
Male	1	2	-	1	1	4	-	1	1	-	-	-
Female	6	1	2	-	4	2	5	3	5	1	-	-
White, both sexes	6	1	1	1	5	6	3	4	4	-	-	-
Male	1	1	-	1	1	4	-	1	-	-	-	-
Female	5	-	1	-	4	2	3	3	4	-	-	-
Black, both sexes	1	2	1	-	-	-	2	-	2	1	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	1	1	1	-	-	-	2	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	33	33	34	25	21	44	51	44	33	13	3	-
Male	13	12	7	8	8	21	13	11	7	4	1	-
Female	20	21	27	17	13	23	38	33	26	9	2	-
White, both sexes	20	22	18	20	14	32	36	34	27	11	2	-
Male	10	7	3	8	5	16	9	8	4	4	-	-
Female	10	15	15	12	9	16	27	26	23	7	2	-
Black, both sexes	9	7	12	2	3	1	5	2	-	1	-	-
Male	1	4	3	-	-	1	2	-	-	-	-	-
Female	8	3	9	2	3	-	3	2	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	4	4	3	4	11	10	8	6	1	1	-
Male	2	1	1	-	3	4	2	3	3	-	1	-
Female	1	3	3	3	1	7	8	5	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Colorado												
All races, both sexes	5	3	1	2	2	2	8	4	5	-	-	-
Male	2	1	-	2	1	-	1	1	-	-	-	-
Female	3	2	1	-	1	2	7	3	5	-	-	-
White, both sexes	4	2	1	2	2	2	8	4	4	-	-	-
Male	1	1	-	2	1	-	1	1	-	-	-	-
Female	3	1	1	-	1	2	7	3	4	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Connecticut												
All races, both sexes	5	2	6	3	3	2	2	3	5	2	-	-
Male	3	-	1	1	-	-	1	1	-	-	-	-
Female	2	2	5	2	3	2	1	2	5	2	-	-
White, both sexes	3	2	6	2	2	2	2	2	5	2	-	-
Male	1	-	1	-	-	-	1	1	-	-	-	-
Female	2	2	5	2	2	2	1	1	5	2	-	-
Black, both sexes	1	-	-	1	1	-	-	1	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Delaware												
All races, both sexes	-	1	1	-	-	1	-	-	1	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	-	1	-	-	-	1	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	2	1	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	2	1	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Florida												
All races, both sexes	10	15	7	8	19	8	18	18	4	4	-	-
Male	4	3	3	1	8	2	6	6	1	-	-	-
Female	6	12	4	7	11	6	12	12	3	4	-	-
White, both sexes	6	9	4	5	13	7	15	14	4	4	-	-
Male	3	2	2	1	4	2	5	4	1	-	-	-
Female	3	7	2	4	9	5	10	10	3	4	-	-
Black, both sexes	3	6	3	3	6	1	2	4	-	-	-	-
Male	1	1	1	-	4	-	-	2	-	-	-	-
Female	2	5	2	3	2	1	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	9	12	6	7	8	5	13	5	5	1	-	-
Male	2	6	-	4	2	2	2	1	-	1	-	-
Female	7	6	6	3	6	3	11	4	5	-	-	-
White, both sexes	5	5	4	4	2	3	9	3	4	1	-	-
Male	1	2	-	2	-	2	1	1	-	1	-	-
Female	4	3	4	2	2	1	8	2	4	-	-	-
Black, both sexes	4	7	2	2	6	2	3	1	-	-	-	-
Male	1	4	-	1	2	-	-	-	-	-	-	-
Female	3	3	2	1	4	2	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	1	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Hawaii												
All races, both sexes	5	2	1	5	2	-	2	-	1	-	-	-
Male	3	2	-	3	-	-	2	-	-	-	-	-
Female	2	-	1	2	2	-	-	-	1	-	-	-
White, both sexes	1	-	-	3	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	2	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	1	2	2	-	2	-	1	-	-	-
Male	3	2	-	2	-	-	2	-	-	-	-	-
Female	1	-	1	-	2	-	-	-	1	-	-	-
Idaho												
All races, both sexes	-	3	2	-	-	2	-	3	2	3	-	-
Male	-	1	2	-	-	2	-	-	-	1	-	-
Female	-	2	-	-	-	-	-	3	2	2	-	-
White, both sexes	-	3	2	-	-	2	-	3	2	3	-	-
Male	-	1	2	-	-	2	-	-	-	1	-	-
Female	-	2	-	-	-	-	-	3	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Illinois												
All races, both sexes	26	8	8	7	5	13	5	11	11	3	-	-
Male	9	3	1	2	3	3	2	5	2	2	-	-
Female	17	5	7	5	2	10	3	6	9	1	-	-
White, both sexes	10	2	5	4	2	9	5	9	9	3	-	-
Male	2	1	-	1	1	2	2	5	2	2	-	-
Female	8	1	5	3	1	7	3	4	7	1	-	-
Black, both sexes	15	6	3	3	3	3	-	2	2	-	-	-
Male	7	2	1	1	2	1	-	-	-	-	-	-
Female	8	4	2	2	1	2	-	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Indiana												
All races, both sexes	3	7	3	5	2	7	6	5	5	3	-	-
Male	-	2	1	-	-	2	1	1	-	-	-	-
Female	3	5	2	5	2	5	5	4	5	3	-	-
White, both sexes	1	5	3	4	2	6	5	5	5	3	-	-
Male	-	2	1	-	-	2	1	1	-	-	-	-
Female	1	3	2	4	2	4	4	4	5	3	-	-
Black, both sexes	2	1	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Iowa												
All races, both sexes	3	1	1	-	1	3	9	4	4	-	-	-
Male	2	-	-	-	-	1	1	1	1	-	-	-
Female	1	1	1	-	1	2	8	3	3	-	-	-
White, both sexes	2	1	1	-	1	3	9	4	4	-	-	-
Male	1	-	-	-	-	1	1	1	1	-	-	-
Female	1	1	1	-	1	2	8	3	3	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	-	2	3	2	2	-	5	2	2	-	-
Male	1	-	1	1	-	-	-	1	1	-	-	-
Female	2	-	1	2	2	2	-	4	1	2	-	-
White, both sexes	3	-	2	3	2	1	-	5	2	1	-	-
Male	1	-	1	1	-	-	-	1	1	-	-	-
Female	2	-	1	2	2	1	-	4	1	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Kentucky												
All races, both sexes	7	2	2	6	5	3	5	6	3	2	-	-
Male	2	1	-	1	1	-	3	1	1	-	-	-
Female	5	1	2	5	4	3	2	5	2	2	-	-
White, both sexes	6	2	1	5	4	3	5	6	3	2	-	-
Male	2	1	-	1	1	-	3	1	1	-	-	-
Female	4	1	1	4	3	3	2	5	2	2	-	-
Black, both sexes	1	-	1	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	5	4	3	1	5	3	3	1	-	-	-
Male	-	3	-	1	-	3	1	-	-	-	-	-
Female	4	2	4	2	1	2	2	3	1	-	-	-
White, both sexes	3	1	2	1	1	5	2	1	1	-	-	-
Male	-	-	-	-	-	3	1	-	-	-	-	-
Female	3	1	2	1	1	2	1	1	1	-	-	-
Black, both sexes	1	4	2	2	-	-	1	2	-	-	-	-
Male	-	3	-	1	-	-	-	-	-	-	-	-
Female	1	1	2	1	-	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Maine												
All races, both sexes	-	-	-	2	3	-	-	3	3	-	-	-
Male	-	-	-	-	-	-	-	1	2	-	-	-
Female	-	-	-	2	3	-	-	2	1	-	-	-
White, both sexes	-	-	-	2	3	-	-	3	3	-	-	-
Male	-	-	-	-	-	-	-	1	2	-	-	-
Female	-	-	-	2	3	-	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5	3	3	5	4	1	2	6	4	1	-	-
Male	3	-	2	1	2	1	2	2	2	-	-	-
Female	2	3	1	4	2	-	-	4	2	1	-	-
White, both sexes	2	2	1	1	3	1	-	5	3	1	-	-
Male	-	-	1	-	2	1	-	2	2	-	-	-
Female	2	2	-	1	1	-	-	3	1	1	-	-
Black, both sexes	3	1	2	4	1	-	2	1	1	-	-	-
Male	3	-	1	1	-	-	2	-	-	-	-	-
Female	-	1	1	3	1	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Massachusetts												
All races, both sexes	1	2	3	3	2	5	7	5	3	5	1	-
Male	1	-	-	1	2	-	1	-	1	-	-	-
Female	-	2	3	2	-	5	6	5	2	5	1	-
White, both sexes	-	2	3	3	1	5	6	4	3	4	1	-
Male	-	-	-	1	1	-	1	-	1	-	-	-
Female	-	2	3	2	-	5	5	4	2	4	1	-
Black, both sexes	1	-	-	-	1	-	1	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Michigan												
All races, both sexes	5	2	13	5	9	14	13	9	3	1	-	-
Male	3	1	3	-	4	3	5	1	-	-	-	-
Female	2	1	10	5	5	11	8	8	3	1	-	-
White, both sexes	5	-	6	4	7	13	12	9	3	-	-	-
Male	3	-	2	-	2	3	4	1	-	-	-	-
Female	2	-	4	4	5	10	8	8	3	-	-	-
Black, both sexes	-	2	7	1	2	1	1	-	-	1	-	-
Male	-	1	1	-	2	-	1	-	-	-	-	-
Female	-	1	6	1	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Minnesota												
All races, both sexes	3	5	2	2	1	2	10	3	6	1	1	-
Male	-	2	1	-	-	1	1	1	2	-	-	-
Female	3	3	1	2	1	1	9	2	4	1	1	-
White, both sexes	3	4	2	1	1	2	8	3	5	-	1	-
Male	-	1	1	-	-	1	1	1	2	-	-	-
Female	3	3	1	1	1	1	7	2	3	-	1	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	1	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
Mississippi												
All races, both sexes	3	3	4	1	1	4	4	4	2	2	-	-
Male	-	2	2	-	1	-	2	1	-	1	-	-
Female	3	1	2	1	-	4	2	3	2	1	-	-
White, both sexes	1	-	2	-	1	2	3	2	2	1	-	-
Male	-	-	1	-	1	-	2	-	-	1	-	-
Female	1	-	1	-	-	2	1	2	2	-	-	-
Black, both sexes	2	3	2	1	-	2	1	2	-	1	-	-
Male	-	2	1	-	-	-	-	1	-	-	-	-
Female	2	1	1	1	-	2	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Missouri												
All races, both sexes	4	3	8	5	5	6	2	2	4	2	-	-
Male	3	-	3	1	-	1	-	-	1	-	-	-
Female	1	3	5	4	5	5	2	2	3	2	-	-
White, both sexes	3	3	7	4	4	5	2	2	4	2	-	-
Male	2	-	2	1	-	1	-	-	1	-	-	-
Female	1	3	5	3	4	4	2	2	3	2	-	-
Black, both sexes	1	-	1	1	1	1	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	1	2	1	1	1	2	3	2	1	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	2	1	1	-	2	2	2	1	-	-
White, both sexes	-	1	1	1	1	1	2	3	2	1	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	1	1	1	-	2	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Nebraska												
All races, both sexes	2	1	-	1	1	-	4	4	3	3	-	-
Male	1	-	-	1	1	-	-	1	1	-	-	-
Female	1	1	-	-	-	-	4	3	2	3	-	-
White, both sexes	1	1	-	1	1	-	4	4	3	3	-	-
Male	-	-	-	1	1	-	-	1	1	-	-	-
Female	1	1	-	-	-	-	4	3	2	3	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	4	1	2	-	-	2	1	1	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	3	1	2	-	-	1	-	1	-	-	-
White, both sexes	-	3	1	1	-	-	1	1	1	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	2	1	1	-	-	-	-	1	-	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
New Hampshire												
All races, both sexes	2	-	-	-	2	1	1	1	1	-	-	-
Male	1	-	-	-	1	1	1	-	-	-	-	-
Female	1	-	-	-	1	-	-	1	1	-	-	-
White, both sexes	2	-	-	-	2	1	1	1	1	-	-	-
Male	1	-	-	-	1	1	1	-	-	-	-	-
Female	1	-	-	-	1	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	13	5	6	7	10	7	7	8	3	2	-	-
Male	7	2	2	1	4	1	1	4	-	-	-	-
Female	6	3	4	6	6	6	6	4	3	2	-	-
White, both sexes	9	3	3	4	5	5	7	8	3	1	-	-
Male	5	1	-	-	2	1	1	4	-	-	-	-
Female	4	2	3	4	3	4	6	4	3	1	-	-
Black, both sexes	4	2	3	3	3	1	-	-	-	1	-	-
Male	2	1	2	1	1	-	-	-	-	-	-	-
Female	2	1	1	2	2	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
New Mexico												
All races, both sexes	2	1	2	2	-	-	7	3	-	-	-	-
Male	1	-	-	1	-	-	3	2	-	-	-	-
Female	1	1	2	1	-	-	4	1	-	-	-	-
White, both sexes	2	1	2	2	-	-	6	3	-	-	-	-
Male	1	-	-	1	-	-	2	2	-	-	-	-
Female	1	1	2	1	-	-	4	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	27	19	21	20	10	14	18	22	11	6	-	-
Male	12	7	9	4	2	3	8	5	1	-	-	-
Female	15	12	12	16	8	11	10	17	10	6	-	-
White, both sexes	14	9	14	14	7	12	15	17	11	4	-	-
Male	5	3	5	2	2	2	6	3	1	-	-	-
Female	9	6	9	12	5	10	9	14	10	4	-	-
Black, both sexes	12	10	7	5	2	2	3	4	-	2	-	-
Male	6	4	4	1	-	1	2	1	-	-	-	-
Female	6	6	3	4	2	1	1	3	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	-	-	1	-	-	-	-
Male	1	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
North Carolina												
All races, both sexes	4	5	8	8	17	5	14	8	8	2	-	-
Male	1	3	3	3	8	1	4	2	2	1	-	-
Female	3	2	5	5	9	4	10	6	6	1	-	-
White, both sexes	-	-	3	6	13	3	10	7	7	2	-	-
Male	-	-	1	2	7	-	2	2	2	1	-	-
Female	-	-	2	4	6	3	8	5	5	1	-	-
Black, both sexes	4	5	5	2	4	2	3	1	1	-	-	-
Male	1	3	2	1	1	1	1	-	-	-	-	-
Female	3	2	3	1	3	1	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	3	2	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	2	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	3	2	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Ohio												
All races, both sexes	12	14	12	6	9	15	12	12	15	3	1	-
Male	7	3	4	2	3	5	1	4	2	1	-	-
Female	5	11	8	4	6	10	11	8	13	2	1	-
White, both sexes	9	10	7	5	8	14	12	10	13	3	1	-
Male	5	2	1	1	3	4	1	4	2	1	-	-
Female	4	8	6	4	5	10	11	6	11	2	1	-
Black, both sexes	3	4	5	1	1	1	-	2	2	-	-	-
Male	2	1	3	1	-	1	-	-	-	-	-	-
Female	1	3	2	-	1	-	-	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	2	2	4	4	3	4	3	2	-	-	-
Male	1	1	1	2	-	2	2	-	-	-	-	-
Female	2	1	1	2	4	1	2	3	2	-	-	-
White, both sexes	2	1	2	3	3	3	3	3	2	-	-	-
Male	-	1	1	2	-	2	2	-	-	-	-	-
Female	2	-	1	1	3	1	1	3	2	-	-	-
Black, both sexes	1	1	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Oregon												
All races, both sexes	5	6	4	3	5	5	14	12	6	2	-	-
Male	1	1	2	2	3	1	5	3	3	-	-	-
Female	4	5	2	1	2	4	9	9	3	2	-	-
White, both sexes	4	6	3	2	5	5	14	12	6	2	-	-
Male	-	1	2	1	3	1	5	3	3	-	-	-
Female	4	5	1	1	2	4	9	9	3	2	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	16	10	13	9	8	10	16	10	12	2	1	-
Male	4	6	7	2	2	5	2	1	3	1	-	-
Female	12	4	6	7	6	5	14	9	9	1	1	-
White, both sexes	11	7	9	7	6	9	15	10	11	2	1	-
Male	3	5	3	2	2	5	1	1	3	1	-	-
Female	8	2	6	5	4	4	14	9	8	1	1	-
Black, both sexes	4	3	4	1	2	1	-	-	1	-	-	-
Male	-	1	4	-	-	-	-	-	-	-	-	-
Female	4	2	-	1	2	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Rhode Island												
All races, both sexes	-	-	1	-	-	-	-	3	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	3	1	-	-	-
White, both sexes	-	-	1	-	-	-	-	3	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	12	2	6	4	9	4	7	3	1	1	1	-
Male	5	2	2	2	3	2	3	-	-	-	-	-
Female	7	-	4	2	6	2	4	3	1	1	1	-
White, both sexes	4	-	3	2	6	2	6	3	1	1	1	-
Male	1	-	1	-	2	2	3	-	-	-	-	-
Female	3	-	2	2	4	-	3	3	1	1	1	-
Black, both sexes	8	2	3	2	3	2	1	-	-	-	-	-
Male	4	2	1	2	1	-	-	-	-	-	-	-
Female	4	-	2	-	2	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
South Dakota												
All races, both sexes	-	1	3	1	-	1	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	2	1	-	1	-	-	1	-	-	-
White, both sexes	-	1	3	1	-	1	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	2	1	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	8	5	3	2	8	7	1	7	4	1	1	-
Male	2	2	1	1	3	2	-	2	2	1	-	-
Female	6	3	2	1	5	5	1	5	2	-	1	-
White, both sexes	4	2	1	1	6	6	1	7	4	1	-	-
Male	1	1	-	-	1	2	-	2	2	1	-	-
Female	3	1	1	1	5	4	1	5	2	-	-	-
Black, both sexes	4	3	2	1	2	1	-	-	-	-	1	-
Male	1	1	1	1	2	-	-	-	-	-	-	-
Female	3	2	1	-	-	1	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Texas												
All races, both sexes	10	20	15	17	18	18	21	22	14	4	1	-
Male	6	6	7	7	8	3	2	8	2	1	-	-
Female	4	14	8	10	10	15	19	14	12	3	1	-
White, both sexes	6	14	11	11	13	12	21	20	14	3	-	-
Male	4	4	6	5	7	2	2	8	2	1	-	-
Female	2	10	5	6	6	10	19	12	12	2	-	-
Black, both sexes	4	6	4	5	5	5	-	2	-	1	1	-
Male	2	2	1	2	1	1	-	-	-	-	-	-
Female	2	4	3	3	4	4	-	2	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Utah												
All races, both sexes	-	4	2	-	1	5	7	6	6	1	-	-
Male	-	1	1	-	-	3	2	-	3	-	-	-
Female	-	3	1	-	1	2	5	6	3	1	-	-
White, both sexes	-	4	2	-	1	5	5	6	6	1	-	-
Male	-	1	1	-	-	3	1	-	3	-	-	-
Female	-	3	1	-	1	2	4	6	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Vermont												
All races, both sexes	2	-	-	-	1	1	1	-	1	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	1	-	1	1	-	-
White, both sexes	2	-	-	-	1	1	1	-	1	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	7	7	3	3	5	6	10	9	7	-	-	-
Male	5	3	1	2	-	3	3	3	4	-	-	-
Female	2	4	2	1	5	3	7	6	3	-	-	-
White, both sexes	-	3	1	2	4	4	7	8	6	-	-	-
Male	-	2	-	1	-	1	3	3	3	-	-	-
Female	-	1	1	1	4	3	4	5	3	-	-	-
Black, both sexes	7	4	2	1	1	1	2	1	1	-	-	-
Male	5	1	1	1	-	1	-	-	1	-	-	-
Female	2	3	1	-	1	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Washington												
All races, both sexes	9	5	8	4	8	8	12	5	10	3	1	-
Male	2	1	3	1	3	4	4	1	2	2	1	-
Female	7	4	5	3	5	4	8	4	8	1	-	-
White, both sexes	8	5	7	4	5	7	11	4	9	3	1	-
Male	1	1	3	1	-	4	3	1	2	2	1	-
Female	7	4	4	3	5	3	8	3	7	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	3	1	1	1	1	-	-	-
Male	-	-	-	-	3	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	-	1	1	-	-	-
West Virginia												
All races, both sexes	2	2	2	1	1	1	4	3	2	2	-	-
Male	2	-	-	-	-	-	1	-	-	1	-	-
Female	-	2	2	1	1	1	3	3	2	1	-	-
White, both sexes	2	2	2	-	1	1	4	3	2	2	-	-
Male	2	-	-	-	-	-	1	-	-	1	-	-
Female	-	2	2	-	1	1	3	3	2	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Wisconsin												
All races, both sexes	2	5	3	2	3	9	11	8	8	2	-	-
Male	1	1	2	1	1	2	4	1	2	-	-	-
Female	1	4	1	1	2	7	7	7	6	2	-	-
White, both sexes	1	5	2	2	2	9	11	8	8	1	-	-
Male	-	1	1	1	1	2	4	1	2	-	-	-
Female	1	4	1	1	1	7	7	7	6	1	-	-
Black, both sexes	1	-	1	-	1	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Wyoming												
All races, both sexes	-	2	-	-	-	-	4	2	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	4	2	-	-	-	-
White, both sexes	-	2	-	-	-	-	4	2	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	4	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
United States												
All races, both sexes	973	-	-	-	-	-	3	1	2	1	6	11
Male	284	-	-	-	-	-	3	-	1	-	3	7
Female	689	-	-	-	-	-	-	1	1	1	3	4
White, both sexes	891	-	-	-	-	-	3	1	2	1	3	8
Male	244	-	-	-	-	-	3	-	1	-	1	6
Female	647	-	-	-	-	-	-	1	1	1	2	2
Black, both sexes	35	-	-	-	-	-	-	-	-	-	3	-
Male	22	-	-	-	-	-	-	-	-	-	2	-
Female	13	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	42	-	-	-	-	-	-	-	-	-	-	3
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	2
Alabama												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Arkansas												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	120	-	-	-	-	-	1	-	-	-	-	2
Male	32	-	-	-	-	-	1	-	-	-	-	2
Female	88	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	102	-	-	-	-	-	1	-	-	-	-	2
Male	25	-	-	-	-	-	1	-	-	-	-	2
Female	77	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Colorado												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Florida												
All races, both sexes	71	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Hawaii												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Illinois												
All races, both sexes	40	-	-	-	-	-	-	-	1	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	35	-	-	-	-	-	-	-	1	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Indiana												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Iowa												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Kentucky												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Maine												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Massachusetts												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	19	-	-	-	-	-	1	-	-	-	1	-
Male	9	-	-	-	-	-	1	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	1	-	-	-	-	-
Male	8	-	-	-	-	-	1	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Minnesota												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Missouri												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Nebraska												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
New Hampshire												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
New Mexico												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
North Carolina												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	2	1
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	31	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Ohio												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Oregon												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	49	-	-	-	-	-	-	-	1	-	-	-
Male	18	-	-	-	-	-	-	-	1	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	1	-	-	-
Male	16	-	-	-	-	-	-	-	1	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Rhode Island												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	16	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
South Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Texas												
All races, both sexes	69	-	-	-	-	-	1	-	-	-	-	1
Male	21	-	-	-	-	-	1	-	-	-	-	1
Female	48	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	63	-	-	-	-	-	1	-	-	-	-	1
Male	18	-	-	-	-	-	1	-	-	-	-	1
Female	45	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Vermont												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Washington												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Wisconsin												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
United States												
All races, both sexes	21	23	37	55	98	140	232	202	113	24	4	-
Male	14	11	16	17	30	50	66	49	11	5	1	-
Female	7	12	21	38	68	90	166	153	102	19	3	-
White, both sexes	16	19	33	51	86	128	218	191	105	23	3	-
Male	11	8	13	14	26	43	61	43	9	5	-	-
Female	5	11	20	37	60	85	157	148	96	18	3	-
Black, both sexes	2	3	3	1	6	5	5	3	2	1	1	-
Male	1	3	3	1	3	4	2	2	-	-	1	-
Female	1	-	-	-	3	1	3	1	2	1	-	-
American Indian 1/, both sexes	2	-	1	1	1	-	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	2	5	7	9	8	6	-	-	-
Male	-	-	-	1	1	3	3	4	2	-	-	-
Female	1	1	-	1	4	4	6	4	4	-	-	-
Alabama												
All races, both sexes	-	1	2	-	1	-	3	4	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	1	1	-	1	-	2	4	-	1	-	-
White, both sexes	-	1	2	-	1	-	3	4	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	1	1	-	1	-	2	4	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	1	4	5	3	6	5	3	-	-	-
Male	-	-	1	2	3	1	4	2	1	-	-	-
Female	1	-	-	2	2	2	2	3	2	-	-	-
White, both sexes	1	-	1	3	5	3	6	5	3	-	-	-
Male	-	-	1	1	3	1	4	2	1	-	-	-
Female	1	-	-	2	2	2	2	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Arkansas												
All races, both sexes	-	-	-	1	2	2	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	2	2	1	1	-	-	-	-
White, both sexes	-	-	-	1	2	2	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	2	2	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	2	7	7	10	15	33	25	13	2	2	-
Male	1	-	1	1	4	6	10	2	3	1	-	-
Female	-	2	6	6	6	9	23	23	10	1	2	-
White, both sexes	1	1	7	6	8	12	28	22	10	2	2	-
Male	1	-	1	1	3	4	9	1	1	1	-	-
Female	-	1	6	5	5	8	19	21	9	1	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	2	3	5	3	3	-	-	-
Male	-	-	-	-	1	2	1	1	2	-	-	-
Female	-	1	-	1	1	1	4	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Colorado												
All races, both sexes	-	-	-	1	1	1	3	2	4	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	1	1	1	2	2	3	-	-	-
White, both sexes	-	-	-	1	1	1	2	2	4	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	1	1	1	1	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	1	-	-	5	2	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	4	2	1	-	-
White, both sexes	-	-	-	-	-	-	-	5	2	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	4	2	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Florida												
All races, both sexes	1	3	2	3	5	6	15	22	12	2	-	-
Male	-	2	1	1	-	4	2	10	2	-	-	-
Female	1	1	1	2	5	2	13	12	10	2	-	-
White, both sexes	1	3	2	3	5	6	15	21	12	2	-	-
Male	-	2	1	1	-	4	2	9	2	-	-	-
Female	1	1	1	2	5	2	13	12	10	2	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	1	1	1	3	4	7	7	1	-	-	-
Male	-	1	-	1	-	3	-	1	-	-	-	-
Female	-	-	1	-	3	1	7	6	1	-	-	-
White, both sexes	-	1	1	1	3	4	7	7	1	-	-	-
Male	-	1	-	1	-	3	-	1	-	-	-	-
Female	-	-	1	-	3	1	7	6	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Hawaii												
All races, both sexes	-	-	-	-	2	3	1	3	3	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	2	2	1	2	3	-	-	-
White, both sexes	-	-	-	-	-	2	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	1	3	2	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	2	1	1	2	2	-	-	-
Idaho												
All races, both sexes	-	-	-	1	1	1	-	2	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-
White, both sexes	-	-	-	1	1	1	-	2	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Illinois												
All races, both sexes	1	1	1	2	3	5	10	9	5	1	-	-
Male	1	1	-	-	1	2	6	4	1	1	-	-
Female	-	-	1	2	2	3	4	5	4	-	-	-
White, both sexes	1	1	1	2	3	4	8	9	4	1	-	-
Male	1	1	-	-	1	1	5	4	1	1	-	-
Female	-	-	1	2	2	3	3	5	3	-	-	-
Black, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Indiana												
All races, both sexes	1	1	-	-	3	2	6	7	-	2	-	-
Male	-	-	-	-	1	1	3	1	-	-	-	-
Female	1	1	-	-	2	1	3	6	-	2	-	-
White, both sexes	1	1	-	-	2	2	6	7	-	2	-	-
Male	-	-	-	-	-	1	3	1	-	-	-	-
Female	1	1	-	-	2	1	3	6	-	2	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Iowa												
All races, both sexes	-	1	-	1	3	2	3	2	3	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	2	2	2	2	3	-	-	-
White, both sexes	-	1	-	1	3	2	3	2	3	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	2	2	2	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	1	-	-	2	3	4	5	4	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	1	-	-	2	2	3	5	4	-	-	-
White, both sexes	-	1	-	-	2	3	4	5	4	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	1	-	-	2	2	3	5	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Kentucky												
All races, both sexes	-	-	1	-	6	2	2	4	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	5	2	1	4	-	-	-	-
White, both sexes	-	-	1	-	6	2	2	4	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	5	2	1	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	1	2	2	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	2	1	-	-	-	-	-
White, both sexes	-	-	-	1	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Maine												
All races, both sexes	-	-	-	1	1	1	3	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	3	-	1	-	-	-
White, both sexes	-	-	-	1	1	1	3	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	3	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	1	1	2	-	4	3	2	-	1	-	-
Male	1	-	-	-	-	1	1	1	-	1	-	-
Female	-	1	1	2	-	3	2	1	-	-	-	-
White, both sexes	1	1	1	2	-	3	3	2	-	1	-	-
Male	1	-	-	-	-	-	1	1	-	1	-	-
Female	-	1	1	2	-	3	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Massachusetts												
All races, both sexes	-	1	2	3	-	2	4	10	2	2	-	-
Male	-	1	2	2	-	1	2	2	-	-	-	-
Female	-	-	-	1	-	1	2	8	2	2	-	-
White, both sexes	-	-	2	3	-	2	4	10	1	1	-	-
Male	-	-	2	2	-	1	2	2	-	-	-	-
Female	-	-	-	1	-	1	2	8	1	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	1	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	-	-	-	1	5	3	4	2	-	-	-
Male	2	-	-	-	-	2	1	2	-	-	-	-
Female	-	-	-	-	1	3	2	2	2	-	-	-
White, both sexes	2	-	-	-	-	5	3	4	2	-	-	-
Male	2	-	-	-	-	2	1	2	-	-	-	-
Female	-	-	-	-	-	3	2	2	2	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Minnesota												
All races, both sexes	1	-	-	-	-	5	4	4	1	1	-	-
Male	1	-	-	-	-	4	1	-	-	-	-	-
Female	-	-	-	-	-	1	3	4	1	1	-	-
White, both sexes	1	-	-	-	-	5	4	4	1	1	-	-
Male	1	-	-	-	-	4	1	-	-	-	-	-
Female	-	-	-	-	-	1	3	4	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	1	1	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	1	-	3	-	-	-	-	-
White, both sexes	-	-	1	1	1	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	1	-	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Missouri												
All races, both sexes	-	1	-	1	4	3	8	3	3	1	-	-
Male	-	1	-	-	4	3	2	2	-	1	-	-
Female	-	-	-	1	-	-	6	1	3	-	-	-
White, both sexes	-	1	-	1	4	3	8	3	3	1	-	-
Male	-	1	-	-	4	3	2	2	-	1	-	-
Female	-	-	-	1	-	-	6	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Nebraska												
All races, both sexes	-	-	1	2	1	-	2	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	1	-	2	1	1	-	-	-
White, both sexes	-	-	1	2	1	-	2	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	1	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
New Hampshire												
All races, both sexes	-	-	-	-	1	1	3	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	3	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	3	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	1	1	1	3	2	5	4	3	-	1	-
Male	1	-	1	-	1	-	2	1	-	-	1	-
Female	-	1	-	1	2	2	3	3	3	-	-	-
White, both sexes	1	1	-	1	3	2	5	4	3	-	-	-
Male	1	-	-	-	1	-	2	1	-	-	-	-
Female	-	1	-	1	2	2	3	3	3	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	1	-
Male	-	-	1	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
New Mexico												
All races, both sexes	-	-	1	1	1	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	1	1	-	2	-	1	-	-	-
White, both sexes	-	-	-	1	-	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	1	-	4	2	5	10	9	4	-	-	-
Male	1	1	-	3	1	1	4	4	-	-	-	-
Female	2	-	-	1	1	4	6	5	4	-	-	-
White, both sexes	2	-	-	3	2	4	10	8	4	-	-	-
Male	1	-	-	2	1	-	4	3	-	-	-	-
Female	1	-	-	1	1	4	6	5	4	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
North Carolina												
All races, both sexes	1	2	2	2	1	5	7	6	4	1	-	-
Male	1	-	1	-	1	-	-	2	-	-	-	-
Female	-	2	1	2	-	5	7	4	4	1	-	-
White, both sexes	1	2	1	2	1	5	6	6	4	1	-	-
Male	1	-	-	-	1	-	-	2	-	-	-	-
Female	-	2	1	2	-	5	6	4	4	1	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Ohio												
All races, both sexes	1	-	2	1	3	7	6	5	3	-	-	-
Male	1	-	2	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	3	6	5	4	3	-	-	-
White, both sexes	1	-	2	1	3	5	6	5	3	-	-	-
Male	1	-	2	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	3	5	5	4	3	-	-	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	2	2	2	-	-	2	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	2	1	-	-	2	-	-
White, both sexes	-	-	-	-	2	2	2	-	-	2	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	2	1	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Oregon												
All races, both sexes	1	-	-	-	3	3	1	6	3	1	-	-
Male	-	-	-	-	1	3	1	2	-	-	-	-
Female	1	-	-	-	2	-	-	4	3	1	-	-
White, both sexes	1	-	-	-	3	3	1	6	3	1	-	-
Male	-	-	-	-	1	3	1	2	-	-	-	-
Female	1	-	-	-	2	-	-	4	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	-	3	2	1	9	9	9	11	2	-	-
Male	2	-	1	2	-	4	4	2	1	1	-	-
Female	-	-	2	-	1	5	5	7	10	1	-	-
White, both sexes	1	-	3	2	1	9	9	8	10	2	-	-
Male	1	-	1	2	-	4	4	1	1	1	-	-
Female	-	-	2	-	1	5	5	7	9	1	-	-
Black, both sexes	1	-	-	-	-	-	-	1	1	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Rhode Island												
All races, both sexes	-	1	-	1	1	2	2	2	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	1	-	1	1	1	1	2	1	-	-	-
White, both sexes	-	1	-	1	1	2	2	2	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	1	-	1	1	1	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	1	1	-	2	3	4	2	1	1	-	-
Male	-	1	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	3	4	1	1	1	-	-
White, both sexes	-	-	-	-	2	3	4	2	1	1	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	3	4	1	1	1	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
South Dakota												
All races, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	2	4	1	4	8	1	2	2	-	-
Male	-	-	-	-	1	2	1	-	1	-	-	-
Female	-	-	2	4	-	2	7	1	1	2	-	-
White, both sexes	-	-	2	4	1	4	8	1	2	2	-	-
Male	-	-	-	-	1	2	1	-	1	-	-	-
Female	-	-	2	4	-	2	7	1	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Texas												
All races, both sexes	-	1	2	5	9	10	18	16	5	1	-	-
Male	-	-	2	2	3	3	4	5	-	-	-	-
Female	-	1	-	3	6	7	14	11	5	1	-	-
White, both sexes	-	1	2	4	7	10	17	14	5	1	-	-
Male	-	-	2	1	3	3	3	4	-	-	-	-
Female	-	1	-	3	4	7	14	10	5	1	-	-
Black, both sexes	-	-	-	1	1	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Vermont												
All races, both sexes	-	-	1	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	1	-	-	-
White, both sexes	-	-	1	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	1	1	1	3	3	7	2	4	-	1	-
Male	-	1	1	-	1	1	2	-	-	-	-	-
Female	-	-	-	1	2	2	5	2	4	-	1	-
White, both sexes	-	1	1	1	2	2	7	2	4	-	1	-
Male	-	1	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	2	2	5	2	4	-	1	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Washington												
All races, both sexes	-	1	-	-	2	5	9	4	7	-	-	-
Male	-	1	-	-	-	1	3	-	1	-	-	-
Female	-	-	-	-	2	4	6	4	6	-	-	-
White, both sexes	-	1	-	-	2	4	8	4	7	-	-	-
Male	-	1	-	-	-	1	2	-	1	-	-	-
Female	-	-	-	-	2	3	6	4	6	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	1	2	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	2	1	-	-	-	-
White, both sexes	-	-	-	-	1	2	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Wisconsin												
All races, both sexes	-	-	-	-	1	1	6	4	2	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	1	-	4	4	2	-	-	-
White, both sexes	-	-	-	-	1	1	6	4	2	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	1	-	4	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	2	2	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	2	2	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
United States												
All races, both sexes	17,899	13	10	6	7	12	19	35	45	54	118	194
Male	9,854	7	4	4	7	4	11	22	32	35	71	113
Female	8,045	6	6	2	-	8	8	13	13	19	47	81
White, both sexes	16,085	8	7	5	5	7	12	24	35	42	90	156
Male	8,865	3	2	3	5	2	8	14	25	25	57	94
Female	7,220	5	5	2	-	5	4	10	10	17	33	62
Black, both sexes	1,493	5	-	1	2	3	7	11	9	10	27	34
Male	807	4	-	1	2	1	3	8	6	8	14	15
Female	686	1	-	-	-	2	4	3	3	2	13	19
American Indian 1/, both sexes	75	-	-	-	-	1	-	-	1	1	1	2
Male	44	-	-	-	-	-	-	-	1	1	-	2
Female	31	-	-	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	246	-	3	-	-	1	-	-	-	1	-	2
Male	138	-	2	-	-	1	-	-	-	1	-	2
Female	108	-	1	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	300	-	-	-	-	-	2	1	1	2	1	5
Male	163	-	-	-	-	-	2	-	1	2	-	4
Female	137	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	245	-	-	-	-	-	-	-	1	1	1	4
Male	134	-	-	-	-	-	-	-	1	1	-	3
Female	111	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	54	-	-	-	-	-	2	1	-	1	-	1
Male	29	-	-	-	-	-	2	-	-	1	-	1
Female	25	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Alaska												
All races, both sexes	13	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	238	-	-	-	-	-	-	-	2	1	-	4
Male	137	-	-	-	-	-	-	-	2	1	-	3
Female	101	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	226	-	-	-	-	-	-	-	2	1	-	4
Male	129	-	-	-	-	-	-	-	2	1	-	3
Female	97	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Arkansas												
All races, both sexes	203	-	-	-	-	-	-	-	-	2	1	2
Male	111	-	-	-	-	-	-	-	-	2	1	2
Female	92	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	180	-	-	-	-	-	-	-	-	2	1	1
Male	100	-	-	-	-	-	-	-	-	2	1	1
Female	80	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	935	1	1	-	1	1	-	2	1	2	2	11
Male	514	-	-	-	1	1	-	1	1	2	2	7
Female	421	1	1	-	-	-	-	1	-	-	-	4
White, both sexes	844	1	-	-	1	-	-	1	1	2	2	9
Male	468	-	-	-	1	-	-	-	1	2	2	6
Female	376	1	-	-	-	-	-	1	-	-	-	3
Black, both sexes	40	-	-	-	-	-	-	1	-	-	-	1
Male	22	-	-	-	-	-	-	1	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	-	1	-	-	1	-	-	-	-	-	1
Male	24	-	-	-	-	1	-	-	-	-	-	1
Female	25	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Colorado												
All races, both sexes	266	1	-	-	-	-	-	-	-	-	-	1
Male	149	-	-	-	-	-	-	-	-	-	-	1
Female	117	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	254	1	-	-	-	-	-	-	-	-	-	-
Male	141	-	-	-	-	-	-	-	-	-	-	-
Female	113	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	334	-	-	-	-	-	-	-	1	-	3	2
Male	168	-	-	-	-	-	-	-	1	-	2	2
Female	166	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	320	-	-	-	-	-	-	-	1	-	3	2
Male	161	-	-	-	-	-	-	-	1	-	2	2
Female	159	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Delaware												
All races, both sexes	58	-	-	-	-	-	-	-	-	1	1	-
Male	34	-	-	-	-	-	-	-	-	1	-	-
Female	24	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	51	-	-	-	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	27	-	-	-	1	-	-	-	-	-	-	-
Male	11	-	-	-	1	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	1	-	-	-	-	-	-	-
Male	7	-	-	-	1	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Florida												
All races, both sexes	967	-	-	-	-	-	3	3	1	3	8	12
Male	573	-	-	-	-	-	1	3	1	3	5	6
Female	394	-	-	-	-	-	2	-	-	-	3	6
White, both sexes	873	-	-	-	-	-	2	3	1	3	6	11
Male	512	-	-	-	-	-	1	3	1	3	3	6
Female	361	-	-	-	-	-	1	-	-	-	3	5
Black, both sexes	89	-	-	-	-	-	1	-	-	-	2	1
Male	57	-	-	-	-	-	-	-	-	-	2	-
Female	32	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	475	-	2	-	-	2	1	2	3	-	3	4
Male	272	-	1	-	-	1	1	2	-	-	1	3
Female	203	-	1	-	-	1	-	-	3	-	2	1
White, both sexes	375	-	1	-	-	1	1	2	2	-	2	4
Male	225	-	-	-	-	1	1	2	-	-	1	3
Female	150	-	1	-	-	-	-	-	2	-	1	1
Black, both sexes	97	-	-	-	-	1	-	-	1	-	1	-
Male	45	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	1	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Hawaii												
All races, both sexes	127	-	1	-	-	-	-	-	-	1	-	1
Male	73	-	1	-	-	-	-	-	-	1	-	1
Female	54	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	1	-	-
Male	14	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	103	-	1	-	-	-	-	-	-	-	-	1
Male	59	-	1	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	65	-	-	-	-	-	-	-	-	-	-	1
Male	39	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	-	-	-	-	1
Male	39	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Illinois												
All races, both sexes	918	1	-	-	-	-	1	-	2	1	6	2
Male	455	1	-	-	-	-	1	-	1	1	3	1
Female	463	-	-	-	-	-	-	-	1	-	3	1
White, both sexes	857	-	-	-	-	-	1	-	2	1	5	2
Male	426	-	-	-	-	-	1	-	1	1	3	1
Female	431	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	50	1	-	-	-	-	-	-	-	-	1	-
Male	24	1	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	477	-	-	-	-	1	-	-	2	2	3	10
Male	249	-	-	-	-	-	-	-	1	2	1	4
Female	228	-	-	-	-	1	-	-	1	-	2	6
White, both sexes	455	-	-	-	-	1	-	-	2	2	3	8
Male	237	-	-	-	-	-	-	-	1	2	1	3
Female	218	-	-	-	-	1	-	-	1	-	2	5
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Iowa												
All races, both sexes	283	1	-	1	-	-	-	2	-	3	-	1
Male	147	1	-	1	-	-	-	-	-	1	-	-
Female	136	-	-	-	-	-	-	2	-	2	-	1
White, both sexes	277	1	-	1	-	-	-	2	-	3	-	1
Male	143	1	-	1	-	-	-	-	-	1	-	-
Female	134	-	-	-	-	-	-	2	-	2	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	172	-	-	-	-	-	-	-	-	2	-	1
Male	85	-	-	-	-	-	-	-	-	2	-	1
Female	87	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	166	-	-	-	-	-	-	-	-	1	-	1
Male	81	-	-	-	-	-	-	-	-	1	-	1
Female	85	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Kentucky												
All races, both sexes	395	1	1	-	-	-	-	-	2	1	3	3
Male	214	-	1	-	-	-	-	-	1	-	2	2
Female	181	1	-	-	-	-	-	-	1	1	1	1
White, both sexes	372	1	1	-	-	-	-	-	2	1	1	3
Male	202	-	1	-	-	-	-	-	1	-	1	2
Female	170	1	-	-	-	-	-	-	1	1	-	1
Black, both sexes	22	-	-	-	-	-	-	-	-	-	2	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	220	-	-	-	1	-	-	-	1	3	2	1
Male	113	-	-	-	1	-	-	-	1	2	1	1
Female	107	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	164	-	-	-	1	-	-	-	1	2	1	-
Male	87	-	-	-	1	-	-	-	1	1	-	-
Female	77	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	55	-	-	-	-	-	-	-	-	1	1	1
Male	26	-	-	-	-	-	-	-	-	1	1	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Maine												
All races, both sexes	99	-	-	-	-	-	-	1	-	1	-	-
Male	55	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	98	-	-	-	-	-	-	1	-	1	-	-
Male	54	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	381	-	-	-	-	-	1	1	1	1	4	5
Male	206	-	-	-	-	-	-	1	1	1	1	3
Female	175	-	-	-	-	-	1	-	-	-	3	2
White, both sexes	298	-	-	-	-	-	1	1	-	1	3	3
Male	165	-	-	-	-	-	-	1	-	1	1	2
Female	133	-	-	-	-	-	1	-	-	-	2	1
Black, both sexes	77	-	-	-	-	-	-	-	1	-	1	2
Male	38	-	-	-	-	-	-	-	1	-	-	1
Female	39	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Massachusetts												
All races, both sexes	538	-	1	-	-	-	-	-	-	-	1	3
Male	287	-	1	-	-	-	-	-	-	-	1	1
Female	251	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	519	-	1	-	-	-	-	-	-	-	1	3
Male	277	-	1	-	-	-	-	-	-	-	1	1
Female	242	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	564	-	-	-	-	-	1	-	3	2	7	5
Male	313	-	-	-	-	-	-	-	2	1	5	4
Female	251	-	-	-	-	-	1	-	1	1	2	1
White, both sexes	510	-	-	-	-	-	-	-	3	-	7	3
Male	277	-	-	-	-	-	-	-	2	-	5	2
Female	233	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	51	-	-	-	-	-	1	-	-	1	-	2
Male	35	-	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Minnesota												
All races, both sexes	242	-	-	-	-	-	-	1	-	1	1	1
Male	141	-	-	-	-	-	-	1	-	1	1	1
Female	101	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	237	-	-	-	-	-	-	1	-	1	1	1
Male	137	-	-	-	-	-	-	1	-	1	1	1
Female	100	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	206	2	-	1	-	-	-	3	1	-	3	3
Male	118	1	-	1	-	-	-	2	1	-	2	1
Female	88	1	-	-	-	-	-	1	-	-	1	2
White, both sexes	159	-	-	-	-	-	-	2	1	-	2	2
Male	96	-	-	-	-	-	-	1	1	-	2	1
Female	63	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	46	2	-	1	-	-	-	1	-	-	1	1
Male	21	1	-	1	-	-	-	1	-	-	-	-
Female	25	1	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Missouri												
All races, both sexes	428	-	1	1	1	-	-	-	1	-	3	4
Male	226	-	-	-	1	-	-	-	-	-	1	3
Female	202	-	1	1	-	-	-	-	1	-	2	1
White, both sexes	397	-	1	1	-	-	-	-	1	-	2	3
Male	209	-	-	-	-	-	-	-	-	-	-	2
Female	188	-	1	1	-	-	-	-	1	-	2	1
Black, both sexes	29	-	-	-	1	-	-	-	-	-	1	1
Male	16	-	-	-	1	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	77	-	1	-	-	-	-	-	1	-	1	-
Male	39	-	-	-	-	-	-	-	1	-	-	-
Female	38	-	1	-	-	-	-	-	-	-	1	-
White, both sexes	74	-	1	-	-	-	-	-	-	-	1	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	1	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Nebraska												
All races, both sexes	119	-	-	-	-	1	-	-	1	1	1	1
Male	65	-	-	-	-	-	-	-	1	1	-	1
Female	54	-	-	-	-	1	-	-	-	-	1	-
White, both sexes	113	-	-	-	-	1	-	-	-	1	1	1
Male	61	-	-	-	-	-	-	-	-	1	-	1
Female	52	-	-	-	-	1	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	119	-	-	-	-	-	1	-	-	-	-	3
Male	68	-	-	-	-	-	1	-	-	-	-	2
Female	51	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	109	-	-	-	-	-	1	-	-	-	-	2
Male	64	-	-	-	-	-	1	-	-	-	-	2
Female	45	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
New Hampshire												
All races, both sexes	63	-	-	-	-	-	-	1	-	1	-	-
Male	28	-	-	-	-	-	-	1	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	63	-	-	-	-	-	-	1	-	1	-	-
Male	28	-	-	-	-	-	-	1	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	540	-	-	1	-	1	-	2	-	1	5	8
Male	295	-	-	1	-	-	-	1	-	1	4	4
Female	245	-	-	-	-	1	-	1	-	-	1	4
White, both sexes	477	-	-	1	-	1	-	2	-	-	5	3
Male	263	-	-	1	-	-	-	1	-	-	4	2
Female	214	-	-	-	-	1	-	1	-	-	1	1
Black, both sexes	59	-	-	-	-	-	-	-	-	-	-	5
Male	28	-	-	-	-	-	-	-	-	-	-	2
Female	31	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
New Mexico												
All races, both sexes	75	-	-	-	-	-	-	-	1	-	1	2
Male	41	-	-	-	-	-	-	-	1	-	-	2
Female	34	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	65	-	-	-	-	-	-	-	1	-	1	2
Male	39	-	-	-	-	-	-	-	1	-	-	2
Female	26	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	736	3	-	1	-	1	-	-	4	-	4	11
Male	411	2	-	-	-	1	-	-	3	-	3	6
Female	325	1	-	1	-	-	-	-	1	-	1	5
White, both sexes	689	1	-	1	-	1	-	-	2	-	3	11
Male	383	-	-	-	-	1	-	-	2	-	2	6
Female	306	1	-	1	-	-	-	-	-	-	1	5
Black, both sexes	40	2	-	-	-	-	-	-	2	-	1	-
Male	24	2	-	-	-	-	-	-	1	-	1	-
Female	16	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
North Carolina												
All races, both sexes	681	-	-	-	-	1	1	-	3	3	10	8
Male	366	-	-	-	-	-	-	-	2	2	7	3
Female	315	-	-	-	-	1	1	-	1	1	3	5
White, both sexes	570	-	-	-	-	1	-	-	2	-	6	6
Male	299	-	-	-	-	-	-	-	1	-	4	3
Female	271	-	-	-	-	1	-	-	1	-	2	3
Black, both sexes	108	-	-	-	-	-	1	-	1	3	4	2
Male	64	-	-	-	-	-	-	-	1	2	3	-
Female	44	-	-	-	-	-	1	-	-	1	1	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	53	-	-	-	-	-	-	-	-	-	1	1
Male	30	-	-	-	-	-	-	-	-	-	1	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	51	-	-	-	-	-	-	-	-	-	1	-
Male	29	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Ohio												
All races, both sexes	853	-	1	-	-	1	-	1	2	3	1	9
Male	466	-	-	-	-	-	-	1	2	-	1	6
Female	387	-	1	-	-	1	-	-	-	3	-	3
White, both sexes	776	-	1	-	-	-	-	-	1	3	1	8
Male	424	-	-	-	-	-	-	-	1	-	1	6
Female	352	-	1	-	-	-	-	-	-	3	-	2
Black, both sexes	76	-	-	-	-	1	-	1	1	-	-	1
Male	42	-	-	-	-	-	-	1	1	-	-	-
Female	34	-	-	-	-	1	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	214	-	-	-	-	-	-	1	2	-	3	2
Male	102	-	-	-	-	-	-	-	1	-	1	1
Female	112	-	-	-	-	-	-	1	1	-	2	1
White, both sexes	197	-	-	-	-	-	-	1	1	-	3	2
Male	95	-	-	-	-	-	-	-	-	-	1	1
Female	102	-	-	-	-	-	-	1	1	-	2	1
Black, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Oregon												
All races, both sexes	183	-	-	-	-	-	2	-	-	-	-	2
Male	98	-	-	-	-	-	1	-	-	-	-	1
Female	85	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	176	-	-	-	-	-	2	-	-	-	-	2
Male	92	-	-	-	-	-	1	-	-	-	-	1
Female	84	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,272	1	-	1	-	1	2	2	1	2	9	17
Male	749	1	-	1	-	1	1	1	1	-	6	8
Female	523	-	-	-	-	-	1	1	-	2	3	9
White, both sexes	1,162	1	-	1	-	-	2	1	1	2	7	13
Male	698	1	-	1	-	-	1	1	1	-	5	7
Female	464	-	-	-	-	-	1	-	-	2	2	6
Black, both sexes	103	-	-	-	-	1	-	1	-	-	2	4
Male	47	-	-	-	-	1	-	-	-	-	1	1
Female	56	-	-	-	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Rhode Island												
All races, both sexes	89	-	-	-	-	-	-	1	-	-	-	1
Male	51	-	-	-	-	-	-	-	-	-	-	1
Female	38	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	86	-	-	-	-	-	-	1	-	-	-	1
Male	49	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	294	-	-	-	-	-	-	2	2	2	1	4
Male	163	-	-	-	-	-	-	1	-	1	-	2
Female	131	-	-	-	-	-	-	1	2	1	1	2
White, both sexes	224	-	-	-	-	-	-	1	1	1	-	3
Male	123	-	-	-	-	-	-	-	-	-	-	2
Female	101	-	-	-	-	-	-	1	1	1	-	1
Black, both sexes	70	-	-	-	-	-	-	1	1	1	1	1
Male	40	-	-	-	-	-	-	1	-	1	-	-
Female	30	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
South Dakota												
All races, both sexes	46	-	-	-	-	-	-	-	-	1	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	44	-	-	-	-	-	-	-	-	1	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	473	-	-	-	-	-	1	2	1	-	4	8
Male	249	-	-	-	-	-	-	1	1	-	3	5
Female	224	-	-	-	-	-	1	1	-	-	1	3
White, both sexes	431	-	-	-	-	-	-	-	1	-	3	7
Male	226	-	-	-	-	-	-	-	1	-	3	4
Female	205	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	41	-	-	-	-	-	1	2	-	-	1	1
Male	22	-	-	-	-	-	-	1	-	-	-	1
Female	19	-	-	-	-	-	1	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Texas												
All races, both sexes	1,245	2	1	-	2	-	1	3	1	6	14	14
Male	684	1	-	-	2	-	1	3	1	4	10	9
Female	561	1	1	-	-	-	-	-	-	2	4	5
White, both sexes	1,137	2	1	-	2	-	1	1	1	6	10	13
Male	617	1	-	-	2	-	1	1	1	4	7	8
Female	520	1	1	-	-	-	-	-	-	2	3	5
Black, both sexes	102	-	-	-	-	-	-	2	-	-	3	1
Male	62	-	-	-	-	-	-	2	-	-	3	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	98	-	-	-	1	-	1	1	-	-	-	-
Male	58	-	-	-	1	-	1	1	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	97	-	-	-	1	-	1	1	-	-	-	-
Male	57	-	-	-	1	-	1	1	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Vermont												
All races, both sexes	41	-	-	-	-	1	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	1	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	610	-	-	-	-	-	1	-	1	2	3	9
Male	343	-	-	-	-	-	1	-	1	2	2	5
Female	267	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	487	-	-	-	-	-	-	-	1	1	1	6
Male	276	-	-	-	-	-	-	-	1	1	1	4
Female	211	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	113	-	-	-	-	-	1	-	-	1	2	3
Male	61	-	-	-	-	-	1	-	-	1	1	1
Female	52	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Washington												
All races, both sexes	454	-	-	-	-	1	-	-	1	3	4	5
Male	233	-	-	-	-	-	-	-	1	1	2	2
Female	221	-	-	-	-	1	-	-	-	2	2	3
White, both sexes	420	-	-	-	-	-	-	-	1	3	3	5
Male	213	-	-	-	-	-	-	-	1	1	2	2
Female	207	-	-	-	-	-	-	-	-	2	1	3
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	7	-	-	-	-	1	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	226	-	-	-	-	-	-	1	-	-	2	1
Male	165	-	-	-	-	-	-	1	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	219	-	-	-	-	-	-	-	-	-	1	1
Male	159	-	-	-	-	-	-	-	-	-	-	-
Female	60	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	7	-	-	-	-	-	-	1	-	-	1	-
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Wisconsin												
All races, both sexes	375	-	-	-	-	-	-	1	-	-	2	6
Male	226	-	-	-	-	-	-	1	-	-	2	2
Female	149	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	363	-	-	-	-	-	-	1	-	-	2	4
Male	219	-	-	-	-	-	-	1	-	-	2	1
Female	144	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	32	-	-	-	-	-	-	1	1	-	-	-
Male	17	-	-	-	-	-	-	-	1	-	-	-
Female	15	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	1	1	-	-	-
Male	17	-	-	-	-	-	-	-	1	-	-	-
Female	15	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
United States												
All races, both sexes	271	389	527	728	1,231	2,133	3,573	4,215	2,996	1,098	225	-
Male	179	249	346	483	799	1,330	2,098	2,223	1,388	382	67	-
Female	92	140	181	245	432	803	1,475	1,992	1,608	716	158	-
White, both sexes	225	320	455	614	1,045	1,881	3,269	3,919	2,757	1,017	192	-
Male	147	206	298	406	688	1,171	1,933	2,072	1,289	360	57	-
Female	78	114	157	208	357	710	1,336	1,847	1,468	657	135	-
Black, both sexes	45	61	64	96	164	204	245	239	178	64	24	-
Male	31	37	44	67	97	131	135	116	68	13	6	-
Female	14	24	20	29	67	73	110	123	110	51	18	-
American Indian 1/, both sexes	1	4	4	14	4	15	3	11	12	-	1	-
Male	1	3	2	8	3	8	1	8	5	-	1	-
Female	-	1	2	6	1	7	2	3	7	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	4	4	18	33	56	46	49	17	8	-
Male	-	3	2	2	11	20	29	27	26	9	3	-
Female	-	1	2	2	7	13	27	19	23	8	5	-
Alabama												
All races, both sexes	6	14	2	10	25	33	57	76	51	9	5	-
Male	4	10	2	6	16	20	31	44	16	4	1	-
Female	2	4	-	4	9	13	26	32	35	5	4	-
White, both sexes	4	11	2	8	21	25	47	66	43	7	4	-
Male	3	8	2	5	14	13	27	39	13	4	1	-
Female	1	3	-	3	7	12	20	27	30	3	3	-
Black, both sexes	2	3	-	1	4	8	10	10	8	2	1	-
Male	1	2	-	1	2	7	4	5	3	-	-	-
Female	1	1	-	-	2	1	6	5	5	2	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Alaska												
All races, both sexes	-	-	1	-	1	3	4	1	2	-	-	-
Male	-	-	-	-	-	3	3	-	-	-	-	-
Female	-	-	1	-	1	-	1	1	2	-	-	-
White, both sexes	-	-	1	-	1	3	3	1	-	-	-	-
Male	-	-	-	-	-	3	3	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	4	8	13	21	32	55	54	26	11	2	-
Male	3	2	4	6	16	12	35	29	16	6	2	-
Female	2	2	4	7	5	20	20	25	10	5	-	-
White, both sexes	5	2	8	13	21	29	55	50	25	10	1	-
Male	3	1	4	6	16	10	35	26	15	6	1	-
Female	2	1	4	7	5	19	20	24	10	4	-	-
Black, both sexes	-	1	-	-	-	1	-	-	-	1	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	2	-	3	1	-	1	-
Male	-	-	-	-	-	1	-	3	1	-	1	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Arkansas												
All races, both sexes	4	6	8	5	8	25	43	45	35	16	3	-
Male	2	2	7	4	3	15	29	25	13	6	-	-
Female	2	4	1	1	5	10	14	20	22	10	3	-
White, both sexes	2	4	6	4	5	22	40	43	33	15	2	-
Male	1	2	5	3	1	14	27	24	13	6	-	-
Female	1	2	1	1	4	8	13	19	20	9	2	-
Black, both sexes	2	2	2	1	3	3	3	2	2	1	1	-
Male	1	-	2	1	2	1	2	1	-	-	-	-
Female	1	2	-	-	1	2	1	1	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	13	17	34	40	52	120	198	215	140	67	17	-
Male	8	7	22	24	32	68	108	121	76	26	7	-
Female	5	10	12	16	20	52	90	94	64	41	10	-
White, both sexes	12	15	29	36	42	108	183	199	125	64	14	-
Male	7	7	19	23	26	62	101	113	67	25	6	-
Female	5	8	10	13	16	46	82	86	58	39	8	-
Black, both sexes	1	2	4	3	8	4	5	5	6	-	-	-
Male	1	-	3	1	5	2	4	2	3	-	-	-
Female	-	2	1	2	3	2	1	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	8	9	11	9	3	3	-
Male	-	-	-	-	1	4	3	6	6	1	1	-
Female	-	-	1	-	1	4	6	5	3	2	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Colorado												
All races, both sexes	5	5	9	5	15	36	49	68	55	14	3	-
Male	4	3	6	2	10	24	32	37	24	4	2	-
Female	1	2	3	3	5	12	17	31	31	10	1	-
White, both sexes	4	5	8	5	11	35	47	66	55	14	3	-
Male	3	3	5	2	7	24	31	36	24	4	2	-
Female	1	2	3	3	4	11	16	30	31	10	1	-
Black, both sexes	-	-	1	-	2	1	1	2	-	-	-	-
Male	-	-	1	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	1	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	1	5	10	10	18	24	74	84	70	30	2	-
Male	-	3	4	6	7	15	44	41	33	10	-	-
Female	1	2	6	4	11	9	30	43	37	20	2	-
White, both sexes	1	5	9	9	15	22	71	82	68	30	2	-
Male	-	3	4	5	6	13	41	41	33	10	-	-
Female	1	2	5	4	9	9	30	41	35	20	2	-
Black, both sexes	-	-	1	1	3	2	3	2	2	-	-	-
Male	-	-	-	1	1	2	3	-	-	-	-	-
Female	-	-	1	-	2	-	-	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Delaware												
All races, both sexes	1	1	3	1	2	6	12	15	13	2	-	-
Male	1	1	3	-	2	6	9	5	6	-	-	-
Female	-	-	-	1	-	-	3	10	7	2	-	-
White, both sexes	1	1	1	1	2	5	12	14	12	2	-	-
Male	1	1	1	-	2	5	9	4	5	-	-	-
Female	-	-	-	1	-	-	3	10	7	2	-	-
Black, both sexes	-	-	2	-	-	1	-	1	1	-	-	-
Male	-	-	2	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	-	1	-	3	4	9	5	3	-	-
Male	-	1	-	-	-	1	2	3	1	2	-	-
Female	-	-	-	1	-	2	2	6	4	1	-	-
White, both sexes	-	-	-	-	-	1	1	1	1	2	-	-
Male	-	-	-	-	-	1	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	-	1	-	1	-	2	2	8	4	1	-	-
Male	-	1	-	-	-	-	-	3	1	1	-	-
Female	-	-	-	1	-	2	2	5	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Florida												
All races, both sexes	9	27	22	38	59	107	208	231	163	62	11	-
Male	4	24	15	24	38	80	131	131	78	26	3	-
Female	5	3	7	14	21	27	77	100	85	36	8	-
White, both sexes	7	20	19	27	52	96	190	220	150	57	9	-
Male	3	17	14	15	33	71	120	124	71	24	3	-
Female	4	3	5	12	19	25	70	96	79	33	6	-
Black, both sexes	2	7	3	11	7	9	17	10	12	5	2	-
Male	1	7	1	9	5	7	11	6	6	2	-	-
Female	1	-	2	2	2	2	6	4	6	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	9	7	16	26	33	60	96	106	71	27	7	-
Male	5	4	11	22	23	41	59	51	34	10	3	-
Female	4	3	5	4	10	19	37	55	37	17	4	-
White, both sexes	7	2	12	22	23	47	76	89	60	18	6	-
Male	4	2	9	18	17	31	49	46	31	7	3	-
Female	3	-	3	4	6	16	27	43	29	11	3	-
Black, both sexes	2	5	4	4	10	13	19	17	10	9	1	-
Male	1	2	2	4	6	10	10	5	2	3	-	-
Female	1	3	2	-	4	3	9	12	8	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Hawaii												
All races, both sexes	-	3	3	5	5	12	29	21	31	13	2	-
Male	-	3	3	3	4	7	17	10	14	7	2	-
Female	-	-	-	2	1	5	12	11	17	6	-	-
White, both sexes	-	-	1	1	-	3	6	5	7	-	-	-
Male	-	-	1	1	-	2	3	3	3	-	-	-
Female	-	-	-	-	-	1	3	2	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	2	4	5	9	23	16	24	13	2	-
Male	-	3	2	2	4	5	14	7	11	7	2	-
Female	-	-	-	2	1	4	9	9	13	6	-	-
Idaho												
All races, both sexes	3	5	4	1	4	8	13	14	11	1	-	-
Male	1	4	2	1	4	5	4	11	6	-	-	-
Female	2	1	2	-	-	3	9	3	5	1	-	-
White, both sexes	3	5	4	1	4	8	13	14	11	1	-	-
Male	1	4	2	1	4	5	4	11	6	-	-	-
Female	2	1	2	-	-	3	9	3	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Illinois												
All races, both sexes	13	18	29	29	54	98	182	241	171	58	12	-
Male	10	11	20	15	30	50	93	112	84	21	1	-
Female	3	7	9	14	24	48	89	129	87	37	11	-
White, both sexes	11	17	26	28	43	86	171	230	168	56	10	-
Male	9	11	18	14	23	43	89	108	83	20	1	-
Female	2	6	8	14	20	43	82	122	85	36	9	-
Black, both sexes	2	1	2	1	10	9	7	10	3	2	1	-
Male	1	-	2	1	6	4	3	4	1	1	-	-
Female	1	1	-	-	4	5	4	6	2	1	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	3	4	1	-	-	1	-
Male	-	-	-	-	1	3	1	-	-	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	1	-
Indiana												
All races, both sexes	9	15	15	23	38	54	85	113	74	27	6	-
Male	6	10	11	12	23	30	51	58	31	7	2	-
Female	3	5	4	11	15	24	34	55	43	20	4	-
White, both sexes	9	14	14	21	37	51	82	110	71	26	4	-
Male	6	9	11	12	22	28	49	57	29	6	1	-
Female	3	5	3	9	15	23	33	53	42	20	3	-
Black, both sexes	-	1	1	2	1	3	3	3	3	1	2	-
Male	-	1	-	-	1	2	2	1	2	1	1	-
Female	-	-	1	2	-	1	1	2	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Iowa												
All races, both sexes	2	1	7	9	20	27	41	79	64	23	2	-
Male	1	-	5	7	10	13	25	38	31	14	-	-
Female	1	1	2	2	10	14	16	41	33	9	2	-
White, both sexes	2	1	7	7	20	26	40	79	62	23	2	-
Male	1	-	5	6	10	12	24	38	30	14	-	-
Female	1	1	2	1	10	14	16	41	32	9	2	-
Black, both sexes	-	-	-	1	-	1	1	-	2	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	4	-	4	8	3	16	31	43	36	20	4	-
Male	2	-	1	7	3	9	13	24	18	5	-	-
Female	2	-	3	1	-	7	18	19	18	15	4	-
White, both sexes	4	-	4	8	3	15	30	42	34	20	4	-
Male	2	-	1	7	3	8	13	23	17	5	-	-
Female	2	-	3	1	-	7	17	19	17	15	4	-
Black, both sexes	-	-	-	-	-	1	1	-	2	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Kentucky												
All races, both sexes	3	8	12	22	28	56	76	84	74	19	2	-
Male	2	8	8	16	20	41	39	36	36	1	1	-
Female	1	-	4	6	8	15	37	48	38	18	1	-
White, both sexes	3	8	11	19	28	52	74	80	70	16	2	-
Male	2	8	7	14	20	38	38	34	34	1	1	-
Female	1	-	4	5	8	14	36	46	36	15	1	-
Black, both sexes	-	-	1	3	-	4	1	4	4	3	-	-
Male	-	-	1	2	-	3	1	2	2	-	-	-
Female	-	-	-	1	-	1	-	2	2	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Louisiana												
All races, both sexes	3	4	12	11	19	30	48	39	35	9	2	-
Male	3	3	5	4	12	22	26	14	13	5	-	-
Female	-	1	7	7	7	8	22	25	22	4	2	-
White, both sexes	3	3	9	9	11	21	39	29	26	8	1	-
Male	3	2	4	3	7	17	23	11	10	4	-	-
Female	-	1	5	6	4	4	16	18	16	4	1	-
Black, both sexes	-	1	3	2	8	9	9	9	9	1	1	-
Male	-	1	1	1	5	5	3	3	3	1	-	-
Female	-	-	2	1	3	4	6	6	6	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Maine												
All races, both sexes	-	1	1	4	8	9	19	27	17	10	1	-
Male	-	1	1	4	6	6	10	16	8	2	1	-
Female	-	-	-	-	2	3	9	11	9	8	-	-
White, both sexes	-	1	1	3	8	9	19	27	17	10	1	-
Male	-	1	1	3	6	6	10	16	8	2	1	-
Female	-	-	-	-	2	3	9	11	9	8	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	10	12	14	10	37	48	68	90	54	21	4	-
Male	5	5	11	6	21	32	41	47	26	5	-	-
Female	5	7	3	4	16	16	27	43	28	16	4	-
White, both sexes	6	6	9	4	22	36	56	80	47	20	3	-
Male	2	3	8	2	16	25	32	42	25	5	-	-
Female	4	3	1	2	6	11	24	38	22	15	3	-
Black, both sexes	4	6	5	6	14	9	10	10	7	1	1	-
Male	3	2	3	4	5	6	7	5	1	-	-	-
Female	1	4	2	2	9	3	3	5	6	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	3	2	-	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Massachusetts												
All races, both sexes	5	5	7	13	31	74	119	121	104	43	11	-
Male	2	3	7	7	14	54	68	62	45	17	5	-
Female	3	2	-	6	17	20	51	59	59	26	6	-
White, both sexes	5	5	7	12	30	72	115	116	100	42	10	-
Male	2	3	7	6	13	53	65	59	44	17	5	-
Female	3	2	-	6	17	19	50	57	56	25	5	-
Black, both sexes	-	-	-	1	-	1	2	3	4	1	1	-
Male	-	-	-	1	-	-	2	1	1	-	-	-
Female	-	-	-	-	-	1	-	2	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	2	2	-	-	-	-
Male	-	-	-	-	1	1	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Michigan												
All races, both sexes	12	13	13	28	33	78	109	129	95	30	6	-
Male	9	9	11	18	24	46	66	69	36	11	2	-
Female	3	4	2	10	9	32	43	60	59	19	4	-
White, both sexes	7	11	11	26	27	72	98	120	89	30	6	-
Male	5	7	10	17	20	41	58	64	33	11	2	-
Female	2	4	1	9	7	31	40	56	56	19	4	-
Black, both sexes	5	2	2	2	6	6	11	9	4	-	-	-
Male	4	2	1	1	4	5	8	5	3	-	-	-
Female	1	-	1	1	2	1	3	4	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Minnesota												
All races, both sexes	5	3	9	6	14	35	36	57	52	16	5	-
Male	5	2	6	6	11	20	22	34	25	5	1	-
Female	-	1	3	-	3	15	14	23	27	11	4	-
White, both sexes	5	3	9	5	12	34	35	57	52	16	5	-
Male	5	2	6	5	9	20	21	34	25	5	1	-
Female	-	1	3	-	3	14	14	23	27	11	4	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	7	9	10	15	29	36	45	23	13	2	-
Male	3	4	8	8	7	22	24	23	9	2	-	-
Female	1	3	1	2	8	7	12	22	14	11	2	-
White, both sexes	3	4	7	7	12	23	28	37	19	10	2	-
Male	2	1	6	6	6	18	22	19	9	2	-	-
Female	1	3	1	1	6	5	6	18	10	8	2	-
Black, both sexes	1	3	2	3	3	5	8	8	4	3	-	-
Male	1	3	2	2	1	3	2	4	-	-	-	-
Female	-	-	-	1	2	2	6	4	4	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Missouri												
All races, both sexes	11	9	10	13	29	55	84	101	78	21	6	-
Male	5	2	7	7	21	34	45	53	39	6	2	-
Female	6	7	3	6	8	21	39	48	39	15	4	-
White, both sexes	10	8	10	11	27	50	79	97	70	21	6	-
Male	4	2	7	5	20	30	44	51	36	6	2	-
Female	6	6	3	6	7	20	35	46	34	15	4	-
Black, both sexes	1	1	-	2	2	4	5	4	7	-	-	-
Male	1	-	-	2	1	3	1	2	3	-	-	-
Female	-	1	-	-	1	1	4	2	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Montana												
All races, both sexes	-	1	2	4	5	9	14	11	21	7	-	-
Male	-	1	2	3	4	4	10	3	8	3	-	-
Female	-	-	-	1	1	5	4	8	13	4	-	-
White, both sexes	-	1	2	3	4	9	14	11	21	7	-	-
Male	-	1	2	2	3	4	10	3	8	3	-	-
Female	-	-	-	1	1	5	4	8	13	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Nebraska												
All races, both sexes	-	2	1	6	2	15	19	37	19	11	2	-
Male	-	2	1	5	1	11	12	14	11	4	1	-
Female	-	-	-	1	1	4	7	23	8	7	1	-
White, both sexes	-	2	-	4	2	13	19	37	19	11	2	-
Male	-	2	-	4	1	10	12	14	11	4	1	-
Female	-	-	-	-	1	3	7	23	8	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	2	-	2	-	-	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	2	1	10	11	15	32	17	17	6	-	-
Male	3	2	1	6	6	11	22	7	5	2	-	-
Female	1	-	-	4	5	4	10	10	12	4	-	-
White, both sexes	4	2	1	9	8	14	30	17	15	6	-	-
Male	3	2	1	5	5	10	21	7	5	2	-	-
Female	1	-	-	4	3	4	9	10	10	4	-	-
Black, both sexes	-	-	-	-	2	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
New Hampshire												
All races, both sexes	1	1	4	3	3	1	15	16	12	5	-	-
Male	1	-	2	-	3	-	5	9	7	-	-	-
Female	-	1	2	3	-	1	10	7	5	5	-	-
White, both sexes	1	1	4	3	3	1	15	16	12	5	-	-
Male	1	-	2	-	3	-	5	9	7	-	-	-
Female	-	1	2	3	-	1	10	7	5	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	10	13	12	11	34	61	112	150	88	27	4	-
Male	6	6	9	7	23	37	68	81	39	8	-	-
Female	4	7	3	4	11	24	44	69	49	19	4	-
White, both sexes	4	12	9	8	30	51	103	136	83	25	4	-
Male	2	6	7	6	22	29	63	74	38	8	-	-
Female	2	6	2	2	8	22	40	62	45	17	4	-
Black, both sexes	6	1	3	3	4	9	9	12	5	2	-	-
Male	4	-	2	1	1	7	5	5	1	-	-	-
Female	2	1	1	2	3	2	4	7	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
New Mexico												
All races, both sexes	1	2	3	6	7	14	13	12	8	3	2	-
Male	-	1	3	4	2	5	9	8	3	2	1	-
Female	1	1	-	2	5	9	4	4	5	1	1	-
White, both sexes	1	1	3	5	5	11	13	11	6	3	2	-
Male	-	-	3	4	2	5	9	8	2	2	1	-
Female	1	1	-	1	3	6	4	3	4	1	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	1	3	-	1	2	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	1	3	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	11	15	25	26	45	72	148	201	107	53	9	-
Male	7	8	14	20	28	49	89	115	47	16	3	-
Female	4	7	11	6	17	23	59	86	60	37	6	-
White, both sexes	10	13	22	25	41	71	140	191	102	48	7	-
Male	7	6	12	19	26	48	85	108	43	16	2	-
Female	3	7	10	6	15	23	55	83	59	32	5	-
Black, both sexes	1	2	3	1	4	1	7	6	4	5	1	-
Male	-	2	2	1	2	1	4	4	3	-	1	-
Female	1	-	1	-	2	-	3	2	1	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	4	1	-	1	-
Male	-	-	-	-	-	-	-	3	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
North Carolina												
All races, both sexes	11	18	32	43	55	77	133	143	99	39	5	-
Male	7	13	20	27	35	44	75	69	53	8	1	-
Female	4	5	12	16	20	33	58	74	46	31	4	-
White, both sexes	8	10	26	36	42	63	120	127	84	35	4	-
Male	4	6	15	25	25	37	69	58	44	8	-	-
Female	4	4	11	11	17	26	51	69	40	27	4	-
Black, both sexes	3	7	5	7	13	13	13	16	15	4	1	-
Male	3	6	4	2	10	6	6	11	9	-	1	-
Female	-	1	1	5	3	7	7	5	6	4	-	-
American Indian 1/, both sexes	-	1	1	-	-	1	-	-	-	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	1	2	2	7	11	14	7	5	1	-
Male	-	-	1	1	2	4	7	6	4	3	-	-
Female	-	1	-	1	-	3	4	8	3	2	1	-
White, both sexes	-	1	1	1	2	7	11	14	7	5	1	-
Male	-	-	1	1	2	4	7	6	4	3	-	-
Female	-	1	-	-	-	3	4	8	3	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Ohio												
All races, both sexes	12	13	21	35	69	105	146	209	160	59	6	-
Male	9	10	16	25	49	77	87	99	63	20	1	-
Female	3	3	5	10	20	28	59	110	97	39	5	-
White, both sexes	11	11	18	32	61	91	135	193	150	55	5	-
Male	8	9	13	22	44	67	80	95	59	19	-	-
Female	3	2	5	10	17	24	55	98	91	36	5	-
Black, both sexes	1	2	3	3	7	14	11	16	10	4	1	-
Male	1	1	3	3	5	10	7	4	4	1	1	-
Female	-	1	-	-	2	4	4	12	6	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	2	6	14	15	27	40	49	38	10	5	-
Male	-	2	2	9	7	16	20	24	15	2	2	-
Female	-	-	4	5	8	11	20	25	23	8	3	-
White, both sexes	-	2	5	11	14	25	39	47	33	10	4	-
Male	-	2	2	7	6	16	19	23	14	2	2	-
Female	-	-	3	4	8	9	20	24	19	8	2	-
Black, both sexes	-	-	-	1	-	1	1	-	1	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	1	2	1	1	-	2	4	-	-	-
Male	-	-	-	2	1	-	-	1	1	-	-	-
Female	-	-	1	-	-	1	-	1	3	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Oregon												
All races, both sexes	7	2	3	8	16	16	32	38	41	12	4	-
Male	5	1	1	4	7	9	24	21	17	6	1	-
Female	2	1	2	4	9	7	8	17	24	6	3	-
White, both sexes	6	2	3	8	16	14	32	36	39	12	4	-
Male	4	1	1	4	7	7	24	20	15	6	1	-
Female	2	1	2	4	9	7	8	16	24	6	3	-
Black, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	1	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	18	28	27	38	78	160	291	295	227	53	21	-
Male	15	20	18	23	48	97	185	183	117	15	8	-
Female	3	8	9	15	30	63	106	112	110	38	13	-
White, both sexes	16	27	25	29	68	141	268	283	207	51	19	-
Male	13	19	16	18	42	89	175	177	110	15	7	-
Female	3	8	9	11	26	52	93	106	97	36	12	-
Black, both sexes	2	1	2	9	9	19	21	11	17	2	2	-
Male	2	1	2	5	5	8	9	6	5	-	1	-
Female	-	-	-	4	4	11	12	5	12	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	2	1	3	-	-	-
Male	-	-	-	-	1	-	1	-	2	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Rhode Island												
All races, both sexes	-	3	2	5	6	12	16	26	12	4	1	-
Male	-	1	1	4	5	8	10	12	8	-	1	-
Female	-	2	1	1	1	4	6	14	4	4	-	-
White, both sexes	-	3	2	5	6	10	16	25	12	4	1	-
Male	-	1	1	4	5	6	10	12	8	-	1	-
Female	-	2	1	1	1	4	6	13	4	4	-	-
Black, both sexes	-	-	-	-	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	9	7	15	34	40	67	68	31	7	4	-
Male	1	5	5	14	28	28	33	30	11	2	2	-
Female	-	4	2	1	6	12	34	38	20	5	2	-
White, both sexes	1	5	4	10	23	29	54	56	26	7	3	-
Male	1	2	4	10	20	21	28	22	9	2	2	-
Female	-	3	-	-	3	8	26	34	17	5	1	-
Black, both sexes	-	4	3	5	11	11	13	12	5	-	1	-
Male	-	3	1	4	8	7	5	8	2	-	-	-
Female	-	1	2	1	3	4	8	4	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
South Dakota												
All races, both sexes	-	-	-	2	3	6	4	16	6	8	-	-
Male	-	-	-	1	2	5	3	9	3	3	-	-
Female	-	-	-	1	1	1	1	7	3	5	-	-
White, both sexes	-	-	-	2	3	5	4	16	5	8	-	-
Male	-	-	-	1	2	4	3	9	3	3	-	-
Female	-	-	-	1	1	1	1	7	2	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	15	19	22	31	56	90	123	67	21	7	-
Male	4	8	11	14	23	32	52	54	31	9	1	-
Female	2	7	8	8	8	24	38	69	36	12	6	-
White, both sexes	4	14	16	19	27	52	86	116	64	17	5	-
Male	2	8	8	11	20	30	50	50	29	9	1	-
Female	2	6	8	8	7	22	36	66	35	8	4	-
Black, both sexes	2	1	3	3	4	4	4	7	2	4	2	-
Male	2	-	3	3	3	2	2	4	1	-	-	-
Female	-	1	-	-	1	2	2	3	1	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Texas												
All races, both sexes	22	28	39	56	98	134	271	260	179	94	20	-
Male	16	19	24	40	63	72	155	143	85	31	5	-
Female	6	9	15	16	35	62	116	117	94	63	15	-
White, both sexes	19	25	35	45	90	126	245	241	170	86	18	-
Male	15	19	22	31	59	67	135	131	80	29	4	-
Female	4	6	13	14	31	59	110	110	90	57	14	-
Black, both sexes	3	3	4	11	8	8	26	14	9	8	2	-
Male	1	-	2	9	4	5	20	7	5	2	1	-
Female	2	3	2	2	4	3	6	7	4	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	4	-	-	-	-
Male	-	-	-	-	-	-	-	4	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	4	8	3	-	13	10	18	22	13	4	-	-
Male	2	4	1	-	8	3	10	18	7	2	-	-
Female	2	4	2	-	5	7	8	4	6	2	-	-
White, both sexes	4	8	3	-	13	10	18	21	13	4	-	-
Male	2	4	1	-	8	3	10	17	7	2	-	-
Female	2	4	2	-	5	7	8	4	6	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Vermont												
All races, both sexes	-	-	1	1	1	3	11	8	8	6	1	-
Male	-	-	-	1	1	2	6	5	3	2	1	-
Female	-	-	1	-	-	1	5	3	5	4	-	-
White, both sexes	-	-	1	1	1	3	11	8	8	6	1	-
Male	-	-	-	1	1	2	6	5	3	2	1	-
Female	-	-	1	-	-	1	5	3	5	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	10	12	17	26	59	92	104	136	97	34	7	-
Male	8	5	12	21	40	61	62	70	36	16	1	-
Female	2	7	5	5	19	31	42	66	61	18	6	-
White, both sexes	7	8	14	19	44	69	83	117	83	30	4	-
Male	6	4	9	14	31	44	52	60	33	15	1	-
Female	1	4	5	5	13	25	31	57	50	15	3	-
Black, both sexes	3	3	3	7	15	20	18	19	13	3	2	-
Male	2	1	3	7	9	15	8	10	2	-	-	-
Female	1	2	-	-	6	5	10	9	11	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	2	3	-	-	1	1	-
Male	-	-	-	-	-	1	2	-	-	1	-	-
Female	-	1	-	-	-	1	1	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Washington												
All races, both sexes	6	10	17	24	27	34	80	113	86	36	7	-
Male	2	7	7	14	21	15	39	65	43	12	2	-
Female	4	3	10	10	6	19	41	48	43	24	5	-
White, both sexes	5	8	16	22	24	31	73	105	81	36	7	-
Male	2	5	7	13	20	14	33	59	40	12	2	-
Female	3	3	9	9	4	17	40	46	41	24	5	-
Black, both sexes	1	1	1	-	1	-	3	5	1	-	-	-
Male	-	1	-	-	1	-	2	4	1	-	-	-
Female	1	-	1	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	2	-	1	-	1	1	-	-	-
Male	-	1	-	1	-	-	-	1	1	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	2	4	2	3	-	-	-
Male	-	-	-	-	-	1	4	1	1	-	-	-
Female	-	-	-	-	2	1	-	1	2	-	-	-
West Virginia												
All races, both sexes	5	6	6	16	23	43	38	45	30	8	2	-
Male	5	6	5	15	21	33	27	29	18	4	1	-
Female	-	-	1	1	2	10	11	16	12	4	1	-
White, both sexes	5	6	6	16	21	42	38	44	30	7	2	-
Male	5	6	5	15	19	32	27	28	18	3	1	-
Female	-	-	1	1	2	10	11	16	12	4	1	-
Black, both sexes	-	-	-	-	2	1	-	1	-	1	-	-
Male	-	-	-	-	2	1	-	1	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Wisconsin												
All races, both sexes	5	7	15	14	17	44	86	90	66	20	2	-
Male	3	6	9	10	12	30	56	52	33	10	-	-
Female	2	1	6	4	5	14	30	38	33	10	2	-
White, both sexes	5	7	13	14	15	40	86	89	65	20	2	-
Male	3	6	8	10	11	28	56	51	32	10	-	-
Female	2	1	5	4	4	12	30	38	33	10	2	-
Black, both sexes	-	-	1	-	2	4	-	1	1	-	-	-
Male	-	-	1	-	1	2	-	1	1	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	5	2	6	8	7	1	-	-
Male	-	-	1	-	3	1	5	3	3	-	-	-
Female	-	-	-	-	2	1	1	5	4	1	-	-
White, both sexes	-	-	1	-	5	2	6	8	7	1	-	-
Male	-	-	1	-	3	1	5	3	3	-	-	-
Female	-	-	-	-	2	1	1	5	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
United States												
All races, both sexes	924	2	1	-	-	1	-	-	1	1	2	8
Male	887	2	1	-	-	-	-	-	1	1	1	4
Female	37	-	-	-	-	1	-	-	-	-	1	4
White, both sexes	868	1	1	-	-	-	-	-	1	1	2	7
Male	835	1	1	-	-	-	-	-	1	1	1	3
Female	33	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	49	1	-	-	-	1	-	-	-	-	-	1
Male	46	1	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Arkansas												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	1
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	44	-	-	-	-	-	-	-	-	-	-	1
Male	40	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Colorado												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Delaware												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Florida												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	38	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	38	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Hawaii												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Illinois												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Kentucky												
All races, both sexes	45	-	1	-	-	-	-	-	-	-	-	-
Male	45	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	45	-	1	-	-	-	-	-	-	-	-	-
Male	45	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Maine												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Massachusetts												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Minnesota												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Missouri												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Nebraska												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
New Mexico												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	21	1	-	-	-	-	-	-	-	-	-	-
Male	21	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
North Carolina												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	1	-
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Ohio												
All races, both sexes	44	-	-	-	-	1	-	-	-	-	-	1
Male	40	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	1
Male	39	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Oregon												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	137	-	-	-	-	-	-	-	-	-	-	-
Male	136	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	136	-	-	-	-	-	-	-	-	-	-	-
Male	135	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
South Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Texas												
All races, both sexes	44	1	-	-	-	-	-	-	-	-	-	1
Male	41	1	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	1	-	-	-	-	-	-	-	-	-	1
Male	39	1	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Vermont												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	43	-	-	-	-	-	-	-	-	-	-	1
Male	43	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	1
Male	42	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Washington												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	102	-	-	-	-	-	-	-	-	-	-	-
Male	101	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	98	-	-	-	-	-	-	-	-	-	-	-
Male	97	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
Wisconsin												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
United States												
All races, both sexes	12	22	28	60	123	163	229	170	92	7	2	-
Male	12	22	28	57	117	161	222	162	89	6	1	-
Female	-	-	-	3	6	2	7	8	3	1	1	-
White, both sexes	9	17	26	57	116	155	215	163	88	7	2	-
Male	9	17	26	54	110	153	210	155	86	6	1	-
Female	-	-	-	3	6	2	5	8	2	1	1	-
Black, both sexes	3	5	1	2	6	7	12	6	4	-	-	-
Male	3	5	1	2	6	7	11	6	3	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	1	1	1	1	1	-	-	-	-
Male	-	-	1	1	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	2	1	-	6	3	7	2	5	-	-	-
Male	1	2	1	-	6	2	6	2	5	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	1	1	1	-	6	2	5	2	3	-	-	-
Male	1	1	1	-	6	1	4	2	3	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	1	-	-	-	1	2	-	2	-	-	-
Male	-	1	-	-	-	1	2	-	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Alaska												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	1	3	-	1	3	-	-	-	-
Male	-	-	-	1	3	-	1	3	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	3	-	1	2	-	-	-	-
Male	-	-	-	1	3	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Arkansas												
All races, both sexes	-	-	2	2	1	1	2	-	1	1	-	-
Male	-	-	2	2	1	1	2	-	1	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	2	2	-	1	2	-	1	1	-	-
Male	-	-	2	2	-	1	2	-	1	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	2	3	3	8	12	12	4	1	-	-
Male	-	-	2	2	3	8	11	11	3	1	-	-
Female	-	-	-	1	-	-	1	1	1	-	-	-
White, both sexes	-	-	2	3	3	8	11	11	4	1	-	-
Male	-	-	2	2	3	8	11	10	3	1	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Colorado												
All races, both sexes	-	-	1	1	1	2	6	2	3	-	-	-
Male	-	-	1	1	1	2	6	2	3	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	2	6	2	3	-	-	-
Male	-	-	-	1	1	2	6	2	3	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	1	2	2	1	-	-
Male	-	-	-	-	-	-	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	-	-	-	-	-	-	1	2	2	1	-	-
Male	-	-	-	-	-	-	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Delaware												
All races, both sexes	-	-	-	-	1	1	4	-	2	-	-	-
Male	-	-	-	-	1	1	4	-	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	4	-	2	-	-	-
Male	-	-	-	-	1	1	4	-	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Florida												
All races, both sexes	-	3	1	3	6	5	9	10	2	-	-	-
Male	-	3	1	3	6	5	9	9	2	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	3	1	3	6	5	9	10	2	-	-	-
Male	-	3	1	3	6	5	9	9	2	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	1	1	2	1	2	1	-	-	-
Male	-	-	-	1	1	2	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	2	1	2	1	-	-	-
Male	-	-	-	1	-	2	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Hawaii												
All races, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	1	1	-	2	2	-	-	-
Male	-	-	-	-	1	1	-	2	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	2	2	-	-	-
Male	-	-	-	-	1	1	-	2	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Illinois												
All races, both sexes	-	1	-	3	2	3	5	3	1	-	-	-
Male	-	1	-	2	2	3	5	3	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	3	2	3	5	3	1	-	-	-
Male	-	1	-	2	2	3	5	3	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	1	1	-	1	1	2	1	1	-	-	-
Male	-	1	1	-	-	1	2	1	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	1	1	-	1	1	2	1	1	-	-	-
Male	-	1	1	-	-	1	2	1	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Kentucky												
All races, both sexes	1	4	1	6	9	7	8	5	3	-	-	-
Male	1	4	1	6	9	7	8	5	3	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	4	1	6	9	7	8	5	3	-	-	-
Male	1	4	1	6	9	7	8	5	3	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	1	1	4	4	-	-	-	-	-
Male	-	-	-	1	1	4	4	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	3	4	-	-	-	-	-
Male	-	-	-	-	1	3	4	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Maine												
All races, both sexes	-	-	-	2	2	-	1	2	1	-	-	-
Male	-	-	-	2	2	-	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	2	-	1	2	1	-	-	-
Male	-	-	-	1	2	-	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	2	-	-	4	4	4	4	-	1	-	-
Male	1	2	-	-	4	4	3	4	-	1	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	3	3	4	4	-	1	-	-
Male	1	-	-	-	3	3	3	4	-	1	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	2	-	-	1	1	-	-	-	-	-	-
Male	-	2	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Massachusetts												
All races, both sexes	-	-	-	-	1	2	3	2	1	-	-	-
Male	-	-	-	-	1	2	3	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	2	3	2	1	-	-	-
Male	-	-	-	-	1	2	3	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	3	2	5	6	4	2	-	-	-
Male	1	-	-	3	2	5	6	3	2	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	2	2	4	4	3	2	-	-	-
Male	-	-	-	2	2	4	4	2	2	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	1	-	1	2	1	-	-	-	-
Male	1	-	-	1	-	1	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Minnesota												
All races, both sexes	1	-	-	-	3	-	4	2	1	-	-	-
Male	1	-	-	-	3	-	4	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	3	-	4	2	1	-	-	-
Male	1	-	-	-	3	-	4	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	2	2	3	4	4	2	-	-	-	-
Male	-	1	2	2	3	4	3	2	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	2	2	3	4	2	1	-	-	-	-
Male	-	-	2	2	3	4	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	2	1	-	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Missouri												
All races, both sexes	-	-	1	-	2	1	2	1	-	-	-	-
Male	-	-	1	-	2	1	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	2	1	2	1	-	-	-	-
Male	-	-	1	-	2	1	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Nebraska												
All races, both sexes	-	-	1	-	1	2	2	1	-	-	-	-
Male	-	-	1	-	1	2	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	2	2	1	-	-	-	-
Male	-	-	-	-	1	2	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	1	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
New Hampshire												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	-	-	-	3	3	11	4	2	-	-	-
Male	1	-	-	-	3	3	9	4	2	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	3	3	9	4	2	-	-	-
Male	-	-	-	-	3	3	7	4	2	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
New Mexico												
All races, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	1	2	3	8	4	1	1	-	-
Male	-	-	-	1	2	3	8	4	1	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	2	3	8	4	1	1	-	-
Male	-	-	-	1	2	3	8	4	1	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
North Carolina												
All races, both sexes	1	2	1	1	4	2	4	3	1	-	-	-
Male	1	2	1	1	4	2	4	3	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	1	1	3	2	3	3	1	-	-	-
Male	-	1	1	1	3	2	3	3	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	1	-	1	-	-	-	-	-
Male	1	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	1	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	1	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Ohio												
All races, both sexes	-	2	2	3	7	11	3	10	4	-	-	-
Male	-	2	2	2	6	11	3	10	4	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
White, both sexes	-	2	2	3	7	11	3	9	4	-	-	-
Male	-	2	2	2	6	11	3	9	4	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	3	1	1	4	1	1	-	-	-
Male	-	-	-	3	1	1	4	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	3	-	1	4	1	1	-	-	-
Male	-	-	-	3	-	1	4	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Oregon												
All races, both sexes	-	-	-	1	-	1	8	1	1	-	1	-
Male	-	-	-	1	-	1	8	-	1	-	1	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	8	1	1	-	1	-
Male	-	-	-	1	-	-	8	-	1	-	1	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	3	3	6	22	39	34	28	1	1	-
Male	-	-	3	3	6	22	39	34	28	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	3	3	6	21	39	34	28	1	1	-
Male	-	-	3	3	6	21	39	34	28	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Rhode Island												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	2	4	4	2	6	-	-	-	-
Male	-	-	-	2	4	4	2	4	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	2	4	4	2	6	-	-	-	-
Male	-	-	-	2	4	4	2	4	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
South Dakota												
All races, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	1	1	1	2	2	7	2	3	-	-	-
Male	-	1	1	1	2	2	7	2	3	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	1	1	2	2	7	2	3	-	-	-
Male	-	1	1	1	2	2	7	2	3	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Texas												
All races, both sexes	1	-	-	4	8	12	8	7	2	-	-	-
Male	1	-	-	4	6	12	8	6	2	-	-	-
Female	-	-	-	-	2	-	-	1	-	-	-	-
White, both sexes	1	-	-	4	8	12	6	7	2	-	-	-
Male	1	-	-	4	6	12	6	6	2	-	-	-
Female	-	-	-	-	2	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Vermont												
All races, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	3	1	6	10	11	7	4	-	-	-
Male	-	-	3	1	6	10	11	7	4	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	3	1	6	10	10	7	4	-	-	-
Male	-	-	3	1	6	10	10	7	4	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Washington												
All races, both sexes	-	-	-	-	5	1	4	4	-	-	-	-
Male	-	-	-	-	5	1	4	4	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	5	1	3	4	-	-	-	-
Male	-	-	-	-	5	1	3	4	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	4	2	4	9	15	24	18	18	8	-	-	-
Male	4	2	4	9	14	24	18	18	8	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	4	2	4	9	13	23	18	17	8	-	-	-
Male	4	2	4	9	12	23	18	17	8	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	2	1	-	1	-	-	-	-
Male	-	-	-	-	2	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
Wisconsin												
All races, both sexes	-	1	-	-	2	5	5	-	1	-	-	-
Male	-	1	-	-	1	5	5	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	2	4	5	-	1	-	-	-
Male	-	1	-	-	1	4	5	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
United States												
All races, both sexes	16,887	11	9	6	7	11	19	34	44	52	114	184
Male	8,916	5	3	4	7	4	11	22	31	34	69	109
Female	7,971	6	6	2	-	7	8	12	13	18	45	75
White, both sexes	15,136	7	6	5	5	7	12	23	34	40	86	147
Male	7,984	2	1	3	5	2	8	14	24	24	55	91
Female	7,152	5	5	2	-	5	4	9	10	16	31	56
Black, both sexes	1,439	4	-	1	2	2	7	11	9	10	27	33
Male	757	3	-	1	2	1	3	8	6	8	14	14
Female	682	1	-	-	-	1	4	3	3	2	13	19
American Indian 1/, both sexes	68	-	-	-	-	1	-	-	1	1	1	2
Male	39	-	-	-	-	-	-	-	1	1	-	2
Female	29	-	-	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	244	-	3	-	-	1	-	-	-	1	-	2
Male	136	-	2	-	-	1	-	-	-	1	-	2
Female	108	-	1	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	273	-	-	-	-	-	2	1	1	2	1	5
Male	138	-	-	-	-	-	2	-	1	2	-	4
Female	135	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	224	-	-	-	-	-	-	-	1	1	1	4
Male	115	-	-	-	-	-	-	-	1	1	-	3
Female	109	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	48	-	-	-	-	-	2	1	-	1	-	1
Male	23	-	-	-	-	-	2	-	-	1	-	1
Female	25	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Alaska												
All races, both sexes	12	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	230	-	-	-	-	-	-	-	2	1	-	4
Male	129	-	-	-	-	-	-	-	2	1	-	3
Female	101	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	219	-	-	-	-	-	-	-	2	1	-	4
Male	122	-	-	-	-	-	-	-	2	1	-	3
Female	97	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Arkansas												
All races, both sexes	191	-	-	-	-	-	-	-	-	2	-	2
Male	99	-	-	-	-	-	-	-	-	2	-	2
Female	92	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	169	-	-	-	-	-	-	-	-	2	-	1
Male	89	-	-	-	-	-	-	-	-	2	-	1
Female	80	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	888	1	1	-	1	1	-	2	1	2	2	10
Male	472	-	-	-	1	1	-	1	1	2	2	7
Female	416	1	1	-	-	-	-	1	-	-	-	3
White, both sexes	799	1	-	-	1	-	-	1	1	2	2	8
Male	427	-	-	-	1	-	-	-	1	2	2	6
Female	372	1	-	-	-	-	-	1	-	-	-	2
Black, both sexes	39	-	-	-	-	-	-	1	-	-	-	1
Male	21	-	-	-	-	-	-	1	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	-	1	-	-	1	-	-	-	-	-	1
Male	24	-	-	-	-	1	-	-	-	-	-	1
Female	25	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Colorado												
All races, both sexes	249	1	-	-	-	-	-	-	-	-	-	1
Male	132	-	-	-	-	-	-	-	-	-	-	1
Female	117	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	238	1	-	-	-	-	-	-	-	-	-	-
Male	125	-	-	-	-	-	-	-	-	-	-	-
Female	113	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	327	-	-	-	-	-	-	-	1	-	3	2
Male	163	-	-	-	-	-	-	-	1	-	2	2
Female	164	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	313	-	-	-	-	-	-	-	1	-	3	2
Male	156	-	-	-	-	-	-	-	1	-	2	2
Female	157	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Delaware												
All races, both sexes	50	-	-	-	-	-	-	-	-	1	1	-
Male	26	-	-	-	-	-	-	-	-	1	-	-
Female	24	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	25	-	-	-	1	-	-	-	-	-	-	-
Male	9	-	-	-	1	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	18	-	-	-	1	-	-	-	-	-	-	-
Male	5	-	-	-	1	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Florida												
All races, both sexes	923	-	-	-	-	-	3	3	1	3	8	11
Male	534	-	-	-	-	-	1	3	1	3	5	6
Female	389	-	-	-	-	-	2	-	-	-	3	5
White, both sexes	829	-	-	-	-	-	2	3	1	3	6	10
Male	473	-	-	-	-	-	1	3	1	3	3	6
Female	356	-	-	-	-	-	1	-	-	-	3	4
Black, both sexes	89	-	-	-	-	-	1	-	-	-	2	1
Male	57	-	-	-	-	-	-	-	-	-	2	-
Female	32	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	466	-	2	-	-	2	1	2	3	-	3	4
Male	263	-	1	-	-	1	1	2	-	-	1	3
Female	203	-	1	-	-	1	-	-	3	-	2	1
White, both sexes	367	-	1	-	-	1	1	2	2	-	2	4
Male	217	-	-	-	-	1	1	2	-	-	1	3
Female	150	-	1	-	-	-	-	-	2	-	1	1
Black, both sexes	96	-	-	-	-	1	-	-	1	-	1	-
Male	44	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	1	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Hawaii												
All races, both sexes	123	-	1	-	-	-	-	-	-	1	-	1
Male	69	-	1	-	-	-	-	-	-	1	-	1
Female	54	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	102	-	1	-	-	-	-	-	-	-	-	1
Male	58	-	1	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	59	-	-	-	-	-	-	-	-	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	59	-	-	-	-	-	-	-	-	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Illinois												
All races, both sexes	899	1	-	-	-	-	1	-	2	1	6	2
Male	438	1	-	-	-	-	1	-	1	1	3	1
Female	461	-	-	-	-	-	-	-	1	-	3	1
White, both sexes	838	-	-	-	-	-	1	-	2	1	5	2
Male	409	-	-	-	-	-	1	-	1	1	3	1
Female	429	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	50	1	-	-	-	-	-	-	-	-	1	-
Male	24	1	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	464	-	-	-	-	1	-	-	2	2	3	10
Male	238	-	-	-	-	-	-	-	1	2	1	4
Female	226	-	-	-	-	1	-	-	1	-	2	6
White, both sexes	442	-	-	-	-	1	-	-	2	2	3	8
Male	226	-	-	-	-	-	-	-	1	2	1	3
Female	216	-	-	-	-	1	-	-	1	-	2	5
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Iowa												
All races, both sexes	283	1	-	1	-	-	-	2	-	3	-	1
Male	147	1	-	1	-	-	-	-	-	1	-	-
Female	136	-	-	-	-	-	-	2	-	2	-	1
White, both sexes	277	1	-	1	-	-	-	2	-	3	-	1
Male	143	1	-	1	-	-	-	-	-	1	-	-
Female	134	-	-	-	-	-	-	2	-	2	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	167	-	-	-	-	-	-	-	-	2	-	1
Male	81	-	-	-	-	-	-	-	-	2	-	1
Female	86	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	162	-	-	-	-	-	-	-	-	1	-	1
Male	77	-	-	-	-	-	-	-	-	1	-	1
Female	85	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Kentucky												
All races, both sexes	350	1	-	-	-	-	-	-	2	1	3	3
Male	169	-	-	-	-	-	-	-	1	-	2	2
Female	181	1	-	-	-	-	-	-	1	1	1	1
White, both sexes	327	1	-	-	-	-	-	-	2	1	1	3
Male	157	-	-	-	-	-	-	-	1	-	1	2
Female	170	1	-	-	-	-	-	-	1	1	-	1
Black, both sexes	22	-	-	-	-	-	-	-	-	-	2	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	209	-	-	-	1	-	-	-	1	2	2	1
Male	103	-	-	-	1	-	-	-	1	2	1	1
Female	106	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	155	-	-	-	1	-	-	-	1	1	1	-
Male	79	-	-	-	1	-	-	-	1	1	-	-
Female	76	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	53	-	-	-	-	-	-	-	-	1	1	1
Male	24	-	-	-	-	-	-	-	-	1	1	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Maine												
All races, both sexes	91	-	-	-	-	-	-	1	-	1	-	-
Male	47	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	91	-	-	-	-	-	-	1	-	1	-	-
Male	47	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	359	-	-	-	-	-	1	1	1	1	4	5
Male	186	-	-	-	-	-	-	1	1	1	1	3
Female	173	-	-	-	-	-	1	-	-	-	3	2
White, both sexes	280	-	-	-	-	-	1	1	-	1	3	3
Male	149	-	-	-	-	-	-	1	-	1	1	2
Female	131	-	-	-	-	-	1	-	-	-	2	1
Black, both sexes	73	-	-	-	-	-	-	-	1	-	1	2
Male	34	-	-	-	-	-	-	-	1	-	-	1
Female	39	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Massachusetts												
All races, both sexes	529	-	1	-	-	-	-	-	-	-	1	3
Male	278	-	1	-	-	-	-	-	-	-	1	1
Female	251	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	510	-	1	-	-	-	-	-	-	-	1	3
Male	268	-	1	-	-	-	-	-	-	-	1	1
Female	242	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	533	-	-	-	-	-	1	-	3	2	6	5
Male	284	-	-	-	-	-	-	-	2	1	5	4
Female	249	-	-	-	-	-	1	-	1	1	1	1
White, both sexes	486	-	-	-	-	-	-	-	3	-	6	3
Male	255	-	-	-	-	-	-	-	2	-	5	2
Female	231	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	44	-	-	-	-	-	1	-	-	1	-	2
Male	28	-	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Minnesota												
All races, both sexes	228	-	-	-	-	-	-	1	-	-	1	1
Male	127	-	-	-	-	-	-	1	-	-	1	1
Female	101	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	223	-	-	-	-	-	-	1	-	-	1	1
Male	123	-	-	-	-	-	-	1	-	-	1	1
Female	100	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	186	2	-	1	-	-	-	3	1	-	3	3
Male	101	1	-	1	-	-	-	2	1	-	2	1
Female	85	1	-	-	-	-	-	1	-	-	1	2
White, both sexes	144	-	-	-	-	-	-	2	1	-	2	2
Male	82	-	-	-	-	-	-	1	1	-	2	1
Female	62	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	41	2	-	1	-	-	-	1	-	-	1	1
Male	18	1	-	1	-	-	-	1	-	-	-	-
Female	23	1	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Missouri												
All races, both sexes	418	-	1	1	1	-	-	-	1	-	2	4
Male	218	-	-	-	1	-	-	-	-	-	1	3
Female	200	-	1	1	-	-	-	-	1	-	1	1
White, both sexes	387	-	1	1	-	-	-	-	1	-	1	3
Male	201	-	-	-	-	-	-	-	-	-	-	2
Female	186	-	1	1	-	-	-	-	1	-	1	1
Black, both sexes	29	-	-	-	1	-	-	-	-	-	1	1
Male	16	-	-	-	1	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	72	-	1	-	-	-	-	-	1	-	1	-
Male	36	-	-	-	-	-	-	-	1	-	-	-
Female	36	-	1	-	-	-	-	-	-	-	1	-
White, both sexes	69	-	1	-	-	-	-	-	-	-	1	-
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	1	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Nebraska												
All races, both sexes	111	-	-	-	-	1	-	-	1	1	1	1
Male	58	-	-	-	-	-	-	-	1	1	-	1
Female	53	-	-	-	-	1	-	-	-	-	1	-
White, both sexes	106	-	-	-	-	1	-	-	-	1	1	1
Male	55	-	-	-	-	-	-	-	-	1	-	1
Female	51	-	-	-	-	1	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	115	-	-	-	-	-	1	-	-	-	-	3
Male	64	-	-	-	-	-	1	-	-	-	-	2
Female	51	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	105	-	-	-	-	-	1	-	-	-	-	2
Male	60	-	-	-	-	-	1	-	-	-	-	2
Female	45	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
New Hampshire												
All races, both sexes	62	-	-	-	-	-	-	1	-	1	-	-
Male	27	-	-	-	-	-	-	1	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	62	-	-	-	-	-	-	1	-	1	-	-
Male	27	-	-	-	-	-	-	1	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	513	-	-	1	-	1	-	2	-	1	5	7
Male	272	-	-	1	-	-	-	1	-	1	4	3
Female	241	-	-	-	-	1	-	1	-	-	1	4
White, both sexes	454	-	-	1	-	1	-	2	-	-	5	3
Male	244	-	-	1	-	-	-	1	-	-	4	2
Female	210	-	-	-	-	1	-	1	-	-	1	1
Black, both sexes	55	-	-	-	-	-	-	-	-	-	-	4
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
New Mexico												
All races, both sexes	70	-	-	-	-	-	-	-	1	-	1	2
Male	38	-	-	-	-	-	-	-	1	-	-	2
Female	32	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	61	-	-	-	-	-	-	-	1	-	1	2
Male	36	-	-	-	-	-	-	-	1	-	-	2
Female	25	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	711	2	-	1	-	1	-	-	4	-	4	11
Male	387	1	-	-	-	1	-	-	3	-	3	6
Female	324	1	-	1	-	-	-	-	1	-	1	5
White, both sexes	665	1	-	1	-	1	-	-	2	-	3	11
Male	360	-	-	-	-	1	-	-	2	-	2	6
Female	305	1	-	1	-	-	-	-	-	-	1	5
Black, both sexes	39	1	-	-	-	-	-	-	2	-	1	-
Male	23	1	-	-	-	-	-	-	1	-	1	-
Female	16	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
North Carolina												
All races, both sexes	656	-	-	-	-	1	1	-	3	3	9	8
Male	344	-	-	-	-	-	-	-	2	2	6	3
Female	312	-	-	-	-	1	1	-	1	1	3	5
White, both sexes	550	-	-	-	-	1	-	-	2	-	5	6
Male	282	-	-	-	-	-	-	-	1	-	3	3
Female	268	-	-	-	-	1	-	-	1	-	2	3
Black, both sexes	103	-	-	-	-	-	1	-	1	3	4	2
Male	59	-	-	-	-	-	-	-	1	2	3	-
Female	44	-	-	-	-	-	1	-	-	1	1	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	1	1
Male	27	-	-	-	-	-	-	-	-	-	1	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	48	-	-	-	-	-	-	-	-	-	1	-
Male	26	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Ohio												
All races, both sexes	805	-	1	-	-	-	-	1	2	3	1	8
Male	424	-	-	-	-	-	-	1	2	-	1	6
Female	381	-	1	-	-	-	-	-	-	3	-	2
White, both sexes	731	-	1	-	-	-	-	-	1	3	1	7
Male	384	-	-	-	-	-	-	-	1	-	1	6
Female	347	-	1	-	-	-	-	-	-	3	-	1
Black, both sexes	73	-	-	-	-	-	-	1	1	-	-	1
Male	40	-	-	-	-	-	-	1	1	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	200	-	-	-	-	-	-	-	2	-	3	1
Male	89	-	-	-	-	-	-	-	1	-	1	-
Female	111	-	-	-	-	-	-	-	1	-	2	1
White, both sexes	184	-	-	-	-	-	-	-	1	-	3	1
Male	83	-	-	-	-	-	-	-	-	-	1	-
Female	101	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Oregon												
All races, both sexes	166	-	-	-	-	-	2	-	-	-	-	2
Male	85	-	-	-	-	-	1	-	-	-	-	1
Female	81	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	160	-	-	-	-	-	2	-	-	-	-	2
Male	80	-	-	-	-	-	1	-	-	-	-	1
Female	80	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,129	1	-	1	-	1	2	2	1	2	9	17
Male	608	1	-	1	-	1	1	1	1	-	6	8
Female	521	-	-	-	-	-	1	1	-	2	3	9
White, both sexes	1,021	1	-	1	-	-	2	1	1	2	7	13
Male	559	1	-	1	-	-	1	1	1	-	5	7
Female	462	-	-	-	-	-	1	-	-	2	2	6
Black, both sexes	101	-	-	-	-	1	-	1	-	-	2	4
Male	45	-	-	-	-	1	-	-	-	-	1	1
Female	56	-	-	-	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Rhode Island												
All races, both sexes	86	-	-	-	-	-	-	1	-	-	-	1
Male	49	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	83	-	-	-	-	-	-	1	-	-	-	1
Male	47	-	-	-	-	-	-	-	-	-	-	1
Female	36	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	276	-	-	-	-	-	-	2	2	2	1	4
Male	147	-	-	-	-	-	-	1	-	1	-	2
Female	129	-	-	-	-	-	-	1	2	1	1	2
White, both sexes	206	-	-	-	-	-	-	1	1	1	-	3
Male	107	-	-	-	-	-	-	-	-	-	-	2
Female	99	-	-	-	-	-	-	1	1	1	-	1
Black, both sexes	70	-	-	-	-	-	-	1	1	1	1	1
Male	40	-	-	-	-	-	-	1	-	1	-	-
Female	30	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
South Dakota												
All races, both sexes	44	-	-	-	-	-	-	-	-	1	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	1	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	453	-	-	-	-	-	1	2	1	-	4	8
Male	230	-	-	-	-	-	-	1	1	-	3	5
Female	223	-	-	-	-	-	1	1	-	-	1	3
White, both sexes	411	-	-	-	-	-	-	-	1	-	3	7
Male	207	-	-	-	-	-	-	-	1	-	3	4
Female	204	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	41	-	-	-	-	-	1	2	-	-	1	1
Male	22	-	-	-	-	-	-	1	-	-	-	1
Female	19	-	-	-	-	-	1	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Texas												
All races, both sexes	1,197	1	1	-	2	-	1	3	1	6	14	13
Male	641	-	-	-	2	-	1	3	1	4	10	8
Female	556	1	1	-	-	-	-	-	-	2	4	5
White, both sexes	1,091	1	1	-	2	-	1	1	1	6	10	12
Male	576	-	-	-	2	-	1	1	1	4	7	7
Female	515	1	1	-	-	-	-	-	-	2	3	5
Black, both sexes	100	-	-	-	-	-	-	2	-	-	3	1
Male	60	-	-	-	-	-	-	2	-	-	3	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	94	-	-	-	1	-	1	1	-	-	-	-
Male	55	-	-	-	1	-	1	1	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	93	-	-	-	1	-	1	1	-	-	-	-
Male	54	-	-	-	1	-	1	1	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Vermont												
All races, both sexes	37	-	-	-	-	1	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	37	-	-	-	-	1	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	564	-	-	-	-	-	1	-	1	2	3	8
Male	297	-	-	-	-	-	1	-	1	2	2	4
Female	267	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	442	-	-	-	-	-	-	-	1	1	1	5
Male	231	-	-	-	-	-	-	-	1	1	1	3
Female	211	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	112	-	-	-	-	-	1	-	-	1	2	3
Male	60	-	-	-	-	-	1	-	-	1	1	1
Female	52	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Washington												
All races, both sexes	438	-	-	-	-	1	-	-	1	3	4	4
Male	218	-	-	-	-	-	-	-	1	1	2	2
Female	220	-	-	-	-	1	-	-	-	2	2	2
White, both sexes	405	-	-	-	-	-	-	-	1	3	3	4
Male	199	-	-	-	-	-	-	-	1	1	2	2
Female	206	-	-	-	-	-	-	-	-	2	1	2
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	7	-	-	-	-	1	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	123	-	-	-	-	-	-	1	-	-	2	-
Male	64	-	-	-	-	-	-	1	-	-	-	-
Female	59	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	120	-	-	-	-	-	-	-	-	-	1	-
Male	62	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	1	-	-	1	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Wisconsin												
All races, both sexes	354	-	-	-	-	-	-	1	-	-	2	5
Male	210	-	-	-	-	-	-	1	-	-	2	2
Female	144	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	343	-	-	-	-	-	-	1	-	-	2	3
Male	204	-	-	-	-	-	-	1	-	-	2	1
Female	139	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	29	-	-	-	-	-	-	1	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	1	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
United States												
All races, both sexes	253	359	496	658	1,092	1,956	3,328	4,037	2,903	1,091	223	-
Male	163	222	315	420	672	1,162	1,867	2,056	1,298	376	66	-
Female	90	137	181	238	420	794	1,461	1,981	1,605	715	157	-
White, both sexes	212	295	426	551	913	1,713	3,038	3,748	2,668	1,010	190	-
Male	136	184	269	348	568	1,012	1,714	1,912	1,202	354	56	-
Female	76	111	157	203	345	701	1,324	1,836	1,466	656	134	-
Black, both sexes	40	56	63	92	158	196	233	233	174	64	24	-
Male	26	32	43	64	91	123	124	110	65	13	6	-
Female	14	24	20	28	67	73	109	123	109	51	18	-
American Indian 1/, both sexes	1	4	3	12	3	14	2	10	12	-	1	-
Male	1	3	1	7	2	7	1	7	5	-	1	-
Female	-	1	2	5	1	7	1	3	7	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	4	3	18	33	55	46	49	17	8	-
Male	-	3	2	1	11	20	28	27	26	9	3	-
Female	-	1	2	2	7	13	27	19	23	8	5	-
Alabama												
All races, both sexes	5	12	1	10	19	30	50	74	46	9	5	-
Male	3	8	1	6	10	18	25	42	11	4	1	-
Female	2	4	-	4	9	12	25	32	35	5	4	-
White, both sexes	3	10	1	8	15	23	42	64	40	7	4	-
Male	2	7	1	5	8	12	23	37	10	4	1	-
Female	1	3	-	3	7	11	19	27	30	3	3	-
Black, both sexes	2	2	-	1	4	7	8	10	6	2	1	-
Male	1	1	-	1	2	6	2	5	1	-	-	-
Female	1	1	-	-	2	1	6	5	5	2	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Alaska												
All races, both sexes	-	-	1	-	1	2	4	1	2	-	-	-
Male	-	-	-	-	-	2	3	-	-	-	-	-
Female	-	-	1	-	1	-	1	1	2	-	-	-
White, both sexes	-	-	1	-	1	2	3	1	-	-	-	-
Male	-	-	-	-	-	2	3	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	4	8	12	18	32	54	51	26	11	2	-
Male	3	2	4	5	13	12	34	26	16	6	2	-
Female	2	2	4	7	5	20	20	25	10	5	-	-
White, both sexes	5	2	8	12	18	29	54	48	25	10	1	-
Male	3	1	4	5	13	10	34	24	15	6	1	-
Female	2	1	4	7	5	19	20	24	10	4	-	-
Black, both sexes	-	1	-	-	-	1	-	-	-	1	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	2	-	2	1	-	1	-
Male	-	-	-	-	-	1	-	2	1	-	1	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Arkansas												
All races, both sexes	4	6	6	3	7	24	40	45	34	15	3	-
Male	2	2	5	2	2	14	26	25	12	5	-	-
Female	2	4	1	1	5	10	14	20	22	10	3	-
White, both sexes	2	4	4	2	5	21	37	43	32	14	2	-
Male	1	2	3	1	1	13	24	24	12	5	-	-
Female	1	2	1	1	4	8	13	19	20	9	2	-
Black, both sexes	2	2	2	1	2	3	3	2	2	1	1	-
Male	1	-	2	1	1	1	2	1	-	-	-	-
Female	1	2	-	-	1	2	1	1	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	13	17	32	37	48	112	186	203	136	66	17	-
Male	8	7	20	22	28	60	97	110	73	25	7	-
Female	5	10	12	15	20	52	89	93	63	41	10	-
White, both sexes	12	15	27	33	38	100	172	188	121	63	14	-
Male	7	7	17	21	22	54	90	103	64	24	6	-
Female	5	8	10	12	16	46	82	85	57	39	8	-
Black, both sexes	1	2	4	3	8	4	5	4	6	-	-	-
Male	1	-	3	1	5	2	4	1	3	-	-	-
Female	-	2	1	2	3	2	1	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	8	9	11	9	3	3	-
Male	-	-	-	-	1	4	3	6	6	1	1	-
Female	-	-	1	-	1	4	6	5	3	2	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Colorado												
All races, both sexes	5	5	7	4	14	34	43	66	52	14	3	-
Male	4	3	4	1	9	22	26	35	21	4	2	-
Female	1	2	3	3	5	12	17	31	31	10	1	-
White, both sexes	4	5	7	4	10	33	41	64	52	14	3	-
Male	3	3	4	1	6	22	25	34	21	4	2	-
Female	1	2	3	3	4	11	16	30	31	10	1	-
Black, both sexes	-	-	-	-	2	1	1	2	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	1	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	1	4	10	10	18	24	73	82	68	29	2	-
Male	-	2	4	6	7	15	43	39	32	10	-	-
Female	1	2	6	4	11	9	30	43	36	19	2	-
White, both sexes	1	4	9	9	15	22	70	80	66	29	2	-
Male	-	2	4	5	6	13	40	39	32	10	-	-
Female	1	2	5	4	9	9	30	41	34	19	2	-
Black, both sexes	-	-	1	1	3	2	3	2	2	-	-	-
Male	-	-	-	1	1	2	3	-	-	-	-	-
Female	-	-	1	-	2	-	-	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Delaware												
All races, both sexes	1	1	3	1	1	5	8	15	11	2	-	-
Male	1	1	3	-	1	5	5	5	4	-	-	-
Female	-	-	-	1	-	-	3	10	7	2	-	-
White, both sexes	1	1	1	1	1	4	8	14	10	2	-	-
Male	1	1	1	-	1	4	5	4	3	-	-	-
Female	-	-	-	1	-	-	3	10	7	2	-	-
Black, both sexes	-	-	2	-	-	1	-	1	1	-	-	-
Male	-	-	2	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	-	1	-	3	4	8	4	3	-	-
Male	-	1	-	-	-	1	2	2	-	2	-	-
Female	-	-	-	1	-	2	2	6	4	1	-	-
White, both sexes	-	-	-	-	-	1	1	1	1	2	-	-
Male	-	-	-	-	-	1	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	-	1	-	1	-	2	2	7	3	1	-	-
Male	-	1	-	-	-	-	-	2	-	1	-	-
Female	-	-	-	1	-	2	2	5	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Florida												
All races, both sexes	9	24	21	34	52	102	197	221	161	62	11	-
Male	4	21	14	21	32	75	121	122	76	26	3	-
Female	5	3	7	13	20	27	76	99	85	36	8	-
White, both sexes	7	17	18	23	45	91	179	210	148	57	9	-
Male	3	14	13	12	27	66	110	115	69	24	3	-
Female	4	3	5	11	18	25	69	95	79	33	6	-
Black, both sexes	2	7	3	11	7	9	17	10	12	5	2	-
Male	1	7	1	9	5	7	11	6	6	2	-	-
Female	1	-	2	2	2	2	6	4	6	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	9	7	16	24	32	58	95	104	70	27	7	-
Male	5	4	11	20	22	39	58	49	33	10	3	-
Female	4	3	5	4	10	19	37	55	37	17	4	-
White, both sexes	7	2	12	20	23	45	75	87	59	18	6	-
Male	4	2	9	16	17	29	48	44	30	7	3	-
Female	3	-	3	4	6	16	27	43	29	11	3	-
Black, both sexes	2	5	4	4	9	13	19	17	10	9	1	-
Male	1	2	2	4	5	10	10	5	2	3	-	-
Female	1	3	2	-	4	3	9	12	8	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Hawaii												
All races, both sexes	-	3	3	4	5	11	28	20	31	13	2	-
Male	-	3	3	2	4	6	16	9	14	7	2	-
Female	-	-	-	2	1	5	12	11	17	6	-	-
White, both sexes	-	-	1	1	-	2	5	4	7	-	-	-
Male	-	-	1	1	-	1	2	2	3	-	-	-
Female	-	-	-	-	-	1	3	2	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	2	3	5	9	23	16	24	13	2	-
Male	-	3	2	1	4	5	14	7	11	7	2	-
Female	-	-	-	2	1	4	9	9	13	6	-	-
Idaho												
All races, both sexes	3	5	4	1	3	7	13	12	9	1	-	-
Male	1	4	2	1	3	4	4	9	4	-	-	-
Female	2	1	2	-	-	3	9	3	5	1	-	-
White, both sexes	3	5	4	1	3	7	13	12	9	1	-	-
Male	1	4	2	1	3	4	4	9	4	-	-	-
Female	2	1	2	-	-	3	9	3	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Illinois												
All races, both sexes	13	16	29	26	52	95	177	238	170	58	12	-
Male	10	10	20	13	28	47	88	109	83	21	1	-
Female	3	6	9	13	24	48	89	129	87	37	11	-
White, both sexes	11	15	26	25	41	83	166	227	167	56	10	-
Male	9	10	18	12	21	40	84	105	82	20	1	-
Female	2	5	8	13	20	43	82	122	85	36	9	-
Black, both sexes	2	1	2	1	10	9	7	10	3	2	1	-
Male	1	-	2	1	6	4	3	4	1	1	-	-
Female	1	1	-	-	4	5	4	6	2	1	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	3	4	1	-	-	1	-
Male	-	-	-	-	1	3	1	-	-	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	1	-
Indiana												
All races, both sexes	9	12	14	22	36	52	83	112	73	27	6	-
Male	6	8	10	11	22	28	49	57	30	7	2	-
Female	3	4	4	11	14	24	34	55	43	20	4	-
White, both sexes	9	11	13	20	35	49	80	109	70	26	4	-
Male	6	7	10	11	21	26	47	56	28	6	1	-
Female	3	4	3	9	14	23	33	53	42	20	3	-
Black, both sexes	-	1	1	2	1	3	3	3	3	1	2	-
Male	-	1	-	-	1	2	2	1	2	1	1	-
Female	-	-	1	2	-	1	1	2	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Iowa												
All races, both sexes	2	1	7	9	20	27	41	79	64	23	2	-
Male	1	-	5	7	10	13	25	38	31	14	-	-
Female	1	1	2	2	10	14	16	41	33	9	2	-
White, both sexes	2	1	7	7	20	26	40	79	62	23	2	-
Male	1	-	5	6	10	12	24	38	30	14	-	-
Female	1	1	2	1	10	14	16	41	32	9	2	-
Black, both sexes	-	-	-	1	-	1	1	-	2	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	4	-	4	8	3	15	30	41	35	20	4	-
Male	2	-	1	7	3	8	12	22	18	5	-	-
Female	2	-	3	1	-	7	18	19	17	15	4	-
White, both sexes	4	-	4	8	3	14	29	40	34	20	4	-
Male	2	-	1	7	3	7	12	21	17	5	-	-
Female	2	-	3	1	-	7	17	19	17	15	4	-
Black, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Kentucky												
All races, both sexes	2	4	11	16	19	49	68	79	71	19	2	-
Male	1	4	7	10	11	34	31	31	33	1	1	-
Female	1	-	4	6	8	15	37	48	38	18	1	-
White, both sexes	2	4	10	13	19	45	66	75	67	16	2	-
Male	1	4	6	8	11	31	30	29	31	1	1	-
Female	1	-	4	5	8	14	36	46	36	15	1	-
Black, both sexes	-	-	1	3	-	4	1	4	4	3	-	-
Male	-	-	1	2	-	3	1	2	2	-	-	-
Female	-	-	-	1	-	1	-	2	2	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Louisiana												
All races, both sexes	3	4	12	10	18	26	44	39	35	9	2	-
Male	3	3	5	3	11	18	22	14	13	5	-	-
Female	-	1	7	7	7	8	22	25	22	4	2	-
White, both sexes	3	3	9	9	10	18	35	29	26	8	1	-
Male	3	2	4	3	6	14	19	11	10	4	-	-
Female	-	1	5	6	4	4	16	18	16	4	1	-
Black, both sexes	-	1	3	1	8	8	9	9	9	1	1	-
Male	-	1	1	-	5	4	3	3	3	1	-	-
Female	-	-	2	1	3	4	6	6	6	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Maine												
All races, both sexes	-	1	1	2	6	9	18	25	16	10	1	-
Male	-	1	1	2	4	6	9	14	7	2	1	-
Female	-	-	-	-	2	3	9	11	9	8	-	-
White, both sexes	-	1	1	2	6	9	18	25	16	10	1	-
Male	-	1	1	2	4	6	9	14	7	2	1	-
Female	-	-	-	-	2	3	9	11	9	8	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	9	10	14	10	33	43	63	86	54	20	4	-
Male	4	3	11	6	17	27	38	43	26	4	-	-
Female	5	7	3	4	16	16	25	43	28	16	4	-
White, both sexes	5	6	9	4	19	32	51	76	47	19	3	-
Male	1	3	8	2	13	21	29	38	25	4	-	-
Female	4	3	1	2	6	11	22	38	22	15	3	-
Black, both sexes	4	4	5	6	13	8	10	10	7	1	1	-
Male	3	-	3	4	4	5	7	5	1	-	-	-
Female	1	4	2	2	9	3	3	5	6	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	3	2	-	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Massachusetts												
All races, both sexes	5	5	7	13	30	72	116	119	103	43	11	-
Male	2	3	7	7	13	52	65	60	44	17	5	-
Female	3	2	-	6	17	20	51	59	59	26	6	-
White, both sexes	5	5	7	12	29	70	112	114	99	42	10	-
Male	2	3	7	6	12	51	62	57	43	17	5	-
Female	3	2	-	6	17	19	50	57	56	25	5	-
Black, both sexes	-	-	-	1	-	1	2	3	4	1	1	-
Male	-	-	-	1	-	-	2	1	1	-	-	-
Female	-	-	-	-	-	1	-	2	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	2	2	-	-	-	-
Male	-	-	-	-	1	1	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Michigan												
All races, both sexes	10	11	13	25	29	73	101	125	93	30	6	-
Male	7	7	11	15	20	41	58	66	34	11	2	-
Female	3	4	2	10	9	32	43	59	59	19	4	-
White, both sexes	7	9	11	24	23	68	92	117	87	30	6	-
Male	5	5	10	15	16	37	52	62	31	11	2	-
Female	2	4	1	9	7	31	40	55	56	19	4	-
Black, both sexes	3	2	2	1	6	5	9	8	4	-	-	-
Male	2	2	1	-	4	4	6	4	3	-	-	-
Female	1	-	1	1	2	1	3	4	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Minnesota												
All races, both sexes	4	3	9	6	11	34	31	55	51	16	5	-
Male	4	2	6	6	8	19	17	32	24	5	1	-
Female	-	1	3	-	3	15	14	23	27	11	4	-
White, both sexes	4	3	9	5	9	33	30	55	51	16	5	-
Male	4	2	6	5	6	19	16	32	24	5	1	-
Female	-	1	3	-	3	14	14	23	27	11	4	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	6	7	7	11	25	32	43	23	13	2	-
Male	3	3	6	6	4	18	21	21	9	2	-	-
Female	1	3	1	1	7	7	11	22	14	11	2	-
White, both sexes	3	4	5	5	8	19	26	36	19	10	2	-
Male	2	1	4	4	3	14	20	18	9	2	-	-
Female	1	3	1	1	5	5	6	18	10	8	2	-
Black, both sexes	1	2	2	2	3	5	6	7	4	3	-	-
Male	1	2	2	2	1	3	1	3	-	-	-	-
Female	-	-	-	-	2	2	5	4	4	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Missouri												
All races, both sexes	11	9	9	13	26	53	82	100	78	21	6	-
Male	5	2	6	7	18	33	43	52	39	6	2	-
Female	6	7	3	6	8	20	39	48	39	15	4	-
White, both sexes	10	8	9	11	24	48	77	96	70	21	6	-
Male	4	2	6	5	17	29	42	50	36	6	2	-
Female	6	6	3	6	7	19	35	46	34	15	4	-
Black, both sexes	1	1	-	2	2	4	5	4	7	-	-	-
Male	1	-	-	2	1	3	1	2	3	-	-	-
Female	-	1	-	-	1	1	4	2	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Montana												
All races, both sexes	-	1	2	4	5	8	13	9	20	7	-	-
Male	-	1	2	3	4	4	9	2	7	3	-	-
Female	-	-	-	1	1	4	4	7	13	4	-	-
White, both sexes	-	1	2	3	4	8	13	9	20	7	-	-
Male	-	1	2	2	3	4	9	2	7	3	-	-
Female	-	-	-	1	1	4	4	7	13	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Nebraska												
All races, both sexes	-	2	-	6	1	13	17	35	19	11	2	-
Male	-	2	-	5	-	9	11	12	11	4	1	-
Female	-	-	-	1	1	4	6	23	8	7	1	-
White, both sexes	-	2	-	4	1	11	17	35	19	11	2	-
Male	-	2	-	4	-	8	11	12	11	4	1	-
Female	-	-	-	-	1	3	6	23	8	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	2	1	10	10	14	30	17	17	6	-	-
Male	3	2	1	6	5	10	20	7	5	2	-	-
Female	1	-	-	4	5	4	10	10	12	4	-	-
White, both sexes	4	2	1	9	7	13	28	17	15	6	-	-
Male	3	2	1	5	4	9	19	7	5	2	-	-
Female	1	-	-	4	3	4	9	10	10	4	-	-
Black, both sexes	-	-	-	-	2	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
New Hampshire												
All races, both sexes	1	1	3	3	3	1	15	16	12	5	-	-
Male	1	-	1	-	3	-	5	9	7	-	-	-
Female	-	1	2	3	-	1	10	7	5	5	-	-
White, both sexes	1	1	3	3	3	1	15	16	12	5	-	-
Male	1	-	1	-	3	-	5	9	7	-	-	-
Female	-	1	2	3	-	1	10	7	5	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	9	13	12	11	29	58	101	146	86	27	4	-
Male	5	6	9	7	20	34	59	77	37	8	-	-
Female	4	7	3	4	9	24	42	69	49	19	4	-
White, both sexes	4	12	9	8	25	48	94	132	81	25	4	-
Male	2	6	7	6	19	26	56	70	36	8	-	-
Female	2	6	2	2	6	22	38	62	45	17	4	-
Black, both sexes	5	1	3	3	4	9	7	12	5	2	-	-
Male	3	-	2	1	1	7	3	5	1	-	-	-
Female	2	1	1	2	3	2	4	7	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
New Mexico												
All races, both sexes	-	2	3	4	7	13	12	12	8	3	2	-
Male	-	1	3	3	2	4	8	8	3	2	1	-
Female	-	1	-	1	5	9	4	4	5	1	1	-
White, both sexes	-	1	3	4	5	10	12	11	6	3	2	-
Male	-	-	3	3	2	4	8	8	2	2	1	-
Female	-	1	-	1	3	6	4	3	4	1	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	3	-	1	2	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	3	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	11	15	24	25	42	67	140	197	106	52	9	-
Male	7	8	13	19	25	45	81	111	46	15	3	-
Female	4	7	11	6	17	22	59	86	60	37	6	-
White, both sexes	10	13	21	24	38	66	132	187	101	47	7	-
Male	7	6	11	18	23	44	77	104	42	15	2	-
Female	3	7	10	6	15	22	55	83	59	32	5	-
Black, both sexes	1	2	3	1	4	1	7	6	4	5	1	-
Male	-	2	2	1	2	1	4	4	3	-	1	-
Female	1	-	1	-	2	-	3	2	1	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	4	1	-	1	-
Male	-	-	-	-	-	-	-	3	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
North Carolina												
All races, both sexes	9	16	31	41	50	73	129	140	98	39	5	-
Male	5	11	19	25	31	42	71	66	52	8	1	-
Female	4	5	12	16	19	31	58	74	46	31	4	-
White, both sexes	8	9	25	34	38	59	117	124	83	35	4	-
Male	4	5	14	23	22	35	66	55	43	8	-	-
Female	4	4	11	11	16	24	51	69	40	27	4	-
Black, both sexes	1	6	5	7	12	13	12	16	15	4	1	-
Male	1	5	4	2	9	6	5	11	9	-	1	-
Female	-	1	1	5	3	7	7	5	6	4	-	-
American Indian 1/, both sexes	-	1	1	-	-	1	-	-	-	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	1	2	2	7	11	13	6	4	1	-
Male	-	-	1	1	2	4	7	5	3	2	-	-
Female	-	1	-	1	-	3	4	8	3	2	1	-
White, both sexes	-	1	1	1	2	7	11	13	6	4	1	-
Male	-	-	1	1	2	4	7	5	3	2	-	-
Female	-	1	-	-	-	3	4	8	3	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Ohio												
All races, both sexes	12	10	19	31	62	93	143	198	156	59	6	-
Male	9	7	14	23	43	65	84	89	59	20	1	-
Female	3	3	5	8	19	28	59	109	97	39	5	-
White, both sexes	11	8	16	28	54	80	132	183	146	55	5	-
Male	8	6	11	20	38	56	77	86	55	19	-	-
Female	3	2	5	8	16	24	55	97	91	36	5	-
Black, both sexes	1	2	3	3	7	13	11	15	10	4	1	-
Male	1	1	3	3	5	9	7	3	4	1	1	-
Female	-	1	-	-	2	4	4	12	6	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	2	6	11	13	26	36	48	37	10	5	-
Male	-	2	2	6	5	15	16	23	14	2	2	-
Female	-	-	4	5	8	11	20	25	23	8	3	-
White, both sexes	-	2	5	8	13	24	35	46	32	10	4	-
Male	-	2	2	4	5	15	15	22	13	2	2	-
Female	-	-	3	4	8	9	20	24	19	8	2	-
Black, both sexes	-	-	-	1	-	1	1	-	1	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	1	2	-	1	-	2	4	-	-	-
Male	-	-	-	2	-	-	-	1	1	-	-	-
Female	-	-	1	-	-	1	-	1	3	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Oregon												
All races, both sexes	7	2	3	7	16	13	23	36	40	12	3	-
Male	5	1	1	3	7	7	16	21	16	6	-	-
Female	2	1	2	4	9	6	7	15	24	6	3	-
White, both sexes	6	2	3	7	16	12	23	34	38	12	3	-
Male	4	1	1	3	7	6	16	20	14	6	-	-
Female	2	1	2	4	9	6	7	14	24	6	3	-
Black, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	1	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	17	28	24	34	72	138	252	257	199	52	20	-
Male	14	20	15	19	42	75	146	146	89	14	8	-
Female	3	8	9	15	30	63	106	111	110	38	12	-
White, both sexes	15	27	22	26	62	120	229	245	179	50	18	-
Male	12	19	13	15	36	68	136	140	82	14	7	-
Female	3	8	9	11	26	52	93	105	97	36	11	-
Black, both sexes	2	1	2	8	9	18	21	11	17	2	2	-
Male	2	1	2	4	5	7	9	6	5	-	1	-
Female	-	-	-	4	4	11	12	5	12	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	2	1	3	-	-	-
Male	-	-	-	-	1	-	1	-	2	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Rhode Island												
All races, both sexes	-	2	2	4	6	11	16	26	12	4	1	-
Male	-	1	1	3	5	7	10	12	8	-	1	-
Female	-	1	1	1	1	4	6	14	4	4	-	-
White, both sexes	-	2	2	4	6	9	16	25	12	4	1	-
Male	-	1	1	3	5	5	10	12	8	-	1	-
Female	-	1	1	1	1	4	6	13	4	4	-	-
Black, both sexes	-	-	-	-	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	9	7	13	30	36	65	62	31	7	4	-
Male	1	5	5	12	24	24	31	26	11	2	2	-
Female	-	4	2	1	6	12	34	36	20	5	2	-
White, both sexes	1	5	4	8	19	25	52	50	26	7	3	-
Male	1	2	4	8	16	17	26	18	9	2	2	-
Female	-	3	-	-	3	8	26	32	17	5	1	-
Black, both sexes	-	4	3	5	11	11	13	12	5	-	1	-
Male	-	3	1	4	8	7	5	8	2	-	-	-
Female	-	1	2	1	3	4	8	4	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
South Dakota												
All races, both sexes	-	-	-	1	2	6	4	16	6	8	-	-
Male	-	-	-	-	1	5	3	9	3	3	-	-
Female	-	-	-	1	1	1	1	7	3	5	-	-
White, both sexes	-	-	-	1	2	5	4	16	5	8	-	-
Male	-	-	-	-	1	4	3	9	3	3	-	-
Female	-	-	-	1	1	1	1	7	2	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	14	18	21	29	54	82	121	64	21	7	-
Male	4	7	10	13	21	30	45	52	28	9	1	-
Female	2	7	8	8	8	24	37	69	36	12	6	-
White, both sexes	4	13	15	18	25	50	78	114	61	17	5	-
Male	2	7	7	10	18	28	43	48	26	9	1	-
Female	2	6	8	8	7	22	35	66	35	8	4	-
Black, both sexes	2	1	3	3	4	4	4	7	2	4	2	-
Male	2	-	3	3	3	2	2	4	1	-	-	-
Female	-	1	-	-	1	2	2	3	1	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Texas												
All races, both sexes	20	28	38	52	90	122	261	253	177	94	20	-
Male	14	19	23	36	57	60	147	137	83	31	5	-
Female	6	9	15	16	33	62	114	116	94	63	15	-
White, both sexes	17	25	34	41	82	114	237	234	168	86	18	-
Male	13	19	21	27	53	55	129	125	78	29	4	-
Female	4	6	13	14	29	59	108	109	90	57	14	-
Black, both sexes	3	3	4	11	8	8	24	14	9	8	2	-
Male	1	-	2	9	4	5	18	7	5	2	1	-
Female	2	3	2	2	4	3	6	7	4	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	4	-	-	-	-
Male	-	-	-	-	-	-	-	4	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	3	8	3	-	12	10	16	22	13	4	-	-
Male	2	4	1	-	7	3	8	18	7	2	-	-
Female	1	4	2	-	5	7	8	4	6	2	-	-
White, both sexes	3	8	3	-	12	10	16	21	13	4	-	-
Male	2	4	1	-	7	3	8	17	7	2	-	-
Female	1	4	2	-	5	7	8	4	6	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Vermont												
All races, both sexes	-	-	1	-	-	3	9	8	8	6	1	-
Male	-	-	-	-	-	2	4	5	3	2	1	-
Female	-	-	1	-	-	1	5	3	5	4	-	-
White, both sexes	-	-	1	-	-	3	9	8	8	6	1	-
Male	-	-	-	-	-	2	4	5	3	2	1	-
Female	-	-	1	-	-	1	5	3	5	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	10	12	14	25	51	82	92	129	93	34	7	-
Male	8	5	9	20	32	51	50	63	32	16	1	-
Female	2	7	5	5	19	31	42	66	61	18	6	-
White, both sexes	7	8	11	18	36	59	72	110	79	30	4	-
Male	6	4	6	13	23	34	41	53	29	15	1	-
Female	1	4	5	5	13	25	31	57	50	15	3	-
Black, both sexes	3	3	3	7	15	20	17	19	13	3	2	-
Male	2	1	3	7	9	15	7	10	2	-	-	-
Female	1	2	-	-	6	5	10	9	11	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	2	3	-	-	1	1	-
Male	-	-	-	-	-	1	2	-	-	1	-	-
Female	-	1	-	-	-	1	1	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Washington												
All races, both sexes	6	10	17	24	22	33	76	109	85	36	7	-
Male	2	7	7	14	16	14	35	61	42	12	2	-
Female	4	3	10	10	6	19	41	48	43	24	5	-
White, both sexes	5	8	16	22	19	30	70	101	80	36	7	-
Male	2	5	7	13	15	13	30	55	39	12	2	-
Female	3	3	9	9	4	17	40	46	41	24	5	-
Black, both sexes	1	1	1	-	1	-	3	5	1	-	-	-
Male	-	1	-	-	1	-	2	4	1	-	-	-
Female	1	-	1	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	2	-	1	-	1	1	-	-	-
Male	-	1	-	1	-	-	-	1	1	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	2	3	2	3	-	-	-
Male	-	-	-	-	-	1	3	1	1	-	-	-
Female	-	-	-	-	2	1	-	1	2	-	-	-
West Virginia												
All races, both sexes	1	4	2	7	8	19	20	27	22	8	2	-
Male	1	4	1	6	7	9	9	11	10	4	1	-
Female	-	-	1	1	1	10	11	16	12	4	1	-
White, both sexes	1	4	2	7	8	19	20	27	22	7	2	-
Male	1	4	1	6	7	9	9	11	10	3	1	-
Female	-	-	1	1	1	10	11	16	12	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Wisconsin												
All races, both sexes	5	6	15	14	13	38	78	90	65	20	2	-
Male	3	5	9	10	10	25	49	52	32	10	-	-
Female	2	1	6	4	3	13	29	38	33	10	2	-
White, both sexes	5	6	13	14	11	35	78	89	64	20	2	-
Male	3	5	8	10	9	24	49	51	31	10	-	-
Female	2	1	5	4	2	11	29	38	33	10	2	-
Black, both sexes	-	-	1	-	2	3	-	1	1	-	-	-
Male	-	-	1	-	1	1	-	1	1	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	5	1	6	7	7	1	-	-
Male	-	-	1	-	3	1	5	2	3	-	-	-
Female	-	-	-	-	2	-	1	5	4	1	-	-
White, both sexes	-	-	1	-	5	1	6	7	7	1	-	-
Male	-	-	1	-	3	1	5	2	3	-	-	-
Female	-	-	-	-	2	-	1	5	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
United States												
All races, both sexes	16,880	10	8	6	7	11	19	34	44	52	114	184
Male	8,912	5	2	4	7	4	11	22	31	34	69	109
Female	7,968	5	6	2	-	7	8	12	13	18	45	75
White, both sexes	15,131	6	6	5	5	7	12	23	34	40	86	147
Male	7,982	2	1	3	5	2	8	14	24	24	55	91
Female	7,149	4	5	2	-	5	4	9	10	16	31	56
Black, both sexes	1,438	4	-	1	2	2	7	11	9	10	27	33
Male	756	3	-	1	2	1	3	8	6	8	14	14
Female	682	1	-	-	-	1	4	3	3	2	13	19
American Indian 1/, both sexes	68	-	-	-	-	1	-	-	1	1	1	2
Male	39	-	-	-	-	-	-	-	1	1	-	2
Female	29	-	-	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	243	-	2	-	-	1	-	-	-	1	-	2
Male	135	-	1	-	-	1	-	-	-	1	-	2
Female	108	-	1	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	273	-	-	-	-	-	2	1	1	2	1	5
Male	138	-	-	-	-	-	2	-	1	2	-	4
Female	135	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	224	-	-	-	-	-	-	-	1	1	1	4
Male	115	-	-	-	-	-	-	-	1	1	-	3
Female	109	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	48	-	-	-	-	-	2	1	-	1	-	1
Male	23	-	-	-	-	-	2	-	-	1	-	1
Female	25	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Alaska												
All races, both sexes	12	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	230	-	-	-	-	-	-	-	2	1	-	4
Male	129	-	-	-	-	-	-	-	2	1	-	3
Female	101	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	219	-	-	-	-	-	-	-	2	1	-	4
Male	122	-	-	-	-	-	-	-	2	1	-	3
Female	97	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Arkansas												
All races, both sexes	191	-	-	-	-	-	-	-	-	2	-	2
Male	99	-	-	-	-	-	-	-	-	2	-	2
Female	92	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	169	-	-	-	-	-	-	-	-	2	-	1
Male	89	-	-	-	-	-	-	-	-	2	-	1
Female	80	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	887	-	1	-	1	1	-	2	1	2	2	10
Male	472	-	-	-	1	1	-	1	1	2	2	7
Female	415	-	1	-	-	-	-	1	-	-	-	3
White, both sexes	798	-	-	-	1	-	-	1	1	2	2	8
Male	427	-	-	-	1	-	-	-	1	2	2	6
Female	371	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	39	-	-	-	-	-	-	1	-	-	-	1
Male	21	-	-	-	-	-	-	1	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	-	1	-	-	1	-	-	-	-	-	1
Male	24	-	-	-	-	1	-	-	-	-	-	1
Female	25	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Colorado												
All races, both sexes	249	1	-	-	-	-	-	-	-	-	-	1
Male	132	-	-	-	-	-	-	-	-	-	-	1
Female	117	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	238	1	-	-	-	-	-	-	-	-	-	-
Male	125	-	-	-	-	-	-	-	-	-	-	-
Female	113	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	327	-	-	-	-	-	-	-	1	-	3	2
Male	163	-	-	-	-	-	-	-	1	-	2	2
Female	164	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	313	-	-	-	-	-	-	-	1	-	3	2
Male	156	-	-	-	-	-	-	-	1	-	2	2
Female	157	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Delaware												
All races, both sexes	50	-	-	-	-	-	-	-	-	1	1	-
Male	26	-	-	-	-	-	-	-	-	1	-	-
Female	24	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	25	-	-	-	1	-	-	-	-	-	-	-
Male	9	-	-	-	1	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	18	-	-	-	1	-	-	-	-	-	-	-
Male	5	-	-	-	1	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Florida												
All races, both sexes	923	-	-	-	-	-	3	3	1	3	8	11
Male	534	-	-	-	-	-	1	3	1	3	5	6
Female	389	-	-	-	-	-	2	-	-	-	3	5
White, both sexes	829	-	-	-	-	-	2	3	1	3	6	10
Male	473	-	-	-	-	-	1	3	1	3	3	6
Female	356	-	-	-	-	-	1	-	-	-	3	4
Black, both sexes	89	-	-	-	-	-	1	-	-	-	2	1
Male	57	-	-	-	-	-	-	-	-	-	2	-
Female	32	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	466	-	2	-	-	2	1	2	3	-	3	4
Male	263	-	1	-	-	1	1	2	-	-	1	3
Female	203	-	1	-	-	1	-	-	3	-	2	1
White, both sexes	367	-	1	-	-	1	1	2	2	-	2	4
Male	217	-	-	-	-	1	1	2	-	-	1	3
Female	150	-	1	-	-	-	-	-	2	-	1	1
Black, both sexes	96	-	-	-	-	1	-	-	1	-	1	-
Male	44	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	1	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Hawaii												
All races, both sexes	122	-	-	-	-	-	-	-	-	1	-	1
Male	68	-	-	-	-	-	-	-	-	1	-	1
Female	54	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	101	-	-	-	-	-	-	-	-	-	-	1
Male	57	-	-	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	59	-	-	-	-	-	-	-	-	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	59	-	-	-	-	-	-	-	-	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Illinois												
All races, both sexes	899	1	-	-	-	-	1	-	2	1	6	2
Male	438	1	-	-	-	-	1	-	1	1	3	1
Female	461	-	-	-	-	-	-	-	1	-	3	1
White, both sexes	838	-	-	-	-	-	1	-	2	1	5	2
Male	409	-	-	-	-	-	1	-	1	1	3	1
Female	429	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	50	1	-	-	-	-	-	-	-	-	1	-
Male	24	1	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	464	-	-	-	-	1	-	-	2	2	3	10
Male	238	-	-	-	-	-	-	-	1	2	1	4
Female	226	-	-	-	-	1	-	-	1	-	2	6
White, both sexes	442	-	-	-	-	1	-	-	2	2	3	8
Male	226	-	-	-	-	-	-	-	1	2	1	3
Female	216	-	-	-	-	1	-	-	1	-	2	5
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Iowa												
All races, both sexes	283	1	-	1	-	-	-	2	-	3	-	1
Male	147	1	-	1	-	-	-	-	-	1	-	-
Female	136	-	-	-	-	-	-	2	-	2	-	1
White, both sexes	277	1	-	1	-	-	-	2	-	3	-	1
Male	143	1	-	1	-	-	-	-	-	1	-	-
Female	134	-	-	-	-	-	-	2	-	2	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	167	-	-	-	-	-	-	-	-	2	-	1
Male	81	-	-	-	-	-	-	-	-	2	-	1
Female	86	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	162	-	-	-	-	-	-	-	-	1	-	1
Male	77	-	-	-	-	-	-	-	-	1	-	1
Female	85	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Kentucky												
All races, both sexes	350	1	-	-	-	-	-	-	2	1	3	3
Male	169	-	-	-	-	-	-	-	1	-	2	2
Female	181	1	-	-	-	-	-	-	1	1	1	1
White, both sexes	327	1	-	-	-	-	-	-	2	1	1	3
Male	157	-	-	-	-	-	-	-	1	-	1	2
Female	170	1	-	-	-	-	-	-	1	1	-	1
Black, both sexes	22	-	-	-	-	-	-	-	-	-	2	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	209	-	-	-	1	-	-	-	1	2	2	1
Male	103	-	-	-	1	-	-	-	1	2	1	1
Female	106	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	155	-	-	-	1	-	-	-	1	1	1	-
Male	79	-	-	-	1	-	-	-	1	1	-	-
Female	76	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	53	-	-	-	-	-	-	-	-	1	1	1
Male	24	-	-	-	-	-	-	-	-	1	1	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Maine												
All races, both sexes	91	-	-	-	-	-	-	1	-	1	-	-
Male	47	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	91	-	-	-	-	-	-	1	-	1	-	-
Male	47	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	359	-	-	-	-	-	1	1	1	1	4	5
Male	186	-	-	-	-	-	-	1	1	1	1	3
Female	173	-	-	-	-	-	1	-	-	-	3	2
White, both sexes	280	-	-	-	-	-	1	1	-	1	3	3
Male	149	-	-	-	-	-	-	1	-	1	1	2
Female	131	-	-	-	-	-	1	-	-	-	2	1
Black, both sexes	73	-	-	-	-	-	-	-	1	-	1	2
Male	34	-	-	-	-	-	-	-	1	-	-	1
Female	39	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Massachusetts												
All races, both sexes	529	-	1	-	-	-	-	-	-	-	1	3
Male	278	-	1	-	-	-	-	-	-	-	1	1
Female	251	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	510	-	1	-	-	-	-	-	-	-	1	3
Male	268	-	1	-	-	-	-	-	-	-	1	1
Female	242	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	533	-	-	-	-	-	1	-	3	2	6	5
Male	284	-	-	-	-	-	-	-	2	1	5	4
Female	249	-	-	-	-	-	1	-	1	1	1	1
White, both sexes	486	-	-	-	-	-	-	-	3	-	6	3
Male	255	-	-	-	-	-	-	-	2	-	5	2
Female	231	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	44	-	-	-	-	-	1	-	-	1	-	2
Male	28	-	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Minnesota												
All races, both sexes	228	-	-	-	-	-	-	1	-	-	1	1
Male	127	-	-	-	-	-	-	1	-	-	1	1
Female	101	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	223	-	-	-	-	-	-	1	-	-	1	1
Male	123	-	-	-	-	-	-	1	-	-	1	1
Female	100	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	186	2	-	1	-	-	-	3	1	-	3	3
Male	101	1	-	1	-	-	-	2	1	-	2	1
Female	85	1	-	-	-	-	-	1	-	-	1	2
White, both sexes	144	-	-	-	-	-	-	2	1	-	2	2
Male	82	-	-	-	-	-	-	1	1	-	2	1
Female	62	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	41	2	-	1	-	-	-	1	-	-	1	1
Male	18	1	-	1	-	-	-	1	-	-	-	-
Female	23	1	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Missouri												
All races, both sexes	418	-	1	1	1	-	-	-	1	-	2	4
Male	218	-	-	-	1	-	-	-	-	-	1	3
Female	200	-	1	1	-	-	-	-	1	-	1	1
White, both sexes	387	-	1	1	-	-	-	-	1	-	1	3
Male	201	-	-	-	-	-	-	-	-	-	-	2
Female	186	-	1	1	-	-	-	-	1	-	1	1
Black, both sexes	29	-	-	-	1	-	-	-	-	-	1	1
Male	16	-	-	-	1	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	72	-	1	-	-	-	-	-	1	-	1	-
Male	36	-	-	-	-	-	-	-	1	-	-	-
Female	36	-	1	-	-	-	-	-	-	-	1	-
White, both sexes	69	-	1	-	-	-	-	-	-	-	1	-
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	1	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Nebraska												
All races, both sexes	111	-	-	-	-	1	-	-	1	1	1	1
Male	58	-	-	-	-	-	-	-	1	1	-	1
Female	53	-	-	-	-	1	-	-	-	-	1	-
White, both sexes	106	-	-	-	-	1	-	-	-	1	1	1
Male	55	-	-	-	-	-	-	-	-	1	-	1
Female	51	-	-	-	-	1	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	115	-	-	-	-	-	1	-	-	-	-	3
Male	64	-	-	-	-	-	1	-	-	-	-	2
Female	51	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	105	-	-	-	-	-	1	-	-	-	-	2
Male	60	-	-	-	-	-	1	-	-	-	-	2
Female	45	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
New Hampshire												
All races, both sexes	62	-	-	-	-	-	-	1	-	1	-	-
Male	27	-	-	-	-	-	-	1	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	62	-	-	-	-	-	-	1	-	1	-	-
Male	27	-	-	-	-	-	-	1	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	513	-	-	1	-	1	-	2	-	1	5	7
Male	272	-	-	1	-	-	-	1	-	1	4	3
Female	241	-	-	-	-	1	-	1	-	-	1	4
White, both sexes	454	-	-	1	-	1	-	2	-	-	5	3
Male	244	-	-	1	-	-	-	1	-	-	4	2
Female	210	-	-	-	-	1	-	1	-	-	1	1
Black, both sexes	55	-	-	-	-	-	-	-	-	-	-	4
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
New Mexico												
All races, both sexes	70	-	-	-	-	-	-	-	1	-	1	2
Male	38	-	-	-	-	-	-	-	1	-	-	2
Female	32	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	61	-	-	-	-	-	-	-	1	-	1	2
Male	36	-	-	-	-	-	-	-	1	-	-	2
Female	25	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	711	2	-	1	-	1	-	-	4	-	4	11
Male	387	1	-	-	-	1	-	-	3	-	3	6
Female	324	1	-	1	-	-	-	-	1	-	1	5
White, both sexes	665	1	-	1	-	1	-	-	2	-	3	11
Male	360	-	-	-	-	1	-	-	2	-	2	6
Female	305	1	-	1	-	-	-	-	-	-	1	5
Black, both sexes	39	1	-	-	-	-	-	-	2	-	1	-
Male	23	1	-	-	-	-	-	-	1	-	1	-
Female	16	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
North Carolina												
All races, both sexes	655	-	-	-	-	1	1	-	3	3	9	8
Male	343	-	-	-	-	-	-	-	2	2	6	3
Female	312	-	-	-	-	1	1	-	1	1	3	5
White, both sexes	550	-	-	-	-	1	-	-	2	-	5	6
Male	282	-	-	-	-	-	-	-	1	-	3	3
Female	268	-	-	-	-	1	-	-	1	-	2	3
Black, both sexes	102	-	-	-	-	-	1	-	1	3	4	2
Male	58	-	-	-	-	-	-	-	1	2	3	-
Female	44	-	-	-	-	-	1	-	-	1	1	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	1	1
Male	27	-	-	-	-	-	-	-	-	-	1	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	48	-	-	-	-	-	-	-	-	-	1	-
Male	26	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Ohio												
All races, both sexes	805	-	1	-	-	-	-	1	2	3	1	8
Male	424	-	-	-	-	-	-	1	2	-	1	6
Female	381	-	1	-	-	-	-	-	-	3	-	2
White, both sexes	731	-	1	-	-	-	-	-	1	3	1	7
Male	384	-	-	-	-	-	-	-	1	-	1	6
Female	347	-	1	-	-	-	-	-	-	3	-	1
Black, both sexes	73	-	-	-	-	-	-	1	1	-	-	1
Male	40	-	-	-	-	-	-	1	1	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	200	-	-	-	-	-	-	-	2	-	3	1
Male	89	-	-	-	-	-	-	-	1	-	1	-
Female	111	-	-	-	-	-	-	-	1	-	2	1
White, both sexes	184	-	-	-	-	-	-	-	1	-	3	1
Male	83	-	-	-	-	-	-	-	-	-	1	-
Female	101	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Oregon												
All races, both sexes	166	-	-	-	-	-	2	-	-	-	-	2
Male	85	-	-	-	-	-	1	-	-	-	-	1
Female	81	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	160	-	-	-	-	-	2	-	-	-	-	2
Male	80	-	-	-	-	-	1	-	-	-	-	1
Female	80	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,129	1	-	1	-	1	2	2	1	2	9	17
Male	608	1	-	1	-	1	1	1	1	-	6	8
Female	521	-	-	-	-	-	1	1	-	2	3	9
White, both sexes	1,021	1	-	1	-	-	2	1	1	2	7	13
Male	559	1	-	1	-	-	1	1	1	-	5	7
Female	462	-	-	-	-	-	1	-	-	2	2	6
Black, both sexes	101	-	-	-	-	1	-	1	-	-	2	4
Male	45	-	-	-	-	1	-	-	-	-	1	1
Female	56	-	-	-	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Rhode Island												
All races, both sexes	86	-	-	-	-	-	-	1	-	-	-	1
Male	49	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	83	-	-	-	-	-	-	1	-	-	-	1
Male	47	-	-	-	-	-	-	-	-	-	-	1
Female	36	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	276	-	-	-	-	-	-	2	2	2	1	4
Male	147	-	-	-	-	-	-	1	-	1	-	2
Female	129	-	-	-	-	-	-	1	2	1	1	2
White, both sexes	206	-	-	-	-	-	-	1	1	1	-	3
Male	107	-	-	-	-	-	-	-	-	-	-	2
Female	99	-	-	-	-	-	-	1	1	1	-	1
Black, both sexes	70	-	-	-	-	-	-	1	1	1	1	1
Male	40	-	-	-	-	-	-	1	-	1	-	-
Female	30	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
South Dakota												
All races, both sexes	44	-	-	-	-	-	-	-	-	1	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	1	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	453	-	-	-	-	-	1	2	1	-	4	8
Male	230	-	-	-	-	-	-	1	1	-	3	5
Female	223	-	-	-	-	-	1	1	-	-	1	3
White, both sexes	411	-	-	-	-	-	-	-	1	-	3	7
Male	207	-	-	-	-	-	-	-	1	-	3	4
Female	204	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	41	-	-	-	-	-	1	2	-	-	1	1
Male	22	-	-	-	-	-	-	1	-	-	-	1
Female	19	-	-	-	-	-	1	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Texas												
All races, both sexes	1,196	1	1	-	2	-	1	3	1	6	14	13
Male	641	-	-	-	2	-	1	3	1	4	10	8
Female	555	1	1	-	-	-	-	-	-	2	4	5
White, both sexes	1,090	1	1	-	2	-	1	1	1	6	10	12
Male	576	-	-	-	2	-	1	1	1	4	7	7
Female	514	1	1	-	-	-	-	-	-	2	3	5
Black, both sexes	100	-	-	-	-	-	-	2	-	-	3	1
Male	60	-	-	-	-	-	-	2	-	-	3	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	94	-	-	-	1	-	1	1	-	-	-	-
Male	55	-	-	-	1	-	1	1	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	93	-	-	-	1	-	1	1	-	-	-	-
Male	54	-	-	-	1	-	1	1	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Vermont												
All races, both sexes	37	-	-	-	-	1	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	37	-	-	-	-	1	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	564	-	-	-	-	-	1	-	1	2	3	8
Male	297	-	-	-	-	-	1	-	1	2	2	4
Female	267	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	442	-	-	-	-	-	-	-	1	1	1	5
Male	231	-	-	-	-	-	-	-	1	1	1	3
Female	211	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	112	-	-	-	-	-	1	-	-	1	2	3
Male	60	-	-	-	-	-	1	-	-	1	1	1
Female	52	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Washington												
All races, both sexes	438	-	-	-	-	1	-	-	1	3	4	4
Male	218	-	-	-	-	-	-	-	1	1	2	2
Female	220	-	-	-	-	1	-	-	-	2	2	2
White, both sexes	405	-	-	-	-	-	-	-	1	3	3	4
Male	199	-	-	-	-	-	-	-	1	1	2	2
Female	206	-	-	-	-	-	-	-	-	2	1	2
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	7	-	-	-	-	1	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	123	-	-	-	-	-	-	1	-	-	2	-
Male	64	-	-	-	-	-	-	1	-	-	-	-
Female	59	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	120	-	-	-	-	-	-	-	-	-	1	-
Male	62	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	1	-	-	1	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Wisconsin												
All races, both sexes	351	-	-	-	-	-	-	1	-	-	2	5
Male	208	-	-	-	-	-	-	1	-	-	2	2
Female	143	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	340	-	-	-	-	-	-	1	-	-	2	3
Male	202	-	-	-	-	-	-	1	-	-	2	1
Female	138	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	29	-	-	-	-	-	-	1	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	1	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
United States												
All races, both sexes	253	357	495	658	1,092	1,955	3,328	4,037	2,902	1,091	223	-
Male	163	221	314	420	672	1,161	1,867	2,056	1,298	376	66	-
Female	90	136	181	238	420	794	1,461	1,981	1,604	715	157	-
White, both sexes	212	294	425	551	913	1,712	3,038	3,748	2,667	1,010	190	-
Male	136	184	268	348	568	1,011	1,714	1,912	1,202	354	56	-
Female	76	110	157	203	345	701	1,324	1,836	1,465	656	134	-
Black, both sexes	40	55	63	92	158	196	233	233	174	64	24	-
Male	26	31	43	64	91	123	124	110	65	13	6	-
Female	14	24	20	28	67	73	109	123	109	51	18	-
American Indian 1/, both sexes	1	4	3	12	3	14	2	10	12	-	1	-
Male	1	3	1	7	2	7	1	7	5	-	1	-
Female	-	1	2	5	1	7	1	3	7	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	4	3	18	33	55	46	49	17	8	-
Male	-	3	2	1	11	20	28	27	26	9	3	-
Female	-	1	2	2	7	13	27	19	23	8	5	-
Alabama												
All races, both sexes	5	12	1	10	19	30	50	74	46	9	5	-
Male	3	8	1	6	10	18	25	42	11	4	1	-
Female	2	4	-	4	9	12	25	32	35	5	4	-
White, both sexes	3	10	1	8	15	23	42	64	40	7	4	-
Male	2	7	1	5	8	12	23	37	10	4	1	-
Female	1	3	-	3	7	11	19	27	30	3	3	-
Black, both sexes	2	2	-	1	4	7	8	10	6	2	1	-
Male	1	1	-	1	2	6	2	5	1	-	-	-
Female	1	1	-	-	2	1	6	5	5	2	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Alaska												
All races, both sexes	-	-	1	-	1	2	4	1	2	-	-	-
Male	-	-	-	-	-	2	3	-	-	-	-	-
Female	-	-	1	-	1	-	1	1	2	-	-	-
White, both sexes	-	-	1	-	1	2	3	1	-	-	-	-
Male	-	-	-	-	-	2	3	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	4	8	12	18	32	54	51	26	11	2	-
Male	3	2	4	5	13	12	34	26	16	6	2	-
Female	2	2	4	7	5	20	20	25	10	5	-	-
White, both sexes	5	2	8	12	18	29	54	48	25	10	1	-
Male	3	1	4	5	13	10	34	24	15	6	1	-
Female	2	1	4	7	5	19	20	24	10	4	-	-
Black, both sexes	-	1	-	-	-	1	-	-	-	1	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	2	-	2	1	-	1	-
Male	-	-	-	-	-	1	-	2	1	-	1	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Arkansas												
All races, both sexes	4	6	6	3	7	24	40	45	34	15	3	-
Male	2	2	5	2	2	14	26	25	12	5	-	-
Female	2	4	1	1	5	10	14	20	22	10	3	-
White, both sexes	2	4	4	2	5	21	37	43	32	14	2	-
Male	1	2	3	1	1	13	24	24	12	5	-	-
Female	1	2	1	1	4	8	13	19	20	9	2	-
Black, both sexes	2	2	2	1	2	3	3	2	2	1	1	-
Male	1	-	2	1	1	1	2	1	-	-	-	-
Female	1	2	-	-	1	2	1	1	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	13	17	32	37	48	112	186	203	136	66	17	-
Male	8	7	20	22	28	60	97	110	73	25	7	-
Female	5	10	12	15	20	52	89	93	63	41	10	-
White, both sexes	12	15	27	33	38	100	172	188	121	63	14	-
Male	7	7	17	21	22	54	90	103	64	24	6	-
Female	5	8	10	12	16	46	82	85	57	39	8	-
Black, both sexes	1	2	4	3	8	4	5	4	6	-	-	-
Male	1	-	3	1	5	2	4	1	3	-	-	-
Female	-	2	1	2	3	2	1	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	8	9	11	9	3	3	-
Male	-	-	-	-	1	4	3	6	6	1	1	-
Female	-	-	1	-	1	4	6	5	3	2	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Colorado												
All races, both sexes	5	5	7	4	14	34	43	66	52	14	3	-
Male	4	3	4	1	9	22	26	35	21	4	2	-
Female	1	2	3	3	5	12	17	31	31	10	1	-
White, both sexes	4	5	7	4	10	33	41	64	52	14	3	-
Male	3	3	4	1	6	22	25	34	21	4	2	-
Female	1	2	3	3	4	11	16	30	31	10	1	-
Black, both sexes	-	-	-	-	2	1	1	2	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	1	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	1	4	10	10	18	24	73	82	68	29	2	-
Male	-	2	4	6	7	15	43	39	32	10	-	-
Female	1	2	6	4	11	9	30	43	36	19	2	-
White, both sexes	1	4	9	9	15	22	70	80	66	29	2	-
Male	-	2	4	5	6	13	40	39	32	10	-	-
Female	1	2	5	4	9	9	30	41	34	19	2	-
Black, both sexes	-	-	1	1	3	2	3	2	2	-	-	-
Male	-	-	-	1	1	2	3	-	-	-	-	-
Female	-	-	1	-	2	-	-	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Delaware												
All races, both sexes	1	1	3	1	1	5	8	15	11	2	-	-
Male	1	1	3	-	1	5	5	5	4	-	-	-
Female	-	-	-	1	-	-	3	10	7	2	-	-
White, both sexes	1	1	1	1	1	4	8	14	10	2	-	-
Male	1	1	1	-	1	4	5	4	3	-	-	-
Female	-	-	-	1	-	-	3	10	7	2	-	-
Black, both sexes	-	-	2	-	-	1	-	1	1	-	-	-
Male	-	-	2	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	-	1	-	3	4	8	4	3	-	-
Male	-	1	-	-	-	1	2	2	-	2	-	-
Female	-	-	-	1	-	2	2	6	4	1	-	-
White, both sexes	-	-	-	-	-	1	1	1	1	2	-	-
Male	-	-	-	-	-	1	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	-	1	-	1	-	2	2	7	3	1	-	-
Male	-	1	-	-	-	-	-	2	-	1	-	-
Female	-	-	-	1	-	2	2	5	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Florida												
All races, both sexes	9	24	21	34	52	102	197	221	161	62	11	-
Male	4	21	14	21	32	75	121	122	76	26	3	-
Female	5	3	7	13	20	27	76	99	85	36	8	-
White, both sexes	7	17	18	23	45	91	179	210	148	57	9	-
Male	3	14	13	12	27	66	110	115	69	24	3	-
Female	4	3	5	11	18	25	69	95	79	33	6	-
Black, both sexes	2	7	3	11	7	9	17	10	12	5	2	-
Male	1	7	1	9	5	7	11	6	6	2	-	-
Female	1	-	2	2	2	2	6	4	6	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	9	7	16	24	32	58	95	104	70	27	7	-
Male	5	4	11	20	22	39	58	49	33	10	3	-
Female	4	3	5	4	10	19	37	55	37	17	4	-
White, both sexes	7	2	12	20	23	45	75	87	59	18	6	-
Male	4	2	9	16	17	29	48	44	30	7	3	-
Female	3	-	3	4	6	16	27	43	29	11	3	-
Black, both sexes	2	5	4	4	9	13	19	17	10	9	1	-
Male	1	2	2	4	5	10	10	5	2	3	-	-
Female	1	3	2	-	4	3	9	12	8	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Hawaii												
All races, both sexes	-	3	3	4	5	11	28	20	31	13	2	-
Male	-	3	3	2	4	6	16	9	14	7	2	-
Female	-	-	-	2	1	5	12	11	17	6	-	-
White, both sexes	-	-	1	1	-	2	5	4	7	-	-	-
Male	-	-	1	1	-	1	2	2	3	-	-	-
Female	-	-	-	-	-	1	3	2	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	2	3	5	9	23	16	24	13	2	-
Male	-	3	2	1	4	5	14	7	11	7	2	-
Female	-	-	-	2	1	4	9	9	13	6	-	-
Idaho												
All races, both sexes	3	5	4	1	3	7	13	12	9	1	-	-
Male	1	4	2	1	3	4	4	9	4	-	-	-
Female	2	1	2	-	-	3	9	3	5	1	-	-
White, both sexes	3	5	4	1	3	7	13	12	9	1	-	-
Male	1	4	2	1	3	4	4	9	4	-	-	-
Female	2	1	2	-	-	3	9	3	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Illinois												
All races, both sexes	13	16	29	26	52	95	177	238	170	58	12	-
Male	10	10	20	13	28	47	88	109	83	21	1	-
Female	3	6	9	13	24	48	89	129	87	37	11	-
White, both sexes	11	15	26	25	41	83	166	227	167	56	10	-
Male	9	10	18	12	21	40	84	105	82	20	1	-
Female	2	5	8	13	20	43	82	122	85	36	9	-
Black, both sexes	2	1	2	1	10	9	7	10	3	2	1	-
Male	1	-	2	1	6	4	3	4	1	1	-	-
Female	1	1	-	-	4	5	4	6	2	1	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	3	4	1	-	-	1	-
Male	-	-	-	-	1	3	1	-	-	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	1	-
Indiana												
All races, both sexes	9	12	14	22	36	52	83	112	73	27	6	-
Male	6	8	10	11	22	28	49	57	30	7	2	-
Female	3	4	4	11	14	24	34	55	43	20	4	-
White, both sexes	9	11	13	20	35	49	80	109	70	26	4	-
Male	6	7	10	11	21	26	47	56	28	6	1	-
Female	3	4	3	9	14	23	33	53	42	20	3	-
Black, both sexes	-	1	1	2	1	3	3	3	3	1	2	-
Male	-	1	-	-	1	2	2	1	2	1	1	-
Female	-	-	1	2	-	1	1	2	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Iowa												
All races, both sexes	2	1	7	9	20	27	41	79	64	23	2	-
Male	1	-	5	7	10	13	25	38	31	14	-	-
Female	1	1	2	2	10	14	16	41	33	9	2	-
White, both sexes	2	1	7	7	20	26	40	79	62	23	2	-
Male	1	-	5	6	10	12	24	38	30	14	-	-
Female	1	1	2	1	10	14	16	41	32	9	2	-
Black, both sexes	-	-	-	1	-	1	1	-	2	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	4	-	4	8	3	15	30	41	35	20	4	-
Male	2	-	1	7	3	8	12	22	18	5	-	-
Female	2	-	3	1	-	7	18	19	17	15	4	-
White, both sexes	4	-	4	8	3	14	29	40	34	20	4	-
Male	2	-	1	7	3	7	12	21	17	5	-	-
Female	2	-	3	1	-	7	17	19	17	15	4	-
Black, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Kentucky												
All races, both sexes	2	4	11	16	19	49	68	79	71	19	2	-
Male	1	4	7	10	11	34	31	31	33	1	1	-
Female	1	-	4	6	8	15	37	48	38	18	1	-
White, both sexes	2	4	10	13	19	45	66	75	67	16	2	-
Male	1	4	6	8	11	31	30	29	31	1	1	-
Female	1	-	4	5	8	14	36	46	36	15	1	-
Black, both sexes	-	-	1	3	-	4	1	4	4	3	-	-
Male	-	-	1	2	-	3	1	2	2	-	-	-
Female	-	-	-	1	-	1	-	2	2	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Louisiana												
All races, both sexes	3	4	12	10	18	26	44	39	35	9	2	-
Male	3	3	5	3	11	18	22	14	13	5	-	-
Female	-	1	7	7	7	8	22	25	22	4	2	-
White, both sexes	3	3	9	9	10	18	35	29	26	8	1	-
Male	3	2	4	3	6	14	19	11	10	4	-	-
Female	-	1	5	6	4	4	16	18	16	4	1	-
Black, both sexes	-	1	3	1	8	8	9	9	9	1	1	-
Male	-	1	1	-	5	4	3	3	3	1	-	-
Female	-	-	2	1	3	4	6	6	6	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Maine												
All races, both sexes	-	1	1	2	6	9	18	25	16	10	1	-
Male	-	1	1	2	4	6	9	14	7	2	1	-
Female	-	-	-	-	2	3	9	11	9	8	-	-
White, both sexes	-	1	1	2	6	9	18	25	16	10	1	-
Male	-	1	1	2	4	6	9	14	7	2	1	-
Female	-	-	-	-	2	3	9	11	9	8	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	9	10	14	10	33	43	63	86	54	20	4	-
Male	4	3	11	6	17	27	38	43	26	4	-	-
Female	5	7	3	4	16	16	25	43	28	16	4	-
White, both sexes	5	6	9	4	19	32	51	76	47	19	3	-
Male	1	3	8	2	13	21	29	38	25	4	-	-
Female	4	3	1	2	6	11	22	38	22	15	3	-
Black, both sexes	4	4	5	6	13	8	10	10	7	1	1	-
Male	3	-	3	4	4	5	7	5	1	-	-	-
Female	1	4	2	2	9	3	3	5	6	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	3	2	-	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Massachusetts												
All races, both sexes	5	5	7	13	30	72	116	119	103	43	11	-
Male	2	3	7	7	13	52	65	60	44	17	5	-
Female	3	2	-	6	17	20	51	59	59	26	6	-
White, both sexes	5	5	7	12	29	70	112	114	99	42	10	-
Male	2	3	7	6	12	51	62	57	43	17	5	-
Female	3	2	-	6	17	19	50	57	56	25	5	-
Black, both sexes	-	-	-	1	-	1	2	3	4	1	1	-
Male	-	-	-	1	-	-	2	1	1	-	-	-
Female	-	-	-	-	-	1	-	2	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	2	2	-	-	-	-
Male	-	-	-	-	1	1	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Michigan												
All races, both sexes	10	11	13	25	29	73	101	125	93	30	6	-
Male	7	7	11	15	20	41	58	66	34	11	2	-
Female	3	4	2	10	9	32	43	59	59	19	4	-
White, both sexes	7	9	11	24	23	68	92	117	87	30	6	-
Male	5	5	10	15	16	37	52	62	31	11	2	-
Female	2	4	1	9	7	31	40	55	56	19	4	-
Black, both sexes	3	2	2	1	6	5	9	8	4	-	-	-
Male	2	2	1	-	4	4	6	4	3	-	-	-
Female	1	-	1	1	2	1	3	4	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Minnesota												
All races, both sexes	4	3	9	6	11	34	31	55	51	16	5	-
Male	4	2	6	6	8	19	17	32	24	5	1	-
Female	-	1	3	-	3	15	14	23	27	11	4	-
White, both sexes	4	3	9	5	9	33	30	55	51	16	5	-
Male	4	2	6	5	6	19	16	32	24	5	1	-
Female	-	1	3	-	3	14	14	23	27	11	4	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	6	7	7	11	25	32	43	23	13	2	-
Male	3	3	6	6	4	18	21	21	9	2	-	-
Female	1	3	1	1	7	7	11	22	14	11	2	-
White, both sexes	3	4	5	5	8	19	26	36	19	10	2	-
Male	2	1	4	4	3	14	20	18	9	2	-	-
Female	1	3	1	1	5	5	6	18	10	8	2	-
Black, both sexes	1	2	2	2	3	5	6	7	4	3	-	-
Male	1	2	2	2	1	3	1	3	-	-	-	-
Female	-	-	-	-	2	2	5	4	4	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Missouri												
All races, both sexes	11	9	9	13	26	53	82	100	78	21	6	-
Male	5	2	6	7	18	33	43	52	39	6	2	-
Female	6	7	3	6	8	20	39	48	39	15	4	-
White, both sexes	10	8	9	11	24	48	77	96	70	21	6	-
Male	4	2	6	5	17	29	42	50	36	6	2	-
Female	6	6	3	6	7	19	35	46	34	15	4	-
Black, both sexes	1	1	-	2	2	4	5	4	7	-	-	-
Male	1	-	-	2	1	3	1	2	3	-	-	-
Female	-	1	-	-	1	1	4	2	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Montana												
All races, both sexes	-	1	2	4	5	8	13	9	20	7	-	-
Male	-	1	2	3	4	4	9	2	7	3	-	-
Female	-	-	-	1	1	4	4	7	13	4	-	-
White, both sexes	-	1	2	3	4	8	13	9	20	7	-	-
Male	-	1	2	2	3	4	9	2	7	3	-	-
Female	-	-	-	1	1	4	4	7	13	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Nebraska												
All races, both sexes	-	2	-	6	1	13	17	35	19	11	2	-
Male	-	2	-	5	-	9	11	12	11	4	1	-
Female	-	-	-	1	1	4	6	23	8	7	1	-
White, both sexes	-	2	-	4	1	11	17	35	19	11	2	-
Male	-	2	-	4	-	8	11	12	11	4	1	-
Female	-	-	-	-	1	3	6	23	8	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	2	1	10	10	14	30	17	17	6	-	-
Male	3	2	1	6	5	10	20	7	5	2	-	-
Female	1	-	-	4	5	4	10	10	12	4	-	-
White, both sexes	4	2	1	9	7	13	28	17	15	6	-	-
Male	3	2	1	5	4	9	19	7	5	2	-	-
Female	1	-	-	4	3	4	9	10	10	4	-	-
Black, both sexes	-	-	-	-	2	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
New Hampshire												
All races, both sexes	1	1	3	3	3	1	15	16	12	5	-	-
Male	1	-	1	-	3	-	5	9	7	-	-	-
Female	-	1	2	3	-	1	10	7	5	5	-	-
White, both sexes	1	1	3	3	3	1	15	16	12	5	-	-
Male	1	-	1	-	3	-	5	9	7	-	-	-
Female	-	1	2	3	-	1	10	7	5	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	9	13	12	11	29	58	101	146	86	27	4	-
Male	5	6	9	7	20	34	59	77	37	8	-	-
Female	4	7	3	4	9	24	42	69	49	19	4	-
White, both sexes	4	12	9	8	25	48	94	132	81	25	4	-
Male	2	6	7	6	19	26	56	70	36	8	-	-
Female	2	6	2	2	6	22	38	62	45	17	4	-
Black, both sexes	5	1	3	3	4	9	7	12	5	2	-	-
Male	3	-	2	1	1	7	3	5	1	-	-	-
Female	2	1	1	2	3	2	4	7	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
New Mexico												
All races, both sexes	-	2	3	4	7	13	12	12	8	3	2	-
Male	-	1	3	3	2	4	8	8	3	2	1	-
Female	-	1	-	1	5	9	4	4	5	1	1	-
White, both sexes	-	1	3	4	5	10	12	11	6	3	2	-
Male	-	-	3	3	2	4	8	8	2	2	1	-
Female	-	1	-	1	3	6	4	3	4	1	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	3	-	1	2	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	3	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	11	15	24	25	42	67	140	197	106	52	9	-
Male	7	8	13	19	25	45	81	111	46	15	3	-
Female	4	7	11	6	17	22	59	86	60	37	6	-
White, both sexes	10	13	21	24	38	66	132	187	101	47	7	-
Male	7	6	11	18	23	44	77	104	42	15	2	-
Female	3	7	10	6	15	22	55	83	59	32	5	-
Black, both sexes	1	2	3	1	4	1	7	6	4	5	1	-
Male	-	2	2	1	2	1	4	4	3	-	1	-
Female	1	-	1	-	2	-	3	2	1	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	4	1	-	1	-
Male	-	-	-	-	-	-	-	3	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
North Carolina												
All races, both sexes	9	15	31	41	50	73	129	140	98	39	5	-
Male	5	10	19	25	31	42	71	66	52	8	1	-
Female	4	5	12	16	19	31	58	74	46	31	4	-
White, both sexes	8	9	25	34	38	59	117	124	83	35	4	-
Male	4	5	14	23	22	35	66	55	43	8	-	-
Female	4	4	11	11	16	24	51	69	40	27	4	-
Black, both sexes	1	5	5	7	12	13	12	16	15	4	1	-
Male	1	4	4	2	9	6	5	11	9	-	1	-
Female	-	1	1	5	3	7	7	5	6	4	-	-
American Indian 1/, both sexes	-	1	1	-	-	1	-	-	-	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	1	2	2	7	11	13	6	4	1	-
Male	-	-	1	1	2	4	7	5	3	2	-	-
Female	-	1	-	1	-	3	4	8	3	2	1	-
White, both sexes	-	1	1	1	2	7	11	13	6	4	1	-
Male	-	-	1	1	2	4	7	5	3	2	-	-
Female	-	1	-	-	-	3	4	8	3	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Ohio												
All races, both sexes	12	10	19	31	62	93	143	198	156	59	6	-
Male	9	7	14	23	43	65	84	89	59	20	1	-
Female	3	3	5	8	19	28	59	109	97	39	5	-
White, both sexes	11	8	16	28	54	80	132	183	146	55	5	-
Male	8	6	11	20	38	56	77	86	55	19	-	-
Female	3	2	5	8	16	24	55	97	91	36	5	-
Black, both sexes	1	2	3	3	7	13	11	15	10	4	1	-
Male	1	1	3	3	5	9	7	3	4	1	1	-
Female	-	1	-	-	2	4	4	12	6	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	2	6	11	13	26	36	48	37	10	5	-
Male	-	2	2	6	5	15	16	23	14	2	2	-
Female	-	-	4	5	8	11	20	25	23	8	3	-
White, both sexes	-	2	5	8	13	24	35	46	32	10	4	-
Male	-	2	2	4	5	15	15	22	13	2	2	-
Female	-	-	3	4	8	9	20	24	19	8	2	-
Black, both sexes	-	-	-	1	-	1	1	-	1	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	1	2	-	1	-	2	4	-	-	-
Male	-	-	-	2	-	-	-	1	1	-	-	-
Female	-	-	1	-	-	1	-	1	3	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Oregon												
All races, both sexes	7	2	3	7	16	13	23	36	40	12	3	-
Male	5	1	1	3	7	7	16	21	16	6	-	-
Female	2	1	2	4	9	6	7	15	24	6	3	-
White, both sexes	6	2	3	7	16	12	23	34	38	12	3	-
Male	4	1	1	3	7	6	16	20	14	6	-	-
Female	2	1	2	4	9	6	7	14	24	6	3	-
Black, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	1	2	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	17	28	24	34	72	138	252	257	199	52	20	-
Male	14	20	15	19	42	75	146	146	89	14	8	-
Female	3	8	9	15	30	63	106	111	110	38	12	-
White, both sexes	15	27	22	26	62	120	229	245	179	50	18	-
Male	12	19	13	15	36	68	136	140	82	14	7	-
Female	3	8	9	11	26	52	93	105	97	36	11	-
Black, both sexes	2	1	2	8	9	18	21	11	17	2	2	-
Male	2	1	2	4	5	7	9	6	5	-	1	-
Female	-	-	-	4	4	11	12	5	12	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	2	1	3	-	-	-
Male	-	-	-	-	1	-	1	-	2	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Rhode Island												
All races, both sexes	-	2	2	4	6	11	16	26	12	4	1	-
Male	-	1	1	3	5	7	10	12	8	-	1	-
Female	-	1	1	1	1	4	6	14	4	4	-	-
White, both sexes	-	2	2	4	6	9	16	25	12	4	1	-
Male	-	1	1	3	5	5	10	12	8	-	1	-
Female	-	1	1	1	1	4	6	13	4	4	-	-
Black, both sexes	-	-	-	-	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	9	7	13	30	36	65	62	31	7	4	-
Male	1	5	5	12	24	24	31	26	11	2	2	-
Female	-	4	2	1	6	12	34	36	20	5	2	-
White, both sexes	1	5	4	8	19	25	52	50	26	7	3	-
Male	1	2	4	8	16	17	26	18	9	2	2	-
Female	-	3	-	-	3	8	26	32	17	5	1	-
Black, both sexes	-	4	3	5	11	11	13	12	5	-	1	-
Male	-	3	1	4	8	7	5	8	2	-	-	-
Female	-	1	2	1	3	4	8	4	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
South Dakota												
All races, both sexes	-	-	-	1	2	6	4	16	6	8	-	-
Male	-	-	-	-	1	5	3	9	3	3	-	-
Female	-	-	-	1	1	1	1	7	3	5	-	-
White, both sexes	-	-	-	1	2	5	4	16	5	8	-	-
Male	-	-	-	-	1	4	3	9	3	3	-	-
Female	-	-	-	1	1	1	1	7	2	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	14	18	21	29	54	82	121	64	21	7	-
Male	4	7	10	13	21	30	45	52	28	9	1	-
Female	2	7	8	8	8	24	37	69	36	12	6	-
White, both sexes	4	13	15	18	25	50	78	114	61	17	5	-
Male	2	7	7	10	18	28	43	48	26	9	1	-
Female	2	6	8	8	7	22	35	66	35	8	4	-
Black, both sexes	2	1	3	3	4	4	4	7	2	4	2	-
Male	2	-	3	3	3	2	2	4	1	-	-	-
Female	-	1	-	-	1	2	2	3	1	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Texas												
All races, both sexes	20	27	38	52	90	122	261	253	177	94	20	-
Male	14	19	23	36	57	60	147	137	83	31	5	-
Female	6	8	15	16	33	62	114	116	94	63	15	-
White, both sexes	17	24	34	41	82	114	237	234	168	86	18	-
Male	13	19	21	27	53	55	129	125	78	29	4	-
Female	4	5	13	14	29	59	108	109	90	57	14	-
Black, both sexes	3	3	4	11	8	8	24	14	9	8	2	-
Male	1	-	2	9	4	5	18	7	5	2	1	-
Female	2	3	2	2	4	3	6	7	4	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	4	-	-	-	-
Male	-	-	-	-	-	-	-	4	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	3	8	3	-	12	10	16	22	13	4	-	-
Male	2	4	1	-	7	3	8	18	7	2	-	-
Female	1	4	2	-	5	7	8	4	6	2	-	-
White, both sexes	3	8	3	-	12	10	16	21	13	4	-	-
Male	2	4	1	-	7	3	8	17	7	2	-	-
Female	1	4	2	-	5	7	8	4	6	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Vermont												
All races, both sexes	-	-	1	-	-	3	9	8	8	6	1	-
Male	-	-	-	-	-	2	4	5	3	2	1	-
Female	-	-	1	-	-	1	5	3	5	4	-	-
White, both sexes	-	-	1	-	-	3	9	8	8	6	1	-
Male	-	-	-	-	-	2	4	5	3	2	1	-
Female	-	-	1	-	-	1	5	3	5	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	10	12	14	25	51	82	92	129	93	34	7	-
Male	8	5	9	20	32	51	50	63	32	16	1	-
Female	2	7	5	5	19	31	42	66	61	18	6	-
White, both sexes	7	8	11	18	36	59	72	110	79	30	4	-
Male	6	4	6	13	23	34	41	53	29	15	1	-
Female	1	4	5	5	13	25	31	57	50	15	3	-
Black, both sexes	3	3	3	7	15	20	17	19	13	3	2	-
Male	2	1	3	7	9	15	7	10	2	-	-	-
Female	1	2	-	-	6	5	10	9	11	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	2	3	-	-	1	1	-
Male	-	-	-	-	-	1	2	-	-	1	-	-
Female	-	1	-	-	-	1	1	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Washington												
All races, both sexes	6	10	17	24	22	33	76	109	85	36	7	-
Male	2	7	7	14	16	14	35	61	42	12	2	-
Female	4	3	10	10	6	19	41	48	43	24	5	-
White, both sexes	5	8	16	22	19	30	70	101	80	36	7	-
Male	2	5	7	13	15	13	30	55	39	12	2	-
Female	3	3	9	9	4	17	40	46	41	24	5	-
Black, both sexes	1	1	1	-	1	-	3	5	1	-	-	-
Male	-	1	-	-	1	-	2	4	1	-	-	-
Female	1	-	1	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	2	-	1	-	1	1	-	-	-
Male	-	1	-	1	-	-	-	1	1	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	2	3	2	3	-	-	-
Male	-	-	-	-	-	1	3	1	1	-	-	-
Female	-	-	-	-	2	1	-	1	2	-	-	-
West Virginia												
All races, both sexes	1	4	2	7	8	19	20	27	22	8	2	-
Male	1	4	1	6	7	9	9	11	10	4	1	-
Female	-	-	1	1	1	10	11	16	12	4	1	-
White, both sexes	1	4	2	7	8	19	20	27	22	7	2	-
Male	1	4	1	6	7	9	9	11	10	3	1	-
Female	-	-	1	1	1	10	11	16	12	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Wisconsin												
All races, both sexes	5	6	14	14	13	37	78	90	64	20	2	-
Male	3	5	8	10	10	24	49	52	32	10	-	-
Female	2	1	6	4	3	13	29	38	32	10	2	-
White, both sexes	5	6	12	14	11	34	78	89	63	20	2	-
Male	3	5	7	10	9	23	49	51	31	10	-	-
Female	2	1	5	4	2	11	29	38	32	10	2	-
Black, both sexes	-	-	1	-	2	3	-	1	1	-	-	-
Male	-	-	1	-	1	1	-	1	1	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	5	1	6	7	7	1	-	-
Male	-	-	1	-	3	1	5	2	3	-	-	-
Female	-	-	-	-	2	-	1	5	4	1	-	-
White, both sexes	-	-	1	-	5	1	6	7	7	1	-	-
Male	-	-	1	-	3	1	5	2	3	-	-	-
Female	-	-	-	-	2	-	1	5	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
United States												
All races, both sexes	7	1	1	-	-	-	-	-	-	-	-	-
Male	4	-	1	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Hawaii												
All races, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Texas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Wisconsin												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
United States												
All races, both sexes	-	2	1	-	-	1	-	-	1	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	1	1	-	-	1	-	-	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
North Carolina												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Texas												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Wisconsin												
All races, both sexes	-	-	1	-	-	1	-	-	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	1	-	-	1	-	-	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
United States												
All races, both sexes	88	-	-	-	-	-	-	1	-	1	2	2
Male	51	-	-	-	-	-	-	-	-	-	1	-
Female	37	-	-	-	-	-	-	1	-	1	1	2
White, both sexes	81	-	-	-	-	-	-	1	-	1	2	2
Male	46	-	-	-	-	-	-	-	-	-	1	-
Female	35	-	-	-	-	-	-	1	-	1	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Arkansas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Colorado												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Florida												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Minnesota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Missouri												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Nebraska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
New Mexico												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
North Carolina												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Ohio												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Oregon												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Rhode Island												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Texas												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
Wisconsin												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
United States												
All races, both sexes	6	8	3	10	16	14	16	8	1	-	-	-
Male	4	5	3	6	10	7	9	5	1	-	-	-
Female	2	3	-	4	6	7	7	3	-	-	-	-
White, both sexes	4	8	3	6	16	13	16	8	1	-	-	-
Male	2	5	3	4	10	6	9	5	1	-	-	-
Female	2	3	-	2	6	7	7	3	-	-	-	-
Black, both sexes	2	-	-	2	-	1	-	-	-	-	-	-
Male	2	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Colorado												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Florida												
All races, both sexes	-	-	-	1	1	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	1	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Hawaii												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Illinois												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	2	-	1	1	1	-	-	-	-	-	-
Male	-	1	-	1	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	2	-	1	1	1	-	-	-	-	-	-
Male	-	1	-	1	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	2	-	-	2	-	2	-	-	-	-	-
Male	1	2	-	-	2	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	2	-	-	2	-	2	-	-	-	-	-
Male	-	2	-	-	2	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Minnesota												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Missouri												
All races, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
New Mexico												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	1	-	1	2	-	-	-	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	2	-	-	-	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
North Carolina												
All races, both sexes	1	-	-	1	1	2	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	2	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Ohio												
All races, both sexes	-	1	-	1	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	1	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Oregon												
All races, both sexes	-	-	-	-	-	2	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	2	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	-	1	-	-	-	4	-	-	-	-
Male	1	-	-	1	-	-	-	3	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	4	-	-	-	-
Male	1	-	-	-	-	-	-	3	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Rhode Island												
All races, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Texas												
All races, both sexes	1	-	1	-	-	-	2	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	1	-	1	-	-	-	2	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Vermont												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	2	-	1	-	-	-	-	-
Male	-	-	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	2	-	1	-	-	-	-	-
Male	-	-	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
Wisconsin												
All races, both sexes	-	-	-	-	2	1	3	-	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	2	1	3	-	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
United States												
All races, both sexes	705	10	7	2	3	3	8	4	8	17	38	43
Male	408	3	1	1	-	2	5	1	4	12	20	22
Female	297	7	6	1	3	1	3	3	4	5	18	21
White, both sexes	585	7	4	2	-	2	5	2	4	11	26	29
Male	348	2	-	1	-	2	3	1	1	7	14	15
Female	237	5	4	1	-	-	2	1	3	4	12	14
Black, both sexes	102	3	2	-	3	1	3	2	4	6	12	13
Male	49	1	1	-	-	-	2	-	3	5	6	6
Female	53	2	1	-	3	1	1	2	1	1	6	7
American Indian 1/, both sexes	6	-	1	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	14	-	1	-	-	-	-	1	-	1	1	-
Male	10	-	1	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	11	-	-	-	-	-	-	-	-	1	1	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	1	-	-	-	-	1	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Arkansas												
All races, both sexes	10	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	59	3	1	-	-	-	-	-	2	2	2	4
Male	34	-	-	-	-	-	-	-	1	2	2	3
Female	25	3	1	-	-	-	-	-	1	-	-	1
White, both sexes	52	2	1	-	-	-	-	-	2	2	1	3
Male	31	-	-	-	-	-	-	-	1	2	1	3
Female	21	2	1	-	-	-	-	-	1	-	-	-
Black, both sexes	6	1	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	4	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Colorado												
All races, both sexes	19	-	-	-	-	-	1	-	-	-	2	-
Male	10	-	-	-	-	-	1	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	18	-	-	-	-	-	1	-	-	-	2	-
Male	9	-	-	-	-	-	1	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Delaware												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Florida												
All races, both sexes	40	1	1	-	-	-	-	-	-	1	-	4
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	20	1	1	-	-	-	-	-	-	1	-	2
White, both sexes	35	-	-	-	-	-	-	-	-	1	-	3
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	15	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	5	1	1	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	1	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	3	1
Male	16	-	-	-	-	-	-	-	-	-	2	-
Female	9	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Hawaii												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Illinois												
All races, both sexes	28	1	-	-	-	-	-	-	-	2	1	1
Male	18	1	-	-	-	-	-	-	-	2	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	1	-	-	-	-	-	-	-	-	1	1
Male	14	1	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	2	-	-
Male	4	-	-	-	-	-	-	-	-	2	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	15	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Iowa												
All races, both sexes	11	-	-	-	-	-	-	-	-	1	1	1
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	11	-	-	-	-	-	-	-	-	1	1	1
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	7	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Kentucky												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	13	-	-	-	-	-	1	-	1	-	-	3
Male	8	-	-	-	-	-	-	-	1	-	-	2
Female	5	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	1	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	1	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Maine												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	2	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	2	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	3	-
Male	8	-	-	-	-	-	-	-	-	-	2	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	3	-
Male	5	-	-	-	-	-	-	-	-	-	2	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Massachusetts												
All races, both sexes	13	-	-	-	-	-	-	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	13	-	-	-	-	-	-	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	36	-	1	-	-	-	-	1	-	-	-	3
Male	22	-	-	-	-	-	-	1	-	-	-	3
Female	14	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	1	-	-	-	3
Male	19	-	-	-	-	-	-	1	-	-	-	3
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Minnesota												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	8	-	-	-	1	-	-	1	1	1	-	-
Male	5	-	-	-	-	-	-	-	1	1	-	-
Female	3	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	4	-	-	-	1	-	-	-	1	1	-	-
Male	3	-	-	-	-	-	-	-	1	1	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Missouri												
All races, both sexes	11	-	-	-	-	1	-	-	-	-	1	2
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	1	-	-	-	-	1	2
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	3	-	-	-	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Nebraska												
All races, both sexes	7	-	-	-	-	1	-	-	-	-	-	-
Male	6	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	1	-	-	-	-	-	-
Male	6	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	-	-	-	-	-	1	-	-	2	-	-
Male	7	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	8	-	-	-	-	-	1	-	-	1	-	-
Male	7	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
New Hampshire												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	2
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
New Mexico												
All races, both sexes	4	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	40	-	-	-	-	-	1	-	-	-	2	-
Male	24	-	-	-	-	-	1	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	33	-	-	-	-	-	-	-	-	-	1	-
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	1	-	-	-	1	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
North Carolina												
All races, both sexes	24	1	-	-	1	-	1	-	1	-	-	4
Male	10	1	-	-	-	-	-	-	-	-	-	2
Female	14	-	-	-	1	-	1	-	1	-	-	2
White, both sexes	17	1	-	-	-	-	-	-	-	-	-	2
Male	7	1	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	1	-	1	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	1	-	1	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Ohio												
All races, both sexes	25	-	-	-	-	-	-	1	-	2	1	1
Male	14	-	-	-	-	-	-	-	-	2	1	-
Female	11	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	21	-	-	-	-	-	-	-	-	1	1	1
Male	13	-	-	-	-	-	-	-	-	1	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	1	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	12	-	-	-	-	-	-	-	-	2	1	2
Male	7	-	-	-	-	-	-	-	-	1	-	2
Female	5	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	10	-	-	-	-	-	-	-	-	2	1	2
Male	6	-	-	-	-	-	-	-	-	1	-	2
Female	4	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Oregon												
All races, both sexes	6	1	1	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	6	1	1	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	2	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	2	-
Male	16	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Rhode Island												
All races, both sexes	7	-	1	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	1	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	12	1	-	-	-	-	-	-	-	-	4	1
Male	8	1	-	-	-	-	-	-	-	-	3	-
Female	4	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	7	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	2	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	1	-	-	-	-	-	-	-	-	2	-
Male	3	1	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	3
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	3
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Texas												
All races, both sexes	63	2	-	1	1	1	2	-	-	1	2	2
Male	40	-	-	1	-	1	1	-	-	1	-	2
Female	23	2	-	-	1	-	1	-	-	-	2	-
White, both sexes	55	2	-	1	-	1	2	-	-	-	2	1
Male	35	-	-	1	-	1	1	-	-	-	-	1
Female	20	2	-	-	-	-	1	-	-	-	2	-
Black, both sexes	4	-	-	-	1	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	1
Female	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Vermont												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	19	-	-	-	-	-	-	-	1	-	4	1
Male	12	-	-	-	-	-	-	-	1	-	3	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	3	1
Male	9	-	-	-	-	-	-	-	-	-	2	-
Female	5	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	5	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	1	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Washington												
All races, both sexes	15	-	-	-	-	-	-	-	1	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	14	-	-	-	-	-	-	-	1	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Wisconsin												
All races, both sexes	11	-	-	-	-	-	1	-	-	-	-	1
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	1	-	-	-	-	1
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
United States												
All races, both sexes	57	43	48	66	74	77	89	74	26	7	1	-
Male	38	30	31	44	41	43	52	45	11	2	-	-
Female	19	13	17	22	33	34	37	29	15	5	1	-
White, both sexes	49	34	44	61	60	70	76	68	25	6	-	-
Male	33	25	28	42	36	40	44	42	11	1	-	-
Female	16	9	16	19	24	30	32	26	14	5	-	-
Black, both sexes	7	8	4	3	10	5	10	4	1	-	1	-
Male	4	4	3	1	3	2	6	2	-	-	-	-
Female	3	4	1	2	7	3	4	2	1	-	1	-
American Indian 1/, both sexes	1	-	-	-	1	1	1	-	-	-	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	3	1	2	2	-	1	-	-
Male	-	1	-	1	2	-	1	1	-	1	-	-
Female	-	-	-	1	1	1	1	1	-	-	-	-
Alabama												
All races, both sexes	1	-	-	2	1	1	-	-	1	-	-	-
Male	1	-	-	2	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	-	-	2	1	1	-	-	1	-	-	-
Male	1	-	-	2	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Alaska												
All races, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	1	2	-	3	1	1	1	1	-	-	-
Male	-	1	2	-	1	1	1	1	1	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
White, both sexes	-	1	2	-	3	-	1	1	1	-	-	-
Male	-	1	2	-	1	-	1	1	1	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Arkansas												
All races, both sexes	2	-	1	1	1	-	1	2	-	-	-	-
Male	2	-	-	1	1	-	1	2	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	1	1	-	1	1	-	-	-	-
Male	2	-	-	1	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	8	2	5	4	6	6	7	5	1	1	-	-
Male	5	1	3	4	5	3	2	2	1	-	-	-
Female	3	1	2	-	1	3	5	3	-	1	-	-
White, both sexes	6	2	5	3	6	6	6	5	1	1	-	-
Male	4	1	3	3	5	3	2	2	1	-	-	-
Female	2	1	2	-	1	3	4	3	-	1	-	-
Black, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Colorado												
All races, both sexes	2	2	2	4	2	-	1	3	-	-	-	-
Male	-	2	2	2	-	-	-	2	-	-	-	-
Female	2	-	-	2	2	-	1	1	-	-	-	-
White, both sexes	2	2	2	4	2	-	1	2	-	-	-	-
Male	-	2	2	2	-	-	-	1	-	-	-	-
Female	2	-	-	2	2	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	1	1	-	2	1	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	1	-	-	-	1	1	-	2	1	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Delaware												
All races, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Florida												
All races, both sexes	-	4	2	4	3	6	7	6	1	-	-	-
Male	-	3	2	2	1	3	4	3	-	-	-	-
Female	-	1	-	2	2	3	3	3	1	-	-	-
White, both sexes	-	4	2	4	2	5	7	6	1	-	-	-
Male	-	3	2	2	1	3	4	3	-	-	-	-
Female	-	1	-	2	1	2	3	3	1	-	-	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	3	4	2	1	5	3	-	-	-	-	-
Male	3	2	2	2	1	2	2	-	-	-	-	-
Female	-	1	2	-	-	3	1	-	-	-	-	-
White, both sexes	3	2	3	2	-	5	2	-	-	-	-	-
Male	3	1	1	2	-	2	1	-	-	-	-	-
Female	-	1	2	-	-	3	1	-	-	-	-	-
Black, both sexes	-	1	1	-	1	-	1	-	-	-	-	-
Male	-	1	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Hawaii												
All races, both sexes	-	1	-	-	2	1	-	-	-	1	-	-
Male	-	1	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	2	1	-	-	-	1	-	-
Male	-	1	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Idaho												
All races, both sexes	-	1	-	1	2	-	1	-	-	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
White, both sexes	-	1	-	1	2	-	1	-	-	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Illinois												
All races, both sexes	2	1	2	4	6	2	3	1	2	-	-	-
Male	2	-	1	3	2	2	3	-	1	-	-	-
Female	-	1	1	1	4	-	-	1	1	-	-	-
White, both sexes	1	-	2	4	5	1	3	1	2	-	-	-
Male	1	-	1	3	2	1	3	-	1	-	-	-
Female	-	-	1	1	3	-	-	1	1	-	-	-
Black, both sexes	1	1	-	-	1	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	1	-	2	-	2	4	-	2	-	-	-
Male	1	1	-	2	-	-	3	-	-	-	-	-
Female	1	-	-	-	-	2	1	-	2	-	-	-
White, both sexes	2	1	-	2	-	1	4	-	2	-	-	-
Male	1	1	-	2	-	-	3	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Iowa												
All races, both sexes	-	2	1	-	-	-	2	3	-	-	-	-
Male	-	2	-	-	-	-	1	2	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
White, both sexes	-	2	1	-	-	-	2	3	-	-	-	-
Male	-	2	-	-	-	-	1	2	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	-	-	1	2	1	-	-	-	-	-	-
Male	2	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
White, both sexes	2	-	-	1	-	1	-	-	-	-	-	-
Male	2	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Kentucky												
All races, both sexes	3	-	-	1	-	1	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	2	-	-	1	-	1	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	2	-	-	2	3	-	-	-	-
Male	1	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
White, both sexes	1	-	-	2	-	-	2	2	-	-	-	-
Male	1	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Maine												
All races, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	1	1	2	2	-	3	-	-
Male	1	-	-	-	-	1	2	1	-	1	-	-
Female	-	-	-	-	1	-	-	1	-	2	-	-
White, both sexes	-	-	-	-	-	1	-	2	-	3	-	-
Male	-	-	-	-	-	1	-	1	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	2	-	-
Black, both sexes	1	-	-	-	1	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Massachusetts												
All races, both sexes	-	1	1	1	1	3	1	1	1	-	-	-
Male	-	1	1	-	1	2	-	1	-	-	-	-
Female	-	-	-	1	-	1	1	-	1	-	-	-
White, both sexes	-	1	1	1	1	3	1	1	1	-	-	-
Male	-	1	1	-	1	2	-	1	-	-	-	-
Female	-	-	-	1	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4	2	-	2	3	5	9	2	3	1	-	-
Male	3	-	-	1	2	4	5	1	2	-	-	-
Female	1	2	-	1	1	1	4	1	1	1	-	-
White, both sexes	3	2	-	2	2	4	7	2	3	1	-	-
Male	2	-	-	1	2	3	4	1	2	-	-	-
Female	1	2	-	1	-	1	3	1	1	1	-	-
Black, both sexes	1	-	-	-	1	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Minnesota												
All races, both sexes	-	-	-	-	-	1	3	3	3	-	-	-
Male	-	-	-	-	-	-	1	3	1	-	-	-
Female	-	-	-	-	-	1	2	-	2	-	-	-
White, both sexes	-	-	-	-	-	1	2	3	3	-	-	-
Male	-	-	-	-	-	-	1	3	1	-	-	-
Female	-	-	-	-	-	1	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Missouri												
All races, both sexes	-	1	-	1	3	1	1	-	-	-	-	-
Male	-	-	-	1	3	1	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	3	1	1	-	-	-	-	-
Male	-	-	-	1	3	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Nebraska												
All races, both sexes	1	-	-	1	-	-	2	1	1	-	-	-
Male	1	-	-	1	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	2	1	1	-	-	-
Male	1	-	-	1	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	-	-	2	1	1	1	-	-	-	-
Male	-	1	-	-	2	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	2	1	1	1	-	-	-	-
Male	-	1	-	-	2	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
New Hampshire												
All races, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	2	2	2	1	-	-	-
Male	-	-	-	-	-	1	1	2	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	2	2	2	1	-	-	-
Male	-	-	-	-	-	1	1	2	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
New Mexico												
All races, both sexes	-	-	1	-	-	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	-	5	4	3	8	6	5	1	-	1	-
Male	2	-	4	2	2	5	4	3	-	-	-	-
Female	2	-	1	2	1	3	2	2	1	-	1	-
White, both sexes	4	-	5	3	3	8	5	4	-	-	-	-
Male	2	-	4	2	2	5	3	2	-	-	-	-
Female	2	-	1	1	1	3	2	2	-	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	1	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
North Carolina												
All races, both sexes	1	2	3	1	2	2	2	3	-	-	-	-
Male	-	2	1	-	-	-	1	3	-	-	-	-
Female	1	-	2	1	2	2	1	-	-	-	-	-
White, both sexes	1	1	2	1	2	2	2	3	-	-	-	-
Male	-	1	-	-	-	-	1	3	-	-	-	-
Female	1	-	2	1	2	2	1	-	-	-	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Ohio												
All races, both sexes	3	2	1	3	3	2	3	2	1	-	-	-
Male	2	1	1	2	2	1	1	1	-	-	-	-
Female	1	1	-	1	1	1	2	1	1	-	-	-
White, both sexes	3	2	1	3	2	2	2	2	1	-	-	-
Male	2	1	1	2	2	1	1	1	-	-	-	-
Female	1	1	-	1	-	1	1	1	1	-	-	-
Black, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	1	1	-	-	-	1	1	-	-	-	-
Male	2	1	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	1	1	1	-	-	-	1	1	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Oregon												
All races, both sexes	1	-	-	-	1	2	-	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	2	-	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	3	1	2	4	7	6	4	1	-	-	-
Male	-	1	1	2	3	3	4	2	-	-	-	-
Female	-	2	-	-	1	4	2	2	1	-	-	-
White, both sexes	-	2	1	2	4	7	5	3	1	-	-	-
Male	-	1	1	2	3	3	3	2	-	-	-	-
Female	-	1	-	-	1	4	2	1	1	-	-	-
Black, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Rhode Island												
All races, both sexes	-	-	1	2	1	-	-	1	1	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-
White, both sexes	-	-	1	2	1	-	-	1	1	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	2	1	-	2	-	1	-	-	-	-	-
Male	-	1	1	-	1	-	1	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	2	-	1	-	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	1	-	2	1	4	1	1	1	-	-	-
Male	-	1	-	2	-	2	-	-	1	-	-	-
Female	1	-	-	-	1	2	1	1	-	-	-	-
White, both sexes	1	-	-	1	1	4	1	1	1	-	-	-
Male	-	-	-	1	-	2	-	-	1	-	-	-
Female	1	-	-	-	1	2	1	1	-	-	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Texas												
All races, both sexes	6	3	4	10	8	6	7	5	2	-	-	-
Male	5	3	2	5	5	5	5	3	1	-	-	-
Female	1	-	2	5	3	1	2	2	1	-	-	-
White, both sexes	6	3	4	9	7	6	4	5	2	-	-	-
Male	5	3	2	5	4	5	3	3	1	-	-	-
Female	1	-	2	4	3	1	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	2	-	-	-	-	-
White, both sexes	-	-	1	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Vermont												
All races, both sexes	1	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	2	2	3	1	-	2	-	1	-	-
Male	1	1	1	1	3	-	-	1	-	-	-	-
Female	-	-	1	1	-	1	-	1	-	1	-	-
White, both sexes	1	1	2	1	2	-	-	2	-	1	-	-
Male	1	1	1	1	2	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Washington												
All races, both sexes	2	2	1	2	-	1	3	2	-	-	-	-
Male	2	-	1	1	-	1	1	1	-	-	-	-
Female	-	2	-	1	-	-	2	1	-	-	-	-
White, both sexes	2	2	1	2	-	1	3	1	-	-	-	-
Male	2	-	1	1	-	1	1	1	-	-	-	-
Female	-	2	-	1	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	-	1	-	1	-	-	-	3	-	-	-	-
Male	-	1	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	1	-	1	-	-	-	3	-	-	-	-
Male	-	1	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Wisconsin												
All races, both sexes	-	-	2	1	1	1	1	3	-	-	-	-
Male	-	-	-	1	-	-	1	2	-	-	-	-
Female	-	-	2	-	1	1	-	1	-	-	-	-
White, both sexes	-	-	2	1	-	1	1	3	-	-	-	-
Male	-	-	-	1	-	-	1	2	-	-	-	-
Female	-	-	2	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
United States												
All races, both sexes	1,024	1	-	-	-	-	-	1	1	1	3	18
Male	441	1	-	-	-	-	-	1	1	-	2	12
Female	583	-	-	-	-	-	-	-	-	1	1	6
White, both sexes	915	-	-	-	-	-	-	-	1	1	1	13
Male	392	-	-	-	-	-	-	-	1	-	-	9
Female	523	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	92	1	-	-	-	-	-	1	-	-	2	5
Male	42	1	-	-	-	-	-	1	-	-	2	3
Female	50	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Arkansas												
All races, both sexes	7	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	2
Male	26	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	2
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Colorado												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Delaware												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Florida												
All races, both sexes	45	-	-	-	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	40	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	31	1	-	-	-	-	-	-	-	-	-	1
Male	11	1	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Hawaii												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Illinois												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	1	-
Male	28	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Iowa												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Kentucky												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Maine												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Massachusetts												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	45	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Minnesota												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	17	-	-	-	-	-	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	1	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Missouri												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Nebraska												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
New Hampshire												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	-	2
Male	18	-	-	-	-	-	-	-	-	-	-	2
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	44	-	-	-	-	-	-	-	-	-	-	2
Male	18	-	-	-	-	-	-	-	-	-	-	2
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
New Mexico												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	70	-	-	-	-	-	-	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	61	-	-	-	-	-	-	-	-	-	-	1
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
North Carolina												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	1	5
Male	19	-	-	-	-	-	-	-	-	-	-	3
Female	19	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	29	-	-	-	-	-	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	3
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Ohio												
All races, both sexes	59	-	-	-	-	-	-	-	-	1	-	2
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	36	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	54	-	-	-	-	-	-	-	-	1	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Oregon												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	77	-	-	-	-	-	-	-	-	-	-	-
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	72	-	-	-	-	-	-	-	-	-	-	-
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Rhode Island												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
South Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Texas												
All races, both sexes	61	-	-	-	-	-	-	-	-	-	1	3
Male	26	-	-	-	-	-	-	-	-	-	1	3
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	57	-	-	-	-	-	-	-	-	-	-	3
Male	24	-	-	-	-	-	-	-	-	-	-	3
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Washington												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
Wisconsin												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
United States												
All races, both sexes	17	22	42	49	82	121	210	222	163	62	9	-
Male	10	9	23	21	43	49	92	98	61	17	1	-
Female	7	13	19	28	39	72	118	124	102	45	8	-
White, both sexes	12	18	36	40	70	110	184	211	152	57	9	-
Male	7	9	19	17	38	46	81	94	55	15	1	-
Female	5	9	17	23	32	64	103	117	97	42	8	-
Black, both sexes	5	4	6	6	12	8	19	11	8	4	-	-
Male	3	-	4	3	5	3	7	4	5	1	-	-
Female	2	4	2	3	7	5	12	7	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	-	3	7	-	2	1	-	-
Male	-	-	-	1	-	-	4	-	1	1	-	-
Female	-	-	-	2	-	3	3	-	1	-	-	-
Alabama												
All races, both sexes	-	-	-	1	1	-	4	6	3	1	-	-
Male	-	-	-	1	-	-	-	3	-	1	-	-
Female	-	-	-	-	1	-	4	3	3	-	-	-
White, both sexes	-	-	-	1	1	-	2	5	3	-	-	-
Male	-	-	-	1	-	-	-	3	-	-	-	-
Female	-	-	-	-	1	-	2	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	2	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	1	1	2	1	5	2	2	-	-	-
Male	-	-	1	-	1	-	-	1	2	-	-	-
Female	1	-	-	1	1	1	5	1	-	-	-	-
White, both sexes	1	-	1	1	2	1	5	2	2	-	-	-
Male	-	-	1	-	1	-	-	1	2	-	-	-
Female	1	-	-	1	1	1	5	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Arkansas												
All races, both sexes	-	-	-	-	1	-	3	1	1	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	3	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	2	1	1	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	3	3	4	1	3	6	12	8	2	1	-
Male	1	-	3	3	1	2	3	9	2	1	-	-
Female	-	3	-	1	-	1	3	3	6	1	1	-
White, both sexes	1	2	3	4	-	3	5	11	8	2	1	-
Male	1	-	3	3	-	2	2	8	2	1	-	-
Female	-	2	-	1	-	1	3	3	6	1	1	-
Black, both sexes	-	1	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Colorado												
All races, both sexes	1	-	-	-	-	2	2	3	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	2	3	-	1	-	-
White, both sexes	1	-	-	-	-	1	2	3	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	1	-	4	4	9	1	-	-
Male	-	-	-	-	1	-	1	2	3	-	-	-
Female	-	-	-	-	-	-	3	2	6	1	-	-
White, both sexes	-	-	-	-	1	-	4	4	9	1	-	-
Male	-	-	-	-	1	-	1	2	3	-	-	-
Female	-	-	-	-	-	-	3	2	6	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Delaware												
All races, both sexes	-	-	-	-	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	1	1	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	1	-	-	1	-	1	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Florida												
All races, both sexes	1	2	1	1	5	6	11	9	4	3	2	-
Male	1	2	-	1	-	4	9	7	2	2	-	-
Female	-	-	1	-	5	2	2	2	2	1	2	-
White, both sexes	1	2	1	1	3	5	9	9	4	3	2	-
Male	1	2	-	1	-	4	7	7	2	2	-	-
Female	-	-	1	-	3	1	2	2	2	1	2	-
Black, both sexes	-	-	-	-	2	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Georgia												
All races, both sexes	-	1	3	1	2	3	3	5	6	5	-	-
Male	-	-	2	1	1	-	-	3	2	1	-	-
Female	-	1	1	-	1	3	3	2	4	4	-	-
White, both sexes	-	-	1	-	2	3	3	5	5	5	-	-
Male	-	-	1	-	1	-	-	3	1	1	-	-
Female	-	-	-	-	1	3	3	2	4	4	-	-
Black, both sexes	-	1	2	1	-	-	-	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	1	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Hawaii												
All races, both sexes	-	-	-	1	-	1	4	1	1	1	-	-
Male	-	-	-	1	-	-	2	1	1	1	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	4	-	1	1	-	-
Male	-	-	-	1	-	-	2	-	1	1	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Illinois												
All races, both sexes	-	-	3	3	5	6	8	19	8	1	-	-
Male	-	-	1	2	2	4	6	9	3	-	-	-
Female	-	-	2	1	3	2	2	10	5	1	-	-
White, both sexes	-	-	2	3	4	5	7	18	8	1	-	-
Male	-	-	1	2	2	3	5	8	3	-	-	-
Female	-	-	1	1	2	2	2	10	5	1	-	-
Black, both sexes	-	-	1	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	1	1	2	4	6	1	6	3	-	-
Male	-	-	1	-	1	-	3	-	1	-	-	-
Female	-	-	-	1	1	4	3	1	5	3	-	-
White, both sexes	-	-	1	1	2	4	5	1	5	3	-	-
Male	-	-	1	-	1	-	2	-	-	-	-	-
Female	-	-	-	1	1	4	3	1	5	3	-	-
Black, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Iowa												
All races, both sexes	-	-	-	-	1	1	4	1	1	-	-	-
Male	-	-	-	-	-	1	2	1	1	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	4	1	1	-	-	-
Male	-	-	-	-	-	1	2	1	1	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	2	1	-	2	1	1	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	2	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	2	1	-	2	1	1	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	2	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Kentucky												
All races, both sexes	-	-	2	2	2	1	7	2	2	-	-	-
Male	-	-	1	2	1	1	1	2	-	-	-	-
Female	-	-	1	-	1	-	6	-	2	-	-	-
White, both sexes	-	-	2	1	2	1	7	2	2	-	-	-
Male	-	-	1	1	1	1	1	2	-	-	-	-
Female	-	-	1	-	1	-	6	-	2	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	2	2	1	-	3	2	1	-	-
Male	-	-	-	1	2	1	-	2	1	-	-	-
Female	-	-	-	1	-	-	-	1	1	1	-	-
White, both sexes	-	-	-	2	2	1	-	3	2	1	-	-
Male	-	-	-	1	2	1	-	2	1	-	-	-
Female	-	-	-	1	-	-	-	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Maine												
All races, both sexes	-	-	-	-	-	1	1	2	1	2	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	1	1	1	1	1	-	-
White, both sexes	-	-	-	-	-	1	1	2	1	2	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	1	1	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	1	1	2	2	5	8	6	-	-	-
Male	-	-	1	1	2	2	1	3	2	-	-	-
Female	-	-	-	-	-	-	4	5	4	-	-	-
White, both sexes	-	-	-	1	1	1	4	6	4	-	-	-
Male	-	-	-	1	1	1	1	2	1	-	-	-
Female	-	-	-	-	-	-	3	4	3	-	-	-
Black, both sexes	-	-	1	-	1	1	1	2	1	-	-	-
Male	-	-	1	-	1	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Massachusetts												
All races, both sexes	-	-	-	-	2	3	4	2	6	1	-	-
Male	-	-	-	-	1	1	2	1	5	-	-	-
Female	-	-	-	-	1	2	2	1	1	1	-	-
White, both sexes	-	-	-	-	1	3	3	2	6	1	-	-
Male	-	-	-	-	-	1	2	1	5	-	-	-
Female	-	-	-	-	1	2	1	1	1	1	-	-
Black, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	3	3	3	3	11	6	12	2	2	-	-
Male	-	1	1	-	2	7	3	4	1	2	-	-
Female	-	2	2	3	1	4	3	8	1	-	-	-
White, both sexes	-	2	2	3	2	11	5	12	2	2	-	-
Male	-	1	-	-	1	7	2	4	1	2	-	-
Female	-	1	2	3	1	4	3	8	1	-	-	-
Black, both sexes	-	1	1	-	1	-	1	-	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Minnesota												
All races, both sexes	-	-	-	1	-	1	4	4	2	1	-	-
Male	-	-	-	-	-	1	1	1	-	1	-	-
Female	-	-	-	1	-	-	3	3	2	-	-	-
White, both sexes	-	-	-	1	-	1	4	4	2	1	-	-
Male	-	-	-	-	-	1	1	1	-	1	-	-
Female	-	-	-	1	-	-	3	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	-	1	1	1	2	4	2	1	1	-	-
Male	2	-	-	-	1	-	4	1	-	-	-	-
Female	-	-	1	1	-	2	-	1	1	1	-	-
White, both sexes	1	-	1	1	1	1	4	2	-	-	-	-
Male	1	-	-	-	1	-	4	1	-	-	-	-
Female	-	-	1	1	-	1	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	1	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Missouri												
All races, both sexes	1	2	1	1	1	1	5	7	-	-	-	-
Male	1	1	-	-	1	1	1	2	-	-	-	-
Female	-	1	1	1	-	-	4	5	-	-	-	-
White, both sexes	1	2	1	-	1	1	4	6	-	-	-	-
Male	1	1	-	-	1	1	1	2	-	-	-	-
Female	-	1	1	-	-	-	3	4	-	-	-	-
Black, both sexes	-	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Nebraska												
All races, both sexes	-	-	2	-	1	3	-	3	2	-	-	-
Male	-	-	2	-	-	1	-	2	1	-	-	-
Female	-	-	-	-	1	2	-	1	1	-	-	-
White, both sexes	-	-	2	-	1	3	-	3	2	-	-	-
Male	-	-	2	-	-	1	-	2	1	-	-	-
Female	-	-	-	-	1	2	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	3	1	1	-	2	1	-	1	-	-
Male	1	-	2	-	-	-	1	-	-	1	-	-
Female	-	-	1	1	1	-	1	1	-	-	-	-
White, both sexes	1	-	3	-	1	-	2	1	-	1	-	-
Male	1	-	2	-	-	-	1	-	-	1	-	-
Female	-	-	1	-	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
New Hampshire												
All races, both sexes	-	-	-	2	1	-	1	2	-	-	-	-
Male	-	-	-	1	1	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	2	1	-	1	2	-	-	-	-
Male	-	-	-	1	1	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	1	1	-	3	2	12	13	9	-	-	-
Male	1	-	1	-	1	-	4	5	4	-	-	-
Female	-	1	-	-	2	2	8	8	5	-	-	-
White, both sexes	1	1	1	-	3	2	12	13	9	-	-	-
Male	1	-	1	-	1	-	4	5	4	-	-	-
Female	-	1	-	-	2	2	8	8	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
New Mexico												
All races, both sexes	-	-	-	-	1	-	1	-	2	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	1	1	3	5	11	15	19	6	6	-	-
Male	-	-	1	-	3	5	6	6	4	2	-	-
Female	2	1	-	3	2	6	9	13	2	4	-	-
White, both sexes	1	-	1	1	3	11	13	18	6	6	-	-
Male	-	-	1	-	2	5	6	6	4	2	-	-
Female	1	-	-	1	1	6	7	12	2	4	-	-
Black, both sexes	1	1	-	1	2	-	2	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	1	-	1	1	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
North Carolina												
All races, both sexes	1	-	2	2	3	3	7	7	4	2	1	-
Male	1	-	2	1	1	1	5	3	-	1	1	-
Female	-	-	-	1	2	2	2	4	4	1	-	-
White, both sexes	-	-	1	1	3	3	5	6	4	2	1	-
Male	-	-	1	-	1	1	3	3	-	1	1	-
Female	-	-	-	1	2	2	2	3	4	1	-	-
Black, both sexes	1	-	1	1	-	-	2	1	-	-	-	-
Male	1	-	1	1	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Ohio												
All races, both sexes	1	2	1	-	5	9	11	9	12	5	1	-
Male	-	-	-	-	2	5	5	3	5	2	-	-
Female	1	2	1	-	3	4	6	6	7	3	1	-
White, both sexes	1	2	1	-	5	9	10	7	11	5	1	-
Male	-	-	-	-	2	5	5	2	4	2	-	-
Female	1	2	1	-	3	4	5	5	7	3	1	-
Black, both sexes	-	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	1	-	-	-	-	2	-	2	1	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	1	-	-	-	-	1	-	1	1	-	-
White, both sexes	-	1	-	-	-	-	2	-	2	1	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	1	-	-	-	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Oregon												
All races, both sexes	-	-	2	1	1	1	2	4	1	1	-	-
Male	-	-	-	-	1	-	1	2	-	-	-	-
Female	-	-	2	1	-	1	1	2	1	1	-	-
White, both sexes	-	-	2	1	1	1	2	4	1	1	-	-
Male	-	-	-	-	1	-	1	2	-	-	-	-
Female	-	-	2	1	-	1	1	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	1	3	1	7	11	16	15	15	4	3	-
Male	1	1	1	-	3	3	9	7	6	-	-	-
Female	-	-	2	1	4	8	7	8	9	4	3	-
White, both sexes	-	1	3	1	6	11	15	14	15	3	3	-
Male	-	1	1	-	3	3	9	7	6	-	-	-
Female	-	-	2	1	3	8	6	7	9	3	3	-
Black, both sexes	1	-	-	-	1	-	1	1	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Rhode Island												
All races, both sexes	-	1	-	-	1	2	3	-	1	-	-	-
Male	-	1	-	-	1	-	3	-	1	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
White, both sexes	-	1	-	-	1	1	3	-	1	-	-	-
Male	-	1	-	-	1	-	3	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	1	2	4	3	5	2	1	-	-
Male	-	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	1	1	3	2	4	2	1	-	-
White, both sexes	-	-	-	1	2	2	3	5	2	1	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	1	1	2	2	4	2	1	-	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
South Dakota												
All races, both sexes	-	-	-	1	-	2	-	1	1	-	-	-
Male	-	-	-	1	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	2	-	1	1	-	-	-
Male	-	-	-	1	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	1	1	2	1	2	8	6	2	1	-	-
Male	-	1	-	1	-	-	3	2	-	-	-	-
Female	-	-	1	1	1	2	5	4	2	1	-	-
White, both sexes	-	1	1	2	1	2	8	6	2	1	-	-
Male	-	1	-	1	-	-	3	2	-	-	-	-
Female	-	-	1	1	1	2	5	4	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Texas												
All races, both sexes	-	1	4	4	8	5	9	9	12	4	1	-
Male	-	1	2	2	5	2	3	3	3	1	-	-
Female	-	-	2	2	3	3	6	6	9	3	1	-
White, both sexes	-	1	4	4	7	5	8	9	11	4	1	-
Male	-	1	2	2	5	2	3	3	2	1	-	-
Female	-	-	2	2	2	3	5	6	9	3	1	-
Black, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	1	1	-	2	2	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	2	2	1	-	-
White, both sexes	-	-	-	-	1	1	-	2	2	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	-	2	3	6	6	7	7	3	-	-
Male	-	1	-	-	2	1	1	2	4	-	-	-
Female	1	-	-	2	1	5	5	5	3	3	-	-
White, both sexes	1	1	-	2	2	5	4	7	6	2	-	-
Male	-	1	-	-	2	1	-	2	4	-	-	-
Female	1	-	-	2	-	4	4	5	2	2	-	-
Black, both sexes	-	-	-	-	1	1	2	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Washington												
All races, both sexes	-	-	-	-	2	3	1	2	3	2	-	-
Male	-	-	-	-	2	1	-	2	2	-	-	-
Female	-	-	-	-	-	2	1	-	1	2	-	-
White, both sexes	-	-	-	-	2	3	1	2	3	2	-	-
Male	-	-	-	-	2	1	-	2	2	-	-	-
Female	-	-	-	-	-	2	1	-	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	1	2	1	2	5	5	1	-	-	-
Male	-	-	1	2	-	1	4	3	-	-	-	-
Female	-	-	-	-	1	1	1	2	1	-	-	-
White, both sexes	-	-	1	2	1	2	5	5	1	-	-	-
Male	-	-	1	2	-	1	4	3	-	-	-	-
Female	-	-	-	-	1	1	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Wisconsin												
All races, both sexes	1	1	-	-	-	2	4	2	3	2	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	1	1	-	-	-	1	2	2	3	2	-	-
White, both sexes	1	1	-	-	-	2	4	2	3	2	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	1	1	-	-	-	1	2	2	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
United States												
All races, both sexes	25,176	294	72	27	27	40	38	71	97	156	274	448
Male	12,497	171	49	14	15	24	26	35	38	84	130	226
Female	12,679	123	23	13	12	16	12	36	59	72	144	222
White, both sexes	22,358	157	49	21	19	31	24	58	76	113	198	324
Male	11,136	87	34	10	10	18	15	26	31	62	93	171
Female	11,222	70	15	11	9	13	9	32	45	51	105	153
Black, both sexes	2,139	119	17	5	7	7	14	11	20	34	64	113
Male	1,016	72	11	3	5	5	11	7	6	17	30	50
Female	1,123	47	6	2	2	2	3	4	14	17	34	63
American Indian 1/, both sexes	214	8	2	1	1	1	-	1	-	3	5	4
Male	104	6	2	1	-	-	-	1	-	1	2	2
Female	110	2	-	-	1	1	-	-	-	2	3	2
Asian or Pacific Islander 2/, both sexes	465	10	4	-	-	1	-	1	1	6	7	7
Male	241	6	2	-	-	1	-	1	1	4	5	3
Female	224	4	2	-	-	-	-	-	-	2	2	4
Alabama												
All races, both sexes	546	2	-	1	-	2	-	1	1	5	7	13
Male	267	2	-	-	-	1	-	1	-	3	4	9
Female	279	-	-	1	-	1	-	-	1	2	3	4
White, both sexes	436	1	-	1	-	-	-	1	1	4	6	8
Male	214	1	-	-	-	-	-	1	-	2	3	6
Female	222	-	-	1	-	-	-	-	1	2	3	2
Black, both sexes	108	1	-	-	-	2	-	-	-	1	1	5
Male	53	1	-	-	-	1	-	-	-	1	1	3
Female	55	-	-	-	-	1	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Alaska												
All races, both sexes	43	1	-	-	-	1	1	-	-	-	1	2
Male	22	1	-	-	-	-	-	-	-	-	-	2
Female	21	-	-	-	-	1	1	-	-	-	1	-
White, both sexes	26	-	-	-	-	-	1	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	14	1	-	-	-	1	-	-	-	-	-	1
Male	7	1	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	486	9	2	1	1	1	1	1	1	6	9	10
Male	267	5	2	1	-	-	1	-	-	3	2	6
Female	219	4	-	-	1	1	-	1	1	3	7	4
White, both sexes	444	8	-	1	-	-	-	1	1	4	6	10
Male	245	4	-	1	-	-	-	-	-	2	2	6
Female	199	4	-	-	-	-	-	1	1	2	4	4
Black, both sexes	11	1	1	-	-	1	1	-	-	-	1	-
Male	5	1	1	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	1	-	-	-	-	1	-
American Indian 1/, both sexes	27	-	1	-	1	-	-	-	-	2	2	-
Male	15	-	1	-	-	-	-	-	-	1	-	-
Female	12	-	-	-	1	-	-	-	-	1	2	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Arkansas												
All races, both sexes	255	-	1	1	1	-	1	1	5	2	2	6
Male	131	-	1	1	1	-	-	-	1	2	1	3
Female	124	-	-	-	-	-	1	1	4	-	1	3
White, both sexes	226	-	-	1	1	-	1	1	4	2	2	4
Male	115	-	-	1	1	-	-	-	-	2	1	2
Female	111	-	-	-	-	-	1	1	4	-	1	2
Black, both sexes	25	-	1	-	-	-	-	-	1	-	-	2
Male	14	-	1	-	-	-	-	-	1	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2,011	33	9	2	6	2	3	4	8	13	26	36
Male	1,021	16	5	-	5	2	2	2	3	8	14	18
Female	990	17	4	2	1	-	1	2	5	5	12	18
White, both sexes	1,753	20	6	2	5	2	2	2	6	11	21	27
Male	905	8	3	-	4	2	1	-	3	7	11	16
Female	848	12	3	2	1	-	1	2	3	4	10	11
Black, both sexes	90	8	-	-	1	-	1	1	2	1	3	6
Male	34	5	-	-	1	-	1	1	-	-	1	1
Female	56	3	-	-	-	-	-	-	2	1	2	5
American Indian 1/, both sexes	16	1	-	-	-	-	-	1	-	-	1	-
Male	11	1	-	-	-	-	-	1	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	152	4	3	-	-	-	-	-	-	1	1	3
Male	71	2	2	-	-	-	-	-	-	1	1	1
Female	81	2	1	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Colorado												
All races, both sexes	351	2	2	-	-	-	-	2	2	2	2	7
Male	197	1	2	-	-	-	-	1	2	1	-	3
Female	154	1	-	-	-	-	-	1	-	1	2	4
White, both sexes	333	1	1	-	-	-	-	1	2	1	2	7
Male	186	-	1	-	-	-	-	-	2	1	-	3
Female	147	1	-	-	-	-	-	1	-	-	2	4
Black, both sexes	14	1	1	-	-	-	-	1	-	1	-	-
Male	9	1	1	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	347	2	1	-	-	-	-	1	2	1	3	7
Male	150	-	1	-	-	-	-	-	1	1	3	3
Female	197	2	-	-	-	-	-	1	1	-	-	4
White, both sexes	325	1	1	-	-	-	-	1	2	1	1	6
Male	138	-	1	-	-	-	-	-	1	1	1	3
Female	187	1	-	-	-	-	-	1	1	-	-	3
Black, both sexes	21	1	-	-	-	-	-	-	-	-	2	1
Male	11	-	-	-	-	-	-	-	-	-	2	-
Female	10	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Delaware												
All races, both sexes	80	-	-	1	-	-	-	-	-	1	-	-
Male	45	-	-	1	-	-	-	-	-	1	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	72	-	-	1	-	-	-	-	-	1	-	-
Male	42	-	-	1	-	-	-	-	-	1	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	28	1	-	-	-	-	-	-	-	1	-	1
Male	14	1	-	-	-	-	-	-	-	1	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	19	1	-	-	-	-	-	-	-	1	-	1
Male	8	1	-	-	-	-	-	-	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Florida												
All races, both sexes	1,584	27	4	-	2	5	1	7	4	12	19	24
Male	788	16	3	-	1	4	1	3	2	7	6	12
Female	796	11	1	-	1	1	-	4	2	5	13	12
White, both sexes	1,466	17	1	-	1	5	1	6	4	7	12	17
Male	737	11	1	-	-	4	1	2	2	5	5	7
Female	729	6	-	-	1	1	-	4	2	2	7	10
Black, both sexes	103	9	3	-	1	-	-	1	-	4	7	7
Male	43	4	2	-	1	-	-	1	-	1	1	5
Female	60	5	1	-	-	-	-	-	-	3	6	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	1	-	-	-	-	-	-	-	1	-	-
Male	7	1	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	988	15	-	1	2	2	2	5	7	5	15	25
Male	443	8	-	1	1	2	1	1	4	4	6	11
Female	545	7	-	-	1	-	1	4	3	1	9	14
White, both sexes	800	3	-	-	1	2	2	3	5	5	10	15
Male	375	3	-	-	-	2	1	1	4	4	5	8
Female	425	-	-	-	1	-	1	2	1	1	5	7
Black, both sexes	178	12	-	1	1	-	-	2	2	-	5	10
Male	65	5	-	1	1	-	-	-	-	-	1	3
Female	113	7	-	-	-	-	-	2	2	-	4	7
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Hawaii												
All races, both sexes	116	2	-	-	-	-	-	-	-	2	2	1
Male	64	-	-	-	-	-	-	-	-	2	1	-
Female	52	2	-	-	-	-	-	-	-	-	1	1
White, both sexes	24	1	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	11	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	91	1	-	-	-	-	-	-	-	2	1	1
Male	51	-	-	-	-	-	-	-	-	2	1	-
Female	40	1	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	118	1	-	-	-	1	-	-	-	2	1	-
Male	67	1	-	-	-	-	-	-	-	1	-	-
Female	51	-	-	-	-	1	-	-	-	1	1	-
White, both sexes	116	1	-	-	-	1	-	-	-	1	1	-
Male	67	1	-	-	-	-	-	-	-	1	-	-
Female	49	-	-	-	-	1	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Illinois												
All races, both sexes	1,095	14	3	-	2	1	3	1	2	5	9	24
Male	540	9	3	-	1	-	1	-	1	2	4	15
Female	555	5	-	-	1	1	2	1	1	3	5	9
White, both sexes	921	5	2	-	-	1	2	1	-	3	6	17
Male	459	2	2	-	-	-	1	-	-	2	3	11
Female	462	3	-	-	-	1	1	1	-	1	3	6
Black, both sexes	151	9	1	-	2	-	1	-	2	2	2	7
Male	68	7	1	-	1	-	-	-	1	-	1	4
Female	83	2	-	-	1	-	1	-	1	2	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Indiana												
All races, both sexes	685	11	1	-	-	-	-	1	2	4	6	8
Male	333	7	-	-	-	-	-	-	1	2	4	5
Female	352	4	1	-	-	-	-	1	1	2	2	3
White, both sexes	652	9	1	-	-	-	-	1	2	2	4	6
Male	310	6	-	-	-	-	-	-	1	1	3	3
Female	342	3	1	-	-	-	-	1	1	1	1	3
Black, both sexes	33	2	-	-	-	-	-	-	-	2	2	2
Male	23	1	-	-	-	-	-	-	-	1	1	2
Female	10	1	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Iowa												
All races, both sexes	270	1	2	-	1	1	-	-	-	-	1	6
Male	139	-	2	-	-	1	-	-	-	-	-	2
Female	131	1	-	-	1	-	-	-	-	-	1	4
White, both sexes	264	1	2	-	1	1	-	-	-	-	1	6
Male	137	-	2	-	-	1	-	-	-	-	-	2
Female	127	1	-	-	1	-	-	-	-	-	1	4
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	274	2	-	2	-	-	-	2	1	1	3	2
Male	131	2	-	1	-	-	-	2	-	1	2	1
Female	143	-	-	1	-	-	-	-	1	-	1	1
White, both sexes	255	-	-	2	-	-	-	1	1	-	3	2
Male	120	-	-	1	-	-	-	1	-	-	2	1
Female	135	-	-	1	-	-	-	-	1	-	1	1
Black, both sexes	14	2	-	-	-	-	-	1	-	1	-	-
Male	9	2	-	-	-	-	-	1	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Kentucky												
All races, both sexes	430	4	-	1	-	-	-	2	2	2	6	5
Male	196	3	-	1	-	-	-	-	2	2	3	3
Female	234	1	-	-	-	-	-	2	-	-	3	2
White, both sexes	408	3	-	1	-	-	-	2	2	1	6	4
Male	186	2	-	1	-	-	-	-	2	1	3	3
Female	222	1	-	-	-	-	-	2	-	-	3	1
Black, both sexes	21	1	-	-	-	-	-	-	-	1	-	1
Male	10	1	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	465	7	2	-	-	-	4	2	3	1	11	10
Male	234	5	2	-	-	-	2	1	-	1	6	4
Female	231	2	-	-	-	-	2	1	3	-	5	6
White, both sexes	365	1	1	-	-	-	1	1	2	-	3	8
Male	179	1	1	-	-	-	-	1	-	-	2	4
Female	186	-	-	-	-	-	1	-	2	-	1	4
Black, both sexes	96	6	1	-	-	-	3	1	1	1	7	2
Male	54	4	1	-	-	-	2	-	-	1	4	-
Female	42	2	-	-	-	-	1	1	1	-	3	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Maine												
All races, both sexes	178	1	1	-	-	-	-	-	-	3	3	5
Male	92	1	1	-	-	-	-	-	-	1	2	3
Female	86	-	-	-	-	-	-	-	-	2	1	2
White, both sexes	177	1	-	-	-	-	-	-	-	3	3	5
Male	91	1	-	-	-	-	-	-	-	1	2	3
Female	86	-	-	-	-	-	-	-	-	2	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	393	3	-	-	-	-	-	1	2	2	2	4
Male	211	1	-	-	-	-	-	1	1	1	1	3
Female	182	2	-	-	-	-	-	-	1	1	1	1
White, both sexes	313	1	-	-	-	-	-	-	2	1	1	2
Male	164	-	-	-	-	-	-	-	1	1	-	2
Female	149	1	-	-	-	-	-	-	1	-	1	-
Black, both sexes	74	2	-	-	-	-	-	1	-	1	1	2
Male	45	1	-	-	-	-	-	1	-	-	1	1
Female	29	1	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Massachusetts												
All races, both sexes	733	4	1	2	1	-	1	1	2	5	3	5
Male	338	3	1	1	1	-	1	-	-	2	2	2
Female	395	1	-	1	-	-	-	1	2	3	1	3
White, both sexes	704	3	1	2	1	-	1	1	2	5	3	4
Male	326	2	1	1	1	-	1	-	-	2	2	2
Female	378	1	-	1	-	-	-	1	2	3	1	2
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	1	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	807	9	1	-	2	-	4	-	1	7	7	8
Male	386	4	1	-	1	-	3	-	-	2	2	2
Female	421	5	-	-	1	-	1	-	1	5	5	6
White, both sexes	707	6	1	-	2	-	2	-	1	5	5	5
Male	333	3	1	-	1	-	1	-	-	1	1	2
Female	374	3	-	-	1	-	1	-	1	4	4	3
Black, both sexes	83	3	-	-	-	-	2	-	-	2	1	3
Male	44	1	-	-	-	-	2	-	-	1	1	-
Female	39	2	-	-	-	-	-	-	-	1	-	3
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Minnesota												
All races, both sexes	400	3	1	-	-	-	-	1	2	2	6	4
Male	189	1	1	-	-	-	-	-	1	1	2	2
Female	211	2	-	-	-	-	-	1	1	1	4	2
White, both sexes	377	1	-	-	-	-	-	1	1	2	6	3
Male	176	-	-	-	-	-	-	-	-	1	2	2
Female	201	1	-	-	-	-	-	1	1	1	4	1
Black, both sexes	12	1	1	-	-	-	-	-	-	-	-	1
Male	6	1	1	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	1	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	343	13	2	-	-	1	1	1	-	1	8	5
Male	161	9	2	-	-	1	-	-	-	-	3	1
Female	182	4	-	-	-	-	1	1	-	1	5	4
White, both sexes	249	-	1	-	-	-	-	1	-	-	3	1
Male	113	-	1	-	-	-	-	-	-	-	-	1
Female	136	-	-	-	-	-	-	1	-	-	3	-
Black, both sexes	89	12	1	-	-	1	1	-	-	1	5	4
Male	46	8	1	-	-	1	-	-	-	-	3	-
Female	43	4	-	-	-	-	1	-	-	1	2	4
American Indian 1/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Missouri												
All races, both sexes	530	10	2	-	-	-	-	2	1	2	7	8
Male	278	7	-	-	-	-	-	2	-	-	6	2
Female	252	3	2	-	-	-	-	-	1	2	1	6
White, both sexes	470	3	2	-	-	-	-	2	-	1	4	6
Male	253	2	-	-	-	-	-	2	-	-	3	2
Female	217	1	2	-	-	-	-	-	-	1	1	4
Black, both sexes	52	7	-	-	-	-	-	-	1	1	2	2
Male	23	5	-	-	-	-	-	-	-	-	2	-
Female	29	2	-	-	-	-	-	-	1	1	-	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	115	1	1	-	-	-	-	-	1	1	1	-
Male	58	1	1	-	-	-	-	-	1	1	-	-
Female	57	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	107	-	1	-	-	-	-	-	1	1	1	-
Male	55	-	1	-	-	-	-	-	1	1	-	-
Female	52	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Nebraska												
All races, both sexes	165	-	-	1	-	-	-	-	-	1	2	3
Male	85	-	-	-	-	-	-	-	-	1	1	1
Female	80	-	-	1	-	-	-	-	-	-	1	2
White, both sexes	159	-	-	1	-	-	-	-	-	1	2	1
Male	83	-	-	-	-	-	-	-	-	1	1	-
Female	76	-	-	1	-	-	-	-	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	190	4	1	-	-	-	-	1	-	1	6	5
Male	105	4	-	-	-	-	-	-	-	-	4	3
Female	85	-	1	-	-	-	-	1	-	1	2	2
White, both sexes	173	3	1	-	-	-	-	1	-	-	4	5
Male	96	3	-	-	-	-	-	-	-	-	3	3
Female	77	-	1	-	-	-	-	1	-	-	1	2
Black, both sexes	10	1	-	-	-	-	-	-	-	1	1	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
New Hampshire												
All races, both sexes	123	1	-	-	-	-	-	-	-	-	-	2
Male	68	1	-	-	-	-	-	-	-	-	-	1
Female	55	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	123	1	-	-	-	-	-	-	-	-	-	2
Male	68	1	-	-	-	-	-	-	-	-	-	1
Female	55	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	708	6	2	2	1	-	1	3	-	4	7	11
Male	343	3	2	1	1	-	1	1	-	1	5	4
Female	365	3	-	1	-	-	-	2	-	3	2	7
White, both sexes	618	1	2	-	1	-	1	2	-	4	4	7
Male	303	1	2	-	1	-	1	1	-	1	3	3
Female	315	-	-	-	-	-	-	1	-	3	1	4
Black, both sexes	69	5	-	2	-	-	-	1	-	-	3	4
Male	27	2	-	1	-	-	-	-	-	-	2	1
Female	42	3	-	1	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
New Mexico												
All races, both sexes	177	1	-	-	-	-	-	1	1	1	1	1
Male	93	-	-	-	-	-	-	1	1	-	1	-
Female	84	1	-	-	-	-	-	-	-	1	-	1
White, both sexes	149	1	-	-	-	-	-	-	1	1	1	1
Male	77	-	-	-	-	-	-	-	1	-	1	-
Female	72	1	-	-	-	-	-	-	-	1	-	1
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,182	15	2	2	1	7	4	4	4	9	17	23
Male	578	9	2	2	-	4	4	3	3	8	6	12
Female	604	6	-	-	1	3	-	1	1	1	11	11
White, both sexes	1,044	5	2	1	1	6	4	4	2	5	10	17
Male	500	2	2	1	-	3	4	3	1	5	4	7
Female	544	3	-	-	1	3	-	1	1	-	6	10
Black, both sexes	110	9	-	1	-	-	-	-	2	3	7	5
Male	56	6	-	1	-	-	-	-	2	3	2	4
Female	54	3	-	-	-	-	-	-	-	-	5	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	1	-	-	-	1	-	-	-	1	-	1
Male	20	1	-	-	-	1	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
North Carolina												
All races, both sexes	843	6	4	1	-	2	-	1	5	3	5	25
Male	396	5	2	-	-	1	-	1	1	-	3	13
Female	447	1	2	1	-	1	-	-	4	3	2	12
White, both sexes	704	1	2	1	-	2	-	1	-	3	4	14
Male	334	1	-	-	-	1	-	1	-	-	3	8
Female	370	-	2	1	-	1	-	-	-	3	1	6
Black, both sexes	124	5	2	-	-	-	-	-	5	-	1	10
Male	57	4	2	-	-	-	-	-	1	-	-	5
Female	67	1	-	-	-	-	-	-	4	-	1	5
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	45	1	1	-	-	-	-	-	-	-	1	-
Male	26	1	1	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	41	1	1	-	-	-	-	-	-	-	1	-
Male	24	1	1	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Ohio												
All races, both sexes	1,084	8	1	1	-	2	2	-	4	5	9	29
Male	586	4	-	1	-	1	1	-	-	5	4	15
Female	498	4	1	-	-	1	1	-	4	-	5	14
White, both sexes	987	4	-	1	-	2	2	-	4	2	7	23
Male	526	3	-	1	-	1	1	-	-	2	3	11
Female	461	1	-	-	-	1	1	-	4	-	4	12
Black, both sexes	89	4	-	-	-	-	-	-	-	3	2	6
Male	55	1	-	-	-	-	-	-	-	3	1	4
Female	34	3	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	300	4	-	-	-	-	-	-	1	3	3	5
Male	150	2	-	-	-	-	-	-	1	1	-	4
Female	150	2	-	-	-	-	-	-	-	2	3	1
White, both sexes	265	3	-	-	-	-	-	-	1	3	3	3
Male	135	2	-	-	-	-	-	-	1	1	-	3
Female	130	1	-	-	-	-	-	-	-	2	3	-
Black, both sexes	11	1	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	5	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Oregon												
All races, both sexes	350	3	1	-	-	-	-	-	3	1	-	6
Male	165	2	1	-	-	-	-	-	1	-	-	1
Female	185	1	-	-	-	-	-	-	2	1	-	5
White, both sexes	336	1	1	-	-	-	-	-	3	1	-	5
Male	159	-	1	-	-	-	-	-	1	-	-	-
Female	177	1	-	-	-	-	-	-	2	1	-	5
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,491	8	3	2	2	4	1	4	8	10	11	19
Male	717	4	2	-	1	4	1	2	2	6	6	9
Female	774	4	1	2	1	-	-	2	6	4	5	10
White, both sexes	1,380	5	3	1	1	1	1	4	8	10	7	13
Male	668	1	2	-	1	1	1	2	2	6	3	6
Female	712	4	1	1	-	-	-	2	6	4	4	7
Black, both sexes	106	3	-	1	1	3	-	-	-	-	4	6
Male	47	3	-	-	-	3	-	-	-	-	3	3
Female	59	-	-	1	1	-	-	-	-	-	1	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Rhode Island												
All races, both sexes	102	1	-	-	-	-	-	1	-	1	-	2
Male	50	1	-	-	-	-	-	1	-	-	-	-
Female	52	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	99	1	-	-	-	-	-	1	-	1	-	2
Male	48	1	-	-	-	-	-	1	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	445	7	4	-	-	1	2	3	1	5	1	7
Male	210	4	2	-	-	1	2	1	-	4	-	3
Female	235	3	2	-	-	-	-	2	1	1	1	4
White, both sexes	369	3	2	-	-	1	-	3	1	2	1	2
Male	174	2	2	-	-	1	-	1	-	2	-	1
Female	195	1	-	-	-	-	-	2	1	-	1	1
Black, both sexes	73	4	2	-	-	-	2	-	-	3	-	4
Male	35	2	-	-	-	-	2	-	-	2	-	2
Female	38	2	2	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
South Dakota												
All races, both sexes	65	1	-	1	-	-	-	-	-	-	-	-
Male	31	1	-	1	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	60	1	-	-	-	-	-	-	-	-	-	-
Male	29	1	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	501	7	3	1	-	-	1	2	5	3	8	9
Male	272	3	1	1	-	-	1	2	2	1	4	1
Female	229	4	2	-	-	-	-	-	3	2	4	8
White, both sexes	452	7	1	1	-	-	-	1	4	3	7	5
Male	250	3	1	1	-	-	-	1	2	1	3	1
Female	202	4	-	-	-	-	-	-	2	2	4	4
Black, both sexes	47	-	2	-	-	-	1	1	1	-	1	4
Male	21	-	-	-	-	-	1	1	-	-	1	-
Female	26	-	2	-	-	-	-	-	1	-	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Texas												
All races, both sexes	1,532	20	8	2	3	4	2	7	7	9	19	31
Male	741	12	4	-	1	1	1	1	4	3	9	18
Female	791	8	4	2	2	3	1	6	3	6	10	13
White, both sexes	1,388	16	7	2	2	4	2	7	6	7	17	26
Male	679	11	4	-	-	1	1	1	4	2	9	14
Female	709	5	3	2	2	3	1	6	2	5	8	12
Black, both sexes	117	4	1	-	1	-	-	-	1	2	2	4
Male	49	1	-	-	1	-	-	-	-	1	-	3
Female	68	3	1	-	-	-	-	-	1	1	2	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	154	-	-	-	2	1	-	-	1	-	3	1
Male	70	-	-	-	1	-	-	-	-	-	-	-
Female	84	-	-	-	1	1	-	-	1	-	3	1
White, both sexes	150	-	-	-	2	1	-	-	1	-	3	1
Male	68	-	-	-	1	-	-	-	-	-	-	-
Female	82	-	-	-	1	1	-	-	1	-	3	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Vermont												
All races, both sexes	46	1	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	26	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	1	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	26	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	795	12	3	-	-	1	2	4	4	3	11	16
Male	398	6	3	-	-	1	2	4	3	-	7	10
Female	397	6	-	-	-	-	-	-	1	3	4	6
White, both sexes	660	7	3	-	-	1	-	3	3	1	7	10
Male	325	3	3	-	-	1	-	3	2	-	4	8
Female	335	4	-	-	-	-	-	-	1	1	3	2
Black, both sexes	119	4	-	-	-	-	2	-	1	2	3	6
Male	62	3	-	-	-	-	2	-	1	-	2	2
Female	57	1	-	-	-	-	-	-	-	2	1	4
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	1	-	-	1	-
Male	10	-	-	-	-	-	-	1	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Washington												
All races, both sexes	481	2	2	2	-	-	1	3	2	3	3	12
Male	253	1	1	1	-	-	1	3	-	1	2	8
Female	228	1	1	1	-	-	-	-	2	2	1	4
White, both sexes	445	-	2	2	-	-	1	3	2	2	1	8
Male	237	-	1	1	-	-	1	3	-	1	1	5
Female	208	-	1	1	-	-	-	-	2	1	-	3
Black, both sexes	12	-	-	-	-	-	-	-	-	-	1	3
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	7	2	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	1	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	1	-	-
West Virginia												
All races, both sexes	245	-	-	-	-	1	-	-	-	1	4	6
Male	128	-	-	-	-	-	-	-	-	-	2	4
Female	117	-	-	-	-	1	-	-	-	1	2	2
White, both sexes	243	-	-	-	-	1	-	-	-	1	4	6
Male	128	-	-	-	-	-	-	-	-	-	2	4
Female	115	-	-	-	-	1	-	-	-	1	2	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
Wisconsin												
All races, both sexes	421	7	-	-	-	-	-	-	1	5	2	5
Male	217	4	-	-	-	-	-	-	-	3	1	2
Female	204	3	-	-	-	-	-	-	1	2	1	3
White, both sexes	409	7	-	-	-	-	-	-	-	5	2	5
Male	214	4	-	-	-	-	-	-	-	3	1	2
Female	195	3	-	-	-	-	-	-	-	2	1	3
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	60	1	1	-	-	-	-	1	1	-	1	4
Male	33	-	-	-	-	-	-	1	-	-	1	4
Female	27	1	1	-	-	-	-	-	1	-	-	-
White, both sexes	59	1	1	-	-	-	-	1	1	-	1	4
Male	32	-	-	-	-	-	-	1	-	-	1	4
Female	27	1	1	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
United States												
All races, both sexes	696	1,100	1,460	2,058	2,913	4,077	4,831	3,756	2,012	616	113	-
Male	354	618	815	1,170	1,658	2,219	2,391	1,581	707	153	19	-
Female	342	482	645	888	1,255	1,858	2,440	2,175	1,305	463	94	-
White, both sexes	530	875	1,227	1,794	2,620	3,725	4,481	3,486	1,876	581	93	-
Male	281	520	691	1,036	1,492	2,049	2,230	1,468	658	142	12	-
Female	249	355	536	758	1,128	1,676	2,251	2,018	1,218	439	81	-
Black, both sexes	141	191	185	183	227	246	228	193	91	26	17	-
Male	59	81	99	93	131	119	103	74	31	5	4	-
Female	82	110	86	90	96	127	125	119	60	21	13	-
American Indian 1/, both sexes	14	17	17	29	22	26	31	20	10	1	1	-
Male	9	6	7	15	10	14	14	10	3	-	1	-
Female	5	11	10	14	12	12	17	10	7	1	-	-
Asian or Pacific Islander 2/, both sexes	11	17	31	52	44	80	91	57	35	8	2	-
Male	5	11	18	26	25	37	44	29	15	6	2	-
Female	6	6	13	26	19	43	47	28	20	2	-	-
Alabama												
All races, both sexes	18	34	40	42	79	91	87	69	43	9	2	-
Male	9	21	28	28	41	44	28	31	14	3	-	-
Female	9	13	12	14	38	47	59	38	29	6	2	-
White, both sexes	10	24	29	33	67	75	80	56	32	8	-	-
Male	7	16	20	23	34	36	27	24	11	3	-	-
Female	3	8	9	10	33	39	53	32	21	5	-	-
Black, both sexes	8	9	11	9	12	15	7	13	11	1	2	-
Male	2	5	8	5	7	8	1	7	3	-	-	-
Female	6	4	3	4	5	7	6	6	8	1	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Alaska												
All races, both sexes	2	2	2	4	5	10	5	4	3	-	-	-
Male	2	1	2	3	3	5	2	1	-	-	-	-
Female	-	1	-	1	2	5	3	3	3	-	-	-
White, both sexes	-	1	2	2	3	9	4	-	2	-	-	-
Male	-	1	2	1	2	5	2	-	-	-	-	-
Female	-	-	-	1	1	4	2	-	2	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	2	2	1	1	3	1	-	-	-
Male	1	-	-	2	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	1	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Arizona												
All races, both sexes	15	14	22	54	64	88	85	62	30	8	2	-
Male	9	5	12	29	39	54	50	34	12	2	1	-
Female	6	9	10	25	25	34	35	28	18	6	1	-
White, both sexes	13	13	21	48	60	83	80	59	27	8	1	-
Male	7	5	12	27	38	49	45	34	11	2	-	-
Female	6	8	9	21	22	34	35	25	16	6	1	-
Black, both sexes	1	1	1	-	-	-	1	1	1	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	1	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	1	-	-	5	4	4	4	1	1	-	1	-
Male	1	-	-	2	1	4	4	-	-	-	1	-
Female	-	-	-	3	3	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Arkansas												
All races, both sexes	15	17	14	20	32	29	48	34	14	9	3	-
Male	7	12	8	8	25	16	23	15	4	3	-	-
Female	8	5	6	12	7	13	25	19	10	6	3	-
White, both sexes	13	16	10	16	25	27	47	32	14	7	3	-
Male	7	11	7	6	20	14	23	14	4	2	-	-
Female	6	5	3	10	5	13	24	18	10	5	3	-
Black, both sexes	2	1	3	3	7	1	1	2	-	1	-	-
Male	-	1	1	2	5	1	-	1	-	-	-	-
Female	2	-	2	1	2	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	60	90	101	175	216	292	435	296	151	48	5	-
Male	28	53	55	95	121	159	218	142	60	13	2	-
Female	32	37	46	80	95	133	217	154	91	35	3	-
White, both sexes	49	76	90	149	190	254	385	276	130	47	3	-
Male	26	45	50	82	107	143	198	133	53	12	1	-
Female	23	31	40	67	83	111	187	143	77	35	2	-
Black, both sexes	7	8	2	8	8	12	9	5	6	1	1	-
Male	1	3	1	5	4	4	2	1	2	1	-	-
Female	6	5	1	3	4	8	7	4	4	-	1	-
American Indian 1/, both sexes	2	1	2	2	4	-	-	-	2	-	-	-
Male	1	1	1	1	3	-	-	-	1	-	-	-
Female	1	-	1	1	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	7	16	14	26	41	15	13	-	1	-
Male	-	4	3	7	7	12	18	8	4	-	1	-
Female	2	1	4	9	7	14	23	7	9	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Colorado												
All races, both sexes	8	13	15	21	44	60	72	59	29	9	2	-
Male	7	7	8	17	27	32	40	29	17	3	-	-
Female	1	6	7	4	17	28	32	30	12	6	2	-
White, both sexes	8	13	15	20	39	59	69	56	28	9	2	-
Male	7	7	8	16	24	31	39	28	16	3	-	-
Female	1	6	7	4	15	28	30	28	12	6	2	-
Black, both sexes	-	-	-	1	5	-	2	1	1	-	-	-
Male	-	-	-	1	3	-	1	-	1	-	-	-
Female	-	-	-	-	2	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	10	16	12	26	25	53	73	61	41	11	2	-
Male	5	6	6	14	14	32	28	21	13	1	1	-
Female	5	10	6	12	11	21	45	40	28	10	1	-
White, both sexes	8	15	10	21	23	51	71	60	41	10	2	-
Male	5	5	4	13	12	30	27	20	13	1	1	-
Female	3	10	6	8	11	21	44	40	28	9	1	-
Black, both sexes	2	-	2	5	2	2	2	1	-	1	-	-
Male	-	-	2	1	2	2	1	1	-	-	-	-
Female	2	-	-	4	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Delaware												
All races, both sexes	2	4	3	8	5	15	20	13	7	1	-	-
Male	1	1	3	5	4	10	8	9	2	-	-	-
Female	1	3	-	3	1	5	12	4	5	1	-	-
White, both sexes	2	2	2	6	3	15	19	13	7	1	-	-
Male	1	1	2	4	3	10	8	9	2	-	-	-
Female	1	1	-	2	-	5	11	4	5	1	-	-
Black, both sexes	-	2	1	2	2	-	1	-	-	-	-	-
Male	-	-	1	1	1	-	-	-	-	-	-	-
Female	-	2	-	1	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	2	4	1	3	6	3	3	2	1	-	-
Male	-	2	1	1	1	3	1	-	2	-	-	-
Female	-	-	3	-	2	3	2	3	-	1	-	-
White, both sexes	-	-	-	-	1	5	2	-	1	-	-	-
Male	-	-	-	-	1	3	1	-	1	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
Black, both sexes	-	2	4	1	2	1	1	3	1	1	-	-
Male	-	2	1	1	-	-	-	-	1	-	-	-
Female	-	-	3	-	2	1	1	3	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Florida												
All races, both sexes	31	63	78	126	178	262	315	247	128	37	14	-
Male	15	35	40	64	102	145	160	112	49	10	1	-
Female	16	28	38	62	76	117	155	135	79	27	13	-
White, both sexes	23	48	61	119	173	247	311	239	124	36	14	-
Male	12	30	32	61	100	137	160	109	48	9	1	-
Female	11	18	29	58	73	110	151	130	76	27	13	-
Black, both sexes	8	14	16	6	3	11	2	8	2	1	-	-
Male	3	5	7	3	-	6	-	3	-	1	-	-
Female	5	9	9	3	3	5	2	5	2	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	2	4	2	-	2	-	-	-
Male	-	-	-	-	2	2	-	-	1	-	-	-
Female	-	1	-	1	-	2	2	-	1	-	-	-
Georgia												
All races, both sexes	40	43	73	66	118	182	147	148	70	17	5	-
Male	13	21	39	35	67	92	66	41	22	7	1	-
Female	27	22	34	31	51	90	81	107	48	10	4	-
White, both sexes	24	29	61	54	94	158	127	129	61	14	3	-
Male	9	15	34	29	54	83	58	38	21	6	-	-
Female	15	14	27	25	40	75	69	91	40	8	3	-
Black, both sexes	15	14	12	10	22	23	18	18	8	3	2	-
Male	3	6	5	6	12	9	8	2	1	1	1	-
Female	12	8	7	4	10	14	10	16	7	2	1	-
American Indian 1/, both sexes	1	-	-	-	1	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	1	1	1	1	1	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	2	-	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Hawaii												
All races, both sexes	1	2	10	7	8	17	26	22	12	4	-	-
Male	-	1	7	5	2	11	17	7	7	4	-	-
Female	1	1	3	2	6	6	9	15	5	-	-	-
White, both sexes	-	-	3	1	2	3	5	4	3	1	-	-
Male	-	-	3	1	-	2	3	2	1	1	-	-
Female	-	-	-	-	2	1	2	2	2	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	7	6	6	14	21	18	9	3	-	-
Male	-	1	4	4	2	9	14	5	6	3	-	-
Female	1	-	3	2	4	5	7	13	3	-	-	-
Idaho												
All races, both sexes	2	3	5	10	15	13	29	24	12	-	-	-
Male	1	1	3	5	9	12	15	14	5	-	-	-
Female	1	2	2	5	6	1	14	10	7	-	-	-
White, both sexes	2	2	5	10	15	13	29	24	12	-	-	-
Male	1	1	3	5	9	12	15	14	5	-	-	-
Female	1	1	2	5	6	1	14	10	7	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Illinois												
All races, both sexes	27	51	61	84	132	200	220	145	78	28	5	-
Male	10	23	37	45	82	110	106	62	23	4	2	-
Female	17	28	24	39	50	90	114	83	55	24	3	-
White, both sexes	19	37	50	72	108	178	189	126	73	27	5	-
Male	7	19	32	40	68	99	96	50	21	4	2	-
Female	12	18	18	32	40	79	93	76	52	23	3	-
Black, both sexes	5	14	11	9	22	18	26	14	5	1	-	-
Male	1	4	5	4	13	9	8	7	2	-	-	-
Female	4	10	6	5	9	9	18	7	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	3	2	4	5	4	-	-	-	-
Male	2	-	-	1	1	2	2	4	-	-	-	-
Female	1	-	-	2	1	2	3	-	-	-	-	-
Indiana												
All races, both sexes	18	32	37	62	81	112	134	96	51	24	5	-
Male	9	18	21	41	44	62	65	39	10	5	-	-
Female	9	14	16	21	37	50	69	57	41	19	5	-
White, both sexes	15	31	35	60	77	107	126	96	51	24	5	-
Male	7	18	19	40	41	57	60	39	10	5	-	-
Female	8	13	16	20	36	50	66	57	41	19	5	-
Black, both sexes	3	1	2	2	4	5	8	-	-	-	-	-
Male	2	-	2	1	3	5	5	-	-	-	-	-
Female	1	1	-	1	1	-	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Iowa												
All races, both sexes	1	4	11	16	32	46	60	47	24	15	2	-
Male	1	2	5	12	19	25	40	18	8	4	-	-
Female	-	2	6	4	13	21	20	29	16	11	2	-
White, both sexes	1	4	11	16	29	46	58	46	24	15	2	-
Male	1	2	5	12	17	25	40	18	8	4	-	-
Female	-	2	6	4	12	21	18	28	16	11	2	-
Black, both sexes	-	-	-	-	1	-	2	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	5	11	13	25	25	45	53	55	22	6	1	-
Male	2	6	7	16	8	19	33	22	8	1	-	-
Female	3	5	6	9	17	26	20	33	14	5	1	-
White, both sexes	4	9	11	22	23	45	51	53	21	6	1	-
Male	1	6	5	15	7	19	31	22	8	1	-	-
Female	3	3	6	7	16	26	20	31	13	5	1	-
Black, both sexes	1	2	1	2	1	-	1	2	-	-	-	-
Male	1	-	1	1	1	-	1	-	-	-	-	-
Female	-	2	-	1	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Kentucky												
All races, both sexes	21	16	25	42	55	63	72	69	33	10	2	-
Male	8	11	10	29	31	29	31	21	10	2	-	-
Female	13	5	15	13	24	34	41	48	23	8	2	-
White, both sexes	18	14	24	40	54	61	68	68	32	8	2	-
Male	8	9	10	27	31	28	29	21	10	1	-	-
Female	10	5	14	13	23	33	39	47	22	7	2	-
Black, both sexes	3	2	1	2	1	2	4	1	-	2	-	-
Male	-	2	-	2	-	1	2	-	-	1	-	-
Female	3	-	1	-	1	1	2	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	21	27	42	32	64	65	76	67	22	6	3	-
Male	13	11	26	16	36	31	45	26	8	1	-	-
Female	8	16	16	16	28	34	31	41	14	5	3	-
White, both sexes	14	20	27	22	52	58	65	61	20	6	3	-
Male	7	8	13	11	31	31	39	23	6	1	-	-
Female	7	12	14	11	21	27	26	38	14	5	3	-
Black, both sexes	7	7	14	10	11	6	11	6	2	-	-	-
Male	6	3	12	5	5	-	6	3	2	-	-	-
Female	1	4	2	5	6	6	5	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Maine												
All races, both sexes	3	5	11	12	17	22	37	29	22	5	2	-
Male	2	3	5	9	11	13	20	12	6	3	-	-
Female	1	2	6	3	6	9	17	17	16	2	2	-
White, both sexes	3	5	11	12	17	22	37	29	22	5	2	-
Male	2	3	5	9	11	13	20	12	6	3	-	-
Female	1	2	6	3	6	9	17	17	16	2	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	14	25	21	37	49	59	80	64	28	2	-	-
Male	8	15	12	22	30	31	43	30	11	1	-	-
Female	6	10	9	15	19	28	37	34	17	1	-	-
White, both sexes	5	20	18	26	35	52	71	51	26	2	-	-
Male	3	11	9	17	21	29	36	23	10	1	-	-
Female	2	9	9	9	14	23	35	28	16	1	-	-
Black, both sexes	9	5	3	10	12	6	9	11	2	-	-	-
Male	5	4	3	5	8	2	7	6	1	-	-	-
Female	4	1	-	5	4	4	2	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	1	1	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Massachusetts												
All races, both sexes	10	32	35	47	84	112	140	130	83	32	3	-
Male	6	23	17	30	51	49	55	48	34	12	-	-
Female	4	9	18	17	33	63	85	82	49	20	3	-
White, both sexes	9	30	31	46	81	109	132	127	82	31	3	-
Male	6	23	15	29	49	48	54	46	33	11	-	-
Female	3	7	16	17	32	61	78	81	49	20	3	-
Black, both sexes	1	2	3	-	1	-	6	3	-	1	-	-
Male	-	-	2	-	-	-	1	2	-	1	-	-
Female	1	2	1	-	1	-	5	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	2	2	-	1	-	-	-
Male	-	-	-	1	2	-	-	-	1	-	-	-
Female	-	-	1	-	-	2	2	-	-	-	-	-
Michigan												
All races, both sexes	19	30	52	68	99	144	158	110	67	18	3	-
Male	9	17	31	33	57	79	75	39	26	3	2	-
Female	10	13	21	35	42	65	83	71	41	15	1	-
White, both sexes	11	21	41	62	91	129	151	93	63	16	2	-
Male	4	14	24	31	52	70	71	31	22	3	1	-
Female	7	7	17	31	39	59	80	62	41	13	1	-
Black, both sexes	4	7	8	5	5	15	6	16	4	2	-	-
Male	2	3	5	1	4	9	4	7	4	-	-	-
Female	2	4	3	4	1	6	2	9	-	2	-	-
American Indian 1/, both sexes	3	2	1	1	2	-	-	1	-	-	-	-
Male	2	-	-	1	1	-	-	1	-	-	-	-
Female	1	2	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	-	1	-	1	-	-	-	1	-
Male	1	-	2	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Minnesota												
All races, both sexes	11	15	21	25	36	66	90	60	36	20	1	-
Male	6	11	7	14	25	33	40	26	13	6	-	-
Female	5	4	14	11	11	33	50	34	23	14	1	-
White, both sexes	9	13	18	22	34	63	89	60	34	20	1	-
Male	6	9	5	13	23	31	39	26	13	6	-	-
Female	3	4	13	9	11	32	50	34	21	14	1	-
Black, both sexes	2	2	-	2	-	1	1	-	1	-	-	-
Male	-	2	-	1	-	-	1	-	-	-	-	-
Female	2	-	-	1	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	2	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	2	-	-	1	-	-	-
Male	-	-	1	-	2	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Mississippi												
All races, both sexes	7	25	24	30	46	54	45	47	27	4	2	-
Male	3	10	16	17	26	30	14	19	10	-	-	-
Female	4	15	8	13	20	24	31	28	17	4	2	-
White, both sexes	2	17	15	21	37	45	38	43	21	3	1	-
Male	1	7	11	12	20	25	14	15	6	-	-	-
Female	1	10	4	9	17	20	24	28	15	3	1	-
Black, both sexes	5	8	9	8	9	8	5	4	6	1	1	-
Male	2	3	5	4	6	5	-	4	4	-	-	-
Female	3	5	4	4	3	3	5	-	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Missouri												
All races, both sexes	18	26	34	61	68	88	102	56	33	9	3	-
Male	10	20	19	39	41	46	49	22	14	-	1	-
Female	8	6	15	22	27	42	53	34	19	9	2	-
White, both sexes	14	23	28	54	65	78	97	50	31	9	3	-
Male	9	17	17	35	40	44	47	20	14	-	1	-
Female	5	6	11	19	25	34	50	30	17	9	2	-
Black, both sexes	3	3	5	6	3	8	4	5	2	-	-	-
Male	1	3	2	3	1	2	2	2	-	-	-	-
Female	2	-	3	3	2	6	2	3	2	-	-	-
American Indian 1/, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	1	-	-	-	-
Montana												
All races, both sexes	1	5	2	14	14	16	29	14	11	3	1	-
Male	1	1	2	8	8	9	14	7	4	-	-	-
Female	-	4	-	6	6	7	15	7	7	3	1	-
White, both sexes	-	5	2	13	12	15	27	14	11	3	1	-
Male	-	1	2	8	7	9	14	7	4	-	-	-
Female	-	4	-	5	5	6	13	7	7	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	2	1	2	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Nebraska												
All races, both sexes	6	3	6	6	23	37	33	22	16	6	-	-
Male	2	2	4	3	13	18	21	11	6	2	-	-
Female	4	1	2	3	10	19	12	11	10	4	-	-
White, both sexes	5	3	5	6	22	36	33	22	16	6	-	-
Male	2	2	3	3	13	18	21	11	6	2	-	-
Female	3	1	2	3	9	18	12	11	10	4	-	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	11	14	27	20	25	36	17	11	5	1	-
Male	3	9	7	16	10	18	19	8	4	-	-	-
Female	2	2	7	11	10	7	17	9	7	5	1	-
White, both sexes	5	9	12	25	20	22	33	17	11	4	1	-
Male	3	8	7	14	10	16	17	8	4	-	-	-
Female	2	1	5	11	10	6	16	9	7	4	1	-
Black, both sexes	-	2	2	-	-	1	1	-	-	1	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	1	2	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	2	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
New Hampshire												
All races, both sexes	-	3	12	9	16	16	31	16	10	7	-	-
Male	-	1	5	6	11	11	19	9	3	1	-	-
Female	-	2	7	3	5	5	12	7	7	6	-	-
White, both sexes	-	3	12	9	16	16	31	16	10	7	-	-
Male	-	1	5	6	11	11	19	9	3	1	-	-
Female	-	2	7	3	5	5	12	7	7	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	12	28	28	51	85	121	144	121	64	16	1	-
Male	7	16	18	29	43	63	69	50	24	5	-	-
Female	5	12	10	22	42	58	75	71	40	11	1	-
White, both sexes	10	21	23	42	73	108	133	112	59	14	1	-
Male	5	15	17	24	35	56	65	46	23	4	-	-
Female	5	6	6	18	38	52	68	66	36	10	1	-
Black, both sexes	2	7	4	6	7	11	8	5	3	1	-	-
Male	2	1	1	2	4	5	3	2	1	-	-	-
Female	-	6	3	4	3	6	5	3	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	3	5	2	3	4	2	1	-	-
Male	-	-	-	3	4	2	1	2	-	1	-	-
Female	-	-	1	-	1	-	2	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
New Mexico												
All races, both sexes	5	11	9	15	18	23	41	25	18	6	-	-
Male	4	7	8	8	8	14	19	12	9	1	-	-
Female	1	4	1	7	10	9	22	13	9	5	-	-
White, both sexes	5	9	7	14	17	22	31	19	15	5	-	-
Male	4	6	6	7	7	14	13	9	8	1	-	-
Female	1	3	1	7	10	8	18	10	7	4	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	2	1	-	1	10	5	3	1	-	-
Male	-	1	2	1	-	-	6	2	1	-	-	-
Female	-	1	-	-	-	1	4	3	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	28	40	88	105	146	182	188	186	94	35	2	-
Male	19	19	49	58	88	88	86	76	28	13	1	-
Female	9	21	39	47	58	94	102	110	66	22	1	-
White, both sexes	24	29	68	85	134	173	173	175	92	32	2	-
Male	17	15	37	47	79	83	79	71	26	13	1	-
Female	7	14	31	38	55	90	94	104	66	19	1	-
Black, both sexes	3	10	16	15	10	6	12	8	1	2	-	-
Male	2	3	9	7	7	3	4	2	1	-	-	-
Female	1	7	7	8	3	3	8	6	-	2	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	4	4	1	3	3	3	1	1	-	-
Male	-	1	3	3	1	2	3	3	1	-	-	-
Female	1	-	1	1	-	1	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
North Carolina												
All races, both sexes	25	43	61	76	98	130	151	127	57	17	6	-
Male	16	24	26	40	46	67	76	55	15	4	1	-
Female	9	19	35	36	52	63	75	72	42	13	5	-
White, both sexes	14	28	49	65	83	117	135	113	53	15	4	-
Male	10	17	23	34	38	62	70	48	14	3	1	-
Female	4	11	26	31	45	55	65	65	39	12	3	-
Black, both sexes	10	15	11	8	14	12	13	11	4	1	2	-
Male	5	7	3	4	8	5	6	6	1	-	-	-
Female	5	8	8	4	6	7	7	5	3	1	2	-
American Indian 1/, both sexes	-	-	-	3	1	1	2	3	-	-	-	-
Male	-	-	-	2	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	1	2	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	1	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
North Dakota												
All races, both sexes	1	1	2	5	5	6	10	5	5	1	1	-
Male	-	1	1	3	3	5	6	3	2	-	-	-
Female	1	-	1	2	2	1	4	2	3	1	1	-
White, both sexes	1	1	2	4	5	5	8	5	5	1	1	-
Male	-	1	1	3	3	4	5	3	2	-	-	-
Female	1	-	1	1	2	1	3	2	3	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Ohio												
All races, both sexes	32	48	58	83	119	190	217	159	89	27	1	-
Male	20	36	38	48	74	117	108	72	34	7	1	-
Female	12	12	20	35	45	73	109	87	55	20	-	-
White, both sexes	27	40	51	72	114	172	206	151	82	27	-	-
Male	16	31	32	40	72	108	99	68	31	7	-	-
Female	11	9	19	32	42	64	107	83	51	20	-	-
Black, both sexes	5	8	5	8	5	17	10	8	7	-	1	-
Male	4	5	5	5	2	9	8	4	3	-	1	-
Female	1	3	-	3	3	8	2	4	4	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	3	-	-	1	-	-	-	-	-
Male	-	-	1	3	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	6	21	16	36	24	56	52	43	22	4	4	-
Male	4	10	8	23	8	35	23	19	9	1	2	-
Female	2	11	8	13	16	21	29	24	13	3	2	-
White, both sexes	4	15	14	30	24	48	48	40	21	4	4	-
Male	3	7	8	20	8	32	22	16	9	1	2	-
Female	1	8	6	10	16	16	26	24	12	3	2	-
Black, both sexes	-	1	-	3	-	2	1	1	1	-	-	-
Male	-	1	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	2	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	2	5	2	3	-	6	3	2	-	-	-	-
Male	1	2	-	2	-	2	-	2	-	-	-	-
Female	1	3	2	1	-	4	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Oregon												
All races, both sexes	10	16	15	21	35	63	68	49	53	5	1	-
Male	6	10	6	12	23	31	35	19	17	1	-	-
Female	4	6	9	9	12	32	33	30	36	4	1	-
White, both sexes	10	16	14	20	34	60	66	48	51	5	1	-
Male	6	10	6	12	22	30	34	19	17	1	-	-
Female	4	6	8	8	12	30	32	29	34	4	1	-
Black, both sexes	-	-	1	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	2	-	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Pennsylvania												
All races, both sexes	30	50	77	107	163	241	297	264	137	43	10	-
Male	15	24	45	58	98	137	144	106	44	8	1	-
Female	15	26	32	49	65	104	153	158	93	35	9	-
White, both sexes	22	39	71	103	150	226	284	251	130	43	7	-
Male	12	20	42	57	90	130	137	105	41	8	1	-
Female	10	19	29	46	60	96	147	146	89	35	6	-
Black, both sexes	7	10	6	3	13	14	12	13	7	-	3	-
Male	2	4	3	1	8	7	6	1	3	-	-	-
Female	5	6	3	2	5	7	6	12	4	-	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Rhode Island												
All races, both sexes	5	3	5	9	4	17	23	17	11	3	-	-
Male	2	3	2	5	2	10	12	9	3	-	-	-
Female	3	-	3	4	2	7	11	8	8	3	-	-
White, both sexes	5	2	3	9	4	17	23	17	11	3	-	-
Male	2	2	1	5	2	10	12	9	3	-	-	-
Female	3	-	2	4	2	7	11	8	8	3	-	-
Black, both sexes	-	1	2	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	18	15	32	41	58	74	72	61	29	11	3	-
Male	12	7	17	26	29	41	33	18	9	1	-	-
Female	6	8	15	15	29	33	39	43	20	10	3	-
White, both sexes	15	8	25	34	49	66	65	55	25	10	2	-
Male	10	4	13	21	25	37	29	16	9	1	-	-
Female	5	4	12	13	24	29	36	39	16	9	2	-
Black, both sexes	3	7	7	7	9	7	7	5	4	1	1	-
Male	2	3	4	5	4	4	4	1	-	-	-	-
Female	1	4	3	2	5	3	3	4	4	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
South Dakota												
All races, both sexes	1	6	3	1	7	8	15	13	8	1	-	-
Male	-	2	1	-	5	5	9	5	1	1	-	-
Female	1	4	2	1	2	3	6	8	7	-	-	-
White, both sexes	1	4	2	1	7	7	15	13	8	1	-	-
Male	-	2	1	-	5	4	9	5	1	1	-	-
Female	1	2	1	1	2	3	6	8	7	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	2	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	17	32	37	46	48	77	94	69	31	9	2	-
Male	9	19	24	30	30	48	53	31	9	2	1	-
Female	8	13	13	16	18	29	41	38	22	7	1	-
White, both sexes	11	26	34	44	44	73	89	62	31	8	1	-
Male	5	18	23	30	28	45	48	29	9	2	-	-
Female	6	8	11	14	16	28	41	33	22	6	1	-
Black, both sexes	6	6	2	2	4	4	5	6	-	1	1	-
Male	4	1	-	-	2	3	5	2	-	-	1	-
Female	2	5	2	2	2	1	-	4	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Texas												
All races, both sexes	50	76	107	125	189	245	279	207	103	38	1	-
Male	24	42	59	66	98	129	137	82	39	11	-	-
Female	26	34	48	59	91	116	142	125	64	27	1	-
White, both sexes	39	61	97	112	171	220	265	195	94	37	1	-
Male	20	36	55	58	88	118	131	78	37	11	-	-
Female	19	25	42	54	83	102	134	117	57	26	1	-
Black, both sexes	9	9	9	11	14	19	13	11	6	1	-	-
Male	3	3	3	8	9	8	5	4	-	-	-	-
Female	6	6	6	3	5	11	8	7	6	1	-	-
American Indian 1/, both sexes	-	2	-	1	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	1	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	1	1	4	6	-	1	2	-	-	-
Male	1	2	1	-	1	3	-	-	1	-	-	-
Female	1	2	-	1	3	3	-	1	1	-	-	-
Utah												
All races, both sexes	4	7	13	7	17	25	41	21	9	2	-	-
Male	3	3	8	2	10	14	20	6	3	-	-	-
Female	1	4	5	5	7	11	21	15	6	2	-	-
White, both sexes	4	7	11	7	17	24	40	21	9	2	-	-
Male	3	3	7	2	10	13	20	6	3	-	-	-
Female	1	4	4	5	7	11	20	15	6	2	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Vermont												
All races, both sexes	1	1	3	4	4	9	10	7	3	2	1	-
Male	1	1	-	3	2	7	2	3	1	-	-	-
Female	-	-	3	1	2	2	8	4	2	2	1	-
White, both sexes	1	1	3	4	4	9	10	7	3	2	1	-
Male	1	1	-	3	2	7	2	3	1	-	-	-
Female	-	-	3	1	2	2	8	4	2	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	24	37	43	66	98	132	152	100	61	22	4	-
Male	9	19	32	32	52	67	79	51	17	3	1	-
Female	15	18	11	34	46	65	73	49	44	19	3	-
White, both sexes	18	27	33	48	84	117	133	88	55	20	2	-
Male	6	12	25	25	43	58	70	43	16	3	-	-
Female	12	15	8	23	41	59	63	45	39	17	2	-
Black, both sexes	6	7	10	15	13	14	17	9	6	2	2	-
Male	3	5	7	6	8	9	7	5	1	-	1	-
Female	3	2	3	9	5	5	10	4	5	2	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	3	1	1	2	3	-	-	-	-
Male	-	1	-	1	1	-	2	3	-	-	-	-
Female	-	1	-	2	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Washington												
All races, both sexes	17	21	24	42	49	75	99	80	38	4	2	-
Male	5	12	12	24	30	41	60	37	13	1	-	-
Female	12	9	12	18	19	34	39	43	25	3	2	-
White, both sexes	16	18	21	37	47	67	95	80	38	3	2	-
Male	4	11	12	22	28	39	57	37	13	1	-	-
Female	12	7	9	15	19	28	38	43	25	2	2	-
Black, both sexes	1	2	-	-	2	2	1	-	-	-	-	-
Male	1	-	-	-	2	-	1	-	-	-	-	-
Female	-	2	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	2	2	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	2	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	3	-	5	3	-	-	1	-	-
Male	-	1	-	1	-	1	2	-	-	-	-	-
Female	-	-	1	2	-	4	1	-	-	1	-	-
West Virginia												
All races, both sexes	7	11	15	31	33	38	32	40	19	5	2	-
Male	3	8	7	20	18	21	19	16	9	1	-	-
Female	4	3	8	11	15	17	13	24	10	4	2	-
White, both sexes	7	11	15	31	33	38	32	39	19	5	1	-
Male	3	8	7	20	18	21	19	16	9	1	-	-
Female	4	3	8	11	15	17	13	23	10	4	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2006

[Race categories are consistent with the 1977 Office of Management and Budget(OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data--mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Wisconsin												
All races, both sexes	9	7	23	25	52	74	89	68	40	11	3	-
Male	3	5	11	15	30	48	47	32	14	2	-	-
Female	6	2	12	10	22	26	42	36	26	9	3	-
White, both sexes	7	7	23	21	50	72	89	67	40	11	3	-
Male	3	5	11	14	30	47	47	31	14	2	-	-
Female	4	2	12	7	20	25	42	36	26	9	3	-
Black, both sexes	2	-	-	2	1	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	2	-	-	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	2	1	5	8	3	16	8	5	-	-	-
Male	2	1	-	3	3	3	9	4	2	-	-	-
Female	1	1	1	2	5	-	7	4	3	-	-	-
White, both sexes	3	2	1	4	8	3	16	8	5	-	-	-
Male	2	1	-	2	3	3	9	4	2	-	-	-
Female	1	1	1	2	5	-	7	4	3	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-