Person with disabilities experiencing problems accessing NCHS birth and death tables should contact nchsed@cdc.gov, or call 301-458-4688

Person with disabilities experiencing problems accessing NCHS birth and death tables should contact nchsed@cdc.gov, or call 301-458-4688

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (bas															
on the Internationa Classification of D												70	sian or		
Tenth Revision, Sec		All rad	700		White			Black		λme	rican In			ific Isl	ander 2/
Edition, 2004)	Both	AII IA		Both	WIIICC		Both	DIACK		Both	i i can ii.	ididii 1/	Both		ander 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes		Female
All causes															
All ages		1,201,942		2,077,549			289,971	148,602		14,037	7,630	6,407	44,707	23,382	21,325
Under 1 year	28,527	15,980	12,547	18,403	10,345	8,058	8,858	4,886	3,972	393	242	151	873	507	366
1-4 years	4,631	2,541	2,090	3,233	1,781	1,452	1,144	633	511	94	51	43	160	76	84
5-9 years	2,735	1,551	1,184	1,961	1,111	850	631	369	262	39	20	19	104	51	53
10-14 years	3,414	2,074	1,340	2,499	1,528	971	763	462	301	52	27	25	100	57	43
15-19 years	13,739	9,918	3,821	10,142	7,168	2,974	2,952	2,309	643	291	195	96	354	246	108
20-24 years	21,148	16,152	4,996	15,729	12,007	3,722	4,531	3,484	1,047	373	283	90	515	378	137
25-29 years	20,897	15,181	5,716	15,214	11,114	4,100	4,875	3,506	1,369	327	242	85	481	319	162
30-34 years	22,055	15,007	7,048	15,913	10,969	4,944	5,170	3,413	1,757	381	250	131	591	375	216
35-39 years	31,499	20,131	11,368	23,227	15,107	8,120	7,040	4,268	2,772	497	325	172	735	431	304
40-44 years	51,544	32,019	19,525	38,833	24,663	14,170	10,890	6,253	4,637	755	459	296	1,066	644	422
45-49 years	79,268	49,018	30,250	60,500	38,112	22,388	16,254	9,391	6,863	904	563	341	1,610	952	658
50-54 years	105,763	66,154	39,609	81,469	51,569	29,900	21,165	12,721	8,444	1,033	636	397	2,096	1,228	868
55-59 years	133,053	81,373	51,680	105,462	64,935	40,527	23,782	14,168	9,614	1,089	658	431	2,720	1,612	1,108
60-64 years	148,348	87,614	60,734	121,646	72,226	49,420	22,614	13,030	9,584	1,137	636	501	2,951	1,722	1,229
65-69 years	171,883	98,721	73,162	142,813	82,594	60,219	24,208	13,369	10,839	1,257	687	570	3,605	2,071	1,534
70-74 years	218,210	119,444	98,766	185,765	102,331	83,434	26,918	14,218	12,700	1,195	608	587	4,332	2,287	2,045
75-79 years	297,710	154,227	143,483	261,623	136,662	124,961	29,260	14,144	15,116	1,245	607	638	5,582	2,814	2,768
80-84 years	369,628	173,088	196,540	331,393	156,342	175,051	30,742	13,178	17,564	1,297	549	748	6,196	3,019	3,177
85-89 years	352,112	141,809	210,303	321,805	130,335	191,470	23,915	8,617	15,298	900	366	534	5,492	2,491	3,001
90-94 years	238,362	75,514	162,848	218,724	69,431	149,293	15,551	4,411	11,140	534	171	363	3,553	1,501	2,052
95-99 years	92,236	21,168	71,068	84,172	19,209	64,963	6,617	1,429	5,188	182	38	144	1,265	492	773
100 years & over.	19,282	3,087	16,195	16,844	2,647	14,197	2,059	321	1,738	60	16	44	319	103	216
Age not stated	220	171	49	179	142	37	32	22	10	2	1	1	7	6	1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other salmonella infections (A02) All ages..... 10 Under 1 year.... 3 2 1 1 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 2 2 60-64 years..... 2 1 1 65-69 vears..... 70-74 vears..... 6 75-79 years..... 5 80-84 years..... 4 1 1 85-89 years..... 1 2 90-94 years..... 4 2 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Salmonella gastroenteritis (A02.0) All ages..... 15 Under 1 year.... 3 2 1 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 1 85-89 years..... 90-94 years..... 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Salmonella septicemia (A02.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Localized salmonella infections (A02.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Salmonella infection, unspecified (A02.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Shigellosis (A03) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Shigellosis, unspecified (A03.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other bacterial intestinal infections (A04) All ages.... 6,318 2,440 3,878 5,879 2,252 3,627 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 Q 50-54 years..... 2.2 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.1 2.4 80-84 years..... 1,430 1,360 1,351 1,287 85-89 years..... 90-94 years..... 2.4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other intestinal Escherichia coli infections (A04.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Enterocolitis due to Clostridium difficile (A04.7) All ages.... 6,225 2,408 3,817 5,792 2,224 3,568 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 2.8 1 Ω 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.1 2.4 80-84 years..... 1,411 1,342 1,337 1,273 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International															
Classification of Dis	eases.											Δs	ian or		
Tenth Revision, Second		All rac	es		White			Black		Amer	cican Ind			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both		_,,	Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Other specified bacte	rial intesti	nal infec	tions (A04.8	3)											
All ages	66	24	42	61	21	40	4	3	1	_	_	_	1	_	1
Under 1 year	1	-	1	1	-	1	-	-	-	_	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
50-54 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
70-74 years	5	3	2	4	3	1	-	-	-	-	-	-	1	-	1
75-79 years	19	7	12	19	7	12	-	-	-	-	-	-	-	-	-
80-84 years	13	6	7	12	5	7	1	1	-	-	-	-	-	-	-
85-89 years	10	2	8	10	2	8	-	-	-	-	-	-	-	-	-
90-94 years	8	-	8	7	-	7	1	-	1	-	-	-	-	-	-
95-99 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
100 years & over.	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based															
on the International Classification of Dis	02505											λα	ian or		
Tenth Revision, Second		All rac	100		White			Black		λmer	rican Ind			ic Tela	nder 2/
Edition, 2004)	Both	AII IAC	.Cb	Both	WIIICE		Both	Black		Both	.ican inu	ian i/	Both	IC ISIA	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Bacterial intestinal	infection, u	ınspecifie	ed (A04.9)												
All ages	24	8	16	23	7	16	1	1	_	_	_	_	_	_	_
Under 1 year	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
1-4 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-
75-79 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
80-84 years	6	2	4	6	2	4	-	-	-	-	-	-	-	-	-
85-89 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-
90-94 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

24JUN09

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis													sian or		
Tenth Revision, Secon		All rac	es		White			Black			cican Indi	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both	24.1.	m1.	Both	M . 1 .	m1.	Both	34.1.		Both	M.1		Both	M . 1 .	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male E	emale	sexes		Female
Other bacterial food-	borne intox:	ications (A05)												
All ages	6	3	3	4	2	2	2	1	1	_	_	_	_	_	_
Under 1 year	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
65-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
70-74 years	2	-	2	1	-	1	1	-	1	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise													ian or		
Tenth Revision, Second		All rac	es		White			Black			rican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both	_	_	Both	_	_	Both	_	_	Both	_	_	Both	_	_
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale ———	sexes	Male	Female
Botulism (A05.1)															
All ages	3	2	1	2	1	1	1	1	-	-	_	-	-	_	-
Under 1 year	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
65-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Food-borne Vibrio parahemolyticus intoxication (A05.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis	eases,											As	ian or		
Tenth Revision, Second	d	All rac	es		White			Black		Amer	ican Indi	ian 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
Bacterial food-borne	intoxication	ı, unspeci	fied (A05.9)												
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
70-74 years	_	_	-	-	-	-	_	-	-	_	-	_	-	_	-
75-79 years	_	-	-	-	-	_	-	-	-	_	-	-	-	-	-
0-84 years	1	1	-	1	1	-	_	-	-	_	-	_	-	_	-
5-89 years	_	_	-	-	-	-	_	-	-	_	-	_	-	_	-
0-94 years	_	_	-	-	-	-	_	-	-	_	-	_	-	_	-
95-99 years	_	-	-	-	-	_	-	-	-	_	-	-	-	-	-
100 years & over.	_	_	-	-	-	-	_	-	-	_	-	_	-	_	-
Age not stated	-	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise		-11						-1 1			1		sian or		1 0/
Tenth Revision, Second		All rac	ces	D - 1-1-	White		D - 1-1-	Black			rican Indi	ian 1/		ic Isla	inder 2/
Edition, 2004) and age	Both	Male	Female	Both sexes	Male	Female	Both	Male	Female	Both	Male H	Zomolo	Both sexes	Mala	Female
	sexes	Male	remare	sexes	Male	remare	sexes	мате		sexes	Male i	ellale			remare
Amebiasis (A06)															
All ages	3	1	2	3	1	2	-	_	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Amebic liver abscess (A06.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise								_, ,		_			ian or		
Tenth Revision, Second		All rac	es		White			Black			ican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both	Male	E1-	Both	Male	Female	Both	Male	Female	Both	Male F	1_	Both	Mala	Female
and age	sexes	мате	Female	sexes	мате	remaie	sexes	мате	remaie	sexes	мате г		sexes	мате	
Amebiasis, unspecified	d (A06.9)														
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
20-24 years	-	-	-	-	_	-	-	_	-	_	-	-	-	-	_
25-29 years	-	-	-	-	_	-	-	_	-	_	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
00-94 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other protozoal intestinal diseases (A07) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise		-11						-1 1					ian or		1 0 /
Tenth Revision, Second		All rac	es	Dath	White		Dath	Black		Ame: Both	rican Indi	an 1/		ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	sexes	Male F	emale	Both sexes	Male	Female
Giardiasis [lambliasi:	s] (A07.1)														
All ages	1	-	1	1	_	1	_	_	-	_	_	_	_	_	-
Under 1 year	-	-	-	-	_	-	_	-	-	-	-	-	-	-	-
1-4 years	_	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	_	-	-	-	-	-	_	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis												As	ian or		
Tenth Revision, Secon	d	All rac	es		White			Black		Amer	cican Ind	ian 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Cryptosporidiosis (A0	7.2)														
All ages	2	1	1	_	-	_	1	_	1	-	_	-	1	1	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
40-44 years	1	-	1	-	_	-	1	-	1	-	-	-	-	-	-
45-49 years	1	1	-	_	-	-	-	-	-	-	-	-	1	1	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
Age not stated	-	-	-	_	_	-	-	-	-	-	-	-	-	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Viral and other specified intestinal infections (A08) All ages.... 153 99 Under 1 year.... 6 2 4 3 1 2 1 1-4 years..... 4 3 3 2 1 2 2 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 45-49 years..... 1 1 7 50-54 years..... 3 3 55-59 years..... 4 4 60-64 years..... 4 3 1 65-69 vears..... 6 70-74 vears..... 10 8 75-79 years..... 17 10 7 16 6 80-84 years..... 19 7 12 18 7 11 33 22 22 85-89 years..... 11 33 11 27 1 90-94 years..... 23 2.6 22 4 4 95-99 years..... 14 2 12 13 11 1 1 2 100 years & over. 5 2 3 3 5

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Rotaviral enteritis (A08.0) All ages..... Under 1 year.... 1 2 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acute gastroenteropathy due to Norwalk agent (A08.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Adenoviral enteritis (A08.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other viral enteritis (A08.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Viral intestinal infection, unspecified (A08.4) All ages.... 139 Under 1 year.... 4 2 2 2 1 1 1 1 1-4 years..... 4 2 2 1 1 2 1 2 2 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 45-49 years..... 1 1 1 6 50-54 years..... 2 2 55-59 years..... 4 4 60-64 years..... 3 2 65-69 vears..... 6 70-74 vears..... 10 8 75-79 years..... 17 10 7 16 6 80-84 years..... 17 6 11 16 6 10 30 85-89 years..... 10 20 30 10 20 25 1 90-94 years..... 21 2.4 2.0 4 4 95-99 years..... 13 2 11 12 10 1 1 2 100 years & over. 4 2 2 2 4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified intestinal infections (A08.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Diarrhea and gastroenteritis of infectious origin (A09) All ages.... 132 13 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 3 1 55-59 years..... 4 2 2 60-64 years..... 8 3 5 1 65-69 vears..... 5 3 70-74 vears..... 10 1 75-79 years..... 25 10 14 80-84 years..... 36 12 24 35 11 24 33 30 3 85-89 years..... 12 21 12 18 90-94 years..... 14 2 12 13 11 1 1 2 95-99 years..... 5 5 5 5 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis													ian or		
Tenth Revision, Secon		All rac	es		White			Black			ican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		_ ,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Respiratory tuberculo	sis, not cor	nfirmed ba	cteriologica	ally or hist	ologicall	y (A16)									
All ages	490	335	155	287	182	105	121	91	30	11	7	4	71	55	16
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
25-29 years	6	4	2	2	-	2	2	2	-	1	1	-	1	1	-
30-34 years	8	4	4	3	2	1	4	2	2	1	-	1	-	-	-
35-39 years	5	3	2	3	2	1	2	1	1	-	-	-	-	-	-
40-44 years	17	17	-	10	10	-	6	6	-	-	-	-	1	1	-
45-49 years	34	28	6	17	13	4	10	8	2	4	4	-	3	3	-
50-54 years	33	26	7	15	13	2	14	9	5	-	-	-	4	4	-
55-59 years	35	30	5	16	13	3	16	14	2	-	-	-	3	3	-
60-64 years	48	40	8	28	22	6	14	13	1	-	-	-	6	5	1
65-69 years	41	31	10	24	17	7	10	9	1	1	1	-	6	4	2
70-74 years	43	28	15	21	13	8	16	10	6	1	1	-	5	4	1
75-79 years	54	38	16	37	24	13	7	7	-	1	-	1	9	7	2
80-84 years	78	44	34	51	29	22	12	6	6	2	-	2	13	9	4
85-89 years	51	24	27	32	13	19	5	2	3	-	-	-	14	9	5
90-94 years	30	16	14	22	9	13	2	2	-	-	-	-	6	5	1
95-99 years	4	1	3	3	1	2	1	-	1	-	-	-	-	-	-
100 years & over.	_	_	_	_		_	_	_				_	_		_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Bot.h and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Tuberculosis of lung, without mention of bacteriological or histological confirmation (A16.2) All ages.... 19 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 2 1 40-44 years..... 13 13 2 8 45-49 years..... 22 18 a 19 8 50-54 years..... 15 1 0 2 55-59 years..... 23 19 15 13 4 2 5 1 1 60-64 years..... 32 26 6 17 13 6 5 65-69 vears..... 22 16 15 11 5 6 70-74 vears..... 29 17 12 15 9 6 10 3 75-79 years..... 41 3.0 11 29 19 5 6 80-84 years..... 44 24 20 26 15 11 10 1 7 4 2 85-89 years..... 31 15 16 18 8 10 1 11 6 90-94 years..... 19 11 8 15 8 4 4 95-99 years..... 3 2 3 1 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Classification of Dis Tenth Revision, Secon		All race	ag .		White			Black		Amer	ican Indi		ian or	ic Isla	nder 2/
Edition, 2004)	Both	1111 1400		Both			Both	Diaon		Both	10011 11101	u11 1/	Both	10 1010	11401 27
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Tuberculosis of laryn	x, trachea,	and bronch	nus, without	mention of	bacterio	ological or	histologic	al confi	rmation (A1	6.4)					
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-54 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-59 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-74 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
		_	_	_		_	_	_	_			_	_	_	_
95-99 years	_	_	_	_	_	_	_	_	_	_	_				

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female Male Female sexes Male sexes Male sexes sexes sexes Tuberculous pleurisy, without mention of bacteriological or histological confirmation (A16.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Bot.h Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Primary respiratory tuberculosis without mention of bacteriological or histological confirmation (A16.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Distent Revision, Second Edition, 2004)		All rac	es	Both	White		Both	Black		Amer Both	ican Indi		ian or Pacif Both	ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Respiratory tuberculo	sis unspecif	ied, with	out mention	of bacterio	logical o	or histologi	cal confir	mation (<i>I</i>	116.9)						
All ages	175	120	55	97	59	38	48	37	11	7	4	3	23	20	3
Inder 1 year	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-9 years	1	_	1	1	-	1	-	-	-	-	-	-	-	-	-
0-14 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-29 years	2	1	1	1	-	1	-	-	-	1	1	-	-	-	-
0-34 years	7	3	4	2	1	1	4	2	2	1	-	1	-	-	-
5-39 years	2	1	1	1	1	-	1	-	1	-	-	-	-	-	-
0-44 years	3	3	-	1	1	-	1	1	-	-	-	-	1	1	-
5-49 years	12	10	2	8	6	2	2	2	_	2	2	-	-	-	-
0-54 years	13	10	3	4	4	-	6	3	3	-	-	-	3	3	-
5-59 years	11	10	1	1	_	1	8	8	_	-	-	-	2	2	-
0-64 years	16	14	2	11	9	2	5	5	-	-	-	-	-	-	-
5-69 years	19	15	4	9	6	3	5	5	-	1	1	-	4	3	1
0-74 years	12	10	2	5	4	1	6	5	1	-	-	-	1	1	-
5-79 years	13	8	5	8	5	3	2	2	-	1	-	1	2	1	1
0-84 years	32	20	12	23	14	9	2	1	1	1	-	1	6	5	1
5-89 years	20	9	11	14	5	9	3	1	2	-	-	-	3	3	-
0-94 years	10	5	5	7	2	5	2	2	-	-	-	-	1	1	-
5-99 years	1	_	1	_	_	_	1	-	1	-	-	-	-	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Tuberculosis of nervous system (A17) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 2 1 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Tuberculous meningitis (A17.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other tuberculosis of nervous system (A17.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Tuberculosis of other organs (A18) All ages.... 43 31 20 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 45-49 years..... 5 2 3 2 7 3 50-54 years..... 55-59 years..... 6 4 2 60-64 years..... 2 1 1 65-69 vears..... 10 6 70-74 vears..... 12 75-79 years..... 13 8 80-84 years..... 13 6 85-89 years..... 13 9 11 1 90-94 years..... 6 2 6 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Tuberculosis of bones and joints (A18.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 3 40-44 years..... 2 45-49 years..... 5 2 3 2 6 2 50-54 years..... 6 55-59 years..... 5 3 2 60-64 years..... 2 1 1 65-69 vears..... 10 6 70-74 vears..... 10 6 75-79 years..... 13 8 80-84 years..... 11 6 3 10 85-89 years..... 6 1 90-94 years..... 6 2 4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Tuberculosis of genitourinary system (A18.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

24JUN09

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Tuberculous peripheral lymphadenopathy (A18.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Tuberculosis of intestines, peritoneum, and mesenteric glands (A18.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Tuberculosis of adrenal glands (A18.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Tuberculosis of other specified organs (A18.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Miliary tuberculosis (A19) All ages.... 23 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 3 45-49 years..... 8 1 2 8 3 50-54 years..... 2 55-59 years..... 5 3 2 60-64 years..... 2 1 1 65-69 vears..... 6 5 70-74 vears..... 2 75-79 years..... 80-84 years..... 1 6 85-89 years..... 3 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis	seases,	711	_		White			Dll-		D-11-0-1	.i Tudi		ian or	:- T-1-	
Tenth Revision, Secon		All race	:S	D	wnite		D - 1-1-	Black			ican Indi	an 1/		ic ista	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
acute miliary tubercu	losis of mul	tiple site	es (A19.1)												
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-4 years	_	_	-	_	_	-	-	-	-	-	-	-	-	-	-
-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
0-24 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
5-29 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-39 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
0-44 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
5-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
5-79 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
00 years & over.	_	_	-	-	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acute miliary tuberculosis, unspecified (A19.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Miliary tuberculosis, unspecified (A19.9) All ages.... 23 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 8 1 2 3 50-54 years..... 2 55-59 years..... 5 2 1 60-64 years..... 1 1 65-69 vears..... 5 70-74 vears..... 2 75-79 years..... 80-84 years..... 1 6 85-89 years..... 3 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise	eases,											As	ian or		
Tenth Revision, Second	d	All rac	ces		White			Black		Ame	cican Ind	ian 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male I	Female	sexes	Male	Female
Plague (A20)															
All ages	3	-	3	2	-	2	-	-	-	1	-	1	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	1	-	1	-	-	-	-	-	-	1	-	1	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
00-94 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	_	-	_	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pneumonic plague (A20.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis	eases,												ian or		
Tenth Revision, Secon		All rac	es		White			Black			rican Ind	lian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Septicemic plague (A2	0.7)														
All ages	2	-	2	1	-	1	_	_	-	1	-	1	_	_	-
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	1	-	1	-	-	-	-	-	-	1	-	1	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Brucellosis (A23) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Brucellosis due to Brucella melitensis (A23.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Brucellosis, unspecified (A23.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Leptospirosis (A27) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Leptospirosis, unspecified (A27.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other zoonotic bacterial diseases, not elsewhere classified (A28) All ages..... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 3 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 5 70-74 vears..... 75-79 years..... 80-84 years..... 2 3 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc	eases.											As	ian or		
Tenth Revision, Second		All rac	es		White			Black		Amer	ican Indi			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both		,	Both		,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Pasteurellosis (A28.0))														
All ages	13	4	9	13	4	9	_	_	_	_	_	_	_	_	_
Under 1 year	_	_	_	_	_	_	_	-	_	_	-	-	-	-	-
1-4 years	-	_	_	-	_	_	_	-	_	_	-	-	-	-	-
5-9 years	-	_	-	_	_	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
20-24 years	-	_	-	_	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	_	-	_	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	_	-	_	-	-	-	_	-	-	-	-	-	-	-
45-49 years	2	-	2	2	-	2	-	_	-	_	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
65-69 years	4	2	2	4	2	2	-	_	-	-	-	-	-	-	-
70-74 years	1	-	1	1	-	1	-	_	-	_	-	-	-	-	-
75-79 years	-	_	-	_	_	-	-	-	-	-	-	-	-	-	-
30-84 years	2	_	2	2	_	2	-	-	-	-	-	-	-	-	-
35-89 years	2	_	2	2	_	2	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis		211			referènce			Dl. de					ian or		
Tenth Revision, Secon		All rac	ces	Both	White		Dath	Black		Ame: Both	rican Indi	an I/	Pacii Both	ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	sexes	Male	Female	Both sexes	Male	Female	sexes	Male F	omalo	sexes	Mala	Female
						remare		naic				Ciliaic		Haic	remare
Extraintestinal yersi	niosis (A28.	.2)													
All ages	1	-	1	_	_	_	1	_	1	_	_	_	_	_	_
Under 1 year	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	_	-	_	-	-	-	-	_	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	_	-	_	-	-	-	-	_	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	_	-	_	-	-	-	-	_	-	-	-	-	-	-
75-79 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
95-99 years	1	-	1	_	-	-	1	-	1	-	-	-	-	-	-
100 years & over.	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_		_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All race	es		White			Black			rican Indi			ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
ther specified zoono	tic bacteria	l diseases	s, not else	where classi	fied (A28	3.8)									
All ages	7	3	4	6	2	4	1	1	-	-	-	-	-	-	-
nder 1 year	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
5-69 years	1	-	1	1	_	1	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-79 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-
0-84 years	-	-	-	-	_	-	-	-	-	_	_	-	-	-	-
5-89 years	1	1	-	1	1	-	-	-	-	_	_	-	-	-	-
0-94 years	-	-	-	-	_	-	-	-	-	_	_	-	-	-	-
5-99 years	-	-	-	-	_	-	-	-	-	_	_	-	-	-	-
00 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis		222			rale à le c			D11-					ian or		
Tenth Revision, Second	a Both	All rac	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/	Pacii Both	ic Isla	nder 2/
Edition, 2004) and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Leprosy [Hansen's dis	ease] (A30)														
All ages	1	1	-	_	_	_	1	1	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	_	_	-	_	_	_	-	-	_	-	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Leprosy, unspecified (A30.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Infection due to other mycobacteria (A31) All ages.... 128 194 120 19 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 8 1 3 50-54 years..... 2 3 1 55-59 years..... 14 9 10 5 3 1 60-64 years..... 21 13 8 17 10 65-69 vears..... 22 8 17 3 9 70-74 vears..... 24 5 19 23 19 75-79 years..... 40 2.0 36 18 80-84 years..... 34 13 21 29 9 20 1 3 85-89 years..... 38 8 30 35 28 90-94 years..... 11 2 9 11 2 9 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Pulmonary mycobacterial infection (A31.0) All ages.... 103 161 96 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 7 6 1 2 50-54 years..... 2 2 1 1 55-59 years..... 12 8 5 5 3 60-64 years..... 20 13 16 10 65-69 vears..... 17 11 13 6 6 70-74 vears..... 21 5 16 20 75-79 years..... 31 14 29 16 13 80-84 years..... 28 11 17 25 9 16 2 31 85-89 years..... 34 8 26 24 90-94 years..... 8 2 6 8 2 6 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disc													ian or		
Tenth Revision, Second		All rac	es		White			Black			rican India	an 1/		ic Isla	nder 2/
Edition, 2004)	Both	M . 1 .	T 1 .	Both	M. 1 -	Female	Both	24 . 1 .	Female	Both	W.1		Both	M - 1 -	
and age 	sexes	Male	Female	sexes	Male	remate	sexes	Male	remare	sexes	Male Fe	emale	sexes	мате	Female
Other mycobacterial i	nfections (A	31 8)													
oner my oobaccerrar r		.52.07													
All ages	3	1	2	3	1	2	-	-	-	_	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
5-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-79 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.00 years & over.	_	-	-	-	-	-	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mycobacterial infection, unspecified (A31.9) All ages.... 30 Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 1 55-59 years..... 2 2 60-64 years..... 1 65-69 vears..... 5 70-74 vears..... 3 3 75-79 years..... 8 80-84 years..... 6 85-89 years..... 4 90-94 years..... 3 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Listeriosis (A32) All ages.... 30 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 45-49 years..... 1 50-54 years..... 55-59 years..... 2 2 60-64 years..... 2 1 1 65-69 vears..... 4 70-74 vears..... 4 75-79 years..... 4 80-84 years..... 4 3 2 85-89 years..... 1 90-94 years..... 2 2 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Listerial meningitis and meningoencephalitis (A32.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 45-49 years..... 1 50-54 years..... 55-59 years..... 2 2 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 3 75-79 years..... 3 80-84 years..... 2 1 2 85-89 years..... 1 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis													sian or		
Tenth Revision, Secon		All rac	es		White			Black			rican Ind	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both	M - 1 -	T1-	Both	M . 1 .		Both	24 . 7 .	m1.	Both	M. 1.	1 .	Both	24 . 1 .	1 .
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	remale	sexes		Female
Listerial septicemia	(A32.7)														
All ages	7	3	4	6	3	3	1	_	1	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	_	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	_	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
70-74 years	1	-	1	_	-	-	1	-	1	-	-	-	_	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	_	-	-
80-84 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	_	-	_	-	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other tetanus (A35) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

24JUN09

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise													ian or		
Tenth Revision, Second		All rac	es	_	White		_	Black			rican Indi	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both	1	_ ,	Both	1	_ ,	Both			Both			Both		_ ,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Female	sexes	Male	Female
Whooping cough (A37)															
All ages	9	3	6	7	3	4	2	_	2	-	-	-	-	-	-
Under 1 year	8	2	6	6	2	4	2	-	2	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
70-74 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Whooping cough due to Bordetella pertussis (A37.0) All ages.... Under 1 year.... 3 3 2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis		311			White			Dlasl.		7	T		ian or	:- T-1-	
Tenth Revision, Secon Edition, 2004)	na Both	All rac	es	Both	wnite		Both	Black		Ame: Both	rican Ind	ilan 1/	Pacii Both	ic isia	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Whooping cough, unspe	ecified (A37.	.9)													
All ages	6	3	3	5	3	2	1	_	1	-	_	_	_	_	_
Under 1 year	5	2	3	4	2	2	1	_	1	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise													ian or		
Tenth Revision, Second		All rac	es		White			Black			rican Indi	lan 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both		_ ,	Both		_ 1	Both			Both		_ 1
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Scarlet fever (A38)															
All ages	1	1	_	1	1	-	_	_	-	-	-	_	_	-	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	_	-	-	_	-	-	_	_	-	-	-
00-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_		_	_	_	_		_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Meningococcal infection (A39) All ages.... 67 38 79 50 23 Under 1 year.... 11 1.0 1 10 9 1 1 1 1-4 years..... 14 10 4 13 1 1 2 2 5-9 years..... 4 3 1 10-14 years..... 3 1 2 1 2 15-19 years..... 1.0 14 20-24 years..... 8 6 25-29 years..... 3 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 5 2 3 2 8 50-54 years..... 6 2 55-59 years..... 6 1 5 60-64 years..... 3 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 1 2 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Meningococcal meningitis (A39.0) All ages.... 11 19 12 Under 1 year.... 2 2 1 1 1 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 2 50-54 years..... 1 1 1 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International												_			
Classification of Dise		All rac			White			Black		3	ican Indi		ian or	::- T-1-	nder 2/
Tenth Revision, Second Edition, 2004)	Both	AII rac	es	Both	wille		Both	Black		Both	ican indi	an 1/	Both	ic isia	maer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Mala	Female
															- Ciliare
Waterhouse-Friderichse	en syndrome	(A39.1)													
All ages	29	18	11	24	16	8	5	2	3	_	_	_	_	_	_
Under 1 year	6	5	1	6	5	1	-	-	_	_	_	-	_	_	-
1-4 years	6	4	2	5	3	2	1	1	-	-	-	-	-	-	-
5-9 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
10-14 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
15-19 years	2	-	2	1	-	1	1	-	1	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	1	-	1	-	-	-	1	-	1	_	-	-	-	-	-
50-54 years	4	3	1	3	2	1	1	1	-	_	-	-	-	-	-
55-59 years	2	-	2	2	-	2	-	-	-	_	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	-	-	-	-	_	-	-	-	-	-	-	-	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Acute meningococcemia (A39.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Meningococcemia, unspecified (A39.4) All ages.... 19 13 Under 1 year.... 3 3 3 3 1-4 years..... 6 5 6 5 5-9 years..... 1 10-14 years..... 2 1 1 15-19 years..... 3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 3 2 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other meningococcal infections (A39.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

24JUN09

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Meningococcal infection, unspecified (A39.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Streptococcal septicemia (A40) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.2 80-84 years..... 85-89 years..... 90-94 years..... 2.6 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Septicemia due to streptococcus, group A (A40.0) All ages..... 18 Under 1 year.... 2 2 1 1-4 years..... 2 1 2 1 5-9 years..... 1 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Septicemia due to streptococcus, group B (A40.1) All ages..... 17 11 Under 1 year.... 9 5 6 3 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 55-59 years..... 2 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Septicemia due to streptococcus, group D (A40.2) All ages.... 70 105 Under 1 year.... 2 1 1 1 1 1 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 2 2 6 50-54 years..... 3 3 2 2 55-59 years..... 5 2 3 60-64 years..... 9 5 65-69 vears..... 11 6 5 70-74 vears..... 10 2 8 75-79 years..... 18 10 8 80-84 years..... 25 6 19 2.2 16 17 85-89 years..... 9 8 16 1 1 90-94 years..... 5 2 3 5 2 95-99 years..... 2 2 1 1 4 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Septicemia due to Streptococcus pneumoniae (A40.3) All ages.... 128 27 15 Under 1 year.... 8 3 5 4 2 2 2 1 1 2 2 1-4 years..... 6 2 4 3 1 2 1 1 1 5-9 years..... 1 1 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 4 35-39 years..... 40-44 years..... 3 45-49 years..... 8 5 3 1 17 50-54 years..... 8 9 1 15 55-59 years..... 12 6 5 6 60-64 years..... 11 5 6 11 65-69 vears..... 10 8 70-74 vears..... 6 5 5 75-79 years..... 25 20 3 80-84 years..... 15 7 8 14 1 1 85-89 years..... 6 2 6 90-94 years..... 12 12 5 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other streptococcal septicemia (A40.8) All ages..... 16 12 Under 1 year.... 1 1 1-4 years..... 2 1 1 5-9 years..... 1 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 1 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Streptococcal septicemia, unspecified (A40.9) All ages.... 93 43 50 19 Under 1 year.... 4 3 1 3 2 1 1 1 1-4 years..... 2 1 1 1 1 2 5-9 years..... 1 1 10-14 years..... 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 11 5 Ω 2 7 3 50-54 years..... 3 1 5 55-59 years..... 12 10 5 60-64 years..... 4 3 1 65-69 vears..... 10 3 70-74 vears..... 8 75-79 years..... 17 10 13 80-84 years..... 11 2 9 11 9 12 85-89 years..... 11 11 10 7 90-94 years..... 3 3 4 95-99 years..... 2 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based															
on the International Classification of Disc												3 -	sian or		
Tenth Revision, Second		All rac			White			Black		Am ozo	ican In			iia Tale	nder 2/
Edition, 2004)	a Both	AII rac	ces	Both	wille		Both	Black		Both	ican in	dian i/	Both	IC ISIA	inder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Mala	Female	sexes	Mala	Female
					marc			Marc			maic			marc	
Other septicemia (A41)														
All ages	33,766	15,303	18,463	26,996	12,275	14,721	6,031	2,680	3,351	204	96	108	535	252	283
Under 1 year	243	132	111	141	77	64	93	50	43	3	1	2	6	4	2
1-4 years	75	38	37	52	27	25	17	9	8	2	-	2	4	2	2
5-9 years	35	23	12	24	15	9	10	7	3	-	-	-	1	1	-
10-14 years	43	26	17	34	21	13	9	5	4	-	-	-	-	-	-
15-19 years	45	21	24	34	15	19	9	5	4	1	-	1	1	1	-
20-24 years	87	43	44	52	25	27	29	15	14	2	2	-	4	1	3
25-29 years	119	60	59	74	39	35	38	17	21	3	-	3	4	4	-
30-34 years	157	85	72	101	55	46	47	26	21	3	-	3	6	4	2
35-39 years	315	153	162	200	103	97	104	46	58	5	2	3	6	2	4
40-44 years	533	288	245	356	195	161	164	86	78	8	3	5	5	4	1
45-49 years	870	459	411	569	317	252	277	128	149	13	6	7	11	8	3
50-54 years	1,329	756	573	893	517	376	393	218	175	22	11	11	21	10	11
55-59 years	1,838	1,021	817	1,297	719	578	492	275	217	19	10	9	30	17	13
60-64 years	2,133	1,093	1,040	1,587	809	778	490	251	239	21	13	8	35	20	15
65-69 years	2,585	1,321	1,264	1,969	1,026	943	553	260	293	19	10	9	44	25	19
70-74 years	3,415	1,670	1,745	2,648	1,324	1,324	681	306	375	18	11	7	68	29	39
75-79 years	4,807	2,307	2,500	3,973	1,933	2,040	737	338	399	24	8	16	73	28	45
80-84 years	5,817	2,549	3,268	4,921	2,189	2,732	783	306	477	20	10	10	93	44	49
85-89 years	5,184	2,044	3,140	4,529	1,816	2,713	581	198	383	11	4	7	63	26	37
90-94 years	3,018	955	2,063	2,603	830	1,773	362	102	260	8	5	3	45	18	27
95-99 years	967	230	737	825	201	624	129	26	103	2	-	2	11	3	8
100 years & over.	151	29	122	114	22	92	33	6	27	-	-	-	4	1	3
Age not stated	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Septicemia due to Staphylococcus aureus (A41.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.8 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Septicemia due to other specified staphylococcus (A41.1) All ages..... 18 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 90-94 years..... 2 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Septicemia due to unspecified staphylococcus (A41.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 2.8 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1.0 95-99 years..... 1.3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dise	ases.											As	ian or		
Tenth Revision, Second		All rac	es		White			Black		Amer	ican Indi			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both		-,	Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Septicemia due to Hemo	philus infl	uenzae (A	41.3)												
All ages	10	5	5	10	5	5	_	_	_	_	_	_	_	_	_
Under 1 year	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
10-14 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	_	-	-	_	_	-	-	-	-	-	_
10-44 years	-	-	-	-	_	-	-	_	_	-	-	-	-	-	-
15-49 years	1	1	-	1	1	-	-	-	_	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	1	-	1	1	-	1	-	-	_	-	-	-	-	-	-
70-74 years	1	-	1	1	-	1	-	_	_	-	-	-	-	-	-
75-79 years	-	-	_	-	-	_	-	_	_	-	-	-	-	-	-
30-84 years	1	-	1	1	-	1	-	_	_	-	-	-	-	-	-
85-89 years	1	-	1	1	-	1	-	_	_	-	-	-	-	-	-
90-94 years	-	-	_	-	-	_	-	_	_	-	-	-	-	-	-
95-99 years	-	-	_	-	-	_	-	_	_	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis	eases,												ian or		
Tenth Revision, Secon		All rac	es		White			Black			rican Ind	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both		_ ,	Both		_ 1	Both	1	_ ,	Both			Both	1	_ ,
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Septicemia due to ana	erobes (A41.	. 4)													
All ages	126	40	86	108	35	73	14	5	9	2	_	2	2	_	2
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
10-14 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
25-29 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	-	1	1	-	1	_	-	-	_	-	-	-	-	-
40-44 years	2	1	1	2	1	1	-	-	-	_	_	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	-	-	-	-	-	-	1	-	1	-	-	-
55-59 years	6	2	4	5	2	3	-	-	-	1	-	1	-	-	-
60-64 years	10	5	5	9	4	5	1	1	-	-	-	-	-	-	-
65-69 years	5	4	1	5	4	1	-	-	-	-	-	-	-	-	-
70-74 years	9	1	8	6	1	5	3	-	3	_	_	-	-	-	-
75-79 years	20	6	14	17	4	13	3	2	1	_	_	-	-	-	-
80-84 years	23	5	18	20	5	15	2	-	2	_	_	-	1	-	1
85-89 years	29	10	19	27	10	17	1	-	1	-	-	-	1	-	1
90-94 years	16	4	12	13	3	10	3	1	2	_	_	-	-	-	-
95-99 years	1	1	-	1	1	-	-	-	-	_	_	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
Age not stated	-	-	-	-	-	-	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Septicemia due to other Gram-negative organisms (A41.5) All ages.... Under 1 year.... 2.0 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.8 1 Q a 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise												7	ian or		
Tenth Revision, Second		All rac	00		White			Black		λmer	ican Indi			ic Tela	nder 2/
Edition, 2004)	Both	AII IAC	CS	Both	WIIICE		Both	DIACK		Both	ican inu	Laii I/	Both	IC ISIA	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
Other specified septic	emia (A41.8)													
All ages	87	39	48	63	30	33	21	6	15	1	1	_	2	2	-
Under 1 year	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
1-4 years	2	1	1	1	-	1	-	-	-	-	-	-	1	1	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	-	-	-	-	_	-	-	-	-	1	1	-
35-39 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-
40-44 years	2	2	-	2	2	-	_	-	-	-	-	-	-	-	-
45-49 years	4	2	2	3	2	1	1	-	1	-	-	-	-	-	-
50-54 years	1	-	1	_	-	-	1	-	1	-	-	-	-	-	-
55-59 years	10	5	5	9	4	5	1	1	-	-	-	-	-	-	-
60-64 years	6	2	4	3	1	2	3	1	2	-	-	-	-	-	-
65-69 years	6	6	-	4	4	-	2	2	-	-	-	-	-	-	-
70-74 years	7	5	2	5	3	2	1	1	-	1	1	-	-	-	-
75-79 years	9	1	8	6	1	5	3	-	3	_	-	_	_	-	-
80-84 years	14	5	9	10	5	5	4	-	4	_	-	_	_	-	-
85-89 years	10	5	5	9	5	4	1	-	1	_	-	_	_	-	-
90-94 years	7	-	7	5	-	5	2	-	2	-	-	-	-	-	-
95-99 years	2	1	1	_	-	-	2	1	1	-	-	-	-	-	-
100 years & over.	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Dis													ian or		
Tenth Revision, Secon		All rac	es		White			Black			ican In	dian 1/		ic Isla	inder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Septicemia, unspecifi	ed (A41.9)														
All ages	31,290	14,109	17,181	24,974	11,294	13,680	5,644	2,494	3,150	174	85	89	498	236	262
Under 1 year	191	108	83	109	63	46	76	41	35	1	1	-	5	3	2
1-4 years	62	33	29	41	23	18	17	9	8	2	-	2	2	1	1
5-9 years	31	20	11	22	14	8	8	5	3	-	-	-	1	1	-
10-14 years	33	18	15	26	15	11	7	3	4	-	-	-	-	-	-
15-19 years	38	18	20	27	12	15	9	5	4	1	-	1	1	1	-
20-24 years	77	38	39	47	23	24	25	13	12	1	1	-	4	1	3
25-29 years	108	54	54	68	37	31	33	13	20	3	-	3	4	4	-
30-34 years	144	78	66	93	52	41	43	23	20	3	-	3	5	3	2
35-39 years	290	144	146	183	97	86	97	44	53	4	1	3	6	2	4
10-44 years	485	261	224	318	175	143	156	80	76	7	3	4	4	3	1
15-49 years	786	417	369	504	284	220	261	120	141	11	5	6	10	8	2
50-54 years	1,209	695	514	804	469	335	369	207	162	17	9	8	19	10	9
55-59 years	1,698	938	760	1,193	659	534	464	255	209	14	8	6	27	16	11
60-64 years	1,956	1,006	950	1,449	742	707	454	233	221	19	11	8	34	20	14
65-69 years	2,387	1,207	1,180	1,819	939	880	513	236	277	17	10	7	38	22	16
70-74 years	3,192	1,559	1,633	2,468	1,232	1,236	640	288	352	16	10	6	68	29	39
75-79 years	4,442	2,132	2,310	3,666	1,785	1,881	685	311	374	23	8	15	68	28	40
30-84 years	5,405	2,356	3,049	4,560	2,009	2,551	739	295	444	17	10	7	89	42	47
35-89 years	4,852	1,892	2,960	4,232	1,676	2,556	554	190	364	10	3	7	56	23	33
90-94 years	2,853	896	1,957	2,463	781	1,682	340	94	246	7	5	2	43	16	27
95-99 years	903	210	693	770	185	585	122	23	99	1	-	1	10	2	8
100 years & over.	148	29	119	112	22	90	32	6	26	-	-	_	4	1	3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise	eageg.											As	ian or		
Tenth Revision, Second		All rac	es		White			Black		Amei	rican Indi			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Actinomycosis (A42)															
All ages	7	3	4	4	2	2	3	1	2	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
.0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
10-44 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
15-49 years	1	-	1	_	-	-	1	-	1	_	-	-	-	-	-
50-54 years	-	-	-	-	-	-	_	-	-	-	-	-	_	-	-
55-59 years	-	-	-	_	-	-	-	-	-	_	_	_	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
55-69 years	1	1	-	1	1	-	-	-	-	_	_	_	-	-	-
70-74 years	-	_	-	_	-	-	-	-	_	_	-	-	-	-	-
5-79 years	2	1	1	2	1	1	-	-	_	_	-	-	-	-	-
0-84 years	1	_	1	_	-	-	1	-	1	_	-	-	-	-	-
5-89 years	-	_	-	_	-	-	-	-	_	_	-	-	-	-	-
0-94 years	2	1	1	1	-	1	1	1	_	_	-	-	-	-	-
95-99 years	-	_	_	_	_	_	_	_	_	_	-	_	_	_	-
100 years & over.	_	_	-	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis													ian or		
Tenth Revision, Secon		All rac	es	_	White		_	Black			ican Indi	.an 1/		ic Isla	nder 2/
Edition, 2004)	Both	1	_ ,	Both			Both		_ ,	Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Pulmonary actinomycos	is (A42.0)														
All ages	3	_	3	1	_	1	2	_	2	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years		-	-		-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-		-	-	-	-	-	-
80-84 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	_	-	-	_	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Actinomycotic septicemia (A42.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other forms of actinomycosis (A42.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Diseases,				White Black					Asian or						
Tenth Revision, Secon		All races		Wnite Both			Both	Black		American Indian 1/ Both			Pacific Islander 2/ Both		
Edition, 2004) and age	Both sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Actinomycosis, unspec	eified (A42.9	9)													
All ages	1	1	_	_	_	_	1	1	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	_	-	-	-	-	-	_	_	_	-	_	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Nocardiosis (A43) All ages.... 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 5 50-54 years..... 3 55-59 years..... 3 3 1 60-64 years..... 2 1 65-69 vears..... 2 70-74 vears..... 5 3 75-79 years..... 9 80-84 years..... 15 9 6 1 5 85-89 years..... 5 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International																
Classification of Dis										Asian or						
Tenth Revision, Secon		All rac	es		White			Black		American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both			Both			Both			Both			Both			
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female	
Pulmonary nocardiosis	(A43.0)															
All ages	36	22	14	34	20	14	2	2	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
10-14 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-	
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
35-39 years	-	-	-	_	-	-	-	_	-	_	-	-	-	-	-	
40-44 years	1	1	-	_	-	-	1	1	-	_	-	-	-	-	-	
45-49 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-	
50-54 years	2	1	1	2	1	1	-	_	-	-	-	-	-	-	-	
55-59 years	2	2	-	1	1	-	1	1	-	_	-	-	-	-	-	
60-64 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-	
65-69 years	2	1	1	2	1	1	-	-	-	_	-	-	-	-	-	
70-74 years	4	3	1	4	3	1	-	-	-	_	-	-	-	-	-	
75-79 years	7	4	3	7	4	3	-	-	-	_	-	-	-	-	-	
80-84 years	11	6	5	11	6	5	_	_	-	_	-	-	-	-	-	
85-89 years	4	3	1	4	3	1	_	_	-	_	-	-	-	-	-	
90-94 years	1	-	1	1	-	1	_	_	-	_	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other forms of nocardiosis (A43.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second All races				Miles.							Asian or American Indian 1/ Pacific Islander 2/						
Tenth Revision, Second Edition, 2004)	a Both	All rac	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/	Pacific Islander 2/ Both				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female		
Nocardiosis, unspecif	ied (A43.9)																
All ages	10	6	4	8	5	3	2	1	1	-	-	-	-	-	-		
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
50-54 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-		
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
70-74 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
80-84 years	4	3	1	3	3	-	1	-	1	-	-	-	-	-	-		
85-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
95-99 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-		
100 years & over.	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-		
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second All races								-1 1	Asian or American Indian 1/ Pacific Islander 2/						
Tenth Revision, Second Edition, 2004)	a Both	All rac	es	Both	White		Both	Black		Ame: Both	rican Indi	an 1/	Pacii Both	ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Erysipelas (A46)															
All ages	1	1	_	1	1	-	-	_	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other bacterial diseases, not elsewhere classified (A48) All ages.... 137 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 2 2 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 3 35-39 years..... 9 40-44 years..... 4 1 45-49 years..... 13 3 10 Ω 50-54 years..... 12 9 3 9 12 55-59 years..... 18 13 16 12 5 60-64 years..... 19 11 8 17 10 2 65-69 vears..... 17 15 9 9 8 70-74 vears..... 13 6 75-79 years..... 16 15 10 6 80-84 years..... 9 5 3 1 85-89 years..... 14 10 14 10 90-94 years..... 3 4 1 4 1 3 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Gas gangrene (A48.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Legionnaires' disease (A48.1) All ages..... 30 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 2 2 45-49 years..... 4 2 8 50-54 years..... 1 55-59 years..... 10 8 2 9 2 60-64 years..... 12 5 10 65-69 vears..... 9 5 8 70-74 vears..... 12 5 75-79 years..... 10 9 80-84 years..... 4 1 85-89 years..... 10 3 10 90-94 years..... 2 1 2 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Toxic shock syndrome (A48.3) All ages.... 33 47 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 2 2 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 9 45-49 years..... 1 8 3 50-54 years..... 2 1 55-59 years..... 6 5 1 60-64 years..... 5 2 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 2 2 85-89 years..... 90-94 years..... 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified bacterial diseases (A48.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Bacterial infection of unspecified site (A49) All ages.... 1,125 Under 1 year.... 2.0 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 a 50-54 years..... ρ 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Staphylococcal infection, unspecified (A49.0) All ages.... 201 354 183 171 53 Under 1 year.... 7 5 2 2 1 1 5 4 1 1-4 years..... 3 3 2 2 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 3 25-29 years..... 1 1 30-34 years..... 35-39 years..... 8 40-44 years..... 5 45-49 years..... 8 6 2 6 2 1 2.0 50-54 years..... 1.3 15 1 0 55-59 years..... 21 14 7 17 11 6 60-64 years..... 29 13 16 23 10 13 65-69 years..... 33 19 25 18 7 14 70-74 years..... 35 22 13 30 18 12 75-79 years..... 44 2.0 2.4 38 16 2.2 3 80-84 years..... 76 36 40 71 34 37 2 65 85-89 years..... 66 30 36 30 35 1 90-94 years..... 44 18 26 39 16 23 4 3 95-99 years..... 10 8 2 2 4 6 4 4 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Streptococcal infection, unspecified (A49.1) All ages.... 19 Under 1 year.... 2 2 2 2 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 1 1 20-24 years..... 1 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 2 2 5 50-54 years..... 3 2 1 55-59 years..... 8 3 5 6 60-64 years..... 3 65-69 vears..... 7 70-74 vears..... 7 5 75-79 years..... 10 80-84 years..... 14 8 6 11 3 85-89 years..... 8 5 3 1 90-94 years..... 9 5 8 3 1 1 4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hemophilus influenzae infection, unspecified (A49.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mycoplasma infection, unspecified (A49.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other bacterial infections of unspecified site (A49.8) All ages.... 114 18 10 Under 1 year.... 3 2 1 3 1 1-4 years..... 1 1 5-9 years..... 1 1 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 2 5 50-54 years..... 3 55-59 years..... 9 2 4 4 60-64 years..... 7 3 65-69 vears..... 10 6 9 70-74 vears..... 13 5 8 13 75-79 years..... 17 7 80-84 years..... 27 11 16 25 10 15 1 22 85-89 years..... 10 12 18 10 8 90-94 years..... 12 6 6 11 5 95-99 years..... 3 3 2 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Bacterial infection, unspecified (A49.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.3 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.8 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Congenital syphilis (A50) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Congenital syphilis, unspecified (A50.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Early syphilis (A51) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other secondary syphilis (A51.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Late syphilis (A52) All ages.... 31 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 2 45-49 years..... 1 50-54 years..... 1 1 1 55-59 years..... 2 1 1 1 60-64 years..... 4 2 2 3 65-69 vears..... 2 70-74 vears..... 2 75-79 years..... 2 80-84 years..... 4 2 85-89 years..... 90-94 years..... 1 1 1 1 95-99 years..... 3 1 1 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cardiovascular syphilis (A52.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Symptomatic neurosyphilis (A52.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Neurosyphilis, unspecified (A52.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 2 75-79 years..... 80-84 years..... 3 2 85-89 years..... 90-94 years..... 1 1 1 1 95-99 years..... 3 2 1 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Late syphilis, unspecified (A52.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other and unspecified syphilis (A53) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Latent syphilis, unspecified as early or late (A53.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Syphilis, unspecified (A53.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																	
Classification of Dis	eases,									Asian or							
Tenth Revision, Second	d All races			White			Black		Amer	ican Indi		Pacific Islander 2,					
Edition, 2004)	Both			Both			Both			Both			Both				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female		
Gonococcal infection	(A54)																
All ages	3	2	1	1	_	1	2	2	_	_	_	_	_	_	_		
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
10-14 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-		
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
25-29 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-		
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-		
40-44 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-		
45-49 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-		
50-54 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-		
55-59 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-		
60-64 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-		
65-69 years	1	1	-	_	-	_	1	1	-	_	-	-	_	-	-		
70-74 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-		
75-79 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-		
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
85-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
90-94 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-		
95-99 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-		
100 years & over.	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other gonococcal infections (A54.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Gonococcal infection, unspecified (A54.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other predominantly sexually transmitted diseases, not elsewhere classified (A63) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis									Asian or						
Tenth Revision, Secon		All rac	ces		White			Black		American Indian 1/			Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Anogenital (venereal)	warts (A63.	.0)													
All ages	1	_	1	_	_	_	1	_	1	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	_	-	-	-	_	-	-	_	-	_	-	-	_	-	-
15-19 years	_	-	-	-	_	-	-	-	-	_	-	-	_	-	-
20-24 years	_	-	-	-	_	-	-	-	-	_	-	-	_	-	-
25-29 years	_	-	-	-	_	-	-	-	-	_	-	-	_	-	-
30-34 years	_	-	-	-	_	-	-	-	-	_	-	-	_	-	-
35-39 years	_	-	-	-	_	-	-	-	-	_	_	-	_	-	-
40-44 years	_	-	-	-	_	-	-	-	-	_	_	-	_	-	-
45-49 years	1	_	1	-	-	_	1	-	1	_	_	-	_	_	-
50-54 years	_	_	-	-	-	_	-	-	-	_	_	-	_	_	-
55-59 years	_	_	-	-	-	_	-	-	-	_	_	-	_	_	-
60-64 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
70-74 years	_	-	-	-	-	-	-	_	-	_	_	-	-	-	-
'5-79 years	_	-	_	-	_	_	_	_	_	_	-	_	-	-	_
30-84 years	_	-	_	-	_	_	_	_	_	_	-	_	-	-	_
35-89 years	_	-	_	-	_	_	_	_	_	_	-	_	-	-	_
90-94 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
95-99 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_	_	_	_	_		_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Unspecified sexually transmitted disease (A64) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second All races									Asian or						
		All rac	es		White			Black		American Indian 1/			Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale ————	sexes	Male	Female
Other spirochetal inf	ections (A69	9)													
All ages	5	2	3	5	2	3	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
40-44 years	1	-	1	1	_	1	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	_
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	_
70-74 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Tenth Revision, Second All races											Asian or American Indian 1/ Pacific Islander 2/						
		All rac	es	_	White		_	Black			rican Indi	an 1/		ic Isla	nder 2/		
Edition, 2004)	Both			Both			Both			Both			Both				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female		
Lyme disease (A69.2)																	
All ages	5	2	3	5	2	3	_	_	-	-	-	-	_	_	_		
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
1-4 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-		
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
25-29 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
35-39 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-		
40-44 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
55-59 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-		
60-64 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-		
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
70-74 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
35-89 years	-	_	-	_	_	_	-	-	_	_	-	-	_	-	-		
90-94 years	-	_	-	_	_	_	-	-	_	_	-	-	_	-	-		
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Spotted fever [tick-borne rickettsioses] (A77) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Spotted fever due to Rickettsia rickettsii (A77.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other spotted fevers (A77.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Classification of Diseases, Tenth Revision, Second All races				White Black						Asian or American Indian 1/ Pacific Islander 2/						
Edition, 2004)	Both	THE TUCCE		Both	MILICE		Both	DIACK		Both	rican muran i/		Both	iliaci 2/		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Q fever (A78)																
All ages	2	2	_	2	2	_	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other rickettsioses (A79) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Rickettsiosis, unspecified (A79.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Atypical virus infections of central nervous system (A81) All ages.... 237 117 120 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 1 2 1 1 50-54 years..... 20 8 12 1.8 11 55-59 years..... 34 20 14 33 19 14 60-64 years..... 38 18 20 34 17 17 65-69 vears..... 42 19 23 40 17 23 70-74 vears..... 38 16 22 38 22 75-79 years..... 38 22 16 38 2.2 16 80-84 years..... 18 8 10 17 9 5 85-89 years..... 3 2 5 3 2 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Creutzfeldt-Jakob disease (A81.0) All ages..... 109 104 101 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 1 1 17 50-54 years..... 6 11 11 17 6 55-59 years..... 29 17 12 28 12 16 60-64 years..... 36 18 18 33 17 16 65-69 vears..... 39 19 20 37 17 20 70-74 vears..... 33 14 19 33 19 75-79 years..... 34 21 13 34 13 80-84 years..... 16 8 8 15 85-89 years..... 4 2 2 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Subacute sclerosing panencephalitis (A81.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Progressive multifocal leukoencephalopathy (A81.2) All ages.... 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 3 2 50-54 years..... 1 55-59 years..... 4 2 2 60-64 years..... 1 1 65-69 vears..... 3 3 70-74 vears..... 5 75-79 years..... 4 80-84 years..... 2 2 2 85-89 years..... 90-94 years..... 1 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Atypical virus infection of central nervous system, unspecified (A81.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dise										Asian or						
Tenth Revision, Second		All races		White			Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both			Both		_	Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Rabies (A82)																
All ages	2	1	1	2	1	1	_	-	-	-	-	-	_	_	-	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
15-19 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
75-79 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
80-84 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-89 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
90-94 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
95-99 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Rabies, unspecified (A82.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mosquito-borne viral encephalitis (A83) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Eastern equine encephalitis (A83.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes St. Louis encephalitis (A83.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes California encephalitis (A83.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other viral encephalitis, not elsewhere classified (A85) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Enteroviral encephalitis (A85.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified viral encephalitis (A85.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Unspecified viral encephalitis (A86) All ages.... 110 59 97 10 Under 1 year.... 6 2 4 5 2 3 1 1-4 years..... 5 3 2 2 1 3 3 5-9 years..... 10-14 years..... 2 15-19 years..... 2 2 20-24 years..... 25-29 years..... 3 30-34 years..... 35-39 years..... 2 40-44 years..... 45-49 years..... 2 1 1 2 50-54 years..... 4 2 2 2 55-59 years..... 8 3 5 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 14 6 8 13 75-79 years..... 14 3 11 80-84 years..... 15 9 6 14 5 1 12 85-89 years..... 6 6 10 1 1 90-94 years..... 6 2 4 6 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Viral meningitis (A87) All ages.... 19 30 37 16 Under 1 year.... 4 3 1 1 1 2 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 6 50-54 years..... 3 1 55-59 years..... 2 1 1 60-64 years..... 3 2 1 65-69 vears..... 2 70-74 vears..... 6 5 75-79 years..... 4 80-84 years..... 6 2 1 85-89 years..... 4 3 90-94 years..... 3 3 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Enteroviral meningitis (A87.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Lymphocytic choriomeningitis (A87.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Viral meningitis, unspecified (A87.9) All ages..... 27 18 Under 1 year.... 2 1 1 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 4 2 1 55-59 years..... 1 1 60-64 years..... 3 2 65-69 vears..... 2 70-74 vears..... 6 5 75-79 years..... 4 80-84 years..... 6 1 85-89 years..... 4 3 90-94 years..... 3 3 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Unspecified viral infection of central nervous system (A89) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 55-59 years..... 3 3 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other mosquito-borne viral fevers (A92) All ages..... 123 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 4 3 50-54 years..... 3 2 55-59 years..... 4 2 2 60-64 years..... 4 3 6 3 3 65-69 vears..... 16 13 70-74 vears..... 25 10 15 10 13 75-79 years..... 25 16 9 15 9 80-84 years..... 23 16 21 15 2 1 7 85-89 years..... 14 14 7 90-94 years..... 9 2 a 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes West Nile fever (A92.3) All ages..... 136 123 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 50-54 years..... 3 2 55-59 years..... 4 2 2 60-64 years..... 4 3 6 3 3 65-69 vears..... 16 13 70-74 vears..... 25 10 15 10 13 75-79 years..... 25 16 9 15 9 80-84 years..... 23 16 21 15 2 1 7 85-89 years..... 14 14 7 90-94 years..... 9 2 a 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Unspecified arthropod-borne viral fever (A94) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second All races				White Black						Asian or American Indian 1/ Pacific Islander 2/						
Edition, 2004)	Both	1111 14005		Both	WIIICE		Both	Black		Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Other viral hemorrhag	ic fevers, n	ot elsewh	ere classifi	.ed (A98)												
All ages	8	5	3	7	4	3	_	_	_	1	1	_	_	_	_	
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-24 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
25-29 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	1	1	-	-	-	-	-	-	-	1	1	-	-	-	-	
10-44 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-	
55-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hemorrhagic fever with renal syndrome (A98.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Herpes-viral [herpes simplex] infections (B00) All ages.... 127 20 12 Under 1 year.... 13 6 7 3 3 9 6 1 1 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 4 2 55-59 years..... 9 5 4 6 3 60-64 years..... 13 8 5 12 65-69 vears..... 8 70-74 vears..... 10 6 10 6 75-79 years..... 31 28 15 15 16 80-84 years..... 21 5 16 19 5 14 2 13 11 85-89 years..... 1 12 12 1 90-94 years..... 9 2 7 9 2 95-99 years..... 2 1 1 1 1 100 years & over. 2 2 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Herpes-viral meningitis (B00.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Herpes-viral encephalitis (B00.4) All ages..... Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 3 50-54 years..... 2 2 55-59 years..... 5 2 2 3 60-64 years..... 3 65-69 vears..... 4 70-74 vears..... 8 3 5 75-79 years..... 21 10 11 18 80-84 years..... 16 4 12 14 10 2 8 85-89 years..... 8 8 8 2 90-94 years..... 6 4 6 4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Herpes-viral ocular disease (B00.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disseminated herpes-viral disease (B00.7) All ages..... 12 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 60-64 years..... 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other forms of herpes-viral infection (B00.8) All ages..... 13 Under 1 year.... 2 1 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 1 1 60-64 years..... 3 2 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Herpes-viral infection, unspecified (B00.9) All ages.... 19 10 Under 1 year.... 9 3 6 2 2 1 1 1-4 years..... 1 1 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 2 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 90-94 years..... 3 3 3 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Varicella [chickenpox] (B01) All ages..... 13 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 4 3 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

	Classification of Diseases, Tenth Revision, Second All races									Asian or							
		All rac	es		White		_	Black			rican Indi	.an 1/		ic Isla	nder 2/		
Edition, 2004)	Both		_ ,	Both			Both		_ 1	Both			Both				
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female		
Varicella meningitis	(B01.0)																
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_		
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
75-79 years			-	-	_	-	-	-	-	-	-	-	-	-	-		
30-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Varicella encephalitis (B01.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based	l															
on the International												3				
Classification of Dis Tenth Revision, Secon		All races			White Black						Asian or American Indian 1/ Pacific Islander					
Edition, 2004)	Both	AII Iac	es	Both	WIIICE		Both	Black		Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Varicella pneumonia (B01.2)															
All ages	6	4	2	6	4	2	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	_	-	-	-	-	-	-	-	-	_	-	-	
25-29 years	-	-	-	_	-	-	-	-	-	-	-	-	_	-	-	
30-34 years	-	-	-	_	-	-	-	-	-	-	-	-	_	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
40-44 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
45-49 years	1	1		1	1	-	-	-	-	-	-	-	-	-	-	
50-54 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	_	-	-	-	-	-	-	-	-	_	-	-	
70-74 years	1	1		1	1	-	-	-	-	-	-	-	-	-	-	
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
80-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
90-94 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-	
95-99 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	-	-	-	-	-	-	-	_	_	-	_	-	_	_	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Varicella without complications (B01.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Zoster [herpes zoster] (B02) All ages..... 102 151 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 2 50-54 years..... 3 1 1 55-59 years..... 1 1 60-64 years..... 3 3 65-69 vears..... 6 70-74 vears..... 9 3 6 75-79 years..... 12 7 11 80-84 years..... 32 18 14 32 18 14 26 24 15 85-89 years..... 10 16 9 90-94 years..... 42 13 29 40 12 28 2 1 95-99 years..... 19 3 19 3 16 16 100 years & over. 8 7 7 1 1 6

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Zoster encephalitis (B02.0) All ages..... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 3 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Zoster meningitis (B02.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Zoster with other nervous system involvement (B02.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 8 5 85-89 years..... 2 90-94 years..... 5 95-99 years..... 6 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis												λα	ian or		
Tenth Revision, Secor					White			Black		American Indian 1/ Pacific Islande:					
Edition, 2004)	Both	1111 100	,,,,	Both	,,,,,,		Both	Didon		Both		u11 1/	Both	10 1010	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Zoster ocular disease	(B02 3)														
All ages	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	-	_	-	-	-	_	_	_	_	_	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Disseminated zoster (B02.7) All ages..... 19 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 4 3 75-79 years..... 3 80-84 years..... 3 2 85-89 years..... 90-94 years..... 3 3 3 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Zoster with other complications (B02.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Zoster without complication (B02.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 55-59 years..... 1 1 2 60-64 years..... 2 65-69 vears..... 1 70-74 vears..... 3 75-79 years..... 6 3 80-84 years..... 13 8 5 13 18 11 85-89 years..... 6 12 16 17 90-94 years..... 28 10 18 26 2 1 95-99 years..... 9 11 2 11 9 100 years & over. 6 5 6 5

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Bot.h Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other viral infections characterized by skin and mucous membrane lesions, not elsewhere classified (BO8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Enteroviral vesicular pharyngitis (B08.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Classification of Dise Tenth Revision, Second		All rac	10.5	White Black						Asian or American Indian 1/ Pacific Islander 2/						
Edition, 2004)	a Both	All rac	es	Both	wille		Both	Black		Both	rican ino	iian i/	Both	ic isia	maer 2/	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	
Acute hepatitis A (B1	5)															
All ages	34	17	17	26	14	12	6	3	3	_	_	-	2	_	2	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-	
40-44 years	1	-	1	1	-	1	-	_	-	_	-	-	-	-	-	
45-49 years	2	1	1	1	1	-	1	_	1	_	-	-	-	-	-	
50-54 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-	
55-59 years	6	4	2	5	4	1	1	-	1	-	-	-	-	-	-	
60-64 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-	
65-69 years	6	2	4	6	2	4	-	-	-	-	-	-	-	-	-	
70-74 years	1	-	1	-	-	-	-	-	-	_	-	-	1	-	1	
75-79 years	5	2	3	3	2	1	1	-	1	_	-	-	1	-	1	
80-84 years	3	1	2	3	1	2	-	_	-	-	-	-	-	-	-	
85-89 years	1	-	1	1	_	1	-	_	-	-	-	-	-	-	-	
90-94 years	4	2	2	3	1	2	1	1	-	-	-	-	-	-	-	
95-99 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
100 years & over.	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-	
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hepatitis A without hepatic coma (B15.9) All ages..... 17 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 2 50-54 years..... 2 55-59 years..... 6 4 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 1 2 90-94 years..... 4 2 2 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute hepatitis B (B16) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 2.2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acute hepatitis B without delta-agent with hepatic coma (B16.2) All ages.... 13 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 6 45-49 years..... 2 8 50-54 years..... 3 55-59 years..... 5 3 2 60-64 years..... 4 65-69 vears..... 70-74 vears..... 4 75-79 years..... 80-84 years..... 1 2 2 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Classification of Dis Tenth Revision, Secon		All rac	A Q		White Black							Asian or American Indian 1/ Pacific Islander 2/							
Edition, 2004)	Both	AII IAC	CB	Both	WIIICC		Both	DIACK		Both	ican ind	iiaii i/	Both	.1C 151a	macı z/				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female				
Acute hepatitis B wit	hout delta-a	gent and	without hepa	atic coma (B	16.9)														
All ages	539	374	165	341	251	90	95	58	37	3	2	1	100	63	37				
Under 1 year	-	_	-	_	-	-	_	_	-	-	-	-	-	-	-				
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
5-9 years	-	_	-	_	-	-	_	_	-	-	-	-	-	-	-				
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
5-19 years	1	1	-	-	-	-	-	-	-	-	-	-	1	1	-				
20-24 years	3	3	-	-	-	-	2	2	-	-	-	-	1	1	-				
25-29 years	3	3	-	2	2	-	1	1	-	-	-	-	-	-	-				
30-34 years	4	2	2	1	1	-	1	-	1	-	-	-	2	1	1				
35-39 years	27	19	8	11	7	4	9	5	4	1	1	-	6	6	-				
10-44 years	37	29	8	26	22	4	6	2	4	-	-	-	5	5	-				
15-49 years	71	52	19	51	39	12	11	7	4	-	-	-	9	6	3				
0-54 years	95	74	21	67	57	10	16	8	8	1	1	-	11	8	3				
55-59 years	97	64	33	56	39	17	22	15	7	-	-	-	19	10	9				
50-64 years	49	33	16	31	22	9	6	4	2	-	-	-	12	7	5				
55-69 years	43	33	10	23	17	6	10	8	2	-	-	-	10	8	2				
70-74 years	35	18	17	20	12	8	5	1	4	-	-	-	10	5	5				
5-79 years	37	24	13	28	20	8	3	2	1	-	-	-	6	2	4				
80-84 years	19	11	8	1.2	6	6	2	2	-	-	-	-	5	3	2				
5-89 years	13	8	5	9	7	2	1	1	-	1	-	1	2	-	2				
0-94 years	4	_	4	3	_	3	-	_	-	-	-	-	1	-	1				
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other acute viral hepatitis (B17) All ages..... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 6 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Acute hepatitis C (B17.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 6 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified acute viral hepatitis (B17.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																	
Classification of Dis	eases,									Asian or							
Tenth Revision, Secon	d	All rac	es		White			Black		Amer	ican In	dian 1/	Pacific Islander 2/				
Edition, 2004)	Both			Both			Both			Both			Both				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female		
Chronic viral hepatit	is (B18)																
All ages	6,539	4,273	2,266	5,281	3,495	1,786	972	619	353	98	63	35	188	96	92		
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-		
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-		
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
10-14 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-		
15-19 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-		
20-24 years	3	1	2	3	1	2	-	-	-	_	-	-	-	-	-		
25-29 years	13	4	9	12	4	8	1	_	1	_	-	-	-	-	-		
30-34 years	32	22	10	26	17	9	4	3	1	1	1	-	1	1	-		
35-39 years	95	65	30	73	49	24	11	6	5	7	7	-	4	3	1		
40-44 years	359	250	109	297	207	90	46	30	16	10	8	2	6	5	1		
45-49 years	997	696	301	848	603	245	119	72	47	14	10	4	16	11	5		
50-54 years	1,695	1,241	454	1,410	1,042	368	239	165	74	25	18	7	21	16	5		
55-59 years	1,337	978	359	1,037	760	277	254	187	67	22	13	9	24	18	6		
60-64 years	611	405	206	476	317	159	108	72	36	6	2	4	21	14	7		
65-69 years	405	216	189	301	165	136	79	40	39	6	2	4	19	9	10		
70-74 years	335	155	180	271	127	144	34	19	15	1	-	1	29	9	20		
75-79 years	324	133	191	254	105	149	38	19	19	5	2	3	27	7	20		
80-84 years	199	69	130	160	62	98	25	5	20	1	-	1	13	2	11		
85-89 years	92	26	66	80	25	55	8	1	7	-	-	-	4	-	4		
90-94 years	34	10	24	28	9	19	4	-	4	-	-	-	2	1	1		
95-99 years	6	1	5	3	1	2	2	-	2	-	-	-	1	-	1		
100 years & over.	1	-	1	1	_	1	-	-	-	-	-	-	-	-	-		
Age not stated	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic viral hepatitis B without delta-agent (B18.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 9 40-44 years..... 9 45-49 years..... 14 a 5 Ω 2 2 50-54 years..... 14 3 9 3 11 55-59 years..... 17 12 5 13 10 3 60-64 years..... 14 9 5 4 65-69 vears..... 13 8 5 5 70-74 vears..... 8 75-79 years..... 9 6 80-84 years..... 4 3 2 9 3 85-89 years..... 2 90-94 years..... 4 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Chronic viral hepatitis C (B18.2) All ages.... 6,406 4,184 2,222 5,188 3,433 1,755 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1.399 1,034 50-54 years..... 1.679 1,229 1.8 55-59 years..... 1,319 1,024 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.0 80-84 years..... 85-89 years..... 90-94 years..... 2.4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other chronic viral hepatitis (B18.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Chronic viral hepatitis, unspecified (B18.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Unspecified viral hepatitis (B19) All ages.... 38 33 10 Under 1 year.... 1 1 1 1 1-4 years..... 1 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 2 25-29 years..... 30-34 years..... 1 35-39 years..... 4 40-44 years..... 6 45-49 years..... 11 a 2 2 2 16 9 2 50-54 years..... 13 55-59 years..... 9 2 8 60-64 years..... 6 2 65-69 vears..... 4 3 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 3 85-89 years..... 2 1 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Unspecified viral hepatitis with coma (B19.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Unspecified viral hepatitis without coma (B19.9) All ages.... 30 Under 1 year.... 1 1 1 1 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 1 25-29 years..... 30-34 years..... 1 35-39 years..... 4 40-44 years..... 6 45-49 years..... 11 a 2 2 2 16 9 2 50-54 years..... 13 55-59 years..... 9 2 8 60-64 years..... 5 2 3 65-69 vears..... 3 70-74 vears..... 75-79 years..... 5 80-84 years..... 4 3 2 2 85-89 years..... 3 1 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis													ian or			
Tenth Revision, Secon		All rac	es		White			Black			American Indian 1/			Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	
Human immunodeficienc	y virus [HI	V] disease	resulting	in infectio	us and par	asitic dise	ases (B20)									
All ages	4,597	3,274	1,323	2,031	1,618	413	2,485	1,593	892	38	25	13	43	38	5	
Under 1 year	4	1	3	1	-	1	3	1	2	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-	
5-9 years	2	1	1	-	-	-	2	1	1	-	_	-	-	-	-	
10-14 years	2	1	1	-	-	-	2	1	1	-	-	-	-	-	-	
15-19 years	17	7	10	5	3	2	11	4	7	1	_	1	-	-	_	
20-24 years	67	40	27	25	14	11	41	25	16	-	_	-	1	1	-	
25-29 years	175	117	58	63	49	14	112	68	44	-	_	-	-	-	-	
30-34 years	310	189	121	117	88	29	190	98	92	2	2	-	1	1	-	
35-39 years	607	394	213	276	211	65	312	168	144	10	6	4	9	9	-	
40-44 years	898	646	252	437	345	92	449	293	156	8	5	3	4	3	1	
45-49 years	957	678	279	411	336	75	530	330	200	8	6	2	8	6	2	
50-54 years	771	590	181	348	281	67	405	295	110	7	4	3	11	10	1	
55-59 years	419	330	89	175	149	26	240	177	63	1	1	-	3	3	-	
60-64 years	201	157	44	97	84	13	103	72	31	-	-	-	1	1	-	
65-69 years	99	72	27	39	30	9	57	39	18	1	1	-	2	2	-	
70-74 years	36	28	8	20	16	4	14	11	3	-	-	-	2	1	1	
75-79 years	20	16	4	11	9	2	8	6	2	-	-	-	1	1	-	
80-84 years	10	6	4	5	2	3	5	4	1	-	-	-	-	-	-	
85-89 years	1	1	-	1	1	_	-	_	-	-	-	-	_	-	-	
90-94 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-	
95-99 years	1	-	1	-	-	_	1	-	1	-	-	-	-	-	-	
100 years & over.	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-	
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes HIV disease resulting in mycobacterial infection (B20.0) All ages.... 43 9 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 1 25-29 years..... 9 30-34 years..... 17 12 35-39 years..... 25 19 13 40-44 years..... 17 14 3 9 8 45-49 years..... 22 13 14 17 50-54 years..... 14 3 9 8 55-59 years..... 5 2 3 3 60-64 years..... 4 3 1 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes HIV disease resulting in other bacterial infections (B20.1) All ages.... 1,009 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes HIV disease resulting in cytomegaloviral disease (B20.2) All ages.... 18 16 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 8 6 2 2 5 3 50-54 years..... 2 3 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis	eases,										sian or				
Tenth Revision, Secon		All rac	es		White			Black		American Indian 1/			Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
HIV disease resulting	in other vi	ral infec	tions (B20.	3)											
All ages	650	479	171	364	293	71	276	179	97	7	4	3	3	3	-
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	3	1	2	-	-	-	3	1	2	-	-	-	-	-	-
20-24 years	3	2	1	1	1	-	2	1	1	-	-	-	-	-	-
25-29 years	8	6	2	6	4	2	2	2	-	-	-	-	-	-	-
30-34 years	22	13	9	13	10	3	9	3	6	-	-	-	-	-	-
35-39 years	57	38	19	35	27	8	20	10	10	2	1	1	-	-	-
40-44 years	117	83	34	72	56	16	43	26	17	1	-	1	1	1	-
15-49 years	166	120	46	99	80	19	64	37	27	3	3	-	-	-	-
50-54 years	155	120	35	86	70	16	67	49	18	1	-	1	1	1	-
55-59 years	81	65	16	29	27	2	51	37	14	-	-	-	1	1	-
50-64 years	26	22	4	16	14	2	10	8	2	-	-	-	-	-	-
55-69 years	7	5	2	4	2	2	3	3	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	_	-	-	-
75-79 years	1	1	-	-	-	-	1	1	-	-	-	_	-	-	-
30-84 years	3	2	1	2	1	1	1	1	-	-	-	_	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
00-94 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
Age not stated	_	-	-	-	-	-	_	_	_	-	_	-	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes HIV disease resulting in candidiasis (B20.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 6 3 2 3 50-54 years..... 2 1 55-59 years..... 2 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes HIV disease resulting in other mycoses (B20.5) All ages.... 16 133 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 3 25-29 years..... 17 15 30-34 years..... 25 19 13 11 35-39 years..... 44 31 13 17 22 40-44 years..... 51 36 15 16 13 34 22 45-49 years..... 42 36 6 21 1 Q 21 17 17 50-54 years..... 21 20 3 17 1 55-59 years..... 12 11 5 1 5 60-64 years..... 10 3 6 65-69 vears..... 5 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes HIV disease resulting in Pneumocystis carinii pneumonia (B20.6) All ages.... 155 32 167 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2 1 1 1 1 20-24 years..... 3 3 3 25-29 years..... 21 16 11 30-34 years..... 27 20 7 13 12 14 8 35-39 years..... 62 45 17 33 25 15 10 45 39 40-44 years..... 82 59 23 35 19 16 24 45-49 years..... 64 41 23 29 34 16 1 Ω 2.7 2.4 5 50-54 years..... 51 43 8 21 16 3 3 55-59 years..... 26 18 8 16 11 10 3 60-64 years..... 14 9 5 6 5 8 65-69 vears..... 3 5 3 70-74 vears..... 4 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes HIV disease resulting in multiple infections (B20.7) All ages.... 1,174 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2. 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... QΩ 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes HIV disease resulting in other infectious and parasitic diseases (B20.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 2.4 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes HIV disease resulting in unspecified infectious or parasitic disease (B20.9) All ages.... 20 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 4 40-44 years..... 10 45-49 years..... 8 5 50-54 years..... 4 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Bot.h Both Bot.h Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Human immunodeficiency virus [HIV] disease resulting in malignant neoplasms (B21) All ages.... 366 267 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 3 1 2 2 1 1 15-19 years..... 1 1 1 1 20-24 years..... 10 3 6 25-29 years..... 2.7 16 11 12 30-34 years..... 44 40 31 29 12 10 35-39 years..... 106 90 16 67 60 39 30 99 89 49 40-44 years..... 173 140 33 72 23 88 82 45-49 years..... 157 127 30 6 67 43 24 2 58 50 9 50-54 years..... 125 108 17 64 1 55-59 years..... 94 76 18 43 40 50 35 15 1 1 60-64 years..... 50 3 31 31 19 16 3 30 19 16 10 7 65-69 vears..... 6 70-74 vears..... 10 9 3 75-79 years..... 5 5 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes HIV disease resulting in Kaposi's sarcoma (B21.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 12 12 12 35-39 years..... 13 13 40-44 years..... 12 10 45-49 years..... 2 50-54 years..... 3 55-59 years..... 3 2 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes HIV disease resulting in Burkitt's lymphoma (B21.1) All ages.... 13 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 10 45-49 years..... 2 2 50-54 years..... 4 3 55-59 years..... 2 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes HIV disease resulting in other types of non-Hodgkin's lymphoma (B21.2) All ages.... 192 123 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 1 1 20-24 years..... 3 25-29 years..... 30-34 years..... 19 15 11 35-39 years..... 47 42 5 29 18 79 57 40-44 years..... 93 34 26 48 45-49 years..... 76 64 12 2.8 20 29 5 2.4 2.0 50-54 years..... 34 8 55-59 years..... 28 25 3 17 17 11 60-64 years..... 15 14 11 11 4 1 65-69 vears..... 14 13 9 70-74 vears..... 1 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Bot.h Both and age Female Female Male Female Male Female Male Female sexes Male sexes Male sexes sexes sexes HIV disease resulting in other malignant neoplasms of lymphoid, hematopoietic, and related tissue (B21.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 2 35-39 years..... 12 9 40-44 years..... 10 10 45-49 years..... 12 11 9 8 50-54 years..... 11 55-59 years..... 3 2 1 60-64 years..... 2 2 65-69 vears..... 3 3 70-74 vears..... 2 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes HIV disease resulting in multiple malignant neoplasms (B21.7) All ages.... 19 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... a 45-49 years..... 50-54 years..... 55-59 years..... 4 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes HIV disease resulting in other malignant neoplasms (B21.8) All ages.... 10 142 101 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 4 35-39 years..... 21 16 9 12 9 40-44 years..... 33 21 12 15 18 10 45-49 years..... 39 26 13 1 2 17 2.0 8 12 45 39 50-54 years..... 51 6 11 11 33 6 55-59 years..... 45 35 10 16 13 28 21 60-64 years..... 13 26 24 2 11 11 15 65-69 vears..... 7 3 5 70-74 vears..... 6 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based																
on the International																
Classification of Dise	eases,									Asian or						
Tenth Revision, Second	ì	All rac	es		White			Black		American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both			Both			Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	remale	sexes	Male	Female	
HIV disease resulting	in unspecif	ied malig	mant neoplas	sm (B21.9)												
All ages	57	45	12	34	29	5	23	16	7	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	_	-	_	-	-	-	-	-	-	-	-	-	-	-	
15-19 years	-	_	-	_	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	1	_	1	_	-	-	1	-	1	-	-	-	-	-	-	
25-29 years	-	_	-	_	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	1	1	-	_	-	-	1	1	-	-	-	-	-	-	-	
35-39 years	2	_	2	1	_	1	1	_	1	_	_	_	_	_	_	
40-44 years	6	6	-	1	1	-	5	5	-	-	_	-	-	_	_	
45-49 years	12	11	1	7	7	-	5	4	1	-	-	-	-	-	-	
50-54 years	15	12	3	10	7	3	5	5	-	_	_	-	-	_	-	
55-59 years	9	6	3	5	5	-	4	1	3	-	-	-	-	-	-	
60-64 years	4	4	-	4	4	-	-	-	-	-	-	-	-	-	-	
65-69 years	5	3	2	4	3	1	1	-	1	-	-	-	-	-	-	
70-74 years	1	1	-	1	1	-	-	_	_	-	-	-	-	_	-	
75-79 years	1	1	-	1	1	-	-	_	_	-	-	-	-	_	-	
30-84 years	_	-	-	_	_	_	_	_	_	_	-	_	_	_	-	
85-89 years	_	-	-	_	_	_	_	_	_	_	-	_	_	_	-	
90-94 years	_	-	-	_	_	_	_	_	_	_	-	_	_	_	-	
95-99 years	-	-	-	_	-	-	-	_	_	-	-	-	-	_	-	
100 years & over.	_	_	_	_	_	_	-	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Bot.h Both Bot.h Both Bot.h and age Female Male Female Male Female sexes Male Female sexes Male Female sexes Male sexes sexes Human immunodeficiency virus [HIV] disease resulting in other specified diseases (B22) All ages.... 1,233 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... Q 1 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes HIV disease resulting in encephalopathy (B22.0) All ages.... 12 116 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 15 10 35-39 years..... 22 17 15 40-44 years..... 38 30 8 20 45-49 years..... 33 21 12 12 21 12 50-54 years..... 2.4 20 4 9 15 55-59 years..... 17 13 9 6 60-64 years..... 10 8 2 65-69 vears..... 9 5 70-74 vears..... 8 6 75-79 years..... 3 3 80-84 years..... 4 2 2 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes HIV disease resulting in wasting syndrome (B22.2) All ages.... 10 105 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 1 1 25-29 years..... 1 30-34 years..... 12 7 11 35-39 years..... 26 12 18 10 40-44 years..... 29 22 7 16 5 45-49 years..... 40 2.8 12 12 11 2.8 17 11 11 5 50-54 years..... 22 16 6 6 5 16 55-59 years..... 12 10 2 6 5 60-64 years..... 2 2 65-69 vears..... 4 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes HIV disease resulting in multiple diseases classified elsewhere (B22.7) All ages.... 443 312 131 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 17 11 6 25-29 years..... 26 17 17 9 30-34 years..... 58 50 8 31 29 2.7 21 35-39 years..... 111 95 16 57 48 37 11 80 74 58 30 40-44 years..... 171 133 38 88 45 94 80 30 45-49 years..... 189 144 14 92 62 131 96 35 65 54 11 42 2.4 50-54 years..... 66 55-59 years..... 93 72 21 38 33 5 55 39 16 60-64 years..... 46 40 6 22 20 23 19 26 23 15 13 10 9 65-69 vears..... 3 70-74 vears..... 10 8 5 5 75-79 years..... 2 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Bot.h Both Bot.h Both Bot.h and age Female Male Female Male Female sexes Male Female sexes Male Female sexes Male sexes sexes Human immunodeficiency virus [HIV] disease resulting in other conditions (B23) All ages.... 2,131 1,371 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Bot.h and age Female Male Female Male Female sexes Male Female sexes Male Female sexes Male sexes sexes HIV disease resulting in hematological and immunological abnormalities, not elsewhere classified (B23.2) All ages.... Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 1 20-24 years..... 1 25-29 years..... 15 10 11 30-34 years..... 15 6 9 8 2 35-39 years..... 41 22 19 15 26 13 13 23 17 40-44 years..... 60 37 34 17 45-49 years..... 50 36 14 14 12 2 34 12 10 19 13 50-54 years..... 39 29 15 19 6 55-59 years..... 22 14 8 9 8 13 6 60-64 years..... 8 4 3 3 3 65-69 vears..... 70-74 vears..... 4 75-79 years..... 5 3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes HIV disease resulting in other specified conditions (B23.8) All ages.... 1,866 1,336 1,213 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1.0 20-24 years..... 25-29 years..... 2.0 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis												ħ.a.	ian or		
Tenth Revision, Secon				White			Black			American Indian 1/ Pacific Islander 2/					
Edition, 2004)	Both	1111 140	-05	Both	,,,,,,		Both	224072		Both			Both	.10 1010	ilaci i
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Unspecified human imm	unodeficien	cy virus [HIV] diseas	e (B24)											
All ages	3,317	2,353	964	1,318	1,078	240	1,968	1,253	715	11	8	3	20	14	6
Under 1 year	2	1	1	1	1	-	1	-	1	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
10-14 years	3	1	2	1	-	1	2	1	1	-	-	-	-	-	-
5-19 years	8	5	3	2	2	_	6	3	3	-	-	-	-	-	-
20-24 years	51	27	24	12	9	3	39	18	21	-	-	-	-	-	-
25-29 years	119	61	58	31	21	10	87	40	47	-	-	-	1	-	1
30-34 years	216	128	88	71	47	24	144	80	64	1	1	-	-	-	-
35-39 years	456	296	160	173	133	40	278	160	118	2	1	1	3	2	1
10-44 years	706	493	213	287	239	48	409	247	162	5	3	2	5	4	1
45-49 years	697	518	179	281	247	34	414	269	145	-	-	-	2	2	-
50-54 years	475	366	109	205	165	40	267	199	68	1	1	-	2	1	1
55-59 years	288	220	68	125	104	21	161	114	47	-	-	-	2	2	-
0-64 years	137	112	25	60	50	10	74	60	14	2	2	-	1	-	1
55-69 years	75	58	17	36	31	5	37	25	12	-	-	-	2	2	-
70-74 years	50	42	8	18	17	1	31	24	7	-	-	-	1	1	-
75-79 years	22	16	6	8	6	2	13	10	3	_	-	-	1	-	1
30-84 years	10	7	3	6	5	1	4	2	2	_	-	-	_	-	-
35-89 years	2	2	-	1	1	-	1	1	-	_	-	-	_	-	-
90-94 years	-	-	-	_	_	-	_	-	-	_	-	-	_	-	-
95-99 years	-	-	-	_	_	-	_	-	-	_	-	-	_	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Disc													ian or		
Tenth Revision, Second		All races		_	White		_	Black		American Indian 1/			Pacific Islander 2		
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	omala	Both sexes	Mala	Female
	SEACS	Maie	remaie		mare	remare	SCACS	maie	remare	SCACS	Male r		BEACS		remare
Cytomegaloviral diseas	se (B25)														
All ages	118	58	60	89	46	43	22	8	14	_	_	_	7	4	3
Under 1 year	16	9	7	11	7	4	4	1	3	-	-	-	1	1	-
1-4 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
5-9 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
l0-14 years	6	2	4	5	2	3	1	_	1	-	-	-	-	-	-
.5-19 years	9	2	7	6	1	5	3	1	2	-	-	-	-	-	-
20-24 years	2	1	1	1	1	-	1	_	1	-	-	-	-	-	-
25-29 years	2	1	1	2	1	1	-	_	-	-	-	-	-	-	-
30-34 years	1	-	1	-	-	-	1	_	1	-	-	-	-	-	-
35-39 years	4	3	1	3	3	-	1	_	1	-	-	-	-	-	-
10-44 years	4	3	1	1	1	-	2	2	-	-	-	-	1	-	1
15-49 years	6	2	4	4	2	2	2	_	2	-	-	-	-	-	-
50-54 years	8	4	4	7	3	4	1	1	-	-	-	-	-	-	-
55-59 years	8	2	6	6	2	4	2	_	2	-	-	-	-	-	-
50-64 years	10	7	3	5	3	2	2	2	-	-	-	-	3	2	1
65-69 years	9	3	6	9	3	6	-	_	-	-	-	-	-	-	-
70-74 years	9	7	2	9	7	2	-	_	-	-	-	-	-	-	-
5-79 years	7	6	1	7	6	1	-	_	-	-	-	-	-	-	-
0-84 years	10	3	7	7	2	5	1	_	1	-	-	-	2	1	1
5-89 years	2	-	2	2	-	2	-	_	-	-	-	-	-	-	-
00-94 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Cytomegaloviral pneumonitis (B25.0) All ages..... 23 13 Under 1 year.... 5 2 3 1 1 3 1-4 years..... 5-9 years..... 2 10-14 years..... 1 15-19 years..... 3 2 1 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 2 2 5 2 50-54 years..... 3 3 1 55-59 years..... 5 2 3 60-64 years..... 8 6 2 65-69 vears..... 4 70-74 vears..... 5 5 75-79 years..... 2 80-84 years..... 3 3 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second All races									sian or						
		All rac	es		White			Black			rican Ind	ian 1/	Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Cytomegaloviral hepat	itis (B25.1))													
All ages	4	_	4	4	_	4	-	_	_	-	_	-	-	_	-
Under 1 year	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
30-34 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
35-39 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
40-44 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
45-49 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
50-54 years	_	-	-	-	-	-	-	-	-	-	-	-	_	-	-
55-59 years	_	-	-	-	-	-	-	-	-	-	-	-	_	-	-
60-64 years	_	-	-	-	-	-	-	-	-	-	-	-	_	-	-
65-69 years	1	-	1	1	-	1	-	-	-	-	-	-	_	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other cytomegaloviral diseases (B25.8) All ages.... Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Cytomegaloviral disease, unspecified (B25.9) All ages.... 40 19 Under 1 year.... 9 6 3 8 5 3 1-4 years..... 1 3 2 5-9 years..... 10-14 years..... 3 1 2 15-19 years..... 2 2 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 2 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 2 1 1 65-69 vears..... 3 70-74 vears..... 3 75-79 years..... 4 80-84 years..... 6 3 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Mumps (B26) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Mumps without complication (B26.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Infectious mononucleosis (B27) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 2 2 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Gamma-herpes-viral mononucleosis (B27.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 2 2 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis	seases,											As	ian or		
Tenth Revision, Secon	nd	All rac	es		White			Black		American Indian 1/			Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Infectious mononucleo	osis, unspeci	ified (B27	.9)												
All ages	3	1	2	3	1	2	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	_	_	-	_	_	_	_	_	-	-	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other viral diseases, not elsewhere classified (B33) All ages.... Under 1 year.... 2 2 1 1 1 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 3 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 2 2 6 50-54 years..... 3 3 55-59 years..... 5 2 3 60-64 years..... 4 1 3 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 2 80-84 years..... 3 85-89 years..... 90-94 years..... 3 2 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Disc Tenth Revision, Second		ses, All races			White Black							Asian or American Indian 1/ Pacific Islander 2/						
Tenth Revision, Second Edition, 2004)	Both	All rac	es	Both	willte		Both	Black		Both	ican indi	lan 1/	Both	IC ISIA	nder 2/			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female			
Viral carditis (B33.2))																	
All ages	43	21	22	35	18	17	4	2	2	_	_	_	4	1	3			
Jnder 1 year	2	-	2	1	-	1	_	-	-	-	-	-	1	-	1			
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
5-9 years	1	-	1	1	-	1	_	-	-	-	-	-	-	-	-			
l0-14 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-			
5-19 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-			
0-24 years	3	2	1	2	2	-	1	-	1	-	-	-	-	-	-			
25-29 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-			
30-34 years	1	1	-	-	-	-	-	-	-	-	-	-	1	1	-			
35-39 years	4	4	-	3	3	-	1	1	-	-	-	-	-	-	-			
10-44 years	1	1	-	1	1	-	_	-	-	-	-	-	-	-	-			
15-49 years	3	1	2	2	1	1	-	-	-	-	-	-	1	-	1			
50-54 years	6	3	3	5	3	2	1	-	1	-	-	-	-	-	-			
55-59 years	5	2	3	4	1	3	1	1	-	-	-	-	-	-	-			
0-64 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-			
5-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-			
70-74 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-			
5-79 years	2	-	2	1	-	1	-	-	-	-	-	-	1	-	1			
0-84 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-			
5-89 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-			
0-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-			
95-99 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-			
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_			

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Retrovirus infections, not elsewhere classified (B33.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hantavirus (cardio)-pulmonary syndrome [HPS][HCPS] (B33.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified viral diseases (B33.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Viral infection of unspecified site (B34) All ages.... 136 221 113 108 52 Under 1 year.... 70 41 29 44 2.6 18 22 11 11 1-4 years..... 36 21 15 26 16 10 7 4 3 5 5 5-9 years..... 4 1 10-14 years..... 5 1 15-19 years..... 10 1.0 6 20-24 years..... 2 2 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 8 4 5 2 3 7 50-54 years..... 5 2 3 55-59 years..... 14 12 10 60-64 years..... 5 3 2 3 2 65-69 vears..... 4 4 70-74 vears..... 15 8 7 75-79 years..... 9 5 8 80-84 years..... 24 12 12 22 11 11 22 85-89 years..... 24 8 16 8 14 1 1 90-94 years..... 16 10 16 10 6 6 95-99 years..... 5 5 5 5 100 years & over. 2 1 2 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases,						Asian or American Indian 1/ Pacific Islander 2/									
Tenth Revision, Secon Edition, 2004)	a Both	All races		White			Black			Amer Both	rican Indi	lan I/	Pacific Islander 2/ Both		
and age	sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Adenovirus infection,	unspecified	d (B34.0)													
All ages	15	12	3	11	9	2	3	2	1	-	-	-	1	1	-
Under 1 year	8	6	2	6	5	1	2	1	1	_	-	-	-	-	-
1-4 years	4	3	1	4	3	1	-	-	-	_	-	-	-	-	-
5-9 years	1	1	-	-	-	-	1	1	-	_	-	-	-	-	-
10-14 years	1	1	-	-	-	-	-	-	-	_	-	-	1	1	-
15-19 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	_	-	-	-	_	_	_	_	_	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Enterovirus infection, unspecified (B34.1) All ages.... Under 1 year.... 6 2 3 3 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Parvovirus infection, unspecified (B34.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other viral infections of unspecified site (B34.8) All ages.... 16 Under 1 year.... 24 13 11 17 10 7 1-4 years..... 12 5 10 3 1 1 5-9 years..... 1 10-14 years..... 2 1 1 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 90-94 years..... 2 1 1 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Viral infection, unspecified (B34.9) All ages.... 102 102 161 38 17 Under 1 year.... 32 2.0 12 18 11 12 5 2 2 1-4 years..... 18 10 8 10 5 5 7 3 3 3 5-9 years..... 3 10-14 years..... 4 1 3 15-19 years..... 3 20-24 years..... 1 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 8 4 4 5 3 6 50-54 years..... 2 2 2 55-59 years..... 12 9 3 10 3 60-64 years..... 5 3 2 3 2 65-69 vears..... 3 3 70-74 vears..... 13 75-79 years..... 9 5 8 80-84 years..... 20 9 11 18 10 85-89 years..... 23 16 21 14 1 1 90-94 years..... 14 5 9 14 9 5 95-99 years..... 4 4 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Candidiasis (B37) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... Ω 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Candidal stomatitis (B37.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pulmonary candidiasis (B37.1) All ages.... 12 12 Under 1 year.... 1 1 1-4 years..... 1 1 1 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 3 75-79 years..... 2 80-84 years..... 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Candidiasis of skin and nail (B37.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Candidiasis of other urogenital sites (B37.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 4 75-79 years..... 2 80-84 years..... 3 2 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis	seases,	711										Asian or						
Tenth Revision, Secon		All rac	All races		White			Black			American Indian 1/			Pacific Islander 2/				
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male Fe	emale	Both sexes	Male	Female			
Candidal meningitis (B37.5)																	
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_			
Under 1 year	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-			
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
.0-14 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
20-24 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
25-29 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-			
35-39 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
10-44 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
15-49 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
55-69 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-			
70-74 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
5-79 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
0-84 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
5-89 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
0-94 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
95-99 years	_	_	-	_	_	-	_	-	_	_	-	-	_	-	-			
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_			

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Candidal endocarditis (B37.6) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Candidal septicemia (B37.7) All ages.... 177 93 134 67 38 Under 1 year.... 12 6 6 5 3 2 6 3 3 1 1-4 years..... 3 1 2 3 2 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 6 4 2 2 1 7 3 50-54 years..... 2 5 2 3 55-59 years..... 13 9 11 4 60-64 years..... 5 3 2 4 65-69 vears..... 17 8 70-74 vears..... 25 16 6 75-79 years..... 2.2 10 12 20 12 80-84 years..... 24 12 12 19 10 5 22 3 85-89 years..... 9 13 19 11 90-94 years..... 5 4 1 5 1 95-99 years..... 1 1 1 1 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis	eases,												ian or		
Tenth Revision, Secon					White			Black			rican Ind	ian 1/	Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Candidiasis of other	sites (B37.8	3)													
All ages	33	17	16	28	14	14	4	2	2	-	-	-	1	1	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	2	1	1	-	-	-	2	1	1	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
50-54 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-
55-59 years	2	1	1	1	-	1	-	-	-	-	-	-	1	1	-
60-64 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
65-69 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
70-74 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
75-79 years	6	2	4	5	2	3	1	-	1	-	-	-	-	-	-
80-84 years	6	3	3	6	3	3	-	-	-	-	-	-	-	-	-
85-89 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Candidiasis, unspecified (B37.9) All ages..... 11 Under 1 year.... 2 2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 3 75-79 years..... 3 80-84 years..... 6 85-89 years..... 90-94 years..... 3 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Coccidioidomycosis (B38) All ages.... 110 79 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 1 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 16 10 45-49 years..... 4 3 1 2 50-54 years..... ρ 3 3 11 55-59 years..... 7 2 3 5 60-64 years..... 11 8 3 65-69 vears..... 7 6 70-74 vears..... 10 6 75-79 years..... 11 9 3 80-84 years..... 7 85-89 years..... 9 2 90-94 years..... 2 1 1 95-99 years..... 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acute pulmonary coccidioidomycosis (B38.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Chronic pulmonary coccidioidomycosis (B38.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Pulmonary coccidioidomycosis, unspecified (B38.2) All ages.... 30 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 3 3 60-64 years..... 2 1 1 65-69 vears..... 2 70-74 vears..... 6 75-79 years..... 80-84 years..... 1 6 85-89 years..... 90-94 years..... 1 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Coccidioidomycosis meningitis (B38.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 3 40-44 years..... 45-49 years..... 2 50-54 years..... 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disseminated coccidioidomycosis (B38.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 3 1 2 60-64 years..... 5 3 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Coccidioidomycosis, unspecified (B38.9) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Histoplasmosis (B39) All ages.... 19 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 2 50-54 years..... 3 2 55-59 years..... 60-64 years..... 3 65-69 vears..... 3 70-74 vears..... 8 5 75-79 years..... 80-84 years..... 4 2 85-89 years..... 2 2 90-94 years..... 95-99 years..... 1 1 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Chronic pulmonary histoplasmosis capsulati (B39.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Pulmonary histoplasmosis capsulati, unspecified (B39.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based																
on the International Classification of Dise	22505										λα	ian or				
Tenth Revision, Second		ases, All races			White			Black		Asian or American Indian 1/ Pacific Islander 2						
Edition, 2004)	Both	1111 100		Both	***************************************		Both	Diaon		Both	,		Both	ilidel 2,		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female	
Histoplasmosis, unspec	cified (B39.	9)														
All ages	25	17	8	21	13	8	3	3	_	_	-	-	1	1	-	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-	
50-54 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	5	4	1	5	4	1	-	-	-	-	-	-	-	-	-	
65-69 years	3	3	-	3	3	-	-	-	-	-	-	-	-	-	-	
70-74 years	6	3	3	6	3	3	-	-	-	-	-	-	-	-	-	
'5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
80-84 years	3	1	2	2	-	2	-	-	-	-	-	-	1	1	-	
35-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	_	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Blastomycosis (B40) All ages.... 19 9 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 3 2 1 55-59 years..... 4 4 60-64 years..... 2 1 65-69 vears..... 6 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Pulmonary blastomycosis, unspecified (B40.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Disseminated blastomycosis (B40.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Blastomycosis, unspecified (B40.9) All ages..... 10 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Sporotrichosis (B42) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	seases,	711										Asian or American Indian 1/ Pacific Islander 2/						
		All races		Both	White		Black			Both	rican indi	an 1/						
Edition, 2004) and age	Both sexes	Male	Female	sexes	Male	Female	Both sexes	Male	Female	sexes	Male F	emale	Both sexes	Male	Female			
Disseminated sporotri	chosis (B42.	.7)																
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_			
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
5-9 years	_	-	-	_	-	-	-	-	_	-	-	-	-	-	-			
.0-14 years	_	-	-	_	-	-	-	-	_	-	-	-	-	-	-			
.5-19 years	_	-	-	_	-	-	-	-	_	-	-	-	-	-	-			
0-24 years	_	-	-	_	-	-	-	-	_	-	-	-	-	-	-			
25-29 years	_	-	-	_	-	-	-	-	_	-	-	-	-	-	-			
0-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-			
35-39 years	_	-	-	_	-	-	-	-	_	-	-	-	-	-	-			
10-44 years	_	-	-	_	-	-	-	-	_	-	-	-	-	-	-			
15-49 years	_	-	-	-	_	-	-	_	_	-	_	_	-	-	-			
0-54 years	_	-	-	-	_	-	-	_	_	-	_	_	-	-	-			
55-59 years	_	-	-	-	_	-	-	_	_	-	_	_	-	-	-			
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
55-69 years	_	-	-	-	_	-	-	_	_	-	_	_	-	-	-			
70-74 years	-	-	-	-	-	-	-	_	_	_	-	-	-	-	-			
5-79 years	1	1	-	1	1	-	-	_	_	_	-	-	-	-	-			
0-84 years	-	-	-	_	-	-	-	_	_	_	-	-	-	-	-			
5-89 years	-	-	-	_	-	-	-	_	_	_	-	-	-	-	-			
0-94 years	-	-	-	_	-	-	-	_	_	_	-	-	-	-	-			
95-99 years	_	_	-	_	_	-	_	_	-	-	-	_	-	_	-			
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_			

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other forms of sporotrichosis (B42.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Aspergillosis (B44) All ages.... 132 106 42 10 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 7 40-44 years..... 45-49 years..... 11 6 5 50-54 years..... 11 ρ 3 1 0 55-59 years..... 18 14 3 11 1 60-64 years..... 14 10 11 1 65-69 vears..... 8 10 3 14 6 70-74 vears..... 22 16 6 20 75-79 years..... 12 11 5 80-84 years..... 7 5 1 1 3 85-89 years..... 2 3 1 90-94 years..... 1 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Invasive pulmonary aspergillosis (B44.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other pulmonary aspergillosis (B44.1) All ages..... 32 59 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 2 1 1 2 6 50-54 years..... 2 5 1 55-59 years..... 12 10 5 60-64 years..... 6 2 6 65-69 vears..... 10 70-74 vears..... 10 75-79 years..... 9 80-84 years..... 6 2 1 1 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Disseminated aspergillosis (B44.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other forms of aspergillosis (B44.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Aspergillosis, unspecified (B44.9) All ages..... 13 30 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 8 3 50-54 years..... 3 55-59 years..... 2 2 60-64 years..... 6 5 1 65-69 vears..... 3 70-74 vears..... 8 75-79 years..... 2 80-84 years..... 85-89 years..... 1 1 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Classification of Dise Tenth Revision, Second		All rac	100		White			Black		λmer	ican Indi		sian or	ic Tela	nder 2/
Edition, 2004)	Both	AII IAC	.05	Both	WIIICC		Both	DIACK		Both	ican ina.	Idii I/	Both	.10 1514	inder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
Cryptococcosis (B45)															
All ages	100	68	32	66	46	20	29	19	10	2	2	-	3	1	2
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
30-34 years	4	4	-	2	2	-	1	1	-	1	1	-	-	-	-
35-39 years	7	6	1	3	3	-	4	3	1	-	-	-	-	-	-
40-44 years	8	6	2	4	3	1	3	2	1	1	1	-	-	-	-
45-49 years	8	5	3	6	4	2	2	1	1	-	-	-	-	-	-
50-54 years	15	11	4	10	7	3	4	4	-	-	-	-	1	-	1
55-59 years	9	5	4	5	3	2	4	2	2	-	-	-	-	-	-
60-64 years	7	3	4	2	1	1	4	2	2	-	-	-	1	-	1
65-69 years	8	7	1	7	6	1	1	1	-	-	-	-	-	-	-
70-74 years	9	5	4	6	3	3	2	1	1	-	-	-	1	1	-
75-79 years	7	5	2	6	4	2	1	1	-	-	-	-	-	-	-
80-84 years	9	4	5	9	4	5	-	-	-	-	-	-	-	-	-
85-89 years	6	5	1	5	5	-	1	-	1	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dise													sian or		
Tenth Revision, Second		All rac	es		White			Black			rican Indi	an 1/		ic Isla	nder 2/
Edition, 2004) and age	Both	Male	Female	Both sexes	Male	Female	Both	Male	Female	Both	Male F	omala	Both sexes	Molo	Female
	sexes	мате	remare		Maie	remaie	sexes	мате	remare	sexes	Maie r		sexes	Male	remare
Pulmonary cryptococcos	sis (B45.0)														
All ages	11	7	4	8	5	3	3	2	1	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
45-49 years	2	1	1	1	1	-	1	-	1	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	-	-	-	-	-	-	_	-	-	-	-	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Cerebral cryptococcosis (B45.1) All ages..... 20 12 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 4 1 50-54 years..... 13 1.0 3 55-59 years..... 6 2 4 60-64 years..... 4 2 2 65-69 vears..... 6 70-74 vears..... 5 75-79 years..... 80-84 years..... 4 3 2 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Disseminated cryptococcosis (B45.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cryptococcosis, unspecified (B45.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Zygomycosis (B46) All ages.... 12 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 1 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pulmonary mucormycosis (B46.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Rhinocerebral mucormycosis (B46.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Gastrointestinal mucormycosis (B46.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis								-1 1					sian or		1 0/
Tenth Revision, Secon		All rac	es		White			Black			rican Ind	lian 1/		ic Isla	nder 2/
Edition, 2004)	Both		- 1	Both			Both		- 1	Both		_ ,	Both		_ ,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	remale	sexes		Female
Disseminated mucormyc	osis (B46.4)	ı													
All ages	1	1	_	1	1	-	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	_	-	_	-	-	-	-	-	-	-	-	_	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	_	-	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Mucormycosis, unspecified (B46.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other zygomycoses (B46.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis		711			White			D]l-		2	rican India		ian or	::- T-1-	
Tenth Revision, Secon		All rac	ces	Dath	wnite		Dath	Black			rican India	an 1/		ic isia	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male Fe	emale	Both sexes	Male	Female
Mycetoma (B47)															
All ages	4	2	2	3	2	1	1	_	1	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	3	1	2	2	1	1	1	-	1	-	-	-	-	-	-
50-54 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Mycetoma, unspecified (B47.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Unspecified mycosis (B49) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... Ω 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 2.4 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Plasmodium falciparum malaria (B50) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Plasmodium falciparum malaria, unspecified (B50.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Unspecified malaria (B54) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Chagas' disease (B57) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis													ian or		
Tenth Revision, Second		All rac	es		White			Black			rican Indi	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Female	sexes	Male	Female
Chagas' disease (chro	nic) with he	eart invol	vement (B57.	2)											
All ages	3	2	1	2	1	1	1	1	-	_	-	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	_	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	-	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise		211			rale de la			Dl. de					sian or		0 /
Tenth Revision, Second		All rac	es	Both	White		Dath	Black			rican Indi	ian I/		ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	Tomalo	Both sexes	Mala	Female
Toxoplasmosis (B58)															
All ages	13	8	5	9	6	3	4	2	2	_	-	-	_	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
35-39 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
40-44 years	3	-	3	2	-	2	1	-	1	-	-	-	-	-	-
45-49 years	3	3	-	2	2	-	1	1	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Toxoplasma meningoencephalitis (B58.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Toxoplasmosis with other organ involvement (B58.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Toxoplasmosis, unspecified (B58.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Pneumocystosis (B59) All ages.... 148 104 43 38 18 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 1 25-29 years..... 30-34 years..... 3 35-39 years..... 11 40-44 years..... 19 13 45-49 years..... 16 1.0 6 1 0 6 8 50-54 years..... 16 1.0 ρ 55-59 years..... 11 7 9 4 5 60-64 years..... 21 12 9 19 11 65-69 vears..... 10 8 2 70-74 vears..... 15 8 75-79 years..... 8 1 80-84 years..... 4 2 2 85-89 years..... 4 2 1 90-94 years..... 1 1 1 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis													sian or		
Tenth Revision, Secon		All rac	es	_	White		_	Black			rican Ind	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female 	sexes	Male	Female
Other protozoal disea	ses, not els	sewhere cl	assified (Be	50)											
All ages	6	4	2	6	4	2	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
10-14 years	1	1	-	1	1	-	-	-	-	-	_	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	_
25-29 years	-	-	-	-	-	-	-	_	-	-	_	-	_	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
50-54 years	1	-	1	1	-	1	-	_	-	-	_	-	_	-	-
55-59 years	-	-	-	-	-	-	-	_	-	-	_	-	_	-	-
60-64 years	-	-	-	-	-	-	-	_	-	-	_	-	_	-	-
65-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
85-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International															
Classification of Dise		211			meta di sala			D11-		3			ian or		/
Tenth Revision, Second Edition, 2004)	Both	All rac	ces	Both	White		Both	Black		Amer Both	ican Indi	an 1/	Pacii Both	ic ista	nder 2/
		Male	Female		Male	Female		Male	Female		Male F	1 -		M-1-	Female
and age	sexes	мате	remaie	sexes	мате	remaie	sexes	мате	remaie	sexes	Male F		sexes		remare
Babesiosis (B60.0)															
All ages	5	3	2	5	3	2	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
75-79 years	_	_	-	_	_	-	-	-	-	_	-	-	-	-	-
30-84 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
35-89 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
95-99 years	_	_	-	_	_	-	-	-	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Negleriasis (B60.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cysticercosis (B69) All ages..... 12 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Cysticercosis of central nervous system (B69.0) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Trichinellosis (B75) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Strongyloidiasis (B78) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disseminated strongyloidiasis (B78.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Strongyloidiasis, unspecified (B78.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other intestinal helminthiases, not elsewhere classified (B81) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Intestinal capillariasis (B81.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dis	eases,											As	ian or		
Tenth Revision, Secon	.d	All rac	es		White			Black		Amei	rican Indi	an 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Trichostrongyliasis (B81.2)														
All ages	1	1	_	_	_	_	1	1	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	_	-	-	-	-	_	-	_	-	-	-	-	-	-	-
10-44 years	-	-	-	_	-	_	-	-	_	_	-	-	-	-	-
45-49 years	_	-	-	-	-	_	-	_	-	-	-	-	-	-	-
50-54 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	_	-	-	-	-	_	-	_	-	-	-	-	-	-	-
75-79 years	_	-	-	-	-	_	-	_	-	-	-	-	-	-	-
30-84 years	_	-	-	-	-	_	-	_	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	_	-	-	-	-	_	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	_	-	_	-	-	_	_	-	-	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis		All rac			White			Black		3	rican Indi		ian or	::- T-1-	nder 2/
Tenth Revision, Secon Edition, 2004)	Both	All rac	es	Both	willte		Both	Black		Both	rican indi	an 1/	Both	ic isia	maer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Mixed intestinal helm	inthiases (E	381.4)													
All ages	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
Under 1 year	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	-	_	-	-
5-9 years	_	-	-	_	_	_	-	-	_	_	_	-	_	_	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	_	-	-	_	-	-
35-39 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	_
40-44 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	_
45-49 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	_
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other helminthiases (B83) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Helminthiasis, unspecified (B83.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Scabies (B86) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc	eases,											As	ian or		
Tenth Revision, Second		All rac	es		White			Black		Amer	ican Indi			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Myiasis (B87)															
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-		-	-	-	-	_	-	_	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-		-	-	-	-	_	-	_	_	-	-	-	-
10-14 years	-	-		-	-	-	-	_	-	_	_	-	-	-	-
15-19 years	-	-		-	-	-	-	_	-	_	_	-	-	-	-
20-24 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	_	_	_	-	-	-	-	-	-	-
15-49 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-		-	-	-	-	_	-	_	_	-	-	-	-
50-64 years	-	-		-	-	-	-	_	-	_	_	-	-	-	-
55-69 years	-	-		-	-	-	-	_	-	_	_	-	-	-	-
70-74 years	-	-	-	_	-	_	_	-	-	-	-	-	-	-	-
'5-79 years	_	_	-	_	_	_	_	-	-	_	-	-	-	-	-
0-84 years	-	-	-	_	-	_	_	_	_	-	-	-	-	-	-
35-89 years	-	-	-	_	-	_	_	_	_	-	-	-	-	-	-
0-94 years	1	_	1	1	_	1	_	-	-	_	-	-	-	-	-
95-99 years	_	_	-	_	_	_	_	-	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	100		White			Black		Δme	rican Indi		ian or	ic Isla	nder 2/
Edition, 2004)	Both	1111 140	.00	Both	WILLCC		Both	Diach		Both	ican inai	an i	Both	.10 1010	naci 27
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Myiasis, unspecified	(B87.9)														
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	_	-	-	_	-	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Unspecified parasitic disease (B89) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Sequelae of tuberculosis (B90) All ages..... 21 21 28 14 1 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 8 80-84 years..... 8 2 2 85-89 years..... 12 5 10 90-94 years..... 5 2 3 3 5 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Sequelae of respiratory and unspecified tuberculosis (B90.9) All ages..... 14 14 1 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 1 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 8 80-84 years..... 8 2 2 85-89 years..... 12 5 10 90-94 years..... 5 2 3 3 5 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Sequelae of poliomyelitis (B91) All ages..... 159 185 338 157 181 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 1 1 2 1 6 50-54 years..... 5 5 1 6 1 55-59 years..... 18 7 18 11 11 60-64 years..... 36 15 21 36 15 21 65-69 vears..... 44 23 21 43 23 20 70-74 vears..... 39 22 17 37 16 75-79 years..... 69 34 35 34 35 80-84 years..... 63 26 37 61 25 36 28 85-89 years..... 40 11 29 39 11 1 90-94 years..... 23 11 12 23 11 12 95-99 years..... 3 3 3 3 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis	eases,												ian or		
Tenth Revision, Secon		All rac	es		White			Black			ican In	dian 1/		ic Isla	ınder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Sequelae of other and	unspecified	l infectio	us and paras	sitic diseas	es (B94)										
All ages	903	475	428	763	400	363	112	58	54	4	3	1	24	14	10
Under 1 year															
1-4 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-9 years	2	1	1	1	-	1	-	-	-	-	-	-	1	1	-
10-14 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
l5-19 years	3	1	2	2	1	1	1	-	1	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	5	3	2	2	2	-	2	1	1	-	-	-	1	-	1
30-34 years	10	4	6	7	2	5	3	2	1	-	-	-	-	-	-
35-39 years	8	4	4	4	2	2	3	1	2	1	1	-	_	-	-
0-44 years	25	16	9	20	12	8	5	4	1	-	-	-	-	-	-
15-49 years	43	25	18	36	21	15	6	3	3	1	1	-	-	-	-
0-54 years	78	57	21	64	46	18	12	10	2	1	-	1	1	1	-
55-59 years	82	52	30	67	43	24	12	6	6	1	1	-	2	2	-
0-64 years	72	38	34	57	32	25	11	4	7	-	-	-	4	2	2
55-69 years	88	46	42	75	39	36	10	6	4	-	-	-	3	1	2
70-74 years	103	59	44	89	52	37	12	6	6	-	-	-	2	1	1
5-79 years	120	56	64	102	48	54	14	7	7	-	-	-	4	1	3
0-84 years	123	61	62	109	53	56	9	4	5	_	_	_	5	4	1
5-89 years	77	31	46	71	28	43	6	3	3	_	_	_	_	-	-
00-94 years	38	13	25	34	12	22	3	-	3	-	-	-	1	1	-
95-99 years	21	4	17	18	3	15	3	1	2	_	_	_	_	-	-
100 years & over.	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Sequelae of viral encephalitis (B94.1) All ages.... 4 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Sequelae of viral hepatitis (B94.2) All ages..... 179 129 50 147 107 40 18 13 1 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 8 3 45-49 years..... 23 14 1 Q 11 49 9 41 34 50-54 years..... 40 6 1 55-59 years..... 36 27 9 29 22 5 1 1 60-64 years..... 15 8 12 2 1 65-69 vears..... 11 70-74 vears..... 13 8 5 75-79 years..... 11 9 80-84 years..... 9 8 1 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Sequelae of other specified infectious and parasitic diseases (B94.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 a 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 2.8 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Sequelae of unspecified infectious or parasitic disease (B94.9) All ages.... 1 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other and unspecified infectious diseases (B99) All ages.... 163 237 141 Under 1 year..... 5 4 1 3 3 2 1 1 1-4 years..... 4 4 3 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 3 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 8 6 2 5 3 7 50-54 years..... 4 3 2 55-59 years..... 16 8 13 3 8 60-64 years..... 17 13 15 3 12 65-69 years..... 13 11 6 7 6 70-74 years..... 26 11 15 19 13 75-79 years..... 32 11 2.1 31 2.0 11 80-84 years..... 43 22 21 37 19 18 6 35 85-89 years..... 40 16 2.4 13 22 5 3 90-94 years..... 35 1.0 25 32 24 3 1 8 2 95-99 years..... 7 18 11 16 10 2 1 1 6 100 years & over. 2 1 1 1 4 4 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Malignant neoplasm of lip (C00) All ages..... 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 3 50-54 years..... 3 55-59 years..... 2 2 60-64 years..... 3 3 65-69 vears..... 2 70-74 vears..... 6 6 75-79 years..... 9 9 80-84 years..... 3 85-89 years..... 9 9 90-94 years..... 10 5 1.0 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis													ian or		
Tenth Revision, Secon	d	All rac	es		White			Black			ican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
External upper lip (C	00.0)														
All ages	3	2	1	3	2	1	_	_	-	_	-	-	_	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
l5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	-	-	-	-	_	_	_	_	-	-	-	-	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis													sian or		
Tenth Revision, Secon		All rac	es		White			Black			rican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
External lower lip (C	00.1)														
All ages	13	12	1	13	12	1	-	_	-	-	-	-	-	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
70-74 years	5	5	-	5	5	-	-	-	-	-	-	-	-	-	-
75-79 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis	eases,	-11						-1 1					sian or		1 0/
Tenth Revision, Secon		All rac	es		White			Black			rican Indi	an 1/		ıc Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
Lip, unspecified, inn	er aspect (C	200.5)													
All ages	1	1	-	1	1	-	-	_	-	-	-	-	-	_	-
Under 1 year	-	_	-	-	-	-	-	-	-	_	-	-	-	-	-
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
0-54 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
5-59 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
5-79 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
0-84 years	_	_	-	-	_	_	_	-	-	_	_	-	_	_	-
5-89 years	-	_	-	-	-	-	-	-	-	_	_	-	-	_	-
0-94 years	-	_	-	-	-	-	-	-	-	_	_	-	-	_	-
5-99 years	-	_	-	-	-	-	-	-	-	_	_	-	-	_	-
.00 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lip, unspecified (C00.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 2 2 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 8 90-94 years..... 3 4 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis	02000											7.0	ian or		
Tenth Revision, Second		All rac	es		White			Black		Amer	ican Indi			ic Isla	nder 2/
Edition, 2004)	Both	1111 100		Both	,,,,,,		Both	Didon		Both	10011 11101		Both	.10 1010	11401 27
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Malignant neoplasm of	base of tor	ngue (C01)													
All ages	144	106	38	130	96	34	12	8	4	_	_	_	2	2	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
40-44 years	7	5	2	7	5	2	-	-	-	-	-	-	-	-	-
45-49 years	13	10	3	11	9	2	2	1	1	-	-	-	-	-	-
50-54 years	18	16	2	16	14	2	1	1	-	-	-	-	1	1	-
55-59 years	30	27	3	27	24	3	3	3	-	-	-	-	-	-	-
60-64 years	15	11	4	11	8	3	3	2	1	-	-	-	1	1	-
65-69 years	14	10	4	14	10	4	-	-	-	-	-	-	-	-	-
70-74 years	19	13	6	19	13	6	-	-	-	-	-	-	-	-	-
75-79 years	14	7	7	12	7	5	2	-	2	-	-	-	-	-	-
80-84 years	9	5	4	8	4	4	1	1	-	-	-	-	-	-	-
85-89 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
90-94 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis															
Tenth Revision, Secon		All rac	0.5		White			Black		7 more	ican Indi		ian or	iia Tala	nder 2/
Edition, 2004)	Both	AII IAC	es	Both	WIIICE		Both	DIACK		Both	Ican Ind	Ian I/	Both	IC ISIA	ilder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
Malignant neoplasm of	other and	unspecifie	d parts of t	congue (C02)										
All ages	1,762	1,147	615	1,541	1,001	540	171	117	54	7	4	3	43	25	18
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
20-24 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
25-29 years	7	6	1	7	6	1	-	-	-	-	-	-	-	-	-
30-34 years	7	4	3	6	3	3	1	1	-	-	-	-	-	-	-
35-39 years	30	19	11	25	14	11	3	3	-	-	-	-	2	2	-
40-44 years	47	31	16	40	25	15	6	5	1	-	-	-	1	1	-
45-49 years	105	84	21	89	73	16	13	9	4	1	-	1	2	2	-
50-54 years	166	130	36	139	113	26	22	13	9	1	1	-	4	3	1
55-59 years	207	156	51	174	136	38	27	16	11	1	1	-	5	3	2
60-64 years	202	154	48	178	139	39	19	12	7	-	-	-	5	3	2
65-69 years	211	153	58	169	122	47	31	26	5	3	1	2	8	4	4
70-74 years	211	131	80	188	113	75	20	16	4	-	-	-	3	2	1
75-79 years	200	122	78	178	109	69	16	11	5	1	1	-	5	1	4
80-84 years	181	91	90	166	84	82	9	4	5	-	-	-	6	3	3
85-89 years	116	42	74	114	41	73	2	1	1	-	-	-	-	-	-
90-94 years	50	17	33	47	16	31	1	-	1	-	-	-	2	1	1
95-99 years	18	7	11	17	7	10	1	-	1	-	-	-	-	-	-
100 years & over.	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Border of tongue (C02.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lingual tonsil (C02.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Tongue, unspecified (C02.9) All ages.... 1,759 1,145 1,538 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 5.0 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Malignant neoplasm of gum (CO3) All ages..... 32 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 4 3 55-59 years..... 5 3 2 60-64 years..... 4 65-69 vears..... 3 70-74 vears..... 5 3 75-79 years..... 15 8 80-84 years..... 10 3 7 9 1 1 7 85-89 years..... 2 5 1 90-94 years..... 2 2 2 1 4 95-99 years..... 2 2 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Upper qum (C03.0) All ages.... 20 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 3 3 65-69 vears..... 70-74 vears..... 75-79 years..... 5 80-84 years..... 4 3 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International															
Classification of Disea	ses,												ian or		
Tenth Revision, Second		All rac	es	_	White		_	Black			rican India	ın 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		_ ,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Lower gum (C03.1)															
All ages	5	4	1	5	4	1	_	_	_	_	_	_	_	_	_
Under 1 year	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-9 years	-	_	_	_	_	_	_	_	_	_	-	_	-	_	-
10-14 years	_	-	-	_	-	_	-	_	_	_	-	-	-	-	-
15-19 years	_	-	-	_	-	_	-	_	_	_	-	-	-	-	-
20-24 years	-	-	-	-	-	_	-	-	-	_	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	_	_	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	100		White			Black		λmar	ican Indi		ian or	ic Tela	nder 2/
Edition, 2004)	Both	AII Iac	.cs	Both	WIIICE		Both	Black		Both	ican indi	all I/	Both	IC ISIA	nuer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Gum, unspecified (CO3	.9)														
All ages	38	14	24	30	11	19	6	3	3	_	_	_	2	_	2
Under 1 year	-	-	-	-	-	-	_	-	-	-	-	_	-	-	-
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-54 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-
5-59 years	4	2	2	3	1	2	1	1	-	-	-	_	-	-	-
0-64 years	-	-	-	-	-	-	_	-	-	-	-	_	-	-	-
5-69 years	2	1	1	2	1	1	_	-	-	-	-	_	-	-	-
0-74 years	4	2	2	2	2	-	1	-	1	-	-	-	1	-	1
5-79 years	10	3	7	9	3	6	-	-	_	-	-	_	1	-	1
0-84 years	4	1	3	3	1	2	1	-	1	-	-	_	-	-	-
5-89 years	4	-	4	4	-	4	-	-	_	-	-	_	-	-	-
0-94 years	2	-	2	1	-	1	1	-	1	-	-	_	-	-	-
5-99 years	1	-	1	1	-	1	-	-	_	-	-	_	-	-	-
100 years & over.	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of floor of mouth (CO4) All ages..... 33 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 9 45-49 years..... 8 1 6 17 50-54 years..... 3 14 15 12 1 55-59 years..... 15 13 2 13 11 60-64 years..... 17 13 12 65-69 vears..... 13 8 11 5 70-74 vears..... 13 12 75-79 years..... 12 12 80-84 years..... 11 6 5 9 1 5 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Floor of mouth, unspecified (C04.9) All ages..... 33 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 9 45-49 years..... 8 1 6 17 50-54 years..... 3 14 15 12 1 55-59 years..... 15 13 2 13 11 60-64 years..... 17 13 12 65-69 vears..... 13 8 11 5 70-74 vears..... 13 12 75-79 years..... 12 12 80-84 years..... 11 6 5 9 1 5 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of palate (CO5) All ages..... 72 122 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 4 8 50-54 years..... 18 10 11 55-59 years..... 12 10 2 10 60-64 years..... 13 6 10 3 65-69 vears..... 17 10 9 70-74 vears..... 17 9 8 13 75-79 years..... 22 9 9 13 80-84 years..... 16 6 10 15 5 10 12 85-89 years..... 6 12 6 90-94 years..... 1.0 9 10 1 95-99 years..... 1 4 3 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Hard palate (C05.0) All ages.... 31 9 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 2 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 2 1 65-69 vears..... 3 3 70-74 vears..... 3 75-79 years..... 5 80-84 years..... 6 2 85-89 years..... 90-94 years..... 3 1 2 1 95-99 years..... 1 1 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Soft palate (C05.1) All ages.... 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 4 55-59 years..... 5 4 1 60-64 years..... 3 3 65-69 vears..... 70-74 vears..... 7 75-79 years..... 3 80-84 years..... 4 2 85-89 years..... 3 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Uvula (C05.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Palate, unspecified (C05.9) All ages.... 30 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 50-54 years..... 12 55-59 years..... 3 3 60-64 years..... 8 5 65-69 vears..... 6 70-74 vears..... 6 75-79 years..... 12 6 80-84 years..... 6 2 85-89 years..... 5 2 3 5 3 90-94 years..... 6 6 95-99 years..... 3 3 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of other and unspecified parts of mouth (CO6) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... ρ 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise					1.			_, ,					sian or		
Tenth Revision, Second		All rac	es	Both	White		D	Black			rican Indi	ian 1/		ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male H	Temale	Both sexes	Male	Female
Cheek mucosa (C06.0)															
All ages	29	12	17	27	12	15	2	_	2	_	-	-	-	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	_	-	-	-	_	-	-
40-44 years	1	-	1	1	-	1	-	-	_	-	-	-	_	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	4	2	2	3	2	1	1	-	1	-	-	-	-	-	-
60-64 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
65-69 years	6	4	2	6	4	2	-	-	-	-	-	-	-	-	-
70-74 years	4	-	4	4	-	4	-	-	-	-	-	-	-	-	-
75-79 years	3	-	3	2	-	2	1	-	1	-	-	-	-	-	-
80-84 years	5	3	2	5	3	2	-	-	-	-	-	-	-	-	-
85-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	_	_	-	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disc	eases.											Δs	ian or		
Tenth Revision, Second		All rac	ces		White			Black		Amei	ican Indi			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Vestibule of mouth (C	06.1)														
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	_	-	_	-	-	-	-	-	-	-	_	-	-	-
55-59 years	-	_	-	_	-	-	-	-	-	-	-	_	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-
30-84 years	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-
85-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	_	-	-	-	-	-	-	-	_	-	-	-
100 years & over.	-	-	-	_	_	-	-	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Retromolar area (C06.2) All ages..... 18 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 4 70-74 vears..... 75-79 years..... 5 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Mouth, unspecified (C06.9) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of parotid gland (CO7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.8 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.1 80-84 years..... 85-89 years..... 90-94 years..... 2.4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of other and unspecified major salivary glands (CO8) All ages.... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 6 40-44 years..... 45-49 years..... 11 5 6 5 2 17 50-54 years..... 3 13 14 12 55-59 years..... 16 12 14 11 60-64 years..... 21 15 19 14 1 65-69 vears..... 21 16 18 13 3 5 3 70-74 vears..... 17 9 8 16 9 75-79 years..... 19 15 12 80-84 years..... 16 9 7 16 9 85-89 years..... 14 8 6 14 90-94 years..... 1.0 10 3 3 95-99 years..... 6 5 1 6 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Submandibular gland (C08.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 5 2 45-49 years..... 3 2 50-54 years..... 2 55-59 years..... 1 1 60-64 years..... 3 2 1 65-69 vears..... 2 70-74 vears..... 2 75-79 years..... 80-84 years..... 3 2 85-89 years..... 90-94 years..... 3 2 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Disc													ian or		
Tenth Revision, Second		All rac	es		White			Black			ican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both		_ ,	Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Sublingual gland (C08	.1)														
All ages	1	-	1	-	_	-	1	-	1	-	-	-	_	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-
5-89 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Major salivary gland, unspecified (C08.9) All ages..... 134 46 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 6 3 3 4 2 15 3 50-54 years..... 11 12 55-59 years..... 15 11 13 10 4 60-64 years..... 18 13 5 16 12 1 65-69 vears..... 19 16 11 3 70-74 vears..... 14 8 6 75-79 years..... 18 11 80-84 years..... 13 6 7 13 6 12 85-89 years..... 8 4 12 7 90-94 years..... 2 5 2 95-99 years..... 5 1 4 5 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of tonsil (CO9) All ages.... 498 156 543 412 131 100 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 24 18 18 16 45-49 years..... 59 52 7 48 41 1.0 1 0 98 77 19 50-54 years..... 81 62 21 17 15 55-59 years..... 104 83 21 90 73 17 14 10 60-64 years..... 83 64 19 62 47 15 17 65-69 years..... 86 62 24 69 48 21 13 70-74 years..... 67 13 58 49 9 6 2 75-79 years..... 64 42 2.2 58 38 80-84 years..... 39 23 16 36 21 15 2 2 20 85-89 years..... 11 9 18 10 8 1 90-94 years..... 3 2 1 3 1 2 95-99 years..... 3 3 1 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Tonsillar fossa (C09.0) All ages..... 20 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 3 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International															
Classification of Dis													ian or		
Tenth Revision, Secon		All rac	es		White			Black			ican Indi	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both		_ 1	Both	1	_ ,	Both		_ 1	Both		_ ,	Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Tonsillar pillar (ant	erior) (post	terior) (C	09.1)												
All ages	3	3	-	3	3	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
80-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Tonsil, unspecified (C09.9) All ages.... 478 153 523 395 128 97 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 23 17 17 15 45-49 years..... 56 49 7 45 38 7 1.0 1 0 94 77 74 59 50-54 years..... 2.0 1.8 17 15 55-59 years..... 101 80 21 87 70 17 14 10 60-64 years..... 82 63 19 61 46 15 17 65-69 years..... 85 62 23 68 48 20 13 70-74 years..... 66 53 13 57 48 9 6 2 75-79 years..... 61 39 2.2 56 36 80-84 years..... 35 20 15 33 19 14 18 85-89 years..... 10 8 16 9 7 1 90-94 years..... 3 2 1 3 2 95-99 years..... 3 3 1 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of oropharynx (C10) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 Ω 1 0 50-54 years..... ρ 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Vallecula (C10.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Oropharynx, unspecified (C10.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 Ω 1 0 50-54 years..... ρ 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of nasopharynx (C11) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... ρ 1 0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Posterior wall of nasopharynx (C11.1) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 3 75-79 years..... 80-84 years..... 1 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Anterior wall of nasopharynx (C11.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Nasopharynx, unspecified (C11.9) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... ρ 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.7 2.0 80-84 years..... 2.7 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of pyriform sinus (C12) All ages.... 17 17 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 50-54 years..... 11 11 55-59 years..... 12 12 6 5 6 60-64 years..... 13 10 3 3 65-69 vears..... 12 9 9 3 70-74 vears..... 17 13 3 75-79 years..... 8 6 80-84 years..... 11 8 3 9 85-89 years..... 4 3 90-94 years..... 3 1 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of hypopharynx (C13) All ages.... 161 122 39 39 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 8 8 6 2 19 50-54 years..... 3 15 16 12 55-59 years..... 32 25 23 16 60-64 years..... 25 23 2 19 17 65-69 vears..... 25 9 23 15 16 70-74 vears..... 23 14 9 18 75-79 years..... 34 2.4 10 28 19 80-84 years..... 21 19 2 19 17 2 10 2 2 85-89 years..... 8 8 90-94 years..... 1 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disc												7 -	sian or		
Tenth Revision, Second		All rac	10.5		White			Black		λmos	ican Indi			ia Talo	nder 2/
Edition, 2004)	Both	AII Iac	.cs	Both	WIIICE		Both	black		Both	ICan Indi	all 1/	Both	IC ISIA	iluel 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Postcricoid region (C	13.0)														
All ages	1	1	_	_	_	_	1	1	_	_	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	_	-	-	-	-	-	-	-	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise	eases.											As	sian or		
Tenth Revision, Second		All rac	es		White			Black		Amer	ican Indi			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Hypopharynx, unspecif:	ied (C13.9)														
All ages	201	155	46	161	122	39	38	31	7	_	_	_	2	2	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-39 years	1	-	1	-	-	-	1	_	1	-	-	-	-	-	-
40-44 years	1	-	1	-	-	-	1	_	1	-	-	-	-	-	-
45-49 years	8	8	-	6	6	-	2	2	-	-	-	-	-	-	-
50-54 years	19	16	3	15	12	3	4	4	-	-	-	-	_	-	-
55-59 years	32	25	7	23	16	7	7	7	-	-	-	-	2	2	-
60-64 years	25	23	2	19	17	2	6	6	-	-	-	-	-	-	-
65-69 years	25	16	9	23	15	8	2	1	1	-	-	-	-	-	-
70-74 years	23	14	9	18	11	7	5	3	2	-	-	-	-	-	-
75-79 years	33	23	10	28	19	9	5	4	1	-	-	-	-	-	-
80-84 years	21	19	2	19	17	2	2	2	-	-	-	-	-	-	-
85-89 years	10	8	2	8	7	1	2	1	1	-	-	-	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	_	_	-	-	-	-	_	-	-
100 years & over.	-	-	-	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Bot.h and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Malignant neoplasm of other and ill-defined sites in the lip, oral cavity, and pharynx (C14) All ages.... 1,604 1,193 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 2.7 2.8 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.4 80-84 years..... 85-89 years..... 90-94 years..... 3.8 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pharynx, unspecified (C14.0) All ages.... 1,604 1,193 1,302 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 2.7 2.8 50-54 years..... 2.1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.4 80-84 years..... 85-89 years..... 90-94 years..... 3.8 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International												_			
Classification of Dis								-1 1		_			ian or	c· - 3	1 0 /
Tenth Revision, Secon		All rac	es	D - 1-1-	White		D - 1-1-	Black			ican In	dian I/		tic Isla	ander 2/
Edition, 2004)	Both	M-l-	Doma la	Both	Mala	E1-	Both	M-1-	E1-	Both	Mala	Esmals.	Both	W-1-	Dama la
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	мате	Female	sexes	Male	Female
Malignant neoplasm of	esophagus	(C15)													
All ages	13,686	10,723	2,963	11,959	9,476	2,483	1,452	1,045	407	66	45	21	209	157	52
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	_	_	-	-	-	-	-	_	-
20-24 years	3	3	-	1	1	-	1	1	-	-	-	-	1	1	-
25-29 years	13	12	1	11	10	1	2	2	-	-	-	-	-	-	-
30-34 years	16	13	3	16	13	3	-	-	-	-	-	-	-	-	-
35-39 years	81	69	12	73	62	11	7	6	1	-	-	-	1	1	-
40-44 years	198	172	26	171	151	20	24	19	5	-	-	-	3	2	1
45-49 years	532	457	75	437	385	52	84	63	21	4	3	1	7	6	1
50-54 years	992	817	175	798	671	127	176	133	43	9	6	3	9	7	2
55-59 years	1,525	1,323	202	1,271	1,134	137	222	163	59	7	5	2	25	21	4
60-64 years	1,764	1,478	286	1,515	1,290	225	218	163	55	11	8	3	20	17	3
65-69 years	1,810	1,490	320	1,559	1,300	259	216	163	53	11	9	2	24	18	6
70-74 years	1,881	1,484	397	1,657	1,315	342	182	137	45	10	9	1	32	23	9
75-79 years	1,984	1,466	518	1,783	1,333	450	150	99	51	10	4	6	41	30	11
80-84 years	1,531	1,119	412	1,411	1,051	360	100	57	43	2	-	2	18	11	7
85-89 years	919	599	320	844	548	296	51	33	18	2	1	1	22	17	5
90-94 years	362	196	166	343	188	155	14	5	9	-	-	-	5	3	2
95-99 years	67	24	43	62	23	39	4	1	3	_	_	-	1	-	1
100 years & over.	7	_	7	6	_	6	1	_	1	_	_	_	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cervical part of esophagus (C15.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Thoracic part of esophagus (C15.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Abdominal part of esophagus (C15.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Upper third of esophagus (C15.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Middle third of esophagus (C15.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Lower third of esophagus (C15.5) All ages..... 10 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 2 2 50-54 years..... 4 4 55-59 years..... 3 13 10 14 11 60-64 years..... 14 13 14 13 65-69 vears..... 11 10 10 70-74 vears..... 5 4 5 75-79 years..... 12 10 80-84 years..... 8 7 8 85-89 years..... 4 3 1 3 90-94 years..... 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Esophagus, unspecified (C15.9) All ages.... 13,596 10,649 2,947 11,873 9,406 2,467 1,449 1,042 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1.511 1,312 1,258 1,124 60-64 years..... 1,746 1,463 1,498 1,276 1.798 1,479 1,549 1,291 65-69 vears..... 70-74 vears..... 1,875 1,479 1,651 1,310 75-79 years..... 1,971 1,455 1,770 1,322 80-84 years..... 1,521 1,110 1,401 1,042 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dis								-1 1		_			sian or		3 0 /
Tenth Revision, Secon		All rac	es		White			Black			ican In	dian 1/		tic Isla	ander 2/
Edition, 2004)	Both	24 - 1 -	W1-	Both	M . 7 .	m1.	Both	34.3.		Both	M . 1 .	m1	Both	M . 1 .	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Malignant neoplasm of	f stomach (C	16)													
All ages	11,345	6,653	4,692	8,582	5,102	3,480	1,916	1,092	824	84	58	26	763	401	362
Under 1 year	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
15-19 years	3	1	2	2	1	1	-	-	-	-	-	-	1	-	1
20-24 years	15	9	6	10	6	4	3	2	1	-	-	-	2	1	1
25-29 years	38	15	23	29	11	18	5	3	2	-	-	-	4	1	3
30-34 years	86	43	43	60	28	32	15	10	5	2	2	-	9	3	6
35-39 years	171	79	92	117	59	58	32	11	21	2	2	-	20	7	13
40-44 years	294	154	140	188	96	92	67	40	27	4	3	1	35	15	20
45-49 years	480	288	192	331	206	125	101	62	39	9	5	4	39	15	24
50-54 years	685	441	244	502	329	173	125	83	42	4	1	3	54	28	26
55-59 years	911	620	291	669	467	202	161	109	52	10	5	5	71	39	32
60-64 years	959	642	317	700	472	228	190	124	66	6	5	1	63	41	22
65-69 years	1,140	743	397	842	561	281	208	131	77	11	9	2	79	42	37
70-74 years	1,322	821	501	994	629	365	227	124	103	9	6	3	92	62	30
75-79 years	1,647	975	672	1,274	772	502	251	151	100	8	4	4	114	48	66
80-84 years	1,696	974	722	1,321	768	553	272	147	125	11	8	3	92	51	41
85-89 years	1,184	584	600	960	473	487	162	72	90	6	6	-	56	33	23
90-94 years	555	210	345	466	180	286	63	17	46	-	_	-	26	13	13
95-99 years	140	47	93	109	40	69	25	5	20	-	_	-	6	2	4
100 years & over.	18	6	12	8	4	4	8	_	8	2	2	-	-	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Cardia (C16.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... R 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.7 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise	eases,											As	sian or		
Tenth Revision, Second	f	All rac	es		White			Black		Amei	rican Indi	an 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Fundus of stomach (C16	6.1)														
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
15-19 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	_
20-24 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	_
30-34 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	_	_	-	-	_	-	_	-	-	-	-	-
40-44 years	-	-	-	_	_	-	-	_	-	_	-	-	-	-	-
45-49 years	_	-	-	_	-	-	-	-	-	_	_	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
70-74 years	_	-	-	_	-	-	-	-	-	_	_	-	-	-	-
75-79 years	1	1	-	1	1	_	_	-	-	-	-	-	_	-	-
30-84 years	_	_	-	-	-	_	_	-	-	-	-	-	_	-	-
35-89 years	_	-	-	_	-	-	-	-	-	_	_	-	-	-	-
90-94 years	_	-	-	_	-	-	-	-	-	_	_	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Disea Tenth Revision, Second	ıses,	All rac	0.5		White			Black		Amos	rican Indi		ian or	ia Talo	nder 2/
	Dath	All rac	es	Dath	white		Both	втаск		Both	rican indi	an 1/	Both	IC ISIA	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Pyloric antrum (C16.3)															
All ages	6	3	3	5	2	3	1	1	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	-	_	-	-	-	-	_	_	_	-	-	-
55-59 years	-	-	-	-	_	-	-	-	-	_	_	_	-	-	-
0-64 years	1	1	-	1	1	-	-	-	-	_	_	_	-	-	-
5-69 years	_	_	-	-	_	_	-	-	-	_	-	-	-	-	-
70-74 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-79 years	_	_	-	-	_	_	_	_	_	_	-	-	_	_	-
0-84 years	1	1	-	1	1	_	_	_	_	_	-	-	_	_	-
5-89 years	3	1	2	2	_	2	1	1	_	_	-	-	_	_	-
0-94 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
95-99 years	1	_	1	1	_	1	_	_	_	_	-	-	_	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	1AG		White			Black		Δmea	rican Indi		ian or	ic Tela	nder 2/
Edition, 2004)	Both	AII IAC	.05	Both	WIIICC		Both	Diack		Both	. ican inai	Lair 1/	Both	.10 1510	iluci 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	remale	sexes	Male	Female
Pylorus (C16.4)															
All ages	11	5	6	9	4	5	1	1	_	_	_	_	1	_	1
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	-	1	-	-	-	-	-	-	-	-	-	1	-	1
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
85-89 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
90-94 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	_	_	_	_	_	_	_	_	_	_	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Lesser curvature of stomach, unspecified (C16.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Greater curvature of stomach, unspecified (C16.6) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Course of death (bosse															
Cause of death (based on the International	1														
Classification of Dis	seases,											As	ian or		
Tenth Revision, Secon	nd	All rac	es		White			Black		Amer	ican In	dian 1/	Paci	fic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Stomach, unspecified	(C16.9)														
All ages	10,610	6,080	4,530	7,889	4,558	3,331	1,883	1,069	814	83	57	26	755	396	359
Under 1 year	-	_	_	_	_	_	-	_	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	2	1	1	1	1	-	-	-	-	-	-	-	1	-	1
20-24 years	12	8	4	8	5	3	2	2	-	-	-	-	2	1	1
25-29 years	36	14	22	27	10	17	5	3	2	-	-	-	4	1	3
30-34 years	85	42	43	59	27	32	15	10	5	2	2	-	9	3	6
35-39 years	163	73	90	111	54	57	32	11	21	2	2	-	18	6	12
40-44 years	278	142	136	174	86	88	66	39	27	4	3	1	34	14	20
45-49 years	438	253	185	290	171	119	100	62	38	9	5	4	39	15	24
50-54 years	631	396	235	456	291	165	117	76	41	4	1	3	54	28	26
55-59 years	815	533	282	576	382	194	158	107	51	10	5	5	71	39	32
60-64 years	870	573	297	616	405	211	187	123	64	5	4	1	62	41	21
65-69 years	1,052	674	378	759	495	264	204	129	75	11	9	2	78	41	37
70-74 years	1,241	756	485	918	569	349	224	121	103	9	6	3	90	60	30
75-79 years	1,545	902	643	1,174	699	475	250	151	99	8	4	4	113	48	65
80-84 years	1,619	911	708	1,249	709	540	267	143	124	11	8	3	92	51	41
85-89 years	1,129	543	586	907	434	473	160	70	90	6	6	-	56	33	23
90-94 years	542	207	335	453	177	276	63	17	46	-	-	-	26	13	13
95-99 years	135	47	88	104	40	64	25	5	20	-	-	-	6	2	4
100 years & over.	17	5	12	7	3	4	8	-	8	2	2	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of small intestine (C17) All ages.... 1,091 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.2 1 0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Duodenum (C17.0) All ages.... 580 301 279 480 218 83 13 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 12 45-49 years..... 14 9 5 10 Ω 2 4 28 50-54 years..... 35 19 7 16 16 12 55-59 years..... 44 20 24 32 17 15 10 2 60-64 years..... 56 32 24 44 17 11 1 65-69 years..... 55 33 22 46 28 18 9 70-74 years..... 85 53 32 66 22 75-79 years..... 83 39 44 72 37 35 8 80-84 years..... 89 51 38 75 46 29 9 5 2 25 85-89 years..... 68 43 61 21 40 3 1 25 90-94 years..... 8 17 2.4 16 1 1 8 95-99 years..... 5 5 1 4 100 years & over. 2 2 1 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Jejunum (C17.1) All ages.... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 6 2 50-54 years..... 3 55-59 years..... 3 3 1 60-64 years..... 9 3 65-69 vears..... 3 70-74 vears..... 3 75-79 years..... 5 80-84 years..... 6 3 2 2 85-89 years..... 2 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Ileum (C17.2) All ages.... 37 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 50-54 years..... 55-59 years..... 4 1 60-64 years..... 2 2 65-69 vears..... 5 70-74 vears..... 8 3 5 75-79 years..... 80-84 years..... 5 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Meckel's diverticulum (C17.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Small intestine, unspecified (C17.9) All ages.... 227 199 369 195 174 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 4 9 40-44 years..... 19 10 45-49 years..... 2.7 14 13 22 12 10 5 2.4 8 50-54 years..... 32 18 14 14 10 55-59 years..... 40 21 19 37 18 19 60-64 years..... 33 15 18 30 15 15 1 65-69 years..... 47 26 21 40 23 17 70-74 years..... 53 33 20 45 19 6 75-79 years..... 68 32 36 60 2.8 32 2 2 80-84 years..... 49 25 24 44 21 23 5 1 20 85-89 years..... 38 23 15 34 14 4 3 1 9 90-94 years..... 3 6 9 3 6 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis													ian or		
Tenth Revision, Secon		All rac	ces		White			Black			ican Ind	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Malignant neoplasm of	E colon (C18)													
All ages	44,061	21,759	22,302	37,121	18,342	18,779	5,816	2,871	2,945	185	99	86	939	447	492
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	1	1	-	-	-	-	-	-	-	-	-	-	1	1	-
15-19 years	8	6	2	6	6	-	1	-	1	-	-	-	1	-	1
20-24 years	27	10	17	23	9	14	3	1	2	1	-	1	-	-	-
25-29 years	81	41	40	60	30	30	17	10	7	-	-	-	4	1	3
30-34 years	156	76	80	111	60	51	35	13	22	3	-	3	7	3	4
35-39 years	308	168	140	225	126	99	70	38	32	2	1	1	11	3	8
40-44 years	654	335	319	481	257	224	145	67	78	1	-	1	27	11	16
45-49 years	1,285	709	576	961	538	423	285	150	135	9	6	3	30	15	15
50-54 years	2,120	1,129	991	1,619	867	752	433	233	200	11	7	4	57	22	35
55-59 years	3,093	1,734	1,359	2,436	1,395	1,041	552	287	265	22	14	8	83	38	45
60-64 years	3,696	2,162	1,534	2,950	1,753	1,197	624	330	294	28	16	12	94	63	31
65-69 years	4,384	2,520	1,864	3,571	2,083	1,488	685	365	320	25	17	8	103	55	48
70-74 years	5,222	2,881	2,341	4,398	2,439	1,959	702	381	321	16	8	8	106	53	53
75-79 years	6,461	3,276	3,185	5,536	2,840	2,696	762	363	399	33	16	17	130	57	73
80-84 years	7,074	3,324	3,750	6,224	2,923	3,301	705	324	381	18	7	11	127	70	57
85-89 years	5,499	2,152	3,347	4,927	1,920	3,007	458	192	266	13	6	7	101	34	67
90-94 years	3,010	976	2,034	2,701	867	1,834	266	93	173	2	1	1	41	15	26
95-99 years	879	235	644	802	208	594	64	22	42	1	-	1	12	5	7
100 years & over.	100	22	78	88	20	68	8	1	7	-	-	-	4	1	3
Age not stated	3	2	1	2	1	1	1	1	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Cecum (C18.0) All ages.... 282 119 163 242 147 38 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 3 2 50-54 years..... 16 9 11 55-59 years..... 10 7 8 2 3 60-64 years..... 16 9 11 5 65-69 years..... 15 5 10 11 70-74 years..... 30 12 18 25 16 75-79 years..... 38 10 28 34 25 80-84 years..... 53 23 30 46 19 27 6 27 41 27 2 85-89 years..... 43 16 14 2 90-94 years..... 41 18 23 3.8 17 21 3 1 95-99 years..... 11 2 9 9 11 2 100 years & over. 2 2 2 2 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Appendix (C18.1) All ages.... 348 173 175 309 153 156 29 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 13 9 10 2 40-44 years..... 11 9 10 2 45-49 years..... 38 2.0 18 35 17 25 38 23 5 50-54 years..... 43 18 15 55-59 years..... 63 33 30 56 28 28 4 3 60-64 years..... 45 19 26 39 18 21 65-69 years..... 42 23 19 39 21 18 70-74 years..... 28 15 13 27 13 75-79 years..... 19 8 11 15 80-84 years..... 16 10 6 14 10 1 1 85-89 years..... 12 6 6 12 6 6 7 2 90-94 years..... 2 5 1 5 1 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Ascending colon (C18.2) All ages.... 295 135 160 253 115 138 25 12 17 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 3 45-49 years..... 6 3 3 5 3 50-54 years..... 11 9 2 9 55-59 years..... 19 9 10 14 2 8 1 60-64 years..... 16 9 7 14 8 6 2 65-69 vears..... 33 18 15 27 13 70-74 vears..... 31 15 16 22 11 11 4 75-79 years..... 39 18 21 34 15 19 1 80-84 years..... 55 28 27 51 26 25 3 2 1 52 47 85-89 years..... 15 37 15 32 1 90-94 years..... 2.4 9 15 23 14 1 1 9 95-99 years..... 3 2 3 1 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise													ian or		
Tenth Revision, Second		All rac	es		White			Black			rican Ind	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Hepatic flexure (C18.3	3)														
All ages	3	-	3	3	_	3	_	-	-	_	-	-	_	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
70-74 years	1	-	1	1	-	1	-	-	-	_	_	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	_	_	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Transverse colon (C18.4) All ages..... 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 5 4 1 60-64 years..... 4 65-69 vears..... 3 3 70-74 vears..... 6 75-79 years..... 80-84 years..... 6 85-89 years..... 90-94 years..... 4 4 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dise	eases,											As	ian or		
Tenth Revision, Second	f	All rac	ces		White			Black		Ame	rican Indi	an 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Splenic flexure (C18.	5)														
All ages	6	4	2	5	3	2	1	1	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-		-	-	-	-	-	-	_	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-		-	-	-	-	-	-	_	-	-	-	-	-
50-54 years	-	-		-	-	-	-	-	-	_	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
60-64 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
70-74 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
75-79 years	-	-		-	-	-	-	-	-	_	-	-	-	-	-
30-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-89 years	1	1		-	-	-	1	1	-	_	-	-	-	-	-
90-94 years	-	-		-	-	-	-	-	-	_	-	-	-	-	-
95-99 years	-	-		-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	-	-	-	_	-	-	_	_	_	-	-	-	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Descending colon (C18.6) All ages.... 30 20 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 2 50-54 years..... 4 2 55-59 years..... 2 2 4 60-64 years..... 5 5 65-69 vears..... 70-74 vears..... 8 6 75-79 years..... 3 80-84 years..... 5 85-89 years..... 3 90-94 years..... 2 2 1 1 4 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Sigmoid colon (C18.7) All ages.... 354 184 170 294 155 139 39 17 20 14 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 9 40-44 years..... 45-49 years..... 10 5 5 a 16 3 50-54 years..... 9 13 55-59 years..... 28 10 18 21 9 12 3 4 60-64 years..... 25 17 8 18 12 6 5 1 65-69 years..... 34 19 27 15 12 15 70-74 years..... 29 18 11 25 16 9 75-79 years..... 48 29 19 44 2.8 16 80-84 years..... 52 24 28 43 2.0 23 6 5 3 39 17 2 85-89 years..... 21 18 35 18 2 90-94 years..... 43 16 27 35 12 23 4 1 4 3 95-99 years..... 12 11 5 6 6 6 1 1 100 years & over. 1 1 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

2

1

1

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International	ì														
Classification of Dis	seases,											As	ian or		
Tenth Revision, Secon	nd	All rac	ces		White			Black		Amer	ican In			fic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Colon, unspecified (C	218.9)														
All ages	42,673	21,091	21,582	35,932	17,776	18,156	5,674	2,793	2,881	182	96	86	885	426	459
Under 1 year	-	-	-	_	-	-	-	-	-	-	-	-	_	-	-
1-4 years	-	_	-	_	-	-	-	_	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
10-14 years	1	1	-	-	-	_	-	-	-	-	-	-	1	1	-
15-19 years	8	6	2	6	6	-	1	_	1	-	-	-	1	-	1
20-24 years	25	8	17	21	7	14	3	1	2	1	-	1	-	-	-
25-29 years	79	39	40	58	28	30	17	10	7	-	-	-	4	1	3
30-34 years	144	67	77	101	53	48	33	11	22	3	-	3	7	3	4
35-39 years	283	152	131	204	112	92	66	36	30	2	1	1	11	3	8
40-44 years	628	327	301	463	251	212	142	66	76	1	-	1	22	10	12
45-49 years	1,227	678	549	910	511	399	280	147	133	9	6	3	28	14	14
50-54 years	2,030	1,080	950	1,548	833	715	416	220	196	11	7	4	55	20	35
55-59 years	2,964	1,669	1,295	2,329	1,341	988	540	281	259	21	13	8	74	34	40
60-64 years	3,583	2,098	1,485	2,859	1,701	1,158	606	320	286	28	16	12	90	61	29
65-69 years	4,250	2,448	1,802	3,458	2,023	1,435	668	355	313	25	17	8	99	53	46
70-74 years	5,088	2,813	2,275	4,285	2,381	1,904	687	374	313	16	8	8	100	50	50
75-79 years	6,302	3,204	3,098	5,396	2,775	2,621	749	358	391	32	15	17	125	56	69
80-84 years	6,883	3,232	3,651	6,056	2,842	3,214	689	317	372	18	7	11	120	66	54
85-89 years	5,341	2,090	3,251	4,784	1,864	2,920	450	187	263	12	5	7	95	34	61
90-94 years	2,887	929	1,958	2,593	827	1,766	255	87	168	2	1	1	37	14	23
95-99 years	850	226	624	774	200	574	63	21	42	1	-	1	12	5	7
100 years & over.	97	22	75	85	20	65	8	1	7	-	-	-	4	1	3

1

1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International Classification of Dis	02505											λα	ian or		
Tenth Revision, Secon		All rac	96		White			Black		Amer	ican Ind			ic Isla	nder 2/
Edition, 2004)	Both	AII IAC	CB	Both	WIIICC		Both	Diack		Both	ican ina.	Idii I/	Both	.10 1514	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Malignant neoplasm of	rectosigmo	id junctio	n (C19)												
All ages	2,566	1,412	1,154	2,171	1,207	964	336	176	160	12	5	7	47	24	23
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	3	3	-	3	3	-	-	-	-	-	-	-	-	-	-
25-29 years	6	2	4	4	1	3	-	-	-	-	-	-	2	1	1
30-34 years	19	11	8	14	8	6	5	3	2	-	-	-	-	-	-
35-39 years	38	16	22	30	9	21	5	4	1	-	-	-	3	3	-
10-44 years	58	23	35	41	16	25	15	6	9	-	-	-	2	1	1
15-49 years	113	66	47	88	50	38	20	13	7	1	-	1	4	3	1
50-54 years	194	111	83	152	88	64	37	20	17	2	1	1	3	2	1
5-59 years	251	152	99	209	130	79	33	19	14	-	-	-	9	3	6
50-64 years	278	170	108	226	145	81	43	21	22	2	-	2	7	4	3
55-69 years	291	180	111	244	153	91	40	21	19	3	2	1	4	4	-
70-74 years	321	192	129	281	168	113	37	23	14	-	-	-	3	1	2
5-79 years	353	197	156	323	182	141	26	14	12	1	-	1	3	1	2
0-84 years	321	164	157	280	143	137	37	19	18	2	2	-	2	-	2
85-89 years	195	89	106	170	79	91	24	9	15	-	-	-	1	1	-
00-94 years	101	32	69	87	28	59	11	4	7	-	-	-	3	-	3
95-99 years	22	3	19	18	3	15	3	-	3	1	-	1	-	-	-
100 years & over.	2	1	1	1	1	-	-	-	-	-	-	-	1	-	1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dis	02505											7.0	ian or		
Tenth Revision, Second		All rac	100		White			Black		Amer	ican In			ic Tela	ınder 2/
Edition, 2004)	Both	AII Iac	.cs	Both	WILLCE		Both	Black		Both	ICan III	ulan 1/	Both	.IC ISIC	ilidel 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Malignant neoplasm of	rectum (C20	O)													
All ages	6,299	3,514	2,785	5,408	3,016	2,392	673	367	306	31	19	12	187	112	75
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	4	1	3	2	1	1	1	_	1	-	-	-	1	-	1
25-29 years	18	11	7	15	8	7	1	1	-	-	-	-	2	2	-
30-34 years	36	22	14	23	14	9	8	5	3	-	-	-	5	3	2
35-39 years	66	33	33	57	29	28	8	3	5	-	-	-	1	1	-
40-44 years	159	98	61	114	67	47	32	21	11	-	-	-	13	10	3
15-49 years	290	163	127	234	129	105	44	28	16	3	2	1	9	4	5
50-54 years	392	254	138	319	208	111	56	34	22	4	2	2	13	10	3
55-59 years	554	349	205	473	298	175	56	36	20	7	4	3	18	11	7
50-64 years	606	395	211	517	332	185	65	47	18	5	3	2	19	13	6
65-69 years	671	422	249	575	369	206	69	39	30	5	3	2	22	11	11
70-74 years	704	416	288	611	362	249	67	40	27	3	2	1	23	12	11
75-79 years	827	471	356	709	412	297	90	44	46	3	2	1	25	13	12
80-84 years	864	452	412	750	391	359	90	46	44	1	1	-	23	14	9
85-89 years	630	268	362	571	247	324	53	16	37	_	-	-	6	5	1
90-94 years	361	127	234	330	120	210	25	5	20	-	-	-	6	2	4
95-99 years	95	27	68	87	24	63	7	2	5	-	-	-	1	1	-
100 years & over.	22	5	17	21	5	16	1	_	1	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of anus and anal canal (C21) All ages.... 387 545 210 335 70 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 4 35-39 years..... 14 11 25 40-44 years..... 33 16 17 10 15 45-49 years..... 49 22 27 40 17 23 7 36 49 30 50-54 years..... 56 2.0 19 6 55-59 years..... 75 32 43 64 30 34 11 60-64 years..... 73 24 49 65 23 42 8 65-69 years..... 53 20 33 51 19 32 70-74 years..... 63 31 32 58 27 31 4 75-79 years..... 64 26 38 58 32 80-84 years..... 60 11 49 54 11 43 4 2 16 50 85-89 years..... 53 37 16 34 3 90-94 years..... 21 5 16 13 5 3 16 3 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Anus, unspecified (C21.0) All ages.... 511 200 311 449 177 272 37 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 13 5 10 40-44 years..... 27 14 13 20 12 45-49 years..... 44 2.0 24 35 15 2.0 7 30 43 17 26 50-54 years..... 48 18 55-59 years..... 62 28 34 52 26 26 10 60-64 years..... 59 18 41 53 17 36 6 65-69 years..... 17 27 44 17 27 44 70-74 years..... 48 22 26 45 20 25 75-79 years..... 51 2.2 29 46 2.2 2.4 80-84 years..... 49 11 38 45 11 34 4 2 85-89 years..... 41 13 28 39 13 26 90-94 years..... 18 5 3 2 13 15 3 12 1 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Anal canal (C21.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 2 5 50-54 years..... 55-59 years..... 5 1 4 60-64 years..... 3 1 65-69 vears..... 6 5 70-74 vears..... 6 75-79 years..... 4 80-84 years..... 2 2 2 85-89 years..... 1 1 90-94 years..... 1 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cloacogenic zone (C21.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Overlapping lesion of rectum, anus, and anal canal (C21.8) All ages.... 20 37 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 3 40-44 years..... 45-49 years..... 3 2 1 50-54 years..... 2 1 1 55-59 years..... 8 3 5 60-64 years..... 3 3 65-69 vears..... 3 70-74 vears..... 8 5 75-79 years..... 7 80-84 years..... 9 9 85-89 years..... 11 9 10 90-94 years..... 2 2 1 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based																	
on the International																	
Classification of Dis										Asian or							
Tenth Revision, Secon		All races		White			Black			American Indian 1/			Pacific Islander 2/				
Edition, 2004)	Both			Both			Both			Both			Both				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female		
Malignant neoplasm of	liver and	intrahepat	ic bile duc	ts (C22)													
All ages	16,525	10,759	5,766	13,103	8,419	4,684	2,187	1,515	672	134	88	46	1,101	737	364		
Under 1 year	2	2	-	1	1	-	-	-	-	-	-	-	1	1	-		
1-4 years	15	10	5	12	8	4	2	1	1	-	-	-	1	1	-		
5-9 years	7	5	2	5	4	1	2	1	1	-	-	-	_	-	-		
10-14 years	9	4	5	9	4	5	-	-	-	-	-	-	_	-	-		
15-19 years	12	4	8	11	4	7	1	-	1	-	-	-	-	-	-		
20-24 years	19	10	9	17	9	8	1	1	-	-	-	-	1	-	1		
25-29 years	31	23	8	17	10	7	7	6	1	-	-	-	7	7	-		
30-34 years	56	31	25	34	14	20	13	10	3	1	-	1	8	7	1		
35-39 years	103	68	35	54	33	21	23	16	7	1	-	1	25	19	6		
40-44 years	258	183	75	170	110	60	50	40	10	1	1	-	37	32	5		
45-49 years	733	558	175	502	385	117	151	105	46	9	9	-	71	59	12		
50-54 years	1,673	1,332	341	1,225	976	249	338	264	74	15	12	3	95	80	15		
55-59 years	2,095	1,640	455	1,521	1,173	348	438	369	69	17	13	4	119	85	34		
60-64 years	1,747	1,252	495	1,335	941	394	264	199	65	16	12	4	132	100	32		
65-69 years	1,789	1,244	545	1,400	966	434	224	158	66	23	17	6	142	103	39		
70-74 years	2,041	1,293	748	1,675	1,070	605	213	138	75	15	5	10	138	80	58		
75-79 years	2,228	1,315	913	1,881	1,128	753	184	101	83	12	10	2	151	76	75		
80-84 years	1,933	1,046	887	1,658	918	740	156	67	89	19	8	11	100	53	47		
85-89 years	1,189	543	646	1,052	483	569	81	32	49	3	-	3	53	28	25		
90-94 years	462	170	292	414	157	257	29	6	23	2	1	1	17	6	11		
95-99 years	116	25	91	103	24	79	10	1	9	-	-	-	3	-	3		
100 years & over.	7	1	6	7	1	6	-	_	-	-	-	-	-	-	-		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Liver cell carcinoma (C22.0) All ages.... 6,815 5,177 1,638 5,116 3,888 1,228 1,053 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 vears..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1,090 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 2.7 80-84 years..... 85-89 years..... 90-94 years..... 8.0 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Intrahepatic bile duct carcinoma (C22.1) All ages.... 3,693 1,802 1,891 3,220 1,570 1,650 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... Ω 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Hepatoblastoma (C22.2) All ages.... 12 18 10 Under 1 year.... 2 2 1 1-4 years..... 13 8 10 4 2 5-9 years..... 2 1 1 10-14 years..... 3 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second All rac				referènce		Asian or American Indian 1/ Pacific Islander 2/									
Edition, 2004)	a Both	All rac	es	Both	White		Both	Black		Ame: Both	rican indi	an 1/	Pacii Both	ic isia	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Angiosarcoma of liver	(C22.3)														
All ages	21	14	7	20	13	7	1	1	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
45-49 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
60-64 years	4	3	1	4	3	1	-	-	-	-	-	-	-	-	-
65-69 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
70-74 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
75-79 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
30-84 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
85-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	-	_	_	-	-	_	_	_	-	_	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other sarcomas of liver (C22.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

24JUN09

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified carcinomas of liver (C22.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Liver, unspecified (C22.9) All ages.... 5,968 3,752 2,216 4,727 2,938 1,789 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 2.4 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of gallbladder (C23) All ages.... 2,000 1,425 1,689 1,190 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 2.6 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Malignant neoplasm of other and unspecified parts of biliary tract (C24) All ages.... 1,427 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 2.2 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Extrahepatic bile duct (C24.0) All ages..... 317 318 563 278 285 19 23 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 45-49 years..... 4 3 1 2 1 1 1 17 2 50-54 years..... 25 8 23 ρ 15 55-59 years..... 46 27 19 41 23 18 2 3 60-64 years..... 44 28 16 35 22 13 4 3 65-69 years..... 44 25 19 35 21 14 70-74 years..... 101 57 44 89 49 40 8 2 75-79 years..... 100 49 87 43 44 10 1 80-84 years..... 100 47 53 97 47 50 3 95 52 85-89 years..... 106 44 62 43 3 31 90-94 years..... 41 9 32 40 1 a 1 95-99 years..... 13 9 10 8 2 1 100 years & over. 2 2 2 2 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Ampulla of Vater (C24.1) All ages..... 158 138 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 2 2 8 2 50-54 years..... 6 2 55-59 years..... 12 8 8 4 60-64 years..... 18 12 6 17 11 1 65-69 vears..... 19 10 9 18 9 70-74 vears..... 18 11 17 75-79 years..... 2.1 15 19 5 6 80-84 years..... 32 17 15 29 16 13 2 1 13 85-89 years..... 3 10 13 3 10 7 90-94 years..... 2 5 6 2 95-99 years..... 1 3 3 1 1 1 1 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Biliary tract, unspecified (C24.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1.8 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based																	
on the International																	
Classification of Dis										Asian or							
Tenth Revision, Secon	d	All rac	es		White			Black			American Indian 1/		Pacific Islander 2				
Edition, 2004)	Both			Both			Both			Both			Both				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female		
Malignant neoplasm of	pancreas (C25)															
All ages	33,454	16,559	16,895	28,677	14,374	14,303	3,868	1,768	2,100	138	55	83	771	362	409		
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
1-4 years	2	1	1	_	-	-	1	-	1	_	-	-	1	1	-		
5-9 years	1	-	1	1	-	1	_	-	-	_	-	-	-	-	-		
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
l5-19 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-		
20-24 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-		
25-29 years	16	5	11	12	4	8	4	1	3	-	-	-	-	-	-		
30-34 years	33	16	17	26	14	12	4	1	3	-	-	-	3	1	2		
35-39 years	123	88	35	96	69	27	22	15	7	_	-	-	5	4	1		
40-44 years	383	220	163	296	175	121	67	37	30	5	3	2	15	5	10		
45-49 years	992	584	408	777	472	305	175	87	88	5	2	3	35	23	12		
50-54 years	1,738	1,077	661	1,393	870	523	299	182	117	3	3	-	43	22	21		
55-59 years	2,795	1,701	1,094	2,292	1,411	881	417	250	167	13	7	6	73	33	40		
60-64 years	3,503	2,034	1,469	2,932	1,729	1,203	481	259	222	21	11	10	69	35	34		
65-69 years	3,914	2,193	1,721	3,312	1,889	1,423	494	250	244	19	9	10	89	45	44		
70-74 years	4,559	2,344	2,215	3,879	2,037	1,842	537	246	291	19	4	15	124	57	67		
75-79 years	5,179	2,412	2,767	4,554	2,165	2,389	508	194	314	14	6	8	103	47	56		
30-84 years	5,081	2,157	2,924	4,527	1,964	2,563	422	144	278	24	4	20	108	45	63		
85-89 years	3,290	1,234	2,056	2,952	1,133	1,819	261	71	190	11	3	8	66	27	39		
90-94 years	1,471	410	1,061	1,295	366	929	143	26	117	4	3	1	29	15	14		
95-99 years	329	75	254	299	71	228	25	3	22	-	-	_	5	1	4		
100 years & over.	40	5	35	29	2	27	8	2	6	_	-	-	3	1	2		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Head of pancreas (C25.0) All ages..... 100 49 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 1 2 50-54 years..... 4 3 1 55-59 years..... 5 2 3 60-64 years..... 8 2 6 65-69 vears..... 7 3 4 70-74 vears..... 3 3 3 75-79 years..... 13 8 80-84 years..... 19 10 9 17 8 2 1 22 7 21 15 85-89 years..... 15 90-94 years..... 1.0 10 6 4 6 95-99 years..... 3 2 3 1 2 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Body of pancreas (C25.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis		311									Asian or American Indian 1/ Pacific Islander 2/							
Tenth Revision, Secon Edition, 2004)	a Both	All rac	es	Both	White		Both	Black		Amei Both	rican indi	ian I/	Pacii Both	ic isia	naer 2/			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male I	Female	sexes	Male	Female			
Tail of pancreas (C25	.2)																	
All ages	15	8	7	13	7	6	1	_	1	_	-	-	1	1	-			
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-			
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
20-24 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-			
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
35-39 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-			
40-44 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-			
45-49 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-			
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-			
60-64 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-			
65-69 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-			
70-74 years	3	3	-	3	3	-	-	-	-	_	-	-	-	-	-			
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-			
30-84 years	2	1	1	1	-	1	-	-	-	-	-	-	1	1	-			
35-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-			
90-94 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-			
95-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-			
100 years & over.	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-			
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_			

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pancreatic duct (C25.3) All ages..... 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 50-54 years..... 55-59 years..... 60-64 years..... 3 65-69 vears..... 4 3 70-74 vears..... 2 75-79 years..... 80-84 years..... 4 3 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Endocrine pancreas (C25.4) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 3 40-44 years..... 45-49 years..... 12 6 2 2 50-54 years..... 10 55-59 years..... 12 10 5 60-64 years..... 9 2 8 1 65-69 vears..... 12 70-74 vears..... 6 75-79 years..... 14 80-84 years..... 7 2 5 6 1 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International	l															
On the international Classification of Dis	eageg											Δα	sian or			
Tenth Revision, Secon		All rac	es		White			Black			American Indian 1/			Pacific Islander 2/		
Edition, 2004)	Both	1111 100	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	Both	,,,,,		Both	224071		Both		41411 17	Both	IIO IDIO	ilidel 2,	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	
Pancreas, unspecified	(C25.9)															
All ages	33,222	16,442	16,780	28,476	14,274	14,202	3,845	1,755	2,090	138	55	83	763	358	405	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	2	1	1	-	-	-	1	-	1	-	-	-	1	1	-	
5-9 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
15-19 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-	
20-24 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-	
25-29 years	15	4	11	11	3	8	4	1	3	-	-	-	-	-	-	
30-34 years	31	16	15	25	14	11	4	1	3	-	-	-	2	1	1	
35-39 years	123	88	35	96	69	27	22	15	7	-	-	-	5	4	1	
40-44 years	378	218	160	294	175	119	64	35	29	5	3	2	15	5	10	
45-49 years	974	574	400	762	463	299	173	86	87	5	2	3	34	23	11	
50-54 years	1,722	1,068	654	1,381	863	518	296	180	116	3	3	-	42	22	20	
55-59 years	2,776	1,692	1,084	2,276	1,404	872	415	249	166	13	7	6	72	32	40	
60-64 years	3,481	2,022	1,459	2,914	1,718	1,196	478	258	220	21	11	10	68	35	33	
65-69 years	3,887	2,178	1,709	3,288	1,876	1,412	492	249	243	19	9	10	88	44	44	
70-74 years	4,544	2,332	2,212	3,865	2,026	1,839	537	246	291	19	4	15	123	56	67	
75-79 years	5,149	2,395	2,754	4,528	2,150	2,378	504	192	312	14	6	8	103	47	56	
80-84 years	5,048	2,143	2,905	4,498	1,953	2,545	419	142	277	24	4	20	107	44	63	
85-89 years	3,262	1,224	2,038	2,925	1,124	1,801	260	70	190	11	3	8	66	27	39	
90-94 years	1,459	405	1,054	1,283	361	922	143	26	117	4	3	1	29	15	14	
95-99 years	326	74	252	296	70	226	25	3	22	-	-	-	5	1	4	
100 years & over.	39	5	34	28	2	26	8	2	6	-	-	-	3	1	2	
Age not stated	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Malignant neoplasm of other and ill-defined digestive organs (C26) All ages.... 1,196 1,018 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.7 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Intestinal tract, part unspecified (C26.0) All ages..... 155 242 102 140 27 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 1 40-44 years..... 9 45-49 years..... 4 5 5 2 15 50-54 years..... 9 2 1.3 55-59 years..... 16 9 7 11 5 5 6 4 60-64 years..... 25 18 23 17 65-69 vears..... 18 12 15 6 70-74 vears..... 32 15 17 29 15 3 75-79 years..... 46 19 2.7 41 17 2.4 80-84 years..... 36 20 16 34 19 15 2 1 35 27 27 85-89 years..... 8 35 8 22 90-94 years..... 16 2.0 6 6 14 95-99 years..... 9 9 1 1 8 8 100 years & over. 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Spleen (C26.1) All ages.... 17 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 6 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Overlapping lesion of digestive system (C26.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Ill-defined sites within the digestive system (C26.9) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 1 Q 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of nasal cavity and middle ear (C30) All ages.... 47 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 4 3 1 55-59 years..... 3 2 1 60-64 years..... 3 1 2 65-69 vears..... 3 70-74 vears..... 9 5 75-79 years..... 80-84 years..... 4 85-89 years..... 8 2 6 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Nasal cavity (C30.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 3 50-54 years..... 1 55-59 years..... 3 2 1 60-64 years..... 3 1 2 65-69 vears..... 3 70-74 vears..... 9 5 75-79 years..... 80-84 years..... 4 5 2 85-89 years..... 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Middle ear (C30.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of accessory sinuses (C31) All ages.... 197 179 326 172 154 43 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 4 40-44 years..... 17 9 45-49 years..... 22 a 13 17 Ω 4 29 9 26 50-54 years..... 20 3 1.8 55-59 years..... 38 28 10 31 24 5 60-64 years..... 44 27 17 38 24 14 65-69 years..... 39 19 20 34 16 18 70-74 years..... 30 18 12 25 11 75-79 years..... 50 19 31 46 18 28 80-84 years..... 47 22 25 41 2.0 21 3 1 35 22 2 85-89 years..... 13 33 13 20 90-94 years..... 13 5 8 13 5 8 95-99 years..... 2 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Maxillary sinus (C31.0) All ages..... 138 114 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 5 50-54 years..... 3 2 55-59 years..... 14 11 1.0 60-64 years..... 20 13 18 12 2 65-69 vears..... 11 11 70-74 vears..... 10 5 8 75-79 years..... 19 15 11 80-84 years..... 23 8 15 17 11 3 15 2 85-89 years..... 4 11 13 9 7 90-94 years..... 3 4 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Ethmoidal sinus (C31.1) All ages.... 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 65-69 vears..... 2 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Disea	ases,	222			referê bi a			pl. d		•			ian or		. 1 0 /
Tenth Revision, Second	D - 1-1-	All rac	es	D - 1-1-	White		D - 1-1-	Black		American Indian 1/			Pacific Islander 2/		
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
Frontal sinus (C31.2)															
All ages	2	2	_	2	2	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
.0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
0-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise Tenth Revision, Second	ases,	ses, All races			White			Black		Asian or American Indian 1/ Pacific Islander 2/						
Edition, 2004)	Both			Both			Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female	
Sphenoidal sinus (C31.	3)															
All ages	8	1	7	8	1	7	_	_	_	_	_	_	_	_	_	
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
L-4 years	_	-	-	_	_	-	-	-	_	_	-	-	_	-	-	
-9 years	_	-	-	_	_	-	-	-	_	_	-	-	_	-	-	
0-14 years	-	_	-	-	-	-	-	_	-	_	-	-	-	-	-	
5-19 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-	
0-24 years	-	_	-	-	-	-	-	_	-	_	-	-	-	-	-	
5-29 years	_	_	-	_	-	-	-	-	_	-	-	-	-	-	-	
0-34 years	_	_	-	_	-	-	-	-	_	-	-	-	-	-	-	
5-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
0-44 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_	
5-49 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
0-54 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_	
5-59 years	-	_	-	_	_	-	-	-	_	_	-	_	-	-	-	
0-64 years	1	_	1	1	_	1	-	-	_	_	-	_	-	-	-	
5-69 years	1	_	1	1	_	1	-	-	_	_	-	_	-	-	-	
0-74 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_	
5-79 years	2	_	2	2	_	2	_	_	_	_	_	_	_	_	_	
0-84 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
5-89 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
0-94 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_	
95-99 years	_	_	_	_	-	_	_	_	_	_	_	_	_	_	_	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Accessory sinus, unspecified (C31.9) All ages.... 116 188 102 23 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 4 40-44 years..... 10 5 45-49 years..... 14 4 10 10 3 20 50-54 years..... 2.1 6 15 14 55-59 years..... 22 17 19 15 3 5 2 60-64 years..... 21 12 9 17 10 65-69 vears..... 25 12 20 9 11 13 70-74 vears..... 17 11 6 15 6 75-79 years..... 2.7 11 2.7 16 16 80-84 years..... 24 14 10 2.4 14 10 17 17 85-89 years..... 6 11 6 11 90-94 years..... 5 2 3 3 5 2 95-99 years..... 2 1 1 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Female Male Female Male Female sexes Male Female sexes sexes Male sexes sexes Malignant neoplasm of larynx (C32) All ages.... 3,821 2,991 3,123 2,420 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Glottis (C32.0) All ages.... 92 19 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 2 50-54 years..... 3 2 1 55-59 years..... 4 3 1 3 60-64 years..... 11 8 3 65-69 vears..... 8 5 3 70-74 vears..... 17 75-79 years..... 9 7 5 80-84 years..... 18 12 6 15 10 2 1 15 85-89 years..... 15 14 14 1 90-94 years..... 3 3 3 3 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Supraglottis (C32.1) All ages.... 193 133 163 106 28 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 45-49 years..... 1.0 3 a 2 50-54 years..... 11 9 R 3 55-59 years..... 34 27 27 21 6 60-64 years..... 29 23 6 21 16 6 65-69 vears..... 24 15 16 8 70-74 vears..... 22 12 10 19 9 10 75-79 years..... 2.8 2.0 8 2.7 19 80-84 years..... 15 11 13 9 2 85-89 years..... 14 8 6 13 90-94 years..... 2 2 1 1 1 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Subglottis (C32.2) All ages..... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 3 1 65-69 vears..... 4 70-74 vears..... 2 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Laryngeal cartilage (C32.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																
Classification of Dis Tenth Revision, Secon		All rac	10.5		White			Black		Asian or American Indian 1/ Pacific Islander 2/						
Edition, 2004)	Both	AII Iac	.cs	Both	WIIICE		Black Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	
Larynx, unspecified (C32.9)															
All ages	3,521	2,774	747	2,865	2,241	624	596	480	116	18	15	3	42	38	4	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-	
35-39 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-	
10-44 years	54	37	17	38	25	13	14	10	4	1	1	-	1	1	-	
15-49 years	155	126	29	112	90	22	42	35	7	1	1	-	-	-	-	
50-54 years	268	220	48	195	158	37	63	52	11	4	4	-	6	6	-	
55-59 years	437	355	82	335	275	60	97	77	20	1	1	-	4	2	2	
50-64 years	459	363	96	377	297	80	76	61	15	2	2	-	4	3	1	
55-69 years	503	418	85	388	319	69	107	91	16	2	2	-	6	6	-	
70-74 years	490	375	115	410	309	101	74	60	14	-	-	-	6	6	-	
5-79 years	471	360	111	409	309	100	58	47	11	-	-	-	4	4	-	
0-84 years	384	297	87	337	261	76	36	28	8	4	2	2	7	6	1	
5-89 years	200	145	55	176	129	47	21	13	8	1	1	-	2	2	-	
90-94 years	78	60	18	68	53	15	6	4	2	2	1	1	2	2	-	
95-99 years	20	16	4	20	16	4	-	-	-	-	-	-	-	-	-	
100 years & over.	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of trachea (C33) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 5 45-49 years..... 2 3 50-54 years..... 2 2 2 55-59 years..... 10 8 2 60-64 years..... 4 2 2 65-69 vears..... 10 6 70-74 vears..... 12 5 75-79 years..... 9 80-84 years..... 5 3 85-89 years..... 90-94 years..... 2 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

the International Lagrange of Both and lung (C34) Ilignant neoplasm of bronchus and lung (C34) Ilages																
Asign of Diseases, mich Revision, Second Roll Permit Revision, Second Roll Permit Revision, Second Roll Roll Roll Roll Roll Roll Roll Rol	Cause of death (base	d														
## Pacific Intervision, Second intervision, 2004) ## Both Both days													_			
Both daily Both Sexes Male Female Both Sexes Male Female Both Sexes Male Female Sexes Male Fem				referê e			73.11-									
Alignant neoplasm of bronchus and lung (C34) 1 ages 158.600 89.243 69.357 138.759 77.444 61.315 16.485 9.819 6.666 635 339 296 2.721 1.641 1.080 1.08			All rac	ces	D - 1-1-	White		D - +1-	Black			ncan In	dian I/		liic Isla	ander 2/
lignant neoplasm of bronchus and lung (C34) 1 ages 158,600 89,243 69,357 138,759 77,444 61,315 16,485 9,819 6,666 635 339 296 2,721 1,641 1,080 der 1 year 1 1 1 1 1 1			M . 1 .			M. 1.	m1 .		M. 1.	m1.		W . 1 .			34.3.	m1.
1 ages	and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	remale	sexes	Male	remale	sexes	Male	remale
Age 1 years 1	Malignant neoplasm o	f bronchus a	and lung ((234)												
4 years	All ages	158,600	89,243	69,357	138,759	77,444	61,315	16,485	9,819	6,666	635	339	296	2,721	1,641	1,080
19 years	Under 1 year	1	1	-	-	_	-	1	1	-	-	-	-	-	-	-
1-14 years	1-4 years	1	1	-	1	1	_	-	-	-	-	-	-	-	-	-
in 19 years	5-9 years	1	1	-	-	_	_	1	1	-	-	-	-	-	-	-
1-24 years	10-14 years	1	-	1		-	-	-	-	-	-	-	-	1	-	1
3-29 years 43 26 17 29 16 13 10 8 2 4 2 2 2 3 3 4 3 4 5 38 22 11 11 7 7 3 4 4 5 3 4 5 3 8 22 11 11 7 7 3 4 4 5 3 4 5 3 8 3 4 5 3 8 3 2 5 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	15-19 years	11	4	7	7	2	5	-	2	_	1	-	1	-	-	-
112 59 53 83 45 38 22 11 11 7 3 4 4 5 38 22 11 11 7 3 4 4 5 39 years 426 216 210 331 169 162 73 37 36 22 10 12 4 5 4 years 1,597 818 779 1,262 647 615 274 139 135 10 5 5 5 12 27 24 5 4 years 4,539 2,430 2,109 3,568 1,898 1,670 852 471 381 24 10 14 95 51 44 15 9 years 8,062 4,762 3,300 6,363 3,726 2,637 1,479 891 588 39 27 12 181 118 63 15 9 years 13,074 7,804 5,270 10,876 6,392 4,484 1,879 1,220 659 74 37 37 245 155 90 164 years 18,229 10,700 7,529 15,610 9,041 6,569 2,243 1,430 813 88 44 44 288 185 103 16 9 years 22,495 13,176 9,319 19,590 11,357 8,233 2,472 1,558 914 112 63 49 321 198 123 16 179 years 25,365 14,506 10,859 22,378 12,713 9,665 2,435 1,469 966 111 59 52 441 265 176 16 16 16 16 16 16 16 16 16 16 16 16 16	20-24 years	16	6	10	11	4	7	5	2	3	-	-	-	-	-	-
3-39 years	25-29 years	43	26	17	29	16	13	10	8	2	-	-	-	4		2
1,597 818 779 1,262 647 615 274 139 135 10 5 5 51 27 24 24 24 24 24 25 25 25 28 28 464 670 253 417 60 21 39 1 1 1 - 21 13 8 20 years & over. 83 28 55 69 24 45 10 9 24 45 10 33 7 11 1 - 21 13 8 20 years & over. 83 28 55 69 24 45 10 9 24 45 10 33 7 11 1 - 21 13 8	30-34 years	112	59	53		45	38	22	11	11	-	-	-	7	3	4
3-49 years 4,539 2,430 2,109 3,568 1,898 1,670 852 471 381 24 10 14 95 51 44 92 92 92 92 92 92 92 92 92 92 92 92 92	35-39 years	426	216						37	36	-	-	-	22		
8,062 4,762 3,300 6,363 3,726 2,637 1,479 891 588 39 27 12 181 118 63 155 9 years 13,074 7,804 5,270 10,876 6,392 4,484 1,879 1,220 659 74 37 37 245 155 90 164 years 18,229 10,700 7,529 15,610 9,041 6,569 2,243 1,430 813 88 44 44 288 185 103 169 years 22,495 13,176 9,319 19,590 11,357 8,233 2,472 1,558 914 112 63 49 321 198 123 1674 years 25,365 14,506 10,859 22,378 12,713 9,665 2,435 1,469 966 111 59 52 441 265 176 16,79 years 27,061 15,154 11,907 24,441 13,633 10,808 2,115 1,214 901 88 54 34 417 253 164 16,89 years 21,657 11,854 9,803 19,648 10,748 8,900 1,601 870 731 57 27 30 351 209 142 16,89 years 11,381 5,741 5,640 10,454 5,270 5,184 704 346 358 22 8 14 201 117 84 16,994 years 3,692 1,668 2,024 3,367 1,505 1,862 246 125 121 7 3 4 72 35 37 169 years 752 288 464 670 253 417 60 21 39 1 1 1 - 21 13 8 10 years & over. 83 28 55 69 24 45 10 3 7 1 1 1 - 3 3 - 3	40-44 years		818								10					
13,074 7,804 5,270 10,876 6,392 4,484 1,879 1,220 659 74 37 37 245 155 90 1,64 years 18,229 10,700 7,529 15,610 9,041 6,569 2,243 1,430 813 88 44 44 288 185 103 1,65 years 22,495 13,176 9,319 19,590 11,357 8,233 2,472 1,558 914 112 63 49 321 198 123 1,74 years 25,365 14,506 10,859 22,378 12,713 9,665 2,435 1,469 966 111 59 52 441 265 176 1,779 years 27,061 15,154 11,907 24,441 13,633 10,808 2,115 1,214 901 88 54 34 417 253 164 1,684 years 21,657 11,854 9,803 19,648 10,748 8,900 1,601 870 731 57 27 30 351 209 142 1,689 years 11,381 5,741 5,640 10,454 5,270 5,184 704 346 358 22 8 14 201 117 84 1,694 years 3,692 1,668 2,024 3,367 1,505 1,862 246 125 121 7 3 4 72 35 37 1,699 years 752 288 464 670 253 417 60 21 39 1 1 1 - 21 13 8 10 years & over. 83 28 55 69 24 45 10 3 7 1 1 1 - 3 3 - 3	45-49 years		2,430	2,109												
18,229 10,700 7,529 15,610 9,041 6,569 2,243 1,430 813 88 44 44 288 185 103 6-69 years 22,495 13,176 9,319 19,590 11,357 8,233 2,472 1,558 914 112 63 49 321 198 123 1274 years 25,365 14,506 10,859 22,378 12,713 9,665 2,435 1,469 966 111 59 52 441 265 176 1279 years 27,061 15,154 11,907 24,441 13,633 10,808 2,115 1,214 901 88 54 34 417 253 164 128 128 128 128 129 129 129 129 129 129 129 129 129 129	50-54 years															
3-69 years 22,495 13,176 9,319 19,590 11,357 8,233 2,472 1,558 914 112 63 49 321 198 123 124 125 125 125 125 125 125 125 125 125 125	55-59 years															
0-74 years 25,365 14,506 10,859 22,378 12,713 9,665 2,435 1,469 966 111 59 52 441 265 176 1779 years 27,061 15,154 11,907 24,441 13,633 10,808 2,115 1,214 901 88 54 34 417 253 164 16-84 years 21,657 11,854 9,803 19,648 10,748 8,900 1,601 870 731 57 27 30 351 209 142 16-89 years 11,381 5,741 5,640 10,454 5,270 5,184 704 346 358 22 8 14 201 117 84 16-94 years 3,692 1,668 2,024 3,367 1,505 1,862 246 125 121 7 3 4 72 35 37 16-99 years 752 288 464 670 253 417 60 21 39 1 1 - 21 13 8 10 years & over. 83 28 55 69 24 45 10 3 7 1 1 1 - 3 - 3	60-64 years															
5-79 years 27,061 15,154 11,907 24,441 13,633 10,808 2,115 1,214 901 88 54 34 417 253 164 184 years 21,657 11,854 9,803 19,648 10,748 8,900 1,601 870 731 57 27 30 351 209 142 18 14 11,381 5,741 5,640 10,454 5,270 5,184 704 346 358 22 8 14 201 117 84 18 18 18 18 18 18 18 18 18 18 18 18 18	65-69 years															
0-84 years 21,657 11,854 9,803 19,648 10,748 8,900 1,601 870 731 57 27 30 351 209 142 16-89 years 11,381 5,741 5,640 10,454 5,270 5,184 704 346 358 22 8 14 201 117 84 10-94 years 3,692 1,668 2,024 3,367 1,505 1,862 246 125 121 7 3 4 72 35 37 16-99 years 752 288 464 670 253 417 60 21 39 1 1 - 21 13 8 10 years & over. 83 28 55 69 24 45 10 3 7 1 1 1 - 3 - 3	70-74 years	25,365	14,506	10,859	22,378	12,713	9,665	2,435	1,469	966	111		52		265	
5-89 years 11,381 5,741 5,640 10,454 5,270 5,184 704 346 358 22 8 14 201 117 84 5-94 years 3,692 1,668 2,024 3,367 1,505 1,862 246 125 121 7 3 4 72 35 37 5-99 years 752 288 464 670 253 417 60 21 39 1 1 - 21 13 8 10 years & over. 83 28 55 69 24 45 10 3 7 1 1 - 3 - 3	75-79 years				24,441			2,115	1,214							
0-94 years 3,692 1,668 2,024 3,367 1,505 1,862 246 125 121 7 3 4 72 35 37 5-99 years 752 288 464 670 253 417 60 21 39 1 1 - 21 13 8 10 years & over. 83 28 55 69 24 45 10 3 7 1 1 - 3 - 3	80-84 years	21,657	11,854	9,803	19,648	10,748	8,900	1,601	870	731	57	27		351	209	142
752 288 464 670 253 417 60 21 39 1 1 - 21 13 8 8 90 years & over. 83 28 55 69 24 45 10 3 7 1 1 - 3 - 3	85-89 years		5,741	5,640				704				-		201		
00 years & over. 83 28 55 69 24 45 10 3 7 1 1 - 3 - 3	90-94 years	3,692	1,668	2,024	3,367	1,505		246	125	121	7	3	4	72	35	37
	95-99 years	752									_	1	-		13	
re not stated 1 - 1 1 - 1	100 years & over.	83	28	55	69	24	45	10	3	7	1	1	-	3	-	3
	Age not stated	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Main bronchus (C34.0) All ages.... 30 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 50-54 years..... 4 55-59 years..... 1 1 60-64 years..... 3 65-69 vears..... 3 3 70-74 vears..... 2 75-79 years..... 80-84 years..... 6 85-89 years..... 3 3 90-94 years..... 1 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Upper lobe, bronchus or lung (C34.1) All ages.... 292 211 447 253 194 13 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 1 2 40-44 years..... 1 45-49 years..... 10 3 a 3 21 50-54 years..... 10 9 11 16 55-59 years..... 26 19 7 23 2 16 1 60-64 years..... 50 28 22 43 24 19 6 65-69 years..... 67 36 31 57 30 27 70-74 years..... 85 54 31 77 49 28 75-79 years..... 105 44 95 53 42 80-84 years..... 71 41 30 65 36 29 4 3 1 2 2 85-89 years..... 43 2.2 21 41 2.0 21 2 1 90-94 years..... 2.0 9 17 10 1 2 11 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Middle lobe, bronchus or lung (C34.2) All ages..... 12 33 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 4 65-69 vears..... 4 70-74 vears..... 6 75-79 years..... 80-84 years..... 3 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Lower lobe, bronchus or lung (C34.3) All ages.... 123 192 107 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 1 1 1 50-54 years..... 1.0 5 9 55-59 years..... 12 7 11 5 60-64 years..... 27 21 6 24 18 2 65-69 vears..... 21 9 20 8 12 70-74 vears..... 39 16 13 75-79 years..... 42 18 2.4 33 13 2.0 80-84 years..... 30 21 9 26 19 7 1 28 27 85-89 years..... 18 10 18 9 1 1 5 1 90-94 years..... 1 4 1 3 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Overlapping lesion of bronchus and lung (C34.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based	l														
on the International															
Classification of Dis	seases,											As	sian or		
Tenth Revision, Secon	nd	All rad	ces		White			Black		Amer	rican In	dian 1/	Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Bronchus or lung, uns	specified (C	234.9)													
All ages	157,807	88,783	69,024	138,055	77,044	61,011	16,423	9,777	6,646	633	338	295	2,696	1,624	1,072
Under 1 year	1	1	-	_	-	-	1	1	-	_	-	-	-	-	-
1-4 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-9 years	1	1	-	_	-	-	1	1	-	_	-	-	-	-	-
10-14 years	1	-	1	_	-	-	-	-	-	_	-	-	1	-	1
l5-19 years	10	4	6	6	2	4	3	2	1	1	-	1	_	-	-
20-24 years	16	6	10	11	4	7	5	2	3	-	-	-	_	-	-
25-29 years	43	26	17	29	16	13	10	8	2	-	-	-	4	2	2
30-34 years	110	57	53	82	44	38	21	10	11	-	-	-	7	3	4
35-39 years	425	216	209	331	169	162	72	37	35	-	-	-	22	10	12
40-44 years	1,594	817	777	1,259	646	613	274	139	135	10	5	5	51	27	24
45-49 years	4,524	2,421	2,103	3,555	1,891	1,664	850	469	381	24	10	14	95	51	44
50-54 years	8,026	4,742	3,284	6,334	3,712	2,622	1,473	885	588	39	27	12	180	118	62
55-59 years	13,034	7,779	5,255	10,840	6,371	4,469	1,876	1,217	659	74	37	37	244	154	90
60-64 years	18,144	10,644	7,500	15,535	8,992	6,543	2,235	1,425	810	87	43	44	287	184	103
65-69 years	22,399	13,126	9,273	19,505	11,314	8,191	2,465	1,553	912	112	63	49	317	196	121
70-74 years	25,232	14,422	10,810	22,259	12,637	9,622	2,422	1,462	960	111	59	52	440	264	176
75-79 years	26,902	15,067	11,835	24,301	13,559	10,742	2,101	1,204	897	87	54	33	413	250	163
80-84 years	21,542	11,783	9,759	19,545	10,686	8,859	1,596	867	729	57	27	30	344	203	141
85-89 years	11,304	5,699	5,605	10,381	5,230	5,151	703	346	357	22	8	14	198	115	83
90-94 years	3,665	1,655	2,010	3,344	1,493	1,851	245	125	120	7	3	4	69	34	35
95-99 years	749	288	461	667	253	414	60	21	39	1	1	-	21	13	8
100 years & over.	83	28	55	69	24	45	10	3	7	1	1	-	3	-	3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of thymus (C37) All ages.... 149 71 29 13 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 1 1 20-24 years..... 1 25-29 years..... 30-34 years..... 1 35-39 years..... 9 40-44 years..... 11 3 45-49 years..... 13 4 9 a 2 1 50-54 years..... 19 10 12 14 1 55-59 years..... 20 5 18 14 4 15 60-64 years..... 19 11 8 16 11 65-69 vears..... 23 10 18 11 13 70-74 vears..... 16 9 75-79 years..... 25 13 3 16 80-84 years..... 14 7 13 6 1 85-89 years..... 12 3 9 12 9 90-94 years..... 1 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of heart, mediastinum, and pleura (C38) All ages.... 110 101 27 13 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 9 40-44 years..... 9 2 45-49 years..... 16 a 12 15 10 5 50-54 years..... 5 1.0 2 3 55-59 years..... 14 10 11 1 4 3 60-64 years..... 18 9 9 16 8 65-69 vears..... 18 4 15 12 70-74 vears..... 31 13 18 24 16 75-79 years..... 36 18 18 35 18 1 80-84 years..... 29 16 13 25 14 11 22 85-89 years..... 24 13 11 13 9 90-94 years..... 16 3 13 15 12 1 3 95-99 years..... 4 4 3 3 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Heart (C38.0) All ages.... 87 39 43 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 10 5 8 3 50-54 years..... 55-59 years..... 4 2 2 60-64 years..... 3 65-69 vears..... 9 9 70-74 vears..... 7 5 75-79 years..... 11 80-84 years..... 6 2 2 2 85-89 years..... 8 6 90-94 years..... 4 4 95-99 years..... 2 2 1 1 1 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International															
On the International Classification of Dis	eageg											Δα	ian or		
Tenth Revision, Second		All rac	es		White			Black		Amer	ican Indi			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	remale	sexes	Male	Female
Mediastinum, part uns	pecified (C3	8.3)													
All ages	98	52	46	82	45	37	10	2	8	1	1	_	5	4	1
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	_	-	_	-	-	_	-	_	_	_	-	-	_	-
5-9 years	-	_	-	_	-	-	_	-	_	_	_	-	-	_	-
10-14 years	1	-	1	-	_	-	-	-	-	-	_	-	1	-	1
5-19 years	2	2	-	1	1	-	-	-	-	-	_	-	1	1	-
0-24 years	2	1	1	2	1	1	-	-	-	-	_	-	-	-	-
25-29 years	1	1	-	1	1	-	-	-	-	-	_	-	-	-	-
30-34 years	3	3	-	3	3	-	-	-	-	-	_	-	-	-	-
35-39 years	2	2	_	2	2	_	_	_	_	_	_	_	_	_	_
10-44 years	4	4	_	4	4	_	_	_	_	_	_	_	_	_	_
5-49 years	5	3	2	4	2	2	_	_	_	1	1	_	_	_	_
0-54 years	5	3	2	3	3	_	2	_	2	_	-	_	-	_	-
55-59 years	8	6	2	6	5	1	1	-	1	_	-	-	1	1	-
0-64 years	9	4	5	8	4	4	1	-	1	_	-	-	-	-	-
55-69 years	6	3	3	5	2	3	1	1	_	_	-	-	-	-	-
70-74 years	12	4	8	9	2	7	2	1	1	_	_	_	1	1	_
5-79 years	16	10	6	15	9	6	_	_	_	_	_	_	1	1	_
0-84 years	9	3	6	7	3	4	2	_	2	_	_	_	_	_	_
5-89 years	6	2	4	6	2	4	_	_	_	_	_	_	_	_	_
0-94 years	7	1	6	6	1	5	1	_	1	_	_	_	_	_	_
5-99 years	_	_	_	_	_	_	_	_	_	_	-	_	-	_	-
.00 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pleura (C38.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 2 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 2 2 65-69 vears..... 3 70-74 vears..... 12 5 75-79 years..... 9 9 80-84 years..... 14 9 5 13 85-89 years..... 10 9 1 9 90-94 years..... 5 2 3 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Bot.h Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of other and ill-defined sites in the respiratory system and intrathoracic organs (C39) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 1 1 50-54 years..... 3 55-59 years..... 3 1 60-64 years..... 8 65-69 vears..... 13 6 70-74 vears..... 3 75-79 years..... 9 80-84 years..... 8 6 2 2 85-89 years..... 3 2 90-94 years..... 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Upper respiratory tract, part unspecified (C39.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Overlapping lesion of respiratory and intrathoracic organs (C39.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Ill-defined sites within the respiratory system (C39.9) All ages.... 47 30 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 6 50-54 years..... 55-59 years..... 3 60-64 years..... 6 5 65-69 vears..... 12 6 6 70-74 vears..... 3 75-79 years..... 80-84 years..... 2 2 3 85-89 years..... 90-94 years..... 2 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of bone and articular cartilage of limbs (C40) All ages.... 30 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 5 3 2 1 15-19 years..... 20-24 years..... 12 10 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 1 50-54 years..... 3 1 55-59 years..... 6 2 4 6 60-64 years..... 9 6 3 65-69 vears..... 2 70-74 vears..... 8 75-79 years..... 10 80-84 years..... 8 85-89 years..... 6 5 90-94 years..... 3 1 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Scapula and long bones of upper limb (C40.0) All ages.... 19 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 3 75-79 years..... 2 80-84 years..... 2 85-89 years..... 90-94 years..... 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Long bones of lower limb (C40.2) All ages..... 37 25 30 20 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 3 2 1 55-59 years..... 6 2 2 4 60-64 years..... 4 3 65-69 vears..... 70-74 vears..... 5 75-79 years..... 6 3 80-84 years..... 6 2 85-89 years..... 90-94 years..... 1 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Short bones of lower limb (C40.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Bone and articular cartilage of limb, unspecified (C40.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Bot.h Both Bot.h Both Bot.h and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Malignant neoplasm of bone and articular cartilage of other and unspecified sites (C41) All ages.... 1,249 1,054 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 3.0 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Bones of skull and face (C41.0) All ages..... 19 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 45-49 years..... 1 50-54 years..... 3 1 55-59 years..... 1 1 9 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 4 3 90-94 years..... 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mandible (C41.1) All ages.... 105 47 37 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 2 2 3 50-54 years..... 4 3 3 1 55-59 years..... 8 2 1 5 60-64 years..... 5 1 3 65-69 vears..... 8 3 5 3 5 70-74 vears..... 14 5 75-79 years..... 13 5 12 80-84 years..... 20 13 19 12 1 5 2 85-89 years..... 14 7 11 1 90-94 years..... 6 2 4 6 95-99 years..... 2 2 2 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis					1.					_			sian or		
Tenth Revision, Secon		All rac	es	D - 1-1-	White		D - 1-1-	Black			ican Ind	lian 1/		ic Isla	nder 2/
Edition, 2004) and age	Both	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Fomalo	Both sexes	Mala	Female
	sexes	mare	remaie	sexes	мате	remare	sexes	мате	remare	sexes	Male	reliate	sexes	Male	remaie
Vertebral column (C41	.2)														
All ages	108	57	51	94	50	44	9	4	5	3	2	1	2	1	1
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	1	1	-	1	1	-	-	-	-	-	_	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
10-14 years	4	2	2	3	1	2	1	1	-	-	_	-	-	-	-
15-19 years	1	-	1	1	-	1	-	-	-	-	_	-	-	-	-
20-24 years	3	2	1	2	1	1	1	1	-	-	_	-	-	-	-
25-29 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-34 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
35-39 years	3	1	2	3	1	2	-	-	-	-	_	-	-	-	-
40-44 years	2	1	1	1	1	-	1	-	1	-	_	-	-	-	-
45-49 years	6	4	2	4	3	1	1	1	-	1	-	1	-	-	-
50-54 years	5	4	1	5	4	1	-	-	-	-	-	-	-	-	-
55-59 years	5	5	-	4	4	-	-	-	-	1	1	-	-	-	-
60-64 years	8	5	3	7	4	3	1	1	-	-	-	-	-	-	-
65-69 years	8	4	4	7	4	3	1	-	1	-	-	-	-	-	-
70-74 years	16	7	9	13	6	7	1	-	1	-	-	-	2	1	1
75-79 years	10	6	4	10	6	4	-	-	-	-	-	-	-	-	-
80-84 years	13	5	8	12	5	7	1	-	1	-	-	-	-	-	-
85-89 years	8	3	5	8	3	5	-	-	-	-	-	-	-	-	-
90-94 years	9	2	7	9	2	7	-	-	-	-	-	-	-	-	-
95-99 years	3	2	1	1	1	-	1	-	1	1	1	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	-	-	-	-	-	-	_	-	_	_	_	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Ribs, sternum, and clavicle (C41.3) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 3 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Pelvic bones, sacrum, and coccyx (C41.4) All ages.... 29 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 45-49 years..... 1 50-54 years..... 4 3 1 55-59 years..... 2 2 60-64 years..... 5 3 2 65-69 vears..... 5 70-74 vears..... 6 75-79 years..... 4 3 80-84 years..... 6 5 85-89 years..... 3 90-94 years..... 2 1 1 95-99 years..... 1 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Bone and articular cartilage, unspecified (C41.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 2.7 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 2.0 35-39 years..... 40-44 years..... 45-49 years..... 2.2 50-54 years..... 2.0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.2 2.7 80-84 years..... 85-89 years..... 90-94 years..... 1.0 3.0 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International															
Classification of Dis													ian or		
Tenth Revision, Secon		All rac	es		White			Black			rican In	dian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Malignant melanoma of	skin (C43)														
All ages	8,441	5,477	2,964	8,250	5,381	2,869	126	66	60	21	12	9	44	18	26
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	2	1	1	-	-	-	1	1	-	-	-	-	1	-	1
15-19 years	11	7	4	10	7	3	_	-	-	-	-	-	1	-	1
20-24 years	36	21	15	35	20	15	1	1	-	-	-	-	-	-	-
25-29 years	56	37	19	54	36	18	1	-	1	-	-	-	1	1	-
30-34 years	131	74	57	129	72	57	2	2	-	-	-	-	-	-	-
35-39 years	191	124	67	185	121	64	4	2	2	1	1	-	1	_	1
40-44 years	314	184	130	301	179	122	7	3	4	2	_	2	4	2	2
45-49 years	510	311	199	506	309	197	3	1	2	_	_	-	1	1	-
50-54 years	658	434	224	642	426	216	9	6	3	2	_	2	5	2	3
55-59 years	818	546	272	800	539	261	13	6	7	1	-	1	4	1	3
60-64 years	824	576	248	805	560	245	13	11	2	3	3	-	3	2	1
65-69 years	855	578	277	843	573	270	5	2	3	1	1	-	6	2	4
70-74 years	925	608	317	908	601	307	12	4	8	2	1	1	3	2	1
75-79 years	1,034	703	331	1,010	689	321	16	11	5	2	1	1	6	2	4
80-84 years	1,007	652	355	985	643	342	18	6	12	2	2	-	2	1	1
85-89 years	710	437	273	694	429	265	7	4	3	4	3	1	5	1	4
90-94 years	279	153	126	266	147	119	12	5	7	_	_	-	1	1	-
95-99 years	64	27	37	61	26	35	2	1	1	1	_	1	_	_	-
100 years & over.	14	2	12	14	2	12	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Malignant melanoma of lip (C43.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant melanoma of eyelid, including canthus (C43.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant melanoma of ear and external auricular canal (C43.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 55-59 years..... 3 3 60-64 years..... 1 1 65-69 vears..... 3 3 70-74 vears..... 2 75-79 years..... 3 80-84 years..... 6 85-89 years..... 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis													ian or		
Tenth Revision, Secon		All races			White		Black				rican Indi	an 1/	Pacific Islander 2		
Edition, 2004)	Both	Male	Eamala.	Both	Male	E1-	Both	Male	Esmal s	Both	Male F		Both	Mala	Female
and age	sexes	мате	Female	sexes	мате	Female	sexes	мате	Female	sexes	Maie F	emaie	sexes	мате	remaie
Malignant melanoma of	other and u	nspecifie	d parts of f	Tace (C43.3)											
All ages	87	62	25	86	62	24	_	_	_	_	_	_	1	-	1
Jnder 1 year	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	5	4	1	5	4	1	-	-	-	-	-	-	-	-	-
50-54 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
55-59 years	5	4	1	5	4	1	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
70-74 years	7	6	1	7	6	1	-	-	-	-	-	-	-	-	-
75-79 years	21	14	7	21	14	7	-	-	-	-	-	-	-	-	-
30-84 years	14	11	3	14	11	3	_	-	-	-	-	-	-	-	-
85-89 years	18	15	3	17	15	2	_	-	-	-	-	-	1	-	1
90-94 years	7	2	5	7	2	5	-	-	-	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	_	-	-	-	-	-	-	-	-
100 years & over.	2	_	2	2	-	2	-	_	-	-	-	-	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant melanoma of scalp and neck (C43.4) All ages.... 121 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 6 1 9 50-54 years..... 6 3 55-59 years..... 5 5 5 5 60-64 years..... 8 8 1 65-69 vears..... 14 10 70-74 vears..... 13 10 13 10 75-79 years..... 15 11 15 11 80-84 years..... 16 15 15 14 15 7 85-89 years..... 8 15 90-94 years..... 6 5 1 6 5 1 95-99 years..... 1 1 1 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant melanoma of trunk (C43.5) All ages..... 115 164 114 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 1 25-29 years..... 30-34 years..... 1 35-39 years..... 9 40-44 years..... 3 9 45-49 years..... 17 1.0 17 50-54 years..... 19 15 19 15 55-59 years..... 24 19 24 19 5 60-64 years..... 15 14 1 15 14 65-69 vears..... 12 12 8 70-74 vears..... 17 13 17 13 75-79 years..... 15 9 15 9 6 80-84 years..... 15 11 15 11 10 85-89 years..... 4 6 10 90-94 years..... 5 2 3 4 3 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant melanoma of upper limb, including shoulder (C43.6) All ages.... 40 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 8 3 5 2 2 50-54 years..... 1 2 55-59 years..... 14 10 10 14 60-64 years..... 13 9 13 65-69 vears..... 9 5 9 70-74 vears..... 13 10 3 13 75-79 years..... 13 10 13 1.0 80-84 years..... 10 8 2 1.0 8 5 85-89 years..... 12 11 7 90-94 years..... 5 2 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant melanoma of lower limb, including hip (C43.7) All ages.... 105 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 8 8 4 60-64 years..... 8 65-69 vears..... 15 8 70-74 vears..... 5 5 75-79 years..... 11 80-84 years..... 17 6 11 17 11 7 85-89 years..... 13 6 12 6 90-94 years..... 3 4 1 2 2 95-99 years..... 3 2 3 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Malignant melanoma of skin, unspecified (C43.9) All ages.... 7,808 5,058 2,750 7,629 4,971 2,658 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1.0 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other malignant neoplasms of skin (C44) All ages.... 2,627 2,453 1,746 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 2.4 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Skin of lip (C44.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Skin of eyelid, including canthus (C44.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Skin of ear and external auricular canal (C44.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 2 1 55-59 years..... 4 4 60-64 years..... 3 1 65-69 vears..... 6 5 70-74 vears..... 4 75-79 years..... 13 13 80-84 years..... 21 18 21 18 18 85-89 years..... 14 16 13 90-94 years..... 14 9 5 14 9 95-99 years..... 5 5 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Skin of other and unspecified parts of face (C44.3) All ages.... 147 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 4 4 5 50-54 years..... 3 3 5 55-59 years..... 15 14 13 14 1 60-64 years..... 15 14 1 15 14 65-69 vears..... 15 10 5 15 10 70-74 vears..... 17 14 3 17 3 75-79 years..... 31 19 12 31 19 12 80-84 years..... 35 23 12 34 23 11 35 34 20 85-89 years..... 2.0 15 14 30 90-94 years..... 2.0 1.0 3.0 2.0 1.0 95-99 years..... 16 2 14 16 14 2 100 years & over. 6 3 3 3 6 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Skin of scalp and neck (C44.4) All ages..... 1,096 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.8 2.7 80-84 years..... 85-89 years..... 90-94 years..... 5.8 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Skin of trunk (C44.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 3 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 5 3 2 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 80-84 years..... 9 5 6 85-89 years..... 2 90-94 years..... 10 9 8 1 1 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis												Δο	sian or		
Tenth Revision, Secon			All races		White			Black			rican Ind		Pacific Islander 2/		
Edition, 2004)	Both	1111 140		Both	,,,,,,		Both	214011		Both	10011 1110		Both	10 1010	ilaci 2,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Skin of upper limb, i	ncluding sho	oulder (C4	4.6)												
All ages	36	19	17	34	19	15	1	_	1	1	_	1	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	_	-	-	-	-	-	-	-	_	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
65-69 years	4	3	1	4	3	1	-	-	-	-	-	-	-	-	-
70-74 years	2	1	1	2	1	1	-	-	-	_	-	-	-	-	-
75-79 years	3	3	-	3	3	-	-	-	-	_	-	-	-	-	-
30-84 years	5	3	2	5	3	2	-	-	-	-	-	-	_	-	-
85-89 years	10	4	6	9	4	5	1	-	1	-	-	-	_	-	-
90-94 years	4	1	3	4	1	3	-	-	-	-	-	-	_	-	-
95-99 years	4	1	3	3	1	2	-	-	-	1	-	1	_	-	-
100 years & over.	1	_	1	1	-	1	-	-	-	_	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based															
on the International															
Classification of Dis	eases,											As	ian or		
Tenth Revision, Second			es		White		Black			Amer	ican Indi	an 1/	Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Skin of lower limb, i		(044.7)													
Skill of lower limb, i	neruarng mp	(044.7)													
All ages	48	15	33	48	15	33	-	-	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
50-54 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
55-59 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
60-64 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-
65-69 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
70-74 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
75-79 years	6	3	3	6	3	3	-	-	-	-	-	-	-	-	-
80-84 years	7	4	3	7	4	3	-	-	-	-	-	-	-	-	-
85-89 years	11	2	9	11	2	9	-	-	-	-	-	-	-	-	-
90-94 years	9	-	9	9	-	9	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of skin, unspecified (C44.9) All ages.... 1,068 298 1,025 289 32 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 7 40-44 years..... 45-49 years..... 2.0 16 16 12 29 26 2.0 50-54 years..... 2.3 3 6 55-59 years..... 59 49 10 53 44 9 6 5 60-64 years..... 78 60 18 73 55 18 65-69 years..... 79 13 77 65 12 66 70-74 years..... 107 83 24 102 79 23 75-79 years..... 154 116 38 150 114 36 80-84 years..... 188 137 51 179 131 48 2 195 193 130 85-89 years..... 132 63 63 1 90-94 years..... 92 53 39 92 53 39 95-99 years..... 27 44 45 18 18 26 1 1 100 years & over. 2 5 2 5

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																			
Classification of Dise		es, All races										Asian or							
Tenth Revision, Second		All rac	es		White			Black		American Indian 1/			Pacific Islander 2,						
Edition, 2004)	Both			Both			Both			Both			Both						
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female				
Mesothelioma (C45)																			
All ages	2,452	1,973	479	2,310	1,866	444	121	90	31	8	7	1	13	10	3				
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
15-19 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-				
20-24 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-				
25-29 years	5	3	2	4	2	2	1	1	-	-	-	-	-	-	-				
30-34 years	6	4	2	6	4	2	-	-	-	-	-	-	-	-	-				
35-39 years	8	4	4	6	3	3	2	1	1	-	-	-	-	-	-				
40-44 years	31	18	13	30	17	13	1	1	-	-	-	-	-	-	-				
45-49 years	39	25	14	35	23	12	4	2	2	-	-	-	-	-	-				
50-54 years	63	37	26	51	32	19	8	3	5	1	1	-	3	1	2				
55-59 years	119	89	30	108	80	28	8	6	2	-	-	-	3	3	-				
60-64 years	214	180	34	198	170	28	14	9	5	1	1	-	1	-	1				
65-69 years	301	246	55	287	237	50	13	8	5	1	1	-	-	-	-				
70-74 years	404	331	73	379	311	68	23	18	5	1	1	-	1	1	-				
75-79 years	470	408	62	446	387	59	17	15	2	4	3	1	3	3	-				
80-84 years	471	393	78	451	375	76	19	17	2	-	-	-	1	1	-				
85-89 years	256	195	61	247	187	60	9	8	1	-	-	-	-	-	-				
90-94 years	57	34	23	54	32	22	2	1	1	-	-	-	1	1	-				
95-99 years	6	5	1	6	5	1	-	-	-	-	-	-	-	-	-				
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
Age not stated	_	-	-	_	-	-	-	_	-	-	-	-	-	-	-				

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mesothelioma of pleura (C45.0) All ages..... 137 132 36 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 3 5 50-54 years..... 5 55-59 years..... 13 8 13 5 60-64 years..... 14 10 11 65-69 vears..... 27 20 27 70-74 vears..... 26 21 5 19 75-79 years..... 35 30 35 30 80-84 years..... 33 30 3 32 29 13 85-89 years..... 9 4 12 8 1 90-94 years..... 3 3 3 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mesothelioma of peritoneum (C45.1) All ages..... 32 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 5 6 50-54 years..... 6 55-59 years..... 6 2 4 5 60-64 years..... 5 1 65-69 vears..... 10 9 70-74 vears..... 14 13 75-79 years..... 14 8 6 80-84 years..... 6 4 2 2 85-89 years..... 3 1 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Mesothelioma of pericardium (C45.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mesothelioma of other sites (C45.7) All ages.... 242 289 226 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 5 8 50-54 years..... 5 3 55-59 years..... 8 1 6 5 60-64 years..... 25 23 2 22 20 2 65-69 vears..... 25 17 8 17 7 70-74 vears..... 58 43 15 55 14 75-79 years..... 58 48 10 54 45 9 3 80-84 years..... 62 49 13 61 48 13 35 85-89 years..... 28 7 34 2.7 1 90-94 years..... 8 3 5 5 8 3 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Mesothelioma, unspecified (C45.9) All ages..... 1,886 1,548 338 1,780 1,467 313 92 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 2 35-39 years..... 2 9 40-44 years..... 19 10 18 9 45-49 years..... 22 16 6 21 16 5 1 35 25 8 50-54 years..... 44 29 15 1 0 55-59 years..... 92 72 20 84 65 19 6 60-64 years..... 170 143 27 160 137 23 8 5 3 1 65-69 years..... 239 200 39 227 192 35 11 70-74 years..... 306 255 51 287 240 47 17 13 75-79 years..... 363 322 41 346 307 39 12 11 80-84 years..... 369 309 60 351 293 58 17 15 1 205 156 198 150 7 85-89 years..... 49 48 6 1 1 90-94 years..... 45 2.7 18 42 25 17 2 1 1 95-99 years..... 4 3 3 1 4 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Kaposi's sarcoma (C46) All ages.... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 90-94 years..... 3 1 95-99 years..... 2 1 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Kaposi's sarcoma of other sites (C46.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Kaposi's sarcoma, unspecified (C46.9) All ages..... 30 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 90-94 years..... 8 2 1 95-99 years..... 2 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of peripheral nerves and autonomic nervous system (C47) All ages.... 29 12 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 5 3 5 3 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 9 50-54 years..... 6 3 1 55-59 years..... 10 6 1 60-64 years..... 2 1 1 65-69 vears..... 3 70-74 vears..... 3 75-79 years..... 2 80-84 years..... 3 2 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Peripheral nerves of head, face, and neck (C47.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Classification of Dis													ian or		
Tenth Revision, Secon		All races		White			Black			American Indian 1/			Pacific Islander 2		
Edition, 2004)	Both		- 1	Both		_ 1	Both		_ 1	Both		_ 1	Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Peripheral nerves of	upper limb,	including	shoulder (247.1)											
All ages	3	2	1	3	2	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
60-64 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	1	1	-	1	1	-	_	-	-	_	_	_	-	_	-
85-89 years	1	1	-	1	1	-	_	-	-	_	_	_	-	_	-
90-94 years	_	_	-	-	-	-	_	-	-	_	_	_	-	_	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	-	-	-	-	_	-	-	_	_	_	-	_	-
Age not stated	_			_								_			_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Peripheral nerves of lower limb, including hip (C47.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Peripheral nerves of thorax (C47.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Peripheral nerves of pelvis (C47.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Peripheral nerves of trunk, unspecified (C47.6) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Peripheral nerves and autonomic nervous system, unspecified (C47.9) All ages.... 12 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 3 2 3 2 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 6 45-49 years..... 2 9 50-54 years..... 6 3 1 55-59 years..... 10 6 1 60-64 years..... 2 1 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of retroperitoneum and peritoneum (C48) All ages.... 140 677 12 11 Under 1 year..... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 4 40-44 years..... 11 10 45-49 years..... 17 a 8 15 Ω 7 2 1 49 8 50-54 years..... 58 2.3 35 20 29 3 55-59 years..... 81 20 61 76 18 58 4 60-64 years..... 83 14 69 74 13 61 1 65-69 years..... 111 21 90 106 20 70-74 years..... 124 14 110 114 13 101 75-79 years..... 148 16 132 136 15 121 80-84 years..... 143 15 128 135 15 120 5 5 72 65 85-89 years..... 66 59 4 90-94 years..... 2.4 3 21 2.4 21 3 95-99 years..... 4 4 4 4 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Retroperitoneum (C48.0) All ages.... 190 172 17 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 9 6 3 Ω 5 15 3 50-54 years..... 14 1 12 12 55-59 years..... 23 12 21 11 10 11 60-64 years..... 15 8 7 14 8 6 1 65-69 vears..... 24 13 24 13 70-74 vears..... 19 6 13 16 10 75-79 years..... 2.3 13 10 22 10 80-84 years..... 32 13 19 29 13 16 3 85-89 years..... 11 2 9 10 2 8 1 1 90-94 years..... 8 3 5 8 3 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Specified parts of peritoneum (C48.1) All ages..... 29 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 2 50-54 years..... 2 55-59 years..... 2 2 60-64 years..... 1 1 65-69 vears..... 6 70-74 vears..... 8 3 75-79 years..... 10 10 80-84 years..... 7 6 5 85-89 years..... 5 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Peritoneum, unspecified (C48.2) All ages..... 612 605 569 32 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 1 40-44 years..... 7 45-49 years..... 2 5 6 4 1 50-54 years..... 41 36 29 34 55-59 years..... 56 6 50 53 48 3 60-64 years..... 67 5 62 59 55 1 65-69 vears..... 80 78 76 74 70-74 vears..... 97 5 92 92 88 75-79 years..... 115 112 105 102 80-84 years..... 104 103 99 98 2 5 51 52 47 2 85-89 years..... 56 90-94 years..... 15 15 15 15 95-99 years..... 3 3 3 3 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Disc	eases,											As	ian or		
Tenth Revision, Second	d	All rac	es		White			Black		Amer	ican Ind	ian 1/	Pacif	fic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Malignant neoplasm of	other conne	ective and	soft tissu	e (C49)											
All ages	3,785	1,928	1,857	3,271	1,701	1,570	398	172	226	20	8	12	96	47	49
Under 1 year	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
1-4 years	22	13	9	16	10	6	4	2	2	-	-	-	2	1	1
5-9 years	30	21	9	24	17	7	5	3	2	-	-	-	1	1	-
10-14 years	20	12	8	15	9	6	5	3	2	-	-	-	-	-	-
15-19 years	52	31	21	42	25	17	9	5	4	-	-	-	1	1	-
20-24 years	55	33	22	43	28	15	11	4	7	1	1	-	-	-	-
25-29 years	75	42	33	55	31	24	13	8	5	3	1	2	4	2	2
30-34 years	80	51	29	61	37	24	14	10	4	2	2	-	3	2	1
35-39 years	82	43	39	72	38	34	8	4	4	-	-	-	2	1	1
40-44 years	177	91	86	140	70	70	31	17	14	2	1	1	4	3	1
45-49 years	228	114	114	179	97	82	42	14	28	_	_	_	7	3	4
50-54 years	289	133	156	242	113	129	33	15	18	3	1	2	11	4	7
55-59 years	344	165	179	284	147	137	48	16	32	1	-	1	11	2	9
60-64 years	355	172	183	313	152	161	35	18	17	1	-	1	6	2	4
65-69 years	353	203	150	300	180	120	36	14	22	2	1	1	15	8	7
70-74 years	384	184	200	340	171	169	40	10	30	-	-	_	4	3	1
75-79 years	422	224	198	378	200	178	29	16	13	2	-	2	13	8	5
80-84 years	417	212	205	393	200	193	16	7	9	1	1	_	7	4	3
85-89 years	264	122	142	246	116	130	13	5	8	1	-	1	4	1	3
90-94 years	110	54	56	105	52	53	4	1	3	-	-	_	1	1	-
95-99 years	18	5	13	16	5	11	2	_	2	-	-	_	-	_	-
100 years & over.	7	2	5	6	2	4	_	_	_	1	_	1	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Connective and soft tissue of head, face, and neck (C49.0) All ages.... 37 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 2 2 2 50-54 years..... 4 3 1 55-59 years..... 6 2 6 60-64 years..... 5 3 2 65-69 vears..... 8 70-74 vears..... 13 8 5 75-79 years..... 12 1.0 80-84 years..... 13 12 11 11 7 85-89 years..... 13 6 13 90-94 years..... 1.0 5 5 1.0 5 95-99 years..... 3 2 3 1 1 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Connective and soft tissue of upper limb, including shoulder (C49.1) All ages.... 12 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 2 3 50-54 years..... 3 55-59 years..... 2 1 1 60-64 years..... 8 1 65-69 vears..... 4 70-74 vears..... 5 75-79 years..... 5 80-84 years..... 10 2 85-89 years..... 4 2 90-94 years..... 3 3 3 3 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Connective and soft tissue of lower limb, including hip (C49.2) All ages.... 86 29 12 11 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 4 40-44 years..... 8 9 45-49 years..... 8 1 6 2 1 2 50-54 years..... 11 9 2 9 55-59 years..... 14 7 11 2 1 60-64 years..... 17 10 13 3 65-69 vears..... 16 9 13 70-74 vears..... 20 9 16 9 75-79 years..... 26 11 2.0 15 80-84 years..... 25 10 15 23 9 14 1 26 7 25 85-89 years..... 19 19 90-94 years..... 1.0 3 7 1 1 a 3 6 95-99 years..... 2 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dis	eases.											As	ian or		
Tenth Revision, Secon		All rac	es		White			Black		Amer	ican Indi			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Connective and soft t	issue of tho	rax (C49.	3)												
All ages	89	49	40	80	40	40	6	6	_	_	_	_	3	3	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	_	-	-	_	-	_	-	-	_	-	-	-	-	-	-
15-19 years	_	-	-	_	-	_	-	-	_	-	-	-	-	-	-
20-24 years	1	1	-	1	1	_	-	-	_	-	-	-	-	-	-
25-29 years	2	1	1	1	-	1	1	1	_	-	-	-	-	-	-
30-34 years	_	-	-	_	-	_	-	-	_	-	-	-	-	-	-
35-39 years	_	_	-	_	-	_	-	-	_	_	-	-	-	-	-
10-44 years	2	1	1	2	1	1	-	-	_	_	-	-	-	-	-
5-49 years	5	3	2	4	2	2	1	1	_	_	-	_	-	-	-
0-54 years	4	1	3	4	1	3	-	-	_	_	-	_	-	-	-
55-59 years	6	4	2	5	3	2	1	1	_	_	-	_	-	-	-
60-64 years	7	4	3	7	4	3	-	-	-	-	-	-	-	-	-
55-69 years	7	5	2	6	4	2	-	-	-	-	-	-	1	1	-
70-74 years	10	5	5	7	2	5	1	1	_	_	-	_	2	2	-
5-79 years	14	8	6	13	7	6	1	1	_	_	-	_	-	-	-
0-84 years	14	7	7	13	6	7	1	1	_	_	-	-	-	_	-
35-89 years	10	7	3	10	7	3	-	_	_	_	-	-	-	_	-
90-94 years	4	1	3	4	1	3	-	_	_	_	-	-	-	_	-
95-99 years	3	1	2	3	1	2	-	_	_	_	-	-	-	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Connective and soft tissue of abdomen (C49.4) All ages.... 117 231 132 99 28 12 10 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 1 25-29 years..... 30-34 years..... 35-39 years..... 1 5 40-44 years..... 2 45-49 years..... 10 4 6 Ω 17 3 50-54 years..... 6 9 3 3 11 55-59 years..... 30 7 23 19 5 23 4 2 1 60-64 years..... 32 20 12 30 19 11 1 65-69 years..... 27 18 9 23 7 70-74 vears..... 26 10 16 10 14 75-79 years..... 36 21 15 30 16 14 80-84 years..... 43 26 17 40 2.4 16 3 2 1 2 85-89 years..... 2.0 9 11 18 8 10 1 90-94 years..... 13 5 8 13 8 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Connective and soft tissue of pelvis (C49.5) All ages.... 116 68 10 Under 1 year.... 1-4 years..... 1 1 5-9 years..... 10-14 years..... 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 6 50-54 years..... 2 3 55-59 years..... 4 3 5 60-64 years..... 6 1 5 5 65-69 vears..... 18 8 7 10 3 70-74 vears..... 18 11 17 75-79 years..... 2.3 15 19 12 80-84 years..... 21 14 21 14 14 85-89 years..... 8 11 5 90-94 years..... 3 2 1 3 1 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Classification of Dise		211			rale de la			D1l-		3			sian or	(4 m = m.)	1 O /
Tenth Revision, Second	ı Both	All race	es	Both	White		Both	Black		Amer Both	ican Indi	an I/	Pacii Both	ic Isla	nder 2/
Edition, 2004) and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Connective and soft ti	ssue of tru	ınk, unspe	cified (C49.	6)											
All ages	16	9	7	14	8	6	1	-	1	_	-	-	1	1	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	_	_	_	_	_	_	-	-	-	-	-	-	-	-	-
50-54 years	2	1	1	2	1	1	_	-	-	-	_	-	-	-	-
55-59 years	1	-	1	1	_	1	_	-	-	_	-	-	-	-	-
60-64 years	- 5	-	- 1	_	- 3	- 1	-	-	-	_	-	_	- 1	- 1	-
65-69 years	5	4	1	4	-	1	-	_	-	-	_		1	1	-
70-74 years	1	- 1	1	1	- 1	1 1	-	-	-	_	-	-	_	-	-
75-79 years	2	_	1	2	1	_	_	_	_	_	-	_	_	-	_
80-84 years	1	1	- 1	1	_	- 1	_	_		_	-	_	_	-	_
85-89 years	2	2	1	2	2	1	_	_		-	_	-	_	-	-
90-94 years 95-99 years	۷	2	_	∠ _	2	-	_	_	-	-	_	-	_	-	-
7)=77 VEALS	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Overlapping lesion of connective and soft tissue (C49.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Connective and soft tissue, unspecified (C49.9) All ages.... 2,916 1,469 1,447 2,534 1,313 1,221 Under 1 year.... 1-4 years..... 2.7 5-9 years..... 10-14 years..... 15-19 years..... 2.7 2.3 2.0 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 50-54 years..... 2.7 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

1

1

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Bot.h Both Bot.h Both Bot.h and age Female Female Male Female sexes Male Female sexes Male sexes Male sexes Male Female sexes Malignant neoplasm of breast (C50) All ages.... 41,210 389 40,821 34,464 327 34,137 5,752 58 5,694 160 160 834 830 Under 1 year..... 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 1 1 1 1 20-24 vears..... 12 11 5 5 6 25-29 years..... 74 74 46 46 23 23 30-34 years..... 294 291 199 196 85 85 2 8 9 35-39 years..... 795 794 550 550 211 210 9 25 25 8 40-44 years..... 1,557 5 1,552 1,142 1,138 365 364 8 42 42 1,929 597 45-49 years..... 2,641 17 2,624 1.941 12 602 5 17 17 81 81 647 50-54 years..... 3,474 26 3,448 2,675 2,659 657 10 17 125 125 16 17 4,496 38 4,458 3,578 30 3,548 762 8 754 24 24 132 132 55-59 years..... 60-64 years..... 4,375 50 4,325 3,642 42 3,600 613 6 607 18 18 102 2 100 3,939 3,899 3,347 3,313 494 15 77 77 65-69 years..... 40 500 6 15 70-74 years..... 4.022 46 3,976 3,456 39 3,417 472 465 15 15 79 79 75-79 years..... 4,658 55 4,603 4,118 50 4,068 466 462 10 10 64 63 80-84 years..... 4,688 56 4,632 4,198 49 4.149 442 6 436 10 10 38 1 37 273 33 85-89 years..... 3,375 39 3,336 3,060 38 3,022 274 1 8 8 33 1,976 1,778 179 1.985 9 1.786 178 5 15 15 90-94 years..... 8 1 5 95-99 years..... 702 7 701 610 610 85 84 7 1 1 120 2 118 109 2 107 10 10 1 100 years & over.

1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Nipple and areola (C50.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dis		311			relation			D11-		3			sian or		- 1- 0/
Tenth Revision, Secon		All rac	es	D. 1-1-	White		D - 1-1-	Black			rican Indi	lan 1/		ic isia	nder 2/
Edition, 2004)	Both	Mala	Esmals.	Both	M-1-	Female	Both	Mala	Esmal.	Both	Mala E		Both	M-1-	E1-
and age	sexes	Male	Female	sexes	Male		sexes	Male	Female	sexes	Male F		sexes	Male	Female
Lower-inner quadrant	of breast (0	250.3)													
211	1		1	1		1									
All ages	1	-	1	1	_	1	-	-	-	-	_	_	-	-	-
Under 1 year	-	-			_	-	-	-	-	-	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	_	-	-	_
10-14 years	-	-	-	-	-	-	-	_	-	_	-	_	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	_
20-24 years	-	-	-	-	-	-	-	_	-	-	-	_	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	_	-	-	-
35-39 years	-	_	-	-	_	-	-	_	-	_	_	-	-	-	-
40-44 years	-	_	-	-	_	-	-	_	-	_	_	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
60-64 years	-	_	-	-	-	-	-	_	-	_	_	-	-	-	-
65-69 years	-	_	-	-	-	-	-	_	-	_	_	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
75-79 years	-	-			-		-	-	-	-	-	-	-	-	-
80-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based															
on the International															
Classification of Dis													ian or		
Tenth Revision, Secon		All rac	es		White			Black			ican Ind	lian 1/		ic Isla	nder 2/
Edition, 2004)	Both	_	_	Both	_	_	Both	_	_	Both	_	_	Both	_	_
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male ———	Female	sexes	Male	Female
Breast, unspecified	(C50.9)														
All ages	41,206	389	40,817	34,460	327	34,133	5,752	58	5,694	160	_	160	834	4	830
Under 1 year	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
1-4 years	1	-	1	1	-	1	-	_	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	_	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
20-24 years	12	1	11	5	-	5	6	1	5	-	-	-	1	-	1
25-29 years	74	-	74	46	-	46	23	-	23	1	-	1	4	-	4
30-34 years	294	3	291	199	3	196	85	-	85	2	-	2	8	-	8
35-39 years	795	1	794	550	-	550	211	1	210	9	-	9	25	-	25
40-44 years	1,557	5	1,552	1,142	4	1,138	365	1	364	8	-	8	42	-	42
45-49 years	2,641	17	2,624	1,941	12	1,929	602	5	597	17	-	17	81	-	81
50-54 years	3,474	26	3,448	2,675	16	2,659	657	10	647	17	-	17	125	-	125
55-59 years	4,496	38	4,458	3,578	30	3,548	762	8	754	24	-	24	132	-	132
60-64 years	4,375	50	4,325	3,642	42	3,600	613	6	607	18	-	18	102	2	100
65-69 years	3,939	40	3,899	3,347	34	3,313	500	6	494	15	-	15	77	-	77
70-74 years	4,022	46	3,976	3,456	39	3,417	472	7	465	15	-	15	79	-	79
75-79 years	4,658	55	4,603	4,118	50	4,068	466	4	462	10	-	10	64	1	63
80-84 years	4,686	56	4,630	4,196	49	4,147	442	6	436	10	-	10	38	1	37
85-89 years	3,374	39	3,335	3,059	38	3,021	274	1	273	8	-	8	33	-	33
90-94 years	1,984	9	1,975	1,785	8	1,777	179	1	178	5	-	5	15	-	15
95-99 years	702	1	701	610	-	610	85	1	84	-	-	-	7	-	7
100 years & over.	120	2	118	109	2	107	10	-	10	1	-	1	-	-	-
Age not stated	1	-	1	-	-	-	-	_	-	_	-	-	1	-	1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of vulva (C51) All ages.... 862 810 810 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 7 40-44 years..... 11 11 10 10 45-49 years..... 17 17 14 14 3 3 34 50-54 years..... 38 38 34 55-59 years..... 57 57 50 50 6 6 1 60-64 years..... 55 55 50 50 1 73 73 69 69 65-69 vears..... . . . 70-74 vears..... 75 71 71 3 75-79 years..... 111 111 107 107 4 80-84 years..... 142 142 131 131 11 11 142 2 85-89 years..... 142 138 138 2 90-94 years..... 89 89 88 88 1 95-99 years..... 31 31 31 31 100 years & over. 9 9 7 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise	eases,											As	ian or		
Tenth Revision, Second		All rac	es		White			Black			rican In	dian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Labium majus (C51.0)															
All ages	10		10	8		8	2		2	_		_	_		_
Under 1 year	-		-	-		-	-		-	-		-	-		-
1-4 years	-		-	-		-	-		-	-		-	-		-
5-9 years	-		-	-		-	-		-	-		-	-		-
10-14 years	-		-	-		-	_		-	-		-	_		-
l5-19 years	-		-	-		-	-		-	-		-	-		-
20-24 years	-		-	-		-	-		-	-		-	-		-
25-29 years	-		-	-		-	-		-	-		-	-		-
30-34 years	-		-	-		-	-		-	-		-	-		-
35-39 years	-		-	_		_	-		-	-		-	-		-
40-44 years	-		-	-		-	_		-	-		-	_		-
45-49 years	1		1	-		-	1		1	-		-	-		-
50-54 years	-		-	-		-	-		-	-		-	-		-
55-59 years	-		-	-		-	-		-	-		-	-		-
50-64 years	1		1	1		1	-		-	-		-	-		-
65-69 years	1		1	1		1	-		-	-		-	-		-
70-74 years	1		1	1		1	_		_	_		-	_		-
75-79 years	_		_	_		-	_		_	_		-	_		-
30-84 years	3		3	2		2	1		1	-		-	_		_
35-89 years	2		2	2		2	_		_	-		-	_		_
90-94 years	1		1	1		1	_		_	_		_	_		_
95-99 years	_		_	_		_	_		_	_		_	_		_
100 years & over.	_		_	_		_	_		_	_		_	_		_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dise	ases.											As	sian or		
Tenth Revision, Second	abeb,	All rac	es		White			Black		Amer	rican In	dian 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Clitoris (C51.2)															
All ages	1		1	1		1	_		_	_		_	_		_
Under 1 year	-		-	-		-	-		-	-		-	-		-
1-4 years	-		-	-		-	-		-	-		-	-		-
5-9 years	-		-	-		-	-		-	-		-	-		-
10-14 years	-		-	-		-	-		-	-		-	-		-
15-19 years	-		-	-		-	-		-	-		-	-		-
20-24 years	-		-	-		-	-		-	-		-	-		-
25-29 years	-		-	-		-	-		-	-		-	-		-
30-34 years	-		-	-		-	-		-	-		-	-		-
35-39 years	_		_	_		_	_		_	_		_	_		_
40-44 years	_		_	_		_	_		_	_		_	_		_
45-49 years	_		_	_		_	_		_	_		_	_		_
50-54 years	_		_	_		_	_		_	_		_	_		_
55-59 years	-		-	_		_	-		-	-		-	-		-
60-64 years	-		-	_		_	-		-	-		-	-		-
65-69 years	_		-	_		_	_		-	_		-	-		-
70-74 years	_		_	_		_	_		_	_		_	_		_
75-79 years	_		_	_		_	_		_	_		_	_		_
30-84 years	_		_	_		_	_		_	_		_	_		_
85-89 years	1		1	1		1	_		_	_		_	_		_
00-94 years	_		_	_		_	_		_	_		_	_		_
95-99 years	_		_	_		_	_		_	_		_	_		_
100 years & over.	_		_	_		_	_		_	_		_	_		_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Vulva, unspecified (C51.9) All ages.... 851 851 801 801 40 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 7 40-44 years..... 11 11 10 10 45-49 years..... 16 16 14 14 2 2 34 50-54 years..... 38 38 34 55-59 years..... 57 57 50 50 6 6 1 60-64 years..... 54 54 49 49 1 72 72 68 65-69 vears..... 68 . . . 70-74 vears..... 74 70 70 3 3 75-79 years..... 111 111 107 107 4 80-84 years..... 139 139 129 129 10 10 2 85-89 years..... 139 139 135 135 2 88 87 87 1 90-94 years..... 88 95-99 years..... 31 31 31 31 9 9 7 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of vagina (C52) All ages.... 396 324 324 10 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 8 8 45-49 years..... 2.0 2.0 15 15 2 3 3 50-54 years..... 14 14 9 9 3 2 55-59 years..... 34 34 29 29 5 5 60-64 years..... 32 32 27 1 . . . 35 35 28 28 65-69 vears..... 70-74 years..... 34 30 30 75-79 years..... 55 55 47 47 8 80-84 years..... 62 62 51 51 8 8 45 85-89 years..... 56 56 45 11 11 90-94 years..... 33 33 2.7 27 6 6 95-99 years..... 5 5 5 5 100 years & over. 1 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Bot.h Both Bot.h Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Malignant neoplasm of cervix uteri (C53) All ages.... 3,976 3,976 3,014 3,014 784 784 26 26 152 152 . . . Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 1 15-19 years..... . . . 3 20-24 years..... 8 3 25-29 years..... 48 48 37 37 11 11 30-34 years..... 133 133 89 89 31 31 10 10 35-39 years..... 248 248 196 196 44 44 7 7 40-44 years..... 379 379 270 270 93 93 14 14 45-49 years..... 469 469 359 359 89 89 3 3 18 1.8 377 377 95 50-54 years..... 494 494 95 3 3 19 19 55-59 years..... 461 461 371 371 65 65 20 20 5 5 60-64 years..... 360 360 283 283 63 63 2 2 12 12 317 317 235 235 67 67 12 12 65-69 vears..... 3 . . . 70-74 vears..... 287 287 208 208 68 68 3 8 75-79 years..... 263 263 198 198 49 49 1 15 15 80-84 years..... 230 230 185 185 38 38 7 164 42 85-89 years..... 164 114 114 42 8 90 90 65 23 23 90-94 years..... 65 2 95-99 years..... 19 19 17 17 2 2 5 100 years & over. 5 1 1 . . . 4 . . . 4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Endocervix (C53.0) All ages.... 18 13 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 1 60-64 years..... 1 1 65-69 years..... 70-74 years..... 75-79 years..... 2 80-84 years..... 1 3 85-89 years..... 3 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Bot.h Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Overlapping lesion of cervix uteri (C53.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... . . . 20-24 years..... . . . 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Bot.h Both Bot.h Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Cervix uteri, unspecified (C53.9) All ages.... 3,957 3,957 3,000 3,000 780 780 25 25 152 152 . . . Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 3 20-24 years..... 8 5 3 25-29 years..... 48 48 37 37 11 11 30-34 years..... 133 133 89 89 31 31 10 10 35-39 years..... 248 248 196 196 44 7 7 40-44 years..... 375 375 267 267 92 92 14 14 45-49 years..... 469 469 359 359 89 89 3 3 18 1.8 375 375 95 492 492 95 3 3 19 19 50-54 years..... 55-59 years..... 460 460 371 371 65 65 20 20 4 4 60-64 years..... 359 359 282 282 63 63 2 2 12 12 315 315 234 234 66 12 12 65-69 vears..... 66 3 . . . 70-74 vears..... 285 285 206 206 68 68 3 8 75-79 years..... 261 261 196 196 49 49 1 15 15 80-84 years..... 229 229 185 185 37 37 7 160 85-89 years..... 160 111 111 41 41 8 90 90 65 23 23 90-94 years..... 65 2 95-99 years..... 19 19 17 17 2 2 5 100 years & over. 5 1 1 . . . 4 . . . 4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of corpus uteri (C54) All ages.... 3,449 3,449 2,809 2,809 550 550 15 15 75 75 Under 1 year.... 1 1 1 1 1-4 years..... . . . 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 8 8 7 1 35-39 years..... 21 21 17 17 7 40-44 years..... 48 48 40 40 7 45-49 years..... 85 85 65 65 12 12 1 50-54 years..... 158 158 135 135 16 16 2 5 55-59 years..... 338 338 262 262 61 61 15 15 60-64 years..... 414 414 309 309 91 91 14 14 343 343 98 98 65-69 vears..... 453 453 9 70-74 vears..... 508 508 401 401 95 95 4 8 75-79 years..... 490 490 417 417 65 65 8 80-84 years..... 446 446 379 379 61 61 85-89 years..... 292 292 261 261 29 29 . . . 1 90-94 years..... 138 138 130 130 6 2 6 95-99 years..... 44 44 38 38 5 1 1 5 3 3 100 years & over. 3 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Endometrium (C54.1) All ages.... 3,407 3,407 2,774 2,774 543 543 15 15 75 75 Under 1 year..... 1 1 1 1 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 8 8 7 1 35-39 years..... 20 20 7 40-44 years..... 47 47 39 39 7 45-49 years..... 84 84 64 64 12 12 1 50-54 years..... 155 133 133 15 15 2 5 155 55-59 years..... 335 335 259 259 61 61 15 15 60-64 years..... 409 409 306 306 89 89 14 14 . . . 337 337 97 97 65-69 vears..... 446 446 9 70-74 vears..... 504 504 397 397 95 95 4 8 75-79 years..... 485 485 413 413 64 64 8 80-84 years..... 438 438 372 372 60 60 28 85-89 years..... 290 290 260 260 28 . . . 1 136 136 128 128 2 90-94 years..... 6 6 95-99 years..... 38 44 44 38 5 1 5 1 3 100 years & over. 3 3 3 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise								-1 1		_			sian or	1	1 0/
Tenth Revision, Second		All rac	es	D	White		D	Black			rican Ind	dian 1/		fic Isla	nder 2/
Edition, 2004)	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Mala	Female	Both	Mala	Female
and age 	sexes	мате	remaie	sexes	мате	remaie	sexes		remare	sexes	мате	remare	sexes		remale
Myometrium (C54.2)															
All ages	1		1	1		1	_		-	-		-	-		-
Under 1 year	-		-	-		-	-		-	-		-	-		-
1-4 years	-		-	-		-	-		-	-		-	-		-
5-9 years	-		-	-		-	-		-	-		-	-		-
l0-14 years	-		-	-		-	-		-	-		-	-		-
15-19 years	-		-	-		-	-		-	-		-	-		-
20-24 years	-		-	-		-	-		-	-		-	-		-
25-29 years	-		-	-		-	-		-	-		-	-		-
30-34 years	-		-	-		-	-		-	-		-	-		-
35-39 years	-		-	-		-	-		-	-		-	-		-
40-44 years	-		-	-		-	-		-	-		-	-		-
45-49 years	-		-	-		-	-		-	-		-	-		-
50-54 years	-		-	_		-	-		-	-		-	_		-
55-59 years	1		1	1		1	-		-	-		-	_		-
50-64 years	-		-	_		-	-		-	-		-	_		-
65-69 years	-		-	_		-	-		-	-		-	-		-
70-74 years	-		-	-		-	-		-	-		-	-		-
75-79 years	_		-	_		-	-		-	_		-	-		-
30-84 years	_		-	_		-	-		-	_		-	-		-
35-89 years	_		-	_		-	-		-	_		-	-		-
90-94 years	_		-	_		-	-		-	_		-	-		-
95-99 years	_		-	_		-	-		-	_		-	-		-
100 years & over.	-		-	_		-	-		-	_		-	_		-
Age not stated	_		_	_		_	_			_		_	_		_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Corpus uteri, unspecified (C54.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 3 50-54 years..... 3 55-59 years..... 2 2 60-64 years..... 5 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 8 8 1 2 85-89 years..... 2 1 90-94 years..... 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Malignant neoplasm of uterus, part unspecified (C55) All ages.... 3,935 3,935 3,145 3,145 690 690 15 15 85 85 Under 1 year..... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 6 30-34 years..... 11 11 8 35-39 years..... 25 25 22 22 8 40-44 years..... 51 51 39 39 45-49 years..... 135 135 101 101 2.8 2.8 1 5 . . . 33 33 50-54 years..... 221 221 173 173 1 14 55-59 years..... 353 353 271 271 73 73 3 3 6 6 60-64 years..... 417 417 318 318 89 89 2 2 8 328 328 111 65-69 vears..... 455 455 111 16 . . . 70-74 vears..... 481 481 356 356 111 111 13 13 . . . 75-79 years..... 562 562 450 450 100 100 8 80-84 years..... 551 551 476 476 69 69 85-89 years..... 386 386 345 345 36 36 . . . 1 196 196 175 175 21 21 90-94 years..... 95-99 years..... 73 73 70 70 2 2 1 1 10 10 9 100 years & over. 9 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Bot.h Both Bot.h Both Bot.h and age Female Male Female Male Female sexes Male Female sexes Male Female sexes Male sexes sexes Malignant neoplasm of ovary (C56) All ages.... 14,857 14,857 13,289 13,289 1,185 1,185 62 62 321 321 . . . Under 1 year..... 1 1 1 1 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 3 3 3 3 15-19 years..... 2 6 6 20-24 years..... 17 17 16 16 1 25-29 years..... 26 26 20 20 2 30-34 years..... 53 53 47 47 35-39 years..... 116 116 100 100 11 11 5 40-44 years..... 285 285 235 235 38 38 11 11 45-49 years..... 629 629 530 530 76 76 3 3 2.0 20 1,037 9 1.037 881 881 104 9 43 43 50-54 years..... 104 1,395 1,395 1,218 1,218 128 128 6 43 43 55-59 years..... 6 60-64 years..... 1,653 1,653 1,465 1,465 146 146 4 4 38 38 1,504 156 22 22 65-69 vears..... 1,695 1,695 1,504 156 13 13 70-74 vears..... 1,878 1,878 1,708 1,708 130 130 2 2 38 38 75-79 years..... 2,092 2,092 1,892 1,892 151 151 9 40 40 . . . 80-84 years..... 1,969 1,969 1.820 1,820 105 105 11 11 33 33 85-89 years..... 1,306 1,306 1,199 1,199 88 88 2 2 17 17 554 554 2.7 2.7 90-94 years..... 522 522 5 5 95-99 years..... 133 133 119 119 1 12 12 1 1 1 2 1 100 years & over. 8 8 5 5 2 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Malignant neoplasm of other and unspecified female genital organs (C57) All ages.... 304 46 10 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... . . . 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 3 40-44 years..... 11 11 45-49 years..... 6 6 4 2 2 50-54 years..... 3.0 30 21 21 55-59 years..... 31 31 26 26 4 1 60-64 years..... 31 31 26 26 47 47 35 35 10 10 65-69 vears..... 70-74 years..... 56 56 51 5 5 75-79 years..... 50 50 46 46 80-84 years..... 33 33 30 30 3 85-89 years..... 40 40 34 34 3 3 90-94 years..... 21 21 18 18 2 2 1 95-99 years..... 4 3 4 3 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Fallopian tube (C57.0) All ages.... 156 156 136 136 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 2 2 1 50-54 years..... 3 3 14 14 1.0 1 0 55-59 years..... 15 15 14 14 1 60-64 years..... 15 15 13 13 2 25 25 21 65-69 vears..... 70-74 years..... 35 35 33 33 75-79 years..... 18 18 18 18 80-84 years..... 11 11 9 9 9 85-89 years..... 9 1 90-94 years..... 5 5 1 1 4 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Uterine adnexa, unspecified (C57.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified female genital organs (C57.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... . . . 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Overlapping lesion of female genital organs (C57.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... . . . 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise		All rac			White			Black		3	T.	As dian 1/	ian or	::. T.l.	nder 2/
Tenth Revision, Second Edition, 2004)	a Both	AII rac	es	Both	wille		Both	Black		Both	Tean II.	dian 1/	Both	ic isia	inder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Female genital organ,	unspecified	l (C57.9)													
All ages	196		196	160		160	27		27	2		2	7		7
Under 1 year	-		-	-		-	-		-	-		-	-		-
1-4 years	-		-	-		-	-		-	-		-	-		-
5-9 years	-		-	-		-	-		-	-		-	-		-
10-14 years	-		-	-		-	-		-	-		-	-		-
5-19 years	-		-	-		-	-		-	-		-	-		-
0-24 years	-		-	-		-	-		-	-		-	-		-
25-29 years	-		-	-		-	-		-	-		-	-		-
30-34 years	-		-	-		-	-		-	-		-	-		-
35-39 years	2		2	1		1	1		1	-		-	-		-
40-44 years	6		6	5		5	-		-	-		-	1		1
45-49 years	3		3	2		2	1		1	-		-	-		-
50-54 years	14		14	10		10	3		3	-		-	1		1
55-59 years	14		14	12		12	2		2	-		-	-		-
60-64 years	15		15	12		12	2		2	-		-	1		1
65-69 years	19		19	11		11	8		8	-		-	-		-
70-74 years	20		20	17		17	3		3	-		-	-		-
75-79 years	31		31	27		27	3		3	1		1	-		_
30-84 years	22		22	21		21	1		1	-		-	-		_
35-89 years	31		31	26		26	2		2	1		1	2		2
90-94 years	15		15	13		13	1		1	-		-	1		1
95-99 years	4		4	3		3	-		-	-		-	1		1
100 years & over.	_		_	_		_	_		_	_		_	_		_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of placenta (C58) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 2 2 50-54 years..... 1 1 1 1 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of penis (C60) All ages.... 245 211 211 32 32 . . . Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 3 3 3 2 45-49 years..... 12 12 10 1 0 50-54 years..... 12 12 9 9 3 3 . . . 55-59 years..... 25 25 22 22 3 60-64 years..... 36 36 28 28 8 8 32 32 26 26 65-69 vears..... 5 . . . 70-74 vears..... 25 25 22 22 . . . 75-79 years..... 2.7 2.7 26 80-84 years..... 35 35 32 32 2 19 18 85-89 years..... 19 18 90-94 years..... 11 2 11 9 9 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female Male Female sexes Male sexes Male sexes sexes sexes Penis, unspecified (C60.9) All ages.... 245 245 211 211 32 32 . . . Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 3 3 3 2 45-49 years..... 12 12 10 1 0 50-54 years..... 12 12 9 9 3 3 . . . 55-59 years..... 25 25 22 22 3 60-64 years..... 36 36 28 28 8 8 32 32 26 26 65-69 vears..... 5 . . . 70-74 vears..... 25 25 22 22 3 . . . 75-79 years..... 2.7 2.7 26 80-84 years..... 35 35 32 32 2 19 85-89 years..... 19 18 18 90-94 years..... 11 2 11 9 9 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based	Ė														
on the International												_			
Classification of Dis Tenth Revision, Secon		All rac			White			Black		3	ican In		sian or	ic Isla	
Edition, 2004)	Both	AII rac	es	Both	willte		Both	Black		Both	ican in	dian 1/	Both	ic isia	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
		Hare	- Ciliare												
Malignant neoplasm of	f prostate (C61)													
All ages	28,372	28,372		23,202	23,202		4,698	4,698		104	104		368	368	
Under 1 year	-	-		-	-		-	-		-	-		-	-	
1-4 years	-	-		-	-		-	-		-	-		-	-	
5-9 years	-	-		-	-		-	-		-	-		-	-	
10-14 years	1	1		1	1		-	-		-	-		-	-	
15-19 years	-	-		-	-		-	-		-	-		-	-	
20-24 years	1	1		-	-		1	1		-	-		-	-	
25-29 years	3	3		1	1		2	2		-	-		-	-	
30-34 years	2	2		1	1		1	1		-	-		-	-	
35-39 years	3	3		2	2		1	1		-	-		_	-	
40-44 years	37	37		25	25		11	11		-	-		1	1	
45-49 years	93	93		66	66		26	26		1	1		_	-	
50-54 years	312	312		222	222		90	90		-	-		_	-	
55-59 years	833	833		599	599		219	219		7	7		8	8	
60-64 years	1,440	1,440		1,084	1,084		322	322		8	8		26	26	
65-69 years	2,209	2,209		1,649	1,649		511	511		14	14		35	35	
70-74 years	3,394	3,394		2,588	2,588		742	742		17	17		47	47	
75-79 years	4,924	4,924		4,001	4,001		849	849		14	14		60	60	
80-84 years	6,233	6,233		5,261	5,261		870	870		24	24		78	78	
85-89 years	5,377	5,377		4,671	4,671		639	639		9	9		58	58	
90-94 years	2,743	2,743		2,398	2,398		300	300		7	7		38	38	
95-99 years	691	691		578	578		94	94		3	3		16	16	
100 years & over.	76	76		55	55		20	20		-	-		1	1	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Malignant neoplasm of testis (C62) All ages.... 358 326 326 23 23 . . . Under 1 year.... 1-4 years..... 1 1 1 5-9 years..... . . . 10-14 years..... 2 2 2 2 15-19 years..... 5 1 1 20-24 years..... 36 36 33 33 1 25-29 years..... 46 46 30-34 years..... 44 44 37 37 35-39 years..... 46 37 37 40-44 years..... 39 39 36 36 29 45-49 years..... 30 3.0 29 1 26 26 50-54 years..... 28 28 1 1 . . . 55-59 years..... 26 26 24 24 2 60-64 years..... 16 16 16 16 . . . 11 11 11 65-69 vears..... 11 . . . 70-74 vears..... 8 8 . . . 75-79 years..... 6 6 80-84 years..... 4 4 85-89 years..... 5 5 90-94 years..... 4 4 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Testis, unspecified (C62.9) All ages.... 358 358 326 326 23 23 . . . Under 1 year.... 1-4 years..... 1 1 1 5-9 years..... . . . 10-14 years..... 2 2 2 2 15-19 years..... 5 1 1 20-24 years..... 36 36 33 33 1 25-29 years..... 46 46 30-34 years..... 44 44 37 37 35-39 years..... 46 37 37 40-44 years..... 39 39 36 36 45-49 years..... 30 3.0 29 29 1 26 26 2.8 28 50-54 years..... 1 1 . . . 55-59 years..... 26 26 24 24 2 60-64 years..... 16 16 16 16 . . . 11 11 11 65-69 vears..... 11 . . . 70-74 vears..... 8 8 . . . 75-79 years..... 6 6 80-84 years..... 4 4 85-89 years..... 5 5 90-94 years..... 4 4 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of other and unspecified male genital organs (C63) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 2 2 . . . 55-59 years..... 3 3 60-64 years..... 2 2 5 5 65-69 vears..... 70-74 vears..... 7 75-79 years..... 8 80-84 years..... 3 3 . . . 85-89 years..... 9 9 90-94 years..... 3 3 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise	eases,										A:	sian or	
Tenth Revision, Second	i	All rac	es		White			Black			rican Indian 1/		ic Islander 2/
Edition, 2004)	Both			Both			Both			Both		Both	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Female	sexes	Male Female
Epididymis (C63.0)													
All ages	3	3		3	3		-	-		-		_	
Under 1 year	-	-		-	-		-	-		-		-	
1-4 years	-	-		-	-		-	-		-		-	
5-9 years	-	-		-	-		-	-		-		-	
10-14 years	-	-		-	-		-	-		-		-	
15-19 years	-	-		-	-		-	-		-		-	
20-24 years	-	-		-	-		-	-		-		-	
25-29 years	-	-		-	-		-	-		-		-	
30-34 years	-	-		-	-		-	-		-		-	
35-39 years	-	-		-	-		-	-		-		-	
40-44 years	-	-		-	-		-	-		-		-	
45-49 years	1	1		1	1		-	-		-		-	
50-54 years	-	-		-	-		-	-		-		-	
55-59 years	2	2		2	2		-	-		-		-	
60-64 years	-	-		-	-		-	-		-		-	
65-69 years	-	-		-	-		-	-		-		-	
70-74 years	-	-		-	-		-	-		-		-	
75-79 years	-	-		-	-		-	-		-		-	
80-84 years	-	-		-	-		-	-		-		-	
85-89 years	-	-		-	-		-	-		-		-	
90-94 years	-	-		-	-		-	-		-		-	
95-99 years	-	-		-	-		-	-		-		-	
100 years & over.	-	-		-	-		-	-		-		-	
Age not stated	_	_		_	-		_	-		-		-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise												sian or	
Tenth Revision, Second		All rac	ces		White			Black			rican Indian 1/		ic Islander 2/
Edition, 2004)	Both	_	_	Both	_	_	Both	_	_	Both		Both	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Female	sexes	Male Female
Spermatic cord (C63.1)													
All ages	2	2		2	2		-	_		-		-	
Under 1 year	-	-		-	-		-	-		-		-	
1-4 years	-	-		-	-		-	-		-		_	
5-9 years	-	-		-	-		-	-		-		-	
10-14 years	-	-		-	-		-	-		-		-	
15-19 years	-	-		-	-		-	-		-		-	
20-24 years	-	-		-	-		-	-		-		-	
25-29 years	-	-		-	-		-	-		-		-	
30-34 years	-	-		-	-		-	-		-		-	
35-39 years	-	-		-	-		-	-		-		-	
40-44 years	-	-		-	-		-	-		-		-	
45-49 years	-	-		-	-		-	-		-		-	
50-54 years	-	-		-	-		-	-		-		-	
55-59 years	-	-		-	-		-	-		-		-	
60-64 years	-	-		-	-		-	-		-		-	
65-69 years	-	-		-	-		-	-		-		-	
70-74 years	-	-		-	-		-	-		-		-	
75-79 years	-	-		-	-		-	-		-		-	
80-84 years	1	1		1	1		-	-		-		-	
85-89 years	1	1		1	1		-	-		-		-	
90-94 years	-	-		-	-		-	-		-		-	
95-99 years	-	-		-	-		-	-		-		-	
100 years & over.	-	-		-	-		-	-		-		-	
Age not stated	_	_		_	_		_	_		_		_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Scrotum (C63.2) All ages.... 15 15 13 13 Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 . . . 55-59 years..... 1 1 60-64 years..... 1 1 4 65-69 vears..... 70-74 vears..... 3 3 75-79 years..... 2 80-84 years..... 1 1 . . . 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified male genital organs (C63.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... . . . 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... . . . 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... . . . 85-89 years..... . . . 90-94 years..... 95-99 years..... 100 years & over. . . . Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Male genital organ, unspecified (C63.9) All ages..... . . . Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 . . . 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 4 . . . 75-79 years..... 6 80-84 years..... 1 1 . . . 8 85-89 years..... 8 90-94 years..... 2 2 2 2 95-99 years..... 1 1 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based	l														
on the International												_			
Classification of Dis		311			rate to a			D11-		3			ian or	:	- 1 0 /
Tenth Revision, Secon	ia Both	All rac	es	Both	White		Both	Black		Amer Both	ican Ind	alan I/	Both	ic isia	inder 2/
Edition, 2004) and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Mala	Female	sexes	Mala	Female
	Seves	maie	remare	Seves	Male	remare	SCACS		remare	Seves		remare	SEACS		remare
Malignant neoplasm of	kidney, ex	cept renal	pelvis (C6	4)											
All ages	12,214	7,608	4,606	10,805	6,725	4,080	1,132	710	422	79	52	27	198	121	77
Under 1 year	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
1-4 years	11	4	7	10	4	6	-	-	-	1	-	1	-	-	-
5-9 years	23	14	9	19	11	8	4	3	1	-	-	-	-	-	-
10-14 years	7	4	3	3	-	3	4	4	-	-	-	-	-	-	-
L5-19 years	11	5	6	7	3	4	4	2	2	-	-	-	-	-	-
20-24 years	9	6	3	4	2	2	4	3	1	_	_	-	1	1	-
25-29 years	19	12	7	7	3	4	10	8	2	-	-	-	2	1	1
30-34 years	40	25	15	24	11	13	14	12	2	_	_	-	2	2	-
35-39 years	77	48	29	56	32	24	17	13	4	2	2	_	2	1	1
10-44 years	189	140	49	145	109	36	37	29	8	2	1	1	5	1	4
15-49 years	450	313	137	374	258	116	68	49	19	1	-	1	7	6	1
50-54 years	759	540	219	624	454	170	116	74	42	7	4	3	12	8	4
55-59 years	1,158	828	330	1,005	710	295	129	97	32	7	7	-	17	14	3
60-64 years	1,265	883	382	1,122	784	338	108	75	33	11	6	5	24	18	6
65-69 years	1,522	1,037	485	1,347	915	432	132	93	39	11	7	4	32	22	10
70-74 years	1,521	940	581	1,349	840	509	133	75	58	13	10	3	26	15	11
75-79 years	1,742	1,060	682	1,592	979	613	114	60	54	7	6	1	29	15	14
80-84 years	1,608	873	735	1,468	802	666	109	58	51	11	5	6	20	8	12
85-89 years	1,156	602	554	1,062	555	507	80	39	41	2	2	_	12	6	6
90-94 years	509	225	284	463	209	254	37	12	25	3	2	1	6	2	4
95-99 years	126	46	80	113	41	72	11	4	7	1	-	1	1	1	-
100 years & over.	10	1	9	9	1	8	1	_	1	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of renal pelvis (C65) All ages..... 156 70 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 2 2 3 50-54 years..... 2 3 1 2 1 55-59 years..... 20 10 10 20 10 10 60-64 years..... 19 12 7 17 11 6 65-69 vears..... 12 9 3 11 3 70-74 vears..... 30 18 29 75-79 years..... 30 15 15 28 13 15 80-84 years..... 32 16 16 31 16 15 1 1 9 85-89 years..... 5 4 9 5 4 90-94 years..... 5 1 4 5 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of ureter (C66) All ages..... 189 172 338 180 158 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 45-49 years..... 3 2 1 3 2 8 50-54 years..... 8 6 55-59 years..... 27 18 9 26 18 60-64 years..... 16 10 6 15 10 5 65-69 vears..... 31 20 28 19 9 70-74 vears..... 44 25 19 44 25 19 75-79 years..... 69 38 31 67 37 30 80-84 years..... 72 36 36 69 35 34 2 1 48 22 43 25 3 85-89 years..... 26 18 3 29 7 90-94 years..... 22 2.6 19 1 1 95-99 years..... 10 3 7 9 1 3 6 1 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Malignant neoplasm of bladder (C67) All ages.... 13,474 9,412 4,062 12,286 8,735 3,551 1,007 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 vears..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 50-54 years..... 1.0 55-59 years..... 60-64 years..... 1,166 1,051 65-69 vears..... 70-74 vears..... 1,644 1,202 1,471 1,108 75-79 years..... 2,216 1,602 2,034 1,507 80-84 years..... 2,658 1,870 2,461 1,763 85-89 years..... 2,138 1,421 1,995 1,349 90-94 years..... 1,098 1.022 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Trigone of bladder (C67.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dise													ian or		
Tenth Revision, Second		All rac	es	_	White		_	Black			rican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both	1	- 1	Both		- 1	Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale 	sexes	Male	Female
Dome of bladder (C67.	1)														
All ages	1	1	_	1	1	-	-	_	-	-	-	-	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise	ases,											As	ian or		
Tenth Revision, Second		All rac	es		White			Black		Ame	rican Indi	an 1/	Pacif	fic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Bladder neck (C67.5)															
All ages	4	2	2	4	2	2	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	_
l5-19 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	_	_	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	_	_	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	-	_	-	_	-	-	-	-	-	-	-	-
10-44 years	-	-	-	_	_	-	-	-	_	-	_	-	-	-	-
45-49 years	-	-	-	_	-	-	-	_	-	-	-	-	_	-	-
50-54 years	-	-	-	_	-	-	-	_	-	-	-	-	_	-	-
55-59 years	-	-	-	_	-	-	-	_	-	-	-	-	_	-	-
0-64 years	1	-	1	1	-	1	-	_	-	-	-	-	_	-	-
55-69 years	-	-	-	_	-	-	-	_	-	-	-	-	_	-	-
70-74 years	-	-	-	_	-	_	-	_	-	_	_	-	-	_	-
75-79 years	-	-	-	_	-	_	-	_	-	_	_	-	-	_	-
30-84 years	1	-	1	1	-	1	-	_	-	_	_	-	-	_	-
35-89 years	1	1	-	1	1	_	-	_	-	_	_	-	-	_	-
90-94 years	1	1	-	1	1	_	-	_	-	_	_	-	-	_	-
95-99 years	-	-	-	-	-	_	-	-	-	_	_	-	_	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Disc								_, ,					sian or		
Tenth Revision, Second		All rac	es		White			Black			rican Ind	lian 1/		ic Isla	nder 2/
Edition, 2004)	Both	1	- 1	Both			Both		_ 1	Both		_ ,	Both		_ 1
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	remale	sexes		Female
Ureteric orifice (C67	.6)														
All ages	1	_	1	1	-	1	-	-	-	-	-	-	-	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	_	-	-
80-84 years	-	-	-	_	-	-	-	-	-	-	-	-	_	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Urachus (C67.7) All ages.... 21 10 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 3 50-54 years..... 1 55-59 years..... 4 2 2 60-64 years..... 3 2 1 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 80-84 years..... 2 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Bladder, unspecified (C67.9) All ages.... 13,446 9,397 4,049 12,263 8,723 3,540 1,004 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 50-54 years..... 1.0 55-59 years..... 60-64 years..... 1,164 1,049 65-69 vears..... 70-74 vears..... 1,639 1,198 1,467 1,105 75-79 years..... 2,214 1,602 2,032 1,507 80-84 years..... 2,655 1,869 2,459 1,762 85-89 years..... 2,137 1,420 1,994 1,348 1,097 1.021 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of other and unspecified urinary organs (C68) All ages.... 124 37 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 11 2 45-49 years..... 6 5 17 50-54 years..... 7 R 3 6 10 11 55-59 years..... 30 18 12 22 15 6 2 60-64 years..... 32 19 13 22 17 5 1 65-69 vears..... 43 23 20 32 22 70-74 years..... 48 35 13 46 12 75-79 years..... 79 48 31 72 43 29 3 80-84 years..... 79 46 33 72 43 29 2 56 50 85-89 years..... 40 16 39 11 90-94 years..... 31 19 12 29 17 12 1 95-99 years..... 6 3 3 5 2 3 1 1 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Urethra (C68.0) All ages.... 191 108 142 49 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 2 2 4 15 5 50-54 years..... 5 10 10 55-59 years..... 16 9 7 10 5 60-64 years..... 19 9 10 10 3 65-69 vears..... 31 17 21 70-74 vears..... 21 14 7 75-79 years..... 35 19 31 17 16 80-84 years..... 23 14 9 18 11 7 3 2 1 1 85-89 years..... 14 10 12 10 2 1 90-94 years..... 7 1 1 6 1 5 4 95-99 years..... 2 1 1 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Overlapping lesion of urinary organs (C68.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Urinary organ, unspecified (C68.9) All ages..... 160 153 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 5 45-49 years..... 1 2 50-54 years..... 2 1 55-59 years..... 9 14 5 12 60-64 years..... 13 10 3 12 10 65-69 vears..... 12 9 3 11 9 70-74 vears..... 27 6 26 75-79 years..... 44 29 15 41 15 80-84 years..... 56 32 24 54 32 22 1 42 38 29 3 85-89 years..... 30 12 9 3 90-94 years..... 23 13 10 23 13 10 95-99 years..... 4 2 2 3 2 1 100 years & over. 1 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of eye and adnexa (C69) All ages.... 111 105 106 Under 1 year.... 1 1 1 1 1-4 years..... 2 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 45-49 years..... 6 3 3 6 2 2 17 50-54 years..... 8 9 16 55-59 years..... 26 19 23 16 60-64 years..... 18 9 9 17 9 65-69 vears..... 23 15 8 23 70-74 vears..... 32 18 31 17 75-79 years..... 2.7 18 9 2.7 18 9 80-84 years..... 31 13 18 31 13 18 10 10 85-89 years..... 5 5 5 5 90-94 years..... 9 5 4 9 95-99 years..... 4 4 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Retina (C69.2) All ages.... Under 1 year.... 1-4 years..... 2 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Choroid (C69.3) All ages.... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 3 3 60-64 years..... 3 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise	ases,											As	ian or		
Tenth Revision, Second		All rac	es		White			Black		Amei	rican Indi			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Ciliary body (C69.4)															
All ages	6	4	2	6	4	2	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-
20-24 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
30-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	-	_	-	-	-	_	_	-	-	-	-	-	-	-	-
40-44 years	-	_	-	-	-	_	_	-	-	-	-	-	-	-	-
45-49 years	1	1	-	1	1	_	_	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	_	-	-	_	-	-	-	-
65-69 years	2	2	-	2	2	-	-	_	-	-	_	-	-	-	-
70-74 years	1	-	1	1	-	1	_	-	-	-	-	-	-	-	-
75-79 years	_	_	-	_	_	_	_	-	-	-	-	-	-	_	-
30-84 years	_	_	-	_	_	_	_	-	-	-	-	-	-	_	-
35-89 years	1	1	-	1	1	_	_	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	_	_	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lacrimal gland and duct (C69.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Orbit (C69.6) All ages.... 19 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 2 1 1 60-64 years..... 2 1 1 65-69 vears..... 3 70-74 vears..... 8 6 75-79 years..... 8 6 80-84 years..... 10 5 10 85-89 years..... 1 2 90-94 years..... 3 1 1 95-99 years..... 2 2 2 2 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Eye, unspecified (C69.9) All ages..... 139 136 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 2 2 5 2 3 50-54 years..... 12 5 11 55-59 years..... 20 6 18 12 14 60-64 years..... 12 8 4 12 65-69 vears..... 13 13 6 70-74 vears..... 20 10 10 20 10 75-79 years..... 18 7 18 11 7 11 80-84 years..... 16 5 11 16 5 11 85-89 years..... 8 3 5 8 5 90-94 years..... 6 3 3 6 3 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasm of meninges (C70) All ages..... 59 93 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 2 45-49 years..... 1.0 Ω 2 10 50-54 years..... 6 1 9 55-59 years..... 15 10 12 3 5 60-64 years..... 11 5 6 9 65-69 vears..... 5 70-74 vears..... 15 8 75-79 years..... 16 12 80-84 years..... 9 5 4 85-89 years..... 9 5 9 1 90-94 years..... 3 4 95-99 years..... 2 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cerebral meninges (C70.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 4 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Meninges, unspecified (C70.9) All ages.... 47 11 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 2 45-49 years..... 10 9 50-54 years..... 5 9 55-59 years..... 11 60-64 years..... 11 65-69 vears..... 5 70-74 vears..... 15 8 75-79 years..... 13 3 10 12 80-84 years..... 8 4 85-89 years..... 9 5 9 1 90-94 years..... 3 4 95-99 years..... 2 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Malignant neoplasm of brain (C71) All ages.... 12,670 7,088 5,582 11,664 6,554 5,110 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 vears..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 1.138 1.043 2.0 55-59 years..... 1,454 1,323 60-64 years..... 1,430 1,348 1,329 1,233 65-69 years..... 2.4 70-74 years..... 1,406 1,312 75-79 years..... 1,344 1,282 80-84 years..... 1,091 1.027 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based															
on the International															
Classification of Dise												As	ian or		
Tenth Revision, Second			White			Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Cerebrum, except lobes	s and ventri	cles (C71	.0)												
All ages	173	92	81	157	84	73	10	6	4	1	1	_	5	1	4
Under 1 year	1	-	1	1	_	1	_	-	_	_	-	-	-	_	-
1-4 years	4	3	1	1	_	1	3	3	_	_	-	-	-	_	-
5-9 years	4	-	4	4	_	4	_	-	_	_	-	-	-	_	-
10-14 years	6	4	2	4	2	2	1	1	_	-	-	-	1	1	-
.5-19 years	7	3	4	7	3	4	-	-	_	-	-	-	-	-	-
0-24 years	3	-	3	3	-	3	_	-	_	-	-	-	-	-	-
25-29 years	1	-	1	1	-	1	-	-	_	-	-	-	-	-	-
0-34 years	3	2	1	2	1	1	-	-	_	1	1	-	-	-	-
35-39 years	5	3	2	4	3	1	_	-	_	-	-	-	1	-	1
0-44 years	5	4	1	5	4	1	_	-	_	-	-	-	-	-	-
5-49 years	5	4	1	5	4	1	_	-	_	-	-	-	-	-	-
0-54 years	11	6	5	9	6	3	1	-	1	-	-	-	1	-	1
5-59 years	13	6	7	12	6	6	_	-	_	-	-	-	1	-	1
50-64 years	21	13	8	19	12	7	2	1	1	-	-	-	-	-	-
55-69 years	17	11	6	17	11	6	-	-	-	-	-	-	-	-	-
70-74 years	18	12	6	16	12	4	1	-	1	-	-	-	1	-	1
5-79 years	15	5	10	15	5	10	_	-	_	_	-	-	-	_	-
0-84 years	13	8	5	12	7	5	1	1	_	_	-	-	-	_	-
5-89 years	13	6	7	12	6	6	1	-	1	_	-	-	-	_	-
90-94 years	6	2	4	6	2	4	_	-	_	-	-	-	-	-	-
95-99 years	2	-	2	2	-	2	_	-	_	-	-	-	-	-	-
100 years & over.	-	-	_	_	-	_	_	_	_	-	-	-	-	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Frontal lobe (C71.1) All ages.... 42 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 5 40-44 years..... 45-49 years..... 10 5 50-54 years..... 1 55-59 years..... 4 3 60-64 years..... 9 5 65-69 vears..... 6 70-74 vears..... 8 75-79 years..... 9 80-84 years..... 10 5 1 5 85-89 years..... 5 2 90-94 years..... 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Temporal lobe (C71.2) All ages.... 32 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 5 3 2 5 2 9 50-54 years..... 5 R 55-59 years..... 16 15 11 11 5 60-64 years..... 11 5 6 10 65-69 vears..... 9 70-74 vears..... 11 75-79 years..... 9 9 80-84 years..... 12 5 12 85-89 years..... 4 3 4 90-94 years..... 3 3 4 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Parietal lobe (C71.3) All ages.... 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 6 3 60-64 years..... 3 65-69 vears..... 5 70-74 vears..... 75-79 years..... 8 80-84 years..... 5 85-89 years..... 3 1 90-94 years..... 1 1 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Occipital lobe (C71.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cerebral ventricle (C71.5) All ages.... 10 Under 1 year.... 2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dise							Asian or								
		All rac			White		Black			American Indian 1/			Pacific Islander 2/		
Edition, 2004)	Both			Both			Both		_	Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Cerebellum (C71.6)															
All ages	160	98	62	142	91	51	13	5	8	1	_	1	4	2	2
Under 1 year	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
1-4 years	29	19	10	25	17	8	2	1	1	-	-	-	2	1	1
5-9 years	32	22	10	29	21	8	3	1	2	-	-	-	-	-	-
10-14 years	17	12	5	15	11	4	1	1	-	-	-	-	1	-	1
15-19 years	10	8	2	9	8	1	1	-	1	-	-	-	-	-	-
20-24 years	8	5	3	6	3	3	2	2	-	-	-	-	-	-	-
25-29 years	7	4	3	6	4	2	1	-	1	-	-	-	-	-	-
30-34 years	6	3	3	6	3	3	-	-	-	-	-	-	-	-	-
35-39 years	7	5	2	7	5	2	-	-	-	-	-	-	-	-	-
40-44 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
45-49 years	7	4	3	6	4	2	-	-	-	1	-	1	-	-	-
50-54 years	7	2	5	5	2	3	2	-	2	-	-	-	-	-	-
55-59 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
60-64 years	7	4	3	6	3	3	-	-	-	-	-	-	1	1	-
65-69 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	3	1	2	2	1	1	1	-	1	-	-	-	-	-	-
30-84 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-
85-89 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	-	_	_	_	-	-	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Brain stem (C71.7) All ages.... 230 120 110 184 100 34 19 10 Under 1 year.... 1-4 years..... 22 13 16 72 36 36 53 28 25 8 3 5-9 years..... 16 10-14 years..... 2.8 16 12 2.0 12 8 6 2 15-19 years..... 9 3 13 4 9 3 1 20-24 years..... 4 3 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 13 45-49 years..... 9 5 a 50-54 years..... 1 0 3 55-59 years..... 10 5 5 60-64 years..... 2 2 65-69 vears..... 6 5 70-74 vears..... 5 3 75-79 years..... 80-84 years..... 5 5 2 85-89 years..... 3 3 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Overlapping lesion of brain (C71.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 3 50-54 years..... 2 55-59 years..... 4 1 3 60-64 years..... 4 2 65-69 vears..... 70-74 vears..... 75-79 years..... 4 80-84 years..... 5 2 3 2 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Brain, unspecified (C71.9) All ages.... 11,834 6,622 5,212 10,925 6,132 4,793 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 2.4 15-19 years..... 2.1 20-24 vears..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 2.7 50-54 years..... 1.086 1.000 55-59 years..... 1,394 1,266 60-64 years..... 1,367 1,290 1,278 1,186 65-69 years..... 70-74 years..... 1,355 1,263 75-79 years..... 1,294 1,233 80-84 years..... 1,034 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Bot.h Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Malignant neoplasm of spinal cord, cranial nerves, and other parts of central nervous system (C72) All ages.... Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 6 1 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 2 1 1 50-54 years..... 5 6 11 11 55-59 years..... 8 5 2 3 6 60-64 years..... 7 5 2 65-69 vears..... 13 70-74 vears..... 13 6 11 75-79 years..... 7 80-84 years..... 9 6 9 85-89 years..... 2 90-94 years..... 4 3 3 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Spinal cord (C72.0) All ages.... 29 39 13 Under 1 year..... 1 1 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 50-54 years..... 3 3 55-59 years..... 3 2 60-64 years..... 3 2 1 65-69 vears..... 6 70-74 vears..... 5 3 75-79 years..... 80-84 years..... 4 85-89 years..... 3 90-94 years..... 2 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cauda equina (C72.1) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Optic nerve (C72.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other and unspecified cranial nerves (C72.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Overlapping lesion of brain and other parts of central nervous system (C72.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Central nervous system, unspecified (C72.9) All ages.... 47 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 3 1 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2 55-59 years..... 2 2 4 60-64 years..... 4 3 1 65-69 vears..... 5 3 70-74 vears..... 5 75-79 years..... 80-84 years..... 5 85-89 years..... 3 2 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of thyroid gland (C73) All ages.... 1,518 1,321 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of adrenal gland (C74) All ages.... Under 1 year..... 1-4 years..... 2.4 2.0 2.7 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 a a 2.7 50-54 years..... 2.2 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Cortex of adrenal gland (C74.0) All ages.... 29 55 29 Under 1 year.... 1-4 years..... 1 5-9 years..... 1 1 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 45-49 years..... 4 1 3 3 2 2 50-54 years..... 14 2 12 10 12 55-59 years..... 5 3 2 2 60-64 years..... 8 5 3 65-69 vears..... 4 70-74 vears..... 2 75-79 years..... 80-84 years..... 1 3 85-89 years..... 3 2 90-94 years..... 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Medulla of adrenal gland (C74.1) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 3 50-54 years..... 2 1 55-59 years..... 2 5 60-64 years..... 2 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 3 3 80-84 years..... 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Adrenal gland, unspecified (C74.9) All ages.... Under 1 year..... 1-4 years..... 2.0 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.2 ρ R 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.8 2.7 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Malignant neoplasm of other endocrine glands and related structures (C75) All ages.... 67 19 10 Under 1 year.... 2 2 2 2 1-4 years..... 2 1 1 1 5-9 years..... 1 1 1 10-14 years..... 15-19 years..... 1 1 20-24 years..... 3 25-29 years..... 30-34 years..... 4 35-39 years..... 4 40-44 years..... 9 45-49 years..... 5 1 16 50-54 years..... 1 0 12 12 55-59 years..... 10 9 4 6 60-64 years..... 15 6 9 13 1 65-69 vears..... 8 5 3 70-74 vears..... 11 10 10 9 75-79 years..... 15 11 80-84 years..... 9 6 8 5 1 85-89 years..... 11 5 6 10 1 90-94 years..... 5 2 2 1 3 3 95-99 years..... 2 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Parathyroid gland (C75.0) All ages..... 17 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 55-59 years..... 2 1 1 60-64 years..... 3 2 1 65-69 vears..... 70-74 vears..... 75-79 years..... 3 3 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pituitary gland (C75.1) All ages.... 35 20 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 2 50-54 years..... 4 2 55-59 years..... 2 2 60-64 years..... 4 1 3 65-69 vears..... 70-74 vears..... 3 75-79 years..... 80-84 years..... 5 2 85-89 years..... 90-94 years..... 3 2 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Craniopharyngeal duct (C75.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pineal gland (C75.3) All ages.... 22 15 Under 1 year..... 2 2 2 1-4 years..... 2 1 1 1 5-9 years..... 1 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 2 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Carotid body (C75.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Aortic body and other paraganglia (C75.5) All ages.... 13 10 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 5 50-54 years..... 5 55-59 years..... 1 1 60-64 years..... 4 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Endocrine gland, unspecified (C75.9) All ages..... 19 19 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 5 2 3 60-64 years..... 4 2 2 65-69 vears..... 5 70-74 vears..... 3 75-79 years..... 80-84 years..... 3 85-89 years..... 4 3 3 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Malignant neoplasm of other and ill-defined sites (C76) All ages.... 1,734 1,508 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Head, face, and neck (C76.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.8 50-54 years..... 2.3 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.8 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Thorax (C76.1) All ages.... 35 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 5 50-54 years..... 5 55-59 years..... 1 1 1 60-64 years..... 5 65-69 vears..... 6 3 70-74 vears..... 13 6 9 75-79 years..... 13 6 80-84 years..... 13 9 13 13 85-89 years..... 6 12 90-94 years..... 6 5 6 95-99 years..... 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Abdomen (C76.2) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.4 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.7 80-84 years..... 85-89 years..... 90-94 years..... 1.0 95-99 years..... 2.7 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Pelvis (C76.3) All ages.... 234 32 202 209 183 18 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 7 50-54 years..... 55-59 years..... 6 6 6 6 60-64 years..... 14 11 10 8 65-69 vears..... 22 18 18 3 70-74 vears..... 22 18 19 15 75-79 years..... 36 2.7 30 2.4 4 3 80-84 years..... 41 2 39 41 39 36 85-89 years..... 33 33 30 90-94 years..... 26 2 24 2.6 24 95-99 years..... 9 9 9 9 100 years & over. 1 1 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Upper limb (C76.4) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 3 90-94 years..... 1 95-99 years..... 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lower limb (C76.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 5 2 85-89 years..... 90-94 years..... 3 1 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other ill-defined sites (C76.7) All ages..... 103 38 10 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 1 1 1 8 50-54 years..... 55-59 years..... 7 2 5 5 60-64 years..... 9 5 9 65-69 vears..... 12 5 70-74 vears..... 13 9 75-79 years..... 17 10 12 3 80-84 years..... 24 14 10 2.1 13 2 3 85-89 years..... 18 11 7 15 10 5 7 90-94 years..... 2 5 1 1 6 2 95-99 years..... 3 2 1 1 2 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Overlapping lesion of other and ill-defined sites (C76.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 2 90-94 years..... 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Secondary and unspecified malignant neoplasm of lymph nodes (C77) All ages.... 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 2 55-59 years..... 1 1 60-64 years..... 4 3 1 65-69 vears..... 2 70-74 vears..... 6 3 3 75-79 years..... 4 80-84 years..... 2 85-89 years..... 3 90-94 years..... 4 2 2 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis	eases,											As	sian or		
Tenth Revision, Second		All rac	es		White			Black		Amer	rican Indi			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Lymph nodes of head,	face, and ne	eck (C77.0)												
All ages	6	4	2	6	4	2	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
65-69 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
70-74 years	1	-	1	1	_	1	-	-	-	_	_	-	-	-	-
75-79 years	2	2	-	2	2	-	-	-	-	_	_	-	-	-	-
30-84 years	-	-	-	-	_	-	-	-	-	_	_	-	-	-	-
35-89 years	1	1	-	1	1	-	-	-	-	_	_	-	-	-	-
90-94 years	-	-	-	-	_	-	-	-	-	_	_	-	-	-	-
95-99 years	-	-	-	-	_	-	-	-	-	_	_	-	-	-	-
100 years & over.	-	-	-	-	_	-	-	-	-	_	_	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Intrathoracic lymph nodes (C77.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Intra-abdominal lymph nodes (C77.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Axillary and upper limb lymph nodes (C77.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Inguinal and lower limb lymph nodes (C77.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lymph node, unspecified (C77.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 4 3 1 65-69 vears..... 70-74 vears..... 4 75-79 years..... 2 80-84 years..... 3 85-89 years..... 6 3 90-94 years..... 3 1 2 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Dise	eases,											As	ian or		
Tenth Revision, Second	i	All rac	es		White			Black		Amer	ican Ind	lian 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Secondary malignant ne	eoplasm of r	respirator	y and diges	tive organs	(C78)										
All ages	3,231	1,567	1,664	2,678	1,282	1,396	404	218	186	23	9	14	126	58	68
Under 1 year	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
1-4 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
10-14 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
L5-19 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
20-24 years	3	3	-	1	1	-	1	1	-	1	1	_	-	-	-
25-29 years	5	2	3	3	1	2	2	1	1	-	-	_	-	-	-
30-34 years	7	7	-	7	7	-	-	-	-	-	-	_	-	-	-
35-39 years	21	9	12	13	7	6	5	1	4	-	-	-	3	1	2
40-44 years	50	31	19	36	23	13	9	6	3	-	-	-	5	2	3
15-49 years	94	58	36	70	41	29	18	14	4	1	-	1	5	3	2
50-54 years	175	112	63	123	73	50	40	30	10	2	2	_	10	7	3
55-59 years	210	135	75	150	101	49	45	24	21	1	1	_	14	9	5
60-64 years	269	166	103	206	131	75	48	30	18	3	1	2	12	4	8
65-69 years	300	161	139	248	133	115	35	20	15	5	1	4	12	7	5
70-74 years	412	194	218	345	160	185	49	26	23	3	2	1	15	6	9
75-79 years	513	253	260	441	221	220	49	23	26	2	1	1	21	8	13
30-84 years	525	236	289	461	207	254	50	26	24	2	_	2	12	3	9
35-89 years	394	137	257	354	121	233	27	9	18	_	_	-	13	7	6
90-94 years	197	52	145	170	45	125	21	6	15	3	-	3	3	1	2
95-99 years	47	8	39	41	7	34	5	1	4	-	-	-	1	-	1
100 years & over.	5	1	4	5	1	4	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Secondary malignant neoplasm of lung (C78.0) All ages.... 192 322 151 171 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 2 4 15 50-54 years..... 6 9 14 55-59 years..... 19 10 9 15 8 60-64 years..... 31 18 13 23 11 12 65-69 vears..... 33 16 17 26 12 70-74 vears..... 51 29 22 44 26 18 75-79 years..... 73 44 29 66 39 2.7 80-84 years..... 77 37 40 69 32 37 6 5 1 1 32 85-89 years..... 8 24 29 22 1 1 90-94 years..... 23 15 21 14 1 8 95-99 years..... 7 7 7 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Secondary malignant neoplasm of mediastinum (C78.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Tenth Revision, Second Roltin, 2004) Both and age sexes Male Female Both Both and age sexes Male Female Both Both and age sexes Male Female Both Both Both and age sexes Male Female Both Both Both Both Both Both Both Both	Cause of death (based on the International Classification of Dis												As	ian or		
Secondary malignant neoplasm of pleura (C78.2) ***Rate** Sample Sample	Tenth Revision, Second	d	All rac	es		White			Black		Amei	rican Indi	ian 1/	Pacif	ic Isla	nder 2/
Secondary malignant neoplasm of pleura (C78.2) All ages		Both			Both			Both			Both			Both		
All ages	and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Female	sexes	Male	Female
Under 1 years	Secondary malignant n	eoplasm of p	oleura (C7	8.2)												
1-4 years	All ages	138	62	76	121	54	67	14	7	7	_	_	_	3	1	2
5-9 years	Under 1 year	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
10-14 years		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years 1 1 1 - 1 1 1		-	-	-	-	-	-	-	-	-	-	-	-	-		-
45-49 years		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years 1 1 1 - 1 1 1		1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years 1 1 1 - 1 1 1		_	_		_	_	-	-	-	-	-	-	-	-	-	-
60-64 years 2 2 2 - 2 2		1	1		1	1				_	_	-	-	_		_
65-69 years 6 3 3 3 5 2 3 1 1 1		1	1		1	1				_	_	_	-	_		_
70-74 years 18 5 13 13 3 10 4 2 2 1 - 1 75-79 years 19 9 10 17 9 8 2 - 2 80-84 years 28 15 13 23 12 11 3 2 1 2 1 1 98-89 years 35 17 18 32 16 16 3 1 2 90-94 years 20 8 12 19 7 12 1 1 95-99 years 6 - 6 6 - 6 6 - 6 6			2					_	-		_	_	-	_		-
75-79 years 19 9 10 17 9 8 2 - 2 80-84 years 28 15 13 23 12 11 3 2 1 2 1 1 1 85-89 years 35 17 18 32 16 16 3 1 2 90-94 years 20 8 12 19 7 12 1 1 95-99 years 6 - 6 6 - 6 6 - 6 6		-	3		-	_		1	1		_	_		_		
80-84 years 28 15 13 23 12 11 3 2 1 2 1 1 85-89 years 35 17 18 32 16 16 3 1 2 90-94 years 20 8 12 19 7 12 1 1 95-99 years 6 - 6 6 - 6 6 6			5					_			_	-		1		_
85-89 years 35 17 18 32 16 16 3 1 2			-								_	-		_		
90-94 years 20 8 12 19 7 12 1 1 95-99 years 6 - 6 6 - 6									2		_	-		2		
95-99 years 6 - 6 6 - 6								3	1	2	_	-	-	_	-	-
			8			,		1	1	_	_	_	-	_	-	-
	95-99 years 100 years & over.	6	_	6	6	_	6	_	-	_	_	_	-	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Secondary malignant neoplasm of other and unspecified respiratory organs (C78.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Secondary malignant neoplasm of small intestine (C78.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis	seases,											As	ian or		
Tenth Revision, Secor	nd	All race	es		White			Black		Ame	rican Indi	an 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Secondary malignant r	neoplasm of l	arge intes	stine and re	ctum (C78.5)										
All ages	14	6	8	14	6	8	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	_
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
75-79 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
80-84 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
85-89 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
				_			_	_				_			_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Secondary malignant neoplasm of retroperitoneum and peritoneum (C78.6) All ages.... 120 281 12 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 6 40-44 years..... 3 45-49 years..... 14 6 8 13 6 24 50-54 years..... 29 12 1 0 17 14 55-59 years..... 27 8 19 22 7 15 4 1 60-64 years..... 39 18 21 31 16 15 2 65-69 years..... 42 29 37 10 27 13 70-74 years..... 58 14 51 10 41 3 75-79 years..... 85 2.0 65 77 18 59 80-84 years..... 79 25 54 69 23 46 8 2 41 85-89 years..... 34 39 6 33 1 90-94 years..... 23 6 17 21 15 2 2 6 95-99 years..... 4 3 3 1 4 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Secondary malignant neoplasm of liver (C78.7) All ages.... 2,170 1,134 1,036 1,761 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 0 2.7 50-54 years..... ρ 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Secondary malignant neoplasm of other and unspecified digestive organs (C78.8) All ages.... 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 50-54 years..... 3 3 1 55-59 years..... 7 2 5 60-64 years..... 12 5 65-69 vears..... 6 70-74 vears..... 8 3 5 75-79 years..... 7 80-84 years..... 14 5 2 3 85-89 years..... 5 4 1 90-94 years..... 1 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based															
on the International Classification of Dis	eageg											Δα	ian or		
Tenth Revision, Secon		All rac	es		White			Black		Amer	ican Ind			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Secondary malignant n	eoplasm of d	other site	s (C79)												
All ages	4,354	2,776	1,578	3,615	2,297	1,318	642	423	219	22	10	12	75	46	29
Under 1 year	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
1-4 years	2	-	2	1	-	1	-	-	-	-	_	-	1	-	1
5-9 years	3	1	2	2	-	2	1	1	-	-	_	-	-	-	-
10-14 years	1	1	-	1	1	-	-	-	-	-	_	-	-	-	-
15-19 years	5	3	2	2	1	1	3	2	1	-	_	-	-	-	-
20-24 years	5	4	1	5	4	1	-	_	-	-	-	-	-	-	-
25-29 years	13	9	4	10	8	2	3	1	2	-	-	-	-	-	-
30-34 years	7	4	3	5	3	2	2	1	1	-	-	-	-	-	-
35-39 years	36	17	19	21	9	12	11	6	5	1	_	1	3	2	1
40-44 years	104	69	35	82	56	26	22	13	9	-	-	-	-	-	-
45-49 years	212	159	53	158	119	39	46	35	11	2	1	1	6	4	2
50-54 years	342	262	80	249	193	56	83	63	20	1	-	1	9	6	3
55-59 years	492	380	112	371	290	81	113	84	29	3	2	1	5	4	1
60-64 years	482	356	126	389	290	99	79	55	24	6	4	2	8	7	1
65-69 years	472	325	147	379	261	118	85	59	26	3	1	2	5	4	1
70-74 years	506	318	188	432	274	158	62	40	22	3	1	2	9	3	6
75-79 years	567	341	226	495	300	195	60	33	27	1	1	-	11	7	4
80-84 years	515	280	235	480	262	218	28	14	14	-	_	-	7	4	3
85-89 years	353	166	187	316	150	166	27	13	14	2	_	2	8	3	5
90-94 years	185	67	118	170	63	107	12	2	10	-	-	-	3	2	1
95-99 years	44	12	32	40	11	29	4	1	3	-	-	-	-	-	-
100 years & over.	8	2	6	7	2	5	1	-	1	-	-	-	-	-	-
Age not stated	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Secondary malignant neoplasm of kidney and renal pelvis (C79.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 2 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Secondary malignant neoplasm of bladder and other and unspecified urinary organs (C79.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 2 2 60-64 years..... 3 2 65-69 vears..... 70-74 vears..... 4 75-79 years..... 80-84 years..... 5 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis	eases,	211			White			Black			7		sian or		d 2 /
Tenth Revision, Secon		All race	es	D - 1-1-	wnite		D - 1-1-	втаск			rican Indi	an I/		ic isia	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
Secondary malignant n	eoplasm of s	kin (C79.2	2)												
All ages	10	7	3	9	6	3	_	_	_	_	_	_	1	1	_
Jnder 1 year	-	-	-	-	_	-	-	-	-	-	_	-	-	-	-
-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-19 years	-	-	-	-	_	-	-	-	-	-	_	-	-	-	-
0-24 years	-	-	-	-	_	-	-	-	-	-	_	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	1	1	-	-	-	-	-	-	-	-	-	-	1	1	-
0-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	2	2	-	2	2	-	-	-	-	-	_	-	-	-	-
5-79 years	1	-	1	1	_	1	-	-	-	-	_	-	-	-	-
0-84 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
5-89 years	3	3	-	3	3	-	-	-	-	_	-	-	-	-	-
0-94 years	1	-	1	1	-	1	-	_	-	-	-	-	-	-	-
5-99 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
00 years & over.	_	-	-	_	-	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Secondary malignant neoplasm of brain and cerebral meninges (C79.3) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 1 0 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Secondary malignant neoplasm of other and unspecified parts of nervous system (C79.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 80-84 years..... 2 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Secondary malignant neoplasm of bone and bone marrow (C79.5) All ages.... 202 171 59 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 1 30-34 years..... 1 35-39 years..... 40-44 years..... 3 2 6 45-49 years..... 14 11 3 Ω 2 50-54 years..... 13 1.0 3 3 11 R 1 55-59 years..... 26 19 7 20 14 6 5 60-64 years..... 36 25 11 29 20 65-69 years..... 28 17 11 21 14 7 70-74 years..... 48 21 20 6 75-79 years..... 66 40 26 53 33 2.0 6 80-84 years..... 88 50 38 82 46 36 4 3 1 2 1 71 27 61 37 85-89 years..... 44 2.4 5 90-94 years..... 38 13 25 33 11 22 4 3 1 1 1 95-99 years..... 7 5 1 6 6 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis- Tenth Revision, Second		All rac	10.5		White			Black		Amor	rican Ind		ian or	ia Tala	nder 2/
Edition, 2004)	a Both	All rac	es	Both	willte		Both	Black		Both	rican ino	ııan ı/	Both	ic isia	naer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Secondary malignant n	eoplasm of d	ovary (C79	.6)												
All ages	7		7	5		5	1		1	-		_	1		1
Under 1 year	-		-	-		-	_		-	-		-	-		-
1-4 years	-		-	-		-	-		-	-		-	-		-
5-9 years	-		-	-		-	-		-	-		-	-		-
10-14 years	-		-	-		-	-		-	-		-	-		-
15-19 years	-		-	-		-	-		-	-		-	-		-
20-24 years	-		-	-		-	-		-	-		-	-		-
25-29 years	-		-	-		-	-		-	-		-	-		-
30-34 years	-		-	-		-	-		-	-		-	-		-
35-39 years	-		-	-		-	-		-	-		-	-		-
40-44 years	-		-	-		-	-		-	-		-	-		-
45-49 years	1		1	-		-	1		1	-		-	-		-
50-54 years	1		1	-		-	-		-	-		-	1		1
55-59 years	-		-	-		-	-		-	-		-	-		-
60-64 years	1		1	1		1	-		-	-		-	-		-
65-69 years	2		2	2		2	-		-	-		-	-		-
70-74 years	1		1	1		1	_		-	_		-	-		-
75-79 years	_		_	_		_	_		-	-		-	_		-
80-84 years	1		1	1		1	_		-	-		-	_		-
85-89 years	_		_	_		_	_		-	-		-	_		-
90-94 years	_		_	_		_	_		-	-		-	_		-
95-99 years	_		_	_		_	_		-	-		-	_		-
100 years & over.	_		_	_		_	_		-	-		-	_		-
Age not stated	_		_	_		_	_		_	_		_	_		_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Secondary malignant neoplasm of adrenal gland (C79.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International												•			
Classification of Dis Tenth Revision, Secon		All rac	0.5		White			Black		Amore	ican Indi		ian or	ia Talo	nder 2/
Edition, 2004)	.a Both	All rac	es	Both	wille		Both	Black		Both	ican indi	Lan 1/	Both	ic ista	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Temale	sexes	Male	Female
Secondary malignant n	eoplasm of	other spec	ified sites	(C79.8)											
All ages	3,132	2,165	967	2,578	1,774	804	489	349	140	16	9	7	49	33	16
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
20-24 years	5	4	1	5	4	1	-	-	-	-	-	-	-	-	-
25-29 years	7	5	2	6	5	1	1	-	1	-	-	-	-	-	-
30-34 years	6	4	2	4	3	1	2	1	1	-	-	-	-	-	-
35-39 years	25	14	11	14	8	6	8	4	4	1	-	1	2	2	-
40-44 years	82	57	25	62	45	17	20	12	8	-	-	-	-	-	-
45-49 years	162	133	29	125	102	23	32	27	5	1	1	-	4	3	1
50-54 years	294	236	58	217	177	40	72	56	16	-	-	-	5	3	2
55-59 years	416	336	80	311	254	57	97	76	21	3	2	1	5	4	1
60-64 years	377	294	83	307	240	67	59	45	14	4	3	1	7	6	1
65-69 years	384	280	104	310	226	84	67	50	17	3	1	2	4	3	1
70-74 years	361	243	118	306	208	98	46	31	15	3	1	2	6	3	3
75-79 years	376	236	140	325	204	121	42	26	16	1	1	-	8	5	3
80-84 years	308	166	142	286	154	132	18	9	9	-	-	-	4	3	1
85-89 years	203	107	96	184	97	87	16	9	7	-	-	-	3	1	2
90-94 years	97	39	58	91	38	53	5	1	4	-	-	-	1	-	1
95-99 years	24	8	16	21	7	14	3	1	2	-	-	-	-	-	-
100 years & over.	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International	l														
Classification of Dis	seases.											As	ian or		
Tenth Revision, Secon		All rac	es		White			Black		Amer	ican Ind			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Malignant neoplasm wi	thout speci	fication c	of site (C80)											
All ages	25,989	12,875	13,114	22,536	11,254	11,282	2,858	1,358	1,500	139	53	86	456	210	246
Under 1 year	7	4	3	7	4	3	-	-	-	-	-	-	-	-	-
1-4 years	7	4	3	5	3	2	1	1	-	-	-	-	1	-	1
5-9 years	6	1	5	3	-	3	2	1	1	1	-	1	-	-	-
10-14 years	10	4	6	6	3	3	4	1	3	-	-	-	-	-	-
15-19 years	20	16	4	16	12	4	2	2	-	-	-	-	2	2	-
20-24 years	42	25	17	35	22	13	5	2	3	-	-	-	2	1	1
25-29 years	72	39	33	51	31	20	16	6	10	-	-	-	5	2	3
30-34 years	91	41	50	67	30	37	15	6	9	2	1	1	7	4	3
35-39 years	174	72	102	137	60	77	25	10	15	3	1	2	9	1	8
40-44 years	443	195	248	336	156	180	86	33	53	5	1	4	16	5	11
45-49 years	851	433	418	681	339	342	142	83	59	6	2	4	22	9	13
50-54 years	1,362	738	624	1,084	589	495	238	129	109	14	5	9	26	15	11
55-59 years	2,085	1,164	921	1,713	961	752	322	175	147	11	3	8	39	25	14
60-64 years	2,464	1,402	1,062	2,063	1,199	864	343	174	169	18	10	8	40	19	21
65-69 years	2,859	1,570	1,289	2,420	1,356	1,064	367	179	188	16	6	10	56	29	27
70-74 years	3,351	1,761	1,590	2,928	1,556	1,372	363	182	181	20	7	13	40	16	24
75-79 years	3,821	1,930	1,891	3,406	1,751	1,655	337	148	189	18	8	10	60	23	37
80-84 years	3,816	1,776	2,040	3,473	1,625	1,848	269	118	151	15	6	9	59	27	32
85-89 years	2,768	1,152	1,616	2,507	1,040	1,467	203	83	120	7	3	4	51	26	25
90-94 years	1,341	446	895	1,246	427	819	79	16	63	2	-	2	14	3	11
95-99 years	360	95	265	317	85	232	37	8	29	-	-	-	6	2	4
100 years & over.	38	6	32	34	4	30	2	1	1	1	_	1	1	1	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Hodgkin's disease (C81) All ages.... 1,327 1,160 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1.3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 0 a 50-54 years..... 2.4 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lymphocytic predominance (C81.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Nodular sclerosis (C81.1) All ages..... 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Diseases, Tenth Revision, Second All races								Asian or								
		All races		White			Black				rican Indi	an 1/	Pacific Islander 2,			
Edition, 2004)	Both	_	_	Both	_	_	Both	_	_	Both	_	_	Both	_	_	
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Mixed cellularity (C8	1.2)															
All ages	5	3	2	5	3	2	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	_	-	-	-	-	-	-	_	-	_	-	-	
15-19 years	-	-	-	_	-	-	-	-	-	-	_	-	_	-	-	
20-24 years	-	-	-	_	-	-	-	-	-	-	_	-	_	-	-	
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
35-39 years	-	-	-	_	_	-	-	_	-	-	-	-	-	-	-	
40-44 years	-	-	-	_	_	-	-	_	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
50-54 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	_	
60-64 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
70-74 years	-	-	-	_	-	-	-	_	-	_	-	-	-	-	-	
75-79 years	1	_	1	1	-	1	-	-	-	_	_	-	_	_	-	
30-84 years	2	1	1	2	1	1	-	_	-	_	_	-	_	_	-	
85-89 years	1	1	-	1	1	_	-	-	-	_	_	-	_	_	-	
90-94 years	_	-	-	_	-	_	-	_	-	_	_	-	_	_	-	
95-99 years	_	_	-	-	-	_	-	-	-	_	_	-	_	_	-	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lymphocytic depletion (C81.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other Hodgkin's disease (C81.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Hodgkin's disease, unspecified (C81.9) All ages.... 1,303 1,137 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1.3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 Ω a 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Follicular [nodular] non-Hodgkin's lymphoma (C82) All ages.... 267 135 132 16 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 3 9 45-49 years..... 5 Ω 5 3 15 50-54 years..... ρ 1 14 55-59 years..... 20 14 19 6 14 5 1 60-64 years..... 35 18 17 30 16 14 65-69 vears..... 29 15 28 14 70-74 vears..... 47 27 20 18 3 75-79 years..... 41 18 23 39 17 2.2 80-84 years..... 48 23 25 46 23 23 1 25 25 17 85-89 years..... 8 17 8 1 90-94 years..... 8 3 5 1 3 4 95-99 years..... 3 3 3 1 4 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Small cleaved cell, follicular (C82.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mixed small cleaved and large cell, follicular (C82.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Large cell, follicular (C82.2) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other types of follicular non-Hodgkin's lymphoma (C82.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Follicular non-Hodgkin's lymphoma, unspecified (C82.9) All ages.... 246 124 122 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 3 45-49 years..... 8 4 3 50-54 years..... 15 ρ 1 14 55-59 years..... 19 18 14 5 14 1 60-64 years..... 32 15 17 28 14 14 65-69 vears..... 26 12 25 13 12 14 70-74 vears..... 42 18 39 23 16 75-79 years..... 38 17 21 36 16 2.0 80-84 years..... 44 20 24 42 2.0 22 1 15 85-89 years..... 23 8 15 23 8 7 90-94 years..... 3 4 3 4 95-99 years..... 4 3 3 3 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Diffuse non-Hodgkin's lymphoma (C83) All ages.... 2,164 1,294 2,010 1,199 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 2.8 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Small cell (diffuse) (C83.0) All ages..... 126 199 110 19 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 7 50-54 years..... 3 3 4 55-59 years..... 13 9 2 10 3 60-64 years..... 9 2 5 2 65-69 vears..... 22 19 18 4 70-74 vears..... 30 17 13 27 13 75-79 years..... 39 23 38 2.2 16 16 80-84 years..... 39 22 17 37 21 16 2 1 30 28 85-89 years..... 13 17 12 16 90-94 years..... 21 15 2.0 1 6 6 14 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Small cleaved cell (diffuse) (C83.1) All ages.... 172 589 423 166 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 3 2 45-49 years..... 12 9 3 11 Ω 31 50-54 years..... 34 29 5 26 55-59 years..... 40 36 38 35 4 3 60-64 years..... 84 66 18 81 63 18 65-69 years..... 66 48 18 62 46 16 70-74 years..... 106 67 39 100 38 75-79 years..... 101 73 28 100 72 28 80-84 years..... 94 68 26 93 67 26 1 51 85-89 years..... 52 31 21 31 20 2 90-94 years..... 19 11 8 17 10 1 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mixed small and large cell (diffuse) (C83.2) All ages..... 16 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 2 1 1 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 2 75-79 years..... 3 80-84 years..... 2 1 2 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Large cell (diffuse) (C83.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 1 0 50-54 years..... 2.2 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 8.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Immunoblastic (diffuse) (C83.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Lymphoblastic (diffuse) (C83.5) All ages.... 25 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 4 1 3 1 15-19 years..... 3 2 1 20-24 years..... 3 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 1 7 50-54 years..... 3 1 55-59 years..... 5 4 1 60-64 years..... 3 3 65-69 vears..... 70-74 vears..... 7 5 75-79 years..... 80-84 years..... 2 1 2 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International	l														
Classification of Dis	seases,					Asian or									
Tenth Revision, Secon		All races			White		Black			Amer	rican Indi		Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Undifferentiated (dif	fuse) (C83.6)													
All ages	3	3	_	3	3	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Cause of death (based on the International Classification of Dis												As	ian or		
Tenth Revision, Secon	ıd	All races		White			Black			American Indian 1/				ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Burkitt's tumor (C83.	7)														
All ages	244	166	78	221	152	69	18	12	6	-	-	-	5	2	3
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	5	5	-	5	5	-	-	-	-	-	-	-	-	-	-
10-14 years	3	3	-	3	3	-	-	-	-	-	-	-	-	-	-
15-19 years	6	5	1	5	4	1	1	1	-	-	-	-	-	-	-
20-24 years	5	5	-	4	4	-	1	1	-	-	-	-	-	-	-
25-29 years	9	5	4	8	4	4	1	1	-	-	-	-	-	-	-
30-34 years	7	3	4	5	3	2	2	-	2	-	-	-	-	-	-
35-39 years	9	7	2	7	6	1	2	1	1	-	-	-	-	-	-
40-44 years	11	8	3	9	7	2	2	1	1	-	-	-	-	-	-
45-49 years	21	11	10	18	9	9	3	2	1	-	-	-	-	-	-
50-54 years	16	13	3	14	11	3	2	2	-	-	-	-	-	-	-
55-59 years	24	16	8	23	15	8	-	-	-	-	-	-	1	1	-
60-64 years	20	14	6	17	13	4	1	_	1	-	-	-	2	1	1
65-69 years	26	19	7	26	19	7	-	_	-	-	-	-	-	-	-
70-74 years	27	18	9	25	17	8	1	1	-	-	-	-	1	-	1
75-79 years	19	10	9	18	9	9	1	1	-	-	-	-	-	-	-
80-84 years	22	14	8	20	13	7	1	1	-	-	-	-	1	-	1
85-89 years	12	8	4	12	8	4	-	_	-	-	-	-	-	-	-
90-94 years	2	2	-	2	2	-	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disc												λα	sian or			
Tenth Revision, Second					White		Black			American Indian 1/ Pacific Islander						
Edition, 2004)	Both	1111 1401		Both	,,,,,,		Both	Diaon		Both	. 10411 11141		Both	10 1010	ilidol 2/	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female	
Other types of diffuse	e non-Hodgki	n's lympho	oma (C83.8)													
All ages	26	10	16	24	9	15	2	1	1	_	_	_	_	_	_	
Inder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-14 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-	
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
0-24 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-	
5-29 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-	
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-39 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-	
0-44 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-	
5-49 years	1	1	-	-	-	_	1	1	_	-	-	-	-	-	-	
0-54 years	-	-	-	-	-	_	_	_	_	-	-	-	-	-	-	
5-59 years	3	1	2	3	1	2	_	_	_	-	-	-	-	-	-	
0-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
5-69 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-	
0-74 years	4	3	1	4	3	1	_	_	_	_	_	-	-	-	-	
5-79 years	4	2	2	3	2	1	1	_	1	_	_	-	-	-	-	
0-84 years	4	2	2	4	2	2	_	_	_	_	_	-	-	-	-	
5-89 years	4	-	4	4	-	4	_	_	_	_	_	-	-	-	-	
0-94 years	1	-	1	1	-	1	_	_	_	_	_	-	-	-	-	
5-99 years	-	-	-	-	-	_	_	_	_	_	_	-	-	-	-	
00 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis	seases,											As	ian or		
Tenth Revision, Secon		All races			White			Black		American Indian 1/			Pacific Islander 2		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale 	sexes	Male	Female
Diffuse non-Hodgkin's	s lymphoma, u	nspecifie	d (C83.9)												
All ages	30	19	11	25	14	11	4	4	_	_	_	_	1	1	_
Under 1 year	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
45-49 years	_	-	-	-	-	-	_	_	-	_	-	-	-	-	-
50-54 years	3	-	3	3	-	3	_	_	-	_	-	-	-	-	-
55-59 years	1	1	-	1	1	-	_	_	-	_	-	-	-	-	-
50-64 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
65-69 years	2	1	1	1	-	1	1	1	-	_	-	-	-	-	-
70-74 years	4	4	-	3	3	-	-	_	-	-	-	-	1	1	-
75-79 years	11	7	4	11	7	4	-	_	-	-	-	-	-	-	-
30-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-89 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
90-94 years	2	-	2	2	-	2	-	_	-	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Peripheral and cutaneous T-cell lymphomas (C84) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 2.6 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mycosis fungoides (C84.0) All ages..... 77 20 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 3 1 2 7 50-54 years..... 6 1 1 55-59 years..... 6 3 3 1 60-64 years..... 5 1 65-69 vears..... 5 3 3 70-74 vears..... 7 6 75-79 years..... 8 11 3 80-84 years..... 17 13 13 11 2 6 2 85-89 years..... 4 5 3 1 90-94 years..... 2 2 4 2 2 95-99 years..... 2 2 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac			White			Black		Asian or American Indian 1/ Pacific Islander 2/							
		All rac	es	Dath	WIIICE		Dath	Black			rican indi	an 1/		IC ISIA	nder 2/		
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	'emale	Both sexes	Male	Female		
Sezary's disease (C84	.1)																
All ages	14	10	4	9	7	2	5	3	2	_	_	_	_	_	_		
Jnder 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-		
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
0-44 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-		
5-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
0-54 years	1	1	-	-	_	-	1	1	-	-	-	-	-	-	-		
55-59 years	1	1	-	-	_	-	1	1	-	-	-	-	-	-	-		
0-64 years	3	3	-	3	3	-	-	-	-	-	-	-	-	-	-		
5-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
0-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
5-79 years	2	1	1	1	1	_	1	-	1	_	_	-	-	-	-		
0-84 years	3	2	1	2	1	1	1	1	-	_	_	-	-	-	-		
5-89 years	1	1	_	1	1	_	_	-	-	_	_	-	-	-	-		
0-94 years	-	_	-	-	_	-	-	-	-	-	-	-	-	-	-		
5-99 years	-	_	-	-	_	-	-	-	-	-	-	-	-	-	-		
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes T-zone lymphoma (C84.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Peripheral T-cell lymphoma (C84.4) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 1 1 1 6 50-54 years..... 2 55-59 years..... 4 3 1 2 60-64 years..... 13 9 11 65-69 vears..... 11 5 6 70-74 vears..... 16 8 8 75-79 years..... 21 12 9 80-84 years..... 9 5 9 5 85-89 years..... 4 3 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other and unspecified T-cell lymphomas (C84.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 0 2.2 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Male Female sexes Male Female sexes Male Female sexes Male sexes Male Female sexes Other and unspecified types of non-Hodgkin's lymphoma (C85) All ages.... 17,179 8,950 8,229 15,606 8,103 7,503 1,161 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2.5 20-24 vears..... 25-29 years..... 2.0 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1,326 1,166 1,620 1,442 65-69 vears..... 70-74 years..... 2,132 1,164 1,958 1,080 75-79 years..... 2,900 1,490 1,410 2,721 1,407 1,314 80-84 years..... 3,084 1,481 1,603 2,912 1,413 1.499 1,305 1,243 85-89 years..... 2,231 2,116 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lymphosarcoma (C85.0) All ages.... 29 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 3 3 60-64 years..... 3 2 65-69 vears..... 70-74 vears..... 2 75-79 years..... 3 80-84 years..... 3 85-89 years..... 4 90-94 years..... 1 1 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes B-cell lymphoma, unspecified (C85.1) All ages.... 1,827 1,640 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified types of non-Hodgkin's lymphoma (C85.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female sexes Male Female sexes Male sexes Male sexes Male Female sexes Non-Hodgkin's lymphoma, unspecified type (C85.9) All ages.... 15,317 7,998 7,319 13,933 7,260 6,673 1,033 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2.4 2.0 20-24 vears..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 50-54 years..... 1.0 55-59 years..... 60-64 years..... 1,176 1,040 1,462 1,302 65-69 vears..... 70-74 years..... 1,915 1,052 1,762 75-79 years..... 2,595 1,327 1,268 2,437 1,255 1,182 2.7 80-84 years..... 2.739 1,312 1,427 2,586 1,251 1,335 1,974 1,136 85-89 years..... 1,877 1,081 90-94 years..... 95-99 years..... 100 years & over. 1.0 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant immunoproliferative diseases (C88) All ages.... 135 312 181 131 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 2 2 2 5 50-54 years..... 2 3 3 55-59 years..... 12 11 6 6 5 60-64 years..... 17 14 3 16 13 3 65-69 vears..... 26 20 23 19 3 6 70-74 vears..... 27 13 14 24 11 13 75-79 years..... 64 42 2.2 62 40 2.2 80-84 years..... 69 41 28 66 39 27 65 37 64 85-89 years..... 28 36 28 90-94 years..... 30 12 18 30 12 18 95-99 years..... 5 2 3 5 2 3 100 years & over. 2 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Waldenstrom's macroglobulinemia (C88.0) All ages..... 135 180 131 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 2 2 2 5 50-54 years..... 2 3 3 55-59 years..... 12 6 11 6 5 60-64 years..... 16 13 3 15 12 3 65-69 vears..... 26 20 23 19 3 6 70-74 vears..... 27 13 14 24 11 13 75-79 years..... 64 42 2.2 62 40 2.2 80-84 years..... 69 41 28 66 39 27 65 37 64 85-89 years..... 28 36 28 90-94 years..... 30 12 18 30 12 18 95-99 years..... 5 2 3 5 2 3 100 years & over. 2 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Immunoproliferative small intestinal disease (C88.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based	l														
on the International												_			
Classification of Dis								-1 1		_			ian or		3 0 /
Tenth Revision, Secon		All rac	es		White			Black			ican In	dian 1/		fic Isla	nder 2/
Edition, 2004)	Both	24.1.		Both	M - 1 -		Both	24 - 7 -	m1.	Both	M . 1 .	m1.	Both	M - 7 -	m1.
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	мате	Female	sexes	мате	Female
Multiple myeloma and	malignant pl	lasma cell	neoplasms	(C90)											
All ages	10,784	5,690	5,094	8,726	4,662	4,064	1,834	920	914	49	25	24	175	83	92
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
25-29 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
30-34 years	10	7	3	6	4	2	3	3	-	-	-	-	1	-	1
35-39 years	28	17	11	15	9	6	13	8	5	-	-	-	-	-	-
10-44 years	121	71	50	92	57	35	27	12	15	1	1	-	1	1	-
15-49 years	240	134	106	153	90	63	74	40	34	1	-	1	12	4	8
50-54 years	482	284	198	338	208	130	128	68	60	6	3	3	10	5	5
55-59 years	771	444	327	572	336	236	182	100	82	2	2	-	15	6	9
0-64 years	989	571	418	741	428	313	221	126	95	9	6	3	18	11	7
55-69 years	1,197	671	526	942	541	401	231	116	115	5	4	1	19	10	9
70-74 years	1,593	865	728	1,279	703	576	272	152	120	11	2	9	31	8	23
'5-79 years	1,841	989	852	1,554	851	703	261	122	139	3	-	3	23	16	7
30-84 years	1,819	913	906	1,572	800	772	222	99	123	9	5	4	16	9	7
35-89 years	1,183	520	663	1,024	460	564	136	50	86	1	1	_	22	9	13
90-94 years	434	179	255	372	155	217	54	19	35	1	1	_	7	4	3
95-99 years	67	22	45	58	18	40	9	4	5	-	-	_	-	-	-
100 years & over.	6	1	5	6	1	5	_	_	_	_	-	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Multiple myeloma (C90.0) All ages.... 10,599 5,585 5,014 8,572 4,576 3,996 1,808 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1,175 65-69 vears..... 70-74 vears..... 1,571 1,264 75-79 years..... 1,818 1,532 80-84 years..... 1.792 1.549 1,170 1,012 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Plasma cell leukemia (C90.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 8 2 45-49 years..... 2 7 50-54 years..... 5 55-59 years..... 5 2 3 5 60-64 years..... 10 3 65-69 vears..... 14 10 70-74 vears..... 8 4 75-79 years..... 4 80-84 years..... 8 5 1 2 85-89 years..... 4 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Plasmacytoma, extramedullary (C90.2) All ages..... 40 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 1 5 50-54 years..... 3 55-59 years..... 16 14 11 5 60-64 years..... 12 6 6 10 65-69 vears..... 8 5 70-74 vears..... 14 8 6 9 75-79 years..... 19 9 10 80-84 years..... 19 10 9 16 2 1 9 85-89 years..... 4 5 8 90-94 years..... 6 5 1 6 95-99 years..... 1 1 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Lymphoid leukemia (C91) All ages.... 6,395 3,748 2,647 5,750 3,352 2,398 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1.0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1.027 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International															
Classification of Dis	22505											7.0	ian or		
Tenth Revision, Second		All rac	100		White			Black		Δmer	ican Indi			ic Isla	nder 2/
Edition, 2004)	Both	AII IAC	.05	Both	WIIICC		Both	DIACK		Both	ican inai	an i/	Both	.IC 151a	iluci 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Acute lymphoblastic le	eukemia (C91	.0)													
All ages	1,393	770	623	1,228	673	555	117	67	50	6	5	1	42	25	17
Under 1 year	9	6	3	8	5	3	1	1	-	-	-	-	-	-	-
1-4 years	44	25	19	40	21	19	1	1	-	1	1	-	2	2	-
5-9 years	52	30	22	43	23	20	8	6	2	-	-	-	1	1	-
10-14 years	69	40	29	57	34	23	7	3	4	-	-	-	5	3	2
15-19 years	92	57	35	77	46	31	12	8	4	-	-	-	3	3	-
20-24 years	107	76	31	89	64	25	15	11	4	1	1	-	2	-	2
25-29 years	64	41	23	62	41	21	2	-	2	-	-	-	-	-	-
30-34 years	69	43	26	58	34	24	7	5	2	1	1	-	3	3	-
35-39 years	62	37	25	53	31	22	7	4	3	-	_	-	2	2	-
40-44 years	66	36	30	57	29	28	9	7	2	-	_	-	_	-	-
45-49 years	77	39	38	71	37	34	4	1	3	_	-	-	2	1	1
50-54 years	80	42	38	70	35	35	6	5	1	1	1	-	3	1	2
55-59 years	95	44	51	85	42	43	8	1	7	1	1	-	1	_	1
60-64 years	89	48	41	78	44	34	5	2	3	-	-	-	6	2	4
65-69 years	86	46	40	75	41	34	7	4	3	1	_	1	3	1	2
70-74 years	87	43	44	80	38	42	5	3	2	_	-	-	2	2	-
75-79 years	86	51	35	77	44	33	6	4	2	_	_	_	3	3	_
80-84 years	79	41	38	76	40	36	2	_	2	_	_	_	1	1	_
85-89 years	50	19	31	43	18	25	4	1	3	_	_	_	3	_	3
90-94 years	18	3	15	17	3	14	1	_	1	_	_	_	_	_	_
95-99 years	11	3	8	11	3	8	-	-	-	_	-	-	_	_	-
100 years & over.	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Female Male Female Male Female sexes Male Female sexes sexes Male sexes sexes Chronic lymphocytic leukemia (C91.1) All ages.... 4,498 2,659 1,839 4,077 2,399 1,678 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.0 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Subacute lymphocytic leukemia (C91.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Prolymphocytic leukemia (C91.3) All ages..... 29 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 4 2 6 50-54 years..... 3 3 1 55-59 years..... 8 5 3 60-64 years..... 6 1 65-69 vears..... 7 5 70-74 vears..... 12 5 75-79 years..... 12 12 80-84 years..... 13 8 5 12 8 85-89 years..... 8 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise	eases,											As	ian or		
Tenth Revision, Second	l	All races			White			Black			ican India	an 1/	Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Hairy-cell leukemia (C	291.4)														
All ages	100	80	20	98	78	20	2	2	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	_	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
.0-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
10-44 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
15-49 years	4	3	1	4	3	1	-	-	-	-	-	-	-	-	-
0-54 years	5	4	1	5	4	1	_	-	-	-	-	-	-	-	-
55-59 years	7	7	-	7	7	-	-	-	-	-	-	-	-	-	-
0-64 years	5	5	-	4	4	-	1	1	-	-	-	-	-	-	-
5-69 years	6	4	2	6	4	2	-	-	-	-	-	-	-	-	-
70-74 years	10	8	2	10	8	2	-	_	-	-	-	-	-	-	-
5-79 years	14	14	-	14	14	-	-	-	-	_	-	-	-	-	-
0-84 years	25	21	4	24	20	4	1	1	-	-	-	-	-	-	-
5-89 years	9	7	2	9	7	2	-	_	-	-	-	-	-	-	-
0-94 years	13	5	8	13	5	8	-	_	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Adult T-cell leukemia (C91.5) All ages.... 63 40 Under 1 year.... 1-4 years..... 5-9 years..... 1 2 10-14 years..... 6 4 5 15-19 years..... 3 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 8 5 3 1 60-64 years..... 9 5 3 65-69 vears..... 5 70-74 vears..... 9 5 75-79 years..... 80-84 years..... 8 5 2 85-89 years..... 3 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Lymphoid leukemia, unspecified (C91.9) All ages.... 103 119 92 30 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 2 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 1 50-54 years..... 1.0 8 2 10 55-59 years..... 9 6 7 3 3 60-64 years..... 14 9 5 10 65-69 vears..... 12 6 11 6 70-74 years..... 21 13 8 20 13 7 75-79 years..... 46 2.8 41 25 4 18 16 80-84 years..... 47 32 15 42 2.8 14 4 3 1 17 85-89 years..... 39 21 18 33 16 6 4 90-94 years..... 23 19 21 18 2 1 4 3 1 95-99 years..... 3 8 5 3 4 2 1 100 years & over. 2 1 2 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female sexes Male Female sexes Male sexes Male sexes Male Female sexes Myeloid leukemia (C92) All ages.... 10,222 5,706 4,516 9,094 5,097 3,997 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2.4 20-24 vears..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1,119 1,001 65-69 years..... 70-74 years..... 1,330 1,202 75-79 years..... 1,528 1,386 80-84 years..... 1,590 1,490 85-89 years..... 90-94 years..... 1.0 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Acute myeloid leukemia (C92.0) All ages.... 8,214 4,553 3,661 7,335 4,099 3,236 Under 1 year.... 1.0 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 vears..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 1,106 75-79 years..... 1,260 1,145 80-84 years..... 1,234 1,157 85-89 years..... 90-94 years..... 1.0 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic myeloid leukemia (C92.1) All ages.... 1,077 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 5.8 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis					1.								ian or		
Tenth Revision, Second		All rac	es	D	White		D - 1-1-	Black			cican Indi	an 1/		ıc Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	omalo	Both sexes	Mala	Female
						remare	BCACB	naic				Ciliaic	BCACB	Haic	remare
Myeloid sarcoma (C92.	3)														
All ages	15	6	9	15	6	9	-	_	-	_	_	-	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
75-79 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
80-84 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
85-89 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	-	-	_	-	-	_	_	_	-	-	-	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acute promyelocytic leukemia (C92.4) All ages.... 121 12 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 1 1 1 20-24 years..... 10 6 25-29 years..... 3 30-34 years..... 10 35-39 years..... 6 40-44 years..... 11 45-49 years..... 14 a 5 12 3 50-54 years..... 15 1 0 12 55-59 years..... 11 11 4 60-64 years..... 9 2 8 65-69 vears..... 6 70-74 vears..... 6 75-79 years..... 12 12 80-84 years..... 14 8 14 85-89 years..... 8 4 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acute myelomonocytic leukemia (C92.5) All ages..... 72 149 67 10 Under 1 year.... 1 1 1 1 1-4 years..... 3 1 3 1 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 2 1 2 50-54 years..... 4 3 1 55-59 years..... 4 4 60-64 years..... 4 3 65-69 vears..... 21 70-74 vears..... 19 12 18 75-79 years..... 25 18 22 15 80-84 years..... 38 19 19 36 17 19 18 85-89 years..... 10 8 18 10 8 9 90-94 years..... 2 a 2 95-99 years..... 3 3 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other myeloid leukemia (C92.7) All ages..... 246 148 365 138 20 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 2 1 3 2 6 50-54 years..... 2 1 55-59 years..... 26 18 8 24 18 6 60-64 years..... 25 15 10 24 14 10 65-69 vears..... 41 30 11 35 25 10 70-74 vears..... 56 38 18 37 17 75-79 years..... 70 51 19 65 47 18 3 80-84 years..... 80 50 30 74 47 27 2 2 59 27 2 85-89 years..... 63 30 33 32 1 17 90-94 years..... 6 11 17 11 6 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Myeloid leukemia, unspecified (C92.9) All ages.... 138 85 190 115 75 21 Under 1 year.... 1 1 1 1 1-4 years..... 4 2 2 4 2 2 5-9 years..... 10-14 years..... 1 15-19 years..... 1 20-24 years..... 3 2 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 4 2 50-54 years..... 18 14 15 12 55-59 years..... 18 12 15 9 3 6 60-64 years..... 16 13 3 14 12 65-69 vears..... 20 12 15 8 70-74 vears..... 23 17 6 17 75-79 years..... 25 14 19 1.0 9 3 11 2 80-84 years..... 37 24 13 35 23 12 1 1 21 2 85-89 years..... 13 8 17 11 6 1 7 90-94 years..... 2 5 2 5 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Monocytic leukemia (C93) All ages..... 119 72 112 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 1 5 50-54 years..... 1 55-59 years..... 6 5 1 5 60-64 years..... 6 2 65-69 vears..... 10 5 5 9 70-74 vears..... 12 3 75-79 years..... 26 16 1.0 80-84 years..... 20 14 6 19 13 6 18 85-89 years..... 8 10 16 9 90-94 years..... 6 2 2 4 6 95-99 years..... 2 1 1 2 100 years & over. 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acute monocytic leukemia (C93.0) All ages..... 79 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 2 50-54 years..... 2 1 1 55-59 years..... 6 5 1 5 60-64 years..... 4 2 2 65-69 vears..... 3 6 70-74 vears..... 11 8 3 75-79 years..... 19 19 13 6 80-84 years..... 13 10 3 12 9 3 12 85-89 years..... 5 10 90-94 years..... 2 2 4 4 2 95-99 years..... 1 1 1 100 years & over. 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Chronic monocytic leukemia (C93.1) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 6 3 80-84 years..... 2 2 85-89 years..... 90-94 years..... 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Monocytic leukemia, unspecified (C93.9) All ages..... 16 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 5 85-89 years..... 4 3 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other leukemias of specified cell type (C94) All ages.... 32 30 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 3 1 2 60-64 years..... 3 65-69 vears..... 4 3 70-74 vears..... 9 5 75-79 years..... 80-84 years..... 6 2 85-89 years..... 8 90-94 years..... 1 1 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acute erythremia and erythroleukemia (C94.0) All ages..... 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 1 60-64 years..... 2 1 1 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 3 3 80-84 years..... 2 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	seases,	All rac	100		White			Black		λma	rican Indi		ian or	ic Tela	nder 2/
Edition, 2004)	Both	AII Iac	.cs	Both	WILLCE		Both	Black		Both	.ican indi	an i/	Both		nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Chronic erythremia (C	294.1)														
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Inder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	_	-	_	-	-	-	-	-	_	-	-	_	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.0-14 years	_	-	-	_	_	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
0-24 years	_	-	-	_	_	-	-	-	-	-	-	-	-	-	-
25-29 years	_	-	-	_	_	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	_	-	-	_	_	-	-	-	-	-	-	-	-	-	-
0-44 years	_	-	-	_	_	-	-	-	-	-	-	-	-	-	-
5-49 years	_	-	-	_	_	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
0-74 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
5-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-84 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acute megakaryoblastic leukemia (C94.2) All ages.... 9 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Mast cell leukemia (C94.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International															
Classification of Dis													sian or		
Tenth Revision, Secon		All races			White		Black				ican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both		Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale ————	sexes	Male	Female
Acute myelofibrosis (C94.5)														
All ages	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
85-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified leukemias (C94.7) All ages..... 13 18 11 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 2 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 3 3 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Leukemia of unspecified cell type (C95) All ages.... 5,155 2,838 2,317 4,648 2,573 2,075 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 2.2 15-19 years..... 2.7 2.0 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 1 0 1.0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Acute leukemia of unspecified cell type (C95.0) All ages.... 2,100 1,142 1,934 1,057 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.6 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 2.7 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Chronic leukemia of unspecified cell type (C95.1) All ages.... 174 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 2 50-54 years..... 4 2 2 55-59 years..... 5 2 3 2 5 60-64 years..... 12 8 11 65-69 vears..... 18 13 5 10 3 70-74 vears..... 22 10 12 18 9 9 75-79 years..... 34 18 16 32 17 80-84 years..... 36 22 14 34 21 13 1 32 29 2 85-89 years..... 14 18 11 18 2 90-94 years..... 18 12 12 6 1.8 6 95-99 years..... 7 3 7 4 3 4 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Subacute leukemia of unspecified cell type (C95.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other leukemia of unspecified cell type (C95.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Leukemia, unspecified (C95.9) All ages.... 2,852 1,588 1,264 2,530 1,420 1,110 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2.0 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 2.7 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Bot.h Both and age Female Male Female Male Female Male Female sexes Male sexes Male Female sexes sexes sexes Other and unspecified malignant neoplasms of lymphoid, hematopoietic, and related tissue (C96) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 2 50-54 years..... 55-59 years..... 6 3 3 5 60-64 years..... 3 3 1 65-69 vears..... 4 3 70-74 vears..... 5 75-79 years..... 13 13 6 80-84 years..... 10 1.0 85-89 years..... 13 9 12 7 90-94 years..... 3 3 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Malignant histiocytosis (C96.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Malignant mast cell tumor (C96.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female Male Female sexes Male sexes Male sexes sexes sexes Other specified malignant neoplasms of lymphoid, hematopoietic, and related tissue (C96.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 5 3 85-89 years..... 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malignant neoplasms of lymphoid, hematopoietic, and related tissue, unspecified (C96.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 2 1 1 60-64 years..... 3 3 65-69 vears..... 3 3 70-74 vears..... 5 75-79 years..... 10 80-84 years..... 4 4 85-89 years..... 5 90-94 years..... 3 3 4 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based															
on the International Classification of Dis-	0200											λα	ian or		
Tenth Revision, Second		All rac	es		White			Black		Amer	ican Ind			ic Isla	nder 2/
Edition, 2004)	Both	1111 100		Both	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		Both	224011		Both	10011 1110		Both	10 1010	27
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Malignant neoplasms o	f independer	nt (primar	y) multiple	sites (C97)										
All ages	4,380	2,494	1,886	3,855	2,187	1,668	454	266	188	15	5	10	56	36	20
Under 1 year		. –	. –	-	, -	-	_	_	-	_	-	_	_	_	_
1-4 years	1	-	1	1	_	1	_	-	-	_	-	-	-	_	-
5-9 years	7	4	3	7	4	3	_	-	-	_	-	-	-	_	-
10-14 years	4	2	2	4	2	2	_	-	-	_	-	-	-	_	-
L5-19 years	9	5	4	8	4	4	1	1	-	-	-	-	-	-	-
20-24 years	9	4	5	7	3	4	1	-	1	_	-	-	1	1	-
25-29 years	10	5	5	7	4	3	3	1	2	-	-	-	-	-	-
30-34 years	13	9	4	12	9	3	1	-	1	-	-	-	-	-	-
35-39 years	15	10	5	14	9	5	-	_	-	_	_	-	1	1	-
10-44 years	40	18	22	37	15	22	3	3	-	_	_	-	-	-	-
15-49 years	107	53	54	88	44	44	14	7	7	_	-	-	5	2	3
50-54 years	135	65	70	116	51	65	16	12	4	1	1	-	2	1	1
55-59 years	251	116	135	210	95	115	37	19	18	1	-	1	3	2	1
0-64 years	330	181	149	271	145	126	52	33	19	_	-	-	7	3	4
55-69 years	424	229	195	363	195	168	50	29	21	4	1	3	7	4	3
70-74 years	559	338	221	496	297	199	53	34	19	1	-	1	9	7	2
5-79 years	677	412	265	590	360	230	81	48	33	1	_	1	5	4	1
0-84 years	824	479	345	741	428	313	74	47	27	4	1	3	5	3	2
35-89 years	590	359	231	541	332	209	38	18	20	3	2	1	8	7	1
90-94 years	292	162	130	264	149	115	25	12	13	_	_	-	3	1	2
95-99 years	71	36	35	68	35	33	3	1	2	_	_	-	_	-	-
100 years & over.	12	7	5	10	6	4	2	1	1	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis												ħ.a.	ian or		
Tenth Revision, Secon		All rac	0.7		White			Black		7mos	cican Indi			ia Tala	nder 2/
Edition, 2004)	Both	AII Iac	es	Both	WIIILE		Both	Black		Both	ican inui	.all 1/	Both	IC ISIA	nuer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	omalo	sexes	Mala	Female
		Maie	remaie		Maie	remare	SCACS	Maic	remare		Maie r	CIIIAIE	beves	Maie	remare
Carcinoma in situ of	amal garritu	ogophogu	a and atoms	ah (D00)											
carcinoma in sicu or	Orar cavity,	esopnagu	s, and scome	icii (Doo)											
All ages	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
l5-19 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	_
40-44 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	_
45-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
70-74 years	_	-	-	_	-	-	-	-	-	_	-	-	-	-	-
75-79 years	2	2	-	2	2	-	-	-	-	_	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
35-89 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
90-94 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
95-99 years	_	-	-	-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Lip, oral cavity, and pharynx (D00.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dise													sian or		
Tenth Revision, Second		All rac	es		White			Black			rican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Esophagus (D00.1)															
All ages	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Stomach (D00.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Carcinoma in situ of other and unspecified digestive organs (D01) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis												As	ian or		
Tenth Revision, Second		All rac	es		White			Black		Amer	ican Ind			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both		,	Both		,
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Liver, gallbladder, a	nd bile duct	s (D01.5)													
All ages	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	_	-	-	-	_	-	-	-	-	-	-
20-24 years	-	-	-	-	_	-	-	-	_	-	-	-	-	-	-
25-29 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	_	-	-	-	_	-	-	-	-	-	-
40-44 years	-	-	-	-	_	-	-	-	_	-	-	-	-	-	-
15-49 years	-	-	-	-	_	-	-	-	_	-	-	-	-	-	-
50-54 years	-	-	-	-	_	-	-	-	_	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
70-74 years	_	-	-	-	-	-	-	-	_	_	_	_	-	_	-
75-79 years	-	-	-	-	-	_	-	-	_	_	-	-	-	-	-
0-84 years	1	-	1	1	-	1	-	-	_	_	_	_	-	_	-
5-89 years	-	-	-	-	-	_	-	-	_	_	-	-	-	-	-
0-94 years	-	-	-	-	-	_	-	-	_	_	-	-	-	-	-
5-99 years	1	1	-	1	1	_	-	-	_	_	-	-	-	-	-
100 years & over.	_	-	-	-	-	_	-	-	_	_	_	_	-	_	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Carcinoma in situ of middle ear and respiratory system (D02) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Discount Party Party Canal	eases,	711			White			Dll-		7	ican Indi		ian or	::- T-1-	d 2 /
Tenth Revision, Second		All rac	es	D . 1-1-	wnite		D - 1-1-	Black			rican Indi	an 1/		ic isia	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
Larynx (D02.0)															
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
.0-14 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
5-59 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	_	-	_	-	_	-	-	-	-	-
5-79 years	_	_	-	_	_	_	-	-	-	-	-	-	-	-	-
0-84 years	_	_	-	_	_	_	-	-	-	-	-	-	-	-	-
5-89 years	_	_	-	_	_	_	-	-	-	-	-	-	-	-	-
0-94 years	_	_	-	_	_	_	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	_	-	_	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Bronchus and lung (D02.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Melanoma in situ (D03) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Melanoma in situ of other sites (D03.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis Tenth Revision, Secon		All rac	10.5		White			Black		λmox	ican Indi		ian or	ic Islan	ndor 2/
Edition, 2004)	Both	AII Iac	es	Both	WIIICE		Both	Black		Both	Ican Indi	all 1/	Both	IC ISIAI	nuer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Carcinoma in situ of	breast (D05)														
All ages	2	-	2	1	-	1	_	_	-	_	_	-	1	_	1
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
15-19 years	-	-	-	_	-	-	-	_	-	-	-	_	-	-	-
20-24 years	-	-	-	_	-	-	-	_	-	-	-	_	-	-	-
25-29 years	1	-	1	-	-	-	-	-	-	-	-	-	1	-	1
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	_	-	-	-	_	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	-	_	-	-	-	_	-	-	-
75-79 years	-	-	-	_	-	-	-	_	-	-	-	_	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
00-94 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Intraductal carcinoma in situ (D05.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Carcinoma in situ of breast, unspecified (D05.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Carcinoma in situ of other and unspecified genital organs (D07) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Prostate (D07.5) All ages.... . . . Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... . . . 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... . . . 85-89 years..... . . . 90-94 years..... 95-99 years..... 100 years & over. . . . Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Carcinoma in situ of other and unspecified sites (D09) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Bladder (D09.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Carcinoma in situ, unspecified (D09.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign neoplasm of major salivary glands (D11) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Parotid gland (D11.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Major salivary gland, unspecified (D11.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign neoplasm of colon, rectum, anus, and anal canal (D12) All ages.... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 65-69 vears..... 70-74 vears..... 3 75-79 years..... 80-84 years..... 3 2 85-89 years..... 90-94 years..... 2 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cecum (D12.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Colon, unspecified (D12.6) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 3 75-79 years..... 3 80-84 years..... 3 85-89 years..... 90-94 years..... 2 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Rectum (D12.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign neoplasm of other and ill-defined parts of digestive system (D13) All ages.... 32 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 2 50-54 years..... 1 55-59 years..... 2 1 1 60-64 years..... 6 2 65-69 vears..... 2 70-74 vears..... 6 75-79 years..... 3 80-84 years..... 8 6 1 1 10 85-89 years..... 12 2 10 12 90-94 years..... 5 1 4 5 4 95-99 years..... 3 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Esophagus (D13.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Stomach (D13.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) Both		All races White Blac								Asian or Black American Indian 1/ Pacific Is							
		All rac	es	D - 1-1-	White		D	Black			rican Indi	an 1/		ic Isla	nder 2/		
Edition, 2004) and age	Botn sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female		
Duodenum (D13.2)																	
All ages	6	3	3	5	2	3	_	_	_	_	_	_	1	1	_		
Under 1 year	-	-	-	_	-	-	-	-	-	_	_	-	_	-	-		
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-		
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-		
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
35-39 years	1	1	-	-	-	-	-	-	-	-	-	-	1	1	-		
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
'5-79 years	-	-	-	_	-	-	-	-	-	_	_	-	_	-	-		
0-84 years	1	1	-	1	1	-	-	-	-	_	_	-	_	-	-		
85-89 years	-	-	-	_	-	-	-	-	-	_	_	-	_	-	-		
90-94 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-		
95-99 years	-	-	-	_	-	-	-	-	-	_	_	-	_	-	-		
100 years & over.	_	-	_	_	_	_	_	_	_	_	_	_	_	_	_		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Liver (D13.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Extrahepatic bile ducts (D13.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pancreas (D13.6) All ages.... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 2 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Endocrine pancreas (D13.7) All ages..... 10 14 Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 50-54 years..... 1 1 55-59 years..... 2 1 1 60-64 years..... 4 2 2 65-69 vears..... 2 70-74 vears..... 75-79 years..... 3 80-84 years..... 4 1 85-89 years..... 8 90-94 years..... 1 1 1 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Ill-defined sites within the digestive system (D13.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign neoplasm of middle ear and respiratory system (D14) All ages.... 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Middle ear, nasal cavity, and accessory sinuses (D14.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Larynx (D14.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) Bot										Asian or American Indian 1/ Pacific Islander 2/						
		All rac	es		White			Black			rican Indi	an 1/		ic Isla	nder 2/	
		Male	Hamala.	Both	Male	Esmals.	Both	Male	Hamala	Both	Mala E		Both	M-l-	Female	
and age 	sexes	мате	Female	sexes	мате	Female	sexes		Female	sexes	Male F	emaie ————	sexes		remaie	
Bronchus and lung (D1	4.3)															
All ages	9	3	6	6	3	3	3	_	3	-	-	-	-	_	-	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-	
l5-19 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
65-69 years	2	1	1	1	1	-	1	-	1	-	-	-	-	-	-	
70-74 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-84 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-	
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Age not stated	_	_	-	_	-	-	-	_	_	_	-	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Classification of Dis Tenth Revision, Secon		All rac	100		White		Black			Asian or American Indian 1/ Pacific Islander 2						
Edition, 2004)	Both	AII IAC	.05	Both	WIIICC		Both	DIACK		Both	ican ind.	ian i/	Both	.10 1514	maci 2/	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female	
Respiratory system, u	nspecified	(D14.4)														
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_	
Under 1 year	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
1-4 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
5-9 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
10-14 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
15-19 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
20-24 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_	
25-29 years	-	_	-	-	-	_	-	_	-	_	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
40-44 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
80-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Age not stated	-	-	_	_	_	_	-	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Benign neoplasm of other and unspecified intrathoracic organs (D15) All ages.... 32 Under 1 year.... 5 1 4 3 3 2 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 6 45-49 years..... 2 50-54 years..... 2 2 55-59 years..... 3 3 60-64 years..... 4 2 65-69 vears..... 3 70-74 vears..... 3 3 75-79 years..... 80-84 years..... 5 1 85-89 years..... 10 1 90-94 years..... 6 6 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International															
Classification of Dis		211			White			D11-				ian or			
Tenth Revision, Secon Edition, 2004)	a Both	All rac	ces	Both	wnite		Both	Black		Amer Both	ican Indi	an 1/	Pacii Both	ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'omalo	sexes	Mala	Female
and age	SCACS	mare	remare	SCACS	mare	remare	Seves	maie	remare	SCACS	Male r		SCACS		
Thymus (D15.0)															
All ages	24	11	13	13	8	5	9	2	7	-	-	-	2	1	1
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	3	1	2	-	-	-	3	1	2	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	2	1	1	-	-	-	2	1	1	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	2	-	2	1	-	1	-	-	-	-	-	-	1	-	1
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
65-69 years	3	1	2	2	1	1	1	-	1	-	-	-	-	-	-
70-74 years	3	3	-	2	2	-	-	-	-	-	-	-	1	1	-
75-79 years	3	3	-	3	3	-	-	-	-	-	-	-	-	-	-
80-84 years	1	-	1	-	-	_	1	_	1	-	-	-	_	-	-
85-89 years	3	1	2	2	1	1	1	_	1	-	-	-	_	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	_	-	-	-	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Heart (D15.1) All ages.... 10 Under 1 year.... 5 1 3 3 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 3 1 65-69 vears..... 4 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 2 85-89 years..... 5 90-94 years..... 6 6 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second All races Edition, 2004) Both							Asian or American Indian 1/ Pacific Islander 2/								
		All rac	es		White		_	Black			ican Indi	an 1/		ic Isla	nder 2/
			_ ,	Both			Both			Both		-	Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale ————	sexes	Male	Female
Other specified intra	thoracic org	gans (D15.	7)												
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
l-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-74 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
0-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign neoplasm of bone and articular cartilage (D16) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Bones of skull and face (D16.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lower jaw bone (D16.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Vertebral column (D16.6) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pelvic bones, sacrum, and coccyx (D16.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Benign lipomatous neoplasm (D17) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 2 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign lipomatous neoplasm of intrathoracic organs (D17.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign lipomatous neoplasm of other sites (D17.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign lipomatous neoplasm, unspecified (D17.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hemangioma and lymphangioma, any site (D18) All ages.... 39 36 10 Under 1 year.... 4 2 2 2 1 1 2 1 1-4 years..... 3 3 2 1 1 5-9 years..... 1 10-14 years..... 1 1 1 15-19 years..... 1 20-24 years..... 1 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 4 1 3 50-54 years..... 2 1 55-59 years..... 2 1 1 60-64 years..... 1 6 65-69 vears..... 3 3 70-74 vears..... 3 75-79 years..... 9 5 80-84 years..... 6 4 2 2 2 85-89 years..... 9 90-94 years..... 4 4 95-99 years..... 1 1 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) Both		233			referènce		Asian or American Indian 1/ Pacific Islander 2/								
		All rac	es	Both	White		Both	Black		Amer Both	rican Ind	ian I/	Pacii Both	ic Isla	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Hemangioma, any site ((D18 0)														
All ages	49	20	29	43	16	27	5	3	2	-	-	-	1	1	-
Under 1 year	2	1	1	1	-	1	1	1	_	-	_	_	-	-	-
1-4 years	1	-	1	_	-	-	1	-	1	-	_	_	-	-	-
5-9 years	-	_	-	_	-	-	-	-	-	-	_	_	-	-	-
10-14 years	1	_	1	1	-	1	-	_	-	_	-	_	-	-	-
15-19 years	_	_	-	-	_	_	-	_	-	_	-	_	-	-	-
20-24 years	_	_	_	-	_	_	-	_	-	_	-	-	-	-	-
25-29 years	6 3	2	4	6 3	2	4	-	_	-	_	-	-	-	-	-
30-34 years	3	2	1	_	2	1	-	_	_	-	_	-	_	-	-
35-39 years	-	-	-	- 1	-	-	-	_	-	_	-	-	-	-	-
40-44 years	1	3	1	3	-	1	- 1	1	_	_	_	-	-	_	-
45-49 years	4	2	1 1	3	2 2	1	_	1	-	_	_	-	-	_	-
50-54 years	3	2	1	1	_	1	_	_	_	_	_	_	-	_	-
55-59 years 60-64 years	Τ.	- 1	4	4	1	3	- 1	_	1		_	_	_	_	_
65-69 years	2	2	4	1	1	<i>3</i>	1	_	_	_	_	_	1	1	_
70-74 years	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
70-74 years 75-79 years	2		3	6	3	± 2	_	_	_	_	_	_	_		_
80-84 years	3	2	1	2	3 1	3 1	- 1	1	_	_	_	_	_	_	_
85-89 years	2	1	4	5	1	4	_	_	_	_	_	_	_	_	_
90-94 years	4	_	4	4	_	4	_	_	_	_	_	_	_	_	_
95-99 years	-	_	-	-	_	-	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
Age not stated	_	_	_	_	_										

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lymphangioma, any site (D18.1) All ages..... 10 Under 1 year.... 2 1 1 1 1 1-4 years..... 2 2 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 1 85-89 years..... 4 90-94 years..... 95-99 years..... 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign neoplasm of mesothelial tissue (D19) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mesothelial tissue of other sites (D19.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign neoplasm of soft tissue of retroperitoneum and peritoneum (D20) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

											As	ian or			
Tenth Revision, Second	i.	All rac	ces		White			Black		Amer	ican Indi		Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Peritoneum (D20.1)															
All ages	2	2	_	2	2	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-14 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	_	-	-	_	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	_	-	-	_	-	_	-	-	-	-	-	-	-	-	-
40-44 years	_	-	-	_	-	_	-	-	-	-	-	-	-	-	-
45-49 years	_	-	-	_	-	_	-	-	-	-	-	-	-	-	-
50-54 years	-	-		-	-	-	-	_	-	-	_	-	-	-	-
55-59 years	-	-		-	-	-	-	_	-	-	_	-	-	-	-
60-64 years	-	-		-	-	-	-	_	-	-	_	-	-	-	-
65-69 years	-	-		-	-	-	-	_	-	-	_	-	-	-	-
70-74 years	_	-	-	_	-	_	-	-	-	_	-	-	-	-	-
75-79 years	1	1	-	1	1	_	-	-	-	_	-	-	-	-	-
30-84 years	_	_	-	_	-	_	-	_	_	-	-	-	-	-	-
35-89 years	1	1	-	1	1	_	-	_	_	-	-	-	-	-	-
90-94 years	_	_	-	_	-	_	-	_	_	-	-	-	-	-	-
95-99 years	_	_	-	_	-	_	-	_	_	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other benign neoplasms of connective and other soft tissue (D21) All ages.... 10 Under 1 year.... 2 2 2 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 2 50-54 years..... 1 1 55-59 years..... 1 1 2 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 3 90-94 years..... 3 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Connective and other soft tissue of head, face, and neck (D21.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Connective and other soft tissue of thorax (D21.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Connective and other soft tissue of abdomen (D21.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Connective and other soft tissue, unspecified (D21.9) All ages.... 15 Under 1 year.... 2 2 2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 1 1 55-59 years..... 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 3 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other benign neoplasms of skin (D23) All ages..... 10 9 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 2 1 1 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 2 3 85-89 years..... 90-94 years..... 95-99 years..... 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dise	eases,											As	ian or		
Tenth Revision, Second	i	All rac	es		White			Black		Ame	ican Indi	an 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Skin of scalp and neck	(D23.4)														
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	_	-	_	_	_	-	-	-	_	_	-	_	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
l5-19 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	_	-	_	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	_	-	_	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
'5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Skin of trunk (D23.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Skin of lower limb, including hip (D23.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Skin, unspecified (D23.9) All ages..... 20 19 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 1 90-94 years..... 95-99 years..... 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Leiomyoma of uterus (D25) All ages..... 30 30 10 10 20 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 3 45-49 years..... 1.0 10 50-54 years..... 4 4 2 55-59 years..... 2 2 60-64 years..... 1 1 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Leiomyoma of uterus, unspecified (D25.9) All ages.... 30 10 10 20 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 3 3 45-49 years..... 1.0 10 50-54 years..... 4 4 2 55-59 years..... 2 2 60-64 years..... 1 1 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other benign neoplasms of uterus (D26) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... . . . 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise		- 1 1						-1 1					ian or		3 04
Tenth Revision, Second		All rac	ces	D - 1-1-	White		D - +1-	Black			rican In	dian 1/		ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Corpus uteri (D26.1)															
All ages	1		1	-		-	_		-	_		-	1		1
Under 1 year	-		-	-		-	-		-	-		-	-		-
1-4 years	-		-	-		-	-		-	-		-	-		-
5-9 years	-		-	-		-	-		-	-		-	_		-
10-14 years	-		-	-		-	-		-	-		-	_		-
.5-19 years	-		-	-		-	-		-	-		-	-		-
0-24 years	-		-	-		-	-		-	-		-	-		-
25-29 years	-		-	-		-	-		-	-		-	-		-
30-34 years	-		-	-		-	-		-	-		-	-		-
35-39 years	-		-	-		-	-		-	-		-	-		-
10-44 years	-		-	-		-	-		-	-		-	-		-
45-49 years	-		-	-		-	-		-	-		-	-		-
50-54 years	-			-		-	-		-	-		-	-		-
55-59 years	-		-	-		-	-		-	-		-	-		-
60-64 years	-			-		-	-		-	-		-	-		-
65-69 years	-			-		-	-		-	-		-	-		-
70-74 years	-		-	-		-	-		_	_		-	_		-
75-79 years	1		1	-		-	-		_	_		-	1		1
30-84 years	-		-	-		-	-		_	_		-	_		-
35-89 years	-		-	-		-	-		_	_		-	_		-
90-94 years	-		-	-		-	-		_	_		-	_		-
95-99 years	-		_	_		-	_		_	-		-	_		-
100 years & over.	_		_	_		_	_		_	_		_	_		_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign neoplasm of ovary (D27) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Bot.h Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign neoplasm of other and unspecified female genital organs (D28) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... . . . 20-24 years..... . . . 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Uterine tubes and ligaments (D28.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... . . . 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign neoplasm of male genital organs (D29) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... . . . 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 1 80-84 years..... . . . 85-89 years..... . . . 90-94 years..... 95-99 years..... 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Testis (D29.2) All ages.... . . . Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... . . . 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 1 80-84 years..... . . . 85-89 years..... . . . 90-94 years..... 95-99 years..... 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis													ian or		
Tenth Revision, Secon		All rac	es	_	White		_	Black			rican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Benign neoplasm of ur	inary organs	s (D30)													
All ages	2	2	_	2	2	_	_	_	-	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
45-49 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	_	-	-	-	_	-	-	-	-	-	_	-	-	-	-
85-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
90-94 years	_	-	-	-	_	-	-	-	-	-	_	-	-	-	-
95-99 years	_	-	-	-	_	-	-	-	-	-	_	-	-	-	-
100 years & over.	_	-	-	-	_	-	-	-	-	-	_	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dise	eases,											As	sian or		
Tenth Revision, Second		All rac	es		White			Black		Amen	cican India			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Kidney (D30.0)															
All ages	2	2	_	2	2	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
l0-14 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
40-44 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
15-49 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
50-64 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
55-69 years	-	-		-	-	-	-	_	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
'5-79 years	_	_	-	_	_	_	_	-	-	_	-	-	_	_	-
0-84 years	-	-	-	_	-	-	-	_	-	_	-	-	-	_	-
35-89 years	1	1	-	1	1	_	_	-	-	_	-	-	_	_	-
0-94 years	-	-	-	_	-	-	-	_	-	_	-	-	-	_	-
95-99 years	-	-	-	_	-	-	-	_	-	_	-	-	-	_	-
100 years & over.	_	_	-	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign neoplasm of eye and adnexa (D31) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cornea (D31.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Ciliary body (D31.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Benign neoplasm of meninges (D32) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 years..... 2.8 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 5.8 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Cerebral meninges (D32.0) All ages..... 207 132 176 113 25 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 2 5 9 50-54 years..... 3 R 6 55-59 years..... 13 9 10 4 60-64 years..... 7 5 1 65-69 vears..... 13 13 8 5 70-74 vears..... 18 75-79 years..... 2.8 10 18 23 15 80-84 years..... 35 12 23 33 12 21 2 39 85-89 years..... 49 18 31 13 26 3 15 90-94 years..... 2 13 15 13 2 95-99 years..... 5 5 5 5 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dise	ases,											As	sian or		
Tenth Revision, Second		All rac	ces		White			Black		Ame	rican Indi	an 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Spinal meninges (D32.1)														
All ages	5	1	4	5	1	4	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	_
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-
75-79 years	_	_	-	_	_	-	-	-	-	_	-	-	_	_	-
30-84 years	3	-	3	3	-	3	-	_	-	_	-	-	-	_	-
35-89 years	_	_	-	_	_	-	-	-	-	_	-	-	_	_	-
90-94 years	1	_	1	1	_	1	-	-	-	_	-	-	_	_	-
95-99 years	_	_	-	_	_	-	-	-	-	_	-	-	_	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Meninges, unspecified (D32.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.7 80-84 years..... 85-89 years..... 2.5 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Benign neoplasm of brain and other parts of central nervous system (D33) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 5 7 50-54 years..... 3 55-59 years..... 11 8 3 10 60-64 years..... 6 2 5 1 65-69 vears..... 7 3 70-74 vears..... 10 5 10 75-79 years..... 11 8 80-84 years..... 15 8 15 12 2 85-89 years..... 10 11 9 2 90-94 years..... 8 2 1 6 6 95-99 years..... 3 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis													sian or		
Tenth Revision, Secon		All rac	es		White			Black			rican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both	W. 1 .		Both	M. 1 -	m1	Both	34.1.		Both	W-1- =	1 .	Both	24 - 7 -	D 1 .
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes		Female
Brain, supratentorial	(D33.0)														
All ages	10	5	5	10	5	5	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	_
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	_
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
50-64 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
55-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-84 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	_	_	-	_	_	_	_	_	_	_	_	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Brain, infratentorial (D33.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Brain, unspecified (D33.2) All ages.... 30 32 30 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 8 5 3 3 50-54 years..... 1 55-59 years..... 3 2 1 60-64 years..... 1 1 65-69 vears..... 4 70-74 vears..... 5 3 75-79 years..... 3 80-84 years..... 10 6 10 7 85-89 years..... 5 90-94 years..... 1 3 2 4 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cranial nerves (D33.3) All ages.... 27 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 2 50-54 years..... 1 1 55-59 years..... 2 2 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 4 75-79 years..... 2 80-84 years..... 3 85-89 years..... 4 90-94 years..... 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Spinal cord (D33.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Central nervous system, unspecified (D33.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dise	eases,											As	sian or		
Tenth Revision, Second		All rac	es		White			Black		Amer	ican Indi			fic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Benign neoplasm of oth	ner and unsp	pecified e	ndocrine gla	ands (D35)											
All ages	126	62	64	92	45	47	34	17	17	_	_	_	_	_	_
Under 1 year	1	-	1	1	-	1	_	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
15-19 years	3	2	1	2	2	-	1	-	1	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-
30-34 years	4	1	3	1	-	1	3	1	2	-	-	-	-	-	-
35-39 years	4	3	1	4	3	1	-	-	-	-	-	-	-	-	-
40-44 years	9	5	4	6	3	3	3	2	1	-	-	-	-	-	-
45-49 years	9	5	4	4	3	1	5	2	3	-	-	-	-	-	-
50-54 years	9	5	4	7	5	2	2	-	2	-	-	-	-	-	-
55-59 years	7	4	3	5	3	2	2	1	1	-	-	-	-	-	-
60-64 years	8	4	4	7	3	4	1	1	-	-	-	-	-	-	-
65-69 years	6	3	3	6	3	3	_	-	-	-	-	-	-	-	-
70-74 years	10	5	5	8	4	4	2	1	1	-	-	-	-	_	-
75-79 years	12	4	8	8	3	5	4	1	3	-	-	-	-	_	-
30-84 years	16	7	9	10	4	6	6	3	3	_	-	_	_	_	-
85-89 years	18	9	9	16	7	9	2	2	_	_	-	_	_	_	-
90-94 years	6	4	2	4	2	2	2	2	_	_	_	_	_	_	_
95-99 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_			_	_				_		_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Adrenal gland (D35.0) All ages.... Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 45-49 years..... 3 1 2 5 50-54 years..... 3 2 1 55-59 years..... 3 2 1 60-64 years..... 3 3 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis Tenth Revision, Secon		All rac	es:		White			Black		Amei	ican Indi		ian or Pacif	ic Isla	ınder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Parathyroid gland (D3	5.1)														
All ages	5	_	5	3	_	3	2	_	2	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	2	-	2	1	-	1	1	-	1	-	-	-	-	-	-
30-84 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
85-89 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pituitary gland (D35.2) All ages.... 78 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 6 4 2 50-54 years..... 4 2 2 55-59 years..... 3 2 4 1 60-64 years..... 5 1 1 65-69 vears..... 3 70-74 vears..... 9 5 75-79 years..... 9 5 80-84 years..... 11 6 2 14 2 85-89 years..... 9 5 12 2 90-94 years..... 5 1 3 2 2 4 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Disea	rahitu.							Asian or American Indian 1/ Pacific Islander 2/							
Tenth Revision, Second		All rac	es	Dath	White		Dath	Black			erican Indian 1/				nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Malo	Female	Both sexes	Mala	Female
		maie	remare	SCACS	Maie	remare		Maie	remare		mare	remare		maie	remare
Pineal gland (D35.4)															
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	_	-	-	-	_	_	_	-	-	-	-	-
95-99 years	-	-	-	_	-	-	-	_	_	_	-	-	-	-	-
100 years & over.	-	_	-	_	-	_	_	-	_	_	_	_	-	-	-
Age not stated	_	_	_	_								_		_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Endocrine gland, unspecified (D35.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Benign neoplasm of other and unspecified sites (D36) All ages.... Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 2 45-49 years..... 2 50-54 years..... 4 3 1 3 55-59 years..... 4 2 2 60-64 years..... 4 2 2 65-69 vears..... 5 70-74 vears..... 5 3 75-79 years..... 3 80-84 years..... 6 2 1 85-89 years..... 9 8 90-94 years..... 1 3 4 1 3 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lymph nodes (D36.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Peripheral nerves and autonomic nervous system (D36.1) All ages.... 13 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 55-59 years..... 1 1 2 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 1 85-89 years..... 4 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis													ian or		
Tenth Revision, Secon		All races		_	White		Black				erican Indian 1/		Pacific Islander 2		
Edition, 2004)	Both			Both			Both			Both	Male Female		Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Other specified sites	(D36.7)														
All ages	8	1	7	8	1	7	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
50-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
70-74 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Benign neoplasm of unspecified site (D36.9) All ages..... 17 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 4 3 55-59 years..... 2 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 2 75-79 years..... 2 80-84 years..... 2 85-89 years..... 90-94 years..... 2 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Neoplasm of uncertain or unknown behavior of oral cavity and digestive organs (D37) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... Ω 50-54 years..... 2.3 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Lip, oral cavity, and pharynx (D37.0) All ages.... 15 9 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 2 50-54 years..... 55-59 years..... 2 2 60-64 years..... 4 65-69 vears..... 2 70-74 vears..... 4 75-79 years..... 80-84 years..... 4 2 85-89 years..... 90-94 years..... 4 3 4 95-99 years..... 2 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Stomach (D37.1) All ages.... 72 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 1 1 50-54 years..... 3 3 1 55-59 years..... 1 1 60-64 years..... 3 65-69 vears..... 4 70-74 vears..... 8 6 75-79 years..... 10 80-84 years..... 15 13 10 85-89 years..... 3 8 90-94 years..... 8 8 2 6 95-99 years..... 3 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Small intestine (D37.2) All ages..... 10 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 1 1 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 80-84 years..... 3 85-89 years..... 3 90-94 years..... 2 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Appendix (D37.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Colon (D37.4) All ages.... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 2 2 2 2 60-64 years..... 6 1 65-69 vears..... 10 3 70-74 vears..... 7 5 6 75-79 years..... 12 6 6 12 80-84 years..... 17 6 11 13 8 7 2 85-89 years..... 13 6 11 6 1 90-94 years..... 19 5 14 19 14 95-99 years..... 6 5 5 1 6 100 years & over. 2 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Rectum (D37.5) All ages.... 27 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 3 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 2 80-84 years..... 6 85-89 years..... 4 90-94 years..... 1 4 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based																	
on the International Classification of Dise	22505											7.0	ian or				
Tenth Revision, Second		All races			White				Black			American Indian 1/ Pacific Islander					
Edition, 2004)	Both	AII IAC	CB	Both			Both	DIACK		Both	ridian indian i		Both	.10 1514	inder 2/		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female		
Liver, gallbladder, ar	nd bile duct	s (D37.6)															
All ages	184	80	104	159	71	88	17	7	10	1	_	1	7	2	5		
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
25-29 years	-	-	-	-	_	-	-	-	-	-	-	_	-	-	-		
30-34 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-		
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-		
40-44 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
45-49 years	3	2	1	1	1	-	2	1	1	-	-	-	-	-	-		
50-54 years	11	8	3	8	6	2	2	1	1	-	-	-	1	1	-		
55-59 years	7	5	2	6	4	2	1	1	-	-	-	-	-	-	-		
60-64 years	12	5	7	11	5	6	-	-	-	-	-	_	1	-	1		
65-69 years	13	10	3	12	9	3	1	1	-	-	-	_	-	-	-		
70-74 years	23	14	9	19	13	6	1	_	1	-	-	-	3	1	2		
75-79 years	25	13	12	23	13	10	2	-	2	-	-	-	-	-	-		
80-84 years	32	8	24	25	6	19	5	2	3	-	-	-	2	-	2		
85-89 years	29	10	19	27	9	18	1	1	-	1	-	1	-	-	-		
90-94 years	19	1	18	19	1	18	-	-	-	-	-	-	-	-	-		
95-99 years	7	2	5	6	2	4	1	-	1	-	-	-	-	-	-		
100 years & over.	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-		
Age not stated	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other digestive organs (D37.7) All ages..... 132 194 118 21 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 5 2 3 5 2 50-54 years..... 13 6 11 55-59 years..... 7 4 3 5 60-64 years..... 9 5 6 65-69 vears..... 14 10 70-74 vears..... 25 14 20 75-79 years..... 2.2 8 14 7 80-84 years..... 40 15 25 36 14 22 42 28 3 85-89 years..... 46 14 32 14 3 22 90-94 years..... 18 2.0 17 2 1 4 3 95-99 years..... 9 8 9 8 1 1 100 years & over. 1 1 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Digestive organ, unspecified (D37.9) All ages..... 17 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 2 2 55-59 years..... 5 60-64 years..... 3 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 3 1 90-94 years..... 6 95-99 years..... 3 2 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Bot.h and age Female Male Female Male Female sexes Male Female sexes Male Female sexes Male sexes sexes Neoplasm of uncertain or unknown behavior of middle ear and respiratory and intrathoracic organs (D38) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 3 45-49 years..... 6 2 4 5 2 3 2 50-54 years..... 11 6 5 9 5 55-59 years..... 26 15 11 21 12 9 5 60-64 years..... 37 44 30 14 28 9 65-69 vears..... 53 28 25 44 23 70-74 years..... 68 34 34 61 30 31 75-79 years..... 90 54 36 78 48 30 7 80-84 years..... 108 52 56 98 46 52 2 121 72 105 61 85-89 years..... 49 44 12 4 1 17 90-94 years..... 67 50 63 47 3 1 16 3 95-99 years..... 17 10 2 2 4 13 14 4 1 100 years & over. 4 3 3 4 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Larynx (D38.0) All ages..... 15 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 4 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 4 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Trachea, bronchus, and lung (D38.1) All ages.... 301 501 240 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 5 1 4 4 1 3 2 50-54 years..... 1.0 6 R 3 5 55-59 years..... 21 12 9 17 10 4 2 60-64 years..... 40 28 12 34 26 8 65-69 years..... 49 25 40 20 2.0 70-74 years..... 65 32 33 58 28 30 75-79 years..... 85 50 35 74 45 29 80-84 years..... 100 48 52 91 43 48 6 3 2 107 93 85-89 years..... 43 64 38 55 10 4 1 90-94 years..... 65 17 48 62 16 46 2 2 1 95-99 years..... 13 2 10 2 2 11 2 8 1 100 years & over. 1 3 4 4 1 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis										Asian or						
Tenth Revision, Secon		All races		White			Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both	1	- 1	Both			Both		_ 1	Both	Male Female		Both		_ ,	
nd age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Pleura (D38.2)																
All ages	5	5	_	5	5	_	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-	
15-19 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	_	
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
35-39 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	_	
40-44 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-	
50-54 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
90-94 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-	
95-99 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Mediastinum (D38.3) All ages.... 31 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 2 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 8 85-89 years..... 5 90-94 years..... 1 1 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dise													sian or			
Tenth Revision, Second		All races		White				Black			American Indian 1/			Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	Pemale	sexes	Male	Female	
Thymus (D38.4)																
All ages	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-	
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-	
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-	
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-	
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other respiratory organs (D38.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Respiratory organ, unspecified (D38.6) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Neoplasm of uncertain or unknown behavior of female genital organs (D39) All ages.... 13 Under 1 year.... 2 2 2 2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 5 50-54 years..... 5 55-59 years..... 6 6 5 1 1 5 60-64 years..... 1 3 65-69 years..... 70-74 years..... 75-79 years..... 8 8 7 80-84 years..... 14 14 13 13 1 1 85-89 years..... 14 14 1.3 13 90-94 years..... 6 6 6 6 95-99 years..... 5 1 1 100 years & over. 3 3 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Uterus (D39.0) All ages.... 28 23 Under 1 year.... 2 2 2 2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 2 50-54 years..... 1 1 . . . 55-59 years..... 3 3 60-64 years..... 1 1 3 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 2 2 85-89 years..... 3 3 90-94 years..... 1 1 95-99 years..... 2 2 100 years & over. 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Ovary (D39.1) All ages.... 60 50 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 55-59 years..... 3 3 2 60-64 years..... 6 6 1 1 65-69 vears..... 70-74 years..... 6 75-79 years..... 80-84 years..... 11 11 10 10 1 85-89 years..... 9 9 8 8 90-94 years..... 4 4 95-99 years..... 3 3 1 1 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Placenta (D39.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... . . . 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other female genital organs (D39.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... . . . 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Female genital organ, unspecified (D39.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Neoplasm of uncertain or unknown behavior of male genital organs (D40) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... . . . 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 8 8 2 . . . 2 85-89 years..... 2 . . . 90-94 years..... 2 2 95-99 years..... 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Prostate (D40.0) All ages.... 16 16 . . . Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... . . . 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... . . . 75-79 years..... 80-84 years..... 8 8 2 . . . 2 85-89 years..... 2 . . . 90-94 years..... 2 2 95-99 years..... 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Neoplasm of uncertain or unknown behavior of urinary organs (D41) All ages.... 79 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 50-54 years..... 1 55-59 years..... 6 2 4 6 60-64 years..... 5 3 2 65-69 vears..... 5 70-74 vears..... 9 5 4 75-79 years..... 18 9 9 5 3 13 2 80-84 years..... 39 2.4 15 36 23 13 1 38 36 2 85-89 years..... 18 20 17 19 3 90-94 years..... 40 17 23 34 15 19 3 3 95-99 years..... 9 14 5 13 9 4 100 years & over. 2 4 3 1 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Kidney (D41.0) All ages.... 31 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 4 75-79 years..... 8 80-84 years..... 13 13 16 85-89 years..... 9 15 90-94 years..... 8 4 95-99 years..... 5 100 years & over. 2 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Renal pelvis (D41.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Ureter (D41.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Disc												As	ian or		
Tenth Revision, Second	d	All rac	es		White			Black			ican India	an 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Fe	emale	sexes	Male	Female
Urethra (D41.3)															
All ages	3	2	1	3	2	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.0-14 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-
.5-19 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-
20-24 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-
25-29 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-
30-34 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
10-44 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
5-49 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-
0-54 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
0-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-69 years	_	_	-	_	_	_	-	-	-	_	-	-	_	-	-
0-74 years	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_
5-79 years	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_
0-84 years	1	1	_	1	1	_	_	_	_	_	_	-	_	_	_
5-89 years	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_
0-94 years	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_
95-99 years	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Bladder (D41.4) All ages.... 107 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 4 2 2 60-64 years..... 2 1 1 65-69 vears..... 4 3 70-74 vears..... 5 3 75-79 years..... 9 5 80-84 years..... 2.4 15 9 21 2 1 21 20 85-89 years..... 10 11 9 11 90-94 years..... 2.8 12 23 10 13 3 3 16 95-99 years..... 8 3 4 4 4 100 years & over. 2 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Urinary organ, unspecified (D41.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,	All race	og.		White			Black		ħmo:	cican Indi		ian or	ia Tala	ınder 2/
Edition, 2004)	Both	AII IACE	25	Both	WIIICE		Both	Black		Both	.ican indi	all 1/	Both	ic isia	ilidel 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Jeoplasm of uncertain	or unknown	behavior o	of meninges	(D42)											
All ages	15	6	9	14	5	9	_	_	-	_	_	_	1	1	_
Inder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-49 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-59 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-74 years	1	1	-	-	-	-	-	-	-	-	-	-	1	1	-
5-79 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-
)-84 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
5-89 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
)-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
00 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cerebral meninges (D42.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Meninges, unspecified (D42.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 2 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Bot.h Both Bot.h Both Bot.h and age Female Male Female Male Female sexes Male Female sexes Male Female sexes Male sexes sexes Neoplasm of uncertain or unknown behavior of brain and central nervous system (D43) All ages.... 2,941 1,407 1,534 1,344 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2.0 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Brain, supratentorial (D43.0) All ages..... 129 72 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 4 35-39 years..... 40-44 years..... 2 45-49 years..... 9 4 5 2 2 50-54 years..... 13 1.0 55-59 years..... 7 2 5 6 60-64 years..... 10 6 3 65-69 vears..... 4 3 70-74 vears..... 9 5 4 75-79 years..... 2.0 11 80-84 years..... 21 8 13 19 12 2 1 23 13 85-89 years..... 16 19 3 90-94 years..... 18 8 10 17 10 1 1 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

on the International Classification of Dis Tenth Revision, Secon		All rac	es		White			Black		Amer	rican Ind		ian or Pacif	ic Isla	nder 2/
Edition, 2004)	Both	1111 140	CD	Both	WIIICC		Both	Diack		Both	rean inc	arum 17	Both	.10 1010	ilidel 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Brain, infratentorial	(D43.1)														
All ages	92	42	50	79	38	41	9	2	7	2	1	1	2	1	1
Under 1 year	1	1	-	1	1	-	-	-	-	-	_	-	-	-	-
1-4 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-9 years	7	2	5	7	2	5	-	-	-	-	-	-	-	-	-
10-14 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-39 years	5	3	2	2	2	-	2	1	1	1	-	1	-	-	-
40-44 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-
45-49 years	6	1	5	4	1	3	1	-	1	-	-	-	1	-	1
50-54 years	5	3	2	4	2	2	-	-	-	-	_	-	1	1	-
55-59 years	1	1	-	1	1	-	-	-	-	-	_	-	-	-	-
60-64 years	8	6	2	8	6	2	-	-	-	-	_	-	-	-	-
65-69 years	6	2	4	6	2	4	-	-	-	-	-	-	-	-	-
70-74 years	3	2	1	2	2	-	1	-	1	-	-	-	-	-	-
75-79 years	12	5	7	9	4	5	2	-	2	1	1	-	-	-	-
80-84 years	15	4	11	15	4	11	-	-	-	-	-	-	-	-	-
85-89 years	6	2	4	5	2	3	1	-	1	-	-	-	-	-	-
90-94 years	6	3	3	5	3	2	1	-	1	-	-	-	-	-	-
95-99 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-
100 years & over.	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Brain, unspecified (D43.2) All ages.... 2,635 1,265 1,370 2,317 1,111 1,206 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 2.0 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... ρ 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Disea	ses,											As	ian or		
Tenth Revision, Second		All rac	es		White			Black		Amea	cican Indi	an 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Cranial nerves (D43.3)															
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Jnder 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
10-44 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-
15-49 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	_	-	-	-	-	_	-	_	-	-	-
'5-79 years	_	_	-	_	_	_	-	-	-	_	-	-	-	-	-
0-84 years	_	_	-	_	_	_	-	-	-	_	-	-	-	-	-
35-89 years	-	-	-	-	_	-	-	-	-	_	-	_	-	-	-
90-94 years	-	-	-	-	_	-	-	-	-	_	-	_	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	-	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Spinal cord (D43.4) All ages.... 33 19 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 2 2 65-69 vears..... 2 70-74 vears..... 4 75-79 years..... 80-84 years..... 8 85-89 years..... 3 3 3 90-94 years..... 95-99 years..... 2 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dise	eases,											As	ian or		
Tenth Revision, Second	d	All rac	es		White			Black			rican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	remale	sexes	Male	Female
Central nervous system	m, unspecifi	ed (D43.9)												
All ages	30	18	12	28	17	11	1	_	1	_	_	_	1	1	_
Under 1 year	1	1	-	-	-	-	-	-	-	_	-	-	1	1	-
1-4 years	2	1	1	2	1	1	-	-	-	_	-	-	-	-	-
5-9 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	_	-	-	-	_	-	-	-	-	-
15-19 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	_
20-24 years	-	-	-	-	-	_	-	-	-	_	-	-	-	-	-
25-29 years	-	-	-	-	-	_	-	-	-	_	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	_	-	_	-	-	-	-	_	-	-
40-44 years	-	-	-	-	-	_	-	_	-	-	-	-	_	-	-
15-49 years	-	-	-	-	-	_	-	-	-	_	-	-	-	-	-
50-54 years	3	1	2	3	1	2	-	-	-	_	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
50-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	_	-	-	-	_	-	-	-	-	-
75-79 years	4	1	3	3	1	2	1	-	1	_	-	-	_	_	-
30-84 years	7	4	3	7	4	3	-	_	-	_	_	-	_	_	-
85-89 years	2	1	1	2	1	1	-	_	-	_	_	-	_	_	-
90-94 years	2	1	1	2	1	1	_	-	-	_	-	-	_	_	-
95-99 years	1	1	-	1	1	_	_	-	-	_	-	-	_	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Neoplasm of uncertain or unknown behavior of endocrine glands (D44) All ages.... 73 43 Under 1 year.... 1-4 years..... 2 5-9 years..... 10-14 years..... 3 1 2 2 2 15-19 years..... 2 2 2 20-24 years..... 3 3 25-29 years..... 30-34 years..... 35-39 years..... 5 40-44 years..... 12 45-49 years..... 14 1 0 4 50-54 years..... 14 11 3 6 55-59 years..... 13 8 9 4 5 5 60-64 years..... 20 13 15 11 65-69 vears..... 17 12 9 70-74 vears..... 4 75-79 years..... 22 14 80-84 years..... 19 8 11 14 4 3 85-89 years..... 10 9 10 9 2 90-94 years..... 12 7 10 2 5 5 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Disea	ases,												sian or		
Tenth Revision, Second		All rac	ces		White			Black			rican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale ————	sexes	Male ———	Female
Thyroid gland (D44.0)															
All ages	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
'5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
00-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Dis												As	ian or		
Tenth Revision, Secon	d	All rac	es		White			Black			cican Indi	an 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Adrenal gland (D44.1)															
All ages	15	10	5	13	8	5	2	2	_	_	_	_	_	_	_
Jnder 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
.0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	_	-	-	-	_	-	-	-	-	-
10-44 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-64 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
55-69 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
5-79 years	3	3	-	3	3	_	-	_	-	_	-	-	-	-	-
0-84 years	1	-	1	1	-	1	-	_	-	_	-	-	-	-	-
5-89 years	1	-	1	1	-	1	-	_	-	_	-	-	-	-	-
0-94 years	1	1	-	1	1	_	-	_	-	_	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	_	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Classification of Dis- Tenth Revision, Second		All rac	105		White			Black		\lambda mos	rican Ind		sian or	ia Tala	nder 2/
Edition, 2004)	a Both	AII Iac	es	Both	WIIICE		Both	Black		Both	ican inu	IIaII I/	Both	IC ISIA	muer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Parathyroid gland (D4	4.2)														
All ages	1	_	1	1	-	1	_	_	_	-	-	-	-	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
15-19 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	_	_	-	-	-	-	-	-
95-99 years	1	-	1	1	-	1	-	_	_	-	-	-	-	-	-
100 years & over.	_	-	-	-	-	-	_	-	_	_	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pituitary gland (D44.3) All ages.... 27 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 6 2 50-54 years..... 3 3 1 55-59 years..... 4 3 1 60-64 years..... 6 5 1 65-69 vears..... 11 70-74 vears..... 3 3 75-79 years..... 12 3 80-84 years..... 9 5 2 1 1 85-89 years..... 8 90-94 years..... 11 4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Craniopharyngeal duct (D44.4) All ages.... 27 10 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 3 1 2 2 2 15-19 years..... 2 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 2 2 5 2 50-54 years..... 1 55-59 years..... 6 2 4 60-64 years..... 6 1 5 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 4 1 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Classification of Dise Tenth Revision, Second		All rac	100		White			Black		λma	rican Ind		ian or	fic Tels	ınder 2/
Edition, 2004)	Both	AII IAC	.cs	Both	WIIICE		Both	Black		Both	ICan Inc	itali 1/	Both	.IC ISIA	ilidel 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Pineal gland (D44.5)															
All ages	11	6	5	10	5	5	-	_	-	_	_	-	1	1	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
40-44 years	1	-	1	1	_	1	-	_	-	-	-	-	-	-	-
45-49 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
50-54 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	2	1	1	1	-	1	-	-	-	-	-	-	1	1	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International												_			
Classification of Dise Tenth Revision, Second		All rac			White			Black		7 m o n	ican Indi		ian or	ia Talo	nder 2/
Edition, 2004)	a Both	All rac	es	Both	wille		Both	Black		Both	Tean Indi	an 1/	Both	IC ISIA	nder 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Carotid body (D44.6)															
All ages	4	2	2	2	2	_	2	_	2	_	_	-	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
50-64 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
'5-79 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
0-84 years	2	1	1	1	1	-	1	-	1	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
00-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis	eases,											As	ian or		
Tenth Revision, Secon	ıd	All rac	es		White			Black		Amea	ican Indi	an 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Aortic body and other	paraganglia	a (D44.7)													
All ages	8	5	3	8	5	3	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	_	-	_	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-
0-24 years	_	-	-	-	-	_	_	_	-	_	-	-	-	-	-
25-29 years	1	-	1	1	-	1	_	_	-	_	-	-	-	-	-
30-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_
0-44 years	1	1	_	1	1	_	_	_	_	_	_	_	_	-	_
15-49 years	1	1	-	1	1	_	_	_	_	_	-	-	-	-	-
50-54 years	1	1	-	1	1	_	_	_	_	_	-	-	-	-	_
55-59 years	-	-	-	-	-	_	_	_	_	_	-	-	-	-	-
60-64 years	1	1	-	1	1	_	_	-	-	-	_	-	_	-	-
65-69 years	_	_	-	-	_	_	_	-	-	-	_	-	_	-	-
70-74 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
75-79 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
0-84 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
15-89 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
00-94 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
95-99 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pluriglandular involvement (D44.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Endocrine gland, unspecified (D44.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 4 2 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Polycythemia vera (D45) All ages..... 281 116 165 264 111 153 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 1 5 50-54 years..... 3 2 55-59 years..... 8 3 5 8 5 60-64 years..... 14 14 65-69 vears..... 13 9 13 9 70-74 vears..... 32 17 15 30 13 75-79 years..... 42 2.0 2.2 38 19 19 80-84 years..... 58 20 38 54 18 36 2 42 27 41 85-89 years..... 15 15 26 90-94 years..... 39 1.0 29 3.8 10 28 95-99 years..... 3 16 12 9 3 4 1.3 4 100 years & over. 1 1 1 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Myelodysplastic syndromes (D46) All ages.... 5,182 2,928 2,254 4,803 2,713 2,090 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 1 Q 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1,172 1.112 1,034 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Refractory anemia with sideroblasts (D46.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Refractory anemia with excess of blasts (D46.2) All ages.... 19 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 3 70-74 vears..... 3 75-79 years..... 80-84 years..... 5 3 85-89 years..... 90-94 years..... 3 3 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

on the International Classification of Dis		-11						-1 1			1.		ian or	1	3 04
Tenth Revision, Secon		All race	:S	D - 1-1-	White		D	Black			ican Indi	an 1/		ic Isla	nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	omalo	Both sexes	Mala	Female
											male r				
Refractory anemia wit	ch excess of	blasts wit	h transform	ation (D46.3	3)										
All ages	2	1	1	2	1	1	-	-	_	-	-	-	-	-	-
Under 1 year	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
75-79 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	1	_	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Refractory anemia, unspecified (D46.4) All ages..... 114 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 3 50-54 years..... 1 55-59 years..... 2 60-64 years..... 65-69 vears..... 9 6 70-74 vears..... 15 8 75-79 years..... 15 13 3 80-84 years..... 22 15 19 14 33 85-89 years..... 13 20 32 13 19 1 90-94 years..... 16 7 9 16 a 95-99 years..... 7 8 2 1 6 2 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International Classification of Diseases, Tenth Revision, Second All races				White Black						Asian or American Indian 1/ Pacific Islander 2/					
		AII IACES		Dath	white		Dath	Black		American Indian 1/ Both		Both			
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Other myelodysplastic	c syndromes (D46.7)													
All ages	5	2	3	5	2	3	-	-	-	-	-	-	-	_	-
Under 1 year	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
'0-74 years	2	-	2	2	-	2	-	_	-	-	-	-	-	-	-
5-79 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Male Female Female Male Female Male Female sexes Male Female sexes sexes Male sexes sexes Myelodysplastic syndrome, unspecified (D46.9) All ages.... 5,028 2,850 2,178 4,662 2,639 2,023 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 1 Q 2.2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1,145 1,088 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Bot.h Both Bot.h Both Bot.h and age Female Male Female Male Female sexes Male Female sexes Male Female sexes Male sexes sexes Other neoplasms of uncertain or unknown behavior of lymphoid, hematopoietic, and related tissue (D47) All ages.... 1,549 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a Ω 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Chronic myeloproliferative disease (D47.1) All ages.... 1,333 1,225 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... R 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Monoclonal gammopathy (D47.2) All ages..... 33 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 3 50-54 years..... 2 1 55-59 years..... 1 1 60-64 years..... 4 2 65-69 vears..... 4 3 70-74 vears..... 6 3 3 75-79 years..... 13 8 11 80-84 years..... 16 9 7 14 1 16 85-89 years..... 9 15 90-94 years..... 8 3 5 3 5 1 95-99 years..... 3 3 1 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Essential (hemorrhagic) thrombocythemia (D47.3) All ages..... 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 5 85-89 years..... 3 90-94 years..... 1 3 4 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Bot.h Both and age Female Female Male Female Male Female Male Female sexes Male sexes Male sexes sexes sexes Other specified neoplasms of uncertain or unknown behavior of lymphoid, hematopoietic, and related tissue (D47.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Bot.h Both and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Neoplasm of uncertain or unknown behavior of lymphoid, hematopoietic, and related tissue, unspecified (D47.9) All ages.... Under 1 year.... 1 1 1 1 1-4 years..... 3 2 2 2 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 2 2 5 50-54 years..... 3 3 1 55-59 years..... 6 2 4 5 60-64 years..... 5 3 2 65-69 vears..... 4 3 70-74 vears..... 7 3 75-79 years..... 10 10 80-84 years..... 23 12 11 2.1 11 10 1 15 85-89 years..... 8 13 6 90-94 years..... 9 5 4 9 95-99 years..... 6 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Neoplasm of uncertain or unknown behavior of other and unspecified sites (D48) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Bone and articular cartilage (D48.0) All ages.... 19 Under 1 year.... 8 3 5 3 5 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 2 1 1 60-64 years..... 4 2 2 65-69 vears..... 2 70-74 vears..... 2 75-79 years..... 80-84 years..... 8 5 85-89 years..... 90-94 years..... 5 95-99 years..... 2 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Connective and other soft tissue (D48.1) All ages.... 109 176 91 13 Under 1 year.... 2 2 1 1 1 1 1-4 years..... 1 1 1 1 5-9 years..... 10-14 years..... 1 1 15-19 years..... 2 1 1 20-24 years..... 3 2 25-29 years..... 30-34 years..... 35-39 years..... 4 7 40-44 years..... 45-49 years..... 15 8 12 2 50-54 years..... 6 3 3 3 6 55-59 years..... 18 8 10 16 60-64 years..... 14 11 3 11 2 2 65-69 vears..... 28 24 13 11 13 15 70-74 vears..... 25 14 11 20 12 8 3 75-79 years..... 34 17 17 32 16 16 80-84 years..... 21 11 10 16 9 1 12 2 85-89 years..... 5 7 9 3 2 90-94 years..... 6 5 1 6 95-99 years..... 3 2 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Peripheral nerves and autonomic nervous system (D48.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 2 50-54 years..... 2 2 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis					Asian or										
Tenth Revision, Second		All rac	es	D	White		D - 1-1-	Black			rican Indi	an 1/	Pacific Isla		nder 2/
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	'omalo	Both sexes	Mala	Female
						remare						Ciliaic		Haic	remare
Retroperitoneum (D48.	3)														
All ages	15	7	8	13	7	6	2	_	2	_	-	-	-	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
70-74 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
75-79 years	5	3	2	5	3	2	-	-	-	-	-	-	-	-	-
80-84 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
85-89 years	4	1	3	3	1	2	1	-	1	-	-	-	-	-	-
90-94 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	-	-	-	_	-	-	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Peritoneum (D48.4) All ages.... 16 15 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 1 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 1 1 65-69 vears..... 70-74 vears..... 4 75-79 years..... 80-84 years..... 2 1 85-89 years..... 90-94 years..... 1 95-99 years..... 1 1 1 1 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Skin (D48.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Breast (D48.6) All ages.... 33 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 3 5 50-54 years..... 5 55-59 years..... 2 2 60-64 years..... 5 5 65-69 vears..... 2 70-74 vears..... 5 5 75-79 years..... 3 80-84 years..... 5 5 85-89 years..... 4 3 1 90-94 years..... 4 4 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified sites (D48.7) All ages.... 179 222 149 33 Under 1 year.... 6 1 6 1 1-4 years..... 2 2 2 2 5-9 years..... 1 10-14 years..... 15-19 years..... 1 20-24 years..... 1 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 2 45-49 years..... 3 1 2 6 50-54 years..... 3 3 3 3 6 55-59 years..... 14 9 5 12 60-64 years..... 15 8 9 5 5 1 65-69 vears..... 11 5 10 6 70-74 vears..... 14 7 12 7 75-79 years..... 2.3 19 12 16 80-84 years..... 46 14 32 36 10 26 2 52 7 45 48 85-89 years..... 6 42 4 3 1 90-94 years..... 34 5 29 32 27 1 1 5 95-99 years..... 19 18 13 12 6 1 6 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Neoplasms of uncertain or unknown behavior, unspecified (D48.9) All ages.... 119 37 18 19 Under 1 year.... 4 3 3 2 1 3 2 1 1 1-4 years..... 4 3 1 2 2 1 5-9 years..... 10-14 years..... 2 15-19 years..... 1 20-24 years..... 3 3 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 45-49 years..... 6 2 4 5 2 2 2.0 50-54 years..... 9 17 9 11 55-59 years..... 20 9 11 16 8 8 1 60-64 years..... 27 13 14 21 11 10 5 65-69 vears..... 20 9 16 7 9 70-74 years..... 24 13 11 20 75-79 years..... 2.8 14 2.4 12 14 80-84 years..... 37 17 20 32 14 18 2 25 25 85-89 years..... 8 17 8 17 2 90-94 years..... 2.4 9 15 22 14 1 8 95-99 years..... 4 4 4 4 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Iron deficiency anemia (D50) All ages..... 117 141 102 18 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 5 2 3 60-64 years..... 3 1 2 65-69 vears..... 9 5 70-74 vears..... 14 7 75-79 years..... 13 11 9 11 80-84 years..... 28 5 23 26 22 2 1 29 18 2 85-89 years..... 31 12 19 11 1 90-94 years..... 33 5 28 29 5 24 4 95-99 years..... 17 12 14 10 3 100 years & over. 5 4 5 4

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Iron deficiency anemia secondary to blood loss (chronic) (D50.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 55-59 years..... 2 2 60-64 years..... 2 1 1 65-69 vears..... 3 70-74 vears..... 8 75-79 years..... 8 80-84 years..... 17 14 16 13 1 18 85-89 years..... 11 17 10 90-94 years..... 16 2 14 13 11 3 3 2 95-99 years..... 7 8 3 5 4 1 1 100 years & over. 2 2 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other iron deficiency anemias (D50.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 3 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Iron deficiency anemia, unspecified (D50.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 3 3 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 6 3 75-79 years..... 3 80-84 years..... 10 8 9 13 85-89 years..... 8 12 8 15 90-94 years..... 3 12 14 3 11 1 95-99 years..... 6 5 5 4 1 1 1 1 100 years & over. 3 2 2 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Vitamin B-12 deficiency anemia (D51) All ages.... 30 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 6 85-89 years..... 12 6 11 2 90-94 years..... 1.0 8 4 6 95-99 years..... 8 100 years & over. 5

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Vitamin B-12 deficiency anemia due to intrinsic factor deficiency (D51.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 6 85-89 years..... 10 5 9 2 90-94 years..... 1.0 6 95-99 years..... 8 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Vitamin B-12 deficiency anemia, unspecified (D51.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Folate deficiency anemia (D52) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Folate deficiency anemia, unspecified (D52.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other nutritional anemias (D53) All ages..... 32 10 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 2 60-64 years..... 4 1 3 65-69 vears..... 4 70-74 vears..... 5 3 75-79 years..... 3 80-84 years..... 4 1 1 85-89 years..... 13 3 10 10 7 90-94 years..... 2 5 3 1 3 95-99 years..... 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

ause of death (based n the International															
Classification of Dis	eases,				Asian or										
Tenth Revision, Secon	.d	All rac	es		White			Black		Amen	rican Indi	ian 1/	Pacif	ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female
Protein deficiency an	emia (D53.0)														
All ages	23	10	13	16	5	11	7	5	2	_	_	_	_	_	_
Under 1 year	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	_	-	-	_	-	-	-	-	-	_	-	-	-	-	-
25-29 years	_	-	-	_	-	-	-	-	-	_	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	_	_	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	_	_	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	_	_	-	-	-	-	_	-	-	_	-	-
50-54 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
55-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
60-64 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
65-69 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-
70-74 years	1	1	-	_	_	-	1	1	-	_	-	-	_	-	-
75-79 years	2	2	-	1	1	-	1	1	-	_	-	-	_	-	-
80-84 years	2	2	-	2	2	-	-	-	-	_	-	-	_	-	-
85-89 years	6	1	5	5	1	4	1	-	1	_	-	-	_	-	-
90-94 years	4	2	2	2	_	2	2	2	-	_	-	-	_	-	-
95-99 years	2	-	2	2	-	2	-	-	-	_	-	-	-	-	-
100 years & over.	1	_	1	_	_	_	1	_	1	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Nutritional anemia, unspecified (D53.9) All ages.... 19 18 13 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 3 1 65-69 vears..... 70-74 vears..... 4 75-79 years..... 2 80-84 years..... 2 1 1 5 85-89 years..... 3 90-94 years..... 3 1 1 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Anemia due to enzyme disorders (D55) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Anemia due to glucose-6-phosphate dehydrogenase [G6PD] deficiency (D55.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Thalassemia (D56) All ages.... 21 Under 1 year.... 5 2 1 2 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 2 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Alpha thalassemia (D56.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Beta thalassemia (D56.1) All ages.... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Disc Tenth Revision, Second	etal a series and a							Asian or American Indian 1/ Pacific Islander 2/							
Edition, 2004)	a Both	All races		Both	White		Black Both			Both	rican indi	lan 1/	Both	IC ISIA	naer 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	emale	sexes	Male	Female
Thalassemia trait (D5)	6.3)														
All ages	1	_	1	_	_	_	_	-	-	_	-	_	1	-	1
Under 1 year	1	-	1	_	-	-	-	-	-	_	-	-	1	-	1
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
10-14 years	_	-	-	_	-	-	-	-	-	_	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	_	-	-	_	-	-	-	-	_	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
45-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	_	-	-	_	-	-	-	-	-	_	-	-	-	-	-
55-59 years	_	-	-	_	-	-	-	-	_	-	-	-	-	-	-
60-64 years	_	-	-	_	-	-	-	-	_	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_		_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or															
Tenth Revision, Secon	d	All races			White			Black		Amer	ican Indi	an 1/	Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Other thalassemias (D	56.8)														
All ages	1	1	_	_	_	_	1	1	_	_	_	_	_	_	_
Under 1 year	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
1-4 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
5-9 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
10-14 years	_	_	-	-	-	-	-	_	-	-	-	-	-	-	-
15-19 years	_	_	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	_	_	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	_	_	-	-	-	-	-	_	-	-	-	-	-	-	-
50-54 years	_	_	-	-	-	-	-	_	-	-	-	-	-	-	-
55-59 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
75-79 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
30-84 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
85-89 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
90-94 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
95-99 years	_	-	-	-	-	-	-	-	-	-	_	-	-	-	-
100 years & over.	_	_	-	_	-	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Thalassemia, unspecified (D56.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Sickle-cell disorders (D57) All ages.... 489 229 260 13 466 219 247 Under 1 year.... 3 3 3 3 1-4 years..... 11 9 2 11 9 2 3 3 5-9 years..... 6 6 3 3 10-14 years..... 2 5 7 2 5 15-19 years..... 2.2 2.0 1.0 12 10 10 20-24 years..... 38 24 14 38 24 25-29 years..... 54 29 25 51 30-34 years..... 47 22 25 47 22 25 31 35-39 years..... 54 21 33 51 20 39 77 39 38 40-44 years..... 79 40 31 45-49 years..... 56 24 32 54 23 2.8 2.7 50-54 years..... 42 14 40 13 55-59 years..... 35 18 17 34 17 17 60-64 years..... 17 10 13 5 8 65-69 vears..... 8 6 6 70-74 vears..... 4 3 75-79 years..... 3 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Sickle-cell anemia with crisis (D57.0) All ages.... 97 Under 1 year.... 2 2 2 1-4 years..... 3 3 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 11 11 30-34 years..... 12 12 35-39 years..... 13 12 40-44 years..... 12 12 45-49 years..... 9 3 6 9 9 9 50-54 years..... R 55-59 years..... 9 5 4 60-64 years..... 4 1 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Sickle-cell anemia without crisis (D57.1) All ages.... 200 16 356 164 192 Under 1 year.... 1 1 1 1 1-4 years..... 8 2 8 6 2 6 3 5-9 years..... 3 6 3 3 10-14 years..... 6 1 5 6 1 5 15-19 years..... 15 9 6 1.3 6 20-24 years..... 27 15 12 27 15 12 25-29 years..... 40 37 19 30-34 years..... 34 13 21 34 13 21 23 35-39 years..... 39 16 37 15 32 31 40-44 years..... 66 34 33 26 25 45-49 years..... 47 21 45 20 19 12 19 50-54 years..... 32 13 31 55-59 years..... 25 14 11 24 13 11 60-64 years..... 11 5 6 11 5 6 65-69 vears..... 5 70-74 vears..... 3 3 75-79 years..... 3 80-84 years..... 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Double heterozygous sickling disorders (D57.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 2 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Sickle-cell trait (D57.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other hereditary hemolytic anemias (D58) All ages..... 116 70 Under 1 year.... 1-4 years..... 2 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 2 2 5 50-54 years..... 2 3 2 55-59 years..... 8 4 4 60-64 years..... 4 1 3 65-69 vears..... 5 3 70-74 vears..... 13 6 75-79 years..... 18 6 12 18 12 80-84 years..... 23 10 13 21 13 2 21 12 85-89 years..... 24 9 15 90-94 years..... 9 5 4 9 4 95-99 years..... 5 1 100 years & over. 3 3 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Hereditary spherocytosis (D58.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis										Asian or						
Tenth Revision, Secon		All races		White			Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both		_ ,	Both			Both		_ 1	Both		_ ,	Both		_ ,	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	F'emale	sexes	Male	Female	
Other hemoglobinopath	ies (D58.2)															
All ages	4	2	2	1	_	1	3	2	1	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-	
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	1	1	-	-	-	-	1	1	-	-	_	-	-	-	-	
40-44 years	1	-	1	1	-	1	-	-	-	-	_	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-	
50-54 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
80-84 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Age not stated	-	-	-	-	_	-	-	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hereditary hemolytic anemia, unspecified (D58.9) All ages.... 114 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 2 45-49 years..... 2 50-54 years..... 3 2 2 1 55-59 years..... 8 4 4 60-64 years..... 4 1 3 65-69 vears..... 5 3 70-74 vears..... 13 6 75-79 years..... 18 6 12 18 12 80-84 years..... 23 10 13 21 13 2 21 12 85-89 years..... 24 9 15 90-94 years..... 9 5 9 4 4 95-99 years..... 5 1 100 years & over. 3 3 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acquired hemolytic anemia (D59) All ages.... 56 87 131 79 Under 1 year.... 1-4 years..... 5 5-9 years..... 10-14 years..... 2 1 1 2 15-19 years..... 1 20-24 years..... 4 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 3 45-49 years..... 3 3 2 50-54 years..... 3 1 55-59 years..... 10 4 6 5 60-64 years..... 9 5 1 1 65-69 vears..... 3 3 70-74 vears..... 14 10 13 75-79 years..... 15 11 15 11 80-84 years..... 2.7 10 17 23 9 14 1 23 23 85-89 years..... 11 12 11 12 90-94 years..... 8 3 5 3 5 8 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other autoimmune hemolytic anemias (D59.1) All ages.... 49 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 1 1 55-59 years..... 3 2 3 1 60-64 years..... 5 2 3 65-69 vears..... 70-74 vears..... 9 5 75-79 years..... 11 8 80-84 years..... 20 8 12 17 10 21 21 85-89 years..... 11 10 11 10 7 90-94 years..... 2 5 2 5 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Drug-induced nonautoimmune hemolytic anemia (D59.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hemolytic-uremic syndrome (D59.3) All ages..... 22 19 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 1 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other nonautoimmune hemolytic anemias (D59.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 3 3 60-64 years..... 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Paroxysmal nocturnal hemoglobinuria [Marchiafava-Micheli] (D59.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Classification of Dis					Asian or										
Tenth Revision, Secon		All races			White			Black		American Indian 1/			Pacific Islander 2		
Edition, 2004)	Both	1		Both			Both			Both			Both		_ 1
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale ————	sexes	Male	Female
Acquired hemolytic an	emia, unspec	cified (D5	9.9)												
All ages	9	6	3	9	6	3	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	_	-	-	-	-	-	-	-	-	_	_	-	-	-	-
15-19 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	_	-	-	-	_	-	-	-	-	-	-
25-29 years	-	-	-	-	_	-	-	-	_	-	-	-	-	-	-
30-34 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
35-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_
0-44 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_
15-49 years	_	_	-	-	-	-	-	_	-	_	-	-	-	-	-
50-54 years	1	1	-	1	1	-	-	_	-	_	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	_	-	_	-	-	-	-	-
60-64 years	_	-	-	_	-	_	-	-	-	_	_	-	-	-	-
65-69 years	1	1	-	1	1	_	-	-	-	_	_	-	-	-	-
70-74 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
'5-79 years	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_
30-84 years	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
35-89 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
00-94 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
95-99 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_	_	_	_	_		_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acquired pure red cell aplasia [erythroblastopenia] (D60) All ages.... 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 85-89 years..... 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acquired pure red cell aplasia, unspecified (D60.9) All ages..... 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 85-89 years..... 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other aplastic anemias (D61) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 2.8 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Constitutional aplastic anemia (D61.0) All ages.... 11 15 Under 1 year.... 1-4 years..... 2 5-9 years..... 1 1 10-14 years..... 3 2 1 1 1 15-19 years..... 3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis					Asian or										
Tenth Revision, Secon		All races			White		_	Black		American Indian 1/			Pacific Islander 2		
Edition, 2004)	Both	1	_ ,	Both			Both		_ ,	Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Idiopathic aplastic a	nemia (D61.3	3)													
All ages	5	1	4	4	1	3	_	_	_	_	_	_	1	_	1
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	-	-	_	-	-	-	_	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	1	1	-	1	1	-	-	_	-	-	-	_	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-84 years	1	-	1	-	-	-	-	-	-	-	-	-	1	-	1
85-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Aplastic anemia, unspecified (D61.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 2.8 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acute posthemorrhagic anemia (D62) All ages..... 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 5 85-89 years..... 8 5 90-94 years..... 1 95-99 years..... 1 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Other anemias (D64) All ages.... 2,015 1,271 1,679 1,062 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 8.5 2.8 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other sideroblastic anemias (D64.3) All ages..... 10 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 8 90-94 years..... 4 4 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other specified anemias (D64.8) All ages..... 16 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 2 2 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 5 75-79 years..... 11 6 80-84 years..... 7 3 2 85-89 years..... 6 3 90-94 years..... 3 2 1 95-99 years..... 1 4 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Anemia, unspecified (D64.9) All ages.... 1,950 1,241 1,620 1,036 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 2.4 1 0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 2.0 75-79 years..... 2.8 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disseminated intravascular coagulation [defibrination syndrome] (D65) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.4 2.2 80-84 years..... 2.4 2.4 85-89 years..... 90-94 years..... 2.0 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hereditary factor VIII deficiency (D66) All ages.... 19 12 Under 1 year.... 1 1 1 1-4 years..... 2 2 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 1 1 1 3 50-54 years..... 55-59 years..... 2 2 1 60-64 years..... 1 65-69 vears..... 4 70-74 vears..... 4 75-79 years..... 9 80-84 years..... 11 3 2 2 1 8 2 85-89 years..... 3 5 90-94 years..... 3 4 95-99 years..... 3 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Hereditary factor IX deficiency (D67) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other coagulation defects (D68) All ages.... 1,148 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Von Willebrand's disease (D68.0) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hereditary deficiency of other clotting factors (D68.2) All ages.... Under 1 year.... 2 2 2 2 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 2 5 50-54 years..... 3 55-59 years..... 1 1 60-64 years..... 3 2 65-69 vears..... 5 70-74 vears..... 3 3 75-79 years..... 80-84 years..... 8 2 85-89 years..... 6 90-94 years..... 2 95-99 years..... 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acquired coagulation factor deficiency (D68.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

24JUN09

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified coagulation defects (D68.8) All ages.... 13 Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 2 1 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 9 40-44 years..... 9 45-49 years..... 5 4 9 50-54 years..... 2 R 2 55-59 years..... 12 8 10 4 60-64 years..... 17 13 16 13 65-69 vears..... 10 9 6 70-74 vears..... 2 75-79 years..... 9 80-84 years..... 10 1 2 85-89 years..... 4 1 90-94 years..... 3 1 2 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Coagulation defect, unspecified (D68.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Purpura and other hemorrhagic conditions (D69) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 2.2 50-54 years..... 2.4 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Allergic purpura (D69.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Qualitative platelet defects (D69.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other nonthrombocytopenic purpura (D69.2) All ages..... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Idiopathic thrombocytopenic purpura (D69.3) All ages.... 152 230 106 124 34 13 Under 1 year.... 1-4 years..... 3 1 5-9 years..... 1 1 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 4 35-39 years..... 40-44 years..... 3 9 45-49 years..... 3 6 2 5 6 50-54 years..... 13 8 55-59 years..... 10 6 4 3 60-64 years..... 15 8 11 5 1 65-69 vears..... 13 6 7 13 7 70-74 vears..... 23 12 22 12 10 75-79 years..... 37 21 16 29 17 12 3 80-84 years..... 57 22 35 54 2.2 32 2 42 40 85-89 years..... 16 26 16 24 1 90-94 years..... 22 9 13 2.0 11 1 1 1 9 95-99 years..... 7 3 3 4 6 3 1 1 100 years & over. 2 2 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other primary thrombocytopenia (D69.4) All ages..... 33 Under 1 year.... 2 2 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 2 5 50-54 years..... 1 55-59 years..... 3 3 60-64 years..... 2 1 65-69 vears..... 70-74 vears..... 4 3 75-79 years..... 9 80-84 years..... 6 5 1 85-89 years..... 4 3 90-94 years..... 5 2 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Secondary thrombocytopenia (D69.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 50-54 years..... 3 2 1 55-59 years..... 2 1 1 60-64 years..... 4 2 2 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 80-84 years..... 9 5 1 85-89 years..... 6 6 90-94 years..... 1 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Thrombocytopenia, unspecified (D69.6) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... R 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.2 80-84 years..... 2.4 85-89 years..... 2.2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified hemorrhagic conditions (D69.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hemorrhagic condition, unspecified (D69.9) All ages.... 19 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 5 50-54 years..... 3 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 3 75-79 years..... 2 80-84 years..... 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Agranulocytosis (D70) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... Ω 2.0 50-54 years..... R 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.4 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Functional disorders of polymorphonuclear neutrophils (D71) All ages.... Under 1 year.... 1 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other disorders of white blood cells (D72) All ages..... 106 10 Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 1 50-54 years..... 4 3 2 1 55-59 years..... 6 2 4 60-64 years..... 3 2 1 3 65-69 vears..... 11 5 6 70-74 vears..... 11 6 5 9 75-79 years..... 19 12 18 80-84 years..... 2.0 8 12 2.0 8 12 22 21 16 85-89 years..... 5 17 90-94 years..... 1.0 5 5 10 5 95-99 years..... 6 6 4 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Genetic anomalies of leukocytes (D72.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Eosinophilia (D72.1) All ages.... Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 1 1 3 50-54 years..... 2 1 55-59 years..... 3 2 1 60-64 years..... 2 1 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 3 1 2 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified disorders of white blood cells (D72.8) All ages.... 49 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 1 1 55-59 years..... 3 60-64 years..... 1 1 65-69 vears..... 8 5 3 70-74 vears..... 10 75-79 years..... 15 9 6 80-84 years..... 17 6 11 17 11 20 19 15 85-89 years..... 4 16 9 90-94 years..... 5 4 9 4 95-99 years..... 6 6 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Diseases of spleen (D73) All ages.... 133 79 122 49 10 Under 1 year.... 2 2 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 3 1 1 3 50-54 years..... 1.0 55-59 years..... 7 2 5 6 1 60-64 years..... 12 9 3 11 65-69 vears..... 11 6 5 10 70-74 vears..... 17 8 17 75-79 years..... 19 18 15 80-84 years..... 20 11 9 2.0 11 9 85-89 years..... 12 4 8 12 7 90-94 years..... 3 4 3 95-99 years..... 2 1 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Hypersplenism (D73.1) All ages.... 19 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 4 70-74 vears..... 9 5 75-79 years..... 5 80-84 years..... 11 3 11 5 85-89 years..... 5 90-94 years..... 3 1 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Chronic congestive splenomegaly (D73.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Abscess of spleen (D73.3) All ages..... 13 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 2 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Infarction of spleen (D73.5) All ages..... 27 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 2 50-54 years..... 4 2 55-59 years..... 3 2 1 60-64 years..... 6 2 65-69 vears..... 4 3 70-74 vears..... 3 75-79 years..... 10 80-84 years..... 3 3 85-89 years..... 3 2 90-94 years..... 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																
Classification of Dise					Asian or											
Tenth Revision, Second	i	All rac	es		White			Black		Ame	rican Indi	an 1/				
Edition, 2004)	Both			Both			Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Other diseases of sple	een (D73.8)															
All ages	3	1	2	3	1	2	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
85-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disease of spleen, unspecified (D73.9) All ages.... 12 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 3 50-54 years..... 2 1 1 55-59 years..... 2 1 1 60-64 years..... 2 2 65-69 vears..... 3 70-74 vears..... 3 75-79 years..... 3 3 80-84 years..... 4 85-89 years..... 1 90-94 years..... 1 95-99 years..... 2 1 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Methemoglobinemia (D74) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other methemoglobinemias (D74.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

24JUN09

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																	
Classification of Diseases, Tenth Revision, Second All races				and to							Asian or American Indian 1/ Pacific Islander 2/						
Edition, 2004)	.a Both	All races		Both	White		Both	Black		Amer Both	ican indi	an 1/	Pacific Islander				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male Female		Both sexes Male		e Female		
															- CHICATE		
Other diseases of blo	od and blood	d-forming	organs (D75)														
All ages	131	48	83	115	41	74	13	7	6	_	_	_	3	_	3		
Under 1 year	3	1	2	2	1	1	-	-	-	-	-	-	1	-	1		
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
5-9 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
10-14 years	3	1	2	2	1	1	1	-	1	-	-	-	-	-	-		
15-19 years	2	-	2	1	-	1	1	-	1	-	-	-	-	-	-		
20-24 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-		
25-29 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
35-39 years	2	1	1	2	1	1	-	-	-	-	-	-	-	-	-		
40-44 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-		
45-49 years	5	2	3	4	2	2	-	-	-	-	-	-	1	-	1		
50-54 years	9	5	4	8	4	4	1	1	-	-	-	-	-	-	-		
55-59 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-		
60-64 years	7	1	6	7	1	6	-	-	-	-	-	-	-	-	-		
65-69 years	3	1	2	3	1	2	-	-	-	-	-	-	-	-	-		
70-74 years	10	7	3	7	6	1	3	1	2	-	-	-	-	-	-		
75-79 years	12	5	7	11	4	7	1	1	-	-	-	-	-	-	-		
80-84 years	26	10	16	24	9	15	2	1	1	-	-	-	-	-	-		
85-89 years	20	6	14	17	5	12	2	1	1	-	-	-	1	-	1		
90-94 years	11	3	8	10	2	8	1	1	-	-	-	-	-	-	-		
95-99 years	4	1	3	4	1	3	-	-	-	-	-	-	-	-	-		
100 years & over.	3	1	2	3	1	2	-	_	-	-	-	-	-	-	-		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																
Classification of Dis									Asian or							
Tenth Revision, Secon		All races			White		Black				rican Indi	an 1/	Pacific Islander 2			
Edition, 2004)	Both			Both			Both			Both			Both			
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale ————	sexes	Male	Female	
Secondary polycythemi	a (D75.1)															
All ages	2	1	1	2	1	1	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-	
15-19 years	-	-	-	_	-	_	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	_	-	-	-	_	-	-	_	-	-	
40-44 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	_	
45-49 years	1	1	-	1	1	_	-	-	-	_	-	-	_	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	-	-	_	-	-	-	_	-	-	_	-	-	
75-79 years	_	_	-	_	_	_	-	-	-	_	-	-	_	-	-	
30-84 years	1	_	1	1	_	1	-	-	-	_	-	-	_	-	-	
35-89 years	_	_	-	_	_	_	-	-	-	_	-	-	_	-	-	
90-94 years	-	-	-	-	-	_	-	-	-	_	-	-	_	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Essential thrombocytosis (D75.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 2 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 4 80-84 years..... 6 5 85-89 years..... 8 8 90-94 years..... 2 1 95-99 years..... 1 1 100 years & over. 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other specified diseases of blood and blood-forming organs (D75.8) All ages.... 29 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 3 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 4 1 3 65-69 vears..... 70-74 vears..... 6 75-79 years..... 4 80-84 years..... 15 8 15 10 85-89 years..... 5 8 7 90-94 years..... 1 6 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disease of blood and blood-forming organs, unspecified (D75.9) All ages.... 13 22 Under 1 year.... 2 2 1 1 1-4 years..... 5-9 years..... 10-14 years..... 3 2 2 1 15-19 years..... 1 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 2 1 1 50-54 years..... 4 2 2 55-59 years..... 2 2 60-64 years..... 2 2 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 2 85-89 years..... 90-94 years..... 2 1 1 2 1 95-99 years..... 2 1 1 2 1 100 years & over. 1

24JUN09

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Disea	ases,											As	ian or		
Tenth Revision, Second				White			Black		Amer	rican Ind			ic Isla	nder 2/	
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Certain diseases involv	ving lympho	oreticular	tissue and	reticulohis	stiocytic	system (D76	5)								
All ages	92	60	32	69	44	25	18	12	6	2	2	_	3	2	1
Under 1 year	16	13	3	10	8	2	5	5	-	-	-	-	1	-	1
1-4 years	11	10	1	8	8	-	2	1	1	1	1	-	-	-	-
5-9 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-
10-14 years	2	1	1	1	-	1	-	-	-	-	-	-	1	1	-
15-19 years	6	4	2	6	4	2	-	-	-	-	-	-	-	-	-
20-24 years	3	-	3	1	-	1	2	-	2	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-
40-44 years	6	3	3	5	2	3	-	-	-	-	-	-	1	1	-
45-49 years	5	3	2	4	3	1	1	-	1	-	-	-	-	-	-
50-54 years	4	2	2	2	1	1	2	1	1	-	-	-	-	-	-
55-59 years	9	7	2	7	5	2	2	2	-	-	-	-	-	-	-
60-64 years	6	6	-	6	6	-	-	-	-	-	-	-	-	-	-
65-69 years	6	3	3	4	1	3	1	1	-	1	1	-	-	-	-
70-74 years	4	2	2	3	2	1	1	-	1	-	-	-	-	-	-
75-79 years	5	1	4	5	1	4	-	-	-	-	-	-	-	-	-
80-84 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
85-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	-	-	-	-	-	-	-	_	_	-	-	-	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Langerhans' cell histiocytosis, not elsewhere classified (D76.0) All ages.... Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 3 2 1 2 2 50-54 years..... 3 1 55-59 years..... 3 2 1 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 2 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hemophagocytic lymphohistiocytosis (D76.1) All ages.... 18 Under 1 year.... 13 11 2 8 1 1 1-4 years..... 3 3 2 5-9 years..... 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hemophagocytic syndrome, infection-associated (D76.2) All ages.... 17 Under 1 year.... 1 1 1 1 1-4 years..... 3 2 2 1 5-9 years..... 1 1 1 10-14 years..... 1 1 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 4 1 60-64 years..... 3 3 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other histiocytosis syndromes (D76.3) All ages..... Under 1 year.... 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Immunodeficiency with predominantly antibody defects (D80) All ages.... 37 17 20 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 2 2 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 50-54 years..... 55-59 years..... 5 1 4 60-64 years..... 3 3 65-69 vears..... 3 70-74 vears..... 3 75-79 years..... 4 80-84 years..... 2 1 2 85-89 years..... 1 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hereditary hypogammaglobulinemia (D80.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Nonfamilial hypogammaglobulinemia (D80.1) All ages..... 13 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 55-59 years..... 4 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Selective deficiency of immunoglobulin A [IgA] (D80.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Selective deficiency of immunoglobulin G [IgG] subclasses (D80.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Selective deficiency of immunoglobulin M [IgM] (D80.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Immunodeficiency with increased immunoglobulin M [IgM] (D80.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Immunodeficiency with predominantly antibody defects, unspecified (D80.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Combined immunodeficiencies (D81) All ages..... 13 Under 1 year.... 6 3 3 4 2 1-4 years..... 4 1 3 3 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases,												As	ian or		
Tenth Revision, Second All races				White		Black			Amei	rican Indi	an 1/	Pacific Islander 2/			
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Adenosine deaminase [ADA] deficie	ency (D81.	3)												
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
10-14 years	_	-	-	_	-	_	-	-	-	-	-	-	-	-	-
l5-19 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-
20-24 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-
25-29 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-
10-44 years	_	-	-	-	-	-	-	_	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
'5-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	-	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Combined immunodeficiency, unspecified (D81.9) All ages.... 12 Under 1 year.... 6 3 3 4 2 1-4 years..... 4 1 3 3 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Immunodeficiency associated with other major defects (D82) All ages.... 10 Under 1 year.... 8 4 6 2 1-4 years..... 4 2 2 3 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Wiskott-Aldrich syndrome (D82.0) All ages..... Under 1 year.... 1-4 years..... 2 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Di George's syndrome (D82.1) All ages..... 9 Under 1 year.... 8 2 1-4 years..... 2 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Immunodeficiency following hereditary defective response to Epstein-Barr virus (D82.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Common variable immunodeficiency (D83) All ages..... 13 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 2 1 2 50-54 years..... 1 1 55-59 years..... 3 1 2 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 2 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Common variable immunodeficiency, unspecified (D83.9) All ages..... 10 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 2 1 50-54 years..... 2 1 1 55-59 years..... 3 2 1 60-64 years..... 65-69 vears..... 70-74 vears..... 2 75-79 years..... 2 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other immunodeficiencies (D84) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 a 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Defects in the complement system (D84.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 2 50-54 years..... 1 55-59 years..... 4 1 3 60-64 years..... 2 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified immunodeficiencies (D84.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Immunodeficiency, unspecified (D84.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Sarcoidosis (D86) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.4 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Sarcoidosis of lung (D86.0) All ages..... 138 36 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 9 40-44 years..... 45-49 years..... 14 4 10 12 3 1.0 50-54 years..... 17 2 15 13 12 55-59 years..... 18 10 8 15 8 60-64 years..... 14 6 8 12 65-69 vears..... 20 2 18 70-74 vears..... 13 12 75-79 years..... 10 80-84 years..... 13 1 12 5 85-89 years..... 1 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Sarcoidosis of lung with sarcoidosis of lymph nodes (D86.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Sarcoidosis of other and combined sites (D86.8) All ages.... 33 16 17 57 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 6 35-39 years..... 10 40-44 years..... 19 13 45-49 years..... 16 9 13 50-54 years..... 14 6 8 55-59 years..... 12 8 4 60-64 years..... 2 1 65-69 vears..... 5 3 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Sarcoidosis, unspecified (D86.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3.0 a 2.7 50-54 years..... R 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other disorders involving the immune mechanism, not elsewhere classified (D89) All ages.... Under 1 year.... 2 2 1 1 1 1 1-4 years..... 1 1 5-9 years..... 10-14 years..... 1 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 9 45-49 years..... 6 3 2 50-54 years..... 14 9 R 6 55-59 years..... 17 10 14 60-64 years..... 13 9 13 65-69 vears..... 8 70-74 vears..... 12 8 10 75-79 years..... 17 17 10 80-84 years..... 18 9 9 16 8 8 1 20 85-89 years..... 8 12 19 11 1 1 90-94 years..... 3 2 2 1 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Polyclonal hypergammaglobulinemia (D89.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cryoglobulinemia (D89.1) All ages..... 17 10 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 1 2 50-54 years..... 1 55-59 years..... 3 2 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 2 75-79 years..... 2 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second All races				White Black						Asian or American Indian 1/ Pacific Islander 2/						
Edition, 2004)	a Both	All races		Both	willte		Both	Black		American Indian I/ Both			Pacific Islander 2/ Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	
Hypergammaglobulinemi	a, unspecifi	ied (D89.2	!)													
All ages	7	2	5	6	2	4	1	_	1	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	_	-	-	-	-	-	_	-	-	_	-	-	
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-	
75-79 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-	
80-84 years	3	-	3	2	-	2	1	-	1	_	_	-	-	-	-	
85-89 years	2	2	-	2	2	-	-	_	-	-	-	-	-	-	-	
90-94 years	1	-	1	1	-	1	-	-	-	_	_	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-	
100 years & over.	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-	
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female Male Female sexes Male sexes Male sexes sexes sexes Other specified disorders involving the immune mechanism, not elsewhere classified (D89.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorder involving the immune mechanism, unspecified (D89.9) All ages.... 45 Under 1 year.... 2 2 1 1 1 1 1-4 years..... 1 1 5-9 years..... 10-14 years..... 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 6 5 1 1 5 50-54 years..... 12 ρ 55-59 years..... 13 9 10 60-64 years..... 11 9 2 11 65-69 vears..... 7 6 70-74 vears..... 10 3 75-79 years..... 14 10 80-84 years..... 14 8 6 13 6 85-89 years..... 18 6 12 17 11 90-94 years..... 2 1 1 1 2 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Iodine-deficiency-related thyroid disorders and allied conditions (E01) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Iodine-deficiency-related (endemic) goiter, unspecified (E01.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other hypothyroidism (E03) All ages.... 1,404 1,116 1,270 1,028 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... Ω 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Congenital hypothyroidism without goiter (E03.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hypothyroidism due to medicaments and other exogenous substances (E03.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Atrophy of thyroid (acquired) (E03.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Myxedema coma (E03.5) All ages.... 22 17 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 2 70-74 vears..... 2 75-79 years..... 80-84 years..... 85-89 years..... 4 3 3 90-94 years..... 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hypothyroidism, unspecified (E03.9) All ages..... 1,378 1,098 1,249 1,011 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other nontoxic goiter (E04) All ages..... 33 30 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 3 70-74 vears..... 6 6 75-79 years..... 80-84 years..... 11 9 8 85-89 years..... 1 90-94 years..... 8 2 1 95-99 years..... 7 2 5 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Nontoxic single thyroid nodule (E04.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Nontoxic multinodular goiter (E04.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 1 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Nontoxic goiter, unspecified (E04.9) All ages..... 30 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 3 3 75-79 years..... 80-84 years..... 85-89 years..... 6 1 90-94 years..... 6 5 2 1 95-99 years..... 3 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Thyrotoxicosis [hyperthyroidism] (E05) All ages.... 222 236 172 19 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 3 25-29 years..... 30-34 years..... 6 3 35-39 years..... 12 40-44 years..... 11 45-49 years..... 16 8 8 50-54 years..... 12 R R 55-59 years..... 15 11 10 1 4 60-64 years..... 20 6 14 19 13 65-69 vears..... 16 12 15 11 70-74 vears..... 17 6 15 6 75-79 years..... 2.7 2.0 19 13 80-84 years..... 43 8 35 38 30 34 25 85-89 years..... 35 9 26 9 1 37 3 90-94 years..... 33 32 28 2 2 4 95-99 years..... 17 16 16 15 1 1 100 years & over. 6 5 5 1 6

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Thyrotoxicosis with diffuse goiter (E05.0) All ages..... 10 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 3 85-89 years..... 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Thyrotoxicosis with toxic multinodular goiter (E05.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Thyroid crisis or storm (E05.5) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 1 1 60-64 years..... 3 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Thyrotoxicosis, unspecified (E05.9) All ages.... 198 209 152 13 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 4 35-39 years..... 12 40-44 years..... 7 45-49 years..... 15 8 50-54 years..... 9 R 2 1 55-59 years..... 13 10 9 3 1 3 60-64 years..... 17 6 11 16 10 65-69 years..... 13 3 10 12 9 70-74 years..... 16 10 6 75-79 years..... 2.4 18 16 6 11 80-84 years..... 37 8 29 32 24 25 33 24 85-89 years..... 34 9 9 1 1 3 90-94 years..... 34 30 29 25 2 2 4 95-99 years..... 17 15 16 16 1 1 100 years & over. 6 5 5 1 6

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Thyroiditis (E06) All ages.... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Autoimmune thyroiditis (E06.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other chronic thyroiditis (E06.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Thyroiditis, unspecified (E06.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other disorders of thyroid (E07) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 2 50-54 years..... 2 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 3 75-79 years..... 80-84 years..... 5 2 2 85-89 years..... 9 6 9 90-94 years..... 10 9 1.0 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified disorders of thyroid (E07.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorder of thyroid, unspecified (E07.9) All ages.... 29 37 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 2 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 3 75-79 years..... 80-84 years..... 5 2 2 85-89 years..... 9 6 9 90-94 years..... 10 9 1.0 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Insulin-dependent diabetes mellitus (E10) All ages.... 5,086 2,510 2,576 4,199 2,097 2,102 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes With coma (E10.0) All ages.... 13 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 50-54 years..... 2 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 3 2 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With ketoacidosis (E10.1) All ages.... 177 101 138 60 37 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 1 1 15-19 years..... 1 3 20-24 years..... 13 8 5 11 25-29 years..... 11 30-34 years..... 13 11 35-39 years..... 17 13 12 40-44 years..... 2.7 17 10 19 13 45-49 years..... 2.4 14 10 20 11 50-54 years..... 13 8 5 9 6 55-59 years..... 15 9 6 12 60-64 years..... 6 3 3 5 65-69 vears..... 9 3 5 4 70-74 vears..... 3 3 75-79 years..... 80-84 years..... 5 3 85-89 years..... 4 3 90-94 years..... 3 1 2 1 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With renal complications (E10.2) All ages.... 135 26 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 12 40-44 years..... 12 11 45-49 years..... 13 10 3 12 1 2 50-54 years..... 14 13 12 11 55-59 years..... 8 3 5 3 4 60-64 years..... 11 8 3 65-69 vears..... 12 8 70-74 vears..... 18 8 10 7 75-79 years..... 2.0 9 11 16 80-84 years..... 13 5 8 11 7 1 1 85-89 years..... 11 5 6 11 6 1 90-94 years..... 2 2 4 3 2 95-99 years..... 3 2 3 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With ophthalmic complications (E10.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With neurological complications (E10.4) All ages.... 27 39 19 20 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 4 2 2 50-54 years..... 2 2 55-59 years..... 8 2 6 4 60-64 years..... 2 2 65-69 vears..... 4 70-74 vears..... 5 3 75-79 years..... 3 80-84 years..... 9 6 3 3 85-89 years..... 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes With peripheral circulatory complications (E10.5) All ages.... 391 197 194 75 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 9 35-39 years..... 40-44 years..... 14 9 5 10 45-49 years..... 16 a 7 11 5 33 22 2.4 9 50-54 years..... 9 11 15 55-59 years..... 36 17 19 29 14 15 4 60-64 years..... 45 24 21 42 23 19 3 65-69 years..... 58 36 22 46 29 17 70-74 years..... 61 34 27 49 27 22 11 75-79 years..... 71 29 42 62 2.8 34 9 80-84 years..... 60 22 38 48 19 29 11 8 25 85-89 years..... 46 2.0 26 42 17 4 3 1 90-94 years..... 16 5 14 5 2 11 9 2 95-99 years..... 5 5 1 4 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With other specified complications (E10.6) All ages.... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 3 45-49 years..... 5 2 50-54 years..... 55-59 years..... 2 1 1 60-64 years..... 3 2 1 65-69 vears..... 70-74 vears..... 75-79 years..... 4 80-84 years..... 6 3 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With multiple complications (E10.7) All ages..... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 7 45-49 years..... 5 2 50-54 years..... 12 5 11 55-59 years..... 15 13 10 11 60-64 years..... 10 8 2 3 65-69 vears..... 14 8 13 6 70-74 vears..... 5 75-79 years..... 10 80-84 years..... 9 2 1 1 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based on the International															
Classification of Dis										ian or					
Tenth Revision, Secon		All rac	es		White			Black			rican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale ————	sexes	Male	Female
With unspecified comp	olications (E	10.8)													
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	_	-	-	_	-	_	-	_	-	-	-	-	-	-	-
1-4 years	_	_	-	_	_	_	_	-	-	_	_	-	-	-	-
5-9 years	_	_	-	_	_	_	_	-	-	_	_	-	-	-	-
10-14 years	_	-	-	_	-	_	-	_	-	-	-	-	-	-	-
l5-19 years	_	-	-	_	-	_	-	_	-	-	-	-	-	-	-
20-24 years	_	-	-	_	-	_	-	_	-	-	-	-	-	-	-
25-29 years	_	-	-	_	-	_	-	-	-	-	-	-	-	-	-
30-34 years	_	-	-	_	-	_	-	-	-	-	-	-	-	-	-
35-39 years	_	_	-	_	-	_	-	-	-	-	-	-	-	-	-
40-44 years	1	1	-	1	1	_	-	-	-	-	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	_	-	-	_	-	_	-	_	-	-	-	-	-	-	-
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	_	-	-	_	-	_	-	_	-	-	-	-	-	-	-
70-74 years	_	_	-	_	_	_	_	-	-	_	_	-	_	-	-
75-79 years	_	-	-	_	-	_	-	_	-	_	-	-	-	-	-
30-84 years	_	-	-	_	-	_	-	_	-	_	-	-	-	-	-
35-89 years	_	-	-	_	-	_	-	_	-	_	-	-	-	-	-
90-94 years	_	_	-	_	_	_	_	-	-	_	_	-	_	-	-
95-99 years	_	-	-	_	-	_	-	_	-	_	-	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based															
on the International															
Classification of Dis	eases,										As	ian or			
Tenth Revision, Second All races			White			Black		Amer	ican In	dian 1/	Pacit	ic Isla	nder 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Without complications	(E10.9)														
All ages	4,054	1,967	2,087	3,353	1,644	1,709	630	286	344	27	12	15	44	25	19
Under 1 year	1	_	1	-	-	-	1	-	1	-	-	-	-	-	-
1-4 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
5-9 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	1	-	1	-	-	-	1	_	1	-	-	-	-	-	-
15-19 years	3	1	2	2	1	1	1	_	1	_	_	_	-	-	-
20-24 years	14	9	5	9	6	3	4	2	2	-	-	-	1	1	-
25-29 years	25	8	17	16	5	11	9	3	6	-	-	-	-	-	-
30-34 years	70	35	35	52	26	26	17	9	8	-	-	-	1	-	1
35-39 years	102	56	46	79	41	38	23	15	8	-	-	-	-	-	-
40-44 years	170	95	75	141	77	64	25	16	9	2	-	2	2	2	-
45-49 years	237	152	85	190	124	66	42	26	16	3	1	2	2	1	1
50-54 years	269	163	106	223	135	88	41	24	17	1	-	1	4	4	-
55-59 years	346	183	163	272	144	128	64	33	31	4	1	3	6	5	1
60-64 years	373	207	166	281	165	116	86	38	48	2	2	-	4	2	2
65-69 years	353	198	155	288	167	121	57	28	29	5	2	3	3	1	2
70-74 years	467	233	234	392	196	196	66	31	35	3	3	-	6	3	3
75-79 years	488	218	270	417	191	226	65	24	41	2	-	2	4	3	1
80-84 years	515	202	313	442	179	263	65	18	47	4	3	1	4	2	2
85-89 years	394	149	245	347	131	216	44	17	27	-	-	-	3	1	2
90-94 years	172	48	124	155	46	109	16	2	14	-	-	-	1	_	1
95-99 years	44	8	36	38	8	30	3	-	3	1	-	1	2	_	2
100 years & over.	9	1	8	8	1	7	_	_	_	_	_	_	1	_	1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

Cause of death (based	ì														
on the International															
Classification of Dis								-1 1		_			sian or		1 0/
Tenth Revision, Secon		All races White Black America Both Both Both Both		ncan In	dian 1/										
Edition, 2004)		M-1-	E1-		Mala	E-m-l-		M-1-	Esmal s		Mala	E-mala	Both	M-1-	Esmal s
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	мате	Female	sexes	мате	Female
Noninsulin-dependent	diabetes me	llitus (E1	.1)												
All ages	16,492	7,948	8,544	13,762	6,725	7,037	2,165	952	1,213	231	101	130	334	170	164
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-
20-24 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
25-29 years	6	2	4	5	2	3	1	-	1	-	-	-	-	-	-
30-34 years	22	13	9	9	6	3	7	4	3	5	2	3	1	1	-
35-39 years	65	37	28	43	23	20	19	11	8	3	3	-	-	-	-
40-44 years	127	82	45	97	65	32	24	14	10	4	3	1	2	-	2
45-49 years	263	164	99	183	119	64	62	30	32	10	8	2	8	7	1
50-54 years	589	374	215	444	290	154	114	66	48	19	10	9	12	8	4
55-59 years	891	531	360	685	409	276	165	96	69	15	11	4	26	15	11
60-64 years	1,242	740	502	977	602	375	202	108	94	34	12	22	29	18	11
65-69 years	1,567	868	699	1,242	704	538	262	130	132	28	14	14	35	20	15
70-74 years	1,908	983	925	1,571	825	746	271	126	145	31	14	17	35	18	17
75-79 years	2,566	1,291	1,275	2,146	1,113	1,033	340	144	196	24	9	15	56	25	31
80-84 years	3,010	1,365	1,645	2,622	1,219	1,403	311	114	197	30	7	23	47	25	22
85-89 years	2,493	995	1,498	2,194	900	1,294	227	66	161	20	6	14	52	23	29
90-94 years	1,294	404	890	1,147	357	790	119	36	83	5	1	4	23	10	13
95-99 years	380	85	295	337	78	259	33	6	27	3	1	2	7	-	7
100 years & over.	67	13	54	59	12	47	7	1	6	_	_	_	1	_	1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes With coma (E11.0) All ages.... 32 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 3 3 2 6 50-54 years..... 2 55-59 years..... 5 2 3 60-64 years..... 3 2 1 65-69 vears..... 3 70-74 vears..... 9 3 6 75-79 years..... 11 80-84 years..... 10 6 8 7 2 85-89 years..... 6 90-94 years..... 2 2 95-99 years..... 2 1 2 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

n the International Lassification of Diseases, enth Revision, Second All races White Black American Indian 1/ Pacific Islander 2															
Tenth Revision, Secon		All races			White			Black			ican Indi	an 1/		ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male 	Female
With ketoacidosis (El	.1.1)														
All ages	104	44	60	76	31	45	22	10	12	4	2	2	2	1	1
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	2	-	2	1	-	1	1	-	1	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	4	3	1	2	2	-	2	1	1	-	-	-	-	-	-
10-44 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
15-49 years	7	2	5	3	1	2	4	1	3	-	-	-	-	-	-
0-54 years	11	6	5	6	3	3	3	2	1	2	1	1	-	-	-
55-59 years	9	4	5	8	3	5	1	1	-	-	-	-	-	-	-
0-64 years	9	6	3	8	6	2	1	-	1	-	-	-	-	-	-
55-69 years	18	9	9	11	4	7	6	5	1	-	-	-	1	-	1
70-74 years	3	1	2	2	1	1	1	-	1	-	-	-	-	-	-
5-79 years	8	2	6	5	1	4	2	-	2	1	1	-	-	-	-
0-84 years	13	4	9	11	4	7	1	-	1	1	-	1	-	-	-
5-89 years	5	2	3	5	2	3	-	-	-	-	-	-	-	-	-
00-94 years	10	3	7	9	2	7	-	_	-	-	-	-	1	1	-
95-99 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
100 years & over.	1	-	1	1	_	1	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With renal complications (E11.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1.0 a 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes With ophthalmic complications (E11.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With neurological complications (E11.4) All ages..... 111 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 4 1 2 8 50-54 years..... 5 3 1 55-59 years..... 10 9 1 1 60-64 years..... 15 12 3 65-69 vears..... 12 6 6 70-74 vears..... 17 5 12 11 75-79 years..... 2.0 10 10 19 9 80-84 years..... 19 8 11 19 8 11 85-89 years..... 11 3 8 9 3 6 9 1 90-94 years..... 8 1 1 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes With peripheral circulatory complications (E11.5) All ages.... 1,290 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... Ω 50-54 years..... 2.2 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.4 80-84 years..... 85-89 years..... 2.4 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With other specified complications (E11.6) All ages.... 36 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 50-54 years..... 1 1 55-59 years..... 3 1 2 60-64 years..... 2 1 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 9 80-84 years..... 11 1 9 85-89 years..... 90-94 years..... 6 95-99 years..... 4 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With multiple complications (E11.7) All ages..... 105 158 19 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 2 2 9 50-54 years..... 3 6 3 55-59 years..... 17 12 15 10 5 60-64 years..... 10 8 2 9 65-69 vears..... 28 17 21 12 9 70-74 vears..... 23 11 12 18 10 75-79 years..... 2.7 14 13 2.0 9 3 80-84 years..... 33 15 18 29 15 14 19 17 2 85-89 years..... 10 9 9 8 1 90-94 years..... 6 2 1 1 4 5 2 3 95-99 years..... 3 2 1 3 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes With unspecified complications (E11.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based	۹														
on the International															
Classification of Diseases,									Asian or						
Tenth Revision, Second		All races		White		Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both	M - 1 -		Both	M - 1 -	m1	Both		_ ,	Both	M . 1 .	F1-	Both	24 . 7 .	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Without complications	s (E11.9)														
All ages	13,728	6,535	7,193	11,505	5,558	5,947	1,764	759	1,005	183	81	102	276	137	139
Under 1 year	-	-	-	-	-	-	_	-	-	-	-	-	_	-	-
1-4 years	-	-	-	-	-	-	_	-	-	-	-	-	_	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
25-29 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
30-34 years	20	12	8	8	5	3	6	4	2	5	2	3	1	1	-
35-39 years	49	27	22	35	19	16	12	6	6	2	2	-	-	-	-
40-44 years	101	65	36	78	51	27	18	11	7	3	3	-	2	-	2
45-49 years	207	129	78	147	95	52	45	22	23	9	7	2	6	5	1
50-54 years	480	305	175	366	242	124	89	47	42	14	8	6	11	8	3
55-59 years	723	438	285	553	335	218	140	82	58	11	9	2	19	12	7
60-64 years	1,026	601	425	811	489	322	160	86	74	31	11	20	24	15	9
65-69 years	1,283	706	577	1,017	579	438	215	99	116	21	10	11	30	18	12
70-74 years	1,568	800	768	1,295	676	619	221	98	123	24	11	13	28	15	13
75-79 years	2,137	1,068	1,069	1,795	927	868	277	115	162	18	6	12	47	20	27
80-84 years	2,518	1,147	1,371	2,204	1,023	1,181	256	100	156	20	5	15	38	19	19
85-89 years	2,104	813	1,291	1,860	741	1,119	185	52	133	18	5	13	41	15	26
90-94 years	1,110	336	774	984	296	688	101	30	71	4	1	3	21	9	12
95-99 years	333	72	261	291	65	226	32	6	26	3	1	2	7	-	7
100 years & over.	63	13	50	56	12	44	6	1	5	-	-	-	1	-	1
Age not stated	1	1	-	1	1	-	_	-	-	-	-	_	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Malnutrition-related diabetes mellitus (E12) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes With multiple complications (E12.7) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other specified diabetes mellitus (E13) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes With renal complications (E13.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With peripheral circulatory complications (E13.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With other specified complications (E13.6) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes With unspecified complications (E13.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Without complications (E13.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based																	
on the International																	
Classification of Diseases,										Asian or							
Tenth Revision, Second All races			White			Black			American Indian 1/			Pacific Islander 2/					
Edition, 2004)	Both			Both			Both			Both			Both				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female		
Unspecified diabetes	mellitus (E	:14)															
All ages	50,860	25,545	25,315	39,235	20,236	18,999	9,851	4,447	5,404	544	249	295	1,230	613	617		
Under 1 year	5	4	1	5	4	1	-	-	-	_	-	-	_	-	-		
1-4 years	6	2	4	4	2	2	2	-	2	_	-	-	-	-	-		
5-9 years	6	4	2	5	3	2	1	1	-	_	-	-	-	-	-		
10-14 years	19	8	11	12	7	5	7	1	6	_	-	-	-	-	-		
15-19 years	34	19	15	20	11	9	13	7	6	-	-	-	1	1	-		
20-24 years	93	53	40	58	30	28	33	21	12	1	1	-	1	1	-		
25-29 years	196	117	79	107	66	41	82	46	36	2	1	1	5	4	1		
30-34 years	298	179	119	177	103	74	109	69	40	7	3	4	5	4	1		
35-39 years	488	300	188	308	191	117	167	99	68	7	5	2	6	5	1		
40-44 years	1,023	638	385	681	439	242	304	179	125	19	8	11	19	12	7		
45-49 years	1,607	1,018	589	1,134	739	395	422	249	173	21	15	6	30	15	15		
50-54 years	2,576	1,546	1,030	1,763	1,098	665	709	390	319	49	22	27	55	36	19		
55-59 years	3,912	2,290	1,622	2,822	1,699	1,123	942	490	452	62	40	22	86	61	25		
60-64 years	4,499	2,625	1,874	3,308	1,994	1,314	1,014	528	486	70	37	33	107	66	41		
65-69 years	5,065	2,781	2,284	3,741	2,154	1,587	1,129	529	600	55	27	28	140	71	69		
70-74 years	5,929	3,171	2,758	4,523	2,522	2,001	1,190	552	638	68	22	46	148	75	73		
75-79 years	7,299	3,665	3,634	5,834	3,057	2,777	1,208	492	716	62	25	37	195	91	104		
80-84 years	7,661	3,454	4,207	6,250	2,946	3,304	1,161	409	752	61	22	39	189	77	112		
85-89 years	6,035	2,447	3,588	5,045	2,115	2,930	804	256	548	33	12	21	153	64	89		
90-94 years	3,036	988	2,048	2,539	859	1,680	414	100	314	23	9	14	60	20	40		
95-99 years	919	217	702	778	184	594	114	26	88	3	_	3	24	7	17		
100 years & over.	154	19	135	121	13	108	26	3	23	1	_	1	6	3	3		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With coma (E14.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.2 R 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes With ketoacidosis (E14.1) All ages.... 1,637 1,116 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.8 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1.0 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes With renal complications (E14.2) All ages.... 1,646 1,244 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 2.0 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes With ophthalmic complications (E14.3) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 2 60-64 years..... 65-69 vears..... 4 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 1 85-89 years..... 90-94 years..... 3 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With neurological complications (E14.4) All ages.... 104 196 106 90 28 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 12 8 10 2 2.0 6 50-54 years..... 9 14 11 55-59 years..... 19 12 7 13 3 60-64 years..... 15 10 5 13 65-69 vears..... 22 11 19 70-74 vears..... 26 13 13 19 9 10 75-79 years..... 38 2.2 16 36 15 80-84 years..... 38 21 17 36 21 15 1 18 2 85-89 years..... 9 9 16 8 8 1 1 90-94 years..... 8 3 5 3 95-99 years..... 2 2 4 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Female Male Female Male Female sexes Male Female sexes sexes Male sexes sexes With peripheral circulatory complications (E14.5) All ages.... 2,316 2,139 3,369 1,833 1,536 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.6 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1.0 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes With other specified complications (E14.6) All ages.... 110 141 29 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 2 7 50-54 years..... 5 1 55-59 years..... 10 3 9 60-64 years..... 15 6 9 14 65-69 vears..... 17 5 12 13 3 70-74 vears..... 16 8 11 75-79 years..... 32 9 2.3 2.7 19 80-84 years..... 33 17 16 28 15 13 3 15 2 85-89 years..... 5 10 12 3 9 90-94 years..... 16 2 14 13 11 2 2 1 2 95-99 years..... 2 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes With multiple complications (E14.7) All ages.... 229 180 303 179 124 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 4 40-44 years..... 6 3 3 45-49 years..... 16 1.0 6 15 1 0 1 1 9 50-54 years..... 2.4 15 14 9 5 1 0 55-59 years..... 43 25 18 31 18 13 10 5 5 60-64 years..... 49 30 19 36 22 14 13 65-69 years..... 42 26 16 29 17 12 11 9 70-74 years..... 57 33 41 27 11 75-79 years..... 50 2.8 2.2 34 2.3 11 14 9 80-84 years..... 53 2.7 26 39 2.2 17 11 8 2 31 17 85-89 years..... 38 19 19 14 6 5 90-94 years..... 18 9 9 16 2 9 2 95-99 years..... 6 2 6 2 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes With unspecified complications (E14.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based																	
on the International																	
Classification of Diseases,										Asian or							
Tenth Revision, Second		All races			White		Black			American Indian 1/				fic Isla	inder 2/		
Edition, 2004)	Both			Both			Both			Both			Both				
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female		
Without complications	(E14.9)																
All ages	41,992	20,898	21,094	32,641	16,705	15,936	7,869	3,463	4,406	444	208	236	1,038	522	516		
Under 1 year	3	2	1	3	2	1	_	-	-	-	-	-	-	-	-		
1-4 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
5-9 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
10-14 years	6	3	3	5	3	2	1	-	1	-	-	-	-	-	-		
15-19 years	14	8	6	9	4	5	4	3	1	-	-	-	1	1	-		
20-24 years	37	19	18	18	9	9	18	9	9	1	1	-	_	-	-		
25-29 years	108	58	50	64	37	27	38	17	21	2	1	1	4	3	1		
30-34 years	178	105	73	99	57	42	70	43	27	6	3	3	3	2	1		
35-39 years	320	193	127	205	126	79	107	60	47	5	4	1	3	3	-		
40-44 years	726	442	284	491	314	177	210	115	95	13	5	8	12	8	4		
15-49 years	1,236	789	447	852	566	286	342	199	143	17	12	5	25	12	13		
50-54 years	2,078	1,257	821	1,428	893	535	562	313	249	40	19	21	48	32	16		
55-59 years	3,214	1,880	1,334	2,325	1,403	922	765	391	374	55	35	20	69	51	18		
60-64 years	3,724	2,178	1,546	2,754	1,669	1,085	821	421	400	60	35	25	89	53	36		
65-69 years	4,230	2,336	1,894	3,144	1,834	1,310	927	422	505	40	20	20	119	60	59		
70-74 years	4,913	2,618	2,295	3,771	2,091	1,680	968	443	525	48	17	31	126	67	59		
75-79 years	6,035	3,029	3,006	4,845	2,539	2,306	969	383	586	53	23	30	168	84	84		
80-84 years	6,482	2,889	3,593	5,306	2,471	2,835	958	333	625	52	16	36	166	69	97		
85-89 years	5,148	2,047	3,101	4,337	1,780	2,557	653	204	449	29	10	19	129	53	76		
90-94 years	2,609	838	1,771	2,202	735	1,467	338	80	258	19	7	12	50	16	34		
95-99 years	793	188	605	673	158	515	95	24	71	3	-	3	22	6	16		
100 years & over.	136	17	119	108	12	96	23	3	20	1	_	1	4	2	2		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Nondiabetic hypoglycemic coma (E15) All ages..... 18 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 2 50-54 years..... 1 55-59 years..... 2 1 1 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 5 2 1 85-89 years..... 90-94 years..... 2 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other disorders of pancreatic internal secretion (E16) All ages.... 170 92 87 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 4 7 40-44 years..... 10 45-49 years..... 16 6 10 11 5 50-54 years..... 2.4 16 8 14 1 0 55-59 years..... 19 7 12 12 6 3 3 4 60-64 years..... 27 14 13 20 10 10 65-69 vears..... 18 12 9 6 5 70-74 vears..... 25 13 12 13 6 75-79 years..... 30 19 19 11 11 80-84 years..... 34 19 15 2.7 13 14 6 1 35 29 85-89 years..... 15 20 14 15 6 5 7 90-94 years..... 3 3 1 6 4 1 3 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Drug-induced hypoglycemia without coma (E16.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other hypoglycemia (E16.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hypoglycemia, unspecified (E16.2) All ages.... 114 140 164 90 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 4 35-39 years..... 4 7 40-44 years..... 10 45-49 years..... 16 6 10 11 5 23 50-54 years..... 15 8 13 1 0 55-59 years..... 18 7 11 11 6 3 3 60-64 years..... 26 14 12 19 65-69 vears..... 16 12 70-74 vears..... 25 13 12 13 6 75-79 years..... 30 19 19 11 11 80-84 years..... 33 18 15 2.7 13 14 5 1 32 27 85-89 years..... 14 18 13 14 7 90-94 years..... 3 3 1 6 1 3 95-99 years..... 2 2 2 2 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Abnormal secretion of gastrin (E16.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorder of pancreatic internal secretion, unspecified (E16.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Hypoparathyroidism (E20) All ages..... 10 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other hypoparathyroidism (E20.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Hypoparathyroidism, unspecified (E20.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hyperparathyroidism and other disorders of parathyroid gland (E21) All ages.... 19 57 23 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 50-54 years..... 3 3 2 1 55-59 years..... 3 3 1 60-64 years..... 10 6 65-69 vears..... 3 70-74 vears..... 5 5 75-79 years..... 16 9 12 13 80-84 years..... 25 8 17 2.0 15 1 1 13 10 2 85-89 years..... 3 10 8 1 90-94 years..... 13 12 4 a 95-99 years..... 7 1 1 6 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Primary hyperparathyroidism (E21.0) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 2 85-89 years..... 90-94 years..... 3 4 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Secondary hyperparathyroidism, not elsewhere classified (E21.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hyperparathyroidism, unspecified (E21.3) All ages.... 46 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 3 50-54 years..... 3 1 55-59 years..... 2 2 60-64 years..... 3 65-69 vears..... 3 70-74 vears..... 5 75-79 years..... 13 3 10 7 80-84 years..... 21 6 15 16 13 1 1 11 2 85-89 years..... 3 8 8 6 1 90-94 years..... 8 2 2 6 95-99 years..... 6 5 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorder of parathyroid gland, unspecified (E21.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hyperfunction of pituitary gland (E22) All ages..... 46 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 2 45-49 years..... 2 3 50-54 years..... 3 2 55-59 years..... 2 1 1 60-64 years..... 2 2 65-69 vears..... 4 3 70-74 vears..... 5 5 5 75-79 years..... 11 80-84 years..... 13 2 11 13 11 16 11 85-89 years..... 11 16 90-94 years..... 8 2 6 6 8 95-99 years..... 5 5 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acromegaly and pituitary gigantism (E22.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 2 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Syndrome of inappropriate secretion of antidiuretic hormone (E22.2) All ages..... 43 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 2 50-54 years..... 2 55-59 years..... 60-64 years..... 1 65-69 vears..... 3 70-74 vears..... 5 5 5 75-79 years..... 11 80-84 years..... 10 9 1.0 9 15 85-89 years..... 11 15 11 90-94 years..... 8 2 6 6 8 95-99 years..... 5 5 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hyperfunction of pituitary gland, unspecified (E22.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hypofunction and other disorders of the pituitary gland (E23) All ages.... 120 30 Under 1 year.... 4 3 3 2 1 3 2 1 1 1-4 years..... 1 1 3 2 1 5-9 years..... 10-14 years..... 1 1 15-19 years..... 3 3 20-24 years..... 4 3 25-29 years..... 1 1 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 7 4 3 3 50-54 years..... 1.0 6 2 55-59 years..... 8 5 3 6 5 1 60-64 years..... 15 2 13 12 10 3 3 65-69 vears..... 14 12 5 5 70-74 vears..... 9 5 9 75-79 years..... 17 14 80-84 years..... 21 11 10 16 8 5 3 2 10 85-89 years..... 5 5 9 1 90-94 years..... 5 3 2 1 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hypopituitarism (E23.0) All ages.... 57 37 47 17 Under 1 year..... 4 2 2 2 1 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 2 2 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 2 45-49 years..... 3 1 2 7 50-54 years..... 5 2 2 55-59 years..... 5 2 5 60-64 years..... 9 2 65-69 vears..... 8 5 3 70-74 vears..... 6 3 3 75-79 years..... 11 3 80-84 years..... 15 9 6 12 3 85-89 years..... 6 1 6 90-94 years..... 3 2 1 3 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Diabetes insipidus (E23.2) All ages.... 19 Under 1 year.... 3 2 1 1 1 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 1 3 50-54 years..... 2 1 55-59 years..... 1 1 60-64 years..... 6 6 65-69 vears..... 6 70-74 vears..... 2 75-79 years..... 80-84 years..... 4 3 1 2 2 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or															
Tenth Revision, Second All races				White		Black			Amer	rican Ind		Pacific Islander 2/			
Edition, 2004)	Both			Both						Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Hypothalamic dysfunct	ion, not els	sewhere cl	assified (E	23.3)											
All ages	5	1	4	4	1	3	1	_	1	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	2	1	1	2	1	1	-	-	-	_	_	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	_
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
85-89 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
90-94 years	1	-	1	1	-	1	-	-	-	_	_	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other disorders of pituitary gland (E23.6) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorder of pituitary gland, unspecified (E23.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 2 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cushing's syndrome (E24) All ages..... 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 5 3 65-69 vears..... 5 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Pituitary-dependent Cushing's disease (E24.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Drug-induced Cushing's syndrome (E24.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other Cushing's syndrome (E24.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cushing's syndrome, unspecified (E24.9) All ages..... 17 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 1 55-59 years..... 2 2 60-64 years..... 5 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Adrenogenital disorders (E25) All ages.... Under 1 year.... 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Congenital adrenogenital disorders associated with enzyme deficiency (E25.0) All ages.... Under 1 year.... 2 2 2 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Hyperaldosteronism (E26) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Primary hyperaldosteronism (E26.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other hyperaldosteronism (E26.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Hyperaldosteronism, unspecified (E26.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other disorders of adrenal gland (E27) All ages.... 149 215 140 Under 1 year.... 2 2 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 1 25-29 years..... 30-34 years..... 35-39 years..... 9 40-44 years..... 45-49 years..... 10 3 Ω 50-54 years..... 13 6 11 55-59 years..... 17 5 15 11 12 60-64 years..... 18 12 6 14 6 65-69 vears..... 14 2 12 14 12 70-74 vears..... 18 9 8 75-79 years..... 2.4 10 22 9 13 14 80-84 years..... 47 13 34 44 11 33 2 1 37 25 85-89 years..... 12 25 36 11 9 90-94 years..... 3 1 1 6 8 3 5 95-99 years..... 2 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International															
Classification of Dise	eases,						As	ian or							
Tenth Revision, Second All races			es		White		Black			American Indian 1/			Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female	sexes	Male	Female
Primary adrenocortica	l insufficie	ncy (E27.	1)												
All ages	112	43	69	106	42	64	3	1	2	3	_	3	_	-	_
Under 1 year	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	_	-	-	-	-	-	-	_	-	-	-	-	-	-
15-19 years	2	2	-	2	2	-	-	-	_	-	-	-	-	-	-
20-24 years	4	1	3	4	1	3	-	-	_	-	-	-	-	-	-
25-29 years	-	_	-	-	-	-	-	-	_	-	-	-	-	-	-
30-34 years	3	2	1	3	2	1	-	-	_	-	-	-	-	-	-
35-39 years	5	4	1	5	4	1	_	-	_	-	-	-	_	-	-
40-44 years	6	1	5	5	1	4	_	_	_	1	_	1	_	_	_
45-49 years	4	-	4	3	-	3	1	-	1	-	-	-	_	_	-
50-54 years	9	3	6	8	3	5	_	-	_	1	-	1	_	_	-
55-59 years	6	2	4	6	2	4	_	-	_	_	_	_	_	_	-
0-64 years	3	3	-	3	3	-	_	-	-	-	-	-	-	-	-
65-69 years	5	2	3	5	2	3	_	-	_	_	_	_	_	_	-
70-74 years	6	4	2	5	3	2	1	1	_	_	_	_	_	_	_
75-79 years	14	6	8	13	6	7	_	_	_	1	_	1	_	_	_
30-84 years	21	4	17	20	4	16	1	_	1	_	_	_	_	_	_
35-89 years	19	6	13	19	6	13	_	_	_	_	_	_	_	_	_
90-94 years	4	2	2	4	2	2	_	_	_	_	_	_	_	_	_
95-99 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Addisonian crisis (E27.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Drug-induced adrenocortical insufficiency (E27.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other and unspecified adrenocortical insufficiency (E27.4) All ages.... Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 5 3 2 3 50-54 years..... 3 55-59 years..... 10 3 60-64 years..... 10 6 3 65-69 vears..... 6 6 70-74 vears..... 8 5 75-79 years..... 7 5 80-84 years..... 19 11 17 11 15 10 85-89 years..... 5 10 14 90-94 years..... 5 3 1 1 4 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified disorders of adrenal gland (E27.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases,												Asian or						
Tenth Revision, Secon		All races		_	White			Black			rican Indi	an 1/	Pacific Islander 2					
Edition, 2004)	Both	1		Both			Both	1	_ ,	Both			Both		_ 1			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female			
Disorder of adrenal g	land, unspec	cified (E2	7.9)															
All ages	20	9	11	17	6	11	3	3	_	_	_	_	_	_	_			
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-			
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
60-64 years	5	3	2	4	2	2	1	1	-	-	-	-	-	-	-			
65-69 years	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-			
70-74 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-			
75-79 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-			
80-84 years	5	1	4	5	1	4	_	-	-	-	_	-	-	-	-			
85-89 years	2	1	1	2	1	1	_	-	-	-	_	-	-	-	-			
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_			

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Ovarian dysfunction (E28) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... . . . 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Polycystic ovarian syndrome (E28.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... . . . 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Bot.h Bot.h and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Primary ovarian failure (E28.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... . . . 25-29 years..... . . . 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Testicular dysfunction (E29) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... . . . 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... . . . 85-89 years..... . . . 90-94 years..... 95-99 years..... 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Testicular hypofunction (E29.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... . . . 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... . . . 55-59 years..... 60-64 years..... 65-69 years..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... . . . 90-94 years..... 95-99 years..... 100 years & over. . . .

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of puberty, not elsewhere classified (E30) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorder of puberty, unspecified (E30.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Polyglandular dysfunction (E31) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Autoimmune polyglandular failure (E31.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Polyglandular dysfunction, unspecified (E31.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Diseases of thymus (E32) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Persistent hyperplasia of thymus (E32.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other diseases of thymus (E32.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Disease of thymus, unspecified (E32.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other endocrine disorders (E34) All ages.... 39 36 Under 1 year.... 11 3 3 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 50-54 years..... 2 1 55-59 years..... 8 3 5 60-64 years..... 1 1 65-69 vears..... 10 8 70-74 vears..... 18 10 75-79 years..... 13 8 13 8 80-84 years..... 11 6 5 9 1 5 85-89 years..... 90-94 years..... 3 2 2 95-99 years..... 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Carcinoid syndrome (E34.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 50-54 years..... 2 1 55-59 years..... 2 5 6 60-64 years..... 1 1 65-69 vears..... 10 70-74 vears..... 18 10 75-79 years..... 12 7 12 7 80-84 years..... 10 8 3 1 85-89 years..... 4 3 90-94 years..... 3 2 2 95-99 years..... 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Short stature, not elsewhere classified (E34.3) All ages.... Under 1 year.... 11 3 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified endocrine disorders (E34.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Endocrine disorder, unspecified (E34.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis-	eases,											As	sian or		
Tenth Revision, Second All races		ces	White			Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Kwashiorkor (E40)															
All ages	7	2	5	7	2	5	-	-	-	-	-	-	_	-	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
55-59 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Nutritional marasmus (E41) All ages..... 19 30 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 1 1 55-59 years..... 4 2 2 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 5 80-84 years..... 14 12 2 10 85-89 years..... 11 9 90-94 years..... 5 2 3 1 4 2 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Unspecified severe protein-energy malnutrition (E43) All ages.... 255 97 158 59 Under 1 year.... 1 1 1 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 1 40-44 years..... 4 45-49 years..... 1.0 5 5 6 2 50-54 years..... 14 6 8 3 11 55-59 years..... 12 7 8 4 5 2 6 60-64 years..... 17 4 13 12 10 65-69 vears..... 15 5 12 5 3 10 70-74 vears..... 39 16 23 26 12 14 75-79 years..... 47 25 2.2 40 19 21 6 80-84 years..... 54 25 29 45 2.0 25 55 17 85-89 years..... 64 20 44 38 6 5 90-94 years..... 29 8 21 23 18 5 1 5 3 95-99 years..... 6 2 4 6 2 4 100 years & over. 2 2 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Protein-energy malnutrition of moderate and mild degree (E44) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis Tenth Revision, Secor		Asian or													
	Both	All races		Both	White		Dath	Black		American Indian 1/			Pacific Islander 2/		
Edition, 2004) and age	sexes	Male	Female	sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes	Male	Female
Moderate protein-ener	gy malnutrit	ion (E44.0	0)												
All ages	5	2	3	5	2	3	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-19 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-29 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-34 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
35-39 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-44 years	-	-	-	-	-	-	-	-	-	-	-	-	_	-	-
15-49 years	_	-	-	-	_	-	-	-	-	-	-	_	-	-	-
50-54 years	_	-	-	-	_	-	-	-	-	-	-	_	-	-	-
5-59 years	_	-	-	-	_	-	-	-	-	-	-	_	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-69 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
0-74 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
5-79 years	2	2	-	2	2	-	-	-	-	_	_	-	-	-	-
0-84 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
5-89 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
0-94 years	1	-	1	1	-	1	-	-	-	_	_	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	_	_	-	-	-	-
100 years & over.	2	_	2	2	_	2	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Retarded development following protein-energy malnutrition (E45) All ages.... Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 5 2 85-89 years..... 12 9 90-94 years..... 6 2 4 95-99 years..... 6 5 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Unspecified protein-energy malnutrition (E46) All ages.... 1,951 1,269 1,658 1,101 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.8 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Thiamin deficiency (E51) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 2 1 60-64 years..... 4 2 65-69 vears..... 70-74 vears..... 3 3 75-79 years..... 80-84 years..... 2 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based																
on the International																
Classification of Diseases,										Asian or						
Tenth Revision, Secon		All races		White			Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both	_	_	Both	_	_	Both		_	Both			Both		_	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale ————	sexes	Male	Female	
Wernicke's encephalop	athy (E51.2)															
All ages	15	11	4	13	11	2	2	_	2	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	_	-	-	_	_	_	_	-	-	_	_	-	_	_	-	
5-9 years	_	-	-	_	_	_	_	-	-	_	_	-	_	_	-	
10-14 years	-	-	-	-	_	_	-	-	-	-	-	_	-	-	-	
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
15-49 years	-	-		-	_	-	-	-	-	-	-	-	-	-	-	
0-54 years	2	2	-	2	2	-	-	-	-	-	-	-	-	-	-	
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
0-64 years	4	2	2	3	2	1	1	-	1	-	-	-	-	-	-	
55-69 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
70-74 years	3	3	-	3	3	-	-	-	-	-	-	-	-	-	-	
'5-79 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-	
0-84 years	2	1	1	2	1	1	-	-	-	-	-	_	-	-	-	
35-89 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	-	_	-	-	-	_	-	-	-	-	-	-	-	-	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Thiamin deficiency, unspecified (E51.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Deficiency of other B group vitamins (E53) All ages.... 38 Under 1 year.... 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 6 3 6 85-89 years..... 15 11 14 10 90-94 years..... 14 3 11 14 11 95-99 years..... 10 2 8 9 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Deficiency of other specified B group vitamins (E53.8) All ages..... 38 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 6 3 6 85-89 years..... 15 11 14 10 90-94 years..... 14 3 11 14 11 95-99 years..... 10 2 8 9 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Vitamin B deficiency, unspecified (E53.9) All ages.... 1 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Vitamin D deficiency (E55) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Rickets, active (E55.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other vitamin deficiencies (E56) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Vitamin deficiency, unspecified (E56.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Deficiency of other nutrient elements (E61) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Iron deficiency (E61.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other nutritional deficiencies (E63) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 5 60-64 years..... 2 3 65-69 vears..... 2 70-74 vears..... 5 3 75-79 years..... 11 6 5 80-84 years..... 19 10 9 17 9 2 20 85-89 years..... 4 16 19 15 90-94 years..... 18 5 13 17 12 1 1 95-99 years..... 10 10 9 9 1 1 100 years & over. 5 5 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Nutritional deficiency, unspecified (E63.9) All ages.... 30 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 5 60-64 years..... 2 3 65-69 vears..... 2 70-74 vears..... 5 3 75-79 years..... 11 6 5 80-84 years..... 19 10 9 17 9 2 20 85-89 years..... 4 16 19 15 90-94 years..... 18 5 13 17 12 1 1 95-99 years..... 10 10 9 9 1 1 100 years & over. 5 5 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Obesity (E66) All ages.... 4,746 2,231 2,515 3,787 1,840 1,947 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 vears..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.8 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Obesity due to excess calories (E66.0) All ages..... 13 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 3 45-49 years..... 6 5 1 5 50-54 years..... 2 3 55-59 years..... 6 2 60-64 years..... 5 3 2 65-69 vears..... 3 3 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Extreme obesity with alveolar hypoventilation (E66.2) All ages.... 114 13 Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 8 40-44 years..... 45-49 years..... 13 8 5 11 3 50-54 years..... 12 R 9 55-59 years..... 20 8 12 17 11 60-64 years..... 21 11 10 19 10 65-69 vears..... 11 7 11 70-74 vears..... 13 6 13 75-79 years..... 10 10 80-84 years..... 10 6 2 85-89 years..... 6 2 6 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Other obesity (E66.8) All ages.... 2,900 1,374 1,526 2,297 1,129 1,168 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1.0 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 8.5 2.8 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Obesity, unspecified (E66.9) All ages.... 1,678 1,345 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of aromatic amino-acid metabolism (E70) All ages.... Under 1 year.... 2 1 1 1 1-4 years..... 2 1 5-9 years..... 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 2 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other hyperphenylalaninemias (E70.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 2 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of tyrosine metabolism (E70.2) All ages.... Under 1 year.... 2 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Albinism (E70.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Female Male Female Male Female Male Female sexes Male sexes Male sexes sexes sexes Disorders of branched-chain amino-acid metabolism and fatty-acid metabolism (E71) All ages.... Under 1 year.... 14 8 6 11 4 3 1-4 years..... 10 3 6 1 3 2 5-9 years..... 1 1 10-14 years..... 5 5 15-19 years..... 3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 2 2 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other disorders of branched-chain amino-acid metabolism (E71.1) All ages.... Under 1 year.... 3 1 1-4 years..... 2 1 5-9 years..... 1 10-14 years..... 1 1 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of fatty-acid metabolism (E71.3) All ages.... 13 30 Under 1 year.... 11 6 5 8 5 3 3 1-4 years..... 8 3 5 5 3 1 5-9 years..... 1 10-14 years..... 4 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 1 55-59 years..... 2 2 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Other disorders of amino-acid metabolism (E72) All ages.... 37 31 12 Under 1 year.... 11 4 9 5 4 2 2 1-4 years..... 4 3 2 1 5-9 years..... 1 10-14 years..... 3 15-19 years..... 3 1 20-24 years..... 25-29 years..... 30-34 years..... 4 35-39 years..... 40-44 years..... 5 2 45-49 years..... 2 6 50-54 years..... 5 1 55-59 years..... 4 3 1 60-64 years..... 3 1 2 65-69 vears..... 5 3 70-74 vears..... 3 3 75-79 years..... 80-84 years..... 2 2 85-89 years..... 3 90-94 years..... 2 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of amino-acid transport (E72.0) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of sulphur-bearing amino-acid metabolism (E72.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 1 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of urea cycle metabolism (E72.2) All ages..... 23 35 Under 1 year.... 5 4 1 4 3 1 1-4 years..... 2 2 2 1 1 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 2 5 50-54 years..... 4 1 55-59 years..... 3 2 1 60-64 years..... 2 1 1 65-69 vears..... 5 3 70-74 vears..... 3 75-79 years..... 4 80-84 years..... 2 2 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of lysine and hydroxylysine metabolism (E72.3) All ages.... Under 1 year.... 1 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of ornithine metabolism (E72.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of glycine metabolism (E72.5) All ages.... Under 1 year.... 2 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																
Classification of Dise	2222											Δα	ian or			
Tenth Revision, Second		All rac	es	White Black						American Indian 1/			Pacific Islander 2/			
Edition, 2004)	Both	1111 140		Both			Both	Black		Both	10011 11101	.011 17	Both		11401 27	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female	
Other specified disord	ders of amir	no-acid me	tabolism (E	72.8)												
All ages	3	1	2	2	-	2	1	1	-	-	-	-	-	-	-	
Under 1 year	-	-	-	_	-	-	-	_	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	2	-	2	2	_	2	-	-	-	-	-	-	-	-	-	
15-19 years	-	-	-	_	_	-	-	-	-	-	-	-	-	-	-	
20-24 years	1	1	-	_	_	-	1	1	-	-	-	-	-	-	-	
25-29 years	-	-	-	_	_	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-	
40-44 years	-	-	-	_	-	-	_	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	_	_	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-	
75-79 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-	
30-84 years	_	-	-	_	_	_	-	-	_	_	-	-	_	_	-	
35-89 years	_	-	-	_	_	_	-	-	_	_	-	-	_	_	-	
90-94 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-	
95-99 years	-	-	-	_	-	_	-	_	-	-	-	-	-	-	-	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis												As	ian or		
enth Revision, Second All rac		es		White			Black		Ame	rican Indi	an 1/	Pacific Islander 2/			
Edition, 2004)	Both						Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Disorder of amino-aci	d metabolism	ı, unspecif	fied (E72.9)												
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	_	_	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	_	_	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	_
20-24 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
75-79 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	_	_	-	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lactose intolerance (E73) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Congenital lactase deficiency (E73.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																
Classification of Dis							ian or									
Tenth Revision, Second		All races			White			Black			rican Indi	lan 1/	Pacific Islander 2,			
Edition, 2004)	Both			Both			Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Other disorders of ca	rbohydrate m	netabolism	(E74)													
All ages	53	23	30	42	21	21	11	2	9	_	_	_	_	_	_	
Under 1 year	14	4	10	7	2	5	7	2	5	_	-	-	-	_	-	
1-4 years	8	3	5	6	3	3	2	-	2	_	-	-	-	_	-	
5-9 years	1	1	-	1	1	_	_	-	-	_	-	-	-	_	-	
10-14 years	_	_	-	-	-	_	_	_	-	-	-	-	-	-	-	
15-19 years	3	2	1	3	2	1	_	_	-	-	-	-	-	-	-	
20-24 years	2	1	1	2	1	1	_	_	-	-	-	-	-	-	-	
25-29 years	1	1	-	1	1	_	-	-	-	_	-	-	-	-	-	
30-34 years	_	-	-	-	-	_	-	-	-	_	-	-	-	-	-	
35-39 years	1	_	1	1	-	1	_	_	-	-	-	-	-	-	-	
40-44 years	2	1	1	2	1	1	_	_	-	-	-	-	-	-	-	
45-49 years	2	1	1	1	1	_	1	-	1	_	-	-	-	-	-	
50-54 years	1	-	1	1	_	1	_	-	-	_	-	-	-	_	-	
55-59 years	3	2	1	3	2	1	_	-	-	_	-	-	-	-	-	
60-64 years	2	1	1	2	1	1	_	-	-	_	-	-	-	-	-	
65-69 years	4	2	2	3	2	1	1	-	1	_	-	-	-	-	-	
70-74 years	4	1	3	4	1	3	_	_	-	_	_	-	-	_	_	
75-79 years	1	-	1	1	_	1	_	-	-	_	-	-	-	_	-	
30-84 years	4	3	1	4	3	1	_	_	-	_	_	-	-	_	_	
35-89 years	_	-	-	_	_	_	_	-	-	_	-	-	-	_	-	
90-94 years	_	_	-	-	-	_	_	_	-	_	_	-	-	_	_	
95-99 years	_	_	-	-	-	_	_	_	-	_	_	-	-	_	_	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based																	
on the International												_					
Classification of Diseases,				and the second s							Asian or American Indian 1/ Pacific Islander 2/						
Tenth Revision, Second	a Both	All races		Both	White		Both	Black		Ame: Both	rican Indi	lan I/	Pacific Islander 2,				
Edition, 2004)		Male	Female		Male	Female		Male	Female		Male F	7a 7 a	Both	M-l-	Female		
and age	sexes	мате	remaie	sexes	мате	remaie	sexes		remare	sexes	maie r	emale	sexes	мате	remaie		
Glycogen storage disea	ase (E74.0)																
All ages	21	8	13	14	6	8	7	2	5	_	_	_	_	_	_		
Under 1 year	8	3	5	3	1	2	5	2	3	-	-	-	-	-	-		
1-4 years	4	1	3	3	1	2	1	-	1	_	_	_	_	_	-		
5-9 years	_	_	-	_	_	_	-	-	-	_	_	_	_	_	-		
10-14 years	_	_	-	_	_	_	-	-	-	_	_	_	_	_	-		
15-19 years	2	2	-	2	2	_	-	-	-	-	-	-	-	-	-		
20-24 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-		
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
35-39 years	1	_	1	1	_	1	-	-	-	_	_	_	_	_	-		
40-44 years	-	-	-	-	-	_	-	-	-	-	-	-	-	-	-		
45-49 years	2	1	1	1	1	_	1	-	1	-	-	-	-	-	-		
50-54 years	1	-	1	1	_	1	-	-	-	-	-	-	-	-	-		
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
60-64 years	1	-	1	1	_	1	-	-	-	-	-	-	-	-	-		
65-69 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-		
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-		
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
90-94 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-		
95-99 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-		
100 years & over.	-	-	-	-	-	_	-	_	_	_	_	-	_	_	_		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of galactose metabolism (E74.2) All ages.... Under 1 year.... 1 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other disorders of intestinal carbohydrate absorption (E74.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 1 60-64 years..... 1 1 65-69 vears..... 4 70-74 vears..... 2 75-79 years..... 2 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of pyruvate metabolism and gluconeogenesis (E74.4) All ages.... Under 1 year.... 5 4 3 1-4 years..... 1 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified disorders of carbohydrate metabolism (E74.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disorders of sphingolipid metabolism and other lipid storage disorders (E75) All ages.... 75 15 Under 1 year.... 1.0 4 6 9 3 6 1 1-4 years..... 57 25 32 46 2.2 24 17 6 11 14 3 3 5-9 years..... 6 8 10-14 years..... 12 7 12 15-19 years..... 13 11 20-24 years..... 3 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 5 3 50-54 years..... 2 55-59 years..... 3 3 60-64 years..... 3 1 65-69 vears..... 5 5 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 85-89 years..... 1 1 90-94 years..... 95-99 years..... 1 100 years & over. 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes GM2 gangliosidosis (E75.0) All ages..... 13 12 13 Under 1 year.... 1-4 years..... 20 11 18 10 5 1 5-9 years..... 5 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other gangliosidosis (E75.1) All ages..... 9 Under 1 year.... 1-4 years..... 1 1 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other sphingolipidosis (E75.2) All ages.... 35 51 71 39 11 Under 1 year.... 3 4 6 2 4 1 1-4 years..... 2.4 17 17 11 4 8 5 1 3 5-9 years..... 4 10-14 years..... 8 3 5 15-19 years..... 8 2 20-24 years..... 3 25-29 years..... 3 30-34 years..... 4 35-39 years..... 40-44 years..... 3 45-49 years..... 3 50-54 years..... 1 1 55-59 years..... 2 2 60-64 years..... 3 1 65-69 vears..... 3 3 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Neuronal ceroid lipofuscinosis (E75.4) All ages.... 16 Under 1 year.... 1-4 years..... 3 2 5-9 years..... 10-14 years..... 4 2 2 15-19 years..... 2 20-24 years..... 3 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other lipid storage disorders (E75.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Lipid storage disorder, unspecified (E75.6) All ages.... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorders of glycosaminoglycan metabolism (E76) All ages.... 15 Under 1 year.... 1-4 years..... 3 5-9 years..... 10-14 years..... 11 3 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Mucopolysaccharidosis, type I (E76.0) All ages.... Under 1 year.... 1-4 years..... 2 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Mucopolysaccharidosis, type II (E76.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other mucopolysaccharidoses (E76.2) All ages..... 9 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 3 3 15-19 years..... 3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mucopolysaccharidosis, unspecified (E76.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of glycoprotein metabolism (E77) All ages..... 19 Under 1 year.... 1-4 years..... 1 2 5-9 years..... 10-14 years..... 1 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 1 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 2 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Defects in posttranslational modification of lysosomal enzymes (E77.0) All ages.... Under 1 year.... 1-4 years..... 2 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Defects in glycoprotein degradation (E77.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other disorders of glycoprotein metabolism (E77.8) All ages..... 19 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 2 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorder of glycoprotein metabolism, unspecified (E77.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or															
Tenth Revision, Second All races				White		Black			American Indian 1/			nan or Pacific Islander 2/			
Edition, 2004)	Both			Both			Both	Diagn	224011		10011 11101011 17		Both		11401 27
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Disorders of lipoprot	ein metaboli	ism and ot	her lipidem	ias (E78)											
All ages	7,414	3,986	3,428	6,637	3,620	3,017	608	273	335	37	20	17	132	73	59
Under 1 year	1	-	1	1	-	1	-	_	-	-	-	_	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
l0-14 years	1	1	-	-	-	-	1	1	-	-	-	_	-	-	-
5-19 years	-	-	-	-	-	-	-	_	-	-	-	_	-	-	-
20-24 years	4	3	1	4	3	1	-	-	-	-	-	-	-	-	-
25-29 years	8	7	1	7	6	1	1	1	-	-	-	-	-	-	-
30-34 years	10	8	2	8	7	1	2	1	1	-	-	-	-	-	-
35-39 years	29	25	4	25	22	3	3	2	1	-	-	-	1	1	-
10-44 years	69	55	14	63	51	12	5	3	2	1	1	-	-	-	-
15-49 years	175	126	49	141	108	33	27	13	14	3	1	2	4	4	-
50-54 years	301	232	69	258	199	59	38	29	9	1	-	1	4	4	-
55-59 years	460	351	109	398	310	88	50	31	19	5	3	2	7	7	-
50-64 years	565	369	196	497	331	166	54	31	23	6	3	3	8	4	4
65-69 years	670	438	232	580	398	182	72	30	42	6	4	2	12	6	6
70-74 years	810	479	331	715	434	281	79	36	43	2	2	-	14	7	7
75-79 years	1,046	545	501	940	498	442	75	33	42	8	5	3	23	9	14
30-84 years	1,312	620	692	1,202	578	624	84	27	57	2	_	2	24	15	9
35-89 years	1,189	486	703	1,103	456	647	66	22	44	2	1	1	18	7	11
90-94 years	558	190	368	513	175	338	30	7	23	-	-	-	15	8	7
95-99 years	171	44	127	158	39	119	12	5	7	-	-	-	1	-	1
100 years & over.	35	7	28	24	5	19	9	1	8	1	-	1	1	1	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Pure hypercholesterolemia (E78.0) All ages..... 1,823 1,604 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.7 50-54 years..... 2.0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Pure hyperglyceridemia (E78.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 2 2 50-54 years..... 2 55-59 years..... 5 4 1 60-64 years..... 4 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Mixed hyperlipidemia (E78.2) All ages..... 15 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 3 2 60-64 years..... 1 1 65-69 vears..... 2 70-74 vears..... 2 75-79 years..... 80-84 years..... 4 2 85-89 years..... 90-94 years..... 2 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other hyperlipidemia (E78.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Hyperlipidemia, unspecified (E78.5) All ages.... 5,486 2,957 2,529 4,939 2,697 2,242 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.2 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2.0 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lipoprotein deficiency (E78.6) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 1 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 1 85-89 years..... 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other disorders of lipoprotein metabolism (E78.8) All ages.... Under 1 year.... 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorder of lipoprotein metabolism, unspecified (E78.9) All ages..... 10 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 2 2 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 6 90-94 years..... 2 2 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of purine and pyrimidine metabolism (E79) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hyperuricemia without signs of inflammatory arthritis and tophaceous disease (E79.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Lesch-Nyhan syndrome (E79.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other disorders of purine and pyrimidine metabolism (E79.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of porphyrin and bilirubin metabolism (E80) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 1 55-59 years..... 1 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Porphyria cutanea tarda (E80.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other porphyria (E80.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Disorders of mineral metabolism (E83) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.0 Ω 50-54 years..... 2.4 1 0 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 2.8 80-84 years..... 2.0 85-89 years..... 90-94 years..... 2.0 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																		
Classification of Diseases,											Asian or							
Tenth Revision, Second		All races			White			Black			American Indian 1/			Pacific Islander 2/				
Edition, 2004)	Both			Both			Both			Both			Both					
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female			
Disorders of copper m	etabolism (E	83.0)																
All ages	19	15	4	13	10	3	3	2	1	_	_	_	3	3	_			
Under 1 year	1	1	-	1	1	-	_	-	-	-	-	-	-	-	-			
1-4 years	4	4	-	2	2	-	1	1	-	_	-	-	1	1	-			
5-9 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-			
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
15-19 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-			
20-24 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-			
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
30-34 years	3	1	2	1	-	1	1	-	1	-	-	-	1	1	-			
35-39 years	3	3	-	3	3	-	-	-	-	-	-	-	-	-	-			
10-44 years	1	1	-	-	-	-	-	-	-	-	-	-	1	1	-			
45-49 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-			
50-54 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-			
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-			
65-69 years	-	-	-	-	_	-	_	-	-	-	-	-	_	-	-			
70-74 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-			
'5-79 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-			
80-84 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-			
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
95-99 years	-	-	-	-	_	-	_	-	-	-	-	-	_	-	-			
100 years & over.	_	-	-	-	-	-	_	-	_	_	_	-	_	-	_			

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of iron metabolism (E83.1) All ages.... 129 197 120 77 14 Under 1 year.... 4 2 2 4 2 2 1-4 years..... 1 1 1 5-9 years..... 1 10-14 years..... 15-19 years..... 1 1 20-24 years..... 3 25-29 years..... 30-34 years..... 1 35-39 years..... 10 2 40-44 years..... 3 45-49 years..... 1.0 3 5 50-54 years..... 10 6 10 55-59 years..... 23 17 21 17 6 60-64 years..... 27 19 8 26 18 65-69 vears..... 18 13 5 18 13 5 70-74 vears..... 22 10 12 22 10 12 75-79 years..... 45 2.2 23 42 2.0 2.2 80-84 years..... 18 11 7 18 11 7 85-89 years..... 14 8 6 14 8 6 90-94 years..... 2 2 2 2 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of phosphorus metabolism (E83.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of magnesium metabolism (E83.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 3 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disorders of calcium metabolism (E83.5) All ages.... 121 152 96 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 1 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 9 45-49 years..... 5 2 2 3 9 50-54 years..... 5 2 55-59 years..... 13 9 8 5 4 3 5 60-64 years..... 16 12 11 65-69 vears..... 12 5 5 7 70-74 vears..... 18 6 12 15 75-79 years..... 16 9 7 13 6 80-84 years..... 31 11 20 2.7 9 18 4 2 24 24 85-89 years..... 10 14 10 14 3 90-94 years..... 21 5 16 18 13 3 5 95-99 years..... 8 8 8 8 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Cystic fibrosis (E84) All ages.... 438 214 224 415 202 213 16 Under 1 year..... 11 1.0 1 11 10 1 1-4 years..... 3 1 2 3 1 2 9 5 9 5-9 years..... 10-14 years..... 25 12 13 2.4 11 13 1 15-19 years..... 30 32 57 26 31 62 20-24 years..... 35 55 25-29 years..... 77 31 30-34 years..... 52 31 21 49 29 2.0 35-39 years..... 33 18 15 31 17 11 40-44 years..... 28 17 11 28 45-49 years..... 14 8 6 12 Ω 5 50-54 years..... 13 10 3 12 1 0 1 55-59 years..... 6 3 3 6 3 60-64 years..... 5 3 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 2 2 2 85-89 years..... 3 1 90-94 years..... 3 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Cystic fibrosis with pulmonary manifestations (E84.0) All ages.... 226 132 12 Under 1 year..... 3 3 3 3 1-4 years..... 2 5-9 years..... 10-14 years..... 18 8 10 18 8 10 15-19 years..... 39 23 35 2.2 16 13 20-24 years..... 51 17 25-29 years..... 39 30-34 years..... 22 11 11 19 10 35-39 years..... 24 14 10 10 40-44 years..... 18 10 8 18 10 45-49 years..... 4 3 2 50-54 years..... 4 3 1 1 55-59 years..... 2 4 2 60-64 years..... 5 1 4 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Cystic fibrosis with intestinal manifestations (E84.1) All ages.... 30 12 Under 1 year.... 5 4 1 5 1 1-4 years..... 2 2 2 5-9 years..... 10-14 years..... 15-19 years..... 2 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 2 45-49 years..... 3 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Cystic fibrosis with other manifestations (E84.8) All ages.... 11 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 1 1 1 15-19 years..... 3 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 1 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Cystic fibrosis, unspecified (E84.9) All ages..... 130 72 Under 1 year.... 2 2 2 2 1-4 years..... 1 1 1 5-9 years..... 4 1 10-14 years..... 4 1 3 3 3 15-19 years..... 16 15 20-24 years..... 29 13 16 25-29 years..... 30 16 30-34 years..... 21 13 8 13 35-39 years..... 6 2 40-44 years..... 45-49 years..... 5 5 9 50-54 years..... 9 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 2 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Amyloidosis (E85) All ages.... Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.8 2.4 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 8.5 80-84 years..... 85-89 years..... 90-94 years..... 2.8 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Neuropathic heredofamilial amyloidosis (E85.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Heredofamilial amyloidosis, unspecified (E85.2) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 50-54 years..... 55-59 years..... 1 60-64 years..... 3 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Secondary systemic amyloidosis (E85.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Organ-limited amyloidosis (E85.4) All ages..... 268 197 399 229 170 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 6 40-44 years..... 45-49 years..... 1.0 6 4 10 6 4 21 50-54 years..... 15 14 6 1 0 55-59 years..... 35 20 28 16 12 6 15 4 60-64 years..... 40 25 15 32 20 12 3 65-69 years..... 48 24 24 40 20 20 70-74 years..... 65 40 25 56 22 6 75-79 years..... 76 45 31 69 41 28 2 80-84 years..... 87 47 40 79 43 36 85-89 years..... 50 31 19 44 28 16 4 2 2 7 90-94 years..... 2.0 13 19 13 1 6 1 95-99 years..... 3 2 1 3 2 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other amyloidosis (E85.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 4 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Amyloidosis, unspecified (E85.9) All ages..... 282 199 398 162 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 5 40-44 years..... 11 45-49 years..... 2.4 18 6 18 14 5 26 22 50-54 years..... 16 10 14 55-59 years..... 39 28 11 28 21 8 3 5 60-64 years..... 56 35 21 48 30 18 8 65-69 years..... 73 39 55 30 25 70-74 years..... 72 43 29 37 9 75-79 years..... 82 47 35 73 42 31 80-84 years..... 56 28 28 50 25 25 4 2 1 29 25 85-89 years..... 15 14 13 12 2 9 90-94 years..... 5 3 1 1 4 8 5 95-99 years..... 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International												3 -	sian or		
Classification of Diseases, Tenth Revision, Second All races					Black		Amer	ican Ind			ic Isla	nder 2/			
Edition, 2004)	Both	All laces		White Both		Both	BIACK		Both	Ican Inu	itali 1/	Both	.IC ISIA	naer 2/	
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
Volume depletion (E86)														
All ages	3,509	1,120	2,389	3,097	958	2,139	361	144	217	12	4	8	39	14	25
Under 1 year	11	6	5	8	4	4	3	2	1	-	-	-	-	-	-
1-4 years	8	3	5	6	2	4	1	1	-	-	-	-	1	-	1
5-9 years	3	2	1	2	2	-	1	-	1	-	-	-	-	-	-
10-14 years	1	-	1	-	-	-	1	-	1	-	-	-	-	-	-
15-19 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
20-24 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
25-29 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-
30-34 years	2	2	-	1	1	-	1	1	-	-	-	-	-	-	-
35-39 years	7	5	2	3	3	-	4	2	2	-	-	-	-	-	-
40-44 years	14	7	7	10	5	5	4	2	2	-	-	-	-	-	-
45-49 years	17	7	10	14	6	8	2	1	1	-	-	-	1	-	1
50-54 years	32	21	11	24	14	10	8	7	1	-	-	-	-	-	-
55-59 years	51	33	18	41	24	17	10	9	1	-	-	-	-	-	-
60-64 years	46	29	17	39	24	15	6	5	1	1	-	1	-	-	-
65-69 years	87	41	46	67	31	36	18	9	9	1	1	-	1	-	1
70-74 years	151	67	84	124	54	70	22	10	12	5	3	2	-	-	-
75-79 years	330	133	197	290	110	180	35	20	15	2	-	2	3	3	-
80-84 years	575	211	364	503	184	319	63	25	38	-	-	-	9	2	7
85-89 years	810	241	569	744	224	520	58	16	42	-	-	-	8	1	7
90-94 years	844	208	636	765	182	583	65	19	46	2	-	2	12	7	5
95-99 years	396	86	310	352	72	280	41	14	27	-	-	-	3	-	3
100 years & over.	114	14	100	94	12	82	18	1	17	1	-	1	1	1	-
Age not stated	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Other disorders of fluid, electrolyte, and acid-base balance (E87) All ages.... 2,625 1,030 1,595 2,121 1,292 Under 1 year.... 2.7 1.0 1.0 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 2.8 1.3 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Hyperosmolality and hypernatremia (E87.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hypo-osmolality and hyponatremia (E87.1) All ages..... 179 219 161 Under 1 year.... 3 2 1 1 1 1 1 1-4 years..... 1 1 1 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 6 2 2 9 50-54 years..... 5 3 55-59 years..... 12 4 5 3 60-64 years..... 11 5 6 5 65-69 vears..... 11 7 70-74 vears..... 16 3 13 15 12 75-79 years..... 2.4 6 18 16 80-84 years..... 39 10 29 38 10 28 1 55 52 3 85-89 years..... 11 44 10 42 7 90-94 years..... 3.8 31 34 28 3 3 6 95-99 years..... 12 3 9 12 3 9 100 years & over. 5 3 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Acidosis (E87.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Alkalosis (E87.3) All ages.... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 2 60-64 years..... 2 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mixed disorder of acid-base balance (E87.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hyperkalemia (E87.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 2.3 80-84 years..... 2.7 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hypokalemia (E87.6) All ages.... 134 117 79 13 Under 1 year.... 1 1 1 1 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 2 45-49 years..... 11 8 2 2 50-54 years..... 11 5 55-59 years..... 9 1 5 4 3 60-64 years..... 10 6 65-69 vears..... 4 3 70-74 years..... 5 75-79 years..... 15 10 13 9 80-84 years..... 13 11 13 11 17 85-89 years..... 2 15 15 13 17 90-94 years..... 13 12 1 1 4 16 95-99 years..... 4 3 2 1 1 3 1 1 100 years & over. 3 3 3 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																
Classification of Diseases,									Asian or							
Tenth Revision, Second			D. 1-1-	White			Black			American Indian 1/			Pacific Islander 2/			
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	remale	Both sexes	Male	Female	
Fluid overload (E87.7)															
All ages	83	41	42	64	32	32	17	7	10	2	2	_	_	_	-	
Under 1 year	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
.0-14 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
5-19 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-	
20-24 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-	
25-29 years	2	1	1	1	-	1	1	1	-	-	-	-	-	-	-	
30-34 years	1	-	1	-	-	-	1	_	1	-	-	-	-	-	-	
35-39 years	-	_	-	_	_	-	-	_	-	-	_	-	-	-	-	
10-44 years	3	1	2	2	1	1	1	_	1	-	_	-	-	-	-	
5-49 years	3	2	1	2	2	-	1	_	1	-	_	-	-	-	-	
50-54 years	3	2	1	2	1	1	1	1	-	-	-	-	-	-	-	
55-59 years	4	1	3	2	-	2	2	1	1	-	-	-	-	-	-	
0-64 years	10	9	1	9	8	1	_	-	-	1	1	-	-	-	-	
5-69 years	3	1	2	2	-	2	_	-	-	1	1	-	-	-	-	
70-74 years	4	3	1	3	2	1	1	1	_	_	-	-	-	-	-	
5-79 years	10	5	5	8	4	4	2	1	1	_	-	-	-	-	-	
0-84 years	14	7	7	11	6	5	3	1	2	_	-	-	-	-	-	
5-89 years	10	2	8	7	1	6	3	1	2	_	-	-	-	-	-	
0-94 years	11	4	7	10	4	6	1	_	1	_	-	-	-	-	-	
95-99 years	4	2	2	4	2	2	-	_	_	_	-	-	-	-	-	
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Other disorders of electrolyte and fluid balance, not elsewhere classified (E87.8) All ages.... Under 1 year..... 2 1 1 2 1 1 1-4 years..... 2 1 1 1 5-9 years..... 1 1 10-14 years..... 15-19 years..... 1 20-24 years..... 1 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 8 45-49 years..... 10 5 5 3 3 50-54 years..... 12 5 11 55-59 years..... 12 6 9 3 6 60-64 years..... 12 8 4 9 3 3 12 7 65-69 vears..... 5 6 70-74 vears..... 24 20 19 16 75-79 years..... 29 9 2.0 20 13 80-84 years..... 55 18 37 48 16 32 7 2 5 73 63 50 85-89 years..... 14 59 13 10 9 90-94 years..... 55 1.0 45 49 42 6 3 95-99 years..... 24 5 19 22 17 2 2 100 years & over. 1 3 2 4 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second All races			White Black						Asian or American Indian 1/ Pacific Islander 2/						
Edition, 2004)	a Both	All rac	races White Both				American Indian 1/ Both			Both					
and age	sexes	Male	Female	sexes	Male	Female	Both sexes	Male	Female	sexes	Male F	Female	sexes	Male	Female
Other metabolic disor	ders (E88)														
All ages	778	367	411	681	321	360	78	35	43	4	4	_	15	7	8
Under 1 year	43	25	18	34	19	15	5	4	1	-	-	-	4	2	2
1-4 years	30	13	17	28	11	17	1	1	-	-	-	-	1	1	-
5-9 years	10	7	3	9	6	3	1	1	-	-	-	-	-	-	-
10-14 years	5	3	2	4	2	2	1	1	-	-	-	-	-	-	-
15-19 years	4	4	-	4	4	-	-	_	-	-	-	-	-	-	-
20-24 years	2	2	-	2	2	-	-	_	-	-	-	-	-	-	-
25-29 years	4	1	3	4	1	3	-	_	-	-	-	-	-	-	-
30-34 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
35-39 years	9	6	3	8	5	3	1	1	-	-	-	-	-	-	-
40-44 years	22	12	10	22	12	10	-	-	-	-	-	-	-	-	-
45-49 years	29	20	9	27	18	9	2	2	-	-	-	-	-	-	-
50-54 years	40	20	20	33	15	18	7	5	2	-	-	-	-	-	-
55-59 years	58	32	26	51	26	25	5	4	1	2	2	-	-	-	-
60-64 years	54	26	28	44	20	24	7	3	4	1	1	-	2	2	-
65-69 years	51	26	25	44	22	22	6	3	3	1	1	-	-	-	-
70-74 years	65	28	37	55	26	29	7	1	6	-	-	-	3	1	2
75-79 years	96	44	52	85	41	44	11	3	8	-	-	-	_	-	-
80-84 years	99	49	50	87	46	41	11	3	8	-	-	-	1	-	1
85-89 years	88	28	60	76	26	50	10	2	8	-	-	-	2	-	2
90-94 years	44	13	31	39	11	28	3	1	2	-	-	-	2	1	1
95-99 years	19	6	13	19	6	13	-	-	-	-	-	-	-	-	-
100 years & over.	2	-	2	2	-	2	-	-	-	-	-	-	-	-	-
Age not stated	-	-	-	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Disorders of plasma-protein metabolism, not elsewhere classified (E88.0) All ages.... 125 10 Under 1 year.... 1-4 years..... 1 5-9 years..... 1 10-14 years..... 15-19 years..... 1 20-24 years..... 1 1 25-29 years..... 30-34 years..... 1 35-39 years..... 9 40-44 years..... 5 9 45-49 years..... 16 11 5 16 11 5 2.0 9 50-54 years..... 1.8 ρ 10 11 55-59 years..... 33 16 17 31 15 16 60-64 years..... 20 13 19 13 65-69 vears..... 20 9 19 10 9 70-74 vears..... 19 10 9 18 9 9 75-79 years..... 25 7 25 18 80-84 years..... 25 12 13 25 12 13 26 24 17 85-89 years..... 8 18 90-94 years..... 11 10 1 1 3 8 3 95-99 years..... 2 2 1 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Lipodystrophy, not elsewhere classified (E88.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Lipomatosis, not elsewhere classified (E88.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified metabolic disorders (E88.8) All ages..... 18 17 Under 1 year.... 2 2 2 2 1-4 years..... 5 5 1 5-9 years..... 1 1 10-14 years..... 1 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 1 2 2 50-54 years..... 1 1 55-59 years..... 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 3 75-79 years..... 2 80-84 years..... 1 1 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Metabolic disorder, unspecified (E88.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.8 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Vascular dementia (F01) All ages.... 10,910 3,724 7,186 9,674 3,242 6,432 1,008 383 625 29 18 199 111 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 2 2 2 9 7 50-54 years..... 2 1 1 55-59 years..... 51 27 37 21 13 8 24 16 5 1 60-64 years..... 74 42 32 52 32 20 22 10 12 204 121 83 153 91 62 42 27 15 8 3 65-69 vears..... 70-74 vears..... 467 253 214 368 193 175 92 56 36 5 3 75-79 years..... 1,094 510 584 937 438 499 131 59 72 13 10 2.3 80-84 years..... 2,264 955 1,309 1.971 827 1,144 241 105 136 46 20 26 3,060 989 1,899 24 28 85-89 years..... 2,071 2,808 909 191 53 138 9 6 52 2,490 1,701 20 90-94 years..... 626 1,864 2,246 545 187 57 130 1 52 32 95-99 years..... 999 767 3 167 832 922 155 67 9 58 2 8 5 195 27 168 171 25 146 20 18 1 3 3 100 years & over. 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Vascular dementia of acute onset (F01.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 3 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Multi-infarct dementia (F01.1) All ages..... 2,348 1,516 2,102 1,363 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases,									Asian or							
Tenth Revision, Secon		All races		White		Black			American Indian 1/				ic Isla	nder 2/		
Edition, 2004)	Both			Both			Both			Both			Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male 1	Female 	sexes	Male	Female	
Mixed cortical and su	abcortical va	ascular de	mentia (F01.	.3)												
All ages	3	1	2	3	1	2	_	_	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
15-19 years	_	-	-	-	_	-	-	_	-	-	-	-	-	-	-	
20-24 years	_	-	-	-	_	-	-	_	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-	
80-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
90-94 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-	
Age not stated	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other vascular dementia (F01.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Vascular dementia, unspecified (F01.9) All ages.... 8,551 2,887 5,664 7,561 2,498 5,063 816 316 500 12 153 89 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 4 3 3 1 1 55-59 years..... 30 22 8 2 14 16 12 10 6 60-64 years..... 51 32 19 38 24 14 13 8 5 156 97 59 119 73 31 65-69 vears..... 46 22 9 5 70-74 vears..... 356 201 155 278 124 75 46 2 1 75-79 years..... 855 399 456 728 336 392 107 52 55 18 11 80-84 years..... 1,767 743 1,024 1.532 641 891 194 82 112 36 17 19 2,399 762 699 1,502 41 25 85-89 years..... 1,637 2,201 153 45 108 16 1,965 1,760 1,345 157 90-94 years..... 483 1,482 415 49 108 1 43 15 2.8 95-99 years..... 805 674 737 2 131 120 617 60 9 51 2 4 6 100 years & over. 162 22 140 142 21 121 17 16 1 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Tenth Revision, Second All races White Black American Indian 1, Edition, 2004) Both Both Both Both and age sexes Male Female Sexes Male Fem	Both e sexes Male Female
Classification of Diseases, Tenth Revision, Second All races White Black American Indian 1, Edition, 2004) Both Both Both Both and age sexes Male Female sexes Male Female sexes Male Female sexes Male Female	/ Pacific Islander 2/ Both e sexes Male Female
Tenth Revision, Second All races White Black American Indian 1, Edition, 2004) Both Both Both Both and age sexes Male Female sexes Male Fe	/ Pacific Islander 2/ Both e sexes Male Female
Edition, 2004) Both Both Both Both and age sexes Male Female sexes Male Female sexes Male Female sexes Male Female	Both e sexes Male Female
and age sexes Male Female sexes Male Female sexes Male Female sexes Male Female	e sexes Male Female
Unspecified dementia (F03)	2 700 202 400
	3 700 202 400
All ages 65,167 19,649 45,518 59,078 17,677 41,401 5,120 1,624 3,496 179 56 12	5 /90 292 498
onder i year	
I I Year D	
5 5 Jeans	
10 14 Years	
15 17 Years	
20-24 years	
25-29 years	
30-34 years 2 2 - 2 2	
55 55 Years	- 1 - 1
40-44 years 14 7 7 14 7 7	
15 15 16415	
	1 1 1 -
	1 4 3 1
	1 5 2 3
	2 10 4 6
	3 22 10 12
	9 70 37 33
80-84 years 12,374 4,616 7,758 11,091 4,119 6,972 1,083 408 675 45 16 29	
85-89 years 17,814 5,461 12,353 16,374 5,014 11,360 1,189 361 828 38 11 2	
90-94 years 16,006 3,757 12,249 14,774 3,487 11,287 1,007 215 792 42 10 32	
95-99 years 7,577 1,244 6,333 6,945 1,137 5,808 521 67 454 17 4 13	
	5 32 6 26
Age not stated	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Delirium, not induced by alcohol and other psychoactive substances (F05) All ages.... 193 27 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 8 4 8 50-54 years..... 2 2 55-59 years..... 6 6 6 60-64 years..... 8 6 2 8 65-69 vears..... 16 9 7 6 70-74 vears..... 23 13 10 22 12 10 75-79 years..... 44 21 23 39 19 2.0 80-84 years..... 67 31 36 67 31 36 94 38 56 85-89 years..... 94 38 56 77 75 90-94 years..... 32 45 32 43 95-99 years..... 20 8 12 20 8 12 100 years & over. 6 2 2 5 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Delirium superimposed on dementia (F05.1) All ages..... 100 97 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 4 70-74 vears..... 3 3 75-79 years..... 18 10 11 15 80-84 years..... 30 10 20 30 10 20 57 25 57 25 85-89 years..... 32 32 22 90-94 years..... 44 2.0 24 42 2.0 95-99 years..... 3 8 3 ρ 11 11 100 years & over. 3 2 3 1 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other delirium (F05.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Delirium, unspecified (F05.9) All ages.... 126 102 208 112 96 20 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 3 45-49 years..... 8 4 2 8 50-54 years..... 6 2 2 55-59 years..... 5 5 5 5 60-64 years..... 5 2 5 65-69 vears..... 12 10 6 6 70-74 vears..... 20 12 8 19 11 75-79 years..... 26 14 12 2.4 10 80-84 years..... 37 21 16 37 21 16 37 37 85-89 years..... 13 24 13 24 90-94 years..... 33 12 21 33 12 21 95-99 years..... 9 5 9 5 4 4 100 years & over. 3 3 2 2

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Bot.h and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Other mental disorders due to brain damage and dysfunction and to physical disease (F06) All ages.... 1,255 800 92 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 7 45-49 years..... 6 50-54 years..... 12 11 55-59 years..... 8 7 1 1 60-64 years..... 14 7 11 5 3 65-69 vears..... 26 8 18 22 17 5 70-74 years..... 50 27 23 42 21 6 75-79 years..... 136 52 124 47 77 80-84 years..... 205 80 125 191 73 118 13 1 307 208 286 94 85-89 years..... 99 192 18 3 15 242 65 90-94 years..... 313 71 292 227 19 14 2 5 95-99 years..... 142 21 121 133 20 113 9 8 1 100 years & over. 30 1 29 26 25 4 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dise Tenth Revision, Second		All rac	es		White Bla						Asian or American Indian 1/ Pacific Islander 2/					
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male Fe	emale	Both sexes	Male	Female	
Organic mood [affectiv	ve] disorder:	s (F06.3)														
All ages	3	_	3	3	_	3	_	_	_	_	_	_	_	_	-	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-14 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
15-49 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	1	-	1	1	_	1	-	-	-	-	-	-	-	-	-	
5-79 years	1	-	1	1	_	1	-	-	-	-	-	-	-	-	-	
0-84 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-	
5-89 years	1	-	1	1	_	1	-	-	-	-	-	-	-	-	-	
0-94 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
100 years & over.	_	-	-	_	-	_	-	_	-	-	-	-	-	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Bot.h and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Unspecified mental disorder due to brain damage and dysfunction and to physical disease (F06.9) All ages.... 1,252 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 1 40-44 years..... 7 45-49 years..... 6 50-54 years..... 12 11 55-59 years..... 8 7 1 1 60-64 years..... 14 7 11 5 3 65-69 vears..... 26 8 18 17 22 5 70-74 years..... 49 27 22 41 20 6 75-79 years..... 135 52 83 123 47 76 80-84 years..... 205 80 125 191 73 118 13 1 306 207 285 94 85-89 years..... 99 191 18 3 15 71 65 90-94 years..... 313 242 292 227 19 14 2 5 95-99 years..... 121 142 21 133 20 113 9 8 1 100 years & over. 30 1 29 26 25 4 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Personality and behavioral disorders due to brain disease, damage, and dysfunction (F07) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Organic personality disorder (F07.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Postconcussional syndrome (F07.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	eases,				White Black						Asian or American Indian 1/ Pacific Island							
Edition, 2004)	Both			Both			Both			Both			Both					
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female			
Unspecified organic p	ersonality a	and behavi	oral disord	er due to br	ain disea	ase, damage,	, and dysfu	nction (F07.9)									
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_			
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
0-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
25-29 years	-	-	-	-	-	-	-	-	_	-	-	-	-	-	-			
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
45-49 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-			
50-54 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-			
55-59 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-			
50-64 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-			
65-69 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-			
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
0-84 years	-	_	_	-	_	-	_	-	_	-	_	-	-	-	-			
35-89 years	-	_	_	-	_	-	_	-	_	-	_	-	-	-	-			
				_	_	_	_	_	_	_	_	_	_	_	_			
	_	_	_															
90-94 years 95-99 years	-	_	_	_	_	_	_	_	_	_	_	_	_	_	_			

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis						Asian or									
Tenth Revision, Secon		All races			White			Black			rican Ind:	ian 1/		ic Isla	nder 2/
Edition, 2004)	Both		_ 1	Both		_ ,	Both			Both			Both		_ 1
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male I	Female	sexes	Male	Female
Unspecified organic o	r symptomati	c mental	disorder (F0	09)											
All ages	8	3	5	7	3	4	1	_	1	_	_	_	_	_	_
Under 1 year	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
10-14 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	-	1	-	-	-	1	-	1	-	-	_	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	_	-	-	-
80-84 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
90-94 years	1	1	-	1	1	-	-	-	-	-	-	_	-	-	-
95-99 years	3	-	3	3	-	3	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	-	-	-	-	-	-	-	-	_	-	-	-
Age not stated	_	_	_	_	_	_	_	_	_	_	_	-	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female sexes Male Female sexes Male sexes Male sexes Male Female sexes Mental and behavioral disorders due to use of alcohol (F10) All ages.... 7,648 5,936 1,712 6,385 4,958 1,427 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2.7 20-24 vears..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1.261 1.077 50-54 years..... 1,363 1.071 1,148 1,179 55-59 years..... 60-64 years..... 65-69 years..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2.7 2.4 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Acute intoxication (F10.0) All ages.... 712 566 146 601 473 128 19 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2.7 22 2.6 1 2.1 20-24 years..... 39 31 8 33 26 25-29 years..... 51 46 45 30-34 years..... 44 37 7 35 31 35-39 years..... 70 55 15 57 12 40-44 years..... 119 97 22 104 85 19 3 45-49 years..... 133 104 29 113 87 26 17 3 2 64 72 56 50-54 years..... 82 18 16 8 6 1 1 55-59 years..... 80 57 23 67 46 21 9 11 60-64 years..... 28 24 4 22 18 4 21 13 8 16 9 65-69 vears..... 70-74 years..... 8 6 7 75-79 years..... 6 3 3 80-84 years..... 2 1 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Harmful use (F10.1) All ages.... 1,888 1,463 1,572 1,222 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Dependence syndrome (F10.2) All ages.... 4,208 3,251 3,504 2,709 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.4 1 Q 2.2 50-54 years..... ٩n 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 2.4 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based															
on the International Classification of Dis												7.0	ion or		
Tenth Revision, Second All races				White			Black		Asian or American Indian 1/ Pacific Islander						
Edition, 2004)	Both	AII IAC	CB	Both	WIIICC		Both	DIACK		Both	ican indi	.an i/	Both	.10 1510	naci 2/
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Withdrawal state (F10	.3)														
All ages	101	86	15	89	75	14	9	8	1	3	3	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	_	-	_	_	-	_	-	_	_	-	-	-	_	-
5-9 years	-	-	-	_	_	-	_	-	-	_	-	-	-	-	-
10-14 years	-	-	-	_	_	-	_	-	-	_	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	_	-	_	-	-	_	-	_	-	_	-	-	-	-
25-29 years	-	_	-	_	-	-	_	-	_	-	_	-	-	-	-
30-34 years	3	2	1	2	1	1	_	-	_	1	1	-	-	-	-
35-39 years	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
10-44 years	13	9	4	12	8	4	1	1	_	_	_	_	_	_	_
15-49 years	8	7	1	7	6	1	1	1	_	_	_	_	_	_	_
0-54 years	13	10	3	10	7	3	3	3	_	_	_	_	_	_	_
55-59 years	16	14	2	14	12	2	_	-	_	2	2	-	-	_	-
0-64 years	13	12	1	11	11	-	2	1	1	-	-	-	-	_	-
55-69 years	8	8	-	7	7	-	1	1	_	-	-	-	-	_	-
70-74 years	7	7	_	6	6	_	1	1	_	_	_	_	_	_	_
5-79 years	10	9	1	10	9	1	_	_	_	_	_	_	_	_	_
0-84 years	6	5	1	6	5	1	_	_	_	_	_	_	_	_	_
15-89 years	3	2	1	3	2	1	_	_	_	_	_	_	_	_	_
0-94 years	_	_	_	_	_	_	_	_	_	_	-	_	_	_	-
95-99 years	-	-	-	-	-	-	_	-	_	-	-	-	-	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Withdrawal state with delirium (F10.4) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 2 40-44 years..... 15 15 45-49 years..... 9 9 Ω Ω 1 16 1.3 2 50-54 years..... 11 55-59 years..... 12 11 10 9 60-64 years..... 8 1 8 65-69 vears..... 12 70-74 vears..... 7 6 75-79 years..... 6 80-84 years..... 4 4 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Psychotic disorder (F10.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Amnesic syndrome (F10.6) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 3 60-64 years..... 3 3 65-69 vears..... 3 70-74 vears..... 75-79 years..... 2 80-84 years..... 4 2 2 85-89 years..... 90-94 years..... 2 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Residual and late-onset psychotic disorder (F10.7) All ages.... 176 128 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 2 1 50-54 years..... 1 1 1 55-59 years..... 7 2 5 3 2 60-64 years..... 14 11 3 10 65-69 vears..... 22 20 16 70-74 vears..... 35 28 30 75-79 years..... 40 31 9 38 30 80-84 years..... 43 29 14 35 2.4 11 6 30 17 13 85-89 years..... 34 20 14 3 2 1 1 90-94 years..... 11 6 5 a 5 4 95-99 years..... 4 3 1 1 3 1 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other mental and behavioral disorders (F10.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Unspecified mental and behavioral disorder (F10.9) All ages.... 333 252 43 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 3 20-24 years..... 3 3 3 25-29 years..... 30-34 years..... 7 35-39 years..... 24 15 9 19 28 40-44 years..... 36 32 30 2 41 45-49 years..... 64 49 15 51 1 0 8 6 4 47 55 41 5 50-54 years..... 64 17 14 5 3 55-59 years..... 58 50 8 43 37 6 13 12 2 60-64 years..... 48 41 39 33 1 65-69 vears..... 31 22 9 28 19 2 9 70-74 vears..... 19 9 10 15 75-79 years..... 13 7 12 6 80-84 years..... 12 9 3 11 1 8 85-89 years..... 3 5 8 90-94 years..... 2 2 4 2 95-99 years..... 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Mental and behavioral disorders due to use of opioids (F11) All ages.... 280 206 37 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1.0 20-24 years..... 31 21 10 29 20 25-29 years..... 29 22 7 30-34 years..... 35 31 32 29 35-39 years..... 25 18 7 23 17 40-44 years..... 55 36 19 51 45-49 years..... 45 34 11 40 30 1 0 4 3 36 2.7 9 50-54 years..... 54 43 11 1.8 55-59 years..... 21 17 16 13 5 4 4 60-64 years..... 6 5 5 1 65-69 vears..... 3 3 70-74 vears..... 75-79 years..... 4 80-84 years..... 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Harmful use (F11.1) All ages.... 106 25 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 3 20-24 years..... 3 25-29 years..... 30-34 years..... 10 9 35-39 years..... 6 40-44 years..... 20 13 45-49 years..... 15 11 12 1 0 14 50-54 years..... 2.3 20 3 16 55-59 years..... 8 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Dependence syndrome (F11.2) All ages.... 29 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 12 8 11 25-29 years..... 9 30-34 years..... 11 10 35-39 years..... 7 3 40-44 years..... 17 45-49 years..... 9 6 3 a 50-54 years..... 14 9 1.0 55-59 years..... 4 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Withdrawal state (F11.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Withdrawal state with delirium (F11.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Psychotic disorder (F11.5) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Residual and late-onset psychotic disorder (F11.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Unspecified mental and behavioral disorder (F11.9) All ages.... 100 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 10 10 25-29 years..... 10 9 30-34 years..... 13 11 35-39 years..... 11 9 10 40-44 years..... 17 12 45-49 years..... 21 17 1 2 14 2 16 50-54 years..... 13 3 9 55-59 years..... 9 7 2 60-64 years..... 5 1 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 85-89 years..... 1 90-94 years..... 1 95-99 years..... 100 years & over. 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mental and behavioral disorders due to use of cannabinoids (F12) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Dependence syndrome (F12.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Unspecified mental and behavioral disorder (F12.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mental and behavioral disorders due to use of sedatives or hypnotics (F13) All ages.... 9 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 1 35-39 years..... 40-44 years..... 45-49 years..... 3 2 50-54 years..... 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Harmful use (F13.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Dependence syndrome (F13.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

ause of death (based n the International															
Classification of Dise	eases,										As	sian or			
Tenth Revision, Second		All races			White			Black		Amen	rican Indi			ic Isla	nder 2/
Edition, 2004)	Both			Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Withdrawal state (F13	.3)														
All ages	1	1	_	1	1	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	_	-	-
40-44 years	-	-	-	_	-	-	-	-	-	-	-	-	_	-	-
45-49 years	_	-	-	-	-	_	-	_	-	_	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	_	-	-	-	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	_	-	_	-	_	-	-	-	-	-
75-79 years	_	_	-	_	_	_	-	-	-	_	-	-	_	-	-
30-84 years	_	-	-	-	_	-	-	-	_	_	-	-	_	-	-
35-89 years	_	_	-	_	_	_	-	-	-	_	-	-	_	-	-
90-94 years	_	-	-	-	_	-	-	-	_	_	-	-	_	-	-
95-99 years	_	-	-	-	_	-	-	-	_	_	-	-	_	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Unspecified mental and behavioral disorder (F13.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Mental and behavioral disorders due to use of cocaine (F14) All ages.... 352 261 340 232 108 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25 20 5 23 19 25-29 years..... 51 38 13 11 30-34 years..... 44 33 11 32 12 35-39 years..... 76 51 25 49 33 23 45 40-44 years..... 145 100 71 17 73 45 43 29 45-49 years..... 135 91 44 58 15 77 48 129 87 51 32 19 2.2 50-54 years..... 42 76 54 55-59 years..... 39 32 7 11 2 28 23 5 60-64 years..... 28 21 3 25 18 65-69 vears..... 13 11 8 70-74 vears..... 6 6 6 75-79 years..... 3 3 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Acute intoxication (F14.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Harmful use (F14.1) All ages.... 383 268 115 205 149 176 118 Under 1 year..... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 1 1 20-24 years..... 12 10 11 25-29 years..... 35 2.7 33 30-34 years..... 21 16 5 17 35-39 years..... 40 27 13 27 20 13 38 2.7 40-44 years..... 82 53 29 43 25 25 45-49 years..... 70 45 40 28 12 30 17 12 70 25 2.7 16 11 13 50-54 years..... 45 29 42 55-59 years..... 22 17 5 15 11 6 1 60-64 years..... 15 14 14 13 1 65-69 vears..... 9 6 70-74 years..... 3 3 3 75-79 years..... 3 3 2 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Dependence syndrome (F14.2) All ages.... 32 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 12 40-44 years..... 19 16 45-49 years..... 16 14 12 8 4 3 50-54 years..... 10 55-59 years..... 2 2 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Withdrawal state with delirium (F14.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Unspecified mental and behavioral disorder (F14.9) All ages.... 99 125 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 6 3 3 25-29 years..... 12 30-34 years..... 15 12 10 35-39 years..... 21 13 40-44 years..... 41 28 13 21 12 9 45-49 years..... 47 3.0 17 14 11 33 1 0 14 34 18 11 2.2 8 50-54 years..... 49 15 3.0 55-59 years..... 15 13 2 2 12 11 3 60-64 years..... 13 6 11 5 65-69 vears..... 3 3 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Bot.h Both Bot.h and age Female Male Female Male Female Male Female sexes Male Female sexes Male sexes sexes sexes Mental and behavioral disorders due to use of other stimulants, including caffeine (F15) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 4 25-29 years..... 30-34 years..... 10 9 35-39 years..... 16 12 40-44 years..... 23 18 45-49 years..... 24 17 24 17 15 50-54 years..... 16 12 11 55-59 years..... 5 4 5 60-64 years..... 4 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Harmful use (F15.1) All ages.... 12 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 3 35-39 years..... 10 40-44 years..... 14 12 45-49 years..... 16 11 16 11 9 9 50-54 years..... 11 11 55-59 years..... 3 3 3 60-64 years..... 3 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Dependence syndrome (F15.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 1 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Unspecified mental and behavioral disorder (F15.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 6 45-49 years..... 5 1 50-54 years..... 4 2 55-59 years..... 2 1 1 60-64 years..... 1 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mental and behavioral disorders due to use of hallucinogens (F16) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Harmful use (F16.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Mental and behavioral disorders due to use of tobacco (F17) All ages.... 1,415 1,269 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2.7 a 45-49 years..... 2.8 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Harmful use (F17.1) All ages.... 176 104 164 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 2 3 5 2 3 16 10 50-54 years..... 14 1 0 55-59 years..... 25 19 22 17 6 5 60-64 years..... 21 15 6 21 15 65-69 vears..... 19 6 18 13 13 5 70-74 vears..... 29 22 7 29 7 75-79 years..... 2.0 8 19 11 12 80-84 years..... 17 9 8 14 6 1 1 85-89 years..... 11 4 10 90-94 years..... 5 2 3 5 3 95-99 years..... 1 1 1 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

ause of death (based n the International lassification of Diseases, Asian or															
Tenth Revision, Second All races			es.		White			Black			ican Indi		Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both	rican maran r		Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Dependence syndrome (F17.2)														
All ages	39	24	15	37	23	14	1	_	1	_	_	_	1	1	_
Under 1 year	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-14 years	-	-	-	-	-	_	_	-	-	-	-	_	-	-	-
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
5-49 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
0-54 years	4	2	2	4	2	2	-	-	-	-	-	-	-	-	-
55-59 years	4	3	1	4	3	1	-	-	-	-	-	-	-	-	-
0-64 years	6	4	2	6	4	2	-	-	-	-	-	-	-	-	-
5-69 years	6	4	2	6	4	2	-	-	-	-	-	-	-	-	-
0-74 years	7	4	3	6	4	2	1	-	1	-	-	_	-	-	-
5-79 years	2	2	-	2	2	_	_	-	-	-	-	_	-	-	-
0-84 years	4	3	1	4	3	1	-	-	-	-	-	-	-	-	-
5-89 years	4	2	2	3	1	2	-	-	-	-	-	-	1	1	-
0-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
00 years & over.	_	-	-	-	_	-	-	_	_	_	-	_	_	-	-

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Unspecified mental and behavioral disorder (F17.9) All ages.... 1,200 1,068 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... a 50-54 years..... 2.0 1 0 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mental and behavioral disorders due to use of volatile solvents (F18) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

ause of death (based n the International Lassification of Diseases, Asian or															
Tenth Revision, Second All races			White Black						Asian or American Indian 1/ Pacific Islander 2/						
Edition, 2004)	Both	THE EUCCO		Both			Both			Both			Both		
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Harmful use (F18.1)															
All ages	3	3	_	3	3	_	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
L5-19 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-34 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
35-39 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
15-49 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-
0-54 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	_	-	-	-	-	-	_	-	-	-	-	-	-	-
70-74 years	-	_	-	-	-	-	-	-	-	_	-	-	-	-	-
5-79 years	-	_	-	-	-	-	-	-	-	_	-	-	-	-	-
0-84 years	_	_	_	_	_	_	_	_	_	_	_	_	-	_	-
5-89 years	-	_	-	-	-	-	-	-	-	_	-	-	-	-	-
0-94 years	-	_	-	-	-	-	-	-	-	_	-	-	-	-	-
95-99 years	_	_	_	_	_	_	_	_	_	_	_	_	-	_	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Dependence syndrome (F18.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Unspecified mental and behavioral disorder (F18.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 1 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Bot.h Both Bot.h Both Bot.h and age Female Male Female Male Female sexes Male Female sexes Male Female sexes Male sexes sexes Mental and behavioral disorders due to multiple-drug use and use of other psychoactive substances (F19) All ages.... 1,600 1,082 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2.7 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Acute intoxication (F19.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or Tenth Revision, Second All races White Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Bot.h Both Both Bot.h and age Female Female Male Female Male Female sexes Male Female sexes Male sexes Male sexes sexes Harmful use (F19.1) All ages.... 1,201 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2.7 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 years..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over. Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Dependence syndrome (F19.2) All ages.... 163 77 197 133 36 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2 2 2 2 20-24 years..... 15 14 13 13 25-29 years..... 19 13 6 30-34 years..... 20 14 6 18 13 35-39 years..... 29 17 12 25 11 14 3 40-44 years..... 45 31 14 42 45-49 years..... 37 23 14 27 16 11 1 0 26 8 50-54 years..... 36 24 12 1.8 ρ 55-59 years..... 23 16 7 16 11 5 60-64 years..... 5 2 5 3 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 85-89 years..... 90-94 years..... 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Withdrawal state (F19.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Residual and late-onset psychotic disorder (F19.7) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Unspecified mental and behavioral disorder (F19.9) All ages.... 131 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 2 1 20-24 years..... 11 3 10 25-29 years..... 9 9 30-34 years..... 10 10 35-39 years..... 20 13 18 40-44 years..... 18 11 18 45-49 years..... 33 20 13 2.6 15 11 18 1.3 5 50-54 years..... 21 15 6 2 55-59 years..... 9 5 4 3 60-64 years..... 4 2 2 65-69 vears..... 3 70-74 vears..... 2 75-79 years..... 80-84 years..... 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Schizophrenia (F20) All ages.... 426 181 245 369 151 218 48 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 10 3 8 45-49 years..... 21 1.0 11 13 50-54 years..... 2.4 16 8 18 12 55-59 years..... 20 12 8 16 10 2 6 60-64 years..... 30 18 12 23 12 11 65-69 years..... 33 20 13 32 19 13 70-74 years..... 46 21 25 40 19 21 3 75-79 years..... 58 2.8 30 52 2.7 25 1 80-84 years..... 70 21 49 65 17 48 58 85-89 years..... 60 18 42 18 40 1 90-94 years..... 29 28 2.7 27 2 1 1 95-99 years..... 9 9 8 8 1 1 100 years & over. 1 3 1 4 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Paranoid schizophrenia (F20.0) All ages..... 19 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 3 2 50-54 years..... 1 55-59 years..... 1 1 60-64 years..... 2 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 3 80-84 years..... 6 85-89 years..... 4 3 90-94 years..... 2 2 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Catatonic schizophrenia (F20.2) All ages..... 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 50-54 years..... 55-59 years..... 1 60-64 years..... 3 1 65-69 vears..... 2 70-74 vears..... 3 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Undifferentiated schizophrenia (F20.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Postschizophrenic depression (F20.4) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Residual schizophrenia (F20.5) All ages..... 19 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 50-54 years..... 1 1 55-59 years..... 60-64 years..... 2 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 9 3 8 85-89 years..... 90-94 years..... 1 1 95-99 years..... 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Schizophrenia, unspecified (F20.9) All ages..... 140 196 293 115 178 37 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 7 40-44 years..... 45-49 years..... 15 8 12 3 5 50-54 years..... 18 5 12 3 1.3 55-59 years..... 17 11 6 13 2 60-64 years..... 22 13 9 17 9 65-69 years..... 28 16 12 27 15 70-74 years..... 38 18 20 34 16 18 75-79 years..... 45 19 26 41 19 2.2 80-84 years..... 54 16 38 50 12 38 4 47 46 85-89 years..... 13 34 13 33 25 90-94 years..... 24 23 23 1 1 95-99 years..... 5 5 5 5 100 years & over. 1 1 4 3 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Persistent delusional disorders (F22) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Delusional disorder (F22.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 4 85-89 years..... 1 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second All races				White Black						Asian or American Indian 1/ Pacific Islander 2/						
Edition, 2004)	Both			Both			Both			Both	ream maram 17		Both			
and age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Acute and transient p	osychotic dis	sorders (F	23)													
All ages	7	5	2	3	2	1	2	2	_	1	1	_	1	_	1	
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
l0-14 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	-	-	-	-	-	-	_	-	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
60-64 years	1	1	-	-	-	-	-	-	-	1	1	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
75-79 years	2	1	1	1	1	-	_	-	-	-	-	-	1	-	1	
80-84 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-	
85-89 years	1	1	-	-	-	-	1	1	-	_	-	-	-	-	-	
90-94 years	1	1	-	1	1	-	-	-	-	_	-	-	-	-	-	
95-99 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
100 years & over.	_	-	_	_	_	_	_	_	_	_	_	_	_	_	_	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Acute and transient psychotic disorder, unspecified (F23.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 1 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Schizoaffective disorders (F25) All ages..... 19 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 1 50-54 years..... 55-59 years..... 1 60-64 years..... 2 1 65-69 vears..... 2 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 85-89 years..... 2 90-94 years..... 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Schizoaffective disorder, mixed type (F25.2) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Schizoaffective disorder, unspecified (F25.9) All ages.... 18 10 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 5 1 50-54 years..... 55-59 years..... 1 2 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 85-89 years..... 2 90-94 years..... 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other nonorganic psychotic disorders (F28) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International																
Classification of Diseases,										Asian or						
Tenth Revision, Secon		All rac	es	_	White		_	Black			rican Indi	ian 1/	Pacific Islander 2/			
Edition, 2004)	Both	_	_	Both	_	_	Both	_	_	Both			Both	_	_	
nd age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male H	Female	sexes	Male	Female	
Unspecified nonorgani	c psychosis	(F29)														
All ages	33	10	23	28	8	20	5	2	3	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-	
1-4 years	_	-	_	_	_	_	-	-	_	_	-	_	_	-	-	
5-9 years	_	-	_	_	_	_	-	-	_	_	-	_	_	-	-	
10-14 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-	
l5-19 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-	
20-24 years	-	-	-	_	-	-	-	-	-	_	-	-	-	-	-	
25-29 years	1	-	1	_	-	-	1	-	1	_	-	-	-	-	-	
30-34 years	1	-	1	1	-	1	-	-	-	_	-	-	-	-	-	
35-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	-	_	
40-44 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-	
15-49 years	_	-	-	-	-	-	-	_	-	_	-	_	-	-	-	
50-54 years	2	1	1	1	-	1	1	1	-	_	-	_	-	-	-	
55-59 years	3	2	1	2	2	-	1	_	1	_	-	_	-	-	-	
0-64 years	3	2	1	2	1	1	1	1	_	_	-	_	-	-	-	
55-69 years	2	1	1	2	1	1	-	-	_	_	-	_	_	-	-	
70-74 years	3	1	2	3	1	2	-	_	-	_	-	_	-	-	-	
75-79 years	6	1	5	5	1	4	1	_	1	_	-	_	-	-	_	
30-84 years	4	1	3	4	1	3	_	_	_	_	_	_	_	_	_	
35-89 years	2	_	2	2	_	2	_	_	_	_	_	_	_	_	_	
90-94 years	4	1	3	4	1	3	_	_	_	_	_	_	_	_	_	
95-99 years	2	_	2	2	_	2	_	_	-	_	-	_	-	-	_	
100 years & over.	_	_	_	_	_	-	-	_	_	_	_	_	_	_	-	

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Manic episode (F30) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseas	ses,	- 1.1						-1 1		Asian or						
Tenth Revision, Second Edition, 2004)	Both	All races		White Both			Black			Ame: Both	rican Indi	ican Indian 1/		Pacific Islander 2/ Both		
and age	sexes	Male	Female	sexes	Male	Female	Both sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female	
Other manic episodes (F.	30.8)															
All ages	1	1	_	_	_	_	1	1	_	_	_	_	_	_	_	
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5-9 years	-	-	-	-	-	-	-	-	-	-	_	-	-	-	-	
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
l5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
20-24 years	1	1	-	-	-	-	1	1	-	-	-	-	-	-	-	
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
50-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
70-74 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
35-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
90-94 years 95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
		_		_			_									

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Manic episode, unspecified (F30.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Bipolar affective disorder (F31) All ages..... 59 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 3 30-34 years..... 35-39 years..... 40-44 years..... 6 2 2 45-49 years..... 7 50-54 years..... 2 5 55-59 years..... 3 2 1 60-64 years..... 8 65-69 vears..... 4 70-74 vears..... 11 75-79 years..... 7 6 80-84 years..... 16 13 15 12 1 85-89 years..... 11 1 10 10 9 1 90-94 years..... 2 2 1 1 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Bipolar affective disorder, unspecified (F31.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 3 30-34 years..... 35-39 years..... 40-44 years..... 6 2 2 45-49 years..... 7 50-54 years..... 2 5 55-59 years..... 3 2 1 60-64 years..... 8 65-69 vears..... 4 70-74 vears..... 11 75-79 years..... 7 6 80-84 years..... 16 13 15 12 1 85-89 years..... 11 1 10 10 9 1 90-94 years..... 2 2 1 1 1 95-99 years..... 1 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Depressive episode (F32) All ages.... 623 187 436 585 174 411 31 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 6 2 4 5 2 50-54 years..... 5 9 11 6 55-59 years..... 15 5 10 11 4 2 60-64 years..... 10 6 9 1 65-69 years..... 12 7 12 7 5 5 70-74 years..... 30 12 18 28 12 16 75-79 years..... 82 26 56 71 50 10 80-84 years..... 124 33 91 123 33 90 1 1 83 79 5 85-89 years..... 136 53 128 49 3 127 95 1 90-94 years..... 25 102 120 25 6 6 95-99 years..... 39 49 38 50 11 11 100 years & over. 10 1 9 10 9 1 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Severe depressive episode without psychotic symptoms (F32.2) All ages.... 17 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 2 55-59 years..... 3 3 60-64 years..... 4 3 65-69 vears..... 4 70-74 vears..... 6 3 3 75-79 years..... 12 9 8 80-84 years..... 20 16 20 16 10 85-89 years..... 11 10 9 9 90-94 years..... 1 1 8 8 95-99 years..... 2 1 1 2 100 years & over. 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Severe depressive episode with psychotic symptoms (F32.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 2 2 2 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis Tenth Revision, Secon	seases,	All rac			White			Black	Asian or American Indian 1/ Pacific Islander 2/						
		All rac	es	Dath	white		Dath	втаск					Pacific Islander 2/		
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	'emale	Both sexes	Male	Female
Other depressive epis	odes (F32.8)														
All ages	2	_	2	2	_	2	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	_	-	-	-	-	-	-	_	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
.5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-49 years	-	-	-	_	_	-	-	_	-	-	_	-	-	-	-
0-54 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
0-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-69 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
70-74 years	-	-	-	_	-	-	-	-	-	_	_	-	-	-	-
5-79 years	-	-	-	_	-	-	-	-	-	_	_	-	-	-	-
0-84 years	-	-	-	_	-	-	-	-	-	_	_	-	-	-	-
5-89 years	1	-	1	1	-	1	-	-	-	_	_	-	-	-	-
00-94 years	-	-	-	_	-	-	-	-	-	_	_	-	-	-	-
95-99 years	1	-	1	1	-	1	-	-	-	_	_	-	-	-	-
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Depressive episode, unspecified (F32.9) All ages.... 375 508 154 354 25 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 1 30-34 years..... 35-39 years..... 40-44 years..... 4 45-49 years..... 5 1 9 2 50-54 years..... 5 3 55-59 years..... 12 5 60-64 years..... 6 3 3 3 65-69 vears..... 8 6 8 70-74 vears..... 23 8 15 13 75-79 years..... 70 48 62 19 43 80-84 years..... 103 28 75 102 28 74 1 1 71 68 85-89 years..... 123 52 116 48 1 90-94 years..... 116 2.4 92 110 2.4 86 5 5 95-99 years..... 46 37 45 9 9 36 100 years & over. 1 8 9 1 8

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Recurrent depressive disorder (F33) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Dis												As	ian or		
Tenth Revision, Secon	nd	All races			White			Black		American Indian 1/			Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
nd age	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	'emale	sexes	Male	Female
Recurrent depressive	disorder, cu	rrent epi:	sode severe	without psy	chotic sy	mptoms (F33	.2)								
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	_	-	-	-	_	_	_	-	-	_	-	-	-	-	-
5-9 years	_	-	-	-	_	_	-	_	-	-	-	-	-	-	-
10-14 years	_	-	-	-	_	_	-	_	-	-	-	-	-	-	-
L5-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
0-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
10-44 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
15-49 years	-	-	-	-	-	-	-	_	-	_	_	-	-	-	-
50-54 years	-	-	-	-	-	-	-	_	-	_	_	-	-	-	-
55-59 years	_	-	-	-	_	_	_	-	-	_	-	-	-	-	_
0-64 years	_	-	-	-	_	_	-	_	-	-	-	-	-	-	-
5-69 years	_	-	-	-	_	_	-	_	-	-	-	-	-	-	-
70-74 years	-	-	-	-	-	-	-	_	-	_	_	-	-	-	-
'5-79 years	_	-	_	_	_	-	_	_	_	-	-	_	-	_	-
0-84 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
35-89 years	1	-	1	1	_	1	_	_	_	-	-	_	-	_	-
00-94 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
95-99 years	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Recurrent depressive disorder, unspecified (F33.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Persistent mood [affective] disorders (F34) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Cyclothymia (F34.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Dysthymia (F34.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Unspecified mood [affective] disorder (F39) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other anxiety disorders (F41) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 2 1 2 2 50-54 years..... 2 55-59 years..... 2 2 60-64 years..... 1 1 65-69 vears..... 3 70-74 vears..... 75-79 years..... 80-84 years..... 5 85-89 years..... 5 90-94 years..... 6 1 5 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Panic disorder [episodic paroxysmal anxiety] (F41.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Generalized anxiety disorder (F41.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 2 90-94 years..... 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mixed anxiety and depressive disorder (F41.2) All ages..... 12 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 3 80-84 years..... 3 85-89 years..... 90-94 years..... 1 1 95-99 years..... 1 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis												Asian or						
Tenth Revision, Secon		All rac	es	Dath	White		Black			American Indian 1/			Pacific Islander 2/ Both					
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	ama la	sexes	Mala	Female			
															- Chiare			
Other specified anxie	ty disorders	s (F41.8)																
All ages	1	_	1	1	_	1	_	_	_	_	_	_	_	_	_			
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
10-14 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-			
l5-19 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-			
20-24 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	_			
25-29 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	_			
30-34 years	-	-	-	-	_	-	-	_	-	-	-	-	-	-	-			
35-39 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-			
40-44 years	-	-	-	-	_	-	-	_	-	_	-	-	-	-	-			
15-49 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-			
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
50-64 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-			
65-69 years	-	-		-	-	-	-	-	-	-	-	-	-	-	-			
70-74 years	-	-	-	-	-	-	-	-	-	_	-	-	-	-	-			
75-79 years	_	_	-	_	-	_	-	-	-	-	_	-	_	-	-			
30-84 years	-	_	-	-	-	_	-	_	_	_	-	-	-	-	-			
35-89 years	-	_	-	-	-	_	-	_	_	_	-	-	-	-	-			
90-94 years	-	_	-	-	-	_	-	_	_	_	-	-	-	-	-			
95-99 years	1	_	1	1	-	1	-	_	_	_	-	-	-	-	-			
100 years & over.	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_			

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Anxiety disorder, unspecified (F41.9) All ages..... 18 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 2 45-49 years..... 50-54 years..... 1 1 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 3 90-94 years..... 3 1 2 2 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Obsessive-compulsive disorder (F42) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Obsessive-compulsive disorder, unspecified (F42.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Reaction to severe stress, and adjustment disorders (F43) All ages.... 70 29 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 3 25-29 years..... 30-34 years..... 1 35-39 years..... 3 40-44 years..... 45-49 years..... 2 1 1 50-54 years..... 1 1 55-59 years..... 8 6 2 60-64 years..... 6 5 1 65-69 vears..... 4 70-74 vears..... 6 3 3 75-79 years..... 4 80-84 years..... 14 11 3 14 11 85-89 years..... 12 4 8 12 7 90-94 years..... 3 4 95-99 years..... 3 3 3 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Acute stress reaction (F43.0) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Posttraumatic stress disorder (F43.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 4 60-64 years..... 3 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 2 85-89 years..... 90-94 years..... 1 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Adjustment disorders (F43.2) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 3 95-99 years..... 1 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other reactions to severe stress (F43.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Reaction to severe stress, unspecified (F43.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 2 1 1 50-54 years..... 1 1 55-59 years..... 2 1 1 60-64 years..... 3 2 1 65-69 vears..... 3 70-74 vears..... 5 3 75-79 years..... 2 80-84 years..... 11 3 7 85-89 years..... 6 90-94 years..... 3 3 2 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Dissociative [conversion] disorders (F44) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other dissociative [conversion] disorders (F44.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Dissociative [conversion] disorder, unspecified (F44.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Somatoform disorders (F45) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Somatoform autonomic dysfunction (F45.3) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other neurotic disorders (F48) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 3 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Neurasthenia (F48.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other specified neurotic disorders (F48.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Neurotic disorder, unspecified (F48.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Eating disorders (F50) All ages.... 129 20 109 120 19 101 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 4 50-54 years..... 2 2 2 2 55-59 years..... 10 10 10 10 60-64 years..... 5 4 5 65-69 vears..... 2 70-74 vears..... 3 3 75-79 years..... 5 3 80-84 years..... 5 5 1 16 85-89 years..... 6 10 14 8 1 90-94 years..... 2.0 5 15 19 15 1 95-99 years..... 15 3 12 15 12 100 years & over. 4 3 4 3

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Anorexia nervosa (F50.0) All ages..... 38 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 2 50-54 years..... 55-59 years..... 8 8 60-64 years..... 3 3 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Bulimia nervosa (F50.2) All ages..... 11 11 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Other eating disorders (F50.8) All ages..... 45 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 1 2 60-64 years..... 1 65-69 vears..... 70-74 vears..... 2 75-79 years..... 3 80-84 years..... 2 5 1 16 85-89 years..... 10 14 8 1 19 90-94 years..... 5 14 18 14 95-99 years..... 14 2 12 14 12 100 years & over. 3 3 2 2 Age not stated...

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Eating disorder, unspecified (F50.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Sexual dysfunction, not caused by organic disorder or disease (F52) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Sexual aversion and lack of sexual enjoyment (F52.1) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Bot.h Both and age Male Female Male Female Male Female sexes Male Female sexes Male Female sexes sexes sexes Psychological and behavioral factors associated with disorders or diseases classified elsewhere (F54) All ages.... 15 Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 3 3 5 3 50-54 years..... 2 2 55-59 years..... 8 3 5 60-64 years..... 3 3 3 65-69 vears..... 11 70-74 vears..... 17 6 13 3 75-79 years..... 2.2 18 18 15 3 80-84 years..... 46 23 23 39 19 20 6 3 32 28 85-89 years..... 16 16 14 14 2 90-94 years..... 2.4 6 18 2.2 17 2 1 5 1 95-99 years..... 7 2 5 5 100 years & over. 1 1 1 1

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Abuse of non-dependence-producing substances (F55) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Specific personality disorders (F60) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Anankastic personality disorder (F60.5) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Habit and impulse disorders (F63) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other habit and impulse disorders (F63.8) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis					Asian or										
Tenth Revision, Secon		All races			White			Black		American Indian 1/			Pacific Islander 2		
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Mild mental retardati	on (F70)														
All ages	3	2	1	3	2	1	_	_	_	_	_	_	_	_	_
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
75-79 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
30-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	_	-	_	_	_	_	_	_	_	_	-	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Dis		Asian or													
Tenth Revision, Secon		All races		White			Black			American Indian 1/			Pacific Islander 2/		
Edition, 2004)	Both			Both			Both			Both			Both		
and age 	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male F	emale	sexes	Male	Female
Moderate mental retar	dation (F71)														
All ages	5	4	1	5	4	1	-	_	_	-	_	-	-	_	-
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5-9 years	-	-	-	-	-	-	-	_	-	-	-	-	-	-	-
10-14 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
15-19 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20-24 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
35-39 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
40-44 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
45-49 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
50-54 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
60-64 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
65-69 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-
70-74 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
75-79 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-
80-84 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
85-89 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
00-94 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
95-99 years	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
100 years & over.	-	-	-	_	-	-	-	-	-	-	-	-	-	-	-
Age not stated	_	-	_	_	_	_	_	_	_	_	_	_	_	_	_

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Severe mental retardation (F72) All ages.... 37 43 72 33 39 Under 1 year.... 1-4 years..... 2 5-9 years..... 10-14 years..... 3 1 15-19 years..... 2 3 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 5 45-49 years..... 2 3 9 50-54 years..... 6 3 55-59 years..... 9 4 5 60-64 years..... 1 6 1 65-69 vears..... 4 70-74 vears..... 6 75-79 years..... 3 80-84 years..... 2 1 1 85-89 years..... 2 2 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Profound mental retardation (F73) All ages..... 30 Under 1 year.... 1-4 years..... 1 5-9 years..... 10-14 years..... 15-19 years..... 1 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 0 45-49 years..... 1 0 3 7 50-54 years..... 5 55-59 years..... 9 6 60-64 years..... 6 3 3 65-69 vears..... 4 70-74 vears..... 75-79 years..... 80-84 years..... 2 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Unspecified mental retardation (F79) All ages.... 183 132 267 158 109 45 Under 1 year.... 1-4 years..... 2 3 5-9 years..... 1 10-14 years..... 2 2 2 2 15-19 years..... 4 20-24 years..... 25-29 years..... 10 30-34 years..... 10 9 35-39 years..... 19 10 9 13 40-44 years..... 25 17 8 23 45-49 years..... 31 2.0 11 23 15 31 18 50-54 years..... 38 21 17 13 3 55-59 years..... 37 28 9 29 25 8 4 60-64 years..... 34 23 11 32 22 10 65-69 years..... 21 12 9 20 11 9 70-74 years..... 15 8 7 11 75-79 years..... 2.1 10 21 1.0 11 11 80-84 years..... 22 12 10 19 9 10 1 85-89 years..... 11 3 8 10 8 1 90-94 years..... 1 1 1 95-99 years..... 2 2 2 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Specific developmental disorders of scholastic skills (F81) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 2 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Specific spelling disorder (F81.1) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Developmental disorder of scholastic skills, unspecified (F81.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

on the International Classification of Diseases, Tenth Revision, Second All races												Asian or						
		All races		Dath	White		Dath	Black		American Indian 1/			Pacific Islander 2,					
Edition, 2004) and age	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male F	emale	Both sexes Male		Female			
Specific developmenta	ıl disorder o	of motor f	unction (F82	2)														
All ages	6	4	2	6	4	2	_	_	-	_	_	-	-	_	_			
Under 1 year	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
1-4 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
5-9 years	3	2	1	3	2	1	-	-	-	-	-	-	-	-	-			
10-14 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
l5-19 years	1	1	-	1	1	-	-	_	-	-	-	-	-	-	-			
20-24 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
25-29 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
30-34 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
35-39 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
10-44 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
15-49 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
50-54 years	1	-	1	1	-	1	-	-	-	-	-	-	-	-	-			
55-59 years	1	1	-	1	1	-	-	-	-	-	-	-	-	-	-			
60-64 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
65-69 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
70-74 years	_	-	-	_	-	-	-	-	-	-	-	-	-	-	-			
75-79 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
0-84 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
85-89 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
90-94 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
95-99 years	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
100 years & over.																		

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Pervasive developmental disorders (F84) All ages.... 30 27 Under 1 year.... 1-4 years..... 3 3 5-9 years..... 10-14 years..... 4 3 15-19 years..... 6 20-24 years..... 10 25-29 years..... 30-34 years..... 1 35-39 years..... 40-44 years..... 45-49 years..... 1 50-54 years..... 55-59 years..... 1 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Childhood autism (F84.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 1 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Rett's syndrome (F84.2) All ages.... 27 27 Under 1 year.... 1-4 years..... 3 3 5-9 years..... 10-14 years..... 3 3 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other childhood disintegrative disorder (F84.3) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Other pervasive developmental disorders (F84.8) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Unspecified disorder of psychological development (F89) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 1 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Hyperkinetic disorders (F90) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Disturbance of activity and attention (F90.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Hyperkinetic disorder, unspecified (F90.9) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Conduct disorders (F91) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Conduct disorder, unspecified (F91.9) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 60-64 years..... 1 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 1 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Female Male Female Male Female Male Female Male Female sexes Male sexes sexes sexes sexes Mental disorder, not otherwise specified (F99) All ages..... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 55-59 years..... 1 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 1 80-84 years..... 85-89 years..... 90-94 years..... 2 1 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White Tenth Revision, Second All races Black American Indian 1/ Pacific Islander 2/ Edition, 2004) Both Both Both Both Bot.h and age Male Female Male Female Male Female Male Female sexes Male Female sexes sexes sexes sexes Bacterial meningitis, not elsewhere classified (G00) All ages.... Under 1 year.... 2.0 2.8 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... Ω 45-49 years..... a 50-54 years..... 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.

Worktable I. Deaths from each cause, by 5-year age groups, race, and sex: United States, 2006

Age not stated...

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data—mortality"]

Cause of death (based on the International Classification of Diseases, Asian or White American Indian 1/ Tenth Revision, Second All races Black Pacific Islander 2/ Edition, 2004) Both Both Both Both Both and age Male Female Male Female Male Female Male Female Male Female sexes sexes sexes sexes sexes Hemophilus meningitis (G00.0) All ages.... Under 1 year.... 1-4 years..... 5-9 years..... 10-14 years..... 15-19 years..... 20-24 years..... 25-29 years..... 30-34 years..... 35-39 years..... 40-44 years..... 45-49 years..... 50-54 years..... 1 55-59 years..... 60-64 years..... 65-69 vears..... 70-74 vears..... 75-79 years..... 80-84 years..... 85-89 years..... 90-94 years..... 95-99 years..... 100 years & over.