Person with disabilities experiencing problems accessing NCHS birth and death tables should contact **nchsed@cdc.gov**, or call 301-458-4688

Person with disabilities experiencing problems accessing NCHS birth and death tables should contact **nchsed@cdc.gov**, or call 301-458-4688

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
All origins 3/, both sexes												
2006	810.4	690.7	28.4	15.2	82.2	106.3	190.2	427.5	890.9	2,062.1	5,115.0	13,253.1
2005	825.9	692.5	29.4	16.3	81.4	104.4	193.3	432.0	906.9	2,137.1	5,260.0	13,798.6
2004	816.5	685.2	29.9	16.8	80.1	102.1	193.5	427.0	910.3	2,164.6	5,275.1	13,823.5
2003	841.9	700.0	31.5	17.0	81.5	103.6	201.6	433.2	940.9	2,255.0	5,463.1	14,593.3
2002	847.3	695.0	31.2	17.4	81.4	103.6	202.9	430.1	952.4	2,314.7	5,556.9	14,828.3
2001	848.5	683.4	33.3	17.3	80.7	105.2	203.6	428.9	964.6	2,353.3	5,582.4	15,112.8
2000	854.0	736.7	32.4	18.0	79.9	101.4	198.9	425.6	992.2	2,399.1	5,666.5	15,524.4
1999	857.0	736.0	34.2	18.6	79.3	102.2	198.0	418.2	1,005.0	2,457.3	5,714.5	15,554.6
All origins 3/, male												
2006	814.8	756.3	30.5	17.6	119.3	146.8	238.7	541.0	1,110.0	2,516.2	6,177.7	14,309.1
2005	827.2	762.3	33.4	18.6	117.8	143.4	243.0	547.8	1,131.0	2,612.2	6,349.8	14,889.4
2004	817.6	753.7	32.4	19.2	114.7	139.5	243.6	543.5	1,128.8	2,644.8	6,394.3	15,031.1
2003	840.3	777.4	35.1	19.8	116.5	141.4	255.0	552.2	1,165.5	2,771.7	6,641.8	15,794.0
2002	846.6	761.5	35.2	20.0	117.3	142.2	257.5	547.5	1,184.0	2,855.3	6,760.5	16,254.5
2001	846.4	749.8	37.0	19.8	117.0	143.7	259.6	545.1	1,192.7	2,911.5	6,833.0	16,744.8
2000	853.0	806.5	35.9	20.9	114.9	138.6	255.2	542.8	1,230.7	2,979.6	6,972.6	17,501.4
1999	859.2	805.0	37.9	21.5	113.1	139.7	254.9	533.1	1,252.0	3,073.7	7,083.3	17,597.2
All origins 3/, female												
2006	806.1	622.0	26.3	12.8	42.8	64.3	141.6	317.7	687.0	1,677.9	4,388.3	12,759.0
2005	824.6	619.4	25.1	13.9	42.7	64.1	143.6	319.9	698.5	1,736.3	4,520.0	13,297.7
2004	815.4	613.4	27.3	14.3	43.6	63.5	143.5	314.3	707.4	1,761.4	4,521.8	13,280.3
2003	843.4	619.1	27.8	14.0	44.4	64.6	148.5	318.4	732.7	1,823.0	4,675.5	14,062.5
2002	848.0	625.3	27.0	14.7	43.7	64.0	148.8	316.9	738.0	1,864.7	4,757.9	14,209.6
2001	850.4	613.9	29.5	14.6	42.6	66.0	148.2	316.8	754.0	1,890.8	4,760.5	14,429.9
2000	855.0	663.4	28.7	15.0	43.1	63.5	143.2	312.5	772.2	1,921.2	4,814.7	14,719.2
1999	854.9	663.6	30.3	15.6	43.7	64.1	141.8	307.6	777.6	1,952.3	4,825.4	14,731.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
Hispanic, both sexes												
2006	300.1	590.6	26.4	14.2	80.4	81.5	134.4	310.7	657.6	1,531.0	3,745.7	9,022.8
2005	307.3	614.1	28.9	13.7	81.2	81.9	138.9	317.8	676.4	1,621.4	3,977.8	9,436.7
2004	296.2	586.9	27.3	14.4	76.5	78.2	138.7	313.5	684.0	1,607.5	3,926.7	9,483.8
2003	305.8	610.1	30.2	15.7	77.1	83.9	144.9	327.8	714.3	1,685.6	4,093.5	10,431.5
2002	302.2	592.7	29.8	15.5	77.2	79.3	145.8	321.3	719.8	1,734.1	4,143.1	10,679.0
2001	306.8	572.7	30.6	14.7	75.6	84.5	154.8	334.4	722.2	1,776.0	4,262.6	11,821.8
2000	303.8	596.3	29.6	15.7	72.8	84.6	157.6	330.2	744.4	1,803.8	4,335.9	11,822.9
1999	305.7	583.8	30.9	15.6	71.3	85.5	158.2	333.2	752.5	1,853.0	4,356.3	12,177.9
Hispanic, male												
2006	323.9	640.7	28.8	16.4	120.7	112.7	176.5	403.8	843.6	1,910.7	4,492.6	9,435.5
2005	334.4	670.2	33.2	15.3	120.4	115.5	182.0	417.4	875.8	2,029.4	4,856.8	10,140.5
2004	321.1	636.5	30.2	16.7	115.3	109.2	184.2	417.8	874.0	1,994.3	4,791.6	9,932.8
2003	330.7	665.5	33.8	18.4	112.9	118.1	189.8	436.4	920.8	2,115.7	4,948.5	11,029.2
2002	328.7	644.0	34.2	17.4	114.4	112.5	192.5	423.4	937.4	2,193.4	5,043.5	11,674.1
2001	332.9	624.4	33.8	16.6	111.5	118.0	208.5	443.9	923.9	2,242.6	5,258.0	12,888.3
2000	331.3	637.1	31.5	17.9	107.7	120.2	211.0	439.0	965.7	2,287.9	5,395.3	13,086.2
1999	332.6	623.4	32.9	17.8	104.1	120.6	215.1	444.4	974.8	2,368.9	5,379.2	13,485.9
Hispanic, female												
2006	274.6	538.3	24.0	11.8	35.2	43.1	87.1	215.3	486.5	1,222.7	3,222.9	8,803.5
2005	278.2	555.4	24.5	12.0	36.6	41.1	90.6	216.4	493.9	1,291.6	3,365.8	9,068.4
2004	269.7	535.1	24.3	12.0	31.8	40.9	88.2	208.3	511.3	1,297.2	3,329.6	9,253.0
2003	279.3	552.4	26.3	12.8	35.5	43.0	95.5	219.1	527.9	1,342.9	3,506.4	10,128.1
2002	274.0	539.1	25.3	13.5	34.1	40.0	94.9	219.8	524.3	1,368.7	3,526.4	10,186.0
2001	279.0	518.9	27.2	12.7	33.7	45.2	97.0	226.7	543.0	1,408.0	3,589.8	11,300.5
2000	274.6	553.6	27.5	13.4	31.7	43.4	100.5	223.8	548.4	1,423.2	3,624.5	11,202.8
1999	277.2	542.3	28.7	13.3	32.9	44.9	97.4	224.9	555.8	1,448.8	3,675.7	11,547.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
Non-Hispanic 4/, both sexes												
2006	897.1	714.6	28.9	15.5	82.3	112.1	199.5	439.7	908.7	2,099.0	5,192.4	13,450.1
2005	911.2	708.4	29.4	16.9	81.2	109.5	201.8	443.2	924.0	2,172.1	5,327.5	13,990.9
2004	899.4	705.6	30.5	17.2	80.6	107.3	201.7	437.4	926.1	2,200.1	5,340.3	13,997.7
2003	924.4	716.2	31.7	17.2	82.0	107.6	209.3	442.1	955.6	2,289.1	5,522.6	14,740.6
2002	928.8	713.0	31.4	17.8	81.9	108.4	210.3	438.8	967.2	2,347.0	5,610.8	14,958.7
2001	926.2	703.5	33.8	17.7	81.4	108.9	209.2	435.6	979.5	2,383.9	5,628.5	15,199.0
2000	929.6	760.4	32.9	18.4	80.9	104.1	202.9	431.8	1,006.5	2,427.9	5,707.6	15,619.8
1999	929.9	760.7	34.8	19.1	80.5	104.7	201.6	423.3	1,019.2	2,485.0	5,753.4	15,631.0
Non-Hispanic 4/, male												
2006	902.8	784.5	30.8	17.8	118.6	155.7	249.7	555.3	1,129.5	2,555.7	6,274.0	14,558.8
2005	912.6	781.7	33.3	19.4	116.9	150.2	252.9	560.5	1,148.7	2,649.4	6,428.1	15,121.2
2004	900.9	778.7	32.9	19.6	114.2	146.6	252.8	554.7	1,145.4	2,683.5	6,472.4	15,260.4
2003	922.9	798.1	35.2	20.0	116.7	146.5	264.2	561.6	1,179.8	2,808.1	6,715.9	15,984.2
2002	928.0	782.6	35.4	20.4	117.3	148.6	266.0	557.0	1,198.1	2,889.1	6,825.9	16,418.0
2001	923.6	773.2	37.6	20.4	117.5	148.5	265.2	551.6	1,207.0	2,944.1	6,888.0	16,866.3
2000	928.1	836.8	36.8	21.5	115.9	141.6	259.2	548.7	1,244.2	3,009.8	7,020.7	17,639.8
1999	932.2	834.9	38.8	22.1	114.6	142.7	258.1	537.3	1,265.6	3,102.8	7,133.0	17,710.4
Non-Hispanic 4/, female												
2006	891.7	641.4	26.9	13.0	44.3	68.8	150.3	328.5	703.2	1,711.4	4,453.8	12,934.8
2005	910.0	631.6	25.3	14.4	43.8	68.9	151.6	330.3	714.6	1,768.0	4,581.2	13,475.2
2004	898.0	628.9	28.1	14.7	45.7	68.1	151.5	324.5	722.2	1,792.8	4,579.4	13,433.1
2003	925.9	630.7	28.0	14.3	46.0	68.9	155.6	327.3	747.2	1,853.7	4,726.4	14,194.2
2002	929.5	640.1	27.2	15.0	45.3	68.5	155.7	325.2	753.0	1,894.4	4,805.4	14,329.1
2001	928.6	630.4	29.9	14.9	44.1	69.6	154.3	324.0	768.7	1,918.4	4,801.8	14,505.6
2000	931.0	680.1	28.8	15.2	45.0	66.8	147.7	319.2	786.8	1,947.5	4,852.4	14,802.9
1999	927.8	682.9	30.6	16.0	45.4	67.1	146.3	313.7	791.7	1,977.5	4,858.2	14,798.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
Non-Hispanic white, both sexes												
2006 5/	968.5	564.2	25.0	14.0	76.1	102.1	184.8	406.6	861.2	2,055.0	5,205.2	13,688.2
2005 5/	981.8	562.6	26.2	15.2	74.8	98.4	186.7	407.1	873.5	2,126.0	5,330.0	14,222.4
2004 5/	967.8	564.4	26.8	15.5	75.2	95.9	185.9	400.8	875.1	2,152.5	5,337.1	14,191.1
2003 5/	993.6	576.5	27.6	15.6	75.8	95.8	190.9	402.3	902.4	2,237.2	5,515.4	14,944.8
2002	997.5	575.9	27.1	16.0	75.8	95.9	191.6	400.1	912.8	2,296.1	5,596.0	15,149.1
2001	991.1	555.4	30.1	16.3	74.1	96.4	188.3	396.0	924.0	2,325.5	5,604.4	15,371.2
2000	993.2	596.5	28.5	17.1	73.6	90.1	181.0	390.5	950.2	2,366.2	5,675.4	15,799.7
1999	990.7	593.1	30.1	17.3	72.4	91.3	177.8	381.5	957.8	2,420.0	5,716.0	15,795.3
Non-Hispanic white, male												
2006 5/	962.0	621.9	26.7	16.2	107.6	141.1	233.1	515.1	1,064.0	2,490.3	6,278.3	14,841.1
2005 5/	970.6	625.7	29.9	17.4	105.8	134.1	236.1	517.2	1,079.6	2,584.5	6,420.4	15,401.3
2004 5/	957.4	625.2	29.0	17.9	104.8	130.9	235.7	510.5	1,076.4	2,617.9	6,461.5	15,489.2
2003 5/	979.1	647.2	30.2	18.1	105.9	129.9	243.8	513.2	1,110.5	2,738.5	6,692.2	16,234.4
2002	983.9	643.5	30.3	18.3	106.7	130.9	244.9	509.9	1,126.5	2,824.1	6,801.7	16,641.9
2001	975.6	611.6	33.7	18.6	105.1	131.5	241.6	502.6	1,136.3	2,869.4	6,851.5	17,055.3
2000	978.5	658.7	32.4	20.0	103.5	123.0	233.9	497.7	1,170.9	2,930.5	6,977.8	17,853.2
1999	979.6	658.1	33.4	19.9	100.8	124.5	230.0	483.7	1,187.4	3,023.2	7,088.0	17,871.2
Non-Hispanic white, female												
2006 5/	974.7	503.7	23.2	11.8	42.9	62.5	136.3	299.8	668.0	1,677.4	4,460.7	13,150.7
2005 5/	992.6	496.5	22.2	12.9	42.2	62.1	137.0	298.7	677.2	1,729.6	4,579.7	13,683.1
2004 5/	977.7	500.7	24.4	13.0	44.3	60.3	136.0	292.9	683.8	1,752.0	4,571.1	13,609.6
2003 5/	1,007.6	502.5	24.9	12.9	44.3	61.2	138.0	293.1	705.0	1,807.9	4,720.5	14,377.2
2002	1,010.6	504.8	23.8	13.6	43.8	60.3	138.3	292.1	710.5	1,846.0	4,787.9	14,504.3
2001	1,006.1	496.4	26.3	13.9	41.9	60.9	134.9	291.0	723.5	1,864.1	4,777.3	14,670.6
2000	1,007.3	530.9	24.4	13.9	42.6	56.8	128.1	285.0	742.1	1,891.0	4,819.3	14,971.7
1999	1,001.3	524.6	26.7	14.7	42.9	57.8	125.6	281.0	741.6	1,915.1	4,817.7	14,967.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
Non-Hispanic black, both sexes												
2006 5/	759.1	1,339.2	44.3	21.9	115.6	185.7	325.2	738.5	1,472.4	2,887.7	5,980.8	12,655.4
2005 5/	774.4	1,323.9	42.5	24.0	115.8	186.0	328.5	761.4	1,513.6	2,979.6	6,218.3	13,239.1
2004 5/	768.8	1,297.2	45.5	24.4	112.8	186.5	332.3	760.8	1,527.9	3,032.3	6,251.7	13,492.2
2003 5/	788.8	1,290.0	48.0	23.8	116.6	188.9	354.2	788.3	1,578.4	3,158.7	6,433.0	14,107.7
2002	792.8	1,280.3	48.4	25.3	116.9	193.9	357.8	782.0	1,592.9	3,204.8	6,584.8	14,386.7
2001	798.1	1,354.4	49.5	24.0	121.4	191.5	365.8	786.9	1,602.5	3,300.7	6,645.3	14,660.8
2000	805.5	1,477.8	51.7	25.0	123.2	194.6	365.1	802.2	1,633.9	3,378.8	6,764.3	14,837.5
1999	812.1	1,510.2	54.6	27.5	125.5	191.1	376.9	796.7	1,681.7	3,454.5	6,805.4	14,839.2
Non-Hispanic black, male												
2006 5/	815.3	1,453.3	48.2	25.8	176.9	266.0	405.6	944.2	1,935.2	3,736.8	7,520.4	13,403.1
2005 5/	825.7	1,451.5	47.6	27.8	177.7	264.0	407.5	969.9	1,993.8	3,814.1	7,771.1	13,978.1
2004 5/	818.7	1,427.6	49.5	26.9	169.4	262.1	409.3	974.9	1,998.6	3,880.2	7,813.2	14,647.3
2003 5/	840.6	1,427.7	55.1	27.9	176.5	266.6	440.0	1,012.8	2,047.7	4,041.1	8,165.7	15,082.2
2002	842.3	1,369.3	55.8	29.8	177.4	274.3	445.7	1,002.9	2,074.5	4,078.9	8,246.2	15,776.9
2001	849.7	1,484.2	55.1	28.6	186.2	267.9	456.3	1,014.6	2,055.4	4,218.9	8,426.8	16,576.2
2000	859.5	1,623.2	56.1	29.1	187.2	270.2	462.7	1,036.2	2,111.4	4,303.9	8,551.0	16,944.8
1999	872.8	1,650.8	61.5	33.1	191.4	267.9	479.5	1,048.1	2,176.7	4,397.8	8,615.3	17,385.3
Non-Hispanic black, female												
2006 5/	708.0	1,220.1	40.3	17.9	52.8	111.4	254.1	562.2	1,100.2	2,281.4	5,107.2	12,350.5
2005 5/	727.6	1,190.8	37.2	20.0	52.6	114.1	258.5	582.4	1,127.1	2,383.1	5,338.1	12,941.6
2004 5/	723.4	1,161.6	41.3	21.8	55.3	117.2	264.1	577.0	1,149.9	2,427.9	5,370.3	13,035.6
2003 5/	741.6	1,146.9	40.7	19.6	56.0	118.0	278.4	595.7	1,201.6	2,531.2	5,454.6	13,730.3
2002	748.0	1,187.5	40.8	20.7	56.1	121.3	280.1	592.7	1,206.3	2,584.8	5,644.9	13,855.4
2001	751.2	1,217.8	43.6	19.3	56.8	122.4	285.7	591.5	1,238.2	2,653.0	5,645.1	13,951.4
2000	756.7	1,327.7	47.1	20.7	60.1	126.5	278.8	601.2	1,249.4	2,728.0	5,758.6	14,054.9
1999	757.3	1,365.4	47.4	21.8	60.5	122.1	286.4	581.7	1,285.1	2,789.2	5,792.0	13,887.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
All origins 3/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
Non-Hispanic 4/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
Non-Hispanic white, both sexes												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
All origins 3/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
Non-Hispanic 4/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
Non-Hispanic white, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)												
All origins 3/, both sexes												
2006	2.2	*	*	*	*	*	0.1	0.3	1.1	4.8	19.0	50.5
2005	1.9	*	*	*	*	*	0.1	0.2	0.8	4.4	15.9	45.8
2004	1.5	*	*	*	*	*	0.1	0.2	0.9	3.3	12.8	34.3
2003	1.1	*	*	*	*	*	*	0.2	0.6	2.4	9.3	24.8
2002	0.9	*	*	*	*	*	*	0.1	0.5	2.0	7.0	21.0
2001	0.5	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.8
2000	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.7	11.8
1999	0.4	*	*	*	*	*	*	0.1	0.2	1.0	3.2	10.3
All origins 3/, male												
2006	1.7	*	*	*	*	*	0.1	0.3	1.3	4.8	18.6	50.0
2005	1.5	*	*	*	*	*	*	0.3	0.9	4.4	16.4	46.1
2004	1.1	*	*	*	*	*	*	0.2	0.9	3.1	12.7	32.0
2003	0.8	*	*	*	*	*	*	0.2	0.6	2.2	9.3	21.4
2002	0.7	*	*	*	*	*	*	0.1	0.6	2.2	7.0	20.4
2001	0.4	*	*	*	*	*	*	*	0.3	1.3	4.6	12.1
2000	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	11.4
1999	0.3	*	*	*	*	*	*	*	0.2	0.9	3.5	9.5
All origins 3/, female												
2006	2.7	*	*	*	*	*	*	0.3	1.0	4.8	19.4	50.8
2005	2.3	*	*	*	*	*	*	0.2	0.8	4.5	15.5	45.6
2004	1.8	*	*	*	*	*	*	0.2	0.8	3.4	12.9	35.4
2003	1.3	*	*	*	*	*	*	0.2	0.6	2.6	9.3	26.3
2002	1.0	*	*	*	*	*	*	0.1	0.5	1.8	6.9	21.3
2001	0.7	*	*	*	*	*	*	0.1	0.3	1.3	4.5	13.1
2000	0.6	*	*	*	*	*	*	*	0.2	1.2	3.6	11.9
1999	0.5	*	*	*	*	*	*	*	0.2	1.0	3.0	10.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)												
Hispanic, both sexes												
2006	0.6	*	*	*	*	*	*	*	1.3	3.3	15.6	24.6
2005	0.4	*	*	*	*	*	*	*	*	2.6	9.5	24.4
2004	0.4	*	*	*	*	*	*	*	*	2.0	7.3	18.6
2003	0.2	*	*	*	*	*	*	*	*	*	4.8	14.5
2002	0.2	*	*	*	*	*	*	*	*	*	5.2	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	0.5	*	*	*	*	*	*	*	*	3.6	13.0	27.1
2005	0.3	*	*	*	*	*	*	*	*	*	8.4	*
2004	0.3	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.8	*	*	*	*	*	*	*	*	3.2	17.5	23.4
2005	0.6	*	*	*	*	*	*	*	*	*	10.2	26.7
2004	0.5	*	*	*	*	*	*	*	*	*	8.4	17.3
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)												
Non-Hispanic 4/, both sexes												
2006	2.5	*	*	*	*	*	0.1	0.3	1.1	4.9	19.2	51.8
2005	2.2	*	*	*	*	*	0.1	0.2	0.9	4.6	16.2	46.7
2004	1.7	*	*	*	*	*	0.1	0.2	0.9	3.4	13.1	35.0
2003	1.2	*	*	*	*	*	*	0.2	0.6	2.5	9.5	25.2
2002	1.0	*	*	*	*	*	*	0.1	0.5	2.1	7.1	21.4
2001	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.8
2000	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.8	11.9
1999	0.4	*	*	*	*	*	*	*	0.2	1.0	3.3	10.4
Non-Hispanic 4/, male												
2006	2.0	*	*	*	*	*	*	0.3	1.2	4.9	18.9	51.2
2005	1.7	*	*	*	*	*	*	0.3	1.0	4.6	16.9	47.4
2004	1.3	*	*	*	*	*	*	0.2	0.9	3.2	13.1	32.4
2003	0.9	*	*	*	*	*	*	0.2	0.6	2.3	9.5	21.4
2002	0.8	*	*	*	*	*	*	0.1	0.6	2.3	7.2	21.0
2001	0.5	*	*	*	*	*	*	*	0.3	1.3	4.7	12.0
2000	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	11.5
1999	0.3	*	*	*	*	*	*	*	0.2	0.9	3.6	9.6
Non-Hispanic 4/, female												
2006	3.0	*	*	*	*	*	*	0.3	1.0	4.9	19.5	52.1
2005	2.6	*	*	*	*	*	*	0.2	0.8	4.6	15.8	46.4
2004	2.0	*	*	*	*	*	*	0.2	0.8	3.5	13.1	36.1
2003	1.5	*	*	*	*	*	*	0.2	0.7	2.7	9.5	26.8
2002	1.1	*	*	*	*	*	*	0.1	0.5	1.8	7.0	21.6
2001	0.7	*	*	*	*	*	*	0.1	0.4	1.3	4.5	13.2
2000	0.6	*	*	*	*	*	*	*	0.2	1.2	3.6	12.1
1999	0.5	*	*	*	*	*	*	*	0.2	1.0	3.1	10.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)												
Non-Hispanic white, both sexes												
2006 5/	2.9	*	*	*	*	*	*	0.3	1.1	5.2	20.6	55.1
2005 5/	2.5	*	*	*	*	*	0.1	0.2	0.8	4.9	17.2	49.7
2004 5/	2.0	*	*	*	*	*	0.1	0.2	0.9	3.6	13.8	37.3
2003 5/	1.4	*	*	*	*	*	*	0.1	0.6	2.7	10.0	26.5
2002	1.1	*	*	*	*	*	*	0.2	0.5	2.2	7.5	22.5
2001	0.7	*	*	*	*	*	*	0.1	0.3	1.4	4.7	13.6
2000	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.0	12.3
1999	0.5	*	*	*	*	*	*	*	0.2	1.0	3.4	11.0
Non-Hispanic white, male												
2006 5/	2.3	*	*	*	*	*	*	0.3	1.2	5.1	19.9	54.4
2005 5/	2.0	*	*	*	*	*	*	0.3	1.0	4.8	17.5	50.9
2004 5/	1.5	*	*	*	*	*	*	0.2	0.9	3.3	13.8	34.5
2003 5/	1.0	*	*	*	*	*	*	0.2	0.6	2.4	10.1	22.7
2002	0.9	*	*	*	*	*	*	0.2	0.5	2.3	7.5	22.1
2001	0.5	*	*	*	*	*	*	*	0.3	1.4	4.7	12.6
2000	0.5	*	*	*	*	*	*	*	0.3	1.3	4.3	12.0
1999	0.4	*	*	*	*	*	*	*	0.2	1.0	3.8	10.0
Non-Hispanic white, female												
2006 5/	3.6	*	*	*	*	*	*	0.3	1.0	5.2	21.0	55.4
2005 5/	3.0	*	*	*	*	*	*	0.2	0.7	4.9	16.9	49.2
2004 5/	2.4	*	*	*	*	*	*	0.2	0.9	3.8	13.9	38.6
2003 5/	1.8	*	*	*	*	*	*	0.1	0.7	2.9	9.9	28.2
2002	1.3	*	*	*	*	*	*	0.1	0.5	2.0	7.5	22.7
2001	0.9	*	*	*	*	*	*	0.2	0.3	1.3	4.7	14.0
2000	0.7	*	*	*	*	*	*	*	0.3	1.2	3.9	12.5
1999	0.6	*	*	*	*	*	*	*	*	1.0	3.2	11.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)												
Non-Hispanic black, both sexes												
2006 5/	1.0	*	*	*	*	*	*	0.6	1.7	4.5	10.5	24.2
2005 5/	0.9	*	*	*	*	*	*	*	1.3	3.7	9.9	23.2
2004 5/	0.6	*	*	*	*	*	*	*	0.9	2.7	8.3	16.1
2003 5/	0.5	*	*	*	*	*	*	*	0.8	1.5	6.7	15.5
2002	0.4	*	*	*	*	*	*	*	1.1	1.9	3.7	13.3
2001	0.3	*	*	*	*	*	*	*	*	1.3	3.0	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	8.3
1999	0.2	*	*	*	*	*	*	*	*	*	2.8	*
Non-Hispanic black, male												
2006 5/	0.8	*	*	*	*	*	*	*	1.9	4.4	11.1	20.2
2005 5/	0.7	*	*	*	*	*	*	*	*	3.6	13.2	*
2004 5/	0.5	*	*	*	*	*	*	*	*	3.0	7.4	*
2003 5/	0.4	*	*	*	*	*	*	*	*	*	6.6	*
2002	0.4	*	*	*	*	*	*	*	*	2.9	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	1.2	*	*	*	*	*	*	*	1.5	4.5	10.1	25.8
2005 5/	1.0	*	*	*	*	*	*	*	1.3	3.8	8.1	25.7
2004 5/	0.7	*	*	*	*	*	*	*	*	2.4	8.9	16.1
2003 5/	0.6	*	*	*	*	*	*	*	*	*	6.7	16.9
2002	0.5	*	*	*	*	*	*	*	*	*	3.6	13.4
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
All origins 3/, both sexes												
2006	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.3	2.1
2005	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.3	2.2
2004	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.3	2.4
2003	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	1.7	2.3
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	2.6
2001	0.3	*	*	*	*	0.1	0.1	0.2	0.5	0.8	1.6	2.8
2000	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.0	2.7
1999	0.3	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	4.2
All origins 3/, male												
2006	0.3	*	*	*	*	*	0.1	0.4	0.5	0.9	1.8	3.1
2005	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	1.7	3.3
2004	0.3	*	*	*	*	*	0.2	0.3	0.4	1.1	1.6	3.8
2003	0.3	*	*	*	*	*	0.1	0.4	0.5	0.9	2.3	3.1
2002	0.3	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	4.0
2001	0.3	*	*	*	*	*	0.1	0.3	0.7	1.2	2.0	4.4
2000	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.8	4.6
1999	0.4	*	*	*	*	*	0.2	0.5	0.8	1.4	2.8	7.1
All origins 3/, female												
2006	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.9	1.6
2005	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.7
2004	0.2	*	*	*	*	*	*	0.1	*	0.4	1.0	1.7
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.9
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.1
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
2000	0.2	*	*	*	*	*	*	*	0.2	0.7	1.5	2.0
1999	0.2	*	*	*	*	*	*	*	0.3	0.6	1.6	3.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
Hispanic, both sexes												
2006	0.2	*	*	*	*	*	*	*	0.9	1.5	2.6	*
2005	0.2	*	*	*	*	*	*	*	*	*	*	*
2004	0.3	*	*	*	*	*	*	*	*	1.8	3.8	*
2003	0.2	*	*	*	*	*	*	*	*	*	3.1	*
2002	0.2	*	*	*	*	*	*	*	*	1.8	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	3.6	*
2000	0.2	*	*	*	*	*	*	*	*	*	4.5	*
1999	0.4	*	*	*	*	*	*	*	1.2	2.3	5.6	*
Hispanic, male												
2006	0.3	*	*	*	*	*	*	*	*	*	*	*
2005	0.3	*	*	*	*	*	*	*	*	*	*	*
2004	0.3	*	*	*	*	*	*	*	*	*	*	*
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
Non-Hispanic 4/, both sexes												
2006	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.0
2005	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.2
2004	0.2	*	*	*	*	*	0.1	0.2	0.2	0.7	1.1	2.2
2003	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.6	2.2
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.8	2.5
2001	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5	2.7
2000	0.3	*	*	*	*	*	0.1	0.2	0.3	0.8	1.9	2.5
1999	0.3	*	*	*	*	*	0.1	0.2	0.5	0.9	1.9	3.9
Non-Hispanic 4/, male												
2006	0.3	*	*	*	*	*	0.1	0.3	0.5	0.8	1.8	2.8
2005	0.3	*	*	*	*	*	*	0.2	0.4	0.8	1.6	3.2
2004	0.3	*	*	*	*	*	0.2	0.2	0.4	1.0	1.3	3.5
2003	0.3	*	*	*	*	*	*	0.4	0.5	0.8	2.1	3.0
2002	0.3	*	*	*	*	*	0.1	0.3	0.6	1.2	2.3	3.8
2001	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	1.9	4.3
2000	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.6	4.3
1999	0.4	*	*	*	*	*	0.2	0.4	0.8	1.3	2.6	6.8
Non-Hispanic 4/, female												
2006	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.6
2005	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.7
2004	0.2	*	*	*	*	*	*	0.1	*	0.4	1.0	1.6
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.8
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	1.9
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.1
2000	0.2	*	*	*	*	*	*	*	*	0.7	1.4	1.8
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	2.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
Non-Hispanic white, both sexes												
2006 5/	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.9	1.5
2005 5/	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.8	1.6
2004 5/	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.8	1.5
2003 5/	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.2	1.7
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.9
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.0
2000	0.2	*	*	*	*	*	*	0.1	0.1	0.5	1.2	2.0
1999	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.3	3.0
Non-Hispanic white, male												
2006 5/	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	1.7
2005 5/	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.3
2004 5/	0.2	*	*	*	*	*	*	0.1	0.2	0.5	0.9	2.4
2003 5/	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.5	2.4
2002	0.2	*	*	*	*	*	*	0.1	0.3	0.8	1.7	2.9
2001	0.2	*	*	*	*	*	*	0.1	0.4	0.6	1.4	2.9
2000	0.2	*	*	*	*	*	*	0.2	*	0.6	1.5	3.6
1999	0.3	*	*	*	*	*	*	0.2	0.4	0.8	1.8	4.9
Non-Hispanic white, female												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	0.6	1.3
2005 5/	0.1	*	*	*	*	*	*	*	*	0.3	0.6	1.3
2004 5/	0.1	*	*	*	*	*	*	*	*	*	0.7	1.1
2003 5/	0.1	*	*	*	*	*	*	*	*	0.3	1.0	1.4
2002	0.1	*	*	*	*	*	*	*	*	*	1.1	1.4
2001	0.2	*	*	*	*	*	*	*	*	0.4	0.9	1.7
2000	0.1	*	*	*	*	*	*	*	*	0.3	1.1	1.4
1999	0.2	*	*	*	*	*	*	*	*	0.3	1.0	2.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
Non-Hispanic black, both sexes												
2006 5/	0.4	*	*	*	*	*	*	0.7	1.1	2.0	2.6	*
2005 5/	0.4	*	*	*	*	*	*	0.5	0.9	1.8	2.9	*
2004 5/	0.5	*	*	*	*	*	0.4	0.7	0.9	2.4	3.1	6.9
2003 5/	0.5	*	*	*	*	*	*	1.0	1.1	1.2	3.4	*
2002	0.5	*	*	*	*	*	0.4	0.8	1.2	2.6	3.9	*
2001	0.6	*	*	*	*	*	*	0.8	1.5	2.7	3.8	6.9
2000	0.7	*	*	*	*	*	0.5	1.0	1.4	3.2	6.4	*
1999	0.7	*	*	*	*	*	0.4	1.0	1.7	2.6	5.6	8.4
Non-Hispanic black, male												
2006 5/	0.6	*	*	*	*	*	*	1.1	2.1	3.2	*	*
2005 5/	0.5	*	*	*	*	*	*	*	*	3.5	*	*
2004 5/	0.6	*	*	*	*	*	*	*	1.6	3.6	*	*
2003 5/	0.6	*	*	*	*	*	*	1.6	*	*	5.8	*
2002	0.7	*	*	*	*	*	*	1.1	2.0	3.7	6.4	*
2001	0.7	*	*	*	*	*	*	1.4	2.4	4.4	*	*
2000	0.9	*	*	*	*	*	*	1.9	2.5	4.2	11.7	*
1999	0.9	*	*	*	*	*	*	1.8	3.0	4.2	7.8	*
Non-Hispanic black, female												
2006 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	3.5	*
2000	0.4	*	*	*	*	*	*	*	*	2.5	3.5	*
1999	0.5	*	*	*	*	*	*	*	*	*	4.4	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
All origins 3/, both sexes												
2006	0.2	*	*	*	*	*	0.1	0.2	0.3	0.4	1.0	1.6
2005	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.9
2004	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	0.9	2.2
2003	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	1.4	1.8
2002	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.5	2.3
2001	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4
2000	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.7	2.3
1999	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.8	3.7
All origins 3/, male												
2006	0.2	*	*	*	*	*	0.1	0.3	0.5	0.7	1.5	2.4
2005	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.3	2.7
2004	0.2	*	*	*	*	*	0.1	0.2	0.3	0.9	1.2	3.7
2003	0.2	*	*	*	*	*	*	0.3	0.4	0.7	1.9	2.5
2002	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	1.9	3.5
2001	0.3	*	*	*	*	*	*	0.3	0.6	0.9	1.7	3.8
2000	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	2.4	4.1
1999	0.4	*	*	*	*	*	0.1	0.4	0.7	1.2	2.5	6.4
All origins 3/, female												
2006	0.1	*	*	*	*	*	*	*	*	0.2	0.6	1.2
2005	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.5
2004	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.5
2003	0.1	*	*	*	*	*	*	0.1	0.2	0.3	1.0	1.5
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.8
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.1	1.8
2000	0.2	*	*	*	*	*	*	*	*	0.5	1.2	1.6
1999	0.2	*	*	*	*	*	*	*	0.2	0.4	1.3	2.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
Hispanic, both sexes												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.2	*	*	*	*	*	*	*	*	*	3.2	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	5.0	*
Hispanic, male												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.2	*	*	*	*	*	*	*	*	*	*	*
2004	0.3	*	*	*	*	*	*	*	*	*	*	*
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
Non-Hispanic 4/, both sexes												
2006	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.5
2005	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.8
2004	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.8	2.0
2003	0.2	*	*	*	*	*	*	0.2	0.3	0.4	1.3	1.7
2002	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.5	2.2
2001	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.2	2.4
2000	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.6	2.2
1999	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.6	3.5
Non-Hispanic 4/, male												
2006	0.2	*	*	*	*	*	*	0.2	0.4	0.6	1.5	2.2
2005	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.7
2004	0.2	*	*	*	*	*	0.1	0.2	0.3	0.8	1.0	3.3
2003	0.2	*	*	*	*	*	*	0.3	0.4	0.6	1.8	2.4
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	3.3
2001	0.3	*	*	*	*	*	*	0.2	0.6	0.8	1.6	3.8
2000	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	2.2	3.8
1999	0.4	*	*	*	*	*	0.1	0.3	0.7	1.1	2.3	6.1
Non-Hispanic 4/, female												
2006	0.1	*	*	*	*	*	*	*	*	*	0.6	1.2
2005	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.4
2004	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.3
2003	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.0	1.5
2002	0.2	*	*	*	*	*	*	*	*	0.4	1.2	1.7
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.0	1.8
2000	0.2	*	*	*	*	*	*	*	*	0.4	1.2	1.5
1999	0.2	*	*	*	*	*	*	*	*	0.4	1.2	2.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
Non-Hispanic white, both sexes												
2006 5/	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.7	1.1
2005 5/	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.6	1.3
2004 5/	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.6	1.4
2003 5/	0.1	*	*	*	*	*	*	0.1	0.2	0.3	1.0	1.3
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.6
2001	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.7
2000	0.1	*	*	*	*	*	*	0.1	*	0.3	1.1	1.8
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.8
Non-Hispanic white, male												
2006 5/	0.1	*	*	*	*	*	*	*	0.2	0.3	1.0	*
2005 5/	0.1	*	*	*	*	*	*	*	0.2	0.3	0.7	1.8
2004 5/	0.1	*	*	*	*	*	*	*	*	0.5	0.7	2.2
2003 5/	0.2	*	*	*	*	*	*	0.1	0.3	0.4	1.2	1.8
2002	0.2	*	*	*	*	*	*	*	0.2	0.6	1.2	2.7
2001	0.2	*	*	*	*	*	*	*	0.4	0.4	1.1	2.4
2000	0.2	*	*	*	*	*	*	0.2	*	0.5	1.3	3.2
1999	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.6	4.8
Non-Hispanic white, female												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	0.5	1.0
2005 5/	0.1	*	*	*	*	*	*	*	*	*	0.6	1.0
2004 5/	0.1	*	*	*	*	*	*	*	*	*	0.5	1.0
2003 5/	0.1	*	*	*	*	*	*	*	*	*	0.8	1.1
2002	0.1	*	*	*	*	*	*	*	*	*	0.9	1.2
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.9	1.2
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
Non-Hispanic black, both sexes												
2006 5/	0.3	*	*	*	*	*	*	0.5	1.0	1.5	*	*
2005 5/	0.3	*	*	*	*	*	*	*	*	1.3	2.0	*
2004 5/	0.4	*	*	*	*	*	*	0.5	*	1.8	2.2	6.3
2003 5/	0.4	*	*	*	*	*	*	0.9	0.9	*	2.7	*
2002	0.4	*	*	*	*	*	*	0.6	1.0	1.9	3.6	*
2001	0.4	*	*	*	*	*	*	0.6	1.2	2.0	3.0	6.2
2000	0.5	*	*	*	*	*	0.4	0.8	0.9	2.7	5.1	*
1999	0.6	*	*	*	*	*	*	0.7	1.5	2.0	4.8	7.1
Non-Hispanic black, male												
2006 5/	0.5	*	*	*	*	*	*	*	1.9	*	*	*
2005 5/	0.4	*	*	*	*	*	*	*	*	2.8	*	*
2004 5/	0.5	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.5	*	*	*	*	*	*	1.4	*	*	*	*
2002	0.6	*	*	*	*	*	*	1.0	1.8	*	6.2	*
2001	0.6	*	*	*	*	*	*	1.2	2.1	3.4	*	*
2000	0.8	*	*	*	*	*	*	1.5	2.1	3.9	9.3	*
1999	0.7	*	*	*	*	*	*	1.3	2.7	3.2	6.9	*
Non-Hispanic black, female												
2006 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	3.7	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
All origins 3/, both sexes												
2006	0.1	*	*	*	*	*	*	0.1	*	0.2	0.3	0.5
2005	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	*
2004	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.3	*
2003	0.1	*	*	*	*	*	*	*	0.1	0.1	0.3	0.4
2002	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	*
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.3	*
2000	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	*
1999	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5
All origins 3/, male												
2006	0.1	*	*	*	*	*	*	0.1	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2004	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.5	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	*
1999	0.1	*	*	*	*	*	*	0.1	*	*	*	*
All origins 3/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2003	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
Hispanic, both sexes												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
Non-Hispanic 4/, both sexes												
2006	0.1	*	*	*	*	*	*	0.1	*	0.2	0.3	0.5
2005	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	*
2004	0.0	*	*	*	*	*	*	*	*	0.1	0.3	*
2003	0.1	*	*	*	*	*	*	*	*	0.1	0.3	*
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
2001	0.1	*	*	*	*	*	*	0.1	*	0.2	0.3	*
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
Non-Hispanic 4/, male												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.5	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2003	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.1	*	*	*	*	*	*	*	*	0.2	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
Non-Hispanic white, both sexes												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	0.1	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Non-Hispanic white, male												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
Non-Hispanic black, both sexes												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
All origins 3/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	0.7	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
Non-Hispanic 4/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
Non-Hispanic white, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
All origins 3/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
Non-Hispanic 4/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
Non-Hispanic white, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
All origins 3/, both sexes												
2006	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2005	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2004	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2003	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002	0.1	*	*	0.0	0.1	*	*	*	*	*	*	*
2001	0.1	*	*	0.1	0.1	*	*	*	*	*	*	*
2000	0.1	0.6	*	*	0.1	*	*	0.1	*	*	*	*
1999	0.1	0.8	0.2	0.1	0.1	0.1	0.1	*	*	*	*	*
All origins 3/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999	0.1	*	*	*	0.1	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
Non-Hispanic 4/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2004	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2003	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2001	0.1	*	*	0.1	0.2	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.1	*	*	0.1	*	*	*	*
1999	0.1	0.8	0.2	0.1	0.1	*	0.1	*	*	*	*	*
Non-Hispanic 4/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
Non-Hispanic white, both sexes												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2003 5/	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2002	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999	0.1	1.0	0.3	*	0.1	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
Non-Hispanic black, both sexes												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
All origins 3/, both sexes												
2006	11.4	6.5	0.5	0.2	0.3	0.7	2.0	5.2	12.8	32.1	82.4	177.3
2005	11.5	7.4	0.5	0.2	0.4	0.8	1.9	5.2	12.9	32.6	81.4	187.3
2004	11.4	6.6	0.5	0.2	0.3	0.8	1.9	5.4	12.9	32.4	81.6	186.7
2003	11.7	6.9	0.5	0.2	0.4	0.8	2.1	5.3	13.1	32.6	85.0	202.5
2002	11.7	7.3	0.5	0.2	0.3	0.8	1.9	5.2	12.6	34.7	86.5	203.0
2001	11.3	7.7	0.7	0.2	0.3	0.7	1.8	5.0	12.3	32.8	82.3	205.9
2000	11.1	7.2	0.6	0.2	0.3	0.7	1.9	4.9	11.9	31.0	80.4	215.7
1999	11.0	7.5	0.6	0.2	0.3	0.7	1.8	4.6	11.4	31.2	79.4	220.7
All origins 3/, male												
2006	10.5	6.8	0.5	0.3	0.3	0.7	2.1	5.9	14.1	34.8	92.7	194.8
2005	10.5	8.1	0.6	0.2	0.4	0.7	1.9	5.6	14.2	36.3	91.4	201.9
2004	10.4	8.0	0.5	0.1	0.3	0.8	2.0	5.9	13.8	35.8	93.3	203.9
2003	10.5	7.4	0.7	0.2	0.4	0.8	2.1	5.8	14.1	36.2	95.0	213.8
2002	10.6	7.7	0.5	0.2	0.3	0.8	2.0	5.7	13.7	38.0	99.0	213.9
2001	10.2	8.4	0.8	0.2	0.4	0.7	2.0	5.4	13.2	36.7	95.5	224.3
2000	9.8	7.1	0.7	0.2	0.3	0.8	2.0	5.6	13.0	34.5	91.9	223.0
1999	9.8	9.0	0.6	0.2	0.3	0.7	2.1	5.1	12.4	35.2	91.9	233.5
All origins 3/, female												
2006	12.3	6.2	0.6	0.2	0.3	0.7	1.9	4.6	11.5	29.7	75.3	169.2
2005	12.5	6.5	0.5	0.2	0.3	0.8	1.9	4.8	11.7	29.5	74.6	180.6
2004	12.3	5.2	0.6	0.2	0.3	0.8	1.9	4.9	12.0	29.6	73.7	179.0
2003	12.9	6.5	0.4	0.2	0.4	0.8	2.0	4.8	12.2	29.5	78.3	197.5
2002	12.9	7.0	0.5	0.2	0.3	0.8	1.8	4.7	11.7	31.9	78.2	198.3
2001	12.4	7.0	0.6	0.1	0.3	0.7	1.7	4.7	11.4	29.5	73.7	198.3
2000	12.3	7.3	0.6	0.1	0.2	0.6	1.9	4.2	11.0	28.1	72.9	212.7
1999	12.2	5.8	0.5	0.2	0.3	0.6	1.6	4.1	10.5	28.0	71.3	215.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
Hispanic, both sexes												
2006	4.1	6.9	*	*	0.3	0.5	1.2	4.0	9.9	24.9	65.1	122.9
2005	4.0	7.1	*	*	*	0.5	1.2	3.7	9.7	25.5	61.0	134.8
2004	4.0	6.1	0.7	*	*	0.5	1.4	4.0	9.7	25.1	63.8	138.0
2003	3.8	5.8	*	*	*	0.5	1.5	3.9	9.1	22.1	62.0	151.8
2002	3.9	7.8	*	*	*	0.5	1.2	3.7	8.4	28.9	63.7	156.3
2001	3.7	6.1	*	*	*	0.3	1.3	3.9	8.2	25.4	62.4	171.3
2000	3.5	8.0	*	*	*	0.5	1.5	3.9	8.5	21.8	59.3	144.7
1999	3.6	5.7	*	*	*	0.4	1.5	3.1	8.0	25.6	61.7	180.3
Hispanic, male												
2006	4.1	7.5	*	*	*	0.5	1.2	4.9	11.3	30.5	76.5	138.7
2005	3.8	7.8	*	*	*	0.5	1.0	4.3	10.6	29.6	74.2	137.0
2004	3.8	7.8	*	*	*	*	1.6	5.2	10.3	27.2	69.7	147.1
2003	3.5	6.6	*	*	*	0.5	1.5	4.9	9.6	25.3	67.1	160.1
2002	3.7	8.0	*	*	*	0.6	1.6	5.2	9.0	30.8	72.2	164.0
2001	3.6	7.0	*	*	*	*	1.5	4.7	9.9	29.8	76.0	160.0
2000	3.4	8.4	*	*	*	0.6	1.7	5.0	10.0	26.6	70.3	157.2
1999	3.5	7.0	*	*	*	*	2.0	3.8	10.2	29.4	75.1	181.2
Hispanic, female												
2006	4.2	6.3	*	*	*	*	1.1	3.0	8.5	20.3	57.1	114.4
2005	4.2	6.4	*	*	*	*	1.3	3.0	8.8	22.2	51.9	133.6
2004	4.2	*	*	*	*	*	1.2	2.8	9.2	23.4	59.8	133.2
2003	4.1	5.0	*	*	*	*	1.3	2.8	8.7	19.6	58.5	147.7
2002	4.1	7.6	*	*	*	*	0.7	2.3	7.8	27.4	57.9	152.5
2001	3.9	5.2	*	*	*	*	1.1	3.1	6.6	21.9	53.2	176.9
2000	3.5	7.7	*	*	*	*	1.3	2.8	7.2	18.1	51.8	138.5
1999	3.7	*	*	*	*	*	1.0	2.4	6.1	22.5	52.7	179.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
Non-Hispanic 4/, both sexes												
2006	12.7	6.3	0.6	0.2	0.3	0.8	2.1	5.4	13.0	32.6	83.3	179.7
2005	12.8	7.4	0.6	0.2	0.4	0.8	2.0	5.4	13.1	33.0	82.5	189.6
2004	12.5	6.7	0.5	0.2	0.4	0.9	2.0	5.5	13.1	32.9	82.5	188.7
2003	12.9	7.2	0.6	0.2	0.4	0.8	2.1	5.4	13.3	33.2	86.0	204.1
2002	12.9	7.1	0.5	0.2	0.3	0.8	2.0	5.3	12.9	34.9	87.3	204.2
2001	12.4	8.1	0.7	0.1	0.4	0.8	1.9	5.1	12.5	33.1	83.0	206.8
2000	12.1	6.9	0.7	0.2	0.3	0.7	2.0	5.0	12.1	31.4	81.0	217.5
1999	12.0	7.9	0.6	0.2	0.3	0.7	1.8	4.7	11.6	31.4	79.8	221.2
Non-Hispanic 4/, male												
2006	11.7	6.5	0.6	0.3	0.3	0.8	2.2	5.9	14.3	35.1	93.7	197.5
2005	11.7	8.2	0.6	0.3	0.4	0.8	2.0	5.8	14.4	36.6	92.2	205.3
2004	11.5	7.9	0.4	0.1	0.4	0.9	2.0	5.9	14.0	36.3	94.6	206.3
2003	11.6	7.6	0.8	0.2	0.4	0.9	2.1	5.8	14.3	36.8	96.2	215.9
2002	11.6	7.3	0.5	0.3	0.3	0.9	2.0	5.7	14.0	38.2	99.8	215.6
2001	11.2	8.8	0.9	0.1	0.4	0.8	2.1	5.5	13.4	37.0	96.1	226.8
2000	10.7	6.7	0.7	0.2	0.3	0.8	2.0	5.6	13.1	34.8	92.3	224.8
1999	10.7	9.4	0.6	0.2	0.4	0.8	2.1	5.2	12.5	35.4	92.2	234.5
Non-Hispanic 4/, female												
2006	13.6	6.2	0.5	0.2	0.4	0.7	2.0	4.8	11.8	30.4	76.2	171.4
2005	13.8	6.6	0.5	0.2	0.3	0.9	2.0	5.0	11.9	29.9	75.9	182.5
2004	13.5	5.4	0.5	0.2	0.3	0.9	2.0	5.1	12.2	30.0	74.3	180.8
2003	14.2	6.8	0.4	0.2	0.4	0.8	2.1	5.0	12.4	30.1	79.1	199.0
2002	14.1	6.8	0.5	0.2	0.3	0.8	2.0	4.9	11.9	32.1	79.0	199.3
2001	13.5	7.4	0.6	0.2	0.3	0.8	1.7	4.8	11.8	29.9	74.4	198.5
2000	13.5	7.0	0.6	0.1	0.2	0.7	1.9	4.3	11.2	28.7	73.6	214.5
1999	13.2	6.2	0.5	0.2	0.3	0.7	1.6	4.2	10.8	28.2	71.8	215.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
Non-Hispanic white, both sexes												
2006 5/	12.7	4.0	0.5	0.2	0.3	0.6	1.7	4.2	11.0	29.0	78.4	173.2
2005 5/	12.7	4.3	0.5	0.2	0.3	0.6	1.6	4.3	11.2	29.8	76.8	179.2
2004 5/	12.5	4.8	0.4	0.2	0.3	0.7	1.6	4.4	11.1	29.3	77.2	179.6
2003 5/	12.9	4.2	0.5	0.2	0.3	0.6	1.7	4.3	11.3	29.3	80.8	192.4
2002	12.9	5.5	0.5	0.2	0.3	0.7	1.6	4.3	10.7	31.0	81.3	192.6
2001	12.3	5.1	0.7	0.1	0.3	0.6	1.4	4.0	10.4	29.4	77.6	195.6
2000	11.9	4.8	0.5	0.2	0.2	0.5	1.5	3.8	10.0	27.7	74.2	204.3
1999	11.7	5.3	0.5	0.2	0.3	0.6	1.4	3.6	9.5	27.3	73.0	206.0
Non-Hispanic white, male												
2006 5/	11.7	4.1	0.5	0.2	0.2	0.6	1.8	4.8	11.9	31.5	88.5	193.3
2005 5/	11.5	5.3	0.6	0.2	0.3	0.6	1.6	4.6	12.3	32.7	86.2	193.6
2004 5/	11.4	5.7	*	0.2	0.3	0.7	1.6	4.8	11.7	32.4	88.5	198.5
2003 5/	11.4	3.9	0.6	0.2	0.4	0.6	1.7	4.6	12.0	32.5	90.5	204.8
2002	11.5	5.6	0.5	0.3	0.3	0.7	1.6	4.6	11.8	34.0	92.1	203.9
2001	11.0	5.8	0.9	*	0.3	0.5	1.7	4.2	11.1	32.5	89.7	213.6
2000	10.4	4.7	0.5	0.2	0.3	0.6	1.6	4.3	10.9	30.9	84.7	211.2
1999	10.2	6.7	0.5	0.2	0.3	0.7	1.5	4.0	9.9	30.5	84.6	218.1
Non-Hispanic white, female												
2006 5/	13.7	3.9	0.4	*	0.3	0.6	1.6	3.7	10.1	26.8	71.4	163.9
2005 5/	13.9	3.3	*	*	0.2	0.7	1.6	4.0	10.1	27.2	70.3	172.6
2004 5/	13.6	3.9	0.5	0.2	0.2	0.7	1.5	4.1	10.4	26.7	69.5	171.1
2003 5/	14.3	4.5	*	0.2	0.3	0.7	1.7	3.9	10.6	26.6	74.3	187.0
2002	14.2	5.4	*	0.2	0.3	0.6	1.7	3.9	9.7	28.5	74.0	187.7
2001	13.6	4.4	0.5	*	0.3	0.6	1.2	3.8	9.9	26.8	69.6	188.1
2000	13.4	4.8	0.6	*	*	0.5	1.5	3.3	9.2	24.9	67.2	201.6
1999	13.1	3.8	0.5	0.2	0.2	0.5	1.2	3.3	9.1	24.5	65.4	201.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
Non-Hispanic black, both sexes												
2006 5/	16.0	15.6	*	0.3	0.6	1.6	5.0	13.4	31.6	70.2	153.3	290.9
2005 5/	16.5	19.9	0.8	0.3	0.7	2.0	4.7	13.4	31.5	68.2	159.9	340.8
2004 5/	16.1	14.2	*	*	0.7	1.9	4.6	13.6	32.2	71.4	157.9	324.7
2003 5/	16.8	18.8	1.0	*	0.7	2.0	5.1	14.0	32.8	74.0	160.0	373.9
2002	16.8	13.1	*	0.3	0.4	2.0	4.6	13.2	33.8	76.0	172.9	370.8
2001	16.4	20.3	0.9	*	0.7	1.9	4.8	13.7	32.6	72.8	161.9	368.6
2000	16.7	16.6	1.1	*	0.6	2.1	4.8	13.9	31.8	72.5	175.1	395.9
1999	16.6	18.6	*	*	0.6	1.6	5.1	13.2	30.8	75.3	169.8	414.2
Non-Hispanic black, male												
2006 5/	14.9	16.3	*	*	0.7	1.7	5.2	15.1	37.9	77.2	179.6	300.5
2005 5/	15.5	20.2	*	*	0.8	1.9	4.7	15.2	36.6	81.9	183.3	379.7
2004 5/	15.1	15.8	*	*	0.7	2.1	4.8	15.2	37.1	82.9	192.5	346.5
2003 5/	15.8	21.7	*	*	*	2.3	5.2	15.8	38.7	86.9	191.3	409.1
2002	15.5	13.9	*	*	*	1.9	5.0	15.0	36.6	87.6	213.7	399.3
2001	15.4	21.4	*	*	0.9	1.9	4.9	15.2	37.4	86.9	196.8	430.8
2000	15.3	16.4	*	*	*	2.4	4.8	16.8	35.9	81.6	208.9	432.1
1999	15.7	20.8	*	*	*	1.7	6.0	15.4	37.9	91.5	202.6	451.6
Non-Hispanic black, female												
2006 5/	17.0	14.8	*	*	0.6	1.6	4.8	12.0	26.5	65.3	138.4	287.0
2005 5/	17.4	19.5	*	*	0.6	2.1	4.7	11.8	27.3	58.4	146.7	325.2
2004 5/	17.0	12.6	*	*	0.8	1.8	4.5	12.3	28.3	63.1	138.3	316.0
2003 5/	17.7	15.8	*	*	0.7	1.7	5.1	12.5	28.2	64.9	142.3	360.3
2002	18.0	12.3	*	*	*	2.0	4.2	11.6	31.6	67.8	149.9	359.9
2001	17.2	19.1	*	*	*	1.9	4.7	12.5	28.8	62.8	142.4	345.6
2000	17.9	16.9	*	*	*	1.8	4.9	11.5	28.5	66.1	156.1	382.5
1999	17.4	16.4	*	*	*	1.6	4.3	11.3	25.1	63.9	151.4	400.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
All origins 3/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
Non-Hispanic 4/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
Non-Hispanic white, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
Non-Hispanic black, both sexes												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
All origins 3/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
Non-Hispanic 4/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
Non-Hispanic white, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
All origins 3/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
Non-Hispanic 4/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
Non-Hispanic white, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
All origins 3/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
Non-Hispanic 4/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
Non-Hispanic white, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
All origins 3/, both sexes												
2006	2.4	*	*	*	*	0.2	1.2	6.7	6.7	4.5	4.6	3.1
2005	1.9	*	*	*	*	0.2	1.2	5.4	4.5	3.5	3.7	2.4
2004	1.8	*	*	*	*	0.2	1.4	5.3	4.3	3.4	3.6	2.7
2003	1.9	*	*	*	*	0.2	1.5	5.5	3.9	3.8	4.2	2.4
2002	2.0	*	*	*	0.1	0.2	1.6	5.8	4.0	4.2	4.6	3.9
2001	2.0	*	*	*	0.1	0.2	1.8	5.3	4.0	4.6	4.9	3.8
2000	1.9	*	*	*	*	0.2	1.7	5.0	3.7	4.6	5.3	4.0
1999	1.7	*	*	*	*	0.2	1.6	4.1	3.3	4.8	5.3	4.7
All origins 3/, male												
2006	3.2	*	*	*	*	0.2	1.7	9.9	9.9	5.0	4.6	3.0
2005	2.5	*	*	*	*	0.2	1.6	8.1	6.7	3.7	3.7	2.6
2004	2.4	*	*	*	*	0.2	1.8	8.0	6.1	4.0	3.6	3.0
2003	2.5	*	*	*	*	0.2	2.1	8.3	5.4	4.2	4.8	2.8
2002	2.7	*	*	*	*	0.3	2.3	8.9	5.5	4.5	5.3	5.0
2001	2.5	*	*	*	*	0.2	2.5	8.2	5.3	4.8	5.3	4.4
2000	2.4	*	*	*	*	0.2	2.4	7.7	5.0	4.8	6.4	5.7
1999	2.2	*	*	*	*	0.3	2.3	6.3	4.3	5.1	6.6	6.9
All origins 3/, female												
2006	1.6	*	*	*	*	0.1	0.8	3.7	3.8	4.0	4.6	3.1
2005	1.3	*	*	*	*	0.1	0.7	2.9	2.5	3.3	3.7	2.3
2004	1.3	*	*	*	*	0.1	1.0	2.7	2.5	3.0	3.5	2.5
2003	1.3	*	*	*	*	0.1	0.9	2.9	2.5	3.5	3.8	2.3
2002	1.4	*	*	*	*	0.2	1.0	2.8	2.7	3.9	4.1	3.3
2001	1.4	*	*	*	*	0.2	1.1	2.5	2.8	4.4	4.6	3.5
2000	1.4	*	*	*	*	0.2	1.1	2.4	2.6	4.5	4.7	3.3
1999	1.3	*	*	*	*	0.2	0.9	2.0	2.4	4.5	4.5	3.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
Hispanic, both sexes												
2006	2.5	*	*	*	*	*	1.6	9.4	13.0	10.8	10.2	8.6
2005	1.9	*	*	*	*	*	1.8	6.9	8.5	8.6	6.6	*
2004	1.7	*	*	*	*	*	1.8	6.8	7.7	7.0	7.5	*
2003	1.8	*	*	*	*	*	1.8	7.0	7.2	8.7	9.2	*
2002	2.0	*	*	*	*	*	1.7	8.0	8.3	8.6	10.5	*
2001	2.1	*	*	*	*	*	2.3	7.8	10.4	9.9	11.0	*
2000	2.0	*	*	*	*	*	2.5	7.0	8.6	11.1	13.0	*
1999	1.9	*	*	*	*	*	2.1	6.8	7.8	12.4	13.5	*
Hispanic, male												
2006	3.2	*	*	*	*	*	2.5	13.9	17.6	12.0	8.2	*
2005	2.4	*	*	*	*	*	2.5	10.3	11.4	9.0	*	*
2004	2.2	*	*	*	*	*	2.6	10.2	10.4	8.0	*	*
2003	2.3	*	*	*	*	*	2.6	10.8	10.4	7.8	7.7	*
2002	2.5	*	*	*	*	*	2.5	12.0	11.6	7.6	10.3	*
2001	2.6	*	*	*	*	*	3.1	12.2	14.2	7.0	10.3	*
2000	2.5	*	*	*	*	*	3.8	10.6	11.7	11.0	12.3	*
1999	2.3	*	*	*	*	*	3.3	10.5	9.8	13.0	*	*
Hispanic, female												
2006	1.8	*	*	*	*	*	*	4.7	8.7	9.7	11.5	*
2005	1.3	*	*	*	*	*	1.0	3.4	5.8	8.2	6.7	*
2004	1.2	*	*	*	*	*	0.9	3.4	5.2	6.2	9.1	*
2003	1.3	*	*	*	*	*	1.0	3.1	4.4	9.4	10.3	*
2002	1.4	*	*	*	*	*	0.9	4.0	5.4	9.4	10.7	*
2001	1.7	*	*	*	*	*	1.4	3.5	7.1	12.3	11.5	*
2000	1.5	*	*	*	*	*	1.0	3.3	5.8	11.3	13.5	*
1999	1.5	*	*	*	*	*	0.8	3.3	6.0	12.0	16.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
Non-Hispanic 4/, both sexes												
2006	2.4	*	*	*	*	0.1	1.2	6.4	6.2	3.9	4.2	2.8
2005	1.9	*	*	*	*	0.2	1.0	5.3	4.2	3.1	3.5	2.2
2004	1.8	*	*	*	*	0.2	1.3	5.2	4.0	3.2	3.3	2.6
2003	1.9	*	*	*	*	0.2	1.4	5.4	3.6	3.4	4.0	2.3
2002	2.0	*	*	*	*	0.2	1.6	5.6	3.6	3.8	4.3	3.7
2001	1.9	*	*	*	*	0.2	1.7	5.0	3.5	4.2	4.6	3.6
2000	1.9	*	*	*	*	0.2	1.6	4.8	3.3	4.2	5.0	3.8
1999	1.7	*	*	*	*	0.2	1.5	3.8	3.0	4.3	5.0	4.7
Non-Hispanic 4/, male												
2006	3.2	*	*	*	*	0.2	1.5	9.3	9.2	4.4	4.4	3.0
2005	2.5	*	*	*	*	0.2	1.4	7.8	6.2	3.3	3.5	2.4
2004	2.5	*	*	*	*	0.2	1.7	7.8	5.8	3.7	3.5	3.0
2003	2.5	*	*	*	*	0.2	2.0	8.0	5.0	3.9	4.7	2.7
2002	2.7	*	*	*	*	0.3	2.3	8.6	4.9	4.3	5.1	4.8
2001	2.5	*	*	*	*	0.2	2.4	7.7	4.6	4.6	5.0	4.3
2000	2.4	*	*	*	*	0.2	2.2	7.4	4.4	4.4	6.1	5.4
1999	2.2	*	*	*	*	0.2	2.2	5.9	3.9	4.7	6.4	6.8
Non-Hispanic 4/, female												
2006	1.6	*	*	*	*	*	0.8	3.6	3.4	3.5	4.2	2.7
2005	1.3	*	*	*	*	0.1	0.7	2.8	2.3	2.9	3.5	2.2
2004	1.3	*	*	*	*	0.2	1.0	2.7	2.3	2.7	3.2	2.4
2003	1.3	*	*	*	*	0.1	0.8	2.8	2.3	3.0	3.5	2.2
2002	1.4	*	*	*	*	0.2	1.0	2.7	2.4	3.5	3.8	3.2
2001	1.4	*	*	*	*	0.2	1.0	2.3	2.5	3.8	4.3	3.3
2000	1.4	*	*	*	*	0.2	1.0	2.4	2.3	4.0	4.3	3.2
1999	1.2	*	*	*	*	0.2	0.9	1.8	2.1	4.0	4.0	3.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
Non-Hispanic white, both sexes												
2006 5/	2.3	*	*	*	*	0.1	1.1	6.3	5.3	3.3	3.6	2.5
2005 5/	1.8	*	*	*	*	0.1	1.0	5.1	3.6	2.4	3.2	2.0
2004 5/	1.8	*	*	*	*	0.1	1.3	4.9	3.4	2.7	3.0	2.4
2003 5/	1.8	*	*	*	*	0.2	1.4	5.2	2.9	2.9	3.6	2.2
2002	1.9	*	*	*	*	0.2	1.6	5.3	3.0	3.3	4.1	3.5
2001	1.8	*	*	*	*	0.2	1.7	4.6	2.9	3.7	4.3	3.3
2000	1.8	*	*	*	*	0.1	1.6	4.3	2.7	3.7	4.7	3.7
1999	1.6	*	*	*	*	0.2	1.4	3.4	2.5	3.7	4.7	4.5
Non-Hispanic white, male												
2006 5/	3.1	*	*	*	*	0.2	1.4	9.3	7.9	3.7	3.9	3.1
2005 5/	2.5	*	*	*	*	*	1.4	7.7	5.5	2.8	3.4	2.2
2004 5/	2.4	*	*	*	*	*	1.7	7.4	4.8	3.2	3.2	2.7
2003 5/	2.4	*	*	*	*	0.2	2.0	7.7	4.2	3.4	4.2	2.7
2002	2.6	*	*	*	*	0.3	2.3	8.2	4.1	3.9	4.9	4.6
2001	2.5	*	*	*	*	0.2	2.3	7.2	4.0	4.3	5.1	4.4
2000	2.3	*	*	*	*	*	2.1	6.7	3.6	3.8	5.9	5.4
1999	2.1	*	*	*	*	0.2	2.1	5.2	3.4	4.3	6.4	6.5
Non-Hispanic white, female												
2006 5/	1.5	*	*	*	*	*	0.8	3.4	2.9	2.9	3.4	2.2
2005 5/	1.2	*	*	*	*	*	0.7	2.6	1.9	2.2	3.1	1.9
2004 5/	1.2	*	*	*	*	*	0.9	2.5	2.0	2.3	2.8	2.3
2003 5/	1.2	*	*	*	*	*	0.8	2.6	1.8	2.5	3.1	2.0
2002	1.3	*	*	*	*	*	0.9	2.5	2.0	2.8	3.5	3.1
2001	1.2	*	*	*	*	*	1.0	2.0	2.0	3.1	3.8	2.9
2000	1.3	*	*	*	*	*	1.0	2.0	1.8	3.6	4.0	3.1
1999	1.1	*	*	*	*	0.2	0.8	1.7	1.6	3.3	3.6	3.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
Non-Hispanic black, both sexes												
2006 5/	2.9	*	*	*	*	*	1.3	7.8	12.5	7.3	6.7	*
2005 5/	2.3	*	*	*	*	*	1.2	7.1	8.5	6.2	5.1	*
2004 5/	2.3	*	*	*	*	*	1.4	7.6	8.8	4.6	5.0	*
2003 5/	2.3	*	*	*	*	*	1.8	7.6	8.2	5.3	5.9	*
2002	2.3	*	*	*	*	0.4	2.0	8.2	7.2	6.0	5.2	*
2001	2.3	*	*	*	*	*	2.2	8.1	6.8	5.6	5.7	*
2000	2.2	*	*	*	*	0.4	2.1	8.0	5.8	5.8	6.3	*
1999	2.1	*	*	*	*	*	2.2	7.0	5.1	6.8	5.9	*
Non-Hispanic black, male												
2006 5/	3.8	*	*	*	*	*	1.7	11.1	20.0	9.3	7.5	*
2005 5/	2.9	*	*	*	*	*	1.5	10.3	12.9	7.3	*	*
2004 5/	3.1	*	*	*	*	*	1.5	11.9	14.1	6.7	*	*
2003 5/	3.1	*	*	*	*	*	2.4	11.7	12.1	5.9	6.1	*
2002	3.1	*	*	*	*	*	2.6	12.6	10.7	7.0	*	*
2001	2.8	*	*	*	*	*	3.0	12.0	9.0	5.0	*	*
2000	2.9	*	*	*	*	*	3.0	12.3	8.1	7.2	7.7	*
1999	2.6	*	*	*	*	*	3.1	11.8	6.3	5.7	*	*
Non-Hispanic black, female												
2006 5/	2.0	*	*	*	*	*	1.0	5.0	6.5	5.9	6.3	*
2005 5/	1.7	*	*	*	*	*	0.9	4.4	5.0	5.4	5.9	*
2004 5/	1.5	*	*	*	*	*	1.4	3.9	4.5	3.1	5.3	*
2003 5/	1.6	*	*	*	*	*	1.2	4.1	5.1	4.9	5.9	*
2002	1.7	*	*	*	*	*	1.5	4.4	4.4	5.2	5.3	*
2001	1.8	*	*	*	*	*	1.4	4.7	5.0	6.0	6.7	*
2000	1.6	*	*	*	*	*	1.4	4.3	3.9	4.7	5.6	*
1999	1.6	*	*	*	*	*	1.4	2.9	4.1	7.5	6.7	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
All origins 3/, both sexes												
2006	4.0	*	*	*	0.5	2.9	9.2	10.1	5.5	2.5	0.8	*
2005	4.2	*	*	*	0.4	3.3	9.9	10.6	5.3	2.3	0.8	*
2004	4.4	*	*	0.1	0.5	3.7	10.9	10.6	5.4	2.4	0.8	*
2003	4.7	*	*	0.1	0.4	4.0	12.0	10.9	5.4	2.4	0.7	*
2002	4.9	*	*	0.1	0.4	4.6	12.7	11.2	5.1	2.2	0.8	*
2001	5.0	*	*	0.1	0.6	5.3	13.0	10.5	5.2	2.1	0.7	*
2000	5.1	*	*	0.1	0.5	6.1	13.1	11.0	5.1	2.2	0.7	*
1999	5.3	*	0.2	0.2	0.5	6.8	13.8	10.7	4.8	2.2	0.6	*
All origins 3/, male												
2006	5.9	*	*	*	0.6	3.5	12.9	15.3	8.9	4.2	1.6	*
2005	6.3	*	*	*	0.4	4.0	14.3	16.4	8.8	4.1	1.4	*
2004	6.6	*	*	*	0.5	4.5	15.7	16.3	9.0	4.0	1.4	*
2003	7.1	*	*	*	0.4	5.1	17.5	17.2	9.1	4.0	1.5	*
2002	7.4	*	*	*	0.4	5.9	18.8	17.7	8.5	3.9	1.4	*
2001	7.6	*	*	0.1	0.5	7.1	19.5	16.8	8.6	3.5	1.5	*
2000	7.9	*	*	0.1	0.5	8.0	19.8	17.8	8.7	3.8	1.3	*
1999	8.2	*	*	0.2	0.5	9.5	21.0	17.5	8.3	3.8	1.0	*
All origins 3/, female												
2006	2.2	*	*	*	0.4	2.3	5.4	5.1	2.3	1.1	0.3	*
2005	2.2	*	*	*	0.3	2.6	5.6	5.1	2.0	0.9	0.4	*
2004	2.4	*	*	*	0.4	2.8	6.2	5.2	2.0	1.0	0.5	*
2003	2.4	*	*	*	0.4	2.8	6.5	4.8	2.1	1.0	0.3	*
2002	2.5	*	*	*	0.4	3.3	6.7	4.8	1.9	0.8	0.3	*
2001	2.5	*	*	*	0.6	3.5	6.7	4.4	2.0	0.9	*	*
2000	2.5	*	*	0.1	0.4	4.2	6.5	4.4	1.8	0.8	0.3	*
1999	2.5	*	*	0.2	0.5	4.1	6.7	4.1	1.6	0.8	0.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
Hispanic, both sexes												
2006	3.6	*	*	*	0.4	2.4	8.8	11.7	7.6	4.7	*	*
2005	3.9	*	*	*	0.4	2.5	9.6	13.0	7.2	4.3	*	*
2004	4.3	*	*	*	0.4	2.8	11.0	14.1	8.4	5.0	*	*
2003	4.7	*	*	*	0.4	3.3	12.3	15.9	9.8	5.2	*	*
2002	4.8	*	*	*	0.4	3.6	13.4	14.7	9.5	3.4	*	*
2001	5.0	*	*	*	0.5	4.4	13.9	15.5	9.2	4.5	*	*
2000	5.4	*	*	*	0.3	5.6	14.9	16.9	9.5	3.8	*	*
1999	5.8	*	*	*	0.5	6.4	17.3	15.4	8.4	4.7	*	*
Hispanic, male												
2006	5.6	*	*	*	*	3.4	12.9	18.6	12.9	7.9	*	*
2005	5.9	*	*	*	0.6	3.3	14.7	20.8	12.6	7.3	*	*
2004	6.4	*	*	*	*	4.1	16.1	22.0	14.4	8.0	*	*
2003	7.1	*	*	*	0.5	4.6	17.7	26.1	15.4	9.2	*	*
2002	7.2	*	*	*	*	5.1	20.1	23.1	15.0	6.9	*	*
2001	7.6	*	*	*	0.6	6.0	20.8	24.9	14.8	8.6	*	*
2000	8.2	*	*	*	*	7.8	22.7	27.5	15.3	7.2	*	*
1999	8.7	*	*	*	0.6	9.3	26.2	24.2	15.4	9.0	*	*
Hispanic, female												
2006	1.6	*	*	*	*	1.2	4.1	4.5	2.8	*	*	*
2005	1.6	*	*	*	*	1.4	4.0	5.2	2.3	*	*	*
2004	1.9	*	*	*	*	1.2	5.3	6.2	2.9	*	*	*
2003	2.2	*	*	*	*	1.8	6.3	5.6	4.6	*	*	*
2002	2.2	*	*	*	*	1.8	6.2	6.3	4.6	*	*	*
2001	2.3	*	*	*	*	2.5	6.5	6.2	4.1	*	*	*
2000	2.4	*	*	*	*	3.0	6.4	6.6	4.4	*	*	*
1999	2.6	*	*	*	*	3.2	7.9	6.7	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
Non-Hispanic 4/, both sexes												
2006	4.1	*	*	*	0.5	3.0	9.2	9.9	5.2	2.3	0.7	*
2005	4.3	*	*	*	0.4	3.5	9.9	10.3	5.1	2.2	0.7	*
2004	4.4	*	*	0.1	0.5	3.9	10.9	10.1	5.0	2.1	0.7	*
2003	4.6	*	*	0.1	0.4	4.1	11.9	10.2	5.0	2.1	0.7	*
2002	4.8	*	*	0.1	0.4	4.8	12.5	10.6	4.7	2.1	0.7	*
2001	4.9	*	*	0.1	0.6	5.4	12.7	9.9	4.8	1.9	0.6	*
2000	5.0	*	*	0.1	0.5	6.2	12.7	10.3	4.7	2.0	0.6	*
1999	5.2	*	*	0.2	0.5	6.8	13.2	10.0	4.5	2.0	0.6	*
Non-Hispanic 4/, male												
2006	6.0	*	*	*	0.6	3.6	12.9	14.8	8.5	3.9	1.4	*
2005	6.3	*	*	*	0.4	4.1	14.1	15.7	8.5	3.8	1.2	*
2004	6.5	*	*	*	0.5	4.6	15.5	15.5	8.5	3.6	1.2	*
2003	7.0	*	*	*	0.4	5.2	17.3	16.0	8.4	3.6	1.4	*
2002	7.3	*	*	*	0.4	6.0	18.4	16.9	7.9	3.7	1.3	*
2001	7.5	*	*	0.1	0.5	7.3	19.0	15.8	8.0	3.1	1.3	*
2000	7.7	*	*	*	0.5	7.9	19.1	16.7	8.0	3.4	1.2	*
1999	8.0	*	*	0.2	0.5	9.4	20.1	16.5	7.6	3.4	1.0	*
Non-Hispanic 4/, female												
2006	2.3	*	*	*	0.4	2.5	5.6	5.1	2.2	1.0	0.3	*
2005	2.3	*	*	*	0.4	2.8	5.8	5.0	2.0	0.8	0.4	*
2004	2.4	*	*	*	0.5	3.1	6.3	5.0	1.9	0.9	0.4	*
2003	2.4	*	*	*	0.4	3.0	6.5	4.7	1.9	0.9	*	*
2002	2.5	*	*	*	0.5	3.6	6.7	4.7	1.7	0.8	0.3	*
2001	2.5	*	*	*	0.7	3.6	6.6	4.2	1.8	0.9	*	*
2000	2.5	*	*	0.1	0.5	4.4	6.4	4.2	1.6	0.8	*	*
1999	2.5	*	*	0.1	0.6	4.2	6.5	3.9	1.6	0.7	0.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
Non-Hispanic white, both sexes												
2006 5/	1.8	*	*	*	0.1	0.9	4.1	4.3	2.2	1.0	0.4	*
2005 5/	1.9	*	*	*	0.1	1.1	4.6	4.5	2.2	0.9	0.4	*
2004 5/	1.9	*	*	*	0.1	1.2	5.1	4.3	2.2	0.9	0.4	*
2003 5/	2.1	*	*	*	0.1	1.5	5.6	4.5	2.2	0.8	0.3	*
2002	2.1	*	*	*	0.1	1.7	5.8	4.6	2.2	0.8	0.3	*
2001	2.2	*	*	*	0.1	2.1	6.0	4.2	2.1	0.7	0.3	*
2000	2.3	*	*	*	0.1	2.4	6.2	4.6	2.2	0.8	0.3	*
1999	2.4	*	*	*	0.1	3.0	6.4	4.6	2.0	0.8	0.3	*
Non-Hispanic white, male												
2006 5/	3.0	*	*	*	*	1.3	6.8	7.4	4.0	1.8	0.7	*
2005 5/	3.1	*	*	*	*	1.6	7.6	7.6	3.8	1.7	0.7	*
2004 5/	3.3	*	*	*	*	1.8	8.4	7.4	4.0	1.7	0.7	*
2003 5/	3.6	*	*	*	*	2.4	9.4	7.8	4.0	1.5	0.8	*
2002	3.7	*	*	*	*	2.7	9.8	8.0	4.0	1.5	0.5	*
2001	3.8	*	*	*	*	3.2	10.4	7.5	3.8	1.4	0.6	*
2000	4.0	*	*	*	0.2	3.7	10.4	8.2	4.0	1.5	0.6	*
1999	4.2	*	*	*	0.2	4.9	10.9	8.2	3.6	1.5	0.6	*
Non-Hispanic white, female												
2006 5/	0.6	*	*	*	*	0.5	1.5	1.3	0.5	0.2	*	*
2005 5/	0.6	*	*	*	*	0.7	1.6	1.5	0.6	*	*	*
2004 5/	0.6	*	*	*	*	0.6	1.8	1.3	0.4	*	*	*
2003 5/	0.6	*	*	*	*	0.7	1.8	1.2	0.5	0.3	*	*
2002	0.6	*	*	*	*	0.8	1.8	1.2	0.5	*	*	*
2001	0.6	*	*	*	*	0.9	1.7	1.0	0.5	*	*	*
2000	0.7	*	*	*	*	1.1	2.0	1.1	0.5	*	*	*
1999	0.7	*	*	*	*	1.2	1.9	1.1	0.4	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
Non-Hispanic black, both sexes												
2006 5/	17.9	*	*	*	2.2	14.3	39.7	48.3	30.8	14.3	4.8	*
2005 5/	18.5	*	*	*	1.7	15.6	42.1	50.5	30.4	14.0	5.0	*
2004 5/	19.4	*	*	*	2.3	17.9	46.0	50.9	29.9	13.6	4.6	*
2003 5/	20.1	*	*	0.5	1.9	18.1	50.5	51.3	29.9	14.4	4.7	*
2002	21.3	*	*	*	2.1	21.5	54.0	54.8	26.3	14.6	5.7	*
2001	21.6	*	*	0.4	2.8	24.0	55.0	51.9	28.0	12.8	5.3	*
2000	22.0	*	*	0.4	2.2	27.2	54.5	53.9	26.3	13.2	4.6	*
1999	22.3	*	*	0.6	2.3	28.3	57.2	52.2	26.4	13.2	3.7	*
Non-Hispanic black, male												
2006 5/	24.4	*	*	*	2.6	16.2	52.3	70.3	50.3	24.1	10.3	*
2005 5/	25.8	*	*	*	1.8	17.9	56.7	76.4	52.0	25.0	8.7	*
2004 5/	26.7	*	*	*	2.4	20.0	62.1	76.3	50.8	23.4	7.7	*
2003 5/	28.5	*	*	*	1.9	21.7	70.0	79.0	51.4	25.3	8.9	*
2002	30.3	*	*	*	2.0	25.5	75.6	86.2	44.8	26.6	12.3	*
2001	30.8	*	*	*	2.4	30.8	77.8	81.4	47.7	21.4	11.4	*
2000	31.7	*	*	*	2.2	33.4	79.0	86.6	46.4	23.8	8.9	*
1999	32.6	*	*	0.7	1.9	37.1	83.5	85.5	45.6	24.0	6.6	*
Non-Hispanic black, female												
2006 5/	12.1	*	*	*	1.9	12.6	28.5	29.4	15.1	7.2	*	*
2005 5/	11.8	*	*	*	1.6	13.4	29.2	28.3	12.9	6.1	*	*
2004 5/	12.8	*	*	*	2.2	15.9	31.6	29.0	13.1	6.6	*	*
2003 5/	12.6	*	*	*	2.0	14.8	33.2	27.6	12.6	6.6	*	*
2002	13.2	*	*	*	2.2	17.9	34.9	28.0	11.5	6.0	*	*
2001	13.3	*	*	*	3.2	17.8	34.8	26.6	12.1	6.7	*	*
2000	13.1	*	*	*	2.3	21.7	32.8	25.8	10.1	5.8	*	*
1999	13.0	*	*	*	2.8	20.3	34.1	23.8	11.0	5.6	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
All origins 3/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
Non-Hispanic 4/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
Non-Hispanic white, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)												
All origins 3/, both sexes												
2006	2.0	4.0	0.4	0.2	0.2	0.3	0.6	1.3	2.7	5.8	11.5	20.4
2005	2.6	4.1	0.4	0.2	0.2	0.3	0.9	3.2	4.2	7.0	13.7	21.2
2004	2.6	4.1	0.4	0.1	0.2	0.4	1.0	3.0	4.3	7.1	13.2	22.5
2003	2.6	4.3	0.5	0.1	0.2	0.4	1.0	2.9	4.1	7.3	13.3	22.5
2002	2.3	5.6	0.4	0.2	0.2	0.4	1.0	2.3	3.3	6.7	12.5	21.7
2001	2.0	4.7	0.4	0.2	0.2	0.4	0.8	1.7	3.0	6.2	10.6	20.1
2000	2.0	4.4	0.5	0.2	0.2	0.4	0.8	1.6	2.9	5.8	11.1	21.2
1999	2.6	5.3	0.5	0.2	0.2	0.6	1.2	2.3	3.8	7.8	14.2	27.0
All origins 3/, male												
2006	2.1	4.5	0.5	0.2	0.2	0.3	0.8	1.6	3.2	6.7	13.9	25.1
2005	2.9	4.8	0.5	0.1	0.2	0.4	1.2	4.5	5.3	8.3	16.4	23.6
2004	2.8	4.8	0.4	0.1	0.3	0.4	1.2	4.1	5.4	8.4	15.1	29.6
2003	2.8	4.2	0.5	0.1	0.2	0.4	1.3	4.0	5.1	8.5	15.8	26.5
2002	2.6	6.1	0.5	0.2	0.2	0.5	1.3	3.2	4.2	8.0	15.4	26.8
2001	2.1	4.9	0.5	0.1	0.2	0.4	1.0	2.2	3.7	6.8	13.2	23.6
2000	2.1	4.8	0.5	0.2	0.3	0.4	1.0	1.9	3.3	7.2	13.6	27.6
1999	2.7	6.1	0.6	0.2	0.2	0.6	1.5	2.7	4.2	9.2	17.2	31.9
All origins 3/, female												
2006	1.9	3.6	0.3	0.2	0.2	0.3	0.5	1.0	2.3	5.1	9.9	18.2
2005	2.3	3.4	0.3	0.2	0.2	0.3	0.6	2.0	3.2	5.9	11.8	20.1
2004	2.3	3.4	0.4	0.1	0.2	0.3	0.8	1.9	3.2	6.0	11.9	19.3
2003	2.3	4.5	0.5	0.1	0.2	0.3	0.7	1.9	3.1	6.3	11.7	20.8
2002	2.1	5.0	0.3	0.2	0.2	0.4	0.7	1.4	2.6	5.6	10.6	19.5
2001	1.9	4.6	0.4	0.2	0.2	0.3	0.6	1.2	2.4	5.8	8.9	18.7
2000	1.9	3.9	0.4	0.2	0.2	0.3	0.6	1.2	2.6	4.7	9.5	18.6
1999	2.5	4.5	0.4	0.2	0.2	0.6	0.9	1.9	3.4	6.7	12.3	25.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)												
Hispanic, both sexes												
2006	0.9	2.8	*	*	0.3	0.3	0.7	1.2	2.1	5.2	7.9	10.2
2005	1.8	3.5	*	*	*	0.4	1.3	4.8	5.5	8.6	13.3	15.4
2004	1.8	3.4	0.6	*	*	0.3	1.3	4.3	6.0	7.6	14.6	18.6
2003	1.7	2.7	*	*	*	0.3	1.3	4.4	5.6	8.4	15.3	13.0
2002	1.5	4.9	*	*	*	0.7	1.2	3.5	4.3	6.4	11.3	17.7
2001	1.1	5.0	*	*	*	0.5	0.8	2.0	3.0	5.9	10.3	15.0
2000	1.2	3.0	*	*	*	0.4	1.0	1.9	3.2	5.8	11.3	19.2
1999	1.5	4.8	*	*	*	0.6	1.4	2.3	3.8	7.7	14.2	26.0
Hispanic, male												
2006	1.1	*	*	*	*	*	0.8	1.8	2.5	6.2	11.1	*
2005	2.0	*	*	*	*	0.5	1.7	7.0	7.2	8.8	13.0	*
2004	2.1	*	*	*	*	*	1.7	6.0	8.3	8.8	17.5	*
2003	2.0	*	*	*	*	*	1.8	6.8	7.7	9.8	16.5	*
2002	1.8	6.3	*	*	*	0.8	1.6	5.2	5.6	7.6	12.3	*
2001	1.3	5.0	*	*	*	0.6	1.1	2.8	4.4	7.6	12.1	*
2000	1.3	*	*	*	*	0.6	1.4	2.1	4.4	5.9	13.3	*
1999	1.5	5.7	*	*	*	0.6	1.9	2.7	3.8	7.9	15.1	*
Hispanic, female												
2006	0.8	*	*	*	*	*	*	*	1.8	4.5	5.8	*
2005	1.6	*	*	*	*	*	0.8	2.6	4.0	8.3	13.5	17.6
2004	1.4	*	*	*	*	*	0.9	2.6	4.0	6.7	12.6	19.4
2003	1.3	*	*	*	*	*	0.7	2.0	3.6	7.4	14.5	*
2002	1.2	*	*	*	*	0.6	*	1.7	3.2	5.3	10.7	*
2001	1.0	5.0	*	*	*	*	*	1.3	*	4.6	9.1	*
2000	1.0	*	*	*	*	*	*	1.8	*	5.6	9.9	*
1999	1.4	*	*	*	*	*	*	1.9	3.8	7.6	13.5	28.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)												
Non-Hispanic 4/, both sexes												
2006	2.1	4.4	0.4	0.2	0.2	0.3	0.6	1.3	2.8	5.9	11.7	20.9
2005	2.7	4.3	0.4	0.2	0.2	0.3	0.9	3.0	4.1	6.9	13.7	21.5
2004	2.7	4.3	0.4	0.1	0.2	0.4	0.9	2.8	4.1	7.0	13.1	22.6
2003	2.7	4.8	0.5	0.1	0.2	0.4	1.0	2.7	3.9	7.2	13.2	22.9
2002	2.4	5.7	0.4	0.2	0.2	0.4	1.0	2.2	3.3	6.7	12.5	21.8
2001	2.1	4.6	0.5	0.2	0.2	0.3	0.8	1.6	3.0	6.2	10.6	20.3
2000	2.1	4.7	0.4	0.2	0.2	0.3	0.7	1.5	2.9	5.8	11.1	21.2
1999	2.8	5.5	0.5	0.2	0.2	0.6	1.2	2.3	3.8	7.8	14.2	27.0
Non-Hispanic 4/, male												
2006	2.2	4.7	0.5	0.2	0.2	0.3	0.8	1.6	3.3	6.7	14.1	25.8
2005	3.0	5.1	0.5	0.1	0.2	0.4	1.1	4.2	5.0	8.3	16.5	24.2
2004	3.0	5.2	0.4	*	0.2	0.4	1.1	3.9	5.2	8.4	15.0	30.2
2003	2.9	4.8	0.5	*	0.2	0.4	1.2	3.6	4.9	8.4	15.8	27.0
2002	2.7	5.9	0.5	0.2	0.3	0.4	1.3	3.0	4.0	8.0	15.6	26.8
2001	2.2	4.8	0.5	0.1	0.2	0.4	1.0	2.1	3.6	6.7	13.2	24.0
2000	2.2	5.1	0.5	0.2	0.3	0.3	0.9	1.9	3.2	7.3	13.5	27.8
1999	2.8	6.2	0.6	0.2	0.3	0.6	1.4	2.6	4.2	9.2	17.2	32.3
Non-Hispanic 4/, female												
2006	2.1	4.1	*	0.2	0.2	0.3	0.4	1.0	2.3	5.2	10.1	18.6
2005	2.5	3.4	*	0.2	0.2	0.2	0.6	1.9	3.1	5.7	11.7	20.2
2004	2.4	3.3	0.3	*	0.2	0.3	0.8	1.8	3.1	5.9	11.9	19.2
2003	2.5	4.8	0.5	*	0.2	0.3	0.7	1.9	3.0	6.2	11.5	21.2
2002	2.2	5.4	*	0.2	0.2	0.3	0.7	1.4	2.5	5.6	10.5	19.7
2001	2.1	4.5	0.4	0.2	0.2	0.3	0.6	1.2	2.5	5.8	8.9	18.8
2000	2.0	4.3	0.4	0.2	0.2	0.4	0.6	1.1	2.6	4.7	9.5	18.6
1999	2.7	4.7	0.4	0.2	0.2	0.6	1.0	1.9	3.3	6.7	12.2	24.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)												
Non-Hispanic white, both sexes												
2006 5/	2.2	2.8	0.4	0.2	0.2	0.2	0.5	1.1	2.6	5.8	11.7	21.1
2005 5/	2.8	3.7	0.4	0.2	0.2	0.2	0.7	2.9	3.8	6.7	13.4	21.5
2004 5/	2.8	3.7	0.3	0.1	0.2	0.3	0.8	2.8	3.8	6.7	12.8	22.6
2003 5/	2.8	3.8	0.4	0.1	0.2	0.3	0.8	2.6	3.6	7.2	12.8	23.4
2002	2.5	4.8	0.4	0.2	0.2	0.3	0.8	1.9	2.9	6.6	12.4	22.2
2001	2.1	3.3	0.4	0.1	0.2	0.3	0.7	1.5	2.8	6.0	10.4	20.2
2000	2.1	3.5	0.4	0.2	0.2	0.2	0.6	1.3	2.6	5.6	10.9	21.0
1999	2.7	4.4	0.5	0.2	0.2	0.4	0.9	1.8	3.3	7.4	13.6	26.4
Non-Hispanic white, male												
2006 5/	2.3	3.1	0.4	*	0.2	0.2	0.6	1.4	3.0	6.7	14.0	26.5
2005 5/	3.1	4.4	0.4	*	*	0.2	1.0	4.1	4.7	8.2	16.2	24.4
2004 5/	3.0	4.6	*	*	0.2	0.3	0.9	3.8	4.8	8.0	14.6	30.8
2003 5/	2.9	3.6	*	*	0.2	0.3	1.1	3.4	4.5	8.2	15.4	27.6
2002	2.7	4.8	0.4	0.2	0.2	0.3	1.0	2.7	3.5	7.9	15.3	27.2
2001	2.2	3.3	0.4	*	0.2	0.3	0.8	1.9	3.3	6.5	13.1	24.2
2000	2.2	3.9	*	*	0.2	0.2	0.7	1.7	3.0	7.0	13.2	27.0
1999	2.7	5.5	0.6	0.2	0.2	0.4	1.2	2.2	3.8	8.8	16.3	30.8
Non-Hispanic white, female												
2006 5/	2.2	2.4	*	0.2	0.2	0.2	0.4	0.9	2.3	5.1	10.2	18.6
2005 5/	2.5	2.9	*	0.2	0.2	0.2	0.5	1.7	3.0	5.5	11.4	20.1
2004 5/	2.5	2.7	*	*	0.2	0.3	0.7	1.8	2.9	5.5	11.6	18.9
2003 5/	2.6	4.0	*	*	0.2	0.3	0.6	1.7	2.8	6.3	11.1	21.6
2002	2.3	4.9	*	0.2	0.2	0.3	0.5	1.2	2.3	5.5	10.5	20.0
2001	2.1	3.4	*	0.2	0.2	0.2	0.5	1.0	2.3	5.7	8.7	18.6
2000	2.0	3.1	*	0.2	0.2	0.2	0.5	1.0	2.3	4.5	9.5	18.6
1999	2.6	3.3	*	*	0.2	0.4	0.6	1.5	2.9	6.3	11.8	24.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)												
Non-Hispanic black, both sexes												
2006 5/	2.2	10.2	*	*	*	0.7	1.3	2.6	4.5	7.4	12.4	22.6
2005 5/	2.6	6.8	*	*	0.4	0.8	1.5	4.2	6.0	7.8	15.2	22.7
2004 5/	2.6	6.8	*	*	*	0.8	1.7	3.4	6.3	8.8	16.8	24.8
2003 5/	2.6	8.4	0.9	*	*	0.6	1.8	4.4	6.1	7.0	16.4	19.9
2002	2.6	9.0	*	*	0.4	0.9	2.2	4.0	5.9	7.4	13.8	21.8
2001	2.3	10.0	*	*	0.4	0.9	1.6	3.0	5.2	8.3	11.3	20.9
2000	2.3	10.4	*	*	0.4	1.0	1.7	3.0	4.9	8.0	13.3	24.2
1999	3.6	10.0	1.0	*	0.4	1.7	3.1	5.6	7.3	11.9	20.9	34.3
Non-Hispanic black, male												
2006 5/	2.3	10.5	*	*	*	0.8	1.8	3.5	6.1	7.9	15.6	21.1
2005 5/	2.9	8.4	*	*	*	0.9	2.1	5.2	8.2	9.4	19.6	23.8
2004 5/	3.0	7.4	*	*	*	0.9	1.9	5.0	9.1	10.7	20.3	30.5
2003 5/	3.0	9.3	*	*	*	0.8	2.2	5.8	8.1	10.6	19.0	23.4
2002	3.0	10.0	*	*	*	1.1	2.6	5.6	8.7	9.1	17.0	28.4
2001	2.5	11.5	*	*	*	*	2.1	4.0	6.4	9.8	12.6	*
2000	2.6	10.6	*	*	*	0.9	2.2	4.0	5.8	10.5	17.9	31.7
1999	3.7	10.0	*	*	*	1.9	3.1	6.3	7.9	14.1	29.4	46.4
Non-Hispanic black, female												
2006 5/	2.1	10.0	*	*	*	*	0.9	1.9	3.1	7.0	10.6	23.2
2005 5/	2.3	*	*	*	*	*	1.0	3.4	4.2	6.7	12.7	22.2
2004 5/	2.3	*	*	*	*	*	1.5	2.1	4.1	7.4	14.8	22.5
2003 5/	2.2	7.4	*	*	*	*	1.4	3.1	4.6	4.4	14.8	18.6
2002	2.2	7.9	*	*	*	*	1.8	2.6	3.7	6.2	11.9	19.2
2001	2.1	8.4	*	*	*	1.0	1.2	2.1	4.3	7.2	10.6	20.5
2000	2.1	10.2	*	*	*	1.0	1.3	2.2	4.1	6.2	10.8	21.4
1999	3.5	10.0	*	*	*	1.5	3.0	4.9	6.7	10.4	16.1	29.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
All origins 3/, both sexes												
2006	187.0	1.8	2.3	2.2	3.9	9.0	31.9	116.3	321.2	727.2	1,263.8	1,606.1
2005	188.7	1.8	2.3	2.5	4.1	9.0	33.2	118.6	326.9	742.7	1,274.8	1,637.7
2004	188.6	1.8	2.5	2.5	4.1	9.1	33.4	119.0	333.4	755.1	1,280.4	1,653.3
2003	191.5	1.9	2.5	2.6	4.0	9.4	35.0	122.2	343.0	770.3	1,302.5	1,698.2
2002	193.2	1.8	2.6	2.6	4.3	9.7	35.8	123.8	351.1	792.1	1,311.9	1,723.9
2001	194.4	1.6	2.7	2.5	4.3	10.1	36.8	126.5	356.5	802.8	1,315.8	1,765.6
2000	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7	816.3	1,335.6	1,819.4
1999	197.0	1.8	2.7	2.5	4.5	10.0	37.1	127.6	374.6	827.1	1,331.5	1,805.8
All origins 3/, male												
2006	196.6	1.8	2.5	2.5	4.6	8.6	27.4	119.0	363.6	870.4	1,631.3	2,248.7
2005	198.9	2.1	2.6	2.7	4.8	8.8	28.9	121.6	369.5	899.1	1,649.7	2,319.3
2004	198.4	1.8	2.6	2.7	4.8	8.6	29.1	124.3	376.7	907.6	1,662.1	2,349.5
2003	201.3	1.7	2.8	2.8	4.6	8.9	30.8	127.4	386.8	931.7	1,695.4	2,413.8
2002	203.8	2.0	2.7	2.9	4.9	9.2	31.5	128.0	399.8	964.8	1,711.3	2,491.1
2001	205.3	1.5	2.9	2.5	5.0	9.3	32.6	130.3	405.2	984.6	1,727.1	2,613.6
2000	207.2	2.6	3.0	2.7	5.1	9.2	32.7	130.9	415.8	1,001.9	1,760.6	2,710.7
1999	208.9	1.9	2.9	2.7	5.2	9.7	33.3	131.7	428.4	1,019.4	1,767.0	2,722.2
All origins 3/, female												
2006	177.6	1.8	2.1	2.0	3.1	9.5	36.4	113.7	281.8	605.9	1,012.5	1,305.5
2005	178.8	1.5	2.0	2.2	3.3	9.1	37.5	115.8	287.4	610.9	1,020.3	1,324.6
2004	179.1	1.9	2.4	2.2	3.4	9.6	37.7	113.8	293.2	627.1	1,023.5	1,340.1
2003	182.0	2.1	2.1	2.4	3.4	9.9	39.1	117.1	302.3	635.3	1,040.1	1,381.9
2002	183.0	1.6	2.4	2.4	3.6	10.2	40.0	119.8	306.0	648.5	1,046.7	1,391.1
2001	183.9	1.8	2.5	2.4	3.5	10.9	41.0	122.7	311.5	652.2	1,045.4	1,410.7
2000	186.2	2.3	2.5	2.2	3.6	10.4	40.4	124.2	321.3	663.6	1,058.5	1,456.4
1999	185.6	1.7	2.5	2.3	3.7	10.3	40.9	123.7	325.0	669.5	1,048.5	1,436.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
Hispanic, both sexes												
2006	60.1	*	2.2	2.3	4.4	7.2	21.6	74.7	198.6	461.9	844.0	1,077.5
2005	61.3	*	2.2	2.5	4.5	6.8	22.1	77.9	203.3	481.3	882.8	1,128.1
2004	59.3	*	2.0	2.8	4.3	6.9	21.8	75.5	209.7	472.0	872.2	1,131.2
2003	60.3	*	3.0	2.9	4.1	7.1	22.9	80.7	214.7	485.2	890.1	1,234.9
2002	59.7	*	2.3	2.9	4.5	6.8	23.2	78.2	215.1	496.0	894.9	1,308.8
2001	60.5	*	2.8	2.5	4.3	7.2	25.6	82.8	223.4	504.0	913.9	1,363.7
2000	59.9	*	2.5	2.9	4.2	7.3	25.2	82.1	221.0	516.2	953.5	1,383.5
1999	59.6	*	2.5	2.7	4.1	7.4	24.4	84.9	225.1	520.5	926.8	1,399.6
Hispanic, male												
2006	60.4	*	2.6	2.3	5.1	6.1	16.0	71.4	224.8	574.8	1,098.4	1,440.1
2005	63.0	*	2.8	2.5	5.3	6.5	17.8	75.9	236.9	603.5	1,161.8	1,601.5
2004	60.9	*	1.6	3.0	5.5	6.4	18.6	77.4	239.0	585.8	1,174.2	1,508.8
2003	61.5	*	3.6	3.2	4.9	6.8	18.2	81.1	246.5	617.6	1,163.9	1,668.6
2002	61.2	*	2.2	2.9	5.2	6.3	18.4	78.4	254.3	622.3	1,190.8	1,869.0
2001	62.2	*	2.9	2.5	5.5	6.3	21.7	81.5	253.5	642.8	1,258.3	1,967.4
2000	61.3	*	2.9	3.5	4.5	6.9	20.1	79.4	253.1	651.2	1,306.4	2,049.7
1999	61.2	*	2.3	2.8	4.6	6.6	19.8	84.6	268.1	668.9	1,252.9	1,945.3
Hispanic, female												
2006	59.7	*	1.8	2.2	3.5	8.5	27.9	78.1	174.4	370.2	665.9	884.9
2005	59.5	*	1.6	2.5	3.6	7.1	27.0	79.9	172.5	382.5	688.5	880.4
2004	57.7	*	2.4	2.5	3.0	7.5	25.4	73.5	183.0	380.7	663.6	937.0
2003	59.1	*	2.4	2.6	3.2	7.4	28.0	80.2	185.9	379.7	702.1	1,014.8
2002	58.1	*	2.4	2.9	3.6	7.5	28.4	78.0	179.8	395.6	692.2	1,031.2
2001	58.7	*	2.7	2.5	2.9	8.3	29.8	84.2	196.7	394.5	681.2	1,068.6
2000	58.5	*	2.2	2.3	3.7	7.8	30.7	84.7	192.5	410.0	716.5	1,056.5
1999	58.0	*	2.7	2.6	3.5	8.3	29.4	85.1	187.2	404.3	709.8	1,136.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
Non-Hispanic 4/, both sexes												
2006	208.7	1.8	2.3	2.2	3.8	9.5	33.7	120.9	331.3	746.7	1,288.4	1,630.6
2005	209.8	1.8	2.3	2.5	4.0	9.5	35.0	122.9	336.9	761.5	1,296.6	1,660.5
2004	209.4	1.7	2.6	2.4	4.0	9.6	35.2	123.4	342.9	774.5	1,301.1	1,674.3
2003	211.9	2.0	2.3	2.6	4.0	9.9	36.7	126.0	352.4	788.9	1,321.6	1,714.7
2002	213.4	1.7	2.6	2.5	4.2	10.3	37.6	128.0	360.9	810.5	1,329.2	1,736.8
2001	213.9	1.6	2.7	2.4	4.2	10.6	38.2	130.3	365.9	820.5	1,331.4	1,776.6
2000	215.6	2.5	2.7	2.4	4.4	10.3	37.9	131.2	376.6	832.8	1,348.9	1,831.1
1999	215.5	1.7	2.7	2.4	4.5	10.4	38.6	131.0	384.6	843.2	1,344.5	1,816.1
Non-Hispanic 4/, male												
2006	221.3	1.7	2.5	2.5	4.5	9.3	29.5	124.4	374.8	891.3	1,662.8	2,289.8
2005	222.8	2.4	2.6	2.7	4.7	9.4	30.8	126.5	379.9	919.5	1,677.1	2,354.9
2004	221.8	1.8	2.8	2.7	4.6	9.1	30.8	129.2	387.0	928.8	1,687.2	2,387.3
2003	224.3	1.9	2.6	2.7	4.5	9.4	32.8	131.9	396.9	951.1	1,720.2	2,444.0
2002	226.7	1.9	2.8	2.8	4.8	9.9	33.5	132.6	410.0	985.1	1,733.2	2,513.2
2001	227.2	1.6	2.9	2.5	4.9	9.9	34.0	134.6	415.3	1,003.8	1,745.1	2,634.8
2000	228.7	2.6	2.9	2.5	5.2	9.7	34.3	135.1	426.6	1,020.2	1,776.0	2,732.4
1999	229.9	1.6	3.0	2.6	5.3	10.3	35.0	135.4	438.8	1,036.8	1,783.7	2,747.5
Non-Hispanic 4/, female												
2006	196.7	1.9	2.2	2.0	3.0	9.7	37.7	117.5	290.8	624.0	1,032.8	1,324.2
2005	197.5	*	2.1	2.2	3.3	9.5	39.2	119.5	296.8	627.7	1,038.6	1,343.6
2004	197.5	1.6	2.4	2.2	3.4	10.0	39.5	117.8	301.9	644.5	1,041.6	1,355.6
2003	200.0	2.1	2.1	2.4	3.4	10.4	40.6	120.5	311.1	652.7	1,055.6	1,394.3
2002	200.8	1.5	2.4	2.2	3.6	10.6	41.6	123.5	315.4	664.7	1,061.5	1,401.8
2001	201.2	1.6	2.5	2.3	3.6	11.3	42.4	126.0	320.0	668.1	1,059.8	1,419.6
2000	203.1	2.4	2.6	2.2	3.6	10.9	41.4	127.3	330.3	678.1	1,070.6	1,466.6
1999	201.9	1.8	2.5	2.2	3.7	10.5	42.1	126.8	334.5	684.2	1,059.4	1,443.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
Non-Hispanic white, both sexes												
2006 5/	227.1	2.1	2.3	2.3	3.7	9.2	32.4	115.8	325.1	748.2	1,297.8	1,643.1
2005 5/	227.7	2.0	2.3	2.5	4.0	9.0	34.0	117.1	329.0	762.3	1,304.0	1,668.6
2004 5/	227.0	1.9	2.7	2.4	3.9	9.3	33.7	117.3	336.2	773.6	1,305.4	1,680.9
2003 5/	229.5	2.2	2.4	2.6	4.0	9.6	35.2	119.4	345.9	788.8	1,322.6	1,717.8
2002	230.9	1.6	2.6	2.5	4.3	10.0	36.2	122.3	353.0	809.2	1,328.4	1,737.9
2001	230.6	1.8	2.9	2.4	4.3	10.3	36.5	124.6	358.3	815.9	1,326.5	1,775.2
2000	232.2	2.6	2.8	2.4	4.3	10.1	35.6	125.1	369.3	827.7	1,342.9	1,826.6
1999	231.5	1.6	3.0	2.3	4.3	10.2	36.6	124.8	375.0	837.1	1,336.4	1,808.3
Non-Hispanic white, male												
2006 5/	239.9	1.9	2.5	2.5	4.5	9.3	28.9	118.7	363.4	883.0	1,662.9	2,300.2
2005 5/	240.7	2.7	2.5	2.7	4.7	9.0	30.5	120.3	366.1	910.4	1,673.7	2,358.3
2004 5/	239.2	2.0	2.8	2.7	4.4	9.3	29.9	121.9	374.6	917.5	1,677.3	2,387.1
2003 5/	241.8	2.1	2.6	2.7	4.6	9.4	31.9	123.8	384.8	942.0	1,707.8	2,441.7
2002	243.8	2.0	2.8	2.7	4.9	9.8	32.5	125.9	395.5	975.3	1,716.5	2,507.7
2001	243.4	1.7	3.1	2.4	4.9	10.0	32.6	127.3	401.7	988.2	1,721.8	2,616.8
2000	244.4	2.9	3.1	2.6	5.0	9.7	32.3	127.2	412.0	1,002.1	1,750.2	2,714.1
1999	245.1	1.7	3.3	2.6	5.0	10.1	33.5	126.7	422.3	1,019.3	1,757.7	2,721.3
Non-Hispanic white, female												
2006 5/	214.7	2.2	2.1	2.0	2.9	9.2	35.9	113.0	288.5	631.3	1,044.4	1,336.7
2005 5/	215.1	*	2.2	2.2	3.3	8.9	37.5	113.9	293.6	634.4	1,049.5	1,353.2
2004 5/	215.3	1.8	2.5	2.1	3.3	9.3	37.5	112.9	299.8	649.8	1,052.0	1,364.5
2003 5/	217.6	2.3	2.2	2.4	3.4	9.8	38.6	115.1	308.9	657.6	1,062.4	1,399.1
2002	218.5	*	2.3	2.3	3.6	10.3	39.9	118.7	312.8	667.7	1,068.3	1,405.4
2001	218.4	1.8	2.8	2.4	3.6	10.7	40.4	121.9	317.3	669.7	1,064.4	1,425.1
2000	220.6	2.4	2.4	2.3	3.5	10.5	38.9	123.0	328.9	681.0	1,075.3	1,468.7
1999	218.4	*	2.7	2.1	3.6	10.3	39.7	123.0	330.5	684.6	1,060.5	1,444.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
Non-Hispanic black, both sexes												
2006 5/	165.5	*	2.5	2.2	4.1	11.9	45.8	174.0	450.1	890.6	1,420.3	1,700.4
2005 5/	167.4	*	2.5	2.6	4.1	12.7	46.2	179.7	466.0	906.1	1,435.5	1,764.4
2004 5/	167.5	*	2.5	2.6	5.0	12.4	48.3	181.8	467.2	931.9	1,465.9	1,794.3
2003 5/	170.0	*	2.3	2.7	4.1	12.6	50.5	189.7	480.9	941.1	1,508.2	1,867.5
2002	171.5	*	3.0	2.6	4.1	12.8	51.5	186.0	494.6	977.2	1,537.4	1,904.3
2001	172.8	*	2.0	2.7	4.4	13.4	52.8	188.7	494.5	1,005.6	1,559.5	1,947.5
2000	174.9	*	2.8	2.3	5.3	12.3	55.3	196.7	505.2	1,019.3	1,582.8	2,015.1
1999	176.9	*	2.0	2.8	5.6	12.6	55.3	196.8	530.5	1,033.6	1,592.9	2,024.0
Non-Hispanic black, male												
2006 5/	179.2	*	2.7	2.5	4.7	10.6	38.1	187.0	557.2	1,178.9	2,015.7	2,577.4
2005 5/	181.8	*	2.8	3.1	4.8	12.3	37.4	190.4	581.2	1,205.6	2,047.4	2,720.4
2004 5/	182.4	*	2.8	2.5	5.6	10.1	39.8	201.8	580.5	1,231.5	2,116.9	2,781.1
2003 5/	184.8	*	2.4	2.9	4.7	10.6	42.9	211.5	597.1	1,242.1	2,175.8	2,859.6
2002	187.8	*	2.7	3.1	4.6	11.7	44.3	202.3	621.8	1,293.5	2,247.6	3,008.4
2001	190.8	*	2.4	3.0	5.1	10.9	45.8	209.6	614.1	1,354.7	2,310.5	3,238.5
2000	194.9	*	2.6	2.6	6.4	10.5	49.6	219.0	637.4	1,382.7	2,372.3	3,290.1
1999	197.4	*	2.0	2.9	6.6	11.9	48.6	225.8	670.9	1,387.0	2,377.4	3,361.1
Non-Hispanic black, female												
2006 5/	153.0	*	2.4	2.0	3.4	13.2	52.7	162.8	363.9	684.6	1,082.5	1,342.7
2005 5/	154.3	*	2.1	2.2	3.5	13.1	54.0	170.6	373.3	692.1	1,088.7	1,379.5
2004 5/	153.9	*	2.2	2.7	4.4	14.5	55.8	164.6	376.3	718.3	1,098.4	1,404.3
2003 5/	156.5	*	2.1	2.4	3.5	14.5	57.2	170.9	387.6	727.0	1,131.2	1,483.2
2002	156.7	*	3.2	2.0	3.6	13.9	57.8	172.1	392.5	752.8	1,135.6	1,482.4
2001	156.5	*	*	2.3	3.8	15.7	59.0	170.7	398.3	759.4	1,137.8	1,469.3
2000	156.9	*	3.0	1.9	4.2	14.0	60.4	177.5	398.8	763.6	1,138.4	1,541.6
1999	158.3	*	2.0	2.7	4.8	13.3	61.3	172.0	417.9	784.2	1,153.6	1,524.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
All origins 3/, both sexes												
2006	2.6	*	*	*	*	0.1	0.6	2.6	5.8	9.6	13.5	16.3
2005	2.6	*	*	*	0.1	0.1	0.6	2.6	6.4	9.8	12.8	17.0
2004	2.7	*	*	*	0.1	0.1	0.6	2.9	6.1	10.1	13.7	18.0
2003	2.7	*	*	*	0.1	0.1	0.6	2.8	6.5	10.2	13.1	18.6
2002	2.7	*	*	*	0.1	0.1	0.6	2.7	6.8	10.4	13.5	18.7
2001	2.7	*	*	*	0.1	0.1	0.6	2.7	6.9	10.4	14.2	18.5
2000	2.7	*	*	*	0.1	0.1	0.6	2.8	6.8	10.7	13.2	19.4
1999	2.7	*	*	*	0.1	0.1	0.6	2.9	6.8	10.3	14.3	19.3
All origins 3/, male												
2006	3.6	*	*	*	*	0.1	0.9	4.1	9.3	14.8	20.4	21.9
2005	3.6	*	*	*	*	0.2	0.9	4.2	10.3	15.3	18.9	22.0
2004	3.7	*	*	*	*	0.2	0.8	4.6	10.0	15.6	20.3	24.9
2003	3.7	*	*	*	0.1	0.2	0.9	4.6	10.6	15.6	19.4	26.9
2002	3.7	*	*	*	*	0.2	0.8	4.4	10.9	15.9	19.6	26.9
2001	3.7	*	*	*	*	0.2	0.9	4.4	11.3	16.0	20.6	25.8
2000	3.6	*	*	*	*	0.1	1.0	4.5	10.9	16.0	19.2	27.1
1999	3.6	*	*	*	0.1	0.1	0.9	4.6	10.8	15.8	20.9	27.1
All origins 3/, female												
2006	1.6	*	*	*	*	*	0.4	1.1	2.4	5.2	8.8	13.7
2005	1.7	*	*	*	*	0.1	0.3	1.2	2.8	5.2	8.7	14.7
2004	1.7	*	*	*	*	*	0.4	1.2	2.5	5.6	9.2	14.9
2003	1.7	*	*	*	*	0.1	0.4	1.0	2.8	5.7	9.0	15.0
2002	1.7	*	*	*	*	*	0.4	1.1	3.0	5.8	9.5	15.1
2001	1.8	*	*	*	*	0.1	0.4	1.2	2.8	5.8	10.0	15.4
2000	1.8	*	*	*	*	0.1	0.3	1.3	2.9	6.3	9.2	16.3
1999	1.8	*	*	*	*	*	0.4	1.3	3.1	5.8	9.9	16.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Hispanic, both sexes												
2006	0.8	*	*	*	*	*	0.3	1.2	2.6	6.9	8.9	12.9
2005	0.7	*	*	*	*	*	*	1.2	3.5	5.4	10.4	*
2004	0.8	*	*	*	*	*	*	1.5	3.6	6.5	8.6	13.8
2003	0.8	*	*	*	*	*	*	1.3	3.6	6.5	9.8	15.0
2002	0.7	*	*	*	*	*	*	1.3	3.3	5.3	9.0	16.1
2001	0.8	*	*	*	*	*	*	1.6	3.7	6.9	10.5	19.4
2000	0.8	*	*	*	*	*	*	1.3	4.4	6.8	9.5	18.6
1999	0.8	*	*	*	*	*	*	1.4	4.0	6.9	13.1	14.0
Hispanic, male												
2006	1.1	*	*	*	*	*	*	1.7	4.7	11.7	15.8	*
2005	1.1	*	*	*	*	*	*	1.9	6.1	10.0	16.0	*
2004	1.1	*	*	*	*	*	*	2.6	6.0	11.0	12.9	*
2003	1.1	*	*	*	*	*	*	2.1	6.5	10.3	16.5	*
2002	1.1	*	*	*	*	*	*	2.1	5.7	9.7	14.8	*
2001	1.2	*	*	*	*	*	*	2.4	6.3	11.6	17.4	*
2000	1.2	*	*	*	*	*	*	2.2	7.7	12.2	13.8	40.3
1999	1.2	*	*	*	*	*	*	2.3	7.2	13.6	18.2	*
Hispanic, female												
2006	0.4	*	*	*	*	*	*	*	*	3.0	*	12.6
2005	0.4	*	*	*	*	*	*	*	*	*	6.5	*
2004	0.5	*	*	*	*	*	*	*	*	2.8	5.7	*
2003	0.5	*	*	*	*	*	*	*	*	3.5	5.3	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	3.1	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	6.6	*
1999	0.5	*	*	*	*	*	*	*	*	*	9.7	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Non-Hispanic 4/, both sexes												
2006	2.9	*	*	*	*	0.1	0.7	2.8	6.0	9.7	13.7	16.5
2005	2.9	*	*	*	0.1	0.2	0.7	2.8	6.6	10.2	12.9	17.6
2004	3.0	*	*	*	0.1	0.1	0.6	3.0	6.3	10.4	13.9	18.2
2003	3.0	*	*	*	0.1	0.2	0.7	2.9	6.8	10.5	13.3	18.7
2002	3.0	*	*	*	0.1	0.2	0.6	2.9	7.1	10.7	13.7	18.7
2001	3.0	*	*	*	0.1	0.2	0.7	2.8	7.1	10.6	14.3	18.4
2000	2.9	*	*	*	0.1	0.1	0.7	3.0	6.9	10.8	13.3	19.5
1999	2.9	*	*	*	0.1	0.1	0.7	3.0	7.0	10.5	14.3	19.4
Non-Hispanic 4/, male												
2006	4.0	*	*	*	*	0.2	0.9	4.4	9.7	15.0	20.6	22.3
2005	4.1	*	*	*	*	0.2	1.0	4.4	10.6	15.7	19.1	22.8
2004	4.1	*	*	*	*	0.2	0.9	4.8	10.3	15.8	20.7	24.9
2003	4.1	*	*	*	*	0.2	1.0	4.8	10.9	16.0	19.5	27.1
2002	4.1	*	*	*	*	0.2	0.8	4.6	11.3	16.2	19.8	26.7
2001	4.1	*	*	*	*	0.2	1.0	4.6	11.6	16.2	20.6	25.8
2000	3.9	*	*	*	*	0.1	1.0	4.6	11.1	16.1	19.4	26.5
1999	4.0	*	*	*	0.1	0.2	1.0	4.8	11.0	15.8	21.0	27.3
Non-Hispanic 4/, female												
2006	1.8	*	*	*	*	*	0.4	1.2	2.6	5.3	9.1	13.8
2005	1.9	*	*	*	*	0.1	0.4	1.2	2.9	5.5	8.8	15.2
2004	1.9	*	*	*	*	*	0.4	1.3	2.6	5.8	9.4	15.1
2003	1.9	*	*	*	*	*	0.4	1.1	2.9	5.9	9.2	15.0
2002	1.9	*	*	*	*	*	0.4	1.2	3.2	6.1	9.7	15.3
2001	1.9	*	*	*	*	0.1	0.4	1.2	2.9	6.0	10.2	15.3
2000	2.0	*	*	*	*	0.1	0.4	1.3	3.0	6.5	9.3	16.6
1999	2.0	*	*	*	*	*	0.4	1.4	3.2	6.0	9.9	16.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Non-Hispanic white, both sexes												
2006 5/	3.0	*	*	*	*	0.1	0.6	2.6	5.7	9.4	13.8	17.1
2005 5/	3.0	*	*	*	*	0.1	0.6	2.5	6.2	9.8	12.9	17.9
2004 5/	3.1	*	*	*	*	0.1	0.5	2.6	5.8	10.2	13.9	18.7
2003 5/	3.1	*	*	*	*	0.1	0.6	2.6	6.3	10.1	13.0	19.1
2002	3.1	*	*	*	*	0.1	0.5	2.5	6.5	10.6	13.9	18.9
2001	3.1	*	*	*	*	0.1	0.5	2.5	6.5	10.2	14.5	18.9
2000	3.0	*	*	*	*	0.1	0.6	2.5	6.1	10.4	13.3	19.8
1999	3.0	*	*	*	*	0.1	0.5	2.4	6.3	10.1	14.2	19.9
Non-Hispanic white, male												
2006 5/	4.1	*	*	*	*	*	0.8	4.1	9.0	14.1	20.4	23.0
2005 5/	4.1	*	*	*	*	*	0.8	3.9	9.7	14.8	18.5	22.9
2004 5/	4.1	*	*	*	*	0.2	0.7	4.0	9.4	15.0	20.4	24.8
2003 5/	4.1	*	*	*	*	0.2	1.0	4.2	9.9	14.9	18.7	27.4
2002	4.1	*	*	*	*	0.2	0.7	4.0	10.2	15.7	19.9	26.8
2001	4.1	*	*	*	*	0.2	0.8	4.0	10.4	15.3	20.5	26.4
2000	3.9	*	*	*	*	*	0.8	3.8	9.7	15.0	18.9	27.4
1999	3.9	*	*	*	*	*	0.8	3.8	9.7	15.1	20.3	27.5
Non-Hispanic white, female												
2006 5/	2.0	*	*	*	*	*	0.4	1.0	2.5	5.3	9.3	14.4
2005 5/	2.0	*	*	*	*	*	0.4	1.1	2.8	5.6	9.1	15.6
2004 5/	2.0	*	*	*	*	*	0.3	1.1	2.4	6.0	9.6	16.0
2003 5/	2.0	*	*	*	*	*	0.2	0.9	2.9	5.9	9.2	15.4
2002	2.1	*	*	*	*	*	0.3	1.0	2.9	6.2	9.8	15.5
2001	2.1	*	*	*	*	*	0.3	1.0	2.8	5.9	10.5	15.7
2000	2.1	*	*	*	*	*	0.4	1.2	2.8	6.5	9.6	16.8
1999	2.1	*	*	*	*	*	0.3	1.1	3.0	5.9	10.1	16.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Non-Hispanic black, both sexes												
2006 5/	2.7	*	*	*	*	*	0.8	4.4	9.7	13.9	13.8	10.1
2005 5/	2.9	*	*	*	*	*	0.9	5.1	10.9	14.2	13.7	12.8
2004 5/	3.1	*	*	*	*	*	1.3	6.2	11.4	14.1	15.3	12.7
2003 5/	3.0	*	*	*	*	*	1.1	5.2	11.5	15.7	15.6	15.5
2002	2.9	*	*	*	*	*	1.1	5.6	12.4	12.9	12.1	16.0
2001	3.0	*	*	*	*	*	1.3	5.3	12.9	15.7	13.9	13.1
2000	3.1	*	*	*	*	*	1.1	6.1	13.7	15.7	15.1	14.9
1999	3.3	*	*	*	*	*	1.2	7.3	13.6	15.4	16.8	14.5
Non-Hispanic black, male												
2006 5/	4.1	*	*	*	*	*	1.3	7.1	17.4	25.5	25.9	*
2005 5/	4.7	*	*	*	*	*	1.6	8.8	20.0	26.7	27.8	21.9
2004 5/	4.9	*	*	*	*	*	1.9	10.7	20.3	27.1	26.2	27.4
2003 5/	4.8	*	*	*	*	*	1.2	9.2	21.8	29.1	28.6	25.6
2002	4.4	*	*	*	*	*	1.6	9.8	22.5	23.0	18.8	26.3
2001	4.6	*	*	*	*	*	1.7	8.9	23.9	27.3	23.4	*
2000	4.9	*	*	*	*	*	1.8	10.7	24.7	28.6	29.0	*
1999	5.0	*	*	*	*	*	1.8	12.3	23.5	25.4	31.9	25.0
Non-Hispanic black, female												
2006 5/	1.3	*	*	*	*	*	*	2.0	3.6	5.7	6.9	8.6
2005 5/	1.2	*	*	*	*	*	*	1.9	3.5	5.2	5.7	9.2
2004 5/	1.4	*	*	*	*	*	0.9	2.3	4.3	4.8	9.2	*
2003 5/	1.4	*	*	*	*	*	1.1	1.8	3.3	6.2	8.3	11.6
2002	1.5	*	*	*	*	*	0.7	2.1	4.3	5.8	8.3	12.1
2001	1.5	*	*	*	*	*	0.9	2.1	4.0	7.5	8.6	10.7
2000	1.5	*	*	*	*	*	*	2.1	4.9	6.6	7.3	14.4
1999	1.7	*	*	*	*	*	0.7	3.0	5.7	8.4	8.4	10.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
All origins 3/, both sexes												
2006	4.6	*	*	*	*	0.1	0.6	3.5	10.4	19.5	26.9	25.6
2005	4.6	*	*	*	*	0.1	0.7	3.6	10.2	19.8	27.4	24.9
2004	4.4	*	*	*	*	0.1	0.7	3.4	10.3	19.2	26.7	25.8
2003	4.4	*	*	*	*	0.1	0.6	3.4	10.5	19.7	26.4	25.0
2002	4.4	*	*	*	*	*	0.7	3.5	10.4	20.2	26.6	24.6
2001	4.4	*	*	*	*	0.1	0.7	3.6	10.1	20.6	26.3	25.8
2000	4.3	*	*	*	*	0.1	0.7	3.6	10.8	20.0	25.6	25.9
1999	4.3	*	*	*	*	0.1	0.7	3.5	10.6	20.1	24.4	26.8
All origins 3/, male												
2006	7.3	*	*	*	*	0.1	1.1	6.0	18.4	34.3	48.8	48.5
2005	7.3	*	*	*	*	0.1	1.2	6.2	17.9	34.9	50.7	46.0
2004	7.0	*	*	*	*	0.1	1.1	5.9	17.9	33.4	48.8	48.5
2003	6.9	*	*	*	*	0.1	1.1	6.0	18.5	34.1	47.8	45.3
2002	6.9	*	*	*	*	*	1.1	6.0	18.2	35.6	48.3	43.4
2001	6.9	*	*	*	*	0.1	1.1	6.3	17.6	36.0	47.4	48.1
2000	6.7	*	*	*	*	0.2	1.2	6.3	18.6	34.1	45.8	47.8
1999	6.6	*	*	*	*	0.1	1.1	6.1	18.3	35.3	44.4	48.0
All origins 3/, female												
2006	2.0	*	*	*	*	*	0.2	1.1	3.0	7.0	12.0	14.9
2005	1.9	*	*	*	*	*	0.2	1.0	3.1	7.0	11.6	15.3
2004	2.0	*	*	*	*	*	0.2	1.0	3.2	7.3	11.9	15.7
2003	2.0	*	*	*	*	*	0.2	1.0	3.2	7.7	12.0	16.1
2002	2.0	*	*	*	*	*	0.2	1.0	3.2	7.4	12.2	16.5
2001	2.0	*	*	*	*	*	0.2	1.0	3.2	7.9	12.4	16.5
2000	2.1	*	*	*	*	*	0.1	1.0	3.5	8.3	12.4	17.1
1999	2.0	*	*	*	*	*	0.2	0.9	3.6	7.6	11.5	18.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
Hispanic, both sexes												
2006	1.2	*	*	*	*	*	0.3	1.5	4.9	9.9	15.9	14.9
2005	1.1	*	*	*	*	*	*	1.6	4.3	11.2	14.5	18.8
2004	1.1	*	*	*	*	*	*	1.0	4.7	10.3	15.0	18.6
2003	1.1	*	*	*	*	*	*	1.3	5.0	10.0	14.4	16.1
2002	1.1	*	*	*	*	*	0.4	1.5	4.1	10.2	15.2	23.3
2001	1.1	*	*	*	*	*	*	1.9	5.2	8.4	18.2	20.0
2000	1.1	*	*	*	*	*	*	1.2	5.0	11.1	19.4	19.9
1999	1.1	*	*	*	*	*	*	1.5	5.2	9.2	18.1	26.7
Hispanic, male												
2006	1.8	*	*	*	*	*	*	2.7	8.6	17.5	28.8	25.9
2005	1.8	*	*	*	*	*	*	2.7	7.8	20.7	26.7	37.4
2004	1.6	*	*	*	*	*	*	1.9	8.6	18.2	27.2	29.4
2003	1.6	*	*	*	*	*	*	2.4	8.0	17.2	28.4	*
2002	1.6	*	*	*	*	*	*	2.7	8.0	18.3	27.1	36.8
2001	1.7	*	*	*	*	*	*	3.6	8.5	16.0	30.4	41.9
2000	1.6	*	*	*	*	*	*	2.2	8.0	19.4	35.4	40.3
1999	1.7	*	*	*	*	*	*	2.7	10.2	16.3	32.8	49.6
Hispanic, female												
2006	0.5	*	*	*	*	*	*	*	1.5	3.7	6.9	*
2005	0.4	*	*	*	*	*	*	*	*	3.6	6.0	*
2004	0.5	*	*	*	*	*	*	*	*	4.0	6.7	*
2003	0.5	*	*	*	*	*	*	*	2.4	4.3	*	*
2002	0.4	*	*	*	*	*	*	*	*	3.6	7.0	16.6
2001	0.5	*	*	*	*	*	*	*	2.2	*	10.0	*
2000	0.5	*	*	*	*	*	*	*	2.4	4.6	8.6	*
1999	0.4	*	*	*	*	*	*	*	*	3.6	8.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
Non-Hispanic 4/, both sexes												
2006	5.2	*	*	*	*	0.1	0.7	3.7	10.9	20.2	27.6	26.1
2005	5.1	*	*	*	*	0.1	0.8	3.8	10.7	20.4	28.2	25.2
2004	5.0	*	*	*	*	0.1	0.7	3.7	10.7	19.9	27.3	26.2
2003	4.9	*	*	*	*	0.1	0.7	3.7	10.9	20.4	27.0	25.4
2002	4.9	*	*	*	*	*	0.7	3.7	10.8	20.8	27.2	24.5
2001	4.9	*	*	*	*	0.1	0.7	3.7	10.4	21.3	26.6	26.0
2000	4.8	*	*	*	*	0.1	0.7	3.8	11.2	20.4	25.8	26.1
1999	4.7	*	*	*	*	0.1	0.7	3.6	11.0	20.6	24.6	26.8
Non-Hispanic 4/, male												
2006	8.3	*	*	*	*	0.1	1.2	6.4	19.2	35.5	50.0	49.8
2005	8.2	*	*	*	*	0.2	1.4	6.6	18.7	35.9	52.2	46.3
2004	7.9	*	*	*	*	0.2	1.2	6.3	18.6	34.5	49.9	49.4
2003	7.8	*	*	*	*	0.1	1.2	6.4	19.3	35.3	48.8	46.3
2002	7.8	*	*	*	*	*	1.2	6.3	18.9	36.6	49.3	43.5
2001	7.7	*	*	*	*	0.1	1.3	6.5	18.2	37.1	48.1	48.4
2000	7.5	*	*	*	*	0.2	1.3	6.6	19.4	34.8	46.1	47.8
1999	7.4	*	*	*	*	0.2	1.2	6.4	18.8	36.2	44.7	48.0
Non-Hispanic 4/, female												
2006	2.2	*	*	*	*	*	0.2	1.2	3.1	7.3	12.3	15.1
2005	2.2	*	*	*	*	*	0.3	1.0	3.2	7.2	11.9	15.5
2004	2.2	*	*	*	*	*	0.2	1.1	3.4	7.6	12.2	15.8
2003	2.2	*	*	*	*	*	0.2	1.1	3.2	7.9	12.4	16.1
2002	2.2	*	*	*	*	*	0.3	1.1	3.4	7.6	12.5	16.3
2001	2.2	*	*	*	*	*	0.3	1.1	3.2	8.2	12.5	16.7
2000	2.3	*	*	*	*	*	0.1	1.0	3.6	8.5	12.6	17.3
1999	2.2	*	*	*	*	*	0.2	0.9	3.8	7.8	11.6	18.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
Non-Hispanic white, both sexes												
2006 5/	5.7	*	*	*	*	0.1	0.8	3.7	10.9	20.6	28.3	27.0
2005 5/	5.6	*	*	*	*	0.1	0.8	3.8	10.6	20.7	28.8	26.2
2004 5/	5.4	*	*	*	*	0.1	0.8	3.6	10.5	19.9	27.8	26.5
2003 5/	5.3	*	*	*	*	0.1	0.7	3.4	10.6	20.3	27.3	26.1
2002	5.3	*	*	*	*	*	0.7	3.6	10.4	20.5	27.3	25.1
2001	5.1	*	*	*	*	0.1	0.8	3.5	9.7	20.5	26.8	26.5
2000	5.1	*	*	*	*	0.1	0.7	3.5	10.4	20.0	25.5	26.7
1999	4.9	*	*	*	*	0.1	0.7	3.3	10.2	20.0	24.4	27.2
Non-Hispanic white, male												
2006 5/	9.2	*	*	*	*	*	1.4	6.4	19.4	36.0	51.7	51.3
2005 5/	9.1	*	*	*	*	0.2	1.4	6.7	18.7	36.4	53.4	49.0
2004 5/	8.7	*	*	*	*	*	1.3	6.3	18.4	34.6	51.0	50.1
2003 5/	8.5	*	*	*	*	*	1.2	6.0	18.8	35.1	49.9	47.9
2002	8.4	*	*	*	*	*	1.3	6.3	18.2	36.3	49.9	44.7
2001	8.1	*	*	*	*	*	1.3	6.1	17.0	36.0	48.6	49.0
2000	8.0	*	*	*	*	0.2	1.4	6.3	18.3	34.3	45.9	49.4
1999	7.7	*	*	*	*	0.2	1.2	6.0	17.5	35.4	44.4	48.0
Non-Hispanic white, female												
2006 5/	2.3	*	*	*	*	*	0.2	1.1	2.8	7.2	12.2	15.7
2005 5/	2.3	*	*	*	*	*	0.2	0.8	3.0	7.1	11.8	15.8
2004 5/	2.3	*	*	*	*	*	0.2	0.9	3.0	7.2	12.0	16.0
2003 5/	2.3	*	*	*	*	*	0.2	0.9	2.8	7.7	12.0	16.5
2002	2.3	*	*	*	*	*	0.2	0.9	3.0	7.1	12.2	16.6
2001	2.3	*	*	*	*	*	0.2	1.0	2.7	7.5	12.4	17.1
2000	2.3	*	*	*	*	*	*	0.8	3.0	8.0	12.0	17.5
1999	2.2	*	*	*	*	*	0.2	0.8	3.3	7.1	11.3	18.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
Non-Hispanic black, both sexes												
2006 5/	3.8	*	*	*	*	*	0.6	5.2	13.9	22.5	24.9	18.3
2005 5/	3.9	*	*	*	*	*	0.9	4.9	14.5	22.9	27.7	17.5
2004 5/	4.1	*	*	*	*	*	0.7	5.3	15.8	25.2	27.4	21.9
2003 5/	4.3	*	*	*	*	*	0.9	6.1	17.3	26.1	28.4	21.4
2002	4.3	*	*	*	*	*	1.0	5.2	17.7	28.9	29.9	20.6
2001	4.7	*	*	*	*	*	1.0	6.1	18.9	32.8	30.0	24.0
2000	4.7	*	*	*	*	*	0.8	6.8	20.2	28.6	34.1	22.6
1999	4.7	*	*	*	*	*	1.0	6.6	20.3	30.2	31.4	23.6
Non-Hispanic black, male												
2006 5/	5.8	*	*	*	*	*	1.0	8.4	23.2	40.6	43.1	35.7
2005 5/	6.0	*	*	*	*	*	1.5	8.0	25.2	41.1	52.1	26.6
2004 5/	6.0	*	*	*	*	*	1.0	8.5	26.0	42.2	47.6	39.6
2003 5/	6.5	*	*	*	*	*	1.4	10.4	29.4	45.2	46.7	38.4
2002	6.5	*	*	*	*	*	1.5	8.4	30.7	49.8	52.2	38.3
2001	7.2	*	*	*	*	*	1.6	10.5	33.0	56.2	55.5	47.4
2000	7.0	*	*	*	*	*	1.2	11.5	34.8	47.6	58.9	38.7
1999	7.2	*	*	*	*	*	1.7	11.5	35.3	51.1	57.2	49.9
Non-Hispanic black, female												
2006 5/	2.0	*	*	*	*	*	*	2.4	6.4	9.6	14.5	11.2
2005 5/	2.0	*	*	*	*	*	*	2.3	5.9	9.9	13.8	13.8
2004 5/	2.4	*	*	*	*	*	*	2.5	7.6	13.1	15.9	14.9
2003 5/	2.4	*	*	*	*	*	*	2.5	7.5	12.5	18.1	14.9
2002	2.4	*	*	*	*	*	*	2.4	7.2	14.0	17.2	13.8
2001	2.4	*	*	*	*	*	*	2.3	7.5	16.2	15.7	15.4
2000	2.6	*	*	*	*	*	*	2.7	8.5	15.2	20.1	16.6
1999	2.4	*	*	*	*	*	*	2.4	8.3	15.5	17.0	13.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
All origins 3/, both sexes												
2006	3.8	*	*	*	*	0.3	1.1	2.7	5.9	13.0	25.6	35.8
2005	3.9	*	*	*	0.0	0.3	1.0	2.6	5.9	14.0	26.6	38.5
2004	4.0	*	*	*	*	0.3	1.1	2.7	6.5	14.9	26.3	42.2
2003	4.2	*	*	*	*	0.3	1.1	3.0	6.7	15.2	27.6	43.6
2002	4.2	*	*	*	0.1	0.3	1.0	2.8	6.9	15.4	28.7	47.6
2001	4.3	*	*	*	0.1	0.3	1.0	2.9	7.0	16.5	30.2	45.4
2000	4.5	*	*	*	*	0.3	1.1	2.9	7.4	16.6	31.0	52.7
1999	4.6	*	*	*	*	0.3	1.1	3.0	7.8	17.1	31.6	50.1
All origins 3/, male												
2006	4.5	*	*	*	*	0.3	1.1	3.4	8.3	18.0	36.8	50.2
2005	4.6	*	*	*	*	0.3	1.2	3.3	8.2	19.7	37.4	54.2
2004	4.8	*	*	*	*	0.3	1.2	3.5	9.1	20.6	37.2	62.9
2003	4.9	*	*	*	*	0.3	1.2	3.8	9.5	21.9	38.4	60.0
2002	5.0	*	*	*	*	0.4	1.1	3.7	9.5	22.5	40.6	67.2
2001	5.1	*	*	*	*	0.4	1.1	3.8	10.2	23.2	43.1	62.6
2000	5.3	*	*	*	*	0.3	1.3	3.8	10.5	23.7	43.9	75.6
1999	5.5	*	*	*	*	0.4	1.3	4.2	11.0	24.2	46.5	71.3
All origins 3/, female												
2006	3.1	*	*	*	*	0.3	1.1	2.0	3.7	8.8	18.0	29.1
2005	3.2	*	*	*	*	0.3	0.8	1.9	3.7	9.2	19.3	31.2
2004	3.3	*	*	*	*	0.3	0.9	2.0	4.1	10.1	19.0	32.9
2003	3.4	*	*	*	*	0.3	1.0	2.1	4.1	9.7	20.3	36.4
2002	3.5	*	*	*	0.1	0.3	0.9	1.9	4.5	9.5	20.8	39.1
2001	3.5	*	*	*	*	0.3	0.9	2.0	4.1	10.9	21.7	38.3
2000	3.7	*	*	*	*	0.3	0.9	2.1	4.6	10.8	22.6	43.4
1999	3.7	*	*	*	*	0.3	1.0	1.9	4.9	11.3	21.9	41.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
Hispanic, both sexes												
2006	2.9	*	*	*	*	0.6	1.7	3.9	10.2	20.4	37.2	47.0
2005	3.0	*	*	*	*	0.5	1.5	4.6	9.3	20.7	44.0	50.9
2004	3.3	*	*	*	*	0.5	2.0	4.6	11.7	24.4	44.0	62.8
2003	3.1	*	*	*	*	0.5	1.6	5.1	11.4	23.8	43.6	62.7
2002	3.1	*	*	*	*	0.5	1.7	4.2	11.0	21.9	47.2	72.1
2001	3.2	*	*	*	*	0.6	1.8	5.4	11.3	26.4	45.6	63.8
2000	3.1	*	*	*	*	0.4	2.0	4.8	9.8	28.4	50.2	65.7
1999	3.1	*	*	*	*	0.6	1.7	4.1	11.7	26.2	49.8	73.7
Hispanic, male												
2006	3.1	*	*	*	*	0.5	1.4	4.5	13.1	27.1	49.3	55.3
2005	3.4	*	*	*	*	0.5	1.5	5.0	12.1	27.6	65.8	76.0
2004	3.6	*	*	*	*	0.5	2.0	4.7	14.8	32.0	66.9	84.1
2003	3.4	*	*	*	*	*	1.5	5.7	14.1	33.2	63.7	73.9
2002	3.4	*	*	*	*	0.7	1.5	4.7	13.7	32.9	61.6	107.1
2001	3.6	*	*	*	*	*	1.9	6.2	15.0	37.1	60.4	95.2
2000	3.6	*	*	*	*	*	2.0	5.0	13.7	43.3	70.8	98.8
1999	3.5	*	*	*	*	0.7	1.6	5.2	16.4	35.6	73.5	75.5
Hispanic, female												
2006	2.7	*	*	*	*	0.7	2.1	3.3	7.5	14.9	28.8	42.5
2005	2.6	*	*	*	*	*	1.4	4.2	6.8	15.2	28.8	37.8
2004	2.9	*	*	*	*	*	2.1	4.5	8.8	18.3	28.2	51.9
2003	2.8	*	*	*	*	*	1.6	4.5	9.0	16.3	29.8	57.0
2002	2.7	*	*	*	*	*	1.9	3.8	8.7	13.2	37.4	54.7
2001	2.8	*	*	*	*	0.7	1.7	4.6	8.1	18.0	35.7	48.4
2000	2.7	*	*	*	*	*	2.1	4.6	6.3	16.8	36.3	49.5
1999	2.7	*	*	*	*	*	1.8	3.0	7.6	18.8	34.0	72.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
Non-Hispanic 4/, both sexes												
2006	3.9	*	*	*	*	0.2	0.9	2.5	5.5	12.4	24.9	35.2
2005	4.0	*	*	*	*	0.2	0.9	2.4	5.6	13.5	25.5	37.8
2004	4.2	*	*	*	*	0.3	0.9	2.5	6.0	14.2	25.3	41.2
2003	4.3	*	*	*	*	0.3	1.0	2.7	6.3	14.6	26.7	42.7
2002	4.4	*	*	*	0.1	0.3	0.9	2.6	6.5	14.9	27.7	46.3
2001	4.5	*	*	*	*	0.2	0.9	2.6	6.6	15.8	29.4	44.7
2000	4.7	*	*	*	*	0.3	0.9	2.7	7.2	15.8	30.1	52.0
1999	4.7	*	*	*	*	0.3	1.0	2.9	7.5	16.5	30.8	49.2
Non-Hispanic 4/, male												
2006	4.8	*	*	*	*	0.2	1.0	3.3	7.9	17.3	36.0	49.9
2005	4.8	*	*	*	*	0.2	1.1	3.1	7.9	19.1	35.6	53.0
2004	5.0	*	*	*	*	0.2	1.1	3.4	8.6	19.7	35.5	61.7
2003	5.2	*	*	*	*	0.3	1.1	3.6	9.2	21.0	36.9	59.2
2002	5.3	*	*	*	*	0.3	1.0	3.6	9.1	21.7	39.4	65.0
2001	5.4	*	*	*	*	0.3	1.0	3.6	9.8	22.3	42.1	61.0
2000	5.5	*	*	*	*	0.3	1.2	3.7	10.3	22.4	42.6	74.5
1999	5.7	*	*	*	*	0.3	1.2	4.0	10.6	23.4	45.3	71.0
Non-Hispanic 4/, female												
2006	3.1	*	*	*	*	0.3	0.9	1.8	3.4	8.3	17.3	28.4
2005	3.3	*	*	*	*	0.3	0.7	1.6	3.4	8.7	18.7	30.9
2004	3.3	*	*	*	*	0.3	0.8	1.7	3.7	9.5	18.4	32.1
2003	3.5	*	*	*	*	0.3	0.9	1.9	3.7	9.2	19.8	35.5
2002	3.5	*	*	*	*	0.3	0.7	1.7	4.1	9.2	19.9	38.3
2001	3.6	*	*	*	*	0.2	0.8	1.7	3.7	10.4	21.0	37.9
2000	3.8	*	*	*	*	0.3	0.7	1.8	4.4	10.4	21.9	42.9
1999	3.8	*	*	*	*	0.3	0.9	1.8	4.7	10.8	21.3	40.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
Non-Hispanic white, both sexes												
2006 5/	3.6	*	*	*	*	0.2	0.7	2.1	4.6	10.4	21.4	31.5
2005 5/	3.7	*	*	*	*	0.1	0.7	1.9	4.6	11.3	22.0	33.8
2004 5/	3.9	*	*	*	*	0.2	0.7	2.0	4.9	11.8	22.3	37.6
2003 5/	4.0	*	*	*	*	0.2	0.8	2.1	5.4	12.6	23.3	37.5
2002	4.1	*	*	*	*	0.2	0.7	2.1	5.4	12.6	24.1	40.7
2001	4.2	*	*	*	*	0.2	0.7	2.1	5.6	13.5	25.8	40.0
2000	4.4	*	*	*	*	0.2	0.7	2.2	6.1	13.5	26.1	46.3
1999	4.4	*	*	*	*	0.2	0.7	2.3	6.1	14.4	27.0	43.5
Non-Hispanic white, male												
2006 5/	4.5	*	*	*	*	*	0.7	2.8	6.6	14.7	31.3	45.2
2005 5/	4.4	*	*	*	*	*	0.8	2.5	6.5	15.9	30.8	47.2
2004 5/	4.7	*	*	*	*	0.2	0.9	2.7	7.1	16.3	31.1	56.9
2003 5/	4.8	*	*	*	*	0.2	0.9	2.8	7.7	18.0	32.4	52.3
2002	5.0	*	*	*	*	0.2	0.8	2.9	7.8	18.6	35.0	58.0
2001	5.0	*	*	*	*	0.2	0.8	2.8	8.2	18.8	37.4	55.7
2000	5.2	*	*	*	*	0.2	0.8	3.0	8.9	19.3	37.4	67.3
1999	5.4	*	*	*	*	0.2	0.9	3.3	8.7	20.6	39.9	62.6
Non-Hispanic white, female												
2006 5/	2.8	*	*	*	*	0.2	0.6	1.4	2.7	6.7	14.5	25.2
2005 5/	3.0	*	*	*	*	*	0.6	1.3	2.8	7.3	15.9	27.7
2004 5/	3.1	*	*	*	*	0.2	0.5	1.2	2.9	8.0	16.3	28.9
2003 5/	3.3	*	*	*	*	0.2	0.6	1.4	3.1	8.0	17.2	31.0
2002	3.2	*	*	*	*	0.2	0.5	1.2	3.2	7.5	16.8	33.3
2001	3.4	*	*	*	*	*	0.6	1.3	3.1	9.0	18.1	33.5
2000	3.5	*	*	*	*	0.2	0.5	1.4	3.6	8.6	18.7	37.9
1999	3.5	*	*	*	*	0.2	0.6	1.4	3.7	9.1	18.6	35.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
Non-Hispanic black, both sexes												
2006 5/	5.0	*	*	*	*	0.4	1.8	4.5	11.1	24.8	52.0	68.2
2005 5/	5.2	*	*	*	*	0.4	1.8	4.6	11.5	27.0	52.6	75.4
2004 5/	5.4	*	*	*	*	0.5	1.7	5.4	12.9	30.8	50.9	71.4
2003 5/	5.5	*	*	*	*	0.4	1.8	5.8	12.9	27.5	55.0	86.0
2002	5.8	*	*	*	*	0.6	1.6	5.5	13.2	31.5	57.7	100.1
2001	5.8	*	*	*	*	0.5	2.2	5.5	13.1	31.0	59.1	89.3
2000	6.1	*	*	*	*	0.7	2.1	5.8	14.1	32.3	64.3	102.7
1999	6.3	*	*	*	*	0.7	2.3	6.0	16.1	31.5	65.1	103.1
Non-Hispanic black, male												
2006 5/	6.0	*	*	*	*	*	2.0	6.2	16.7	34.8	82.0	85.2
2005 5/	6.4	*	*	*	*	*	2.6	6.5	18.3	41.6	72.0	117.0
2004 5/	6.7	*	*	*	*	*	1.9	7.3	19.6	48.2	75.9	99.6
2003 5/	6.8	*	*	*	*	*	2.0	8.5	20.0	44.8	80.5	107.6
2002	6.9	*	*	*	*	0.8	1.9	7.9	19.5	47.3	80.9	136.7
2001	6.9	*	*	*	*	*	2.4	8.0	20.0	49.2	87.9	106.3
2000	7.2	*	*	*	*	0.8	3.0	8.1	20.9	47.0	91.0	138.6
1999	7.6	*	*	*	*	*	2.8	8.7	23.3	46.4	101.6	149.8
Non-Hispanic black, female												
2006 5/	4.1	*	*	*	*	*	1.6	2.9	6.7	17.6	35.0	61.3
2005 5/	4.1	*	*	*	*	*	1.1	3.0	6.0	16.5	41.6	58.6
2004 5/	4.3	*	*	*	*	*	1.5	3.7	7.6	18.4	36.9	60.2
2003 5/	4.4	*	*	*	*	*	1.6	3.5	7.3	15.2	40.6	77.6
2002	4.9	*	*	*	*	*	1.4	3.4	8.1	20.3	44.6	86.1
2001	4.7	*	*	*	*	*	2.0	3.3	7.6	18.1	43.0	83.0
2000	5.2	*	*	*	*	*	1.4	3.9	8.6	22.0	49.3	89.4
1999	5.1	*	*	*	*	*	1.9	3.7	10.2	21.0	44.6	85.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
All origins 3/, both sexes												
2006	17.9	*	*	*	0.1	0.8	3.0	10.4	27.3	61.9	122.8	207.5
2005	18.0	*	*	*	0.1	0.8	3.0	10.2	26.9	63.3	124.5	216.9
2004	18.3	*	*	*	0.1	0.7	3.0	10.3	28.2	64.7	128.9	225.6
2003	19.2	*	*	*	0.1	0.7	3.0	10.9	29.8	70.5	134.7	239.7
2002	19.7	*	*	*	0.1	0.7	3.2	11.4	31.7	73.3	137.4	241.4
2001	20.0	*	*	*	0.1	0.8	3.2	11.6	31.5	75.3	139.4	255.7
2000	20.4	*	*	*	0.1	0.7	3.0	11.2	32.7	76.7	146.0	270.4
1999	20.5	*	*	*	0.1	0.7	3.0	11.1	33.5	77.4	146.7	271.7
All origins 3/, male												
2006	18.3	*	*	*	0.1	0.8	3.2	11.6	33.0	76.8	149.5	234.6
2005	18.4	*	*	*	0.1	0.8	3.2	11.5	32.9	79.6	151.6	245.9
2004	18.6	*	*	*	0.2	0.7	3.2	11.9	34.2	80.6	156.9	254.7
2003	19.6	*	*	*	0.1	0.8	3.2	12.5	36.0	88.0	164.4	273.3
2002	20.1	*	*	*	0.2	0.8	3.6	13.0	39.4	91.3	167.4	275.1
2001	20.2	*	*	*	0.1	0.8	3.5	13.1	38.3	93.7	169.5	298.3
2000	20.6	*	*	*	0.1	0.8	3.4	12.8	39.6	95.4	179.5	315.6
1999	20.6	*	*	*	0.1	0.8	3.0	12.5	40.5	95.1	181.9	324.0
All origins 3/, female												
2006	17.5	*	*	*	0.1	0.8	2.9	9.2	22.1	49.3	104.6	194.9
2005	17.6	*	*	*	0.1	0.7	2.8	9.0	21.2	49.5	106.1	203.6
2004	18.0	*	*	*	0.1	0.6	2.9	8.7	22.7	51.4	110.0	212.5
2003	18.9	*	*	*	*	0.7	2.7	9.3	24.0	55.9	114.8	224.8
2002	19.2	*	*	*	*	0.7	2.8	9.8	24.5	58.3	117.5	226.8
2001	19.8	*	*	*	0.1	0.7	2.9	10.2	25.3	60.1	119.6	237.8
2000	20.2	*	*	*	*	0.6	2.7	9.8	26.4	61.2	124.2	251.9
1999	20.3	*	*	*	0.1	0.6	2.9	9.7	27.0	62.9	123.8	250.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
Hispanic, both sexes												
2006	6.2	*	*	*	*	0.5	2.0	7.5	20.4	45.7	93.7	142.4
2005	6.0	*	*	*	*	0.5	1.9	7.1	18.4	46.5	92.4	146.7
2004	5.9	*	*	*	*	0.6	1.9	7.2	21.2	47.5	87.5	147.5
2003	6.2	*	*	*	*	0.5	2.1	8.2	23.6	49.7	95.5	144.6
2002	6.1	*	*	*	*	0.4	2.1	7.7	23.0	51.2	95.6	165.7
2001	6.1	*	*	*	*	0.3	2.3	7.2	22.5	52.1	96.5	193.8
2000	6.0	*	*	*	*	0.5	2.3	8.4	23.1	49.2	98.2	181.1
1999	6.0	*	*	*	*	0.5	1.7	8.3	21.0	56.1	101.0	176.8
Hispanic, male												
2006	6.3	*	*	*	*	0.5	1.7	7.8	25.5	60.3	112.9	157.9
2005	6.1	*	*	*	*	*	1.8	8.0	23.3	60.3	111.6	176.8
2004	6.2	*	*	*	*	0.5	2.2	8.5	26.6	59.4	114.1	172.3
2003	6.6	*	*	*	*	0.5	2.3	10.0	29.8	64.8	120.4	172.4
2002	6.4	*	*	*	*	*	2.0	9.3	31.1	67.1	121.1	174.0
2001	6.2	*	*	*	*	*	2.7	7.4	28.2	66.5	116.7	257.1
2000	6.2	*	*	*	*	*	2.0	9.0	28.2	63.5	131.3	235.8
1999	6.1	*	*	*	*	*	1.5	9.1	24.1	72.7	134.0	211.4
Hispanic, female												
2006	6.1	*	*	*	*	*	2.3	7.2	15.8	33.7	80.3	134.2
2005	5.8	*	*	*	*	0.7	2.0	6.2	13.9	35.3	79.1	131.0
2004	5.6	*	*	*	*	0.6	1.4	5.8	16.3	38.0	69.1	134.7
2003	5.7	*	*	*	*	*	1.8	6.4	18.0	37.6	78.5	130.5
2002	5.7	*	*	*	*	*	2.1	6.1	15.7	38.6	78.1	161.6
2001	5.9	*	*	*	*	*	1.9	7.1	17.4	40.8	82.8	162.9
2000	5.8	*	*	*	*	*	2.7	7.8	18.5	38.0	75.9	154.3
1999	5.9	*	*	*	*	*	2.0	7.5	18.3	43.0	79.1	160.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
Non-Hispanic 4/, both sexes												
2006	19.9	*	*	*	0.1	0.9	3.2	10.7	27.8	63.1	124.5	210.6
2005	20.0	*	*	*	0.1	0.8	3.1	10.5	27.5	64.4	126.2	220.1
2004	20.3	*	*	*	0.1	0.7	3.2	10.6	28.8	65.9	130.9	228.8
2003	21.3	*	*	*	0.1	0.8	3.1	11.1	30.2	71.9	136.5	243.2
2002	21.7	*	*	*	0.1	0.8	3.3	11.8	32.3	74.6	139.1	244.0
2001	22.0	*	*	*	0.1	0.9	3.3	12.0	32.2	76.7	141.1	257.4
2000	22.4	*	*	*	0.1	0.7	3.1	11.5	33.3	78.2	147.8	273.1
1999	22.4	*	*	*	0.1	0.7	3.1	11.3	34.3	78.5	148.1	274.3
Non-Hispanic 4/, male												
2006	20.4	*	*	*	*	0.9	3.5	12.0	33.5	77.9	151.7	238.4
2005	20.5	*	*	*	0.1	0.9	3.4	11.9	33.7	80.9	153.7	249.2
2004	20.7	*	*	*	0.2	0.8	3.3	12.3	34.8	81.9	159.1	258.5
2003	21.7	*	*	*	*	0.9	3.4	12.7	36.4	89.4	166.3	277.3
2002	22.3	*	*	*	0.2	0.9	3.8	13.4	39.9	92.7	169.3	279.2
2001	22.3	*	*	*	0.1	0.9	3.6	13.6	38.9	95.2	171.8	299.6
2000	22.8	*	*	*	0.1	0.8	3.6	13.1	40.3	97.2	181.3	318.6
1999	22.7	*	*	*	0.1	0.9	3.2	12.8	41.6	96.2	183.2	327.5
Non-Hispanic 4/, female												
2006	19.4	*	*	*	*	0.9	3.0	9.4	22.6	50.4	105.9	197.7
2005	19.4	*	*	*	*	0.7	2.9	9.3	21.8	50.5	107.6	206.8
2004	19.9	*	*	*	*	0.7	3.1	8.9	23.2	52.3	112.0	215.6
2003	20.9	*	*	*	*	0.7	2.8	9.6	24.4	57.2	116.6	228.1
2002	21.2	*	*	*	*	0.7	2.9	10.2	25.2	59.6	119.1	228.8
2001	21.7	*	*	*	*	0.8	3.0	10.5	25.9	61.3	121.0	239.9
2000	22.1	*	*	*	*	0.6	2.7	9.9	26.9	62.6	126.1	254.7
1999	22.2	*	*	*	*	0.6	3.0	9.9	27.6	63.9	125.3	253.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
Non-Hispanic white, both sexes												
2006 5/	21.2	*	*	*	0.1	0.8	3.0	9.9	26.3	61.3	122.0	210.5
2005 5/	21.2	*	*	*	0.1	0.8	3.0	9.6	25.7	62.5	124.0	218.8
2004 5/	21.7	*	*	*	0.1	0.7	3.0	9.8	27.3	63.8	129.4	228.5
2003 5/	22.7	*	*	*	*	0.7	2.8	10.1	28.5	69.9	135.0	242.1
2002	23.2	*	*	*	0.1	0.8	3.0	10.9	30.6	73.0	136.8	243.2
2001	23.5	*	*	*	0.1	0.8	3.1	11.1	30.6	74.7	138.5	256.9
2000	24.0	*	*	*	0.1	0.6	2.8	10.6	31.7	76.5	145.7	271.9
1999	24.0	*	*	*	0.1	0.7	2.9	10.6	32.8	76.7	145.9	273.4
Non-Hispanic white, male												
2006 5/	21.6	*	*	*	*	0.8	3.2	11.1	31.8	75.5	148.1	236.7
2005 5/	21.8	*	*	*	*	0.8	3.3	10.9	31.5	78.5	151.2	248.8
2004 5/	22.2	*	*	*	*	0.7	3.1	11.5	33.3	79.4	156.5	259.6
2003 5/	23.2	*	*	*	*	0.8	3.1	11.6	34.5	87.4	164.5	276.8
2002	23.8	*	*	*	0.2	0.9	3.4	12.5	38.0	91.1	165.8	277.6
2001	23.9	*	*	*	*	0.9	3.3	12.8	37.4	93.2	169.3	298.2
2000	24.2	*	*	*	*	0.8	3.2	12.2	38.2	95.1	178.4	315.8
1999	24.3	*	*	*	*	0.8	3.1	12.1	39.7	94.8	180.6	328.2
Non-Hispanic white, female												
2006 5/	20.7	*	*	*	*	0.7	2.7	8.7	21.0	49.0	103.9	198.2
2005 5/	20.6	*	*	*	*	0.7	2.6	8.3	20.2	48.6	105.2	205.1
2004 5/	21.3	*	*	*	*	0.6	2.8	8.1	21.7	50.3	110.9	214.6
2003 5/	22.2	*	*	*	*	0.7	2.5	8.5	22.8	54.9	115.0	226.8
2002	22.7	*	*	*	*	0.7	2.7	9.4	23.7	57.6	117.3	228.3
2001	23.1	*	*	*	*	0.7	2.8	9.5	24.2	59.0	118.1	239.8
2000	23.8	*	*	*	*	0.5	2.4	9.0	25.5	60.9	124.1	254.1
1999	23.7	*	*	*	*	0.6	2.7	9.0	26.2	61.5	123.2	251.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
Non-Hispanic black, both sexes												
2006 5/	18.1	*	*	*	*	1.3	5.2	17.5	44.2	90.5	171.2	244.7
2005 5/	18.2	*	*	*	*	1.2	4.4	18.1	46.1	92.4	170.7	264.1
2004 5/	17.8	*	*	*	*	1.0	4.6	16.9	45.8	96.4	167.6	262.5
2003 5/	18.8	*	*	*	*	1.0	5.2	19.2	48.9	102.2	175.0	282.9
2002	18.9	*	*	*	*	0.7	5.1	19.0	50.5	100.6	185.9	278.3
2001	19.2	*	*	*	*	1.1	5.0	19.5	50.6	105.9	189.7	289.0
2000	19.4	*	*	*	*	1.0	5.2	20.0	51.1	106.3	192.5	312.9
1999	19.3	*	*	*	*	1.1	4.9	18.1	53.7	107.2	194.5	308.8
Non-Hispanic black, male												
2006 5/	18.9	*	*	*	*	1.3	5.7	20.5	53.0	118.6	223.6	314.3
2005 5/	18.5	*	*	*	*	1.5	4.5	20.2	58.5	116.4	213.6	297.8
2004 5/	18.0	*	*	*	*	1.1	4.7	19.5	54.7	122.8	218.8	277.4
2003 5/	19.0	*	*	*	*	1.0	5.4	22.4	59.9	128.2	217.0	321.8
2002	19.3	*	*	*	*	0.9	6.0	21.8	62.7	123.0	240.9	332.6
2001	19.0	*	*	*	*	1.2	5.8	21.4	59.3	130.7	229.9	352.3
2000	19.8	*	*	*	*	1.1	6.1	22.7	63.7	133.7	244.4	381.6
1999	19.2	*	*	*	*	1.2	4.8	20.6	67.0	127.8	245.8	351.8
Non-Hispanic black, female												
2006 5/	17.3	*	*	*	*	1.3	4.7	14.9	37.2	70.5	141.5	216.4
2005 5/	17.9	*	*	*	*	0.8	4.3	16.3	36.2	75.3	146.3	250.5
2004 5/	17.6	*	*	*	*	1.0	4.4	14.8	38.7	77.6	138.8	256.6
2003 5/	18.7	*	*	*	*	0.9	5.0	16.4	40.1	83.8	151.2	267.8
2002	18.4	*	*	*	*	*	4.3	16.7	40.7	84.7	154.9	257.5
2001	19.4	*	*	*	*	1.0	4.3	17.9	43.6	88.4	167.1	265.6
2000	19.1	*	*	*	*	1.0	4.4	17.6	40.9	87.1	163.2	287.4
1999	19.5	*	*	*	*	0.9	5.0	15.9	43.1	92.7	165.8	292.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
All origins 3/, both sexes												
2006	5.5	*	*	*	0.1	0.2	0.8	5.6	12.2	20.2	31.9	33.5
2005	5.4	*	0.1	*	0.1	0.2	0.9	5.9	11.3	20.1	32.0	32.5
2004	5.2	*	0.1	*	0.1	0.2	0.9	5.6	10.7	19.9	31.3	32.4
2003	5.1	*	0.1	*	0.1	0.2	0.9	5.4	9.9	19.8	30.7	33.4
2002	4.9	*	0.2	*	0.1	0.3	0.9	4.9	9.6	19.6	29.3	35.0
2001	4.7	*	*	*	0.1	0.2	0.9	4.8	9.3	18.8	29.0	34.1
2000	4.6	*	0.1	0.0	0.1	0.2	1.0	4.5	9.3	18.7	28.8	32.7
1999	4.4	*	0.2	*	0.1	0.2	1.0	4.1	8.8	19.0	28.0	30.6
All origins 3/, male												
2006	7.3	*	*	*	*	0.3	1.1	8.9	19.0	29.3	44.6	43.8
2005	7.2	*	*	*	*	0.3	1.2	9.6	17.9	29.0	45.0	44.8
2004	6.8	*	*	*	0.1	0.3	1.3	9.2	16.6	28.3	42.7	41.5
2003	6.6	*	*	*	0.1	0.3	1.3	8.9	14.9	27.9	44.6	47.5
2002	6.4	*	*	*	*	0.3	1.4	8.2	14.5	28.5	41.6	46.2
2001	6.1	*	*	*	*	0.2	1.3	7.7	13.9	26.8	41.0	50.3
2000	5.9	*	*	*	0.1	0.3	1.3	7.2	13.8	27.0	42.1	44.7
1999	5.6	*	0.3	*	0.1	0.2	1.4	6.5	12.9	27.5	39.0	40.1
All origins 3/, female												
2006	3.8	*	*	*	*	0.2	0.5	2.3	5.8	12.6	23.2	28.7
2005	3.7	*	*	*	*	0.1	0.6	2.4	5.2	12.6	23.2	26.8
2004	3.7	*	*	*	*	0.1	0.4	2.2	5.3	12.8	23.6	28.3
2003	3.5	*	*	*	*	0.2	0.5	2.1	5.3	13.0	21.4	27.2
2002	3.4	*	*	*	*	0.2	0.5	1.8	5.1	12.2	21.1	30.2
2001	3.4	*	*	*	*	0.2	0.4	1.9	5.0	12.1	21.1	27.2
2000	3.3	*	*	*	*	0.1	0.6	1.9	5.1	11.8	20.1	27.8
1999	3.3	*	*	*	*	0.2	0.6	1.8	5.1	11.9	20.9	26.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
Hispanic, both sexes												
2006	4.1	*	*	*	*	*	0.8	7.3	17.2	31.5	51.6	55.2
2005	4.2	*	*	*	*	*	0.9	8.0	18.2	33.8	52.1	54.8
2004	3.8	*	*	*	*	*	1.0	7.2	16.6	30.8	48.6	50.9
2003	3.8	*	*	*	*	*	0.8	7.5	16.1	32.5	49.8	51.8
2002	3.5	*	*	*	*	*	0.8	5.9	15.7	34.1	44.2	59.9
2001	3.4	*	*	*	*	*	0.8	6.7	14.1	28.8	52.1	52.5
2000	3.3	*	*	*	*	*	1.2	5.6	13.7	31.1	51.8	58.4
1999	3.2	*	*	*	*	*	0.9	5.7	13.1	31.3	49.8	53.3
Hispanic, male												
2006	5.2	*	*	*	*	*	1.0	11.5	27.5	45.7	69.2	62.0
2005	5.4	*	*	*	*	*	1.1	12.6	28.8	46.9	68.5	71.0
2004	4.9	*	*	*	*	*	1.6	11.2	27.5	42.3	62.9	71.4
2003	4.8	*	*	*	*	*	1.2	12.5	24.3	45.2	66.0	55.4
2002	4.5	*	*	*	*	*	1.1	9.8	24.1	48.9	59.9	75.3
2001	4.3	*	*	*	*	*	1.3	11.1	21.5	41.1	68.0	64.8
2000	4.2	*	*	*	*	*	1.7	9.1	20.5	43.9	74.7	72.6
1999	3.9	*	*	*	*	*	1.1	9.6	20.7	42.8	63.1	66.9
Hispanic, female												
2006	2.9	*	*	*	*	*	*	3.1	7.7	20.0	39.2	51.5
2005	3.0	*	*	*	*	*	0.7	3.3	8.6	23.1	40.7	46.3
2004	2.6	*	*	*	*	*	*	3.1	6.7	21.6	38.8	40.3
2003	2.7	*	*	*	*	*	*	2.5	8.8	22.4	38.7	50.0
2002	2.5	*	*	*	*	*	*	2.1	8.2	22.4	33.4	52.2
2001	2.5	*	*	*	*	*	*	2.3	7.6	19.1	41.4	46.5
2000	2.4	*	*	*	*	*	*	2.2	7.6	21.1	36.3	51.4
1999	2.4	*	*	*	*	*	*	1.9	6.3	22.2	40.9	46.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
Non-Hispanic 4/, both sexes												
2006	5.8	*	*	*	0.1	0.2	0.8	5.3	11.7	19.3	30.6	32.3
2005	5.6	*	*	*	0.1	0.3	0.9	5.7	10.7	19.0	30.8	31.3
2004	5.4	*	0.2	*	0.1	0.2	0.8	5.5	10.2	19.0	30.3	31.5
2003	5.2	*	*	*	0.1	0.2	0.9	5.2	9.4	18.8	29.7	32.6
2002	5.1	*	0.2	*	0.1	0.3	0.9	4.8	9.1	18.6	28.4	33.9
2001	4.9	*	*	*	0.1	0.2	0.9	4.6	8.9	18.1	27.9	33.2
2000	4.8	*	0.2	*	0.1	0.2	0.9	4.3	8.9	17.9	27.7	31.7
1999	4.6	*	0.2	*	0.1	0.2	1.0	3.9	8.4	18.2	27.1	29.7
Non-Hispanic 4/, male												
2006	7.7	*	*	*	*	0.3	1.2	8.5	18.2	27.9	43.0	42.6
2005	7.5	*	*	*	*	0.4	1.2	9.2	16.9	27.6	43.5	43.4
2004	7.1	*	*	*	0.1	0.3	1.3	9.0	15.6	27.2	41.4	40.0
2003	6.9	*	*	*	*	0.3	1.3	8.4	14.1	26.6	43.5	47.1
2002	6.6	*	*	*	*	0.4	1.4	8.0	13.7	27.1	40.5	44.8
2001	6.3	*	*	*	*	0.2	1.3	7.4	13.2	25.8	39.7	49.7
2000	6.2	*	*	*	*	0.3	1.3	6.9	13.3	25.9	40.5	43.4
1999	5.8	*	*	*	0.1	0.3	1.5	6.2	12.2	26.5	37.9	38.7
Non-Hispanic 4/, female												
2006	3.9	*	*	*	*	0.2	0.5	2.3	5.6	12.0	22.2	27.6
2005	3.8	*	*	*	*	0.1	0.5	2.3	4.9	11.8	22.1	25.8
2004	3.8	*	*	*	*	0.2	0.4	2.1	5.2	12.1	22.7	27.7
2003	3.6	*	*	*	*	0.2	0.5	2.1	5.0	12.3	20.4	26.2
2002	3.6	*	*	*	*	0.2	0.5	1.7	4.8	11.4	20.5	29.2
2001	3.5	*	*	*	*	0.2	0.4	1.9	4.8	11.7	20.2	26.4
2000	3.4	*	*	*	*	0.1	0.6	1.9	4.9	11.2	19.4	27.0
1999	3.4	*	*	*	*	0.2	0.5	1.8	4.9	11.3	20.0	26.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
Non-Hispanic white, both sexes												
2006 5/	5.6	*	*	*	0.1	0.2	0.6	4.4	9.9	17.7	29.1	31.7
2005 5/	5.5	*	*	*	*	0.2	0.6	4.8	9.0	17.2	29.2	30.4
2004 5/	5.3	*	*	*	0.1	0.1	0.6	4.5	8.6	17.1	28.8	30.1
2003 5/	5.1	*	*	*	0.1	0.1	0.7	4.2	8.0	17.0	28.2	31.1
2002	5.0	*	*	*	*	0.2	0.7	3.9	7.7	17.0	27.4	32.6
2001	4.8	*	*	*	0.1	0.1	0.6	3.7	7.6	16.6	26.5	32.4
2000	4.6	*	*	*	*	0.1	0.6	3.4	7.5	16.2	26.5	30.8
1999	4.5	*	0.2	*	0.1	0.1	0.7	3.1	7.1	16.8	26.1	28.9
Non-Hispanic white, male												
2006 5/	7.3	*	*	*	*	0.2	0.7	7.1	14.9	25.2	41.2	42.3
2005 5/	7.2	*	*	*	*	0.2	0.8	7.6	14.0	24.8	41.5	43.4
2004 5/	6.8	*	*	*	*	0.2	0.9	7.4	12.8	24.4	40.0	38.0
2003 5/	6.7	*	*	*	*	*	1.0	6.7	11.9	24.3	41.6	45.0
2002	6.4	*	*	*	*	0.2	1.1	6.4	11.5	24.6	38.7	43.6
2001	6.1	*	*	*	*	*	0.9	5.9	11.2	24.2	38.0	48.9
2000	5.9	*	*	*	*	0.2	0.8	5.3	11.0	23.6	39.2	41.6
1999	5.7	*	*	*	*	*	1.1	4.8	10.2	24.6	36.7	39.0
Non-Hispanic white, female												
2006 5/	4.0	*	*	*	*	0.2	0.5	1.9	5.1	11.1	20.6	26.8
2005 5/	3.8	*	*	*	*	*	0.5	2.0	4.3	10.6	20.8	24.5
2004 5/	3.8	*	*	*	*	*	0.4	1.7	4.7	10.9	21.2	26.5
2003 5/	3.6	*	*	*	*	*	0.5	1.7	4.2	10.7	19.2	25.0
2002	3.6	*	*	*	*	0.2	0.4	1.4	4.2	10.4	19.9	27.8
2001	3.4	*	*	*	*	*	0.3	1.5	4.1	10.2	18.9	25.5
2000	3.4	*	*	*	*	*	0.4	1.6	4.3	10.1	18.2	26.5
1999	3.4	*	*	*	*	*	0.4	1.6	4.2	10.2	19.1	24.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
Non-Hispanic black, both sexes												
2006 5/	5.7	*	*	*	*	0.4	1.3	9.6	22.3	24.6	33.4	31.6
2005 5/	5.6	*	*	*	*	0.5	1.5	9.9	20.5	25.4	33.6	28.9
2004 5/	5.4	*	*	*	*	0.4	1.4	9.8	17.7	26.1	34.7	36.6
2003 5/	5.1	*	*	*	*	0.5	1.4	10.0	16.5	23.1	31.7	41.9
2002	4.7	*	*	*	*	0.5	1.5	8.9	15.0	23.3	30.9	38.1
2001	4.6	*	*	*	*	*	1.7	8.9	15.1	21.9	32.3	35.3
2000	4.5	*	*	*	*	0.4	1.8	8.8	14.5	23.7	29.1	31.8
1999	4.3	*	*	*	*	0.4	1.7	7.4	13.4	24.6	29.7	34.9
Non-Hispanic black, male												
2006 5/	8.3	*	*	*	*	*	2.2	15.7	40.6	40.0	45.9	34.8
2005 5/	8.2	*	*	*	*	*	2.2	17.3	36.4	40.2	51.0	31.4
2004 5/	7.6	*	*	*	*	*	2.3	16.9	30.5	41.3	44.2	47.8
2003 5/	7.0	*	*	*	*	*	2.2	17.3	26.4	33.8	44.7	63.9
2002	6.6	*	*	*	*	*	2.3	15.4	24.4	38.2	46.3	41.6
2001	6.1	*	*	*	*	*	2.7	15.1	24.2	30.3	42.6	42.7
2000	6.3	*	*	*	*	*	2.5	15.0	24.1	37.3	40.4	48.1
1999	5.7	*	*	*	*	*	2.5	13.1	20.3	36.8	41.6	30.9
Non-Hispanic black, female												
2006 5/	3.3	*	*	*	*	*	*	4.4	7.5	13.5	26.3	30.3
2005 5/	3.2	*	*	*	*	*	0.9	3.5	7.8	14.8	23.7	28.0
2004 5/	3.4	*	*	*	*	*	*	3.7	7.4	15.2	29.3	32.1
2003 5/	3.3	*	*	*	*	*	*	3.6	8.6	15.5	24.3	33.4
2002	3.0	*	*	*	*	*	0.8	3.3	7.4	12.8	22.2	36.8
2001	3.3	*	*	*	*	*	0.8	3.6	7.8	16.0	26.4	32.5
2000	2.9	*	*	*	*	*	1.2	3.4	6.7	14.1	22.8	25.7
1999	3.1	*	*	*	*	*	0.9	2.5	7.9	15.9	23.1	36.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
All origins 3/, both sexes												
2006	11.2	*	*	*	*	0.1	1.2	6.3	19.9	44.8	78.6	96.9
2005	11.1	*	*	*	*	0.2	1.3	6.5	19.7	44.5	77.7	97.4
2004	10.8	*	*	*	*	0.2	1.2	6.3	19.8	44.3	76.0	97.7
2003	10.6	*	*	*	*	0.2	1.2	6.2	19.1	44.2	75.4	95.2
2002	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.8	75.0	94.5
2001	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.5	74.1	100.1
2000	10.4	*	*	*	*	0.1	1.2	6.3	19.4	44.1	76.5	96.4
1999	10.4	*	*	*	*	0.1	1.3	6.4	19.8	45.0	73.1	99.7
All origins 3/, male												
2006	11.2	*	*	*	*	0.1	1.4	7.8	24.5	52.3	86.2	102.1
2005	11.1	*	*	*	*	0.2	1.5	7.7	24.3	51.8	86.0	102.8
2004	10.9	*	*	*	*	0.2	1.5	8.1	24.1	52.6	84.6	102.0
2003	10.5	*	*	*	*	0.2	1.5	7.7	23.1	50.9	85.4	95.6
2002	10.5	*	*	*	*	0.1	1.6	7.8	23.6	51.4	84.0	100.9
2001	10.3	*	*	*	*	0.1	1.4	7.6	23.6	50.8	84.0	105.1
2000	10.3	*	*	*	*	0.2	1.4	8.0	23.6	50.4	85.9	102.9
1999	10.4	*	*	*	*	0.2	1.6	8.3	23.9	52.4	81.8	107.5
All origins 3/, female												
2006	11.1	*	*	*	*	0.1	0.9	4.9	15.7	38.4	73.4	94.4
2005	11.0	*	*	*	*	0.1	1.1	5.2	15.5	38.2	72.1	94.9
2004	10.7	*	*	*	*	0.1	0.9	4.7	15.8	37.3	70.1	95.8
2003	10.6	*	*	*	*	0.1	1.0	4.8	15.3	38.6	68.8	95.0
2002	10.5	*	*	*	*	*	1.0	4.4	15.1	39.3	69.1	91.7
2001	10.6	*	*	*	*	0.1	1.1	4.7	15.2	39.4	67.6	98.1
2000	10.5	*	*	*	*	0.1	1.0	4.6	15.5	38.8	70.4	93.8
1999	10.5	*	*	*	*	0.1	1.0	4.7	16.0	39.0	67.4	96.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
Hispanic, both sexes												
2006	3.9	*	*	*	*	*	0.7	4.3	13.8	33.7	59.9	77.1
2005	4.1	*	*	*	*	*	0.9	4.4	15.5	36.0	63.8	78.7
2004	3.8	*	*	*	*	*	0.7	4.1	14.4	35.3	62.8	69.9
2003	3.7	*	*	*	*	*	0.8	4.1	14.4	35.1	60.3	79.8
2002	3.5	*	*	*	*	*	0.9	4.0	13.8	34.7	55.5	79.3
2001	3.6	*	*	*	*	*	0.8	3.7	14.2	36.9	61.3	83.8
2000	3.5	*	*	*	*	*	0.7	4.1	13.7	36.9	61.8	82.9
1999	3.3	*	*	*	*	*	0.9	4.6	13.3	32.6	56.7	84.2
Hispanic, male												
2006	3.8	*	*	*	*	*	0.8	5.0	15.4	39.7	68.9	65.4
2005	4.0	*	*	*	*	*	1.1	5.2	19.2	40.5	65.8	77.2
2004	3.7	*	*	*	*	*	1.0	4.7	17.4	42.4	68.3	57.4
2003	3.6	*	*	*	*	*	1.0	5.0	17.8	40.0	67.1	70.8
2002	3.3	*	*	*	*	*	1.2	5.3	15.7	40.9	55.8	60.2
2001	3.4	*	*	*	*	*	0.9	4.3	16.2	42.3	68.9	62.9
2000	3.4	*	*	*	*	*	0.8	5.2	16.6	45.2	62.9	80.6
1999	3.2	*	*	*	*	*	1.2	5.7	15.2	40.0	56.3	86.3
Hispanic, female												
2006	4.1	*	*	*	*	*	*	3.7	12.3	28.9	53.6	83.3
2005	4.2	*	*	*	*	*	0.8	3.5	12.1	32.3	62.3	79.5
2004	3.9	*	*	*	*	*	*	3.5	11.8	29.5	59.0	76.3
2003	3.8	*	*	*	*	*	*	3.2	11.3	31.2	55.6	84.4
2002	3.7	*	*	*	*	*	*	2.8	12.1	29.8	55.4	88.7
2001	3.9	*	*	*	*	*	0.8	3.2	12.5	32.6	56.2	94.0
2000	3.6	*	*	*	*	*	*	3.0	11.1	30.4	61.1	84.1
1999	3.5	*	*	*	*	*	*	3.5	11.6	26.8	56.9	83.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
Non-Hispanic 4/, both sexes												
2006	12.4	*	*	*	*	0.1	1.2	6.5	20.4	45.6	79.7	97.7
2005	12.2	*	*	*	*	0.2	1.3	6.7	20.0	45.0	78.5	98.1
2004	11.9	*	*	*	*	0.2	1.3	6.6	20.2	44.9	76.5	98.9
2003	11.7	*	*	*	*	0.2	1.3	6.4	19.4	44.8	76.1	95.7
2002	11.6	*	*	*	*	0.1	1.4	6.3	19.6	45.4	75.8	94.9
2001	11.5	*	*	*	*	0.1	1.3	6.4	19.6	44.9	74.6	100.6
2000	11.4	*	*	*	*	0.1	1.3	6.5	19.8	44.4	77.0	96.6
1999	11.4	*	*	*	*	0.1	1.4	6.6	20.2	45.6	73.5	100.1
Non-Hispanic 4/, male												
2006	12.6	*	*	*	*	*	1.5	8.1	25.3	53.2	87.1	104.0
2005	12.3	*	*	*	*	0.2	1.6	8.0	24.7	52.6	87.2	104.1
2004	12.1	*	*	*	*	0.2	1.6	8.4	24.6	53.2	85.4	104.1
2003	11.7	*	*	*	*	0.2	1.6	8.0	23.5	51.6	86.2	96.5
2002	11.7	*	*	*	*	0.1	1.7	8.0	24.1	52.1	85.3	102.4
2001	11.4	*	*	*	*	0.2	1.5	7.9	24.2	51.2	84.6	106.4
2000	11.3	*	*	*	*	0.2	1.5	8.3	24.1	50.7	86.7	103.5
1999	11.4	*	*	*	*	0.2	1.7	8.4	24.4	52.9	82.7	108.2
Non-Hispanic 4/, female												
2006	12.3	*	*	*	*	0.1	1.0	5.0	16.0	39.1	74.6	94.8
2005	12.1	*	*	*	*	0.1	1.1	5.4	15.7	38.7	72.6	95.4
2004	11.8	*	*	*	*	*	1.0	4.8	16.1	37.9	70.6	96.5
2003	11.6	*	*	*	*	0.1	1.0	4.9	15.6	39.1	69.3	95.3
2002	11.5	*	*	*	*	*	1.1	4.6	15.4	39.8	69.5	91.7
2001	11.5	*	*	*	*	0.1	1.2	4.9	15.4	39.8	68.0	98.2
2000	11.4	*	*	*	*	*	1.0	4.7	15.8	39.2	70.6	93.8
1999	11.4	*	*	*	*	*	1.0	4.8	16.4	39.6	67.6	96.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
Non-Hispanic white, both sexes												
2006 5/	13.4	*	*	*	*	0.1	1.2	6.3	20.0	44.9	79.5	96.9
2005 5/	13.2	*	*	*	*	0.2	1.3	6.4	19.6	44.5	77.7	97.6
2004 5/	12.8	*	*	*	*	0.2	1.2	6.3	19.5	44.0	76.1	98.1
2003 5/	12.5	*	*	*	*	0.2	1.2	6.0	18.7	44.1	75.4	94.9
2002	12.4	*	*	*	*	0.1	1.4	6.0	19.0	44.3	75.1	94.6
2001	12.3	*	*	*	*	0.1	1.3	6.1	19.3	44.1	73.7	99.4
2000	12.2	*	*	*	*	0.2	1.2	6.3	19.2	43.4	75.7	94.7
1999	12.1	*	*	*	*	0.1	1.3	6.3	19.5	44.7	72.1	98.9
Non-Hispanic white, male												
2006 5/	13.7	*	*	*	*	*	1.5	7.9	24.8	53.0	88.0	105.3
2005 5/	13.3	*	*	*	*	0.2	1.6	7.6	24.2	52.5	86.7	104.3
2004 5/	13.1	*	*	*	*	0.2	1.5	8.1	23.8	52.0	85.8	105.0
2003 5/	12.6	*	*	*	*	0.2	1.5	7.5	22.8	51.4	86.0	96.8
2002	12.6	*	*	*	*	*	1.7	7.7	23.5	51.2	85.4	102.9
2001	12.4	*	*	*	*	0.2	1.5	7.7	24.0	50.3	84.5	108.0
2000	12.2	*	*	*	*	0.2	1.5	8.0	23.4	50.0	86.1	103.0
1999	12.1	*	*	*	*	0.2	1.6	8.0	23.4	52.1	81.6	108.1
Non-Hispanic white, female												
2006 5/	13.1	*	*	*	*	*	0.9	4.7	15.4	38.0	73.6	93.0
2005 5/	13.0	*	*	*	*	*	1.1	5.1	15.2	37.7	71.5	94.5
2004 5/	12.6	*	*	*	*	*	0.9	4.5	15.4	37.1	69.5	95.1
2003 5/	12.4	*	*	*	*	*	1.0	4.6	14.8	38.0	68.1	94.0
2002	12.2	*	*	*	*	*	1.1	4.3	14.7	38.3	68.1	91.0
2001	12.3	*	*	*	*	*	1.2	4.6	14.9	38.8	66.6	95.8
2000	12.2	*	*	*	*	*	1.0	4.5	15.3	37.9	68.9	91.4
1999	12.1	*	*	*	*	*	1.0	4.7	15.7	38.5	65.9	95.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
Non-Hispanic black, both sexes												
2006 5/	10.2	*	*	*	*	*	1.6	9.4	28.5	58.4	93.1	114.7
2005 5/	10.0	*	*	*	*	*	1.6	9.8	27.5	57.7	95.9	111.7
2004 5/	9.8	*	*	*	*	*	1.7	9.5	28.8	59.0	89.1	115.1
2003 5/	9.8	*	*	*	*	*	1.9	10.3	27.7	57.5	94.0	113.3
2002	9.6	*	*	*	*	*	1.8	8.9	28.3	61.7	90.9	108.3
2001	9.4	*	*	*	*	*	1.6	9.2	25.9	59.0	93.3	122.0
2000	9.7	*	*	*	*	*	1.7	9.2	27.9	60.0	98.9	121.2
1999	9.7	*	*	*	*	*	2.0	9.9	28.8	60.4	95.0	119.0
Non-Hispanic black, male												
2006 5/	9.7	*	*	*	*	*	2.0	11.5	36.2	67.3	93.4	92.5
2005 5/	9.8	*	*	*	*	*	1.7	12.1	34.7	65.7	106.0	101.8
2004 5/	9.8	*	*	*	*	*	2.1	11.6	36.5	73.8	91.3	102.6
2003 5/	9.6	*	*	*	*	*	2.5	13.6	33.7	64.8	99.0	106.5
2002	9.3	*	*	*	*	*	2.2	11.6	34.9	69.5	97.3	102.8
2001	8.9	*	*	*	*	*	1.8	11.4	30.8	68.2	100.5	100.5
2000	9.3	*	*	*	*	*	2.1	12.1	34.6	66.9	108.0	110.4
1999	9.7	*	*	*	*	*	2.4	13.6	37.4	68.7	107.6	115.3
Non-Hispanic black, female												
2006 5/	10.6	*	*	*	*	*	1.3	7.5	22.4	52.1	92.9	123.7
2005 5/	10.3	*	*	*	*	*	1.5	7.9	21.7	52.0	90.2	115.7
2004 5/	9.9	*	*	*	*	*	1.2	7.7	22.6	48.5	87.9	120.1
2003 5/	10.0	*	*	*	*	*	1.3	7.4	22.8	52.4	91.3	116.0
2002	9.8	*	*	*	*	*	1.5	6.5	23.0	56.2	87.2	110.4
2001	9.9	*	*	*	*	*	1.4	7.3	21.9	52.4	89.3	130.0
2000	10.0	*	*	*	*	*	1.5	6.8	22.5	55.1	93.8	125.2
1999	9.7	*	*	*	*	*	1.7	6.7	21.9	54.6	87.9	120.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
All origins 3/, both sexes												
2006	1.3	*	*	*	*	*	0.1	1.0	3.1	5.7	7.1	6.3
2005	1.3	*	*	*	*	*	0.1	1.0	3.0	6.2	7.0	6.3
2004	1.2	*	*	*	*	*	0.1	1.1	3.2	5.7	6.6	6.5
2003	1.3	*	*	*	*	*	0.2	1.0	3.4	6.2	6.9	6.5
2002	1.3	*	*	*	*	*	0.2	1.1	3.4	5.9	7.6	5.5
2001	1.3	*	*	*	*	*	0.2	1.1	3.6	6.4	7.2	6.7
2000	1.4	*	*	*	*	*	0.1	1.1	3.6	6.6	8.1	6.5
1999	1.4	*	*	*	*	*	0.2	1.1	3.8	6.7	7.3	7.0
All origins 3/, male												
2006	2.0	*	*	*	*	*	0.2	1.7	5.1	9.8	13.4	14.8
2005	2.0	*	*	*	*	*	0.2	1.5	5.1	10.7	13.1	15.7
2004	2.0	*	*	*	*	*	0.2	1.8	5.4	10.1	13.0	15.6
2003	2.1	*	*	*	*	*	0.3	1.8	5.8	10.8	13.7	15.3
2002	2.1	*	*	*	*	*	0.3	1.8	5.8	10.0	14.9	13.2
2001	2.2	*	*	*	*	*	0.3	1.8	6.1	11.1	14.2	18.3
2000	2.2	*	*	*	*	*	0.2	1.8	6.2	11.4	15.7	17.4
1999	2.2	*	*	*	*	*	0.3	1.9	6.4	11.4	14.1	17.8
All origins 3/, female												
2006	0.5	*	*	*	*	*	*	0.4	1.2	2.2	2.8	2.3
2005	0.5	*	*	*	*	*	*	0.5	1.0	2.4	2.8	1.9
2004	0.5	*	*	*	*	*	*	0.4	1.2	2.0	2.3	2.4
2003	0.5	*	*	*	*	*	*	0.3	1.2	2.4	2.4	2.7
2002	0.5	*	*	*	*	*	*	0.4	1.2	2.5	2.8	2.2
2001	0.5	*	*	*	*	*	0.1	0.4	1.3	2.5	2.6	1.9
2000	0.6	*	*	*	*	*	*	0.4	1.2	2.7	3.1	2.1
1999	0.6	*	*	*	*	*	*	0.3	1.4	2.9	2.8	2.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
Hispanic, both sexes												
2006	0.4	*	*	*	*	*	*	0.6	1.6	3.3	6.8	*
2005	0.4	*	*	*	*	*	*	0.6	1.3	3.9	7.3	*
2004	0.5	*	*	*	*	*	*	0.5	2.0	4.9	6.9	*
2003	0.4	*	*	*	*	*	*	0.5	2.1	3.6	5.9	*
2002	0.4	*	*	*	*	*	*	*	2.2	4.1	5.5	*
2001	0.4	*	*	*	*	*	*	0.6	2.1	5.3	5.2	*
2000	0.4	*	*	*	*	*	*	*	1.7	4.2	6.9	*
1999	0.5	*	*	*	*	*	*	0.8	1.8	5.3	8.3	*
Hispanic, male												
2006	0.7	*	*	*	*	*	*	1.0	3.0	6.7	14.5	*
2005	0.7	*	*	*	*	*	*	1.0	2.5	8.5	15.4	*
2004	0.8	*	*	*	*	*	*	1.0	4.0	10.6	15.0	*
2003	0.7	*	*	*	*	*	*	*	4.3	7.6	11.9	*
2002	0.7	*	*	*	*	*	*	*	4.3	8.0	11.1	*
2001	0.7	*	*	*	*	*	*	*	3.9	10.6	10.7	*
2000	0.7	*	*	*	*	*	*	*	3.5	7.8	14.8	*
1999	0.9	*	*	*	*	*	*	1.6	3.5	10.5	16.2	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
Non-Hispanic 4/, both sexes												
2006	1.4	*	*	*	*	*	0.2	1.1	3.2	5.8	7.1	6.3
2005	1.4	*	*	*	*	*	0.2	1.0	3.2	6.4	7.0	6.2
2004	1.4	*	*	*	*	*	0.2	1.1	3.3	5.7	6.6	6.5
2003	1.4	*	*	*	*	*	0.2	1.1	3.5	6.3	6.9	6.5
2002	1.4	*	*	*	*	*	0.2	1.1	3.5	6.0	7.7	5.4
2001	1.5	*	*	*	*	*	0.2	1.1	3.7	6.4	7.3	6.7
2000	1.5	*	*	*	*	*	0.1	1.2	3.7	6.7	8.1	6.5
1999	1.5	*	*	*	*	*	0.2	1.1	3.9	6.8	7.2	6.8
Non-Hispanic 4/, male												
2006	2.3	*	*	*	*	*	0.2	1.8	5.3	10.0	13.3	14.7
2005	2.3	*	*	*	*	*	0.2	1.6	5.4	10.9	13.0	15.7
2004	2.2	*	*	*	*	*	0.2	1.9	5.5	10.0	12.8	15.6
2003	2.3	*	*	*	*	*	0.3	1.8	5.9	10.9	13.7	15.0
2002	2.3	*	*	*	*	*	0.3	1.9	5.9	10.1	15.0	13.2
2001	2.4	*	*	*	*	*	0.3	1.9	6.1	11.1	14.3	18.4
2000	2.4	*	*	*	*	*	0.2	1.9	6.4	11.5	15.7	17.3
1999	2.4	*	*	*	*	*	0.3	1.9	6.6	11.4	14.0	17.3
Non-Hispanic 4/, female												
2006	0.6	*	*	*	*	*	*	0.4	1.3	2.3	2.9	2.4
2005	0.6	*	*	*	*	*	*	0.5	1.1	2.5	2.9	1.9
2004	0.6	*	*	*	*	*	*	0.4	1.3	2.1	2.4	2.5
2003	0.6	*	*	*	*	*	*	0.3	1.3	2.5	2.4	2.7
2002	0.6	*	*	*	*	*	*	0.4	1.3	2.6	2.9	2.1
2001	0.6	*	*	*	*	*	0.1	0.4	1.4	2.6	2.7	1.8
2000	0.6	*	*	*	*	*	*	0.5	1.2	2.8	3.1	2.1
1999	0.6	*	*	*	*	*	*	0.3	1.4	3.0	2.8	2.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
Non-Hispanic white, both sexes												
2006 5/	1.5	*	*	*	*	*	0.1	1.0	3.0	5.5	7.0	6.2
2005 5/	1.4	*	*	*	*	*	0.1	0.9	2.8	6.0	6.9	5.9
2004 5/	1.4	*	*	*	*	*	0.1	1.0	3.0	5.4	6.3	6.4
2003 5/	1.4	*	*	*	*	*	0.2	0.9	3.2	6.0	6.6	6.1
2002	1.4	*	*	*	*	*	0.1	0.9	3.1	5.8	7.3	5.5
2001	1.5	*	*	*	*	*	0.2	0.9	3.3	6.0	6.9	6.5
2000	1.5	*	*	*	*	*	0.1	1.0	3.2	6.2	7.7	6.5
1999	1.5	*	*	*	*	*	0.1	0.9	3.5	6.4	7.1	6.5
Non-Hispanic white, male												
2006 5/	2.3	*	*	*	*	*	0.2	1.5	4.9	9.1	12.8	14.5
2005 5/	2.3	*	*	*	*	*	0.2	1.4	4.6	10.0	12.7	14.8
2004 5/	2.2	*	*	*	*	*	0.2	1.6	4.9	9.4	12.1	15.0
2003 5/	2.3	*	*	*	*	*	0.3	1.5	5.2	9.9	12.8	14.1
2002	2.3	*	*	*	*	*	0.2	1.5	5.1	9.5	14.2	13.3
2001	2.3	*	*	*	*	*	0.3	1.5	5.5	10.2	13.3	18.3
2000	2.4	*	*	*	*	*	0.2	1.5	5.4	10.5	15.0	17.4
1999	2.3	*	*	*	*	*	0.2	1.6	5.8	10.7	13.6	16.0
Non-Hispanic white, female												
2006 5/	0.7	*	*	*	*	*	*	0.4	1.2	2.4	3.0	2.3
2005 5/	0.6	*	*	*	*	*	*	0.4	1.1	2.5	3.0	1.8
2004 5/	0.6	*	*	*	*	*	*	0.4	1.3	2.0	2.4	2.5
2003 5/	0.6	*	*	*	*	*	*	0.3	1.2	2.5	2.4	2.6
2002	0.6	*	*	*	*	*	*	0.3	1.3	2.6	2.7	2.2
2001	0.6	*	*	*	*	*	*	0.3	1.2	2.4	2.7	1.6
2000	0.6	*	*	*	*	*	*	0.4	1.1	2.6	3.0	2.1
1999	0.7	*	*	*	*	*	*	0.3	1.3	2.9	2.9	2.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
Non-Hispanic black, both sexes												
2006 5/	1.7	*	*	*	*	*	*	2.2	6.0	10.6	10.2	8.2
2005 5/	1.8	*	*	*	*	*	*	2.3	7.2	11.6	9.8	10.4
2004 5/	1.7	*	*	*	*	*	*	2.3	6.6	10.3	11.4	8.1
2003 5/	1.9	*	*	*	*	*	*	2.6	7.7	11.6	11.9	11.3
2002	1.9	*	*	*	*	*	0.5	2.7	7.7	10.6	13.9	*
2001	1.9	*	*	*	*	*	0.4	2.9	7.9	11.5	12.0	8.4
2000	2.0	*	*	*	*	*	*	2.9	9.2	12.9	14.3	7.0
1999	1.9	*	*	*	*	*	0.6	3.0	8.7	11.7	10.0	8.4
Non-Hispanic black, male												
2006 5/	2.8	*	*	*	*	*	*	4.0	10.8	21.2	22.5	19.2
2005 5/	3.0	*	*	*	*	*	*	3.5	14.0	22.5	21.0	29.5
2004 5/	2.9	*	*	*	*	*	*	4.3	12.6	20.2	25.4	23.4
2003 5/	3.3	*	*	*	*	*	*	4.6	14.0	24.0	28.3	27.7
2002	3.1	*	*	*	*	*	0.8	4.6	15.1	19.8	29.6	*
2001	3.1	*	*	*	*	*	*	5.2	14.2	21.4	28.2	*
2000	3.4	*	*	*	*	*	*	5.2	17.5	24.6	29.0	*
1999	3.2	*	*	*	*	*	1.0	5.3	16.3	22.1	22.5	25.0
Non-Hispanic black, female												
2006 5/	0.6	*	*	*	*	*	*	*	2.1	3.0	3.2	*
2005 5/	0.7	*	*	*	*	*	*	1.2	1.7	3.9	3.5	*
2004 5/	0.6	*	*	*	*	*	*	*	1.7	3.1	3.5	*
2003 5/	0.6	*	*	*	*	*	*	0.8	2.6	2.8	*	*
2002	0.7	*	*	*	*	*	*	1.1	1.8	4.1	5.0	*
2001	0.7	*	*	*	*	*	*	*	2.9	4.6	*	*
2000	0.8	*	*	*	*	*	*	1.0	2.4	4.7	6.1	*
1999	0.7	*	*	*	*	*	*	0.9	2.7	4.3	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
All origins 3/, both sexes												
2006	53.0	*	*	*	0.1	0.4	4.6	29.1	99.1	253.1	373.5	300.5
2005	53.7	*	*	*	*	0.3	5.3	29.7	103.3	259.6	375.6	302.3
2004	53.8	*	*	*	0.1	0.4	5.5	29.9	106.9	265.5	373.6	297.5
2003	54.4	*	*	*	0.1	0.4	5.6	30.3	111.0	269.3	377.8	298.9
2002	54.7	*	*	*	0.1	0.4	6.0	30.3	115.3	275.0	377.6	297.2
2001	54.8	*	*	*	0.1	0.4	6.2	30.7	117.7	279.7	371.4	302.7
2000	55.3	*	*	*	0.1	0.5	6.1	31.6	122.4	284.2	370.8	302.1
1999	54.5	*	*	*	0.1	0.4	6.0	31.3	123.4	281.5	362.1	297.0
All origins 3/, male												
2006	60.5	*	*	*	*	0.4	4.7	33.8	121.6	319.4	509.9	457.7
2005	61.8	*	*	*	*	0.4	5.5	35.1	127.6	330.7	515.1	468.0
2004	62.0	*	*	*	*	0.4	5.8	35.8	131.7	337.6	517.3	466.8
2003	62.9	*	*	*	*	0.4	6.1	36.5	136.7	346.6	525.1	475.1
2002	63.7	*	*	*	*	0.4	6.2	36.6	144.0	355.9	527.9	482.2
2001	64.7	*	*	*	*	0.4	6.7	37.4	147.1	370.0	529.5	512.4
2000	65.5	*	*	*	*	0.5	6.9	38.5	154.0	377.9	532.2	521.2
1999	65.4	*	*	*	*	0.5	6.6	38.2	158.0	377.2	527.6	529.2
All origins 3/, female												
2006	45.7	*	*	*	*	0.4	4.5	24.6	78.2	197.0	280.3	226.9
2005	45.9	*	*	*	*	0.3	5.1	24.5	80.7	199.6	280.9	226.2
2004	45.9	*	*	*	*	0.3	5.2	24.2	83.9	205.0	277.0	221.3
2003	46.1	*	*	*	*	0.4	5.1	24.4	87.1	204.8	279.4	221.0
2002	46.0	*	*	*	*	0.4	5.8	24.3	88.8	207.7	277.8	217.0
2001	45.3	*	*	*	*	0.4	5.7	24.3	90.5	204.9	267.5	215.0
2000	45.4	*	*	*	*	0.5	5.3	25.0	93.3	206.9	265.6	212.8
1999	44.1	*	*	*	*	0.4	5.5	24.6	91.5	203.1	254.5	203.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
Hispanic, both sexes												
2006	9.8	*	*	*	*	*	1.5	8.4	31.3	98.8	167.9	160.8
2005	10.5	*	*	*	*	*	1.7	9.5	35.7	102.6	183.3	177.1
2004	10.2	*	*	*	*	0.3	1.4	9.4	35.4	102.6	178.7	193.1
2003	10.4	*	*	*	*	0.3	1.6	10.2	38.1	105.8	183.8	189.2
2002	10.4	*	*	*	*	0.4	1.5	9.9	40.9	110.4	183.7	189.0
2001	10.3	*	*	*	*	*	1.9	10.7	41.0	108.3	182.3	200.7
2000	10.3	*	*	*	*	*	1.8	10.9	39.3	110.6	198.7	213.7
1999	10.4	*	*	*	*	*	2.0	10.4	44.2	112.9	187.7	227.3
Hispanic, male												
2006	12.0	*	*	*	*	*	1.5	10.5	44.8	140.1	254.2	263.9
2005	13.0	*	*	*	*	*	1.9	11.9	52.3	151.4	281.6	269.0
2004	12.7	*	*	*	*	*	1.7	12.6	50.0	151.7	279.7	287.2
2003	12.8	*	*	*	*	*	1.7	12.6	55.4	161.8	277.3	290.9
2002	13.1	*	*	*	*	*	1.5	12.6	60.7	161.7	299.1	307.9
2001	13.1	*	*	*	*	*	2.4	14.2	59.5	163.7	293.8	335.2
2000	13.3	*	*	*	*	*	2.1	14.8	58.6	167.3	327.5	368.8
1999	13.6	*	*	*	*	*	2.3	14.2	68.1	175.3	316.0	362.3
Hispanic, female												
2006	7.5	*	*	*	*	*	1.5	6.3	19.0	65.2	107.6	106.0
2005	7.8	*	*	*	*	*	1.6	7.1	20.4	63.1	114.9	129.0
2004	7.6	*	*	*	*	*	1.0	6.2	22.1	63.1	108.9	144.8
2003	7.8	*	*	*	*	*	1.4	7.9	22.4	61.2	119.6	137.5
2002	7.6	*	*	*	*	*	1.6	7.3	23.2	69.5	104.6	130.1
2001	7.4	*	*	*	*	*	1.4	7.3	24.6	64.6	107.0	135.0
2000	7.2	*	*	*	*	*	1.4	7.1	22.2	66.0	112.3	137.5
1999	7.1	*	*	*	*	*	1.7	6.7	23.1	64.0	102.3	162.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
Non-Hispanic 4/, both sexes												
2006	60.4	*	*	*	0.1	0.4	5.2	31.5	104.9	264.8	385.9	306.9
2005	60.9	*	*	*	*	0.4	5.9	31.9	108.9	271.1	386.5	307.9
2004	60.9	*	*	*	0.1	0.4	6.2	32.0	112.6	277.0	383.9	301.6
2003	61.2	*	*	*	0.1	0.4	6.2	32.3	116.6	280.3	387.2	303.0
2002	61.4	*	*	*	*	0.4	6.7	32.3	121.0	285.7	386.2	301.0
2001	61.3	*	*	*	*	0.4	6.8	32.5	123.3	290.4	379.4	305.8
2000	61.5	*	*	*	*	0.5	6.6	33.4	128.4	294.2	377.4	304.5
1999	60.5	*	*	*	*	0.5	6.5	33.0	128.9	290.7	368.2	298.8
Non-Hispanic 4/, male												
2006	69.3	*	*	*	*	0.5	5.3	36.5	128.0	332.5	525.5	467.4
2005	70.3	*	*	*	*	0.4	6.1	37.6	133.7	343.4	528.5	477.9
2004	70.4	*	*	*	*	0.4	6.5	38.2	138.1	350.1	529.8	474.8
2003	71.1	*	*	*	*	0.4	6.8	38.9	142.9	358.3	537.2	483.1
2002	71.8	*	*	*	*	0.5	6.9	38.9	150.3	368.1	538.0	488.9
2001	72.6	*	*	*	*	0.4	7.3	39.5	153.2	382.4	539.5	518.4
2000	73.2	*	*	*	*	0.6	7.5	40.5	160.6	389.7	539.9	526.2
1999	72.7	*	*	*	*	0.5	7.1	40.1	164.0	387.8	534.9	534.6
Non-Hispanic 4/, female												
2006	51.9	*	*	*	*	0.4	5.0	26.7	83.3	207.4	290.6	232.3
2005	51.9	*	*	*	*	0.3	5.7	26.4	85.8	210.0	290.3	230.3
2004	51.8	*	*	*	*	0.3	5.8	26.1	88.9	215.4	285.8	224.2
2003	51.8	*	*	*	*	0.4	5.6	26.0	92.2	214.9	287.2	223.9
2002	51.6	*	*	*	*	0.4	6.4	25.9	93.9	216.9	285.5	220.0
2001	50.5	*	*	*	*	0.5	6.2	25.8	95.6	213.9	274.3	217.4
2000	50.4	*	*	*	*	0.5	5.8	26.5	98.6	215.4	271.6	214.8
1999	48.8	*	*	*	*	0.4	6.0	26.1	96.5	211.0	260.0	204.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
Non-Hispanic white, both sexes												
2006 5/	66.9	*	*	*	*	0.4	5.2	30.4	105.1	271.5	395.0	313.0
2005 5/	67.3	*	*	*	*	0.3	5.9	31.0	108.6	277.8	394.7	311.3
2004 5/	67.1	*	*	*	*	0.3	6.1	31.0	112.5	283.2	389.6	306.3
2003 5/	67.5	*	*	*	*	0.4	6.1	31.0	117.0	286.5	392.6	306.7
2002	67.7	*	*	*	*	0.4	6.5	31.2	120.6	292.0	392.2	304.8
2001	67.3	*	*	*	*	0.5	6.6	31.5	124.0	295.2	383.0	308.6
2000	67.4	*	*	*	*	0.5	6.2	32.0	129.1	298.0	381.7	306.8
1999	65.9	*	*	*	*	0.5	6.2	31.5	128.9	293.5	370.3	300.5
Non-Hispanic white, male												
2006 5/	75.8	*	*	*	*	0.5	5.3	34.5	124.8	334.0	531.2	474.2
2005 5/	76.8	*	*	*	*	0.4	6.0	36.0	129.7	345.6	534.1	480.2
2004 5/	76.5	*	*	*	*	0.4	6.3	35.8	134.2	351.4	530.6	480.3
2003 5/	77.3	*	*	*	*	0.4	6.5	36.5	139.4	359.6	537.0	486.8
2002	77.8	*	*	*	*	0.5	6.5	36.5	145.7	369.5	538.3	493.3
2001	78.4	*	*	*	*	0.4	6.9	37.0	150.3	381.1	536.5	520.0
2000	78.9	*	*	*	*	0.6	6.7	37.7	157.7	387.3	537.7	527.3
1999	78.1	*	*	*	*	0.5	6.6	37.1	160.1	384.4	531.1	536.9
Non-Hispanic white, female												
2006 5/	58.4	*	*	*	*	0.4	5.1	26.3	86.4	217.3	300.5	237.8
2005 5/	58.2	*	*	*	*	0.3	5.9	26.0	88.5	219.3	298.7	234.1
2004 5/	58.1	*	*	*	*	0.3	5.9	26.2	92.0	224.6	293.5	228.4
2003 5/	58.1	*	*	*	*	0.4	5.7	25.6	95.7	223.9	295.1	227.4
2002	57.9	*	*	*	*	0.3	6.5	26.0	96.8	225.9	294.3	223.4
2001	56.6	*	*	*	*	0.5	6.4	26.0	99.2	222.3	281.3	220.6
2000	56.4	*	*	*	*	0.5	5.7	26.4	102.2	222.9	279.2	218.0
1999	54.3	*	*	*	*	0.5	5.8	26.1	99.5	217.5	265.0	206.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
Non-Hispanic black, both sexes												
2006 5/	43.3	*	*	*	*	0.6	6.3	46.2	131.3	277.8	370.8	268.4
2005 5/	44.0	*	*	*	*	*	7.0	45.5	140.7	283.5	373.4	294.1
2004 5/	44.2	*	*	*	*	0.5	8.0	46.5	144.2	285.3	390.1	281.5
2003 5/	44.5	*	*	*	*	*	8.0	48.5	147.4	293.9	389.4	283.5
2002	44.7	*	*	*	*	0.4	9.0	48.1	155.6	299.5	385.6	274.9
2001	44.6	*	*	*	*	*	9.2	47.5	149.4	311.4	390.9	281.6
2000	45.4	*	*	*	*	0.6	10.4	51.5	155.0	320.2	379.4	291.0
1999	45.6	*	*	*	*	0.4	9.6	51.5	161.1	317.7	390.9	293.3
Non-Hispanic black, male												
2006 5/	54.1	*	*	*	*	*	6.8	58.5	189.5	410.3	574.5	448.1
2005 5/	55.0	*	*	*	*	*	7.9	58.2	203.5	417.3	575.5	503.4
2004 5/	56.3	*	*	*	*	*	9.0	64.3	211.1	419.0	619.8	488.8
2003 5/	57.0	*	*	*	*	*	9.7	65.3	215.1	434.9	632.5	509.3
2002	58.1	*	*	*	*	*	11.0	66.3	228.0	451.4	633.9	490.1
2001	59.2	*	*	*	*	*	11.4	67.5	218.3	484.4	668.0	550.9
2000	60.3	*	*	*	*	*	13.8	72.4	227.1	495.4	649.5	567.1
1999	60.9	*	*	*	*	*	11.7	73.7	241.9	491.0	665.1	580.0
Non-Hispanic black, female												
2006 5/	33.5	*	*	*	*	*	5.9	35.7	84.5	183.1	255.3	195.1
2005 5/	34.0	*	*	*	*	*	6.3	34.6	90.1	187.9	258.8	209.9
2004 5/	33.2	*	*	*	*	*	7.0	31.3	90.5	190.0	260.5	199.6
2003 5/	33.1	*	*	*	*	*	6.5	34.1	93.1	193.5	252.1	196.0
2002	32.6	*	*	*	*	*	7.3	32.5	97.5	191.8	245.1	192.7
2001	31.3	*	*	*	*	*	7.3	30.4	94.0	189.4	235.3	181.8
2000	31.9	*	*	*	*	*	7.4	33.6	97.0	196.9	227.3	188.4
1999	31.8	*	*	*	*	*	7.9	32.5	96.4	195.5	237.4	186.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
All origins 3/, both sexes												
2006	2.8	*	*	*	0.1	0.5	1.2	2.7	5.2	9.4	15.6	20.1
2005	2.8	*	*	*	0.1	0.5	1.2	2.8	5.2	9.4	15.6	19.8
2004	2.7	*	*	*	0.1	0.5	1.2	2.8	5.3	9.3	14.7	17.7
2003	2.7	*	*	*	0.1	0.5	1.3	2.9	5.5	8.7	14.6	17.6
2002	2.6	*	*	*	0.1	0.5	1.3	2.8	5.1	8.8	13.7	17.9
2001	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.2	13.3	17.8
2000	2.6	*	*	*	0.1	0.5	1.4	3.2	5.4	8.9	13.1	18.9
1999	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.0	13.4	16.4
All origins 3/, male												
2006	3.7	*	*	*	0.1	0.5	1.4	3.5	7.4	13.7	25.6	36.7
2005	3.6	*	*	*	0.1	0.7	1.4	3.6	7.0	13.8	24.5	34.9
2004	3.5	*	*	*	*	0.6	1.4	3.7	7.6	13.7	23.7	31.4
2003	3.5	*	*	*	0.1	0.6	1.5	3.8	7.8	12.6	23.3	30.2
2002	3.3	*	*	*	0.1	0.6	1.6	3.8	7.2	13.0	21.2	29.2
2001	3.4	*	*	*	0.1	0.5	1.7	3.9	7.3	13.9	21.3	29.9
2000	3.3	*	*	*	0.1	0.5	1.6	4.2	7.4	12.9	20.8	31.8
1999	3.3	*	*	*	0.1	0.5	1.8	4.0	7.5	13.2	20.9	27.5
All origins 3/, female												
2006	2.0	*	*	*	*	0.4	0.9	1.9	3.2	5.8	8.9	12.4
2005	2.0	*	*	*	*	0.4	1.0	2.1	3.5	5.6	9.5	12.9
2004	1.9	*	*	*	0.1	0.4	1.0	2.0	3.2	5.5	8.7	11.5
2003	1.9	*	*	*	*	0.4	1.0	2.1	3.5	5.4	8.7	12.1
2002	1.9	*	*	*	0.1	0.4	1.1	1.9	3.2	5.4	8.8	12.9
2001	1.9	*	*	*	*	0.4	1.1	2.3	3.3	5.3	8.1	12.8
2000	2.0	*	*	*	0.1	0.4	1.2	2.2	3.5	5.7	8.2	13.7
1999	1.9	*	*	*	*	0.4	1.0	2.2	3.3	5.5	8.5	11.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
Hispanic, both sexes												
2006	0.5	*	*	*	*	*	*	0.8	1.5	2.6	5.5	*
2005	0.4	*	*	*	*	*	*	0.8	1.3	2.9	3.3	*
2004	0.3	*	*	*	*	*	*	0.7	1.4	1.8	3.5	*
2003	0.4	*	*	*	*	*	*	0.9	1.4	3.0	3.9	*
2002	0.4	*	*	*	*	*	0.4	0.7	1.2	*	4.8	*
2001	0.4	*	*	*	*	*	0.4	0.7	1.4	2.5	5.0	*
2000	0.4	*	*	*	*	*	0.5	0.9	1.6	2.4	*	*
1999	0.4	*	*	*	*	*	*	0.7	1.8	2.6	4.2	*
Hispanic, male												
2006	0.5	*	*	*	*	*	*	0.9	1.8	*	6.6	*
2005	0.5	*	*	*	*	*	*	0.9	2.0	4.2	*	*
2004	0.4	*	*	*	*	*	*	*	*	*	*	*
2003	0.4	*	*	*	*	*	*	*	*	3.9	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.4	*	*	*	*	*	*	*	*	*	4.6	*
2005	0.3	*	*	*	*	*	*	*	*	*	*	*
2004	0.3	*	*	*	*	*	*	*	*	*	*	*
2003	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
Non-Hispanic 4/, both sexes												
2006	3.2	*	*	*	0.1	0.5	1.3	2.9	5.5	9.9	16.3	20.8
2005	3.2	*	*	*	0.1	0.6	1.4	3.0	5.5	9.9	16.3	20.5
2004	3.1	*	*	*	0.1	0.6	1.3	3.0	5.6	9.8	15.3	18.3
2003	3.0	*	*	*	0.1	0.6	1.4	3.1	5.9	9.1	15.1	18.1
2002	2.9	*	*	*	0.1	0.6	1.5	3.0	5.4	9.3	14.1	18.2
2001	3.0	*	*	*	0.1	0.5	1.5	3.3	5.6	9.7	13.7	18.3
2000	3.0	*	*	*	0.1	0.5	1.5	3.4	5.7	9.3	13.5	19.3
1999	2.9	*	*	*	0.1	0.6	1.5	3.3	5.6	9.3	13.7	16.7
Non-Hispanic 4/, male												
2006	4.3	*	*	*	0.2	0.6	1.6	3.8	7.9	14.5	26.8	38.0
2005	4.2	*	*	*	0.1	0.8	1.6	3.9	7.4	14.5	25.6	36.2
2004	4.1	*	*	*	*	0.7	1.6	3.9	8.0	14.5	24.8	32.7
2003	4.0	*	*	*	0.1	0.7	1.7	4.0	8.2	13.2	24.2	30.9
2002	3.8	*	*	*	*	0.7	1.8	4.1	7.6	13.7	22.0	30.2
2001	3.9	*	*	*	0.1	0.6	1.9	4.2	7.8	14.6	22.0	31.0
2000	3.7	*	*	*	0.1	0.6	1.7	4.4	7.8	13.4	21.3	32.6
1999	3.7	*	*	*	0.1	0.6	2.0	4.2	7.8	13.7	21.5	28.2
Non-Hispanic 4/, female												
2006	2.2	*	*	*	*	0.4	1.0	2.1	3.4	6.0	9.1	12.8
2005	2.3	*	*	*	*	0.4	1.1	2.2	3.7	5.9	9.9	13.4
2004	2.1	*	*	*	*	0.5	1.1	2.1	3.4	5.8	9.0	11.9
2003	2.2	*	*	*	*	0.4	1.1	2.2	3.6	5.7	9.0	12.4
2002	2.1	*	*	*	*	0.4	1.2	2.0	3.3	5.7	8.9	12.9
2001	2.1	*	*	*	*	0.5	1.2	2.4	3.5	5.6	8.3	12.9
2000	2.2	*	*	*	*	0.4	1.3	2.4	3.7	5.9	8.4	13.9
1999	2.1	*	*	*	*	0.5	1.1	2.3	3.5	5.7	8.7	12.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
Non-Hispanic white, both sexes												
2006 5/	4.0	*	*	*	0.2	0.7	1.6	3.5	6.4	11.5	18.0	22.5
2005 5/	4.0	*	*	*	0.1	0.8	1.7	3.7	6.3	11.3	18.0	22.2
2004 5/	3.8	*	*	*	0.1	0.8	1.7	3.6	6.6	11.2	16.9	19.9
2003 5/	3.8	*	*	*	0.1	0.7	1.8	3.7	6.8	10.4	16.6	19.6
2002	3.6	*	*	*	0.1	0.7	1.8	3.6	6.2	10.5	15.6	19.6
2001	3.7	*	*	*	0.2	0.7	1.9	4.0	6.5	10.8	15.0	19.9
2000	3.6	*	*	*	0.2	0.7	1.8	4.0	6.5	10.4	14.7	20.8
1999	3.5	*	*	*	0.1	0.7	1.9	3.9	6.4	10.5	14.9	17.9
Non-Hispanic white, male												
2006 5/	5.3	*	*	*	0.2	0.8	2.0	4.6	9.0	16.6	29.5	41.3
2005 5/	5.2	*	*	*	*	1.0	2.0	4.6	8.5	16.6	28.3	39.7
2004 5/	5.1	*	*	*	*	0.9	2.0	4.7	9.3	16.6	27.2	35.8
2003 5/	4.9	*	*	*	0.2	0.9	2.1	4.8	9.4	14.9	26.5	33.8
2002	4.7	*	*	*	*	0.9	2.2	4.8	8.6	15.3	24.1	32.9
2001	4.8	*	*	*	0.2	0.8	2.3	5.0	9.0	16.3	24.1	34.1
2000	4.6	*	*	*	0.2	0.8	2.0	5.2	8.9	15.0	23.2	35.7
1999	4.5	*	*	*	*	0.8	2.4	5.0	8.9	15.4	23.3	30.5
Non-Hispanic white, female												
2006 5/	2.7	*	*	*	*	0.6	1.2	2.5	3.9	7.0	10.0	13.8
2005 5/	2.8	*	*	*	*	0.6	1.4	2.7	4.3	6.7	11.0	14.3
2004 5/	2.6	*	*	*	*	0.6	1.4	2.5	4.0	6.6	9.9	12.8
2003 5/	2.6	*	*	*	*	0.5	1.4	2.6	4.2	6.4	9.8	13.3
2002	2.6	*	*	*	*	0.6	1.4	2.4	3.9	6.5	9.8	13.8
2001	2.6	*	*	*	*	0.6	1.5	2.9	4.1	6.2	9.0	13.9
2000	2.7	*	*	*	*	0.6	1.6	2.8	4.2	6.6	9.1	14.7
1999	2.5	*	*	*	*	0.6	1.4	2.8	4.1	6.4	9.4	12.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
Non-Hispanic black, both sexes												
2006 5/	0.3	*	*	*	*	*	*	*	0.8	*	3.4	5.3
2005 5/	0.3	*	*	*	*	*	*	*	0.9	*	2.9	5.5
2004 5/	0.3	*	*	*	*	*	*	*	*	1.6	3.4	*
2003 5/	0.3	*	*	*	*	*	*	*	0.9	1.6	2.9	*
2002	0.3	*	*	*	*	*	*	*	0.8	1.5	*	*
2001	0.3	*	*	*	*	*	*	*	*	2.2	2.6	*
2000	0.4	*	*	*	*	*	*	*	*	1.9	3.6	6.7
1999	0.3	*	*	*	*	*	*	*	*	1.4	2.8	*
Non-Hispanic black, male												
2006 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	2.1	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
All origins 3/, both sexes												
2006	13.8	*	*	*	*	0.9	5.4	14.1	28.1	42.1	71.6	116.7
2005	14.0	*	*	*	*	0.9	5.7	14.7	28.5	43.5	71.9	122.7
2004	14.1	*	*	*	*	1.0	5.7	15.0	29.1	44.9	72.2	124.2
2003	14.4	*	*	*	*	1.0	6.1	15.6	29.6	45.5	74.9	132.5
2002	14.5	*	*	*	*	1.1	6.1	16.1	29.5	46.5	76.5	134.6
2001	14.7	*	*	*	*	1.2	6.3	16.8	30.2	47.6	76.8	134.2
2000	15.0	*	*	*	*	1.2	6.3	16.9	31.2	49.1	78.9	147.4
1999	14.9	*	*	*	*	1.1	6.1	16.9	31.1	49.5	78.9	144.4
All origins 3/, male												
2006	0.3	*	*	*	*	*	*	0.2	0.6	1.0	2.1	3.0
2005	0.3	*	*	*	*	*	*	0.2	0.5	1.3	2.1	2.4
2004	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.1	3.3
2003	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.0	3.9
2002	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.2	3.4
2001	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.3	3.5
2000	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.2
1999	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.0	4.5
All origins 3/, female												
2006	26.9	*	*	*	*	1.8	10.8	27.6	53.7	76.9	119.2	169.9
2005	27.3	*	*	*	*	1.8	11.3	28.7	54.5	79.2	119.2	177.9
2004	27.5	*	*	*	*	2.0	11.3	29.3	55.8	81.6	119.5	178.6
2003	28.2	*	*	*	*	2.1	12.2	30.4	56.6	82.6	123.7	189.4
2002	28.3	*	*	*	*	2.1	12.0	31.4	56.2	84.4	125.9	191.5
2001	28.6	*	*	*	*	2.4	12.4	32.8	57.5	85.8	125.8	188.9
2000	29.2	*	*	*	*	2.3	12.4	33.0	59.3	88.3	128.9	205.7
1999	28.9	*	*	*	*	2.2	12.0	32.9	59.2	88.9	128.9	200.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
Hispanic, both sexes												
2006	4.7	*	*	*	*	0.8	3.7	9.7	19.0	26.3	41.0	58.7
2005	4.5	*	*	*	*	0.5	3.8	9.9	18.2	26.0	43.5	62.5
2004	4.7	*	*	*	*	0.6	3.7	9.9	20.2	29.4	41.2	72.3
2003	4.7	*	*	*	*	0.7	4.3	10.5	20.5	27.1	42.0	77.7
2002	4.5	*	*	*	*	0.7	3.8	11.0	17.6	27.4	43.7	71.0
2001	4.8	*	*	*	*	0.8	4.5	12.0	22.0	28.0	39.5	68.8
2000	4.8	*	*	*	*	0.8	4.3	12.1	21.0	31.3	45.6	71.7
1999	4.6	*	*	*	*	0.7	4.0	12.2	19.6	28.0	43.5	86.3
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	9.6	*	*	*	*	1.8	7.9	19.6	36.4	47.3	68.8	89.9
2005	9.4	*	*	*	*	1.0	8.0	20.0	34.7	46.9	73.3	95.1
2004	9.6	*	*	*	*	1.4	7.7	19.5	38.2	52.1	69.1	108.8
2003	9.8	*	*	*	*	1.6	9.1	21.0	38.8	48.4	70.8	114.8
2002	9.2	*	*	*	*	1.6	7.8	21.6	33.5	48.7	73.1	105.3
2001	9.7	*	*	*	*	1.7	9.2	23.8	41.5	49.3	64.7	102.4
2000	9.7	*	*	*	*	1.8	8.7	23.9	39.1	54.9	74.9	105.8
1999	9.3	*	*	*	*	1.6	8.3	23.9	36.6	49.1	71.8	127.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
Non-Hispanic 4/, both sexes												
2006	15.3	*	*	*	*	0.9	5.7	14.6	28.8	43.2	73.5	119.4
2005	15.6	*	*	*	*	1.0	6.0	15.2	29.3	44.8	73.4	125.4
2004	15.6	*	*	*	*	1.1	6.0	15.5	29.9	45.9	73.9	126.3
2003	15.9	*	*	*	*	1.1	6.4	16.1	30.3	46.7	76.4	134.7
2002	16.0	*	*	*	*	1.1	6.4	16.6	30.3	47.7	78.0	136.8
2001	16.1	*	*	*	*	1.2	6.5	17.2	30.7	48.8	78.4	136.4
2000	16.5	*	*	*	*	1.2	6.5	17.3	31.9	50.1	80.1	149.7
1999	16.3	*	*	*	*	1.2	6.3	17.3	31.9	50.6	80.2	146.2
Non-Hispanic 4/, male												
2006	0.3	*	*	*	*	*	*	0.2	0.6	1.0	2.1	3.2
2005	0.3	*	*	*	*	*	*	0.2	0.5	1.4	2.2	2.6
2004	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.2	3.3
2003	0.3	*	*	*	*	*	*	0.3	0.6	1.1	2.1	3.8
2002	0.3	*	*	*	*	*	*	0.2	0.6	1.0	2.3	3.5
2001	0.3	*	*	*	*	*	*	0.3	0.6	1.5	2.3	3.7
2000	0.3	*	*	*	*	*	*	0.3	0.7	1.4	2.2	4.2
1999	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.1	4.7
Non-Hispanic 4/, female												
2006	29.7	*	*	*	*	1.9	11.2	28.5	55.1	79.1	122.2	173.5
2005	30.1	*	*	*	*	2.0	11.9	29.6	56.1	81.5	121.7	181.5
2004	30.2	*	*	*	*	2.1	11.9	30.3	57.2	83.7	122.1	181.3
2003	30.8	*	*	*	*	2.1	12.6	31.3	57.9	84.9	126.0	192.2
2002	31.0	*	*	*	*	2.2	12.6	32.3	57.9	86.7	128.2	194.4
2001	31.1	*	*	*	*	2.5	12.8	33.6	58.6	88.1	128.3	191.6
2000	31.8	*	*	*	*	2.4	12.8	33.8	60.7	90.3	130.8	208.6
1999	31.4	*	*	*	*	2.3	12.4	33.6	60.7	91.2	130.9	202.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
Non-Hispanic white, both sexes												
2006 5/	16.2	*	*	*	*	0.8	5.0	13.3	27.6	43.1	73.8	119.8
2005 5/	16.3	*	*	*	*	0.8	5.3	13.6	27.9	44.5	74.0	126.3
2004 5/	16.4	*	*	*	*	0.9	5.2	14.0	28.7	45.8	74.7	126.8
2003 5/	16.8	*	*	*	*	0.9	5.7	14.8	29.0	46.3	76.7	134.7
2002	16.9	*	*	*	*	0.9	5.6	15.2	29.4	47.4	78.0	137.7
2001	17.0	*	*	*	*	1.0	5.8	16.0	29.6	48.2	78.7	137.1
2000	17.4	*	*	*	*	1.1	5.8	16.1	30.8	50.3	80.6	149.6
1999	17.1	*	*	*	*	1.0	5.6	16.0	30.6	50.2	80.3	147.8
Non-Hispanic white, male												
2006 5/	0.3	*	*	*	*	*	*	0.2	0.6	1.0	2.1	3.3
2005 5/	0.3	*	*	*	*	*	*	0.2	0.5	1.3	2.1	2.6
2004 5/	0.3	*	*	*	*	*	*	0.1	0.5	1.0	2.2	2.9
2003 5/	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.2	3.9
2002	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.2	3.4
2001	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.2	3.5
2000	0.4	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.0
1999	0.3	*	*	*	*	*	*	0.2	0.6	1.1	2.0	4.5
Non-Hispanic white, female												
2006 5/	31.4	*	*	*	*	1.5	10.0	26.3	53.4	79.6	123.6	174.1
2005 5/	31.8	*	*	*	*	1.7	10.6	26.8	53.9	81.9	123.4	182.9
2004 5/	32.0	*	*	*	*	1.8	10.4	27.7	55.6	84.4	124.2	182.4
2003 5/	32.6	*	*	*	*	1.8	11.3	29.1	56.0	85.1	127.1	192.3
2002	32.9	*	*	*	*	1.9	11.1	30.0	56.7	87.0	128.9	195.8
2001	32.9	*	*	*	*	2.0	11.5	31.6	57.0	88.0	129.4	192.6
2000	33.8	*	*	*	*	2.1	11.6	31.7	59.2	91.4	132.2	208.3
1999	33.3	*	*	*	*	2.0	11.1	31.5	58.9	91.2	131.5	204.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
Non-Hispanic black, both sexes												
2006 5/	15.1	*	*	*	*	2.0	10.5	24.9	43.3	54.6	90.8	144.7
2005 5/	15.5	*	*	*	*	1.9	11.2	27.8	46.3	58.1	83.2	146.1
2004 5/	15.0	*	*	*	*	2.2	11.3	26.7	44.7	57.4	83.1	145.6
2003 5/	15.5	*	*	*	*	2.3	11.4	27.1	46.1	60.1	91.3	163.3
2002	15.4	*	*	*	*	2.4	12.0	27.7	43.7	61.1	95.3	154.0
2001	15.5	*	*	*	*	2.7	11.5	27.9	45.6	64.6	93.0	153.9
2000	15.3	*	*	*	*	2.3	11.5	28.4	46.4	59.4	91.7	178.7
1999	15.5	*	*	*	*	2.4	12.0	29.2	47.1	64.5	95.6	149.1
Non-Hispanic black, male												
2006 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.3	*	*	*	*	*	*	*	*	2.9	*	*
2004 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	28.5	*	*	*	*	3.9	19.8	45.6	77.3	92.3	140.7	202.5
2005 5/	29.4	*	*	*	*	3.7	21.0	51.4	82.8	97.6	128.2	204.1
2004 5/	28.4	*	*	*	*	4.2	21.2	49.3	79.8	96.8	128.5	199.6
2003 5/	29.4	*	*	*	*	4.3	21.4	50.1	82.4	101.6	141.8	224.5
2002	29.1	*	*	*	*	4.5	22.5	51.0	77.8	102.7	147.1	210.7
2001	29.2	*	*	*	*	5.1	21.5	51.3	81.6	108.4	143.0	208.3
2000	28.9	*	*	*	*	4.3	21.5	52.5	82.5	100.1	141.5	241.6
1999	29.2	*	*	*	*	4.5	22.3	53.7	83.7	108.2	147.2	201.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
All origins 3/, both sexes												
2006	1.3	*	*	*	*	0.4	1.4	2.2	2.6	3.2	3.8	5.2
2005	1.3	*	*	*	*	0.4	1.4	2.3	2.6	3.1	4.0	5.4
2004	1.3	*	*	*	*	0.5	1.4	2.0	2.8	3.3	3.7	5.1
2003	1.3	*	*	*	*	0.5	1.4	2.2	2.8	3.1	4.3	5.7
2002	1.4	*	*	*	*	0.5	1.5	2.2	2.7	3.4	4.1	6.2
2001	1.4	*	*	*	*	0.5	1.6	2.4	2.9	3.2	4.6	7.2
2000	1.5	*	*	*	*	0.6	1.5	2.4	3.1	3.7	4.8	7.3
1999	1.5	*	*	*	*	0.5	1.6	2.5	3.2	3.8	4.6	7.9
All origins 3/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2006	2.6	*	*	*	*	0.9	2.9	4.4	5.0	5.9	6.4	7.7
2005	2.6	*	*	*	*	0.9	2.8	4.4	5.1	5.7	6.7	7.8
2004	2.6	*	*	*	*	1.1	2.8	4.0	5.4	6.0	6.2	7.3
2003	2.7	*	*	*	*	1.0	2.8	4.3	5.5	5.6	7.2	8.3
2002	2.7	*	*	*	*	1.1	3.0	4.4	5.2	6.3	6.9	8.9
2001	2.8	*	*	*	*	1.1	3.1	4.8	5.5	5.8	7.7	10.1
2000	2.9	*	*	*	*	1.2	3.1	4.8	5.9	6.8	8.0	10.3
1999	3.0	*	*	*	*	1.1	3.1	4.9	6.2	6.9	7.7	11.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
Hispanic, both sexes												
2006	1.0	*	*	*	*	0.4	1.4	2.8	3.3	4.4	6.6	*
2005	1.0	*	*	*	*	0.3	1.6	2.8	3.8	4.8	4.5	*
2004	1.0	*	*	*	*	0.4	1.6	2.4	4.3	4.3	5.7	*
2003	1.0	*	*	*	*	0.4	1.4	2.9	3.1	3.5	6.6	*
2002	1.0	*	*	*	*	0.4	1.5	2.4	3.6	4.7	6.7	*
2001	1.1	*	*	*	*	0.5	1.9	3.0	4.1	3.4	7.0	*
2000	1.2	*	*	*	*	0.5	2.1	2.8	4.6	5.6	8.5	13.9
1999	1.1	*	*	*	*	0.5	1.8	3.3	3.9	4.9	6.5	*
Hispanic, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2006	2.2	*	*	*	*	1.0	2.9	5.7	6.3	7.9	11.3	*
2005	2.2	*	*	*	*	0.6	3.4	5.7	7.2	8.8	7.7	*
2004	2.2	*	*	*	*	0.9	3.3	4.8	8.2	7.7	9.6	*
2003	2.0	*	*	*	*	0.9	3.0	5.7	5.9	6.3	11.1	*
2002	2.1	*	*	*	*	0.9	3.1	4.7	6.8	8.4	11.2	*
2001	2.3	*	*	*	*	1.1	3.9	6.0	7.7	6.1	11.8	*
2000	2.5	*	*	*	*	1.1	4.4	5.5	8.6	10.0	14.2	20.8
1999	2.3	*	*	*	*	1.1	3.6	6.6	7.4	8.8	10.8	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
Non-Hispanic 4/, both sexes												
2006	1.4	*	*	*	*	0.5	1.4	2.2	2.5	3.1	3.6	5.2
2005	1.4	*	*	*	*	0.5	1.4	2.2	2.5	2.9	3.9	5.3
2004	1.4	*	*	*	*	0.6	1.4	2.0	2.7	3.2	3.6	5.0
2003	1.4	*	*	*	*	0.5	1.4	2.1	2.8	3.0	4.2	5.7
2002	1.4	*	*	*	*	0.5	1.5	2.2	2.6	3.4	4.0	6.1
2001	1.5	*	*	*	*	0.5	1.5	2.4	2.8	3.2	4.5	7.0
2000	1.5	*	*	*	*	0.6	1.5	2.4	3.0	3.6	4.7	7.1
1999	1.6	*	*	*	*	0.5	1.5	2.4	3.2	3.7	4.6	7.9
Non-Hispanic 4/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, female												
2006	2.7	*	*	*	*	0.9	2.9	4.2	4.9	5.7	6.1	7.6
2005	2.7	*	*	*	*	0.9	2.7	4.3	4.9	5.4	6.6	7.7
2004	2.6	*	*	*	*	1.1	2.7	3.9	5.2	5.9	6.0	7.2
2003	2.7	*	*	*	*	1.0	2.8	4.1	5.4	5.6	7.0	8.2
2002	2.8	*	*	*	*	1.1	2.9	4.3	5.1	6.2	6.7	8.7
2001	2.9	*	*	*	*	1.0	3.0	4.6	5.3	5.8	7.5	10.0
2000	3.0	*	*	*	*	1.2	2.9	4.7	5.7	6.6	7.7	9.9
1999	3.0	*	*	*	*	1.1	3.0	4.7	6.1	6.7	7.5	11.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
Non-Hispanic white, both sexes												
2006 5/	1.3	*	*	*	*	0.4	1.3	1.9	2.4	2.6	3.1	4.1
2005 5/	1.3	*	*	*	*	0.5	1.3	1.9	2.1	2.6	3.4	4.7
2004 5/	1.2	*	*	*	*	0.5	1.2	1.8	2.4	2.6	3.1	3.9
2003 5/	1.3	*	*	*	*	0.4	1.3	1.9	2.5	2.5	3.4	4.8
2002	1.3	*	*	*	*	0.5	1.3	1.9	2.3	2.7	3.4	4.9
2001	1.4	*	*	*	*	0.5	1.4	2.1	2.4	2.6	3.8	6.1
2000	1.4	*	*	*	*	0.5	1.3	2.1	2.5	2.9	4.1	6.0
1999	1.4	*	*	*	*	0.6	1.4	2.3	2.7	3.0	3.9	6.7
Non-Hispanic white, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2006 5/	2.5	*	*	*	*	0.8	2.6	3.8	4.6	4.8	5.2	6.1
2005 5/	2.5	*	*	*	*	0.9	2.5	3.8	4.2	4.8	5.7	6.8
2004 5/	2.4	*	*	*	*	1.1	2.5	3.5	4.6	4.9	5.2	5.7
2003 5/	2.5	*	*	*	*	0.9	2.6	3.7	4.9	4.7	5.7	7.0
2002	2.5	*	*	*	*	1.0	2.7	3.8	4.5	5.0	5.7	7.0
2001	2.7	*	*	*	*	1.0	2.8	4.2	4.7	4.9	6.3	8.6
2000	2.7	*	*	*	*	1.1	2.5	4.2	4.8	5.3	6.8	8.4
1999	2.8	*	*	*	*	1.1	2.8	4.5	5.2	5.5	6.5	9.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
Non-Hispanic black, both sexes												
2006 5/	2.1	*	*	*	*	0.8	2.5	3.7	4.1	7.7	8.8	17.5
2005 5/	2.1	*	*	*	*	0.5	2.4	3.8	5.3	6.5	10.7	13.4
2004 5/	2.1	*	*	*	*	0.8	2.3	3.8	5.4	7.7	9.1	16.4
2003 5/	2.2	*	*	*	*	0.9	2.3	3.7	5.7	7.5	11.6	14.9
2002	2.3	*	*	*	*	0.8	2.6	4.2	5.1	9.2	10.7	17.5
2001	2.2	*	*	*	*	0.8	2.5	3.8	5.3	7.9	12.0	19.0
2000	2.5	*	*	*	*	1.0	2.6	4.3	6.4	9.8	12.0	19.7
1999	2.4	*	*	*	*	0.7	2.5	3.6	7.2	10.8	11.3	21.7
Non-Hispanic black, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2006 5/	3.9	*	*	*	*	1.5	4.7	6.8	7.3	13.2	13.7	24.7
2005 5/	4.0	*	*	*	*	1.1	4.6	7.1	9.5	11.2	16.8	18.8
2004 5/	4.0	*	*	*	*	1.6	4.3	7.1	9.8	13.2	14.2	22.9
2003 5/	4.1	*	*	*	*	1.8	4.3	6.9	10.2	12.8	18.1	20.6
2002	4.3	*	*	*	*	1.6	4.9	7.8	9.2	15.8	16.7	24.2
2001	4.2	*	*	*	*	1.6	4.6	7.1	9.6	13.5	18.7	26.1
2000	4.7	*	*	*	*	1.9	5.0	8.0	11.5	16.6	18.8	27.0
1999	4.5	*	*	*	*	1.2	4.8	6.6	13.0	18.3	17.7	29.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
All origins 3/, both sexes												
2006	2.5	*	*	*	*	0.1	0.3	1.4	4.8	10.0	15.7	21.6
2005	2.4	*	*	*	*	0.1	0.3	1.3	4.8	9.7	15.0	22.1
2004	2.4	*	*	*	*	0.0	0.4	1.3	4.4	10.1	15.5	22.3
2003	2.4	*	*	*	*	0.1	0.3	1.3	4.7	10.1	15.2	22.8
2002	2.4	*	*	*	*	0.1	0.3	1.4	4.5	10.0	15.4	24.6
2001	2.4	*	*	*	*	0.1	0.3	1.4	4.4	9.7	16.4	24.7
2000	2.3	*	*	*	*	0.1	0.3	1.3	4.6	9.7	15.8	24.7
1999	2.3	*	*	*	*	0.1	0.3	1.3	4.3	9.8	16.4	24.0
All origins 3/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2006	4.9	*	*	*	*	0.1	0.7	2.7	9.3	18.5	26.4	31.6
2005	4.7	*	*	*	*	0.1	0.7	2.6	9.3	17.9	25.1	32.2
2004	4.7	*	*	*	*	0.1	0.8	2.6	8.6	18.5	25.9	32.3
2003	4.7	*	*	*	*	0.1	0.7	2.6	9.0	18.5	25.3	33.0
2002	4.7	*	*	*	*	0.1	0.6	2.8	8.7	18.4	25.6	35.2
2001	4.7	*	*	*	*	0.2	0.6	2.8	8.4	17.8	27.2	35.0
2000	4.6	*	*	*	*	0.1	0.6	2.6	8.9	17.7	26.2	34.8
1999	4.5	*	*	*	*	0.1	0.6	2.5	8.2	17.8	27.1	33.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
Hispanic, both sexes												
2006	0.9	*	*	*	*	*	0.3	1.2	3.5	6.5	11.6	15.3
2005	0.9	*	*	*	*	*	0.4	1.5	3.4	7.8	11.9	11.1
2004	0.8	*	*	*	*	*	*	0.9	3.5	7.9	10.5	9.5
2003	0.8	*	*	*	*	*	0.4	1.2	2.9	7.0	9.8	20.7
2002	0.9	*	*	*	*	*	0.3	1.1	3.2	9.0	11.0	20.5
2001	0.9	*	*	*	*	*	*	1.4	3.8	7.5	11.7	18.8
2000	0.8	*	*	*	*	*	*	1.1	3.6	7.0	10.7	19.9
1999	0.8	*	*	*	*	*	*	1.1	3.7	8.3	10.6	15.4
Hispanic, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2006	1.8	*	*	*	*	*	0.7	2.4	6.7	11.9	19.7	23.4
2005	1.9	*	*	*	*	*	0.8	3.1	6.5	14.1	20.2	16.9
2004	1.6	*	*	*	*	*	*	1.7	6.7	14.2	17.8	14.4
2003	1.7	*	*	*	*	*	0.9	2.3	5.6	12.5	16.6	31.3
2002	1.8	*	*	*	*	*	0.7	2.1	6.2	16.1	18.5	30.7
2001	1.8	*	*	*	*	*	*	2.8	7.3	13.4	19.6	27.9
2000	1.6	*	*	*	*	*	*	2.2	6.8	12.4	17.8	29.7
1999	1.6	*	*	*	*	*	*	2.2	7.0	14.8	17.7	22.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
Non-Hispanic 4/, both sexes												
2006	2.7	*	*	*	*	0.1	0.3	1.4	4.9	10.3	15.9	21.9
2005	2.6	*	*	*	*	0.1	0.3	1.3	4.9	9.8	15.1	22.6
2004	2.6	*	*	*	*	*	0.4	1.4	4.5	10.2	15.8	22.8
2003	2.6	*	*	*	*	0.1	0.3	1.3	4.8	10.3	15.4	22.9
2002	2.6	*	*	*	*	*	0.3	1.4	4.6	10.1	15.5	24.6
2001	2.6	*	*	*	*	0.1	0.3	1.4	4.4	9.9	16.6	24.8
2000	2.6	*	*	*	*	0.1	0.3	1.4	4.7	9.8	16.0	24.9
1999	2.5	*	*	*	*	0.1	0.3	1.3	4.3	9.9	16.6	24.3
Non-Hispanic 4/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, female												
2006	5.4	*	*	*	*	0.1	0.7	2.8	9.5	19.0	26.8	32.0
2005	5.2	*	*	*	*	0.1	0.6	2.6	9.5	18.2	25.4	32.9
2004	5.2	*	*	*	*	*	0.8	2.7	8.7	18.8	26.4	33.0
2003	5.1	*	*	*	*	0.1	0.7	2.6	9.2	18.9	25.6	33.0
2002	5.1	*	*	*	*	*	0.6	2.8	8.8	18.5	25.8	35.3
2001	5.1	*	*	*	*	0.2	0.6	2.8	8.5	18.1	27.5	35.1
2000	5.0	*	*	*	*	0.1	0.6	2.7	9.0	17.9	26.5	34.9
1999	4.9	*	*	*	*	0.1	0.6	2.5	8.3	18.0	27.4	34.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
Non-Hispanic white, both sexes												
2006 5/	2.8	*	*	*	*	*	0.3	1.3	4.4	9.0	15.1	22.0
2005 5/	2.7	*	*	*	*	*	0.3	1.2	4.4	8.9	14.3	22.2
2004 5/	2.7	*	*	*	*	*	0.4	1.3	4.1	8.9	15.2	23.0
2003 5/	2.7	*	*	*	*	*	0.3	1.3	4.4	9.3	14.5	22.8
2002	2.6	*	*	*	*	*	0.3	1.3	4.1	8.9	14.6	23.8
2001	2.7	*	*	*	*	*	0.3	1.3	4.0	8.9	15.9	24.5
2000	2.6	*	*	*	*	*	0.3	1.3	4.3	8.6	15.0	24.5
1999	2.6	*	*	*	*	*	0.3	1.3	4.0	8.9	16.0	23.6
Non-Hispanic white, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2006 5/	5.5	*	*	*	*	*	0.7	2.6	8.6	16.8	25.5	32.2
2005 5/	5.3	*	*	*	*	*	0.6	2.5	8.6	16.5	24.1	32.4
2004 5/	5.3	*	*	*	*	*	0.8	2.6	8.0	16.6	25.5	33.3
2003 5/	5.2	*	*	*	*	*	0.6	2.5	8.5	17.2	24.4	32.8
2002	5.1	*	*	*	*	*	0.6	2.7	8.0	16.5	24.4	34.1
2001	5.2	*	*	*	*	*	0.6	2.7	7.8	16.4	26.5	34.7
2000	5.1	*	*	*	*	*	0.5	2.7	8.4	15.9	24.8	34.3
1999	5.1	*	*	*	*	*	0.6	2.5	7.7	16.3	26.4	33.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
Non-Hispanic black, both sexes												
2006 5/	3.2	*	*	*	*	*	*	1.8	10.0	23.2	29.2	26.3
2005 5/	3.1	*	*	*	*	*	0.4	1.9	10.0	20.6	27.5	32.0
2004 5/	3.0	*	*	*	*	*	0.5	2.0	8.1	23.4	26.6	26.5
2003 5/	3.0	*	*	*	*	*	0.5	1.7	9.1	21.4	28.0	30.3
2002	3.1	*	*	*	*	*	0.4	2.0	8.6	22.7	28.4	39.6
2001	2.9	*	*	*	*	*	0.4	2.1	7.7	21.6	28.6	32.8
2000	3.0	*	*	*	*	*	0.4	1.7	8.4	22.9	31.1	33.7
1999	2.8	*	*	*	*	*	0.5	1.4	8.1	21.5	27.5	35.9
Non-Hispanic black, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2006 5/	6.2	*	*	*	*	*	*	3.3	18.0	39.7	45.8	37.0
2005 5/	6.0	*	*	*	*	*	0.8	3.5	18.1	35.3	43.1	44.8
2004 5/	5.7	*	*	*	*	*	1.0	3.6	14.7	40.1	41.5	36.9
2003 5/	5.7	*	*	*	*	*	1.0	3.2	16.5	36.7	43.8	42.1
2002	5.8	*	*	*	*	*	0.7	3.6	15.5	38.7	44.4	54.8
2001	5.5	*	*	*	*	*	0.7	4.0	13.9	36.8	44.6	44.9
2000	5.7	*	*	*	*	*	0.7	3.1	15.3	39.1	48.6	46.2
1999	5.4	*	*	*	*	*	0.9	2.7	14.7	36.7	42.9	49.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
All origins 3/, both sexes												
2006	5.0	*	*	*	0.1	0.2	0.9	3.8	9.6	18.9	31.1	37.8
2005	5.0	*	*	*	*	0.2	0.9	4.0	10.0	19.1	30.9	38.1
2004	5.0	*	*	*	0.1	0.2	1.0	4.0	9.6	20.0	32.1	37.7
2003	5.0	*	*	*	0.1	0.2	1.0	4.0	10.4	19.5	32.6	38.4
2002	5.1	*	*	*	0.1	0.2	1.0	4.3	10.3	19.7	33.4	38.1
2001	5.1	*	*	*	0.1	0.3	1.0	4.1	10.3	20.0	33.2	39.6
2000	5.0	*	*	*	0.1	0.3	1.0	4.2	10.1	19.6	33.2	40.9
1999	4.9	*	*	*	0.1	0.2	1.1	4.2	10.2	19.6	31.7	37.6
All origins 3/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2006	9.8	*	*	*	0.1	0.4	1.8	7.6	18.6	34.9	52.4	55.5
2005	9.8	*	*	*	*	0.4	1.9	7.8	19.4	35.3	51.8	55.6
2004	9.9	*	*	*	0.1	0.5	2.1	7.8	18.5	36.8	53.7	54.6
2003	9.9	*	*	*	0.1	0.4	2.0	7.8	20.0	35.7	54.4	55.4
2002	10.0	*	*	*	0.2	0.5	2.1	8.4	19.9	36.1	55.6	54.6
2001	9.9	*	*	*	0.1	0.5	2.1	8.1	19.9	36.6	55.0	56.1
2000	9.8	*	*	*	0.1	0.5	2.1	8.3	19.4	35.7	54.8	57.6
1999	9.6	*	*	*	0.1	0.4	2.2	8.3	19.6	35.7	52.3	52.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
Hispanic, both sexes												
2006	1.7	*	*	*	*	*	0.7	2.6	6.1	12.8	24.0	27.8
2005	1.7	*	*	*	*	*	0.6	3.1	7.1	13.1	19.9	27.4
2004	1.7	*	*	*	*	*	0.8	2.8	6.5	12.9	20.9	28.1
2003	1.6	*	*	*	*	*	0.8	2.7	6.4	12.7	22.0	28.5
2002	1.6	*	*	*	*	*	0.8	3.2	5.6	12.3	24.2	23.8
2001	1.6	*	*	*	*	*	0.8	2.7	7.1	14.4	20.0	30.6
2000	1.6	*	*	*	*	*	0.6	3.0	6.5	14.3	24.9	32.5
1999	1.4	*	*	*	*	*	0.6	2.7	6.1	12.2	20.8	23.9
Hispanic, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2006	3.5	*	*	*	*	*	1.4	5.2	11.6	23.2	40.7	42.5
2005	3.6	*	*	*	*	*	1.4	6.2	13.6	23.7	33.7	41.7
2004	3.4	*	*	*	*	*	1.7	5.5	12.5	23.3	35.3	42.5
2003	3.3	*	*	*	*	*	1.7	5.5	12.2	22.8	37.1	43.0
2002	3.3	*	*	*	*	*	1.7	6.3	10.7	22.1	40.8	35.6
2001	3.4	*	*	*	*	*	1.6	5.3	13.4	25.8	33.5	45.6
2000	3.4	*	*	*	*	*	1.2	5.9	12.2	25.5	41.6	48.5
1999	2.9	*	*	*	*	*	1.2	5.3	11.5	21.7	34.7	35.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
Non-Hispanic 4/, both sexes												
2006	5.5	*	*	*	*	0.2	1.0	4.0	10.0	19.3	31.5	38.3
2005	5.5	*	*	*	*	0.2	1.0	4.1	10.3	19.6	31.5	38.6
2004	5.6	*	*	*	0.1	0.3	1.1	4.1	9.8	20.5	32.7	38.1
2003	5.6	*	*	*	0.1	0.2	1.0	4.1	10.7	19.9	33.1	38.8
2002	5.6	*	*	*	0.1	0.3	1.1	4.4	10.7	20.2	33.8	38.7
2001	5.6	*	*	*	0.1	0.3	1.1	4.3	10.5	20.3	33.7	39.8
2000	5.5	*	*	*	*	0.3	1.1	4.3	10.3	19.9	33.4	41.2
1999	5.3	*	*	*	0.1	0.2	1.2	4.4	10.5	20.0	32.0	38.1
Non-Hispanic 4/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, female												
2006	10.8	*	*	*	*	0.5	1.9	7.8	19.2	35.7	53.1	56.0
2005	10.8	*	*	*	*	0.4	2.0	8.0	19.9	36.2	52.9	56.2
2004	10.8	*	*	*	0.1	0.5	2.1	8.0	19.0	37.7	54.7	55.1
2003	10.9	*	*	*	0.1	0.5	2.1	8.1	20.5	36.6	55.2	55.9
2002	11.0	*	*	*	0.2	0.5	2.1	8.6	20.6	37.0	56.3	55.3
2001	10.9	*	*	*	0.1	0.5	2.1	8.4	20.3	37.3	55.8	56.4
2000	10.7	*	*	*	*	0.5	2.2	8.4	19.9	36.3	55.2	57.8
1999	10.4	*	*	*	0.2	0.4	2.3	8.6	20.1	36.4	52.7	53.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
Non-Hispanic white, both sexes												
2006 5/	6.2	*	*	*	*	0.3	1.0	4.1	10.4	20.3	32.5	39.3
2005 5/	6.2	*	*	*	*	0.2	1.0	4.2	10.7	20.1	32.4	40.0
2004 5/	6.2	*	*	*	*	0.2	1.1	4.2	10.2	21.0	33.9	39.1
2003 5/	6.3	*	*	*	*	0.3	1.1	4.2	11.1	20.5	34.2	39.5
2002	6.3	*	*	*	0.1	0.3	1.1	4.6	11.1	20.9	34.8	39.6
2001	6.2	*	*	*	*	0.3	1.1	4.5	10.8	21.0	34.8	40.6
2000	6.2	*	*	*	*	0.3	1.1	4.5	10.8	20.5	34.5	42.2
1999	6.0	*	*	*	0.1	0.2	1.2	4.5	10.9	20.5	32.7	39.2
Non-Hispanic white, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2006 5/	12.3	*	*	*	*	0.5	2.0	8.2	20.3	37.9	55.1	57.6
2005 5/	12.2	*	*	*	*	0.5	2.0	8.4	20.9	37.5	54.7	58.2
2004 5/	12.2	*	*	*	*	0.5	2.2	8.3	19.8	39.1	57.0	56.5
2003 5/	12.3	*	*	*	*	0.5	2.2	8.4	21.6	38.1	57.2	56.8
2002	12.4	*	*	*	0.2	0.5	2.2	9.1	21.7	38.7	58.0	56.7
2001	12.2	*	*	*	*	0.5	2.2	9.0	21.0	38.8	57.8	57.5
2000	12.0	*	*	*	*	0.6	2.3	8.9	21.0	37.7	57.2	59.2
1999	11.7	*	*	*	0.2	0.4	2.4	9.0	21.2	37.7	54.1	54.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
Non-Hispanic black, both sexes												
2006 5/	3.1	*	*	*	*	*	0.9	3.5	8.6	16.1	25.6	33.7
2005 5/	3.2	*	*	*	*	*	1.0	3.2	8.8	18.9	28.2	31.1
2004 5/	3.4	*	*	*	*	*	0.9	4.0	8.8	20.5	28.0	32.5
2003 5/	3.2	*	*	*	*	*	1.0	3.9	8.8	17.8	27.8	37.2
2002	3.2	*	*	*	*	*	1.1	3.5	9.2	17.4	29.3	35.4
2001	3.2	*	*	*	*	*	1.2	3.1	10.2	18.5	27.2	36.5
2000	3.0	*	*	*	*	*	1.1	3.4	8.6	18.2	27.1	34.7
1999	3.1	*	*	*	*	*	0.9	3.6	9.1	18.7	28.7	29.1
Non-Hispanic black, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2006 5/	5.9	*	*	*	*	*	1.7	6.6	15.5	27.6	40.1	47.5
2005 5/	6.2	*	*	*	*	*	1.9	6.0	15.9	32.4	44.2	43.7
2004 5/	6.5	*	*	*	*	*	1.8	7.5	15.9	35.2	43.8	45.4
2003 5/	6.2	*	*	*	*	*	1.9	7.2	15.9	30.5	43.5	51.6
2002	6.1	*	*	*	*	*	2.0	6.5	16.6	29.8	45.9	48.9
2001	6.1	*	*	*	*	*	2.2	5.7	18.4	31.5	42.5	50.0
2000	5.8	*	*	*	*	*	2.0	6.3	15.6	31.1	42.4	47.5
1999	5.8	*	*	*	*	*	1.6	6.7	16.4	31.9	44.8	40.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)												
All origins 3/, both sexes												
2006	9.5	*	*	*	*	*	0.1	0.9	7.2	29.6	85.5	167.8
2005	9.8	*	*	*	*	*	0.1	0.9	7.1	30.9	89.4	174.6
2004	9.9	*	*	*	*	*	0.1	1.0	7.0	30.9	92.5	180.9
2003	10.2	*	*	*	*	*	0.1	1.0	7.4	32.9	95.5	184.9
2002	10.6	*	*	*	*	*	0.1	1.0	7.6	34.5	100.1	195.0
2001	10.8	*	*	*	*	*	0.1	1.1	7.5	35.7	103.2	200.9
2000	11.0	*	*	*	*	*	0.1	1.0	7.9	37.0	105.7	209.3
1999	11.4	*	*	*	*	*	0.1	1.0	8.2	39.7	109.9	207.5
All origins 3/, male												
2006	19.2	*	*	*	*	*	0.2	1.9	14.9	64.6	210.6	526.4
2005	19.8	*	*	*	*	*	0.1	1.9	14.7	67.6	221.0	554.7
2004	20.1	*	*	*	*	*	0.1	2.1	14.6	67.8	229.8	583.1
2003	20.7	*	*	*	*	*	0.1	2.1	15.4	72.3	238.3	603.3
2002	21.5	*	*	*	*	*	0.1	1.9	15.8	75.9	251.0	644.5
2001	22.0	*	*	*	*	*	0.1	2.1	15.7	78.9	260.2	681.2
2000	22.5	*	*	*	*	*	0.1	2.0	16.6	82.0	267.9	723.1
1999	23.2	*	*	*	*	*	0.1	2.1	17.1	88.2	279.1	722.4
All origins 3/, female												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)												
Hispanic, both sexes												
2006	3.1	*	*	*	*	*	*	0.7	5.7	24.6	67.4	132.2
2005	3.0	*	*	*	*	*	*	0.6	5.3	24.1	67.7	142.5
2004	3.0	*	*	*	*	*	*	0.6	4.9	22.4	79.8	130.3
2003	3.1	*	*	*	*	*	*	0.7	5.3	25.5	74.5	151.8
2002	3.1	*	*	*	*	*	*	0.8	5.4	24.7	78.7	170.7
2001	3.3	*	*	*	*	*	*	0.8	5.8	29.2	87.8	170.1
2000	2.9	*	*	*	*	*	*	0.8	4.6	25.7	79.2	165.9
1999	3.1	*	*	*	*	*	*	0.7	5.6	28.5	80.4	176.1
Hispanic, male												
2006	6.0	*	*	*	*	*	*	1.4	11.8	55.0	163.8	381.2
2005	5.9	*	*	*	*	*	*	1.2	11.0	54.0	165.0	414.7
2004	5.8	*	*	*	*	*	*	1.2	10.3	50.2	195.3	383.9
2003	5.9	*	*	*	*	*	*	1.3	11.2	57.6	182.9	451.0
2002	6.0	*	*	*	*	*	*	1.6	11.3	55.8	193.7	515.4
2001	6.4	*	*	*	*	*	*	1.6	12.4	66.3	217.8	518.0
2000	5.7	*	*	*	*	*	*	1.7	9.7	58.5	197.2	503.9
1999	6.0	*	*	*	*	*	*	1.5	12.0	64.8	201.3	541.3
Hispanic, female												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)												
Non-Hispanic 4/, both sexes												
2006	10.6	*	*	*	*	*	0.1	1.0	7.3	29.9	86.5	169.3
2005	10.9	*	*	*	*	*	0.1	1.0	7.2	31.4	90.5	176.0
2004	11.0	*	*	*	*	*	0.1	1.1	7.2	31.5	93.0	183.0
2003	11.3	*	*	*	*	*	0.1	1.1	7.6	33.3	96.4	186.0
2002	11.7	*	*	*	*	*	0.1	1.0	7.7	35.0	101.0	195.5
2001	11.9	*	*	*	*	*	0.1	1.1	7.6	36.1	103.6	201.6
2000	12.2	*	*	*	*	*	0.1	1.0	8.2	37.6	106.6	210.4
1999	12.5	*	*	*	*	*	0.1	1.0	8.3	40.3	110.9	208.2
Non-Hispanic 4/, male												
2006	21.6	*	*	*	*	*	0.2	2.0	15.2	65.2	213.1	533.4
2005	22.2	*	*	*	*	*	0.1	2.0	15.0	68.4	224.1	561.7
2004	22.5	*	*	*	*	*	0.1	2.2	14.9	68.9	231.4	592.2
2003	23.1	*	*	*	*	*	0.1	2.2	15.8	73.1	240.9	609.4
2002	24.0	*	*	*	*	*	0.2	2.0	16.1	76.9	253.3	648.7
2001	24.3	*	*	*	*	*	0.1	2.2	15.9	79.5	261.4	686.3
2000	25.0	*	*	*	*	*	0.1	2.0	17.0	83.2	270.2	730.7
1999	25.6	*	*	*	*	*	0.2	2.1	17.4	89.4	281.8	728.2
Non-Hispanic 4/, female												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)												
Non-Hispanic white, both sexes												
2006 5/	10.9	*	*	*	*	*	0.1	0.8	6.3	26.1	80.7	163.1
2005 5/	11.1	*	*	*	*	*	*	0.8	6.1	27.6	84.5	167.8
2004 5/	11.3	*	*	*	*	*	*	0.9	6.2	28.1	86.0	174.0
2003 5/	11.6	*	*	*	*	*	*	0.9	6.3	29.8	89.1	178.2
2002	11.9	*	*	*	*	*	0.1	0.8	6.5	30.9	93.1	186.6
2001	12.0	*	*	*	*	*	*	0.9	6.4	31.9	95.1	189.5
2000	12.3	*	*	*	*	*	*	0.8	6.9	32.8	97.6	199.3
1999	12.6	*	*	*	*	*	0.1	0.9	6.9	35.6	102.6	195.4
Non-Hispanic white, male												
2006 5/	22.2	*	*	*	*	*	0.2	1.6	12.9	56.2	196.9	512.8
2005 5/	22.7	*	*	*	*	*	*	1.6	12.5	59.5	207.2	534.8
2004 5/	23.0	*	*	*	*	*	*	1.7	12.7	60.8	212.3	562.6
2003 5/	23.6	*	*	*	*	*	*	1.8	13.0	64.5	221.0	583.0
2002	24.3	*	*	*	*	*	0.2	1.5	13.4	67.1	232.1	618.5
2001	24.6	*	*	*	*	*	*	1.8	13.3	69.6	238.4	645.0
2000	25.2	*	*	*	*	*	*	1.7	14.2	71.7	246.1	693.6
1999	25.9	*	*	*	*	*	0.1	1.8	14.3	78.2	259.4	685.4
Non-Hispanic white, female												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)												
Non-Hispanic black, both sexes												
2006 5/	12.3	*	*	*	*	*	*	2.3	17.3	70.8	171.1	275.5
2005 5/	12.8	*	*	*	*	*	*	2.4	18.3	72.7	177.9	299.6
2004 5/	12.9	*	*	*	*	*	*	2.9	17.5	70.4	190.8	313.7
2003 5/	13.3	*	*	*	*	*	*	2.7	20.1	74.6	198.8	304.6
2002	14.1	*	*	*	*	*	*	2.6	19.5	82.1	212.3	333.3
2001	14.6	*	*	*	*	*	*	2.9	19.5	84.2	221.6	365.8
2000	15.1	*	*	*	*	*	*	2.4	20.8	91.6	230.8	359.6
1999	15.3	*	*	*	*	*	*	2.7	22.0	94.7	227.5	368.6
Non-Hispanic black, male												
2006 5/	25.8	*	*	*	*	*	*	5.0	38.7	169.9	472.7	951.1
2005 5/	26.8	*	*	*	*	*	*	5.3	40.9	174.5	491.7	1,043.8
2004 5/	27.1	*	*	*	*	*	*	6.4	39.3	169.2	528.8	1,107.6
2003 5/	27.9	*	*	*	*	*	*	5.7	45.1	179.5	550.9	1,091.0
2002	29.6	*	*	*	*	*	*	5.7	43.8	197.9	587.6	1,205.5
2001	30.8	*	*	*	*	*	*	6.2	43.7	203.5	616.3	1,353.6
2000	31.7	*	*	*	*	*	*	5.3	46.6	221.7	640.8	1,328.0
1999	32.2	*	*	*	*	*	*	5.8	49.5	229.1	633.8	1,354.9
Non-Hispanic black, female												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
All origins 3/, both sexes												
2006	4.1	*	*	0.1	0.0	0.1	0.6	2.8	7.8	16.3	26.2	34.3
2005	4.2	*	*	0.1	*	0.1	0.7	3.0	8.3	16.9	26.9	32.6
2004	4.2	*	*	0.1	*	0.1	0.6	3.0	8.7	16.7	26.8	32.7
2003	4.2	*	*	0.1	0.1	0.1	0.7	2.9	9.0	16.9	27.0	33.2
2002	4.2	*	*	0.1	0.0	0.1	0.7	3.3	8.8	17.7	26.3	32.5
2001	4.2	*	0.1	0.1	0.1	0.1	0.8	3.3	9.2	17.6	26.3	32.3
2000	4.2	*	*	0.1	0.1	0.1	0.7	3.3	9.0	17.1	26.6	33.8
1999	4.0	*	*	0.1	*	0.1	0.8	3.1	8.9	16.7	24.6	31.4
All origins 3/, male												
2006	5.2	*	*	*	*	0.2	0.9	4.0	11.4	23.1	37.1	52.1
2005	5.3	*	*	*	*	0.2	0.8	4.3	12.0	24.4	37.9	50.6
2004	5.3	*	*	*	*	0.2	0.8	4.3	12.8	23.9	38.2	47.0
2003	5.4	*	*	*	*	0.2	1.0	4.3	13.3	23.7	39.5	48.7
2002	5.4	*	*	*	*	0.2	1.0	4.8	13.0	25.1	37.9	50.1
2001	5.4	*	*	*	*	0.1	1.1	4.9	13.4	25.2	37.4	52.9
2000	5.3	*	*	*	*	0.1	1.0	4.8	13.1	24.2	38.2	55.1
1999	5.0	*	*	*	*	0.2	1.1	4.5	12.9	23.6	35.7	49.4
All origins 3/, female												
2006	3.1	*	*	*	*	0.1	0.4	1.6	4.5	10.5	18.7	25.9
2005	3.1	*	*	*	*	*	0.5	1.8	4.8	10.5	19.4	24.3
2004	3.1	*	*	*	*	*	0.4	1.7	5.0	10.6	19.1	26.3
2003	3.1	*	*	0.1	*	0.1	0.4	1.6	4.9	11.2	18.7	26.4
2002	3.1	*	*	*	*	*	0.4	1.8	4.9	11.6	18.7	24.9
2001	3.1	*	*	0.1	*	0.1	0.5	1.9	5.3	11.3	19.0	23.7
2000	3.1	*	*	*	*	*	0.5	1.9	5.3	11.3	18.9	25.1
1999	3.0	*	*	*	*	*	0.5	1.8	5.1	11.0	17.4	24.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
Hispanic, both sexes												
2006	1.7	*	*	*	*	*	0.4	2.3	6.9	13.2	22.1	23.5
2005	1.8	*	*	*	*	*	0.7	2.8	6.3	15.7	23.4	32.9
2004	1.8	*	*	*	*	*	0.7	3.1	8.2	14.0	23.4	28.1
2003	1.7	*	*	*	*	*	0.5	2.1	8.1	16.2	23.6	26.9
2002	1.7	*	*	*	*	*	0.6	2.8	8.1	12.9	24.4	26.1
2001	1.7	*	*	*	*	*	0.7	3.3	6.9	14.4	23.3	31.9
2000	1.6	*	*	*	*	*	0.6	2.8	7.9	12.3	24.7	25.9
1999	1.6	*	*	*	*	*	0.6	2.8	7.9	13.9	22.5	28.8
Hispanic, male												
2006	2.0	*	*	*	*	*	*	3.3	9.7	19.8	32.3	32.7
2005	2.2	*	*	*	*	*	0.7	3.4	9.3	22.5	33.1	46.1
2004	2.4	*	*	*	*	*	0.8	4.4	11.2	23.3	34.7	43.4
2003	2.1	*	*	*	*	*	0.7	2.8	11.0	22.3	32.2	38.5
2002	2.1	*	*	*	*	*	0.9	3.9	11.3	18.5	39.0	40.2
2001	2.2	*	*	*	*	*	1.2	4.7	9.9	21.4	35.3	57.1
2000	2.0	*	*	*	*	*	1.0	3.9	11.4	17.9	38.4	40.3
1999	2.0	*	*	*	*	*	0.9	4.3	11.8	21.1	29.7	*
Hispanic, female												
2006	1.2	*	*	*	*	*	*	1.3	4.3	7.8	15.0	18.6
2005	1.4	*	*	*	*	*	*	2.2	3.5	10.1	16.7	26.1
2004	1.3	*	*	*	*	*	*	1.7	5.5	6.5	15.6	20.2
2003	1.4	*	*	*	*	*	*	1.5	5.6	11.3	17.6	21.1
2002	1.2	*	*	*	*	*	*	1.7	5.2	8.4	14.3	19.1
2001	1.2	*	*	*	*	*	*	1.9	4.2	8.8	15.1	19.5
2000	1.1	*	*	*	*	*	*	1.8	4.7	7.8	15.5	*
1999	1.2	*	*	*	*	*	*	1.4	4.4	8.3	17.7	23.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
Non-Hispanic 4/, both sexes												
2006	4.6	*	*	0.1	0.1	0.2	0.6	2.9	7.9	16.5	26.4	34.8
2005	4.6	*	*	0.1	*	0.1	0.7	3.0	8.4	16.9	27.1	32.5
2004	4.6	*	*	0.1	*	0.1	0.6	3.0	8.7	16.8	26.9	32.9
2003	4.6	*	*	0.1	*	0.2	0.8	3.0	9.0	16.9	27.2	33.4
2002	4.6	*	*	0.1	*	0.1	0.7	3.3	8.8	18.0	26.4	32.7
2001	4.6	*	*	0.1	0.1	0.1	0.8	3.3	9.4	17.8	26.4	32.3
2000	4.5	*	*	0.1	*	0.1	0.8	3.3	9.1	17.3	26.6	33.9
1999	4.3	*	*	0.1	*	0.1	0.8	3.2	8.9	16.8	24.6	31.4
Non-Hispanic 4/, male												
2006	5.8	*	*	*	*	0.2	0.9	4.1	11.5	23.3	37.3	53.2
2005	5.9	*	*	*	*	0.2	0.8	4.3	12.2	24.4	38.1	50.7
2004	5.8	*	*	*	*	0.2	0.8	4.3	12.9	23.9	38.4	47.1
2003	5.9	*	*	*	*	0.2	1.1	4.4	13.5	23.7	39.8	49.0
2002	5.9	*	*	*	*	0.2	1.0	4.9	13.0	25.5	37.7	50.4
2001	5.9	*	*	*	*	0.1	1.0	4.9	13.7	25.4	37.5	52.6
2000	5.7	*	*	*	*	0.1	1.0	4.8	13.2	24.4	38.1	55.4
1999	5.4	*	*	*	*	0.2	1.1	4.6	13.0	23.7	35.9	49.8
Non-Hispanic 4/, female												
2006	3.4	*	*	*	*	*	0.4	1.7	4.4	10.8	18.9	26.2
2005	3.4	*	*	*	*	*	0.5	1.7	5.0	10.6	19.6	24.2
2004	3.4	*	*	*	*	*	0.4	1.7	4.9	10.9	19.3	26.5
2003	3.4	*	*	*	*	0.1	0.4	1.7	4.9	11.1	18.7	26.6
2002	3.4	*	*	*	*	*	0.4	1.8	4.9	11.8	18.8	25.1
2001	3.4	*	*	0.1	*	0.1	0.6	1.8	5.4	11.4	19.1	23.8
2000	3.4	*	*	*	*	*	0.5	1.9	5.3	11.4	19.1	25.2
1999	3.2	*	*	*	*	*	0.5	1.8	5.2	11.2	17.3	24.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
Non-Hispanic white, both sexes												
2006 5/	5.1	*	*	*	*	0.1	0.6	2.9	8.2	17.1	27.2	35.5
2005 5/	5.1	*	*	*	*	*	0.7	3.1	8.6	17.5	27.8	33.0
2004 5/	5.1	*	*	0.1	*	0.1	0.6	3.1	9.1	17.3	27.5	33.8
2003 5/	5.1	*	*	0.1	*	0.1	0.7	3.1	9.2	17.4	27.8	34.0
2002	5.1	*	*	0.1	*	0.1	0.7	3.4	9.1	18.7	26.8	33.5
2001	5.1	*	*	*	*	0.1	0.7	3.4	9.6	18.4	26.8	32.8
2000	5.0	*	*	*	*	0.1	0.7	3.4	9.3	17.7	26.9	34.6
1999	4.7	*	*	0.1	*	0.1	0.7	3.1	9.2	17.2	24.9	31.5
Non-Hispanic white, male												
2006 5/	6.4	*	*	*	*	*	0.9	4.1	11.7	23.9	38.5	54.6
2005 5/	6.5	*	*	*	*	*	0.8	4.3	12.3	25.1	38.9	52.1
2004 5/	6.5	*	*	*	*	*	0.8	4.5	13.3	24.4	39.1	48.6
2003 5/	6.6	*	*	*	*	*	1.0	4.5	13.7	24.3	40.7	50.4
2002	6.5	*	*	*	*	*	1.0	5.0	13.3	26.3	38.4	52.4
2001	6.5	*	*	*	*	*	1.0	4.9	13.8	26.2	37.7	54.3
2000	6.3	*	*	*	*	*	1.0	4.9	13.5	24.6	38.5	56.9
1999	6.0	*	*	*	*	*	1.0	4.5	13.3	24.0	36.5	50.2
Non-Hispanic white, female												
2006 5/	3.8	*	*	*	*	*	0.3	1.6	4.8	11.2	19.4	26.6
2005 5/	3.8	*	*	*	*	*	0.5	1.8	5.1	11.0	20.1	24.3
2004 5/	3.8	*	*	*	*	*	0.3	1.8	5.1	11.2	19.6	27.2
2003 5/	3.8	*	*	*	*	*	0.4	1.7	4.8	11.6	19.1	26.8
2002	3.8	*	*	*	*	*	0.4	1.8	5.1	12.2	19.0	25.3
2001	3.8	*	*	*	*	*	0.5	1.8	5.5	11.8	19.5	23.8
2000	3.8	*	*	*	*	*	0.5	1.9	5.4	11.8	19.3	25.7
1999	3.5	*	*	*	*	*	0.5	1.8	5.3	11.5	17.3	24.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
Non-Hispanic black, both sexes												
2006 5/	3.0	*	*	*	*	0.4	1.0	3.6	7.5	15.1	22.5	33.7
2005 5/	3.1	*	*	*	*	0.4	0.9	3.3	9.1	15.5	24.4	30.9
2004 5/	2.9	*	*	*	*	*	0.8	3.2	8.2	16.0	26.3	28.8
2003 5/	3.0	*	*	*	*	*	1.1	3.3	9.7	14.8	24.2	34.2
2002	2.9	*	*	*	*	*	0.8	3.6	8.8	15.8	26.5	28.1
2001	3.1	*	*	*	*	*	1.3	3.8	9.7	15.3	26.0	30.0
2000	2.9	*	*	*	*	*	0.9	3.5	9.0	16.8	25.9	29.6
1999	3.0	*	*	*	*	0.4	1.1	3.9	8.7	16.4	23.9	31.7
Non-Hispanic black, male												
2006 5/	3.9	*	*	*	*	0.8	1.6	5.3	12.3	22.9	32.8	49.5
2005 5/	4.0	*	*	*	*	*	1.4	5.3	14.2	24.3	35.9	40.0
2004 5/	3.7	*	*	*	*	*	1.1	4.6	12.4	24.1	38.2	39.6
2003 5/	3.8	*	*	*	*	*	1.7	5.0	14.1	22.3	37.2	44.7
2002	3.6	*	*	*	*	*	1.2	5.0	14.4	22.1	36.6	39.4
2001	4.0	*	*	*	*	*	1.5	5.5	15.3	22.7	40.8	46.2
2000	3.7	*	*	*	*	*	1.0	5.0	13.4	26.8	38.3	49.3
1999	3.7	*	*	*	*	*	1.6	6.0	13.3	23.9	32.5	49.9
Non-Hispanic black, female												
2006 5/	2.1	*	*	*	*	*	*	2.2	3.6	9.5	16.7	27.3
2005 5/	2.2	*	*	*	*	*	*	1.7	5.0	9.2	18.0	27.2
2004 5/	2.3	*	*	*	*	*	*	1.9	4.9	10.3	19.6	24.5
2003 5/	2.3	*	*	*	*	*	0.7	1.8	6.2	9.5	16.8	30.1
2002	2.3	*	*	*	*	*	*	2.5	4.4	11.3	20.7	23.8
2001	2.3	*	*	*	*	*	1.1	2.2	5.3	10.0	17.7	24.0
2000	2.2	*	*	*	*	*	0.8	2.2	5.4	9.7	18.9	22.2
1999	2.2	*	*	*	*	*	*	2.1	5.0	11.0	19.1	24.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
All origins 3/, both sexes												
2006	4.5	*	*	*	*	*	0.2	1.2	5.0	14.9	37.4	67.4
2005	4.5	*	*	*	*	*	0.3	1.2	4.9	14.9	37.2	68.3
2004	4.4	*	*	*	*	*	0.3	1.3	5.0	14.9	37.4	67.9
2003	4.3	*	*	*	*	*	0.2	1.3	5.0	14.9	36.4	64.8
2002	4.4	*	*	*	*	*	0.3	1.3	4.9	15.9	37.0	66.6
2001	4.3	*	*	*	*	*	0.3	1.2	5.0	15.4	35.7	68.9
2000	4.3	*	*	*	*	*	0.2	1.2	5.0	15.5	35.7	69.6
1999	4.3	*	*	*	*	*	0.2	1.3	5.2	15.9	35.3	68.5
All origins 3/, male												
2006	6.4	*	*	*	*	*	0.3	1.8	7.8	23.7	65.5	131.6
2005	6.3	*	*	*	*	*	0.4	1.7	7.6	24.3	65.1	132.1
2004	6.2	*	*	*	*	*	0.3	2.0	7.9	24.0	64.7	133.0
2003	6.0	*	*	*	*	*	0.3	2.0	7.8	23.9	63.1	125.1
2002	6.0	*	*	*	*	*	0.3	1.8	7.7	25.2	63.0	128.9
2001	6.0	*	*	*	*	*	0.3	1.8	7.9	25.2	62.5	134.8
2000	5.9	*	*	*	*	*	0.3	1.7	7.9	25.2	62.6	139.4
1999	5.9	*	*	*	*	*	0.3	2.0	8.2	25.7	62.5	134.8
All origins 3/, female												
2006	2.7	*	*	*	*	*	0.2	0.6	2.3	7.3	18.1	37.4
2005	2.7	*	*	*	*	*	0.2	0.8	2.3	6.9	18.3	39.0
2004	2.7	*	*	*	*	*	0.2	0.6	2.3	7.3	19.0	38.6
2003	2.6	*	*	*	*	*	0.2	0.6	2.4	7.4	18.5	38.1
2002	2.8	*	*	*	*	*	0.2	0.7	2.4	8.2	19.7	39.6
2001	2.7	*	*	*	*	*	0.3	0.6	2.3	7.2	18.1	41.4
2000	2.7	*	*	*	*	*	0.2	0.6	2.4	7.5	18.1	41.2
1999	2.7	*	*	*	*	*	0.2	0.6	2.5	7.8	17.7	41.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
Hispanic, both sexes												
2006	1.0	*	*	*	*	*	*	*	2.3	8.1	19.7	33.3
2005	1.0	*	*	*	*	*	*	0.5	2.2	8.1	20.7	39.4
2004	1.0	*	*	*	*	*	*	0.6	2.7	7.3	20.7	32.3
2003	1.0	*	*	*	*	*	*	0.7	2.6	7.9	20.8	42.0
2002	1.0	*	*	*	*	*	*	0.7	2.9	7.7	18.3	54.9
2001	0.9	*	*	*	*	*	*	0.6	2.2	7.9	18.8	46.3
2000	0.9	*	*	*	*	*	*	*	2.9	6.6	17.6	47.1
1999	0.9	*	*	*	*	*	*	0.7	3.1	8.5	16.5	38.6
Hispanic, male												
2006	1.2	*	*	*	*	*	*	*	3.5	11.0	35.1	54.1
2005	1.4	*	*	*	*	*	*	*	3.7	13.5	34.1	83.4
2004	1.2	*	*	*	*	*	*	1.1	4.3	11.0	28.6	58.8
2003	1.3	*	*	*	*	*	*	*	4.0	13.1	33.4	77.0
2002	1.3	*	*	*	*	*	*	*	4.2	12.8	27.9	100.4
2001	1.2	*	*	*	*	*	*	*	4.3	11.8	30.0	87.6
2000	1.2	*	*	*	*	*	*	*	4.6	10.6	33.9	84.6
1999	1.3	*	*	*	*	*	*	*	5.6	15.2	27.1	77.6
Hispanic, female												
2006	0.7	*	*	*	*	*	*	*	*	5.8	8.9	22.2
2005	0.6	*	*	*	*	*	*	*	*	3.7	11.4	16.3
2004	0.7	*	*	*	*	*	*	*	*	4.4	15.3	18.7
2003	0.7	*	*	*	*	*	*	*	*	3.7	12.1	24.2
2002	0.7	*	*	*	*	*	*	*	*	3.6	11.8	32.3
2001	0.6	*	*	*	*	*	*	*	*	4.9	11.2	26.1
2000	0.5	*	*	*	*	*	*	*	*	3.5	6.6	28.7
1999	0.5	*	*	*	*	*	*	*	*	*	9.4	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
Non-Hispanic 4/, both sexes												
2006	5.1	*	*	*	*	*	0.3	1.3	5.2	15.4	38.4	69.0
2005	5.0	*	*	*	*	*	0.3	1.3	5.1	15.4	38.2	69.6
2004	5.0	*	*	*	*	*	0.3	1.4	5.2	15.4	38.2	69.4
2003	4.8	*	*	*	*	*	0.2	1.3	5.2	15.4	37.1	65.6
2002	4.9	*	*	*	*	*	0.3	1.3	5.1	16.4	37.8	66.9
2001	4.8	*	*	*	*	*	0.3	1.3	5.2	15.8	36.4	69.6
2000	4.7	*	*	*	*	*	0.3	1.2	5.2	16.1	36.3	70.3
1999	4.7	*	*	*	*	*	0.2	1.3	5.4	16.3	36.0	69.3
Non-Hispanic 4/, male												
2006	7.3	*	*	*	*	*	0.4	2.0	8.2	24.7	67.4	135.5
2005	7.1	*	*	*	*	*	0.4	1.8	7.9	25.1	66.9	134.6
2004	7.1	*	*	*	*	*	0.4	2.1	8.1	24.9	66.6	136.5
2003	6.8	*	*	*	*	*	0.3	2.1	8.1	24.7	64.5	127.2
2002	6.8	*	*	*	*	*	0.4	1.9	7.9	26.0	64.7	129.8
2001	6.7	*	*	*	*	*	0.4	1.9	8.1	26.0	63.8	136.4
2000	6.6	*	*	*	*	*	0.3	1.8	8.1	26.1	63.6	141.3
1999	6.6	*	*	*	*	*	0.3	2.0	8.4	26.3	63.8	136.6
Non-Hispanic 4/, female												
2006	3.0	*	*	*	*	*	0.2	0.7	2.4	7.4	18.6	38.1
2005	3.0	*	*	*	*	*	0.2	0.8	2.5	7.2	18.7	40.0
2004	3.0	*	*	*	*	*	0.2	0.7	2.4	7.5	19.1	39.4
2003	2.9	*	*	*	*	*	0.2	0.6	2.5	7.6	18.8	38.6
2002	3.1	*	*	*	*	*	0.2	0.8	2.4	8.5	20.0	39.8
2001	3.0	*	*	*	*	*	0.3	0.6	2.5	7.4	18.3	41.9
2000	3.0	*	*	*	*	*	0.2	0.6	2.5	7.8	18.5	41.6
1999	3.0	*	*	*	*	*	0.2	0.6	2.6	8.1	17.9	42.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
Non-Hispanic white, both sexes												
2006 5/	5.9	*	*	*	*	*	0.3	1.4	5.5	16.1	40.1	71.5
2005 5/	5.8	*	*	*	*	*	0.3	1.3	5.4	16.2	39.8	71.5
2004 5/	5.8	*	*	*	*	*	0.3	1.4	5.5	16.4	40.1	71.1
2003 5/	5.5	*	*	*	*	*	0.2	1.3	5.5	16.3	38.3	67.6
2002	5.6	*	*	*	*	*	0.3	1.4	5.3	17.1	39.1	68.0
2001	5.5	*	*	*	*	*	0.3	1.3	5.4	16.6	37.4	71.5
2000	5.4	*	*	*	*	*	0.3	1.3	5.5	16.8	37.1	72.8
1999	5.4	*	*	*	*	*	0.2	1.3	5.5	16.8	37.1	70.7
Non-Hispanic white, male												
2006 5/	8.6	*	*	*	*	*	0.4	2.1	8.6	26.3	71.1	142.3
2005 5/	8.3	*	*	*	*	*	0.5	1.9	8.4	26.6	70.2	140.2
2004 5/	8.3	*	*	*	*	*	0.3	2.2	8.7	26.7	70.5	143.1
2003 5/	7.9	*	*	*	*	*	0.3	2.2	8.6	26.1	67.2	133.6
2002	7.9	*	*	*	*	*	0.4	2.0	8.5	27.4	67.2	135.2
2001	7.8	*	*	*	*	*	0.4	2.0	8.6	27.6	66.6	143.0
2000	7.7	*	*	*	*	*	0.3	2.0	8.6	27.5	65.9	148.0
1999	7.6	*	*	*	*	*	0.3	2.0	8.7	27.6	66.2	143.0
Non-Hispanic white, female												
2006 5/	3.3	*	*	*	*	*	0.2	0.7	2.5	7.3	18.5	38.5
2005 5/	3.4	*	*	*	*	*	0.2	0.8	2.5	7.3	18.9	40.1
2004 5/	3.3	*	*	*	*	*	0.2	0.7	2.5	7.5	19.4	38.9
2003 5/	3.3	*	*	*	*	*	0.2	0.5	2.5	7.8	18.7	38.6
2002	3.4	*	*	*	*	*	0.2	0.8	2.3	8.4	20.2	39.0
2001	3.2	*	*	*	*	*	0.3	0.6	2.4	7.4	18.1	41.7
2000	3.3	*	*	*	*	*	0.2	0.7	2.6	7.8	18.2	42.5
1999	3.2	*	*	*	*	*	0.2	0.7	2.5	7.8	18.0	41.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
Non-Hispanic black, both sexes												
2006 5/	2.7	*	*	*	*	*	*	1.3	4.9	13.6	31.7	55.7
2005 5/	2.5	*	*	*	*	*	*	1.5	4.7	12.5	29.8	57.6
2004 5/	2.4	*	*	*	*	*	0.5	1.6	4.2	11.9	27.6	61.3
2003 5/	2.5	*	*	*	*	*	0.4	1.7	4.1	12.3	32.8	53.8
2002	2.6	*	*	*	*	*	*	1.1	4.5	14.7	31.2	64.1
2001	2.4	*	*	*	*	*	0.4	1.3	5.1	12.1	31.4	55.6
2000	2.4	*	*	*	*	*	*	1.1	4.2	13.5	34.9	49.6
1999	2.6	*	*	*	*	*	*	1.5	5.6	15.2	29.4	60.8
Non-Hispanic black, male												
2006 5/	3.1	*	*	*	*	*	*	1.9	7.7	18.2	42.8	95.3
2005 5/	3.0	*	*	*	*	*	*	1.7	6.6	18.3	45.7	101.8
2004 5/	2.7	*	*	*	*	*	*	2.3	6.0	15.0	40.2	83.3
2003 5/	2.9	*	*	*	*	*	*	2.2	5.5	17.8	49.6	79.9
2002	2.8	*	*	*	*	*	*	1.3	5.5	18.7	48.1	91.9
2001	2.6	*	*	*	*	*	*	1.8	6.5	16.5	43.2	78.5
2000	2.8	*	*	*	*	*	*	1.4	5.9	17.8	52.8	88.1
1999	3.0	*	*	*	*	*	*	2.6	7.4	19.2	46.6	86.8
Non-Hispanic black, female												
2006 5/	2.3	*	*	*	*	*	*	0.7	2.7	10.4	25.4	39.6
2005 5/	2.1	*	*	*	*	*	*	1.3	3.1	8.4	20.8	39.8
2004 5/	2.2	*	*	*	*	*	*	1.0	2.7	9.7	20.4	52.6
2003 5/	2.1	*	*	*	*	*	*	1.2	2.9	8.3	23.3	43.7
2002	2.4	*	*	*	*	*	*	0.9	3.7	11.8	21.7	53.5
2001	2.3	*	*	*	*	*	*	*	3.9	9.0	24.8	47.1
2000	2.1	*	*	*	*	*	*	*	2.7	10.4	24.8	35.3
1999	2.3	*	*	*	*	*	*	*	4.2	12.3	19.8	51.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
All origins 3/, both sexes												
2006	4.3	*	0.6	0.8	0.5	0.9	2.0	4.5	9.3	14.7	19.0	16.7
2005	4.4	*	0.5	0.8	0.6	0.9	2.2	4.6	9.4	15.8	20.2	15.8
2004	4.4	*	0.6	0.8	0.5	0.9	2.0	4.6	9.5	15.7	19.6	16.6
2003	4.4	*	0.7	0.8	0.6	1.0	2.4	4.8	9.7	15.5	19.5	15.2
2002	4.4	*	0.7	0.8	0.5	1.0	2.2	5.1	9.6	16.2	19.1	16.4
2001	4.4	*	0.7	0.7	0.6	1.1	2.3	5.0	9.7	15.4	20.4	16.4
2000	4.5	0.5	0.7	0.8	0.6	1.0	2.4	5.1	9.6	16.5	20.0	16.9
1999	4.6	*	0.7	0.8	0.6	1.1	2.4	5.1	10.3	16.9	20.4	16.0
All origins 3/, male												
2006	4.9	*	0.7	0.8	0.5	1.0	2.4	5.5	11.4	18.2	24.0	21.2
2005	5.0	*	0.5	0.8	0.7	1.2	2.8	5.6	11.3	19.3	25.3	19.3
2004	4.9	*	0.7	0.8	0.6	1.1	2.5	5.8	11.8	18.4	23.7	19.4
2003	5.0	*	0.7	0.8	0.6	1.1	2.9	6.2	11.4	18.6	24.6	20.0
2002	5.0	*	0.8	0.9	0.7	1.1	2.8	6.2	11.7	20.2	23.8	20.1
2001	5.0	*	0.7	0.7	0.6	1.3	2.8	6.1	12.1	19.2	25.8	20.1
2000	5.1	*	0.7	0.8	0.7	1.3	3.1	6.2	11.8	20.2	24.3	22.4
1999	5.1	*	0.7	0.7	0.7	1.4	2.8	6.2	12.4	20.8	25.5	20.7
All origins 3/, female												
2006	3.7	*	0.6	0.7	0.4	0.8	1.5	3.6	7.3	11.8	15.5	14.6
2005	3.9	*	0.6	0.8	0.4	0.7	1.6	3.6	7.5	12.9	16.7	14.3
2004	3.9	*	0.6	0.8	0.4	0.7	1.6	3.4	7.3	13.4	16.8	15.4
2003	3.9	*	0.6	0.8	0.5	0.9	2.0	3.5	8.1	12.8	16.1	13.1
2002	3.9	*	0.7	0.7	0.4	0.8	1.7	4.1	7.7	12.8	16.0	14.8
2001	3.9	*	0.6	0.8	0.5	0.8	1.7	3.9	7.6	12.3	16.8	14.8
2000	3.9	*	0.6	0.7	0.5	0.7	1.7	4.0	7.6	13.4	17.3	14.7
1999	4.1	*	0.8	0.8	0.5	0.8	1.9	4.1	8.4	13.7	17.1	14.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Hispanic, both sexes												
2006	1.6	*	0.7	0.7	0.3	0.6	1.1	2.9	5.4	7.5	13.0	12.1
2005	1.8	*	0.6	0.7	0.5	0.5	1.3	2.9	5.9	10.2	13.4	15.4
2004	1.6	*	*	0.7	0.5	0.4	1.1	2.4	5.9	9.0	12.4	14.3
2003	1.7	*	0.6	0.8	0.3	0.6	1.1	2.5	5.9	9.5	15.3	*
2002	1.7	*	0.7	0.7	0.4	0.5	0.9	3.4	5.8	10.0	13.7	13.3
2001	1.7	*	0.7	0.5	0.4	0.6	1.4	2.6	5.9	10.3	14.8	*
2000	1.8	*	*	0.9	0.4	0.7	1.5	2.9	6.2	9.8	14.2	15.9
1999	1.7	*	0.9	0.7	*	0.5	1.3	3.0	6.0	10.0	14.0	*
Hispanic, male												
2006	1.7	*	*	0.8	*	0.5	1.5	3.3	6.0	9.2	16.1	*
2005	2.0	*	*	0.5	0.6	0.6	1.6	3.6	7.4	11.5	18.0	*
2004	1.7	*	*	0.6	0.6	*	1.4	2.9	7.2	10.1	15.4	*
2003	1.8	*	*	0.9	*	0.6	1.2	3.2	6.3	11.8	16.5	*
2002	1.8	*	*	0.7	*	*	1.0	4.4	6.6	12.4	16.0	*
2001	1.8	*	*	*	*	0.6	1.4	3.0	6.8	14.4	17.9	*
2000	1.9	*	*	1.2	*	0.8	1.5	2.6	7.7	12.5	18.7	*
1999	1.7	*	*	0.7	*	*	1.8	2.9	7.8	10.5	16.2	*
Hispanic, female												
2006	1.5	*	*	0.7	*	0.6	0.7	2.4	4.8	6.1	10.8	14.4
2005	1.6	*	*	0.9	*	*	1.0	2.2	4.6	9.2	10.2	15.0
2004	1.5	*	*	0.9	*	*	0.8	1.8	4.7	8.1	10.4	15.1
2003	1.6	*	*	0.7	*	*	0.9	1.9	5.6	7.6	14.5	*
2002	1.5	*	*	0.7	*	*	0.9	2.3	5.1	8.1	12.1	*
2001	1.6	*	*	0.6	*	*	1.4	2.2	5.1	7.1	12.7	*
2000	1.7	*	*	0.7	*	*	1.5	3.2	4.9	7.6	11.2	*
1999	1.7	*	*	0.8	*	*	0.8	3.1	4.5	9.6	12.5	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Non-Hispanic 4/, both sexes												
2006	4.8	*	0.6	0.8	0.5	1.0	2.1	4.7	9.6	15.2	19.3	16.9
2005	4.9	*	0.5	0.8	0.6	1.1	2.3	4.8	9.6	16.2	20.6	15.9
2004	4.8	*	0.7	0.8	0.5	1.0	2.2	4.8	9.7	16.2	20.0	16.7
2003	4.9	*	0.7	0.8	0.6	1.1	2.6	5.0	10.0	15.9	19.7	15.4
2002	4.9	*	0.7	0.8	0.5	1.1	2.4	5.3	9.9	16.6	19.4	16.5
2001	4.8	*	0.6	0.8	0.6	1.2	2.4	5.2	10.0	15.7	20.6	16.5
2000	4.9	*	0.7	0.7	0.6	1.1	2.5	5.3	9.8	16.9	20.2	16.9
1999	5.0	*	0.7	0.8	0.6	1.2	2.5	5.3	10.6	17.3	20.6	16.1
Non-Hispanic 4/, male												
2006	5.4	*	0.6	0.8	0.6	1.2	2.6	5.7	11.9	18.8	24.4	21.9
2005	5.5	*	0.5	0.9	0.7	1.4	3.0	5.8	11.6	19.8	25.7	19.4
2004	5.4	*	0.7	0.9	0.6	1.2	2.7	6.1	12.2	19.0	24.2	19.7
2003	5.5	*	0.7	0.8	0.7	1.3	3.1	6.5	11.8	19.1	25.0	20.1
2002	5.5	*	0.8	0.9	0.7	1.3	3.0	6.4	12.0	20.7	24.1	20.2
2001	5.5	*	0.6	0.7	0.7	1.5	3.0	6.4	12.5	19.4	26.1	20.5
2000	5.5	*	0.7	0.8	0.7	1.4	3.3	6.5	12.1	20.6	24.5	22.3
1999	5.6	*	0.7	0.8	0.7	1.5	3.0	6.5	12.7	21.4	25.8	21.3
Non-Hispanic 4/, female												
2006	4.1	*	0.6	0.7	0.5	0.9	1.6	3.7	7.5	12.2	15.8	14.6
2005	4.2	*	0.5	0.8	0.4	0.7	1.7	3.8	7.8	13.1	17.0	14.2
2004	4.2	*	0.6	0.7	0.4	0.8	1.7	3.5	7.5	13.8	17.2	15.4
2003	4.3	*	0.6	0.8	0.5	1.0	2.1	3.7	8.3	13.2	16.2	13.4
2002	4.2	*	0.7	0.7	0.4	0.9	1.8	4.2	7.9	13.1	16.2	14.9
2001	4.2	*	0.6	0.8	0.5	0.8	1.8	4.1	7.7	12.7	17.0	14.9
2000	4.2	*	0.7	0.7	0.5	0.7	1.8	4.1	7.8	13.8	17.5	14.7
1999	4.4	*	0.7	0.8	0.5	0.9	2.0	4.2	8.6	14.0	17.2	14.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Non-Hispanic white, both sexes												
2006 5/	5.5	*	0.6	0.8	0.5	1.2	2.4	5.3	10.5	16.6	20.7	17.7
2005 5/	5.6	*	0.5	0.9	0.6	1.2	2.6	5.3	10.5	17.6	21.8	16.5
2004 5/	5.6	*	0.7	0.8	0.6	1.1	2.5	5.3	10.7	17.6	21.3	17.4
2003 5/	5.6	*	0.7	0.8	0.7	1.3	3.0	5.5	10.8	17.1	20.9	15.9
2002	5.6	*	0.6	0.8	0.6	1.2	2.8	5.9	10.7	17.9	20.4	17.3
2001	5.6	*	0.7	0.8	0.6	1.3	2.7	5.7	10.8	16.8	21.9	17.2
2000	5.6	*	0.7	0.8	0.6	1.2	2.8	5.9	10.7	18.3	21.2	17.5
1999	5.7	*	0.7	0.8	0.6	1.4	2.8	5.8	11.5	18.7	21.7	16.7
Non-Hispanic white, male												
2006 5/	6.3	*	0.7	0.9	0.6	1.4	3.0	6.3	12.9	20.3	26.0	22.9
2005 5/	6.4	*	0.4	0.9	0.7	1.5	3.4	6.3	12.5	21.4	26.9	20.4
2004 5/	6.3	*	0.8	0.9	0.7	1.4	3.0	6.6	13.2	20.8	25.4	21.0
2003 5/	6.3	*	0.8	0.8	0.7	1.5	3.5	7.0	12.8	20.4	26.3	21.0
2002	6.4	*	0.7	0.9	0.8	1.4	3.5	7.1	12.8	22.3	25.2	21.4
2001	6.3	*	0.7	0.7	0.7	1.7	3.5	7.0	13.4	20.9	27.5	20.9
2000	6.3	*	0.7	0.8	0.7	1.5	3.7	7.1	13.1	22.2	25.7	23.5
1999	6.4	*	0.8	0.8	0.7	1.8	3.3	7.1	13.7	23.0	27.2	22.4
Non-Hispanic white, female												
2006 5/	4.8	*	0.5	0.7	0.5	0.9	1.8	4.2	8.3	13.3	17.1	15.3
2005 5/	4.9	*	0.6	0.8	0.5	0.9	1.9	4.2	8.6	14.3	18.2	14.8
2004 5/	4.9	*	0.7	0.7	0.4	0.9	1.9	3.9	8.3	14.9	18.5	15.8
2003 5/	4.9	*	0.6	0.9	0.6	1.1	2.5	4.1	9.0	14.3	17.3	13.7
2002	4.9	*	0.6	0.8	0.4	1.1	2.1	4.7	8.7	14.2	17.1	15.5
2001	4.8	*	0.6	0.9	0.5	0.9	2.0	4.5	8.4	13.4	18.2	15.7
2000	4.9	*	0.7	0.7	0.5	0.9	1.9	4.6	8.4	15.0	18.3	15.1
1999	5.0	*	0.7	0.7	0.5	1.1	2.3	4.6	9.5	15.1	18.1	14.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Non-Hispanic black, both sexes												
2006 5/	2.0	*	*	0.7	0.5	0.5	1.2	2.6	5.4	8.2	9.6	9.6
2005 5/	2.1	*	*	0.8	0.4	0.5	1.2	2.8	5.5	8.5	12.3	9.6
2004 5/	2.0	*	*	0.8	0.5	0.7	1.3	2.9	5.0	7.9	10.1	10.6
2003 5/	2.2	*	*	0.7	0.6	0.6	1.5	3.1	5.6	9.0	10.6	10.7
2002	2.1	*	1.1	0.8	0.4	0.6	1.2	2.8	5.6	8.6	11.9	9.7
2001	2.1	*	*	0.7	0.5	0.5	1.1	3.1	5.9	9.4	9.9	11.9
2000	2.2	*	*	0.8	0.6	0.7	1.6	2.9	5.8	8.2	12.1	10.5
1999	2.2	*	*	0.8	0.7	0.7	1.4	3.3	6.1	8.2	11.7	10.0
Non-Hispanic black, male												
2006 5/	2.3	*	*	0.6	0.6	*	1.5	3.2	7.2	10.4	12.5	*
2005 5/	2.4	*	*	0.8	*	0.8	1.4	3.7	7.2	10.6	17.9	*
2004 5/	2.3	*	*	0.7	*	*	1.9	4.0	6.9	8.1	14.5	*
2003 5/	2.5	*	*	0.8	0.7	*	2.1	4.1	6.8	11.9	15.6	*
2002	2.3	*	*	1.1	*	0.9	1.3	2.7	8.1	10.2	16.1	*
2001	2.2	*	*	0.8	*	*	1.3	3.7	7.9	10.4	12.0	*
2000	2.4	*	*	0.9	0.8	1.1	2.0	3.7	7.2	9.9	13.3	*
1999	2.4	*	*	0.7	0.9	0.9	1.8	4.5	7.4	10.2	12.5	*
Non-Hispanic black, female												
2006 5/	1.8	*	*	0.8	*	*	0.9	2.0	4.0	6.6	7.9	9.0
2005 5/	1.8	*	*	0.7	*	*	1.0	2.0	4.2	7.0	9.1	10.0
2004 5/	1.8	*	*	0.8	*	*	0.8	1.9	3.5	7.8	7.6	11.6
2003 5/	1.9	*	*	0.7	*	*	0.9	2.3	4.7	6.9	7.8	12.0
2002	1.9	*	*	*	*	*	1.1	2.8	3.7	7.5	9.5	10.5
2001	1.9	*	*	*	*	*	1.0	2.5	4.2	8.7	8.8	9.8
2000	1.9	*	*	0.7	*	*	1.2	2.2	4.7	7.1	11.5	10.5
1999	1.9	*	*	0.9	*	*	1.0	2.2	5.1	6.8	11.2	9.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
All origins 3/, both sexes												
2006	18.4	0.8	0.8	0.7	1.6	2.0	3.5	8.5	24.4	65.2	137.8	186.1
2005	18.6	0.6	0.8	0.8	1.6	2.0	3.4	8.7	25.1	67.0	138.7	189.8
2004	18.6	0.6	0.8	0.9	1.5	2.1	3.4	9.0	25.8	68.2	139.8	189.4
2003	19.1	0.8	0.9	1.0	1.5	2.2	3.9	9.3	26.8	70.3	143.3	198.0
2002	19.5	0.7	0.8	0.9	1.7	2.3	3.9	9.9	28.2	73.9	143.6	197.5
2001	19.8	0.7	1.0	0.9	1.8	2.5	4.1	10.4	29.5	74.6	145.2	199.9
2000	20.1	1.0	0.9	0.9	1.8	2.3	4.1	10.7	30.0	76.3	149.2	204.6
1999	20.2	0.8	0.9	0.9	1.9	2.4	4.2	10.8	31.8	77.8	148.0	199.4
All origins 3/, male												
2006	20.3	*	0.9	0.9	2.0	2.4	4.1	10.5	30.8	82.4	183.1	244.7
2005	20.6	*	1.0	1.0	1.9	2.3	4.0	10.4	31.1	86.6	183.0	262.4
2004	20.4	*	0.9	1.0	1.9	2.4	4.2	11.0	32.1	87.4	181.3	254.9
2003	21.0	*	1.0	1.1	1.8	2.6	4.7	11.4	33.9	90.4	184.8	268.7
2002	21.4	*	0.8	1.0	2.0	2.7	4.6	11.9	35.0	95.2	188.0	269.1
2001	21.5	*	1.0	1.0	2.1	2.9	5.1	12.7	36.3	94.7	187.0	275.3
2000	21.6	1.1	1.0	1.1	2.3	2.7	4.8	13.0	36.1	96.8	192.6	281.8
1999	21.7	*	1.0	1.0	2.2	2.8	5.1	13.2	38.8	97.3	189.0	279.0
All origins 3/, female												
2006	16.5	*	0.8	0.6	1.1	1.7	2.9	6.6	18.3	50.7	106.8	158.7
2005	16.6	*	0.6	0.7	1.2	1.8	2.8	7.0	19.6	50.5	108.6	156.4
2004	16.8	*	0.6	0.7	1.2	1.7	2.7	7.0	19.9	52.0	111.8	160.0
2003	17.3	1.1	0.8	0.8	1.2	1.7	3.1	7.2	20.1	53.4	115.6	166.8
2002	17.7	*	0.7	0.8	1.4	1.9	3.2	7.9	21.9	56.2	114.1	166.5
2001	18.1	*	1.0	0.7	1.4	2.1	3.2	8.1	23.1	57.9	117.7	168.4
2000	18.6	*	0.9	0.7	1.4	1.9	3.4	8.5	24.4	59.5	120.9	173.1
1999	18.7	*	0.8	0.7	1.5	2.0	3.4	8.5	25.3	61.9	121.3	167.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Hispanic, both sexes												
2006	7.0	*	0.7	0.8	2.0	2.0	3.1	7.1	18.6	51.3	93.6	100.9
2005	7.3	*	0.8	1.1	2.2	1.9	2.8	8.1	20.9	52.7	97.6	107.8
2004	6.9	*	0.8	1.4	2.0	2.0	3.2	7.7	19.8	48.3	95.4	103.2
2003	7.1	*	1.3	1.3	1.9	2.0	3.4	8.5	20.4	47.1	99.1	129.6
2002	7.3	*	0.9	1.3	2.1	1.9	3.9	7.7	22.1	53.7	100.8	132.5
2001	7.4	*	1.3	1.2	2.2	2.2	3.9	8.2	23.9	52.6	103.3	134.4
2000	7.5	*	1.0	1.1	2.0	1.9	3.9	8.9	25.0	58.8	103.3	143.3
1999	7.5	*	1.0	1.2	2.0	2.1	3.9	10.1	24.2	58.5	101.0	131.9
Hispanic, male												
2006	7.4	*	*	0.8	2.5	2.1	3.5	8.3	22.4	60.5	115.7	129.7
2005	7.8	*	1.1	1.3	2.7	2.2	3.3	9.0	23.7	64.7	124.9	144.5
2004	7.6	*	*	1.8	2.6	2.2	4.1	9.9	23.9	60.8	117.0	119.1
2003	7.4	*	1.5	1.5	2.4	2.4	4.2	9.5	24.3	57.4	114.7	149.3
2002	7.6	*	*	1.3	2.7	2.3	4.4	8.9	26.2	63.3	119.0	180.7
2001	8.3	*	1.4	1.3	2.7	2.7	5.0	9.7	28.9	67.5	142.2	180.9
2000	8.0	*	*	1.4	2.3	2.4	4.3	11.1	30.7	71.3	126.9	197.5
1999	7.6	*	*	1.3	2.1	2.2	4.2	12.0	29.4	68.5	117.8	135.9
Hispanic, female												
2006	6.6	*	*	0.8	1.4	1.8	2.8	6.0	15.1	43.9	78.1	85.7
2005	6.7	*	*	0.9	1.5	1.6	2.4	7.0	18.4	43.0	78.6	88.6
2004	6.2	*	*	1.1	1.2	1.6	2.2	5.6	16.1	38.2	80.5	95.1
2003	6.8	*	*	1.1	1.3	1.5	2.6	7.4	16.8	39.0	88.5	119.5
2002	7.0	*	*	1.3	1.5	1.4	3.3	6.5	18.5	46.1	88.3	108.6
2001	6.5	*	*	1.1	1.5	1.7	2.6	6.7	19.5	40.8	77.1	111.7
2000	6.9	*	*	0.8	1.5	1.4	3.4	6.8	19.8	48.9	87.5	116.7
1999	7.4	*	*	1.0	1.9	2.1	3.4	8.3	19.6	50.6	89.9	130.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Non-Hispanic 4/, both sexes												
2006	20.3	0.8	0.8	0.7	1.5	2.1	3.6	8.7	24.8	66.2	140.4	190.1
2005	20.4	*	0.8	0.8	1.4	2.0	3.5	8.7	25.4	68.0	141.0	193.6
2004	20.5	*	0.8	0.7	1.4	2.1	3.5	9.1	26.2	69.6	142.1	193.0
2003	21.0	0.8	0.8	0.9	1.4	2.2	4.0	9.3	27.2	71.8	145.4	200.5
2002	21.3	0.8	0.7	0.8	1.6	2.4	3.9	10.1	28.6	75.1	145.3	199.8
2001	21.6	0.6	0.9	0.8	1.7	2.5	4.1	10.5	29.8	75.8	146.8	202.1
2000	21.8	1.0	0.9	0.9	1.8	2.4	4.1	10.8	30.3	77.3	150.9	206.5
1999	21.9	0.7	0.9	0.8	1.8	2.4	4.3	10.8	32.3	78.8	149.5	201.4
Non-Hispanic 4/, male												
2006	22.7	*	0.9	0.9	1.9	2.5	4.3	10.7	31.5	83.9	187.1	250.7
2005	22.8	*	1.0	0.9	1.7	2.3	4.1	10.5	31.6	88.2	186.3	268.5
2004	22.6	*	0.9	0.8	1.7	2.4	4.2	11.1	32.7	89.3	184.8	261.1
2003	23.3	*	0.8	1.0	1.6	2.7	4.8	11.6	34.6	92.6	188.2	273.6
2002	23.6	*	0.8	1.0	1.8	2.8	4.7	12.1	35.6	97.0	190.9	272.8
2001	23.5	*	1.0	1.0	1.9	2.9	5.0	12.9	36.8	96.3	188.6	278.8
2000	23.7	*	1.0	1.0	2.2	2.7	4.8	13.1	36.4	98.1	195.2	285.0
1999	23.7	*	0.9	0.9	2.2	2.9	5.2	13.3	39.5	98.7	191.5	284.3
Non-Hispanic 4/, female												
2006	18.1	*	0.8	0.6	1.0	1.6	2.9	6.7	18.6	51.2	108.5	162.0
2005	18.2	*	0.6	0.6	1.1	1.8	2.9	7.0	19.6	51.0	110.3	159.5
2004	18.5	*	0.6	0.6	1.2	1.7	2.8	7.2	20.1	52.9	113.3	162.6
2003	18.9	*	0.7	0.8	1.1	1.7	3.2	7.1	20.4	54.3	116.8	168.4
2002	19.2	*	0.7	0.7	1.4	2.0	3.1	8.0	22.2	56.8	115.1	168.3
2001	19.7	*	0.9	0.7	1.4	2.2	3.3	8.2	23.3	58.9	119.3	170.2
2000	20.1	*	0.9	0.7	1.3	2.0	3.4	8.6	24.7	60.1	122.1	174.8
1999	20.1	*	0.8	0.7	1.4	2.0	3.4	8.5	25.6	62.5	122.3	168.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Non-Hispanic white, both sexes												
2006 5/	22.6	*	0.9	0.7	1.4	2.0	3.4	8.4	24.5	67.6	145.3	196.6
2005 5/	22.8	*	0.7	0.8	1.4	1.8	3.4	8.3	25.3	69.4	146.2	201.5
2004 5/	22.7	*	0.8	0.7	1.4	2.0	3.4	8.8	26.1	70.4	145.9	198.8
2003 5/	23.4	1.0	0.8	0.9	1.4	2.1	3.8	9.0	27.4	73.2	149.3	207.2
2002	23.7	*	0.8	0.8	1.7	2.3	3.8	9.7	28.4	76.7	149.5	206.5
2001	23.8	*	1.1	0.8	1.7	2.4	4.0	10.3	29.5	77.2	150.5	208.2
2000	24.2	1.3	1.0	0.9	1.7	2.3	4.0	10.5	30.3	79.0	155.1	210.7
1999	24.1	*	1.0	0.8	1.8	2.4	4.2	10.6	32.0	80.4	153.0	206.4
Non-Hispanic white, male												
2006 5/	25.2	*	0.9	0.8	1.9	2.5	4.0	10.3	31.0	85.5	193.4	258.6
2005 5/	25.4	*	0.8	0.9	1.7	2.0	4.0	9.9	31.5	90.1	192.5	281.9
2004 5/	25.1	*	0.9	0.9	1.6	2.4	4.1	10.7	32.8	90.3	190.1	270.9
2003 5/	26.0	*	0.9	1.0	1.6	2.6	4.7	11.2	34.9	95.2	194.8	284.6
2002	26.2	*	0.9	1.0	1.9	2.7	4.5	11.8	35.4	99.3	196.8	283.5
2001	25.9	*	1.0	0.8	2.0	2.8	4.7	12.8	36.4	97.9	193.6	288.7
2000	26.2	*	1.1	1.0	2.1	2.6	4.8	12.7	36.3	100.1	201.0	293.8
1999	26.1	*	1.0	0.9	2.2	2.7	5.0	13.1	39.2	100.7	196.1	291.8
Non-Hispanic white, female												
2006 5/	20.2	*	0.9	0.6	1.0	1.6	2.8	6.5	18.2	52.1	111.9	167.7
2005 5/	20.2	*	0.6	0.6	1.1	1.6	2.7	6.6	19.4	51.4	114.3	164.7
2004 5/	20.4	*	0.6	0.6	1.2	1.5	2.6	7.0	19.8	53.4	115.8	166.5
2003 5/	20.8	*	0.7	0.8	1.1	1.6	3.0	6.8	20.3	54.3	118.5	173.1
2002	21.2	*	0.7	0.7	1.4	1.9	3.1	7.6	21.7	57.5	117.8	173.2
2001	21.8	*	1.1	0.7	1.4	2.0	3.2	7.9	23.0	59.5	122.0	174.6
2000	22.3	*	0.9	0.8	1.2	1.9	3.2	8.4	24.6	61.2	125.0	177.2
1999	22.2	*	0.9	0.6	1.3	2.0	3.3	8.3	25.2	63.4	124.7	172.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Non-Hispanic black, both sexes												
2006 5/	13.8	*	*	0.8	1.6	2.5	5.3	12.4	33.5	69.9	118.7	145.5
2005 5/	13.4	*	1.0	0.7	1.6	3.4	4.7	12.6	31.0	69.8	114.7	139.3
2004 5/	14.0	*	*	0.7	1.7	3.1	4.6	12.8	32.9	76.0	125.2	156.3
2003 5/	13.9	*	*	0.9	1.5	3.0	5.4	13.4	31.8	74.3	127.4	152.3
2002	14.1	*	*	0.8	1.5	3.0	4.8	13.9	36.4	77.0	123.5	153.7
2001	14.5	*	*	1.0	1.8	3.5	5.8	13.5	37.1	79.0	127.6	157.3
2000	14.3	*	*	0.7	2.3	3.0	5.4	15.0	36.5	76.2	122.3	177.1
1999	14.5	*	*	0.9	2.4	3.0	5.6	14.1	39.5	77.5	128.2	162.3
Non-Hispanic black, male												
2006 5/	15.5	*	*	1.0	2.0	2.9	6.7	15.8	44.0	92.6	156.0	198.8
2005 5/	14.6	*	*	1.0	1.9	3.9	5.5	16.0	39.3	90.0	154.5	168.4
2004 5/	14.8	*	*	0.7	2.0	3.4	5.4	16.0	40.3	99.5	152.9	197.1
2003 5/	14.5	*	*	0.9	1.7	3.5	6.6	17.1	41.1	88.6	150.3	188.6
2002	15.1	*	*	1.0	1.5	3.8	6.2	16.5	45.2	99.4	155.3	203.5
2001	15.6	*	*	1.3	2.0	4.0	7.6	16.3	46.6	100.7	161.4	206.7
2000	15.2	*	*	0.8	2.8	3.7	5.7	18.2	45.3	97.8	157.4	226.6
1999	15.6	*	*	1.2	2.6	3.9	7.0	17.5	48.4	96.9	162.9	221.1
Non-Hispanic black, female												
2006 5/	12.3	*	*	*	1.3	2.1	3.9	9.5	25.1	53.6	97.5	123.7
2005 5/	12.2	*	*	*	1.3	2.9	4.0	9.7	24.4	55.4	92.2	127.5
2004 5/	13.2	*	*	0.7	1.4	2.8	3.9	10.0	27.0	59.2	109.6	140.1
2003 5/	13.3	*	*	0.8	1.2	2.5	4.4	10.2	24.3	64.1	114.4	138.3
2002	13.1	*	*	*	1.5	2.3	3.5	11.6	29.4	61.2	105.4	134.6
2001	13.5	*	*	*	1.7	3.1	4.2	11.1	29.5	63.7	108.7	139.0
2000	13.5	*	*	*	1.8	2.5	5.2	12.2	29.4	61.0	102.6	158.7
1999	13.6	*	*	0.7	2.1	2.2	4.3	11.1	32.3	63.8	108.7	140.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
All origins 3/, both sexes												
2006	0.4	*	*	*	0.2	0.3	0.4	0.3	0.6	1.1	2.2	2.6
2005	0.4	*	*	*	0.1	0.3	0.3	0.4	0.6	1.1	2.3	2.1
2004	0.4	*	*	*	0.2	0.3	0.3	0.4	0.7	1.2	1.9	2.2
2003	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.2	2.2	2.4
2002	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.7
2001	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.2	2.2	2.2
2000	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.1	2.5
1999	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.5	2.1	2.4
All origins 3/, male												
2006	0.5	*	*	*	0.2	0.4	0.5	0.5	0.8	1.4	2.6	3.7
2005	0.5	*	*	*	0.1	0.3	0.3	0.4	0.8	1.4	2.5	3.1
2004	0.5	*	*	*	0.2	0.3	0.3	0.5	0.9	1.6	2.5	3.2
2003	0.5	*	*	*	0.1	0.4	0.4	0.5	0.9	1.5	2.6	2.6
2002	0.5	*	*	*	0.2	0.4	0.4	0.5	0.8	1.8	2.5	3.6
2001	0.5	*	*	*	0.2	0.4	0.5	0.6	0.7	1.6	2.6	2.6
2000	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.5	3.7
1999	0.6	*	*	*	0.3	0.4	0.5	0.5	1.0	1.9	2.5	3.8
All origins 3/, female												
2006	0.4	*	*	*	0.1	0.3	0.2	0.2	0.4	0.8	2.0	2.1
2005	0.4	*	*	*	0.2	0.3	0.2	0.3	0.4	0.9	2.2	1.6
2004	0.4	*	*	*	0.1	0.3	0.2	0.3	0.5	0.9	1.5	1.7
2003	0.4	*	*	*	0.2	0.4	0.3	0.3	0.5	0.9	1.9	2.3
2002	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.9	1.6	2.3
2001	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.8	2.0	2.0
2000	0.4	*	*	*	0.2	0.3	0.3	0.3	0.6	1.0	1.8	2.0
1999	0.4	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
Hispanic, both sexes												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.3	*	*	*	*	*	*	*	*	2.2	3.3	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	1.8	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	4.5	*
1999	0.3	*	*	*	*	*	*	*	*	2.5	*	*
Hispanic, male												
2006	0.3	*	*	*	*	*	*	*	*	*	*	*
2005	0.4	*	*	*	*	*	*	*	*	*	*	*
2004	0.3	*	*	*	*	*	*	*	*	*	*	*
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.2	*	*	*	*	*	*	*	*	*	*	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
Non-Hispanic 4/, both sexes												
2006	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.1	2.2	2.6
2005	0.4	*	*	*	0.2	0.3	0.3	0.4	0.6	1.0	2.3	2.0
2004	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.2	1.9	2.1
2003	0.5	*	*	*	0.2	0.5	0.4	0.4	0.7	1.2	2.2	2.3
2002	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.8
2001	0.5	*	*	*	0.2	0.5	0.4	0.5	0.5	1.1	2.2	2.2
2000	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.4
1999	0.5	*	*	*	0.3	0.4	0.4	0.4	0.7	1.5	2.1	2.3
Non-Hispanic 4/, male												
2006	0.6	*	*	*	0.2	0.5	0.5	0.5	0.8	1.3	2.5	3.8
2005	0.5	*	*	*	0.1	0.4	0.3	0.4	0.8	1.3	2.4	3.0
2004	0.5	*	*	*	0.2	0.4	0.3	0.5	1.0	1.6	2.5	3.1
2003	0.5	*	*	*	0.1	0.5	0.5	0.5	0.9	1.5	2.5	2.7
2002	0.6	*	*	*	0.2	0.4	0.4	0.5	0.8	1.8	2.4	3.7
2001	0.5	*	*	*	0.2	0.5	0.5	0.6	0.7	1.5	2.6	2.5
2000	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.3	3.7
1999	0.6	*	*	*	0.3	0.5	0.5	0.5	1.0	1.9	2.3	3.7
Non-Hispanic 4/, female												
2006	0.4	*	*	*	0.1	0.3	0.3	0.2	0.4	0.8	2.0	2.1
2005	0.4	*	*	*	0.2	0.3	0.2	0.4	0.4	0.8	2.2	1.6
2004	0.4	*	*	*	0.2	0.4	0.3	0.3	0.5	0.9	1.5	1.6
2003	0.5	*	*	*	0.2	0.4	0.3	0.3	0.6	0.9	1.9	2.2
2002	0.4	*	*	*	0.2	0.4	0.3	0.4	0.5	0.8	1.5	2.4
2001	0.4	*	*	*	0.2	0.4	0.3	0.4	0.5	0.8	2.0	2.0
2000	0.4	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	1.7	2.0
1999	0.5	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
Non-Hispanic white, both sexes												
2006 5/	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.1	2.3	2.7
2005 5/	0.5	*	*	*	0.2	0.3	0.3	0.4	0.6	1.1	2.4	2.1
2004 5/	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.2	2.0	2.1
2003 5/	0.6	*	*	*	0.2	0.5	0.4	0.4	0.7	1.2	2.3	2.5
2002	0.6	*	*	*	0.2	0.5	0.4	0.5	0.6	1.4	2.0	2.9
2001	0.5	*	*	*	0.2	0.5	0.4	0.5	0.6	1.2	2.3	2.3
2000	0.5	*	*	*	0.2	0.5	0.4	0.4	0.6	1.3	2.1	2.6
1999	0.6	*	*	*	0.3	0.4	0.4	0.5	0.8	1.5	2.2	2.3
Non-Hispanic white, male												
2006 5/	0.6	*	*	*	0.2	0.5	0.5	0.5	0.9	1.4	2.6	3.9
2005 5/	0.5	*	*	*	0.1	0.3	0.3	0.4	0.8	1.3	2.5	3.2
2004 5/	0.6	*	*	*	0.2	0.4	0.3	0.5	1.0	1.6	2.6	3.3
2003 5/	0.6	*	*	*	*	0.5	0.5	0.5	0.9	1.5	2.6	2.7
2002	0.6	*	*	*	0.2	0.5	0.4	0.5	0.8	1.9	2.6	4.0
2001	0.6	*	*	*	0.3	0.5	0.5	0.6	0.7	1.6	2.7	2.6
2000	0.6	*	*	*	0.2	0.5	0.4	0.5	0.7	1.6	2.5	3.9
1999	0.7	*	*	*	0.3	0.5	0.5	0.5	1.1	2.0	2.5	3.8
Non-Hispanic white, female												
2006 5/	0.4	*	*	*	*	0.3	0.3	0.3	0.4	0.9	2.1	2.1
2005 5/	0.5	*	*	*	0.2	0.2	0.2	0.4	0.5	0.9	2.4	1.5
2004 5/	0.4	*	*	*	0.2	0.3	0.3	0.3	0.5	0.9	1.6	1.6
2003 5/	0.5	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	2.0	2.4
2002	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.9	1.6	2.5
2001	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.9	2.0	2.3
2000	0.5	*	*	*	0.2	0.4	0.3	0.4	0.6	1.1	1.9	2.1
1999	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	1.2	2.0	1.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
Non-Hispanic black, both sexes												
2006 5/	0.4	*	*	*	*	0.4	0.4	0.5	*	*	*	*
2005 5/	0.4	*	*	*	*	0.7	0.5	0.6	*	*	*	*
2004 5/	0.4	*	*	*	*	0.6	*	0.4	*	*	*	*
2003 5/	0.3	*	*	*	*	0.6	*	*	0.8	*	*	*
2002	0.3	*	*	*	*	0.4	0.4	0.4	0.8	*	*	*
2001	0.4	*	*	*	*	0.8	0.4	*	*	*	*	*
2000	0.3	*	*	*	*	*	0.4	*	*	*	*	*
1999	0.4	*	*	*	*	0.5	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.5	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.5	*	*	*	*	0.9	*	*	*	*	*	*
2004 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.5	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	0.8	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
All origins 3/, both sexes												
2006	6.9	*	*	0.1	0.3	0.5	1.2	3.3	9.1	24.0	53.7	72.3
2005	7.0	*	*	0.1	0.3	0.6	1.3	3.3	9.6	24.8	54.5	75.2
2004	7.1	*	*	0.1	0.3	0.6	1.2	3.7	9.9	26.1	54.6	75.0
2003	7.4	*	*	0.1	0.3	0.6	1.4	3.8	10.8	26.8	56.1	78.8
2002	7.6	*	*	0.1	0.3	0.7	1.5	3.9	11.5	28.8	57.2	77.3
2001	7.8	*	*	0.1	0.4	0.8	1.7	4.4	12.2	29.3	58.6	79.6
2000	8.1	*	*	0.1	0.5	0.7	1.7	4.6	13.1	30.4	61.0	80.2
1999	8.2	*	*	0.1	0.4	0.7	1.8	4.8	13.3	31.7	61.8	77.7
All origins 3/, male												
2006	7.4	*	*	0.1	0.4	0.7	1.5	4.3	11.8	29.7	66.9	89.6
2005	7.6	*	*	0.2	0.4	0.7	1.7	4.4	11.9	31.4	67.4	100.5
2004	7.6	*	*	0.1	0.4	0.8	1.7	4.8	12.3	32.5	67.3	95.8
2003	8.0	*	*	0.1	0.4	0.8	1.8	4.8	13.8	33.9	69.1	103.6
2002	8.1	*	*	0.2	0.4	0.9	2.0	5.0	14.4	36.0	70.7	99.7
2001	8.4	*	*	0.1	0.5	0.9	2.2	5.8	15.0	36.5	71.3	106.6
2000	8.6	*	*	0.1	0.6	0.9	2.1	6.0	15.9	37.4	74.5	105.9
1999	8.6	*	*	0.1	0.5	0.9	2.3	6.2	16.2	38.3	74.7	104.4
All origins 3/, female												
2006	6.4	*	*	*	0.2	0.4	0.9	2.3	6.7	19.2	44.7	64.2
2005	6.5	*	*	*	0.2	0.4	0.9	2.3	7.5	19.3	45.7	63.5
2004	6.6	*	*	*	0.3	0.5	0.8	2.6	7.6	20.7	46.0	65.7
2003	6.8	*	*	*	0.3	0.4	1.0	2.8	8.0	20.9	47.4	67.8
2002	7.1	*	*	*	0.2	0.5	1.1	3.0	8.8	22.8	48.2	67.6
2001	7.3	*	*	*	0.3	0.6	1.1	3.1	9.7	23.3	50.2	68.3
2000	7.6	*	*	*	0.3	0.6	1.2	3.3	10.5	24.7	52.1	69.8
1999	7.7	*	*	*	0.3	0.6	1.2	3.4	10.6	26.4	53.4	66.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
Hispanic, both sexes												
2006	2.6	*	*	*	0.4	0.5	0.9	2.7	8.2	18.9	40.1	40.3
2005	2.6	*	*	*	0.3	0.4	1.0	2.9	7.7	19.5	44.7	44.5
2004	2.6	*	*	*	0.4	0.6	1.2	3.3	7.5	18.7	40.6	43.8
2003	2.6	*	*	*	0.4	0.6	1.4	3.0	8.4	19.1	40.4	54.4
2002	2.8	*	*	*	0.4	0.6	1.3	3.0	9.4	22.7	43.7	50.4
2001	2.7	*	*	*	0.3	0.5	1.4	3.6	10.9	18.3	43.3	59.4
2000	2.9	*	*	*	0.3	0.6	1.6	3.6	11.1	25.7	41.5	59.7
1999	2.8	*	*	*	*	0.6	1.4	4.4	9.3	25.2	42.7	56.1
Hispanic, male												
2006	2.7	*	*	*	0.6	0.5	1.2	3.1	10.2	22.1	47.7	54.1
2005	2.7	*	*	*	*	0.5	1.2	3.2	8.9	23.9	52.8	52.3
2004	2.8	*	*	*	*	0.7	1.8	4.9	8.8	24.0	47.2	50.4
2003	2.8	*	*	*	*	0.8	1.8	3.4	10.5	23.8	45.3	57.0
2002	2.7	*	*	*	*	0.6	1.5	3.6	11.7	26.0	45.5	60.2
2001	3.0	*	*	*	*	0.6	2.0	4.4	13.2	24.4	58.6	68.6
2000	3.1	*	*	*	*	0.7	1.9	4.6	13.9	30.2	48.2	82.6
1999	2.8	*	*	*	*	*	1.7	5.7	11.8	27.5	48.0	60.4
Hispanic, female												
2006	2.4	*	*	*	*	*	0.6	2.2	6.4	16.3	34.7	33.0
2005	2.5	*	*	*	*	*	0.8	2.5	6.5	15.9	39.1	40.4
2004	2.3	*	*	*	*	*	*	1.6	6.4	14.5	36.1	40.3
2003	2.5	*	*	*	*	*	0.9	2.7	6.6	15.3	37.1	53.1
2002	2.8	*	*	*	*	*	1.2	2.3	7.4	20.1	42.4	45.6
2001	2.4	*	*	*	*	*	0.8	2.8	8.8	13.5	32.9	54.9
2000	2.7	*	*	*	*	*	1.3	2.6	8.6	22.2	37.0	48.5
1999	2.8	*	*	*	*	*	1.0	3.1	7.1	23.4	39.2	54.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
Non-Hispanic 4/, both sexes												
2006	7.6	*	*	0.1	0.3	0.6	1.2	3.3	9.2	24.4	54.5	73.9
2005	7.8	*	*	0.1	0.3	0.6	1.3	3.4	9.8	25.2	55.0	76.6
2004	7.9	*	*	0.1	0.3	0.6	1.2	3.7	10.0	26.6	55.3	76.3
2003	8.1	*	*	0.1	0.3	0.6	1.4	3.9	10.9	27.3	56.8	79.6
2002	8.3	*	*	0.1	0.3	0.7	1.5	4.0	11.6	29.2	57.7	78.2
2001	8.6	*	*	0.1	0.4	0.8	1.7	4.5	12.3	29.9	59.2	80.3
2000	8.8	*	*	0.1	0.5	0.8	1.7	4.7	13.2	30.7	61.7	80.9
1999	8.9	*	*	0.1	0.4	0.7	1.8	4.8	13.5	32.1	62.4	78.3
Non-Hispanic 4/, male												
2006	8.3	*	*	0.1	0.4	0.7	1.5	4.4	11.9	30.3	68.1	91.5
2005	8.5	*	*	0.2	0.4	0.7	1.8	4.5	12.2	31.9	68.2	103.0
2004	8.5	*	*	0.1	0.4	0.8	1.6	4.8	12.6	33.1	68.4	97.8
2003	8.8	*	*	0.1	0.4	0.8	1.8	5.0	14.0	34.5	70.3	105.6
2002	9.0	*	*	0.2	0.4	0.9	2.0	5.1	14.6	36.5	71.8	101.4
2001	9.2	*	*	0.1	0.5	0.9	2.3	5.9	15.1	37.2	71.7	108.1
2000	9.4	*	*	0.2	0.7	0.9	2.2	6.1	15.9	37.7	75.6	106.7
1999	9.4	*	*	0.2	0.5	0.9	2.3	6.2	16.5	38.8	75.5	106.0
Non-Hispanic 4/, female												
2006	7.0	*	*	*	0.2	0.4	0.9	2.3	6.7	19.4	45.3	65.7
2005	7.1	*	*	*	0.2	0.5	0.9	2.3	7.6	19.5	46.0	64.6
2004	7.3	*	*	*	0.3	0.4	0.8	2.7	7.7	21.2	46.5	66.7
2003	7.4	*	*	*	0.3	0.5	1.0	2.8	8.1	21.3	47.8	68.2
2002	7.7	*	*	*	0.3	0.5	1.1	3.0	8.9	23.0	48.3	68.2
2001	8.0	*	*	*	0.3	0.6	1.2	3.1	9.7	23.8	50.9	68.7
2000	8.3	*	*	*	0.3	0.6	1.2	3.4	10.6	24.8	52.6	70.4
1999	8.4	*	*	*	0.3	0.5	1.3	3.4	10.8	26.5	53.9	67.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
Non-Hispanic white, both sexes												
2006 5/	8.7	*	*	0.1	0.2	0.5	1.1	3.3	9.3	25.8	57.9	78.2
2005 5/	8.9	*	*	0.1	0.3	0.5	1.3	3.3	10.0	26.6	58.4	81.3
2004 5/	9.0	*	*	0.1	0.3	0.6	1.2	3.7	10.4	28.1	58.5	80.2
2003 5/	9.3	*	*	0.1	0.3	0.5	1.4	3.8	11.4	28.8	60.0	84.3
2002	9.5	*	*	0.2	0.3	0.7	1.5	4.0	12.0	30.6	60.9	83.1
2001	9.7	*	*	0.1	0.3	0.7	1.7	4.5	12.7	31.4	62.3	84.1
2000	10.1	*	*	0.1	0.4	0.7	1.7	4.8	13.8	32.3	65.1	85.1
1999	10.1	*	*	0.1	0.4	0.7	1.7	4.9	14.1	33.9	65.6	82.0
Non-Hispanic white, male												
2006 5/	9.4	*	*	*	0.3	0.6	1.3	4.3	12.0	31.8	72.2	96.2
2005 5/	9.7	*	*	0.2	0.4	0.6	1.7	4.3	12.4	33.6	72.4	109.8
2004 5/	9.6	*	*	*	0.4	0.8	1.7	4.7	12.9	34.6	72.0	103.5
2003 5/	10.1	*	*	*	0.4	0.7	1.8	4.9	14.6	36.6	74.4	112.0
2002	10.2	*	*	0.2	0.4	0.8	1.9	5.1	15.0	38.3	75.6	107.6
2001	10.4	*	*	*	0.4	0.9	2.1	6.0	15.5	38.8	75.2	113.1
2000	10.6	*	*	0.2	0.6	0.8	2.2	6.1	16.5	39.5	79.6	113.0
1999	10.6	*	*	0.2	0.5	0.8	2.2	6.3	17.0	40.8	78.9	110.0
Non-Hispanic white, female												
2006 5/	8.1	*	*	*	*	0.4	0.8	2.3	6.8	20.5	48.0	69.8
2005 5/	8.2	*	*	*	*	0.4	0.8	2.3	7.8	20.6	48.8	68.3
2004 5/	8.4	*	*	*	0.3	0.4	0.8	2.7	8.0	22.5	49.2	69.8
2003 5/	8.5	*	*	*	0.3	0.4	1.0	2.8	8.4	22.1	50.3	72.2
2002	8.8	*	*	*	0.2	0.5	1.1	2.9	9.2	24.1	51.0	72.6
2001	9.2	*	*	*	0.3	0.5	1.2	3.1	10.1	25.2	53.7	72.0
2000	9.5	*	*	*	0.3	0.6	1.1	3.4	11.3	26.3	55.6	73.8
1999	9.6	*	*	*	0.2	0.5	1.2	3.5	11.4	28.1	56.9	70.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
Non-Hispanic black, both sexes												
2006 5/	3.9	*	*	*	0.5	0.9	2.2	4.6	10.2	17.3	28.3	34.5
2005 5/	3.7	*	*	*	0.4	1.1	2.0	4.4	9.2	17.2	27.0	34.4
2004 5/	3.8	*	*	*	0.5	1.1	1.5	4.8	9.1	18.4	28.2	39.4
2003 5/	3.7	*	*	*	0.4	1.1	2.0	4.9	9.1	18.4	28.6	31.2
2002	4.0	*	*	*	0.4	1.2	2.0	4.8	10.8	21.6	30.1	31.2
2001	4.1	*	*	*	0.6	1.2	2.3	5.1	10.9	21.3	30.6	40.0
2000	3.9	*	*	*	0.6	1.1	2.1	5.0	10.8	19.7	27.1	36.6
1999	4.1	*	*	*	0.6	1.1	2.4	4.8	11.2	20.5	32.1	40.4
Non-Hispanic black, male												
2006 5/	4.4	*	*	*	*	1.0	2.9	6.6	13.9	22.1	32.8	43.1
2005 5/	4.2	*	*	*	*	1.3	2.3	6.3	12.3	23.1	31.1	40.9
2004 5/	4.1	*	*	*	0.6	1.4	1.9	6.6	12.0	23.5	33.4	42.7
2003 5/	3.9	*	*	*	*	1.4	2.4	6.7	11.5	20.7	31.2	35.2
2002	4.5	*	*	*	*	1.6	2.9	5.7	13.9	28.3	35.8	43.8
2001	4.7	*	*	*	0.8	1.3	3.4	6.7	13.7	28.2	37.2	55.4
2000	4.3	*	*	*	*	1.2	2.6	7.0	14.8	25.3	31.8	43.4
1999	4.7	*	*	*	*	1.6	3.3	6.6	14.8	26.2	41.0	59.4
Non-Hispanic black, female												
2006 5/	3.4	*	*	*	*	0.7	1.6	2.8	7.2	13.9	25.7	31.0
2005 5/	3.3	*	*	*	*	0.8	1.7	2.7	6.7	13.0	24.6	31.8
2004 5/	3.4	*	*	*	*	0.8	1.2	3.4	6.9	14.8	25.3	38.1
2003 5/	3.5	*	*	*	*	0.9	1.5	3.4	7.1	16.9	27.2	29.7
2002	3.6	*	*	*	*	0.8	1.2	4.0	8.3	16.9	26.9	26.3
2001	3.7	*	*	*	*	1.1	1.4	3.7	8.6	16.5	27.0	34.2
2000	3.5	*	*	*	*	0.9	1.7	3.3	7.5	15.7	24.5	34.0
1999	3.6	*	*	*	*	*	1.7	3.2	8.2	16.5	27.1	33.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
All origins 3/, both sexes												
2006	7.3	0.8	0.8	0.6	1.1	1.1	1.6	3.2	8.9	25.0	52.6	76.9
2005	7.3	0.5	0.7	0.7	1.1	1.1	1.6	3.3	8.8	25.1	52.3	78.7
2004	7.3	0.6	0.8	0.8	1.0	1.0	1.6	3.3	9.2	25.4	52.4	78.8
2003	7.4	0.7	0.8	0.8	1.0	1.1	1.8	3.3	9.3	25.7	53.9	80.3
2002	7.5	0.5	0.7	0.8	1.2	1.2	1.7	3.6	9.8	26.4	53.1	80.5
2001	7.5	0.6	1.0	0.8	1.2	1.3	1.7	3.8	10.0	26.7	53.3	81.4
2000	7.6	1.0	0.9	0.8	1.2	1.2	1.8	3.7	9.9	27.2	54.5	83.6
1999	7.5	0.7	0.8	0.7	1.2	1.2	1.8	3.7	10.6	27.2	52.8	82.4
All origins 3/, male												
2006	8.4	*	0.8	0.7	1.4	1.3	1.8	3.7	11.4	33.1	75.9	105.1
2005	8.4	*	1.0	0.8	1.4	1.3	1.6	3.6	11.3	34.0	75.0	112.4
2004	8.3	*	0.9	0.9	1.3	1.2	1.8	3.8	11.9	34.4	73.2	110.7
2003	8.4	*	0.9	0.9	1.2	1.4	2.0	4.0	11.9	35.1	75.3	111.8
2002	8.5	*	0.7	0.8	1.4	1.4	1.9	4.2	12.2	36.3	74.9	114.5
2001	8.5	*	1.0	0.9	1.4	1.5	1.9	4.2	12.8	35.7	74.2	117.5
2000	8.5	1.1	0.9	0.9	1.5	1.4	1.9	4.3	12.5	37.2	75.7	119.8
1999	8.4	*	0.8	0.8	1.5	1.4	1.9	4.3	13.1	36.8	73.2	119.2
All origins 3/, female												
2006	6.3	*	0.8	0.6	0.8	0.9	1.5	2.7	6.7	18.2	36.7	63.7
2005	6.2	*	0.5	0.6	0.8	1.0	1.5	3.0	6.4	17.5	37.0	63.3
2004	6.3	*	0.6	0.7	0.8	0.9	1.4	2.8	6.8	17.8	38.4	64.5
2003	6.4	1.1	0.8	0.7	0.7	0.8	1.6	2.7	7.0	17.8	39.6	66.4
2002	6.5	*	0.6	0.7	1.0	1.0	1.5	3.1	7.6	18.1	38.7	65.7
2001	6.6	*	1.0	0.7	1.0	1.1	1.5	3.3	7.5	19.2	39.5	66.3
2000	6.7	*	0.9	0.6	0.9	1.0	1.7	3.2	7.6	19.0	40.6	68.8
1999	6.7	*	0.7	0.7	1.0	1.1	1.6	3.2	8.3	19.3	39.6	67.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
Hispanic, both sexes												
2006	2.8	*	0.7	0.7	1.5	1.3	1.7	2.7	5.9	16.9	29.4	35.2
2005	3.0	*	0.7	1.0	1.7	1.3	1.5	3.2	7.0	17.6	28.2	40.6
2004	2.8	*	0.8	1.3	1.5	1.1	1.5	3.0	6.3	15.7	28.6	36.2
2003	2.9	*	1.2	1.1	1.4	1.1	1.7	3.6	6.3	13.9	32.9	47.2
2002	2.8	*	0.8	1.1	1.6	1.1	2.1	2.5	7.2	15.5	27.4	47.1
2001	3.0	*	1.3	1.1	1.7	1.5	1.9	2.9	6.5	17.3	33.4	42.5
2000	2.9	*	0.9	1.1	1.5	1.1	1.9	3.5	7.8	16.9	32.6	49.1
1999	2.9	*	0.9	1.1	1.6	1.4	2.0	3.4	7.0	17.3	31.0	44.2
Hispanic, male												
2006	3.0	*	*	0.7	1.8	1.4	1.6	3.0	7.1	20.9	40.5	42.9
2005	3.4	*	*	1.1	2.2	1.5	1.7	3.4	8.4	22.9	41.8	57.3
2004	3.1	*	*	1.6	2.0	1.3	1.8	3.3	8.1	20.2	37.9	43.4
2003	3.2	*	1.4	1.3	1.7	1.3	1.9	4.1	7.6	18.3	40.6	61.6
2002	3.0	*	*	1.1	2.0	1.4	2.3	2.8	7.2	18.5	38.6	65.3
2001	3.4	*	1.4	1.1	2.2	1.8	2.3	3.3	8.4	19.6	45.6	68.6
2000	3.3	*	*	1.3	1.7	1.5	2.0	4.1	10.2	20.7	42.8	76.6
1999	3.1	*	*	1.1	1.7	1.4	1.9	3.6	8.6	23.5	39.1	49.6
Hispanic, female												
2006	2.6	*	*	0.7	1.2	1.2	1.7	2.5	4.8	13.7	21.7	31.2
2005	2.5	*	*	0.9	1.2	1.1	1.3	3.0	5.7	13.4	18.8	31.9
2004	2.4	*	*	1.0	0.9	0.9	1.3	2.6	4.6	12.1	22.2	32.4
2003	2.6	*	*	1.0	1.0	0.9	1.5	3.2	5.2	10.4	27.7	39.8
2002	2.6	*	*	1.2	1.1	0.7	1.9	2.3	7.3	13.1	19.7	38.1
2001	2.6	*	*	1.1	1.1	1.1	1.4	2.6	4.8	15.6	25.1	29.8
2000	2.6	*	*	0.8	1.3	*	1.8	2.9	5.6	13.9	25.8	35.6
1999	2.8	*	*	1.0	1.5	1.3	2.1	3.2	5.6	12.4	25.7	41.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
Non-Hispanic 4/, both sexes												
2006	8.1	0.7	0.8	0.6	1.0	1.1	1.6	3.3	9.2	25.6	54.0	78.9
2005	8.0	*	0.8	0.6	1.0	1.1	1.5	3.3	8.9	25.6	53.7	80.6
2004	8.0	*	0.8	0.6	0.9	1.0	1.6	3.3	9.5	26.1	53.7	80.6
2003	8.1	0.8	0.7	0.8	0.9	1.1	1.8	3.3	9.6	26.4	54.9	81.6
2002	8.2	*	0.6	0.7	1.1	1.2	1.6	3.7	10.0	27.0	54.3	81.7
2001	8.2	*	0.9	0.7	1.1	1.2	1.7	3.8	10.3	27.2	54.1	82.7
2000	8.2	1.0	0.9	0.7	1.1	1.2	1.7	3.8	10.1	27.8	55.3	84.7
1999	8.2	*	0.7	0.7	1.2	1.2	1.7	3.8	10.8	27.8	53.6	83.7
Non-Hispanic 4/, male												
2006	9.4	*	0.8	0.7	1.3	1.3	1.8	3.8	11.7	34.0	78.1	108.3
2005	9.3	*	1.0	0.7	1.2	1.2	1.6	3.7	11.5	34.8	76.9	115.3
2004	9.2	*	0.9	0.7	1.1	1.2	1.8	3.9	12.2	35.4	75.1	113.9
2003	9.3	*	0.8	0.8	1.1	1.4	2.0	4.0	12.2	36.2	77.0	113.7
2002	9.4	*	0.7	0.8	1.2	1.4	1.9	4.3	12.6	37.3	76.5	116.5
2001	9.3	*	0.9	0.8	1.2	1.4	1.9	4.3	13.1	36.6	75.4	119.3
2000	9.3	*	0.9	0.8	1.4	1.4	1.9	4.3	12.6	38.2	77.0	121.5
1999	9.2	*	0.8	0.8	1.4	1.3	1.9	4.4	13.4	37.5	74.6	122.1
Non-Hispanic 4/, female												
2006	6.9	*	0.8	0.5	0.7	0.9	1.4	2.8	6.8	18.6	37.6	65.2
2005	6.8	*	0.5	0.5	0.8	1.0	1.5	3.0	6.5	17.8	38.0	64.8
2004	6.9	*	0.6	0.6	0.8	0.9	1.4	2.8	7.0	18.2	39.3	65.7
2003	7.0	*	0.7	0.7	0.7	0.8	1.6	2.7	7.1	18.3	40.2	67.4
2002	7.0	*	0.5	0.6	0.9	1.0	1.4	3.2	7.6	18.4	39.5	66.6
2001	7.2	*	0.9	0.6	0.9	1.1	1.5	3.4	7.7	19.4	40.1	67.5
2000	7.2	*	0.9	0.6	0.8	1.0	1.6	3.3	7.7	19.3	41.1	69.8
1999	7.2	*	0.7	0.6	0.9	1.1	1.5	3.2	8.5	19.8	40.0	68.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
Non-Hispanic white, both sexes												
2006 5/	9.2	*	0.9	0.6	1.0	1.1	1.6	3.4	9.4	26.8	56.3	82.0
2005 5/	9.1	*	0.7	0.6	1.0	1.0	1.6	3.3	9.2	26.7	56.1	84.5
2004 5/	9.0	*	0.7	0.7	0.9	1.0	1.6	3.3	9.7	26.8	55.7	83.6
2003 5/	9.1	1.0	0.8	0.8	0.9	1.1	1.8	3.3	9.8	27.5	57.1	84.7
2002	9.1	*	0.7	0.7	1.1	1.2	1.7	3.7	10.0	28.2	56.2	84.4
2001	9.1	*	1.0	0.7	1.1	1.2	1.7	3.8	10.5	28.1	56.0	86.1
2000	9.2	1.3	0.9	0.7	1.1	1.1	1.7	3.7	10.2	29.0	57.5	87.3
1999	9.1	*	0.8	0.7	1.1	1.2	1.7	3.7	11.0	28.6	55.3	86.8
Non-Hispanic white, male												
2006 5/	10.6	*	0.8	0.7	1.3	1.4	1.8	3.9	12.0	35.3	81.0	112.1
2005 5/	10.5	*	0.8	0.7	1.1	1.1	1.6	3.6	11.9	36.3	79.7	121.9
2004 5/	10.3	*	0.8	0.7	1.1	1.2	1.7	3.8	12.5	36.1	78.0	118.8
2003 5/	10.5	*	0.8	0.9	1.1	1.3	1.9	4.0	12.5	37.7	80.6	119.0
2002	10.5	*	0.8	0.7	1.3	1.3	1.9	4.2	12.7	38.7	79.2	121.3
2001	10.3	*	1.0	0.7	1.3	1.4	1.8	4.3	13.3	37.7	78.0	125.3
2000	10.4	*	1.0	0.8	1.3	1.3	1.9	4.2	12.7	39.5	79.9	125.5
1999	10.2	*	0.8	0.7	1.4	1.3	1.9	4.3	13.6	38.6	77.0	127.2
Non-Hispanic white, female												
2006 5/	7.8	*	0.9	0.5	0.7	0.9	1.5	2.9	6.9	19.4	39.1	68.0
2005 5/	7.7	*	0.5	0.5	0.8	0.9	1.6	3.0	6.6	18.5	39.9	67.3
2004 5/	7.7	*	0.6	0.6	0.8	0.8	1.4	2.8	7.0	18.8	40.4	67.8
2003 5/	7.8	*	0.7	0.6	0.7	0.8	1.6	2.6	7.3	18.8	41.3	69.6
2002	7.8	*	0.5	0.6	0.9	1.0	1.5	3.2	7.5	19.3	40.7	68.4
2001	8.0	*	1.1	0.6	0.9	1.0	1.6	3.3	7.8	19.9	41.4	69.8
2000	8.1	*	0.9	0.6	0.8	0.9	1.5	3.2	7.8	20.1	42.7	72.0
1999	8.0	*	0.9	0.6	0.8	1.1	1.5	3.2	8.5	20.3	41.1	70.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
Non-Hispanic black, both sexes												
2006 5/	4.7	*	*	0.6	1.0	1.1	1.9	3.4	9.7	23.1	41.0	56.8
2005 5/	4.5	*	1.0	0.6	1.0	1.6	1.6	4.0	8.6	21.5	39.0	53.0
2004 5/	5.0	*	*	0.6	1.0	1.3	2.1	4.0	10.4	25.4	41.7	60.7
2003 5/	4.8	*	*	0.7	0.9	1.2	2.4	3.8	9.8	23.4	42.1	60.4
2002	4.8	*	*	0.6	0.9	1.3	1.6	4.4	11.5	22.6	42.9	65.3
2001	4.9	*	*	0.9	1.0	1.5	2.1	4.3	10.1	25.0	42.1	57.1
2000	4.9	*	*	0.6	1.4	1.7	2.1	5.0	10.8	23.1	38.8	66.1
1999	4.9	*	*	0.8	1.5	1.3	1.9	4.4	10.8	25.2	41.6	55.9
Non-Hispanic black, male												
2006 5/	5.5	*	*	0.9	1.2	1.2	2.5	3.9	12.8	32.5	60.6	87.0
2005 5/	5.1	*	*	0.7	1.3	1.6	1.9	4.5	10.8	29.2	59.1	63.8
2004 5/	5.5	*	*	0.7	1.1	1.4	2.3	4.9	12.8	37.3	52.8	81.3
2003 5/	5.2	*	*	0.7	1.2	1.6	2.8	4.6	12.6	31.5	54.0	78.8
2002	5.3	*	*	0.7	1.0	1.5	1.8	5.2	14.1	32.2	58.0	87.5
2001	5.4	*	*	1.2	1.1	1.8	2.6	4.6	13.4	34.4	56.1	75.1
2000	5.5	*	*	0.7	1.9	2.0	2.0	5.6	14.0	32.3	55.8	98.6
1999	5.4	*	*	1.0	1.6	1.5	2.1	5.7	13.7	33.4	55.7	78.4
Non-Hispanic black, female												
2006 5/	4.0	*	*	*	0.8	1.1	1.4	2.9	7.3	16.3	29.8	44.5
2005 5/	4.1	*	*	*	0.7	1.6	1.3	3.7	6.7	16.0	27.6	48.6
2004 5/	4.5	*	*	*	0.9	1.2	1.8	3.2	8.5	17.0	35.4	52.6
2003 5/	4.4	*	*	0.8	0.7	0.9	1.9	3.1	7.5	17.6	35.4	53.2
2002	4.4	*	*	*	0.8	1.1	1.4	3.6	9.4	15.8	34.3	56.9
2001	4.4	*	*	*	1.0	1.1	1.7	4.0	7.4	18.4	34.2	50.5
2000	4.4	*	*	*	0.9	1.4	2.2	4.4	8.3	16.5	29.2	54.1
1999	4.4	*	*	0.6	1.3	1.1	1.7	3.3	8.5	19.4	33.8	47.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
All origins 3/, both sexes												
2006	3.7	*	*	*	*	*	0.3	1.7	5.7	15.0	29.1	33.8
2005	3.8	*	*	*	*	*	0.3	1.6	6.1	15.9	29.5	33.6
2004	3.7	*	*	*	*	*	0.3	1.6	5.9	15.4	30.7	33.1
2003	3.9	*	*	*	*	*	0.3	1.7	5.9	16.5	31.1	36.3
2002	4.0	*	*	*	*	*	0.3	1.8	6.3	17.4	31.2	36.7
2001	3.9	*	*	*	*	*	0.3	1.7	6.6	17.4	30.9	36.3
2000	3.9	*	*	*	*	*	0.3	1.9	6.3	17.3	31.5	37.8
1999	3.9	*	*	*	*	0.1	0.3	1.8	7.1	17.2	31.0	36.6
All origins 3/, male												
2006	4.0	*	*	*	*	*	0.4	2.0	6.8	18.1	37.5	45.8
2005	4.1	*	*	*	*	*	0.4	2.0	7.0	19.7	38.0	46.1
2004	3.9	*	*	*	*	*	0.4	1.9	6.9	18.8	38.1	44.6
2003	4.1	*	*	*	*	*	0.4	2.1	7.4	19.8	37.7	50.4
2002	4.2	*	*	*	*	*	0.3	2.2	7.6	21.0	39.7	50.8
2001	4.1	*	*	*	*	*	0.4	2.1	7.8	20.8	38.6	48.0
2000	4.0	*	*	*	*	*	0.4	2.1	6.9	20.4	39.6	51.8
1999	4.1	*	*	*	*	*	0.4	2.1	8.5	20.1	38.3	51.3
All origins 3/, female												
2006	3.4	*	*	*	*	*	0.3	1.4	4.6	12.4	23.3	28.2
2005	3.5	*	*	*	*	*	0.2	1.3	5.2	12.8	23.7	27.9
2004	3.5	*	*	*	*	*	0.3	1.3	5.0	12.5	25.7	28.0
2003	3.7	*	*	*	*	*	0.2	1.4	4.6	13.8	26.7	30.1
2002	3.7	*	*	*	*	*	0.3	1.5	5.1	14.3	25.5	30.6
2001	3.8	*	*	*	*	*	0.3	1.3	5.5	14.5	25.8	31.3
2000	3.8	*	*	*	*	*	0.3	1.6	5.7	14.7	26.3	32.2
1999	3.8	*	*	*	*	*	0.3	1.5	5.9	14.8	26.2	30.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
Hispanic, both sexes												
2006	1.4	*	*	*	*	*	0.3	1.5	4.0	14.2	21.6	22.7
2005	1.4	*	*	*	*	*	*	1.5	5.4	13.4	21.4	20.1
2004	1.4	*	*	*	*	*	*	1.3	5.5	12.4	24.0	18.6
2003	1.3	*	*	*	*	*	*	1.4	5.2	12.6	23.9	24.9
2002	1.5	*	*	*	*	*	*	1.8	5.0	14.3	26.7	33.3
2001	1.4	*	*	*	*	*	*	1.3	5.8	14.9	24.7	29.4
2000	1.4	*	*	*	*	*	*	1.4	5.3	15.0	24.5	31.2
1999	1.4	*	*	*	*	*	*	1.7	6.8	13.4	23.1	28.1
Hispanic, male												
2006	1.4	*	*	*	*	*	*	1.8	4.6	15.9	24.7	30.4
2005	1.3	*	*	*	*	*	*	1.8	4.9	15.2	26.4	29.9
2004	1.3	*	*	*	*	*	*	1.3	6.4	14.1	30.0	*
2003	1.2	*	*	*	*	*	*	1.6	5.8	13.3	25.7	*
2002	1.6	*	*	*	*	*	*	2.1	6.6	17.6	31.6	53.5
2001	1.5	*	*	*	*	*	*	1.4	6.0	20.4	35.8	38.1
2000	1.3	*	*	*	*	*	*	1.5	5.5	18.4	30.0	*
1999	1.3	*	*	*	*	*	*	1.9	7.8	14.7	24.5	*
Hispanic, female												
2006	1.4	*	*	*	*	*	*	1.2	3.4	12.8	19.5	18.6
2005	1.4	*	*	*	*	*	*	1.3	5.8	11.9	17.9	15.0
2004	1.4	*	*	*	*	*	*	1.2	4.7	11.1	19.8	18.7
2003	1.4	*	*	*	*	*	*	1.2	4.7	12.1	22.6	22.7
2002	1.4	*	*	*	*	*	*	1.5	3.5	11.7	23.3	23.2
2001	1.3	*	*	*	*	*	*	1.2	5.6	10.6	17.2	25.1
2000	1.4	*	*	*	*	*	*	*	5.1	12.3	20.8	30.7
1999	1.5	*	*	*	*	*	*	1.5	5.9	12.4	22.2	31.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
Non-Hispanic 4/, both sexes												
2006	4.1	*	*	*	*	*	0.4	1.7	5.8	15.1	29.5	34.4
2005	4.2	*	*	*	*	*	0.3	1.6	6.1	16.1	29.9	34.2
2004	4.1	*	*	*	*	*	0.3	1.6	5.9	15.6	31.0	33.7
2003	4.3	*	*	*	*	*	0.3	1.8	6.0	16.8	31.4	36.7
2002	4.3	*	*	*	*	*	0.3	1.8	6.4	17.5	31.3	36.7
2001	4.3	*	*	*	*	*	0.3	1.7	6.6	17.5	31.1	36.5
2000	4.3	*	*	*	*	*	0.3	1.9	6.4	17.4	31.8	38.0
1999	4.3	*	*	*	*	*	0.4	1.8	7.1	17.4	31.2	36.8
Non-Hispanic 4/, male												
2006	4.5	*	*	*	*	*	0.4	2.0	7.0	18.2	38.2	46.6
2005	4.6	*	*	*	*	*	0.4	2.0	7.2	20.0	38.7	46.9
2004	4.4	*	*	*	*	*	0.4	1.9	7.0	19.2	38.6	45.9
2003	4.6	*	*	*	*	*	0.5	2.2	7.5	20.2	38.2	51.2
2002	4.6	*	*	*	*	*	0.4	2.2	7.6	21.2	40.0	50.6
2001	4.5	*	*	*	*	*	0.4	2.2	7.9	20.8	38.6	48.4
2000	4.4	*	*	*	*	*	0.4	2.2	7.0	20.5	39.9	52.5
1999	4.5	*	*	*	*	*	0.4	2.1	8.5	20.4	38.9	52.2
Non-Hispanic 4/, female												
2006	3.8	*	*	*	*	*	0.3	1.4	4.7	12.4	23.5	28.6
2005	3.8	*	*	*	*	*	0.2	1.3	5.1	12.8	24.0	28.5
2004	3.9	*	*	*	*	*	0.3	1.3	5.0	12.6	25.9	28.3
2003	4.0	*	*	*	*	*	0.2	1.4	4.6	13.8	26.8	30.3
2002	4.0	*	*	*	*	*	0.3	1.5	5.2	14.5	25.5	30.8
2001	4.1	*	*	*	*	*	0.3	1.4	5.5	14.8	26.1	31.5
2000	4.2	*	*	*	*	*	0.3	1.6	5.7	14.8	26.5	32.1
1999	4.1	*	*	*	*	*	0.3	1.5	5.8	15.0	26.3	30.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
Non-Hispanic white, both sexes												
2006 5/	4.2	*	*	*	*	*	0.3	1.4	5.1	13.9	28.5	33.3
2005 5/	4.3	*	*	*	*	*	0.3	1.3	5.4	14.8	29.0	33.5
2004 5/	4.2	*	*	*	*	*	0.3	1.4	5.3	14.2	29.6	32.6
2003 5/	4.4	*	*	*	*	*	0.3	1.4	5.4	15.6	29.8	35.3
2002	4.5	*	*	*	*	*	0.3	1.5	5.6	16.4	30.3	35.7
2001	4.4	*	*	*	*	*	0.3	1.5	5.7	16.3	29.7	35.1
2000	4.3	*	*	*	*	*	0.3	1.6	5.6	16.2	30.2	35.2
1999	4.3	*	*	*	*	*	0.3	1.5	6.1	16.2	29.6	35.0
Non-Hispanic white, male												
2006 5/	4.6	*	*	*	*	*	0.4	1.7	6.1	17.0	37.2	45.9
2005 5/	4.7	*	*	*	*	*	0.4	1.6	6.4	18.7	37.6	46.5
2004 5/	4.5	*	*	*	*	*	0.3	1.6	6.3	17.9	37.2	44.8
2003 5/	4.7	*	*	*	*	*	0.4	1.8	6.8	19.3	37.0	50.5
2002	4.8	*	*	*	*	*	0.3	1.9	6.9	20.2	39.2	50.1
2001	4.6	*	*	*	*	*	0.3	1.9	6.9	19.5	37.4	47.3
2000	4.6	*	*	*	*	*	0.4	1.8	6.3	19.2	38.5	50.9
1999	4.6	*	*	*	*	*	0.3	1.9	7.5	19.2	37.4	50.6
Non-Hispanic white, female												
2006 5/	3.8	*	*	*	*	*	0.2	1.1	4.1	11.2	22.5	27.5
2005 5/	3.9	*	*	*	*	*	0.2	1.0	4.5	11.4	23.1	27.5
2004 5/	3.9	*	*	*	*	*	0.2	1.1	4.3	11.1	24.4	27.1
2003 5/	4.0	*	*	*	*	*	0.2	1.1	4.0	12.4	24.9	28.7
2002	4.1	*	*	*	*	*	0.2	1.1	4.5	13.1	24.4	29.5
2001	4.1	*	*	*	*	*	0.2	1.1	4.7	13.6	24.6	30.0
2000	4.1	*	*	*	*	*	0.2	1.4	4.9	13.6	24.8	28.9
1999	4.1	*	*	*	*	*	0.2	1.2	4.7	13.7	24.6	28.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
Non-Hispanic black, both sexes												
2006 5/	4.8	*	*	*	*	*	0.7	4.0	12.9	28.4	48.0	52.0
2005 5/	4.7	*	*	*	*	*	0.6	3.7	12.7	30.3	47.6	50.2
2004 5/	4.9	*	*	*	*	*	0.7	3.5	12.6	31.0	53.9	54.7
2003 5/	5.0	*	*	*	*	*	0.8	4.3	12.2	31.4	55.8	60.4
2002	4.8	*	*	*	*	*	0.7	4.3	13.4	31.8	49.2	55.7
2001	5.0	*	*	*	*	*	0.9	3.7	15.3	31.8	53.2	59.3
2000	5.2	*	*	*	*	*	0.8	4.5	14.1	32.5	55.6	72.8
1999	5.1	*	*	*	*	*	0.9	4.3	16.8	30.7	53.6	63.4
Non-Hispanic black, male												
2006 5/	5.1	*	*	*	*	*	0.8	4.6	16.3	36.5	60.6	66.0
2005 5/	4.9	*	*	*	*	*	*	4.6	15.0	36.3	62.8	62.8
2004 5/	4.8	*	*	*	*	*	0.8	4.0	14.5	37.1	65.3	72.1
2003 5/	4.9	*	*	*	*	*	0.9	5.3	15.7	34.5	64.1	74.6
2002	4.8	*	*	*	*	*	0.8	4.8	15.9	37.6	60.1	71.1
2001	5.0	*	*	*	*	*	1.1	4.3	18.5	37.2	66.0	72.8
2000	5.0	*	*	*	*	*	*	5.0	15.4	39.1	68.5	81.0
1999	5.0	*	*	*	*	*	1.1	4.4	19.0	35.6	65.0	80.8
Non-Hispanic black, female												
2006 5/	4.6	*	*	*	*	*	0.7	3.4	10.2	22.7	40.9	46.3
2005 5/	4.6	*	*	*	*	*	*	2.9	10.8	26.0	38.9	45.2
2004 5/	4.9	*	*	*	*	*	*	3.1	11.1	26.7	47.5	47.8
2003 5/	5.1	*	*	*	*	*	*	3.4	9.3	29.2	51.2	54.9
2002	4.9	*	*	*	*	*	*	3.8	11.3	27.7	43.1	49.7
2001	5.0	*	*	*	*	*	0.8	3.1	12.7	28.0	46.0	54.3
2000	5.4	*	*	*	*	*	0.9	4.1	13.1	27.9	48.3	69.8
1999	5.3	*	*	*	*	*	0.7	4.3	15.1	27.2	47.3	56.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
All origins 3/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	0.2	0.4
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
2000	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
All origins 3/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
Non-Hispanic 4/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
2000	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Non-Hispanic 4/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
Non-Hispanic white, both sexes												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Non-Hispanic white, male												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
All origins 3/, both sexes												
2006	21.1	0.5	0.7	0.6	1.2	1.8	4.1	12.7	33.5	75.0	140.1	210.1
2005	21.2	*	0.8	0.7	1.4	1.7	4.2	13.1	34.3	75.2	141.6	212.9
2004	21.1	0.6	0.8	0.7	1.3	1.7	4.3	12.7	34.4	76.8	143.0	213.1
2003	21.5	*	0.7	0.7	1.3	1.7	4.2	13.8	35.2	77.9	146.7	224.0
2002	21.8	0.6	0.8	0.7	1.3	1.8	4.5	13.7	36.6	80.8	146.7	230.1
2001	22.0	*	0.8	0.7	1.3	1.8	4.6	14.1	37.3	82.0	148.7	234.5
2000	22.4	0.6	0.8	0.6	1.3	1.8	4.6	14.1	38.3	85.5	152.6	239.8
1999	23.7	*	0.8	0.7	1.3	1.9	5.1	15.0	41.5	91.4	160.8	250.4
All origins 3/, male												
2006	21.9	*	0.7	0.6	1.6	2.2	4.3	14.3	40.0	89.0	173.9	259.6
2005	22.1	*	0.8	0.7	1.7	2.2	4.6	14.9	41.0	90.1	173.1	263.5
2004	21.9	*	0.7	0.8	1.7	2.1	4.6	14.4	40.6	92.5	177.9	260.6
2003	22.3	*	0.8	0.7	1.6	2.1	4.7	15.7	41.5	93.4	180.8	275.0
2002	22.7	*	0.8	0.7	1.6	2.3	5.0	15.9	43.2	98.1	180.0	290.8
2001	22.6	*	0.9	0.7	1.7	2.2	4.9	16.4	43.8	98.5	181.1	295.2
2000	23.0	*	1.0	0.7	1.6	2.1	5.1	16.0	44.9	103.2	187.8	300.7
1999	24.5	*	0.8	0.8	1.7	2.4	5.8	17.3	49.1	110.4	196.1	318.9
All origins 3/, female												
2006	20.2	*	0.6	0.6	0.9	1.4	3.9	11.3	27.5	63.2	117.0	186.9
2005	20.4	*	0.7	0.6	1.1	1.3	3.8	11.3	28.1	62.6	120.2	189.6
2004	20.3	*	0.9	0.6	0.9	1.4	3.9	11.0	28.6	63.7	119.5	191.8
2003	20.8	*	0.5	0.6	0.9	1.3	3.8	11.9	29.3	64.8	123.9	201.4
2002	21.0	*	0.8	0.7	0.9	1.4	4.1	11.6	30.4	66.3	124.6	203.8
2001	21.4	*	0.6	0.6	0.9	1.5	4.2	11.7	31.3	68.4	127.4	209.1
2000	21.9	*	0.7	0.6	1.0	1.4	4.2	12.2	32.2	70.8	129.6	215.0
1999	23.1	*	0.7	0.7	1.0	1.4	4.5	12.8	34.5	75.8	137.9	222.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
Hispanic, both sexes												
2006	7.7	*	0.6	0.6	1.5	1.3	3.1	9.4	24.4	54.4	97.6	145.5
2005	7.6	*	0.6	0.6	1.3	1.7	3.1	8.0	22.7	55.8	108.9	139.5
2004	7.2	*	0.6	0.5	1.3	1.3	2.6	8.9	22.6	52.7	106.5	140.8
2003	7.7	*	0.6	0.7	1.3	1.5	3.3	9.8	23.7	58.7	109.4	168.9
2002	7.8	*	*	0.8	1.6	1.5	3.0	9.5	25.4	60.3	112.8	164.1
2001	7.7	*	*	0.6	1.3	1.2	3.4	9.8	26.0	60.7	110.9	193.8
2000	7.8	*	0.8	0.7	1.2	1.5	2.7	9.2	26.6	64.1	124.6	192.4
1999	8.1	*	*	0.7	1.3	1.6	3.5	10.6	28.8	64.8	122.3	202.1
Hispanic, male												
2006	7.6	*	*	0.6	1.9	1.7	2.7	9.2	27.0	67.1	113.8	163.5
2005	7.8	*	*	0.5	1.6	2.0	3.5	8.5	27.7	67.1	129.6	159.4
2004	7.1	*	*	0.6	1.8	1.6	2.9	9.3	25.3	59.4	131.3	147.1
2003	7.9	*	*	0.7	1.6	2.1	3.3	11.2	27.7	71.0	127.3	194.0
2002	7.8	*	*	0.8	1.9	1.7	3.1	10.4	29.6	69.2	138.7	204.1
2001	7.7	*	*	0.6	1.8	1.5	3.4	10.7	30.2	67.7	140.9	213.3
2000	7.7	*	*	0.7	1.4	1.9	2.9	9.5	29.2	73.6	151.9	231.8
1999	8.0	*	*	0.7	1.8	1.9	3.5	11.1	33.3	77.1	144.9	235.1
Hispanic, female												
2006	7.8	*	*	0.7	1.0	0.9	3.5	9.6	21.9	44.0	86.3	136.0
2005	7.4	*	*	0.6	1.0	1.4	2.6	7.6	18.1	46.7	94.4	129.0
2004	7.3	*	*	*	0.6	1.1	2.4	8.4	20.2	47.3	89.4	137.6
2003	7.6	*	*	0.7	1.1	0.8	3.2	8.4	20.1	48.8	97.2	156.3
2002	7.7	*	*	0.7	1.2	1.1	2.9	8.6	21.6	53.2	95.0	144.2
2001	7.7	*	*	0.6	0.7	0.8	3.5	8.9	22.4	55.3	90.7	184.3
2000	7.9	*	*	0.6	1.1	1.1	2.5	9.0	24.3	56.6	106.3	173.1
1999	8.2	*	*	0.7	0.8	1.2	3.6	10.1	24.7	55.1	107.2	186.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
Non-Hispanic 4/, both sexes												
2006	23.4	*	0.7	0.6	1.2	1.9	4.3	13.1	34.2	76.5	142.6	213.1
2005	23.5	*	0.8	0.7	1.4	1.7	4.3	13.6	35.3	76.6	143.4	216.2
2004	23.3	0.6	0.8	0.7	1.3	1.8	4.5	13.1	35.2	78.5	144.8	215.9
2003	23.7	*	0.7	0.7	1.2	1.8	4.4	14.1	36.0	79.0	148.4	225.8
2002	24.0	*	0.9	0.7	1.2	1.9	4.8	14.1	37.3	81.9	148.0	232.1
2001	24.1	*	0.8	0.7	1.3	2.0	4.7	14.4	38.1	83.1	150.1	235.3
2000	24.4	0.7	0.8	0.6	1.3	1.8	4.8	14.5	39.1	86.5	153.3	241.0
1999	25.8	*	0.8	0.7	1.3	1.9	5.3	15.3	42.3	92.7	161.9	251.6
Non-Hispanic 4/, male												
2006	24.5	*	0.8	0.6	1.5	2.3	4.6	14.8	41.0	90.4	177.4	264.6
2005	24.5	*	0.8	0.8	1.7	2.2	4.7	15.6	42.1	91.7	175.5	268.7
2004	24.4	*	0.8	0.8	1.6	2.2	4.9	14.9	41.7	94.7	180.3	265.7
2003	24.6	*	0.8	0.7	1.5	2.1	4.9	16.1	42.4	94.7	183.4	278.2
2002	25.0	*	0.9	0.7	1.6	2.4	5.3	16.4	44.1	99.7	181.5	294.0
2001	24.9	*	1.0	0.8	1.7	2.4	5.1	16.9	44.7	100.0	182.7	297.9
2000	25.2	*	0.9	0.6	1.6	2.2	5.3	16.5	45.9	104.7	188.7	303.0
1999	26.8	*	1.0	0.8	1.6	2.5	6.1	17.8	50.1	112.0	197.6	321.9
Non-Hispanic 4/, female												
2006	22.2	*	0.6	0.5	0.8	1.5	3.9	11.4	28.0	64.7	118.9	189.1
2005	22.4	*	0.8	0.7	1.1	1.3	3.9	11.7	28.9	63.8	121.6	192.2
2004	22.3	*	0.8	0.6	1.0	1.5	4.1	11.3	29.3	64.8	121.0	193.7
2003	22.8	*	0.6	0.6	0.9	1.5	3.8	12.2	30.0	65.9	125.1	202.7
2002	22.9	*	0.8	0.7	0.9	1.4	4.3	11.8	31.0	67.1	125.8	205.4
2001	23.3	*	0.6	0.6	0.9	1.6	4.3	12.0	31.9	69.1	128.7	209.3
2000	23.7	*	0.8	0.6	1.0	1.4	4.4	12.5	32.8	71.5	130.2	215.9
1999	25.0	*	0.7	0.6	1.1	1.4	4.6	13.0	35.1	76.9	138.8	223.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
Non-Hispanic white, both sexes												
2006 5/	25.6	*	0.6	0.6	1.2	1.9	4.2	12.5	33.5	77.1	145.3	217.1
2005 5/	25.6	*	0.8	0.7	1.4	1.7	4.3	12.9	34.5	76.9	145.8	220.9
2004 5/	25.4	*	0.8	0.7	1.3	1.9	4.4	12.3	34.4	78.7	146.6	220.6
2003 5/	25.9	*	0.6	0.7	1.3	1.7	4.2	13.4	35.4	79.5	150.3	229.9
2002	26.0	*	0.8	0.7	1.2	1.9	4.6	13.5	36.5	81.8	149.1	235.2
2001	26.0	*	0.9	0.7	1.3	2.0	4.5	13.6	36.8	82.5	150.8	238.9
2000	26.5	*	0.8	0.7	1.3	1.9	4.5	13.6	38.2	86.3	153.9	244.9
1999	27.9	*	0.9	0.7	1.3	2.0	5.0	14.6	40.9	92.7	162.3	253.6
Non-Hispanic white, male												
2006 5/	27.0	*	0.7	0.7	1.5	2.4	4.6	14.1	40.3	91.5	180.8	271.8
2005 5/	26.8	*	0.8	0.7	1.7	2.1	4.7	14.8	41.0	91.3	178.8	276.1
2004 5/	26.5	*	0.9	0.8	1.6	2.4	4.7	13.9	40.3	94.4	183.2	272.4
2003 5/	26.8	*	0.7	0.7	1.6	2.1	4.8	15.2	41.5	94.8	186.1	284.6
2002	27.2	*	0.8	0.7	1.5	2.4	5.0	15.7	42.9	100.0	183.2	300.2
2001	26.8	*	1.1	0.7	1.7	2.3	5.0	15.9	42.8	99.3	183.7	302.9
2000	27.2	*	1.0	0.7	1.6	2.3	5.0	15.6	44.1	104.6	189.1	309.5
1999	28.8	*	1.1	0.8	1.5	2.6	5.8	16.6	48.3	111.5	198.9	326.9
Non-Hispanic white, female												
2006 5/	24.3	*	0.5	0.6	0.8	1.4	3.8	11.0	27.0	64.6	120.7	191.5
2005 5/	24.5	*	0.7	0.7	1.1	1.2	3.8	11.1	28.4	64.3	123.1	195.6
2004 5/	24.4	*	0.8	0.5	1.0	1.4	4.0	10.8	28.8	65.2	121.7	197.4
2003 5/	25.0	*	0.6	0.6	0.9	1.3	3.7	11.7	29.6	66.4	126.2	205.8
2002	24.9	*	0.7	0.7	0.9	1.4	4.3	11.3	30.5	66.3	126.3	207.0
2001	25.3	*	0.7	0.6	1.0	1.6	4.0	11.4	31.3	68.3	129.1	212.3
2000	25.8	*	0.6	0.6	1.0	1.4	4.1	11.7	32.6	71.0	130.8	218.9
1999	27.0	*	0.8	0.6	1.1	1.3	4.2	12.6	33.9	77.0	138.3	224.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
Non-Hispanic black, both sexes												
2006 5/	17.5	*	1.0	0.6	1.3	2.2	5.4	18.8	47.5	87.4	139.5	188.7
2005 5/	17.7	*	*	1.0	1.3	2.4	5.6	19.9	47.3	87.2	146.5	184.6
2004 5/	17.8	*	*	0.9	1.6	1.8	6.2	19.8	48.6	91.9	149.1	188.2
2003 5/	17.9	*	0.9	0.8	1.2	2.4	5.9	21.1	49.0	90.0	151.8	206.4
2002	18.7	*	1.1	0.8	1.3	2.4	6.5	20.5	51.8	97.9	159.6	223.2
2001	18.9	*	*	0.7	1.3	2.4	6.1	22.2	54.0	101.8	158.4	218.5
2000	18.9	*	0.9	0.6	1.5	1.7	7.7	22.6	52.8	101.2	163.6	211.1
1999	20.2	*	*	0.8	1.5	2.0	7.6	23.6	60.6	105.9	172.7	244.8
Non-Hispanic black, male												
2006 5/	18.1	*	*	0.6	1.4	2.5	5.6	23.0	57.6	103.9	180.5	216.2
2005 5/	18.6	*	*	1.1	1.5	3.0	5.9	23.2	60.3	112.9	175.5	221.7
2004 5/	19.0	*	*	0.9	2.0	1.9	6.7	24.8	62.5	117.9	185.7	223.5
2003 5/	18.8	*	*	0.8	1.5	2.4	6.6	25.3	62.2	113.6	189.9	239.7
2002	19.4	*	*	0.9	1.8	2.7	7.6	24.5	64.7	119.0	197.8	265.8
2001	20.0	*	*	0.8	1.8	2.8	6.3	27.1	69.0	128.4	195.0	287.6
2000	20.0	*	*	0.7	2.0	1.9	8.7	26.9	69.1	123.8	212.0	253.6
1999	21.3	*	*	0.7	2.0	2.4	8.5	29.7	77.5	134.5	208.6	293.6
Non-Hispanic black, female												
2006 5/	16.9	*	*	*	1.1	2.0	5.3	15.2	39.4	75.7	116.3	177.5
2005 5/	16.8	*	*	0.8	1.1	1.9	5.3	17.1	36.8	68.8	130.1	169.7
2004 5/	16.7	*	*	1.0	1.3	1.8	5.7	15.6	37.4	73.5	128.5	174.3
2003 5/	17.1	*	*	0.8	0.8	2.3	5.3	17.5	38.5	73.2	130.4	193.6
2002	18.0	*	*	0.8	0.8	2.1	5.5	17.1	41.4	83.0	137.9	206.9
2001	17.8	*	*	0.6	0.9	2.0	5.8	18.0	42.0	83.0	137.8	192.9
2000	17.8	*	*	*	1.0	1.5	6.8	18.9	39.6	85.2	136.3	195.4
1999	19.2	*	*	0.9	1.0	1.7	6.9	18.3	47.0	85.8	152.7	226.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
All origins 3/, both sexes												
2006	4.7	1.6	0.4	0.2	0.2	0.4	0.7	1.8	4.2	12.5	36.4	78.0
2005	4.6	1.4	0.3	0.2	0.2	0.4	0.8	1.7	4.3	12.6	35.2	76.7
2004	4.6	1.7	0.3	0.2	0.3	0.4	0.7	1.7	4.5	12.7	35.8	77.7
2003	4.7	1.5	0.3	0.2	0.2	0.4	0.8	1.7	4.4	12.8	36.4	80.5
2002	4.6	1.7	0.4	0.2	0.2	0.4	0.8	1.8	4.5	12.8	36.1	78.7
2001	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.8	5.0	14.1	37.0	82.7
2000	4.8	1.6	0.3	0.2	0.2	0.4	0.8	1.8	5.0	13.8	37.6	84.9
1999	4.7	1.5	0.4	0.2	0.3	0.4	0.8	1.8	4.8	14.6	36.6	82.9
All origins 3/, male												
2006	4.7	1.6	0.3	0.2	0.2	0.3	0.8	2.0	4.8	15.7	47.5	92.3
2005	4.7	1.0	0.4	0.2	0.2	0.4	0.8	1.9	5.2	15.7	45.8	92.7
2004	4.7	1.7	0.3	0.2	0.3	0.5	0.8	1.9	5.4	15.7	45.6	98.9
2003	4.7	1.6	0.4	0.2	0.3	0.5	0.9	1.8	5.1	16.0	46.2	101.7
2002	4.6	1.5	0.4	0.2	0.2	0.4	0.9	1.9	5.3	15.8	45.9	101.1
2001	4.8	1.3	0.4	0.2	0.3	0.4	0.9	2.0	5.5	17.3	48.2	107.6
2000	4.8	1.6	0.4	0.2	0.3	0.4	0.8	2.1	5.6	17.0	47.5	108.9
1999	4.8	1.5	0.4	0.2	0.3	0.5	0.9	2.0	5.8	18.4	46.1	109.2
All origins 3/, female												
2006	4.7	1.6	0.5	0.2	0.2	0.4	0.7	1.6	3.6	9.8	28.8	71.4
2005	4.6	1.8	0.3	0.2	0.2	0.4	0.8	1.5	3.6	10.0	28.0	69.4
2004	4.6	1.8	0.3	0.2	0.2	0.4	0.7	1.5	3.7	10.2	29.2	68.1
2003	4.7	1.5	0.3	0.2	0.2	0.4	0.6	1.5	3.8	10.2	29.9	71.1
2002	4.6	2.0	0.4	0.2	0.2	0.4	0.8	1.7	3.7	10.4	29.6	68.9
2001	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.6	4.4	11.3	29.7	72.3
2000	4.9	1.5	0.3	0.2	0.2	0.4	0.8	1.6	4.5	11.2	31.1	75.2
1999	4.7	1.5	0.4	0.3	0.2	0.4	0.6	1.6	3.9	11.5	30.4	72.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
Hispanic, both sexes												
2006	1.6	*	*	*	0.3	0.3	0.5	1.4	3.5	8.9	25.0	48.5
2005	1.3	*	*	*	*	0.3	0.4	1.3	3.2	8.1	20.8	34.2
2004	1.5	*	*	*	*	0.4	0.5	1.3	3.8	8.7	23.4	48.5
2003	1.5	*	*	*	0.3	0.4	0.4	1.1	3.1	8.7	28.9	48.2
2002	1.5	*	*	0.4	*	*	0.6	1.5	3.0	8.9	24.7	48.8
2001	1.5	*	*	0.3	*	0.3	0.7	1.5	3.6	8.8	23.1	57.5
2000	1.5	*	*	*	*	0.4	0.7	1.4	3.1	9.1	24.3	70.3
1999	1.5	*	*	*	*	0.4	0.6	1.3	3.7	10.5	23.1	51.9
Hispanic, male												
2006	1.6	*	*	*	*	*	*	1.8	4.1	11.2	30.4	53.0
2005	1.4	*	*	*	*	*	*	1.5	3.9	10.7	27.4	37.4
2004	1.5	*	*	*	*	*	*	1.6	4.1	9.7	27.9	57.4
2003	1.5	*	*	*	*	0.5	*	1.2	3.9	10.0	33.7	52.3
2002	1.4	*	*	*	*	*	*	1.4	3.7	10.3	31.6	56.9
2001	1.5	*	*	*	*	*	0.8	1.8	4.0	10.0	30.0	70.5
2000	1.3	*	*	*	*	*	*	*	3.1	9.7	26.6	86.7
1999	1.3	*	*	*	*	*	0.8	1.8	3.9	10.3	24.5	49.6
Hispanic, female												
2006	1.7	*	*	*	*	*	*	0.9	3.0	7.0	21.2	46.1
2005	1.3	*	*	*	*	*	*	1.0	2.5	6.0	16.3	32.6
2004	1.5	*	*	*	*	*	*	1.0	3.5	7.8	20.3	43.9
2003	1.6	*	*	*	*	*	*	1.0	2.3	7.6	25.5	46.1
2002	1.5	*	*	*	*	*	*	1.5	2.4	7.8	20.0	44.8
2001	1.5	*	*	*	*	*	*	1.2	3.3	7.9	18.4	51.2
2000	1.7	*	*	*	*	*	0.9	1.6	3.1	8.6	22.8	62.3
1999	1.6	*	*	*	*	*	*	*	3.5	10.7	22.2	53.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
Non-Hispanic 4/, both sexes												
2006	5.2	1.5	0.4	0.2	0.2	0.4	0.8	1.9	4.2	12.8	37.1	79.5
2005	5.2	1.3	0.3	0.2	0.2	0.4	0.9	1.8	4.4	12.9	36.0	78.6
2004	5.1	1.7	0.4	0.2	0.3	0.4	0.8	1.7	4.6	12.9	36.4	78.9
2003	5.2	1.5	0.4	0.2	0.2	0.4	0.8	1.7	4.5	13.1	36.8	81.8
2002	5.1	1.6	0.4	0.2	0.2	0.5	0.8	1.8	4.6	13.1	36.6	79.8
2001	5.3	1.4	0.4	0.2	0.3	0.4	0.8	1.9	5.0	14.4	37.5	83.5
2000	5.3	1.7	0.4	0.2	0.3	0.4	0.8	1.9	5.2	14.1	38.1	85.2
1999	5.2	1.6	0.4	0.2	0.3	0.4	0.8	1.8	4.8	14.8	37.0	83.8
Non-Hispanic 4/, male												
2006	5.3	1.6	0.3	0.2	0.2	0.3	0.8	2.1	4.9	16.0	48.5	94.5
2005	5.2	*	0.4	0.2	0.3	0.4	0.9	2.0	5.2	16.1	46.8	95.4
2004	5.2	1.5	0.4	0.2	0.3	0.5	0.8	1.9	5.4	16.1	46.5	101.0
2003	5.2	1.6	0.4	0.2	0.2	0.5	0.9	1.9	5.2	16.3	46.7	103.9
2002	5.1	1.3	0.4	0.2	0.3	0.5	0.9	1.9	5.4	16.2	46.4	102.9
2001	5.3	1.5	0.4	0.2	0.3	0.4	0.9	2.1	5.6	17.8	48.9	109.0
2000	5.3	1.8	0.5	0.2	0.3	0.4	0.9	2.2	5.8	17.5	48.3	109.4
1999	5.3	1.6	0.5	0.2	0.3	0.5	0.9	2.0	5.9	18.8	46.9	111.4
Non-Hispanic 4/, female												
2006	5.2	1.4	0.5	0.2	0.2	0.4	0.7	1.7	3.6	10.1	29.2	72.5
2005	5.1	1.8	*	0.2	0.2	0.4	0.9	1.6	3.7	10.3	28.6	71.0
2004	5.0	1.9	0.4	0.2	0.3	0.4	0.7	1.5	3.7	10.3	29.7	69.0
2003	5.1	1.3	*	0.2	0.2	0.4	0.7	1.6	3.9	10.4	30.2	72.1
2002	5.1	2.0	0.4	0.2	0.2	0.4	0.8	1.7	3.8	10.5	30.0	69.8
2001	5.2	1.4	0.4	0.3	0.3	0.4	0.8	1.7	4.5	11.5	30.1	72.9
2000	5.3	1.7	*	0.3	0.2	0.4	0.8	1.6	4.6	11.4	31.4	75.4
1999	5.1	1.7	0.4	0.3	0.2	0.4	0.7	1.7	3.9	11.5	30.6	72.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
Non-Hispanic white, both sexes												
2006 5/	5.9	1.5	0.5	0.2	0.2	0.3	0.7	1.8	4.2	13.2	38.2	81.6
2005 5/	5.8	1.2	0.3	0.2	0.2	0.4	0.9	1.7	4.4	13.2	37.2	81.7
2004 5/	5.7	1.8	0.4	0.2	0.3	0.4	0.7	1.6	4.4	13.2	37.4	81.6
2003 5/	5.8	1.4	0.3	0.2	0.2	0.5	0.8	1.7	4.5	13.4	38.1	84.4
2002	5.7	1.8	0.4	0.2	0.2	0.4	0.8	1.8	4.6	13.4	37.5	81.8
2001	5.9	1.3	0.4	0.2	0.3	0.4	0.8	1.9	5.0	14.7	38.6	84.8
2000	5.9	1.8	0.4	0.2	0.3	0.4	0.8	1.8	5.2	14.2	39.1	87.6
1999	5.7	1.5	0.4	0.3	0.2	0.4	0.7	1.8	4.8	15.0	37.8	86.3
Non-Hispanic white, male												
2006 5/	6.0	*	*	0.2	0.2	0.3	0.8	2.1	4.9	16.5	50.1	96.0
2005 5/	5.9	*	0.4	0.2	0.3	0.5	0.9	1.9	5.2	16.6	48.7	100.0
2004 5/	5.9	*	0.4	0.2	0.3	0.5	0.8	1.8	5.3	16.5	47.7	105.6
2003 5/	5.9	1.7	*	0.2	0.3	0.5	0.9	1.9	5.1	16.8	48.4	107.9
2002	5.8	*	*	0.2	0.2	0.5	1.0	1.9	5.4	16.7	47.6	106.0
2001	6.0	*	*	0.2	0.2	0.5	0.9	2.1	5.7	18.3	50.1	111.4
2000	5.9	1.8	0.5	0.2	0.3	0.4	0.8	2.1	5.8	17.4	49.9	113.1
1999	5.9	*	*	0.2	0.2	0.4	0.9	2.0	5.9	19.2	48.1	116.8
Non-Hispanic white, female												
2006 5/	5.8	*	0.6	*	0.2	0.4	0.6	1.6	3.5	10.3	29.9	74.9
2005 5/	5.7	*	*	0.2	0.2	0.4	0.8	1.5	3.6	10.3	29.3	73.4
2004 5/	5.6	2.0	*	0.2	0.2	0.3	0.6	1.4	3.6	10.4	30.4	70.9
2003 5/	5.8	*	*	*	0.2	0.5	0.6	1.5	4.0	10.4	31.1	74.0
2002	5.6	2.1	*	*	0.2	0.3	0.7	1.6	3.9	10.7	30.7	71.4
2001	5.8	*	*	0.3	0.3	0.4	0.7	1.7	4.5	11.6	30.9	73.7
2000	5.9	1.8	*	0.3	0.2	0.4	0.8	1.5	4.7	11.5	32.1	77.3
1999	5.6	*	*	0.3	0.2	0.4	0.6	1.5	3.7	11.5	31.1	74.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
Non-Hispanic black, both sexes												
2006 5/	3.2	*	*	*	0.4	0.6	1.2	2.6	5.6	11.9	32.2	63.4
2005 5/	3.1	*	*	*	*	0.5	1.4	2.5	5.5	13.1	28.4	55.7
2004 5/	3.2	*	*	0.3	*	0.7	1.2	2.3	6.3	13.0	32.0	57.9
2003 5/	3.0	*	*	*	*	0.5	1.2	2.5	5.3	13.3	29.6	58.9
2002	3.1	*	*	*	*	0.6	1.2	2.5	5.6	12.8	31.8	64.4
2001	3.2	*	*	0.4	0.4	*	1.1	2.1	6.0	14.5	31.9	72.7
2000	3.2	*	*	*	*	0.6	1.1	2.7	5.6	15.1	31.3	64.2
1999	3.3	*	*	*	0.4	0.6	1.2	2.6	6.4	15.9	33.1	62.7
Non-Hispanic black, male												
2006 5/	3.1	*	*	*	*	*	0.9	2.4	6.5	15.1	42.3	82.5
2005 5/	2.8	*	*	*	*	*	1.4	2.7	6.3	14.2	34.5	56.1
2004 5/	3.1	*	*	*	*	*	1.3	2.6	8.0	15.7	39.9	62.0
2003 5/	3.0	*	*	*	*	*	1.2	2.5	6.9	16.5	36.6	68.2
2002	2.9	*	*	*	*	*	1.0	2.2	7.3	14.3	39.3	81.0
2001	3.1	*	*	*	*	*	1.1	1.9	6.6	17.6	42.3	87.8
2000	3.1	*	*	*	*	*	1.2	2.8	7.0	19.5	36.4	76.3
1999	3.1	*	*	*	*	0.8	1.2	2.4	7.3	19.2	38.8	64.2
Non-Hispanic black, female												
2006 5/	3.3	*	*	*	*	*	1.4	2.7	4.9	9.6	26.5	55.7
2005 5/	3.3	*	*	*	*	*	1.5	2.3	4.9	12.3	24.9	55.5
2004 5/	3.3	*	*	*	*	0.8	1.1	2.1	4.9	11.2	27.5	56.2
2003 5/	3.1	*	*	*	*	*	1.2	2.5	4.1	11.1	25.6	55.3
2002	3.4	*	*	*	*	0.8	1.4	2.7	4.3	11.7	27.5	58.1
2001	3.3	*	*	*	*	*	1.0	2.2	5.4	12.2	26.1	67.2
2000	3.3	*	*	*	*	0.8	1.0	2.6	4.5	12.1	28.5	59.7
1999	3.5	*	*	*	*	*	1.2	2.8	5.7	13.6	29.9	62.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
All origins 3/, both sexes												
2006	1.3	*	0.2	0.1	0.2	0.3	0.4	0.5	1.0	2.1	7.5	30.0
2005	1.6	*	0.1	0.1	0.2	0.4	0.4	0.6	1.0	2.7	9.1	36.6
2004	1.6	*	0.2	0.1	0.3	0.4	0.5	0.6	1.1	2.6	8.6	38.1
2003	1.6	0.5	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	8.8	40.7
2002	1.6	*	0.1	0.1	0.3	0.5	0.4	0.6	1.0	2.8	9.1	40.3
2001	1.6	*	0.1	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.0
2000	1.6	*	0.2	0.1	0.2	0.4	0.4	0.6	1.1	2.8	9.5	42.4
1999	1.6	*	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	9.5	43.8
All origins 3/, male												
2006	1.1	*	*	*	0.3	0.3	0.4	0.5	0.9	2.1	7.9	29.6
2005	1.2	*	*	*	0.3	0.4	0.4	0.5	1.0	2.6	9.5	33.5
2004	1.3	*	*	0.1	0.3	0.4	0.5	0.5	1.2	2.7	8.9	36.5
2003	1.3	*	*	0.1	0.2	0.4	0.4	0.6	1.2	3.0	9.3	38.0
2002	1.3	*	*	0.1	0.3	0.5	0.4	0.6	1.0	2.9	8.9	38.3
2001	1.3	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	9.9	40.0
2000	1.2	*	*	0.1	0.2	0.3	0.3	0.6	1.1	2.8	9.5	40.1
1999	1.3	*	*	0.1	0.3	0.4	0.5	0.6	0.9	2.7	10.1	43.4
All origins 3/, female												
2006	1.6	*	*	*	0.2	0.4	0.5	0.6	1.0	2.2	7.1	30.2
2005	1.9	*	*	*	0.2	0.4	0.4	0.7	1.0	2.7	8.9	38.0
2004	1.8	*	*	0.1	0.3	0.4	0.4	0.6	1.0	2.5	8.4	38.8
2003	1.9	*	*	0.1	0.3	0.4	0.4	0.6	0.8	2.3	8.5	41.9
2002	1.9	*	*	0.1	0.2	0.4	0.4	0.7	1.0	2.6	9.2	41.2
2001	2.0	*	*	0.1	0.3	0.3	0.5	0.6	1.0	2.5	9.6	42.9
2000	2.0	*	*	0.1	0.2	0.4	0.5	0.6	1.0	2.8	9.4	43.3
1999	1.9	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.2	44.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
Hispanic, both sexes												
2006	0.4	*	*	*	*	*	*	*	*	1.5	6.1	21.1
2005	0.4	*	*	*	*	*	*	*	*	2.4	6.4	21.4
2004	0.5	*	*	*	*	*	*	*	*	*	6.3	28.1
2003	0.5	*	*	*	*	*	*	*	*	2.0	7.5	29.0
2002	0.5	*	*	*	*	*	*	*	*	3.1	7.3	34.4
2001	0.5	*	*	*	*	*	*	*	*	2.2	7.8	28.8
2000	0.4	*	*	*	*	*	*	*	*	2.2	7.1	17.3
1999	0.5	*	*	*	*	*	*	*	*	2.6	7.3	35.1
Hispanic, male												
2006	0.3	*	*	*	*	*	*	*	*	*	*	*
2005	0.4	*	*	*	*	*	*	*	*	*	7.0	*
2004	0.4	*	*	*	*	*	*	*	*	*	7.2	*
2003	0.4	*	*	*	*	*	*	*	*	*	*	30.8
2002	0.5	*	*	*	*	*	*	*	*	*	*	53.5
2001	0.4	*	*	*	*	*	*	*	*	*	9.4	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.5	*	*	*	*	*	*	*	*	*	6.2	23.4
2005	0.5	*	*	*	*	*	*	*	*	3.0	6.0	21.5
2004	0.6	*	*	*	*	*	*	*	*	*	5.7	30.3
2003	0.6	*	*	*	*	*	*	*	*	*	7.6	28.1
2002	0.5	*	*	*	*	*	*	*	*	3.0	7.3	24.9
2001	0.6	*	*	*	*	*	*	*	*	*	6.6	32.6
2000	0.5	*	*	*	*	*	*	*	*	*	6.9	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	32.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
Non-Hispanic 4/, both sexes												
2006	1.5	*	0.2	0.1	0.2	0.4	0.5	0.6	1.0	2.2	7.5	30.3
2005	1.7	*	*	0.1	0.3	0.5	0.5	0.6	1.0	2.7	9.3	37.2
2004	1.7	*	0.2	0.1	0.3	0.5	0.5	0.6	1.1	2.6	8.7	38.5
2003	1.8	0.6	*	0.1	0.3	0.5	0.4	0.6	1.0	2.7	8.9	41.2
2002	1.8	*	*	0.1	0.3	0.5	0.5	0.7	1.0	2.7	9.2	40.5
2001	1.8	*	*	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.5
2000	1.8	*	0.3	0.1	0.2	0.4	0.5	0.6	1.1	2.8	9.5	43.2
1999	1.8	*	0.2	0.1	0.4	0.4	0.5	0.7	1.0	2.6	9.6	44.0
Non-Hispanic 4/, male												
2006	1.2	*	*	*	0.3	0.4	0.5	0.5	1.0	2.1	8.1	30.2
2005	1.4	*	*	*	0.3	0.5	0.5	0.6	1.0	2.7	9.7	34.1
2004	1.4	*	*	*	0.4	0.5	0.5	0.6	1.2	2.8	9.0	37.1
2003	1.4	*	*	0.1	0.3	0.5	0.5	0.7	1.2	3.1	9.4	38.3
2002	1.4	*	*	0.1	0.4	0.6	0.4	0.6	1.0	2.9	9.0	37.6
2001	1.4	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	10.0	40.8
2000	1.3	*	*	0.1	0.2	0.4	0.4	0.6	1.2	2.8	9.6	41.0
1999	1.4	*	*	0.1	0.4	0.5	0.5	0.7	1.0	2.6	10.1	43.3
Non-Hispanic 4/, female												
2006	1.7	*	*	*	0.2	0.4	0.5	0.6	1.0	2.2	7.2	30.4
2005	2.1	*	*	*	0.3	0.4	0.5	0.7	1.1	2.6	9.0	38.6
2004	2.0	*	*	0.1	0.3	0.4	0.5	0.6	1.0	2.5	8.5	39.1
2003	2.1	*	*	0.1	0.3	0.4	0.4	0.6	0.8	2.3	8.6	42.4
2002	2.1	*	*	0.2	0.2	0.4	0.5	0.7	1.0	2.6	9.3	41.8
2001	2.2	*	*	0.2	0.3	0.4	0.6	0.6	1.0	2.6	9.7	43.1
2000	2.2	*	*	*	0.3	0.4	0.5	0.6	1.0	2.8	9.5	44.1
1999	2.1	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.3	44.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
Non-Hispanic white, both sexes												
2006 5/	1.4	*	*	*	*	*	0.1	0.3	0.7	2.0	7.4	30.0
2005 5/	1.7	*	*	*	*	*	0.1	0.3	0.8	2.3	8.9	37.2
2004 5/	1.6	*	*	*	0.1	0.1	0.2	0.3	0.7	2.5	8.6	38.1
2003 5/	1.6	*	*	*	*	0.1	0.1	0.3	0.6	2.4	8.6	41.0
2002	1.6	*	*	*	0.1	0.1	0.1	0.3	0.7	2.4	8.7	39.9
2001	1.7	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.4	42.1
2000	1.7	*	*	*	*	*	0.1	0.3	0.8	2.5	9.3	42.9
1999	1.6	*	*	*	0.1	*	0.1	0.3	0.6	2.4	9.4	43.2
Non-Hispanic white, male												
2006 5/	1.1	*	*	*	*	*	*	0.2	0.7	2.0	7.9	30.6
2005 5/	1.2	*	*	*	*	*	*	0.3	0.8	2.3	9.2	34.3
2004 5/	1.3	*	*	*	*	*	0.2	0.3	0.8	2.7	8.6	37.3
2003 5/	1.3	*	*	*	*	*	*	0.3	0.7	2.6	9.2	38.1
2002	1.2	*	*	*	*	0.2	*	0.3	0.7	2.5	8.5	36.3
2001	1.3	*	*	*	*	*	*	0.3	0.7	2.6	9.8	41.2
2000	1.2	*	*	*	*	*	0.1	0.3	0.9	2.7	9.3	39.9
1999	1.2	*	*	*	*	*	*	0.3	0.5	2.4	9.9	43.4
Non-Hispanic white, female												
2006 5/	1.7	*	*	*	*	*	*	0.3	0.8	2.0	7.1	29.7
2005 5/	2.1	*	*	*	*	*	*	0.4	0.9	2.3	8.8	38.5
2004 5/	2.0	*	*	*	*	*	*	0.2	0.7	2.3	8.6	38.4
2003 5/	2.0	*	*	*	*	*	*	0.2	0.5	2.1	8.2	42.2
2002	2.1	*	*	*	*	*	*	0.3	0.7	2.4	8.8	41.4
2001	2.1	*	*	*	*	*	0.2	0.3	0.8	2.4	9.2	42.4
2000	2.1	*	*	*	*	*	*	0.3	0.7	2.4	9.4	44.1
1999	2.0	*	*	*	*	*	*	0.3	0.7	2.4	9.1	43.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
Non-Hispanic black, both sexes												
2006 5/	2.4	*	*	*	1.0	2.1	2.6	2.6	3.2	4.2	10.5	39.3
2005 5/	2.7	*	*	*	1.2	2.5	2.6	2.8	3.0	6.3	14.9	43.4
2004 5/	2.7	*	*	0.4	1.5	2.4	2.5	2.8	4.1	4.8	11.8	48.9
2003 5/	2.8	*	*	0.4	1.3	2.2	2.4	3.1	4.3	5.6	14.5	50.3
2002	2.9	*	*	0.4	1.3	2.5	2.5	3.1	3.7	6.1	16.2	52.9
2001	2.8	*	*	0.4	1.2	2.3	2.8	3.1	3.0	5.5	16.1	49.3
2000	2.7	*	*	0.5	1.1	2.4	2.5	3.1	3.5	5.5	13.4	51.8
1999	2.9	*	0.9	0.5	1.5	2.7	2.6	3.2	4.3	5.0	13.4	53.0
Non-Hispanic black, male												
2006 5/	2.1	*	*	*	1.1	2.1	2.5	2.4	3.7	4.1	12.0	30.2
2005 5/	2.5	*	*	*	1.4	2.7	3.0	2.6	2.6	6.6	17.4	38.1
2004 5/	2.6	*	*	*	1.6	2.8	2.9	2.5	4.9	5.0	14.0	43.7
2003 5/	2.8	*	*	*	1.3	2.3	2.6	3.4	5.6	8.0	15.3	50.1
2002	2.7	*	*	*	1.7	2.9	2.5	2.6	4.0	7.5	17.0	58.0
2001	2.6	*	*	*	1.4	3.0	2.5	3.6	3.2	6.8	15.3	42.7
2000	2.4	*	*	*	1.0	2.2	2.1	3.4	3.5	4.6	13.9	56.4
1999	2.7	*	*	*	1.8	2.9	2.8	3.4	5.1	5.6	12.8	44.0
Non-Hispanic black, female												
2006 5/	2.6	*	*	*	0.9	2.0	2.7	2.8	2.8	4.3	9.6	43.0
2005 5/	2.9	*	*	*	1.0	2.3	2.3	2.9	3.4	6.2	13.5	45.6
2004 5/	2.8	*	*	*	1.3	2.0	2.2	3.0	3.3	4.7	10.6	51.0
2003 5/	2.9	*	*	*	1.3	2.1	2.2	2.8	3.3	4.0	14.0	50.3
2002	3.0	*	*	*	1.0	2.0	2.6	3.5	3.4	5.1	15.8	51.0
2001	2.9	*	*	*	1.1	1.6	3.0	2.7	2.9	4.6	16.5	51.8
2000	3.1	*	*	*	1.1	2.5	2.8	2.9	3.5	6.1	13.2	50.2
1999	3.1	*	*	*	1.2	2.4	2.4	3.0	3.6	4.7	13.7	56.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
All origins 3/, both sexes												
2006	24.2	*	*	0.1	0.4	1.7	4.8	13.2	36.2	81.8	166.8	285.2
2005	25.3	*	*	0.1	0.5	1.5	4.7	13.4	37.2	86.8	177.2	312.1
2004	24.9	*	*	0.1	0.4	1.5	4.6	13.4	37.1	87.2	176.9	307.0
2003	25.5	*	*	0.1	0.4	1.6	4.6	13.9	38.5	90.8	181.1	317.5
2002	25.4	*	*	0.1	0.4	1.6	4.8	13.7	37.7	91.4	182.8	320.6
2001	25.1	*	*	0.1	0.4	1.5	4.3	13.6	37.8	91.4	181.4	321.8
2000	24.6	*	*	0.1	0.4	1.6	4.3	13.1	37.8	90.7	179.5	319.7
1999	24.5	*	*	0.1	0.4	1.4	4.3	12.9	38.3	91.8	178.0	317.2
All origins 3/, male												
2006	24.4	*	*	*	0.4	1.8	5.9	16.5	43.8	96.2	194.1	321.0
2005	25.0	*	*	*	0.5	1.8	5.8	16.6	44.9	99.7	202.7	344.4
2004	24.4	*	*	*	0.4	1.7	5.7	16.5	44.0	100.0	202.5	333.9
2003	24.8	*	*	*	0.5	1.9	5.7	16.7	45.0	104.0	207.2	340.7
2002	24.2	*	*	0.1	0.5	1.9	5.9	16.4	44.5	101.4	202.6	348.5
2001	23.5	*	*	0.1	0.4	1.7	5.3	16.4	43.1	101.7	198.9	343.1
2000	22.9	*	*	*	0.5	1.9	5.1	15.3	42.4	99.1	199.6	345.1
1999	22.8	*	*	*	0.4	1.6	5.1	15.6	42.6	99.8	198.2	345.7
All origins 3/, female												
2006	24.0	*	*	*	0.3	1.5	3.7	9.9	29.1	69.7	148.1	268.4
2005	25.7	*	*	0.1	0.4	1.2	3.5	10.3	30.1	75.9	159.9	297.2
2004	25.4	*	*	0.1	0.4	1.3	3.5	10.4	30.7	76.4	159.6	294.9
2003	26.2	*	*	*	0.3	1.4	3.5	11.2	32.4	79.8	163.6	307.2
2002	26.5	*	*	*	0.4	1.3	3.7	11.1	31.4	83.1	169.7	308.4
2001	26.6	*	*	*	0.3	1.3	3.4	11.0	32.9	82.8	169.9	312.9
2000	26.3	*	*	*	0.4	1.2	3.4	11.1	33.6	83.7	166.3	309.4
1999	26.2	*	*	*	0.4	1.3	3.5	10.4	34.3	85.2	164.9	305.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
Hispanic, both sexes												
2006	14.2	*	*	*	*	0.9	2.8	13.2	42.6	118.0	228.9	369.7
2005	15.6	*	*	*	*	0.9	3.4	14.8	49.1	126.4	267.0	404.7
2004	14.5	*	*	*	*	0.8	3.0	13.3	48.3	124.9	258.4	365.8
2003	15.5	*	*	*	*	0.9	3.3	15.5	53.7	131.8	272.1	428.1
2002	15.3	*	*	*	*	0.8	3.5	14.0	52.1	134.7	285.6	437.9
2001	15.3	*	*	*	*	0.7	3.1	14.9	53.2	141.2	279.2	489.6
2000	15.1	*	*	*	*	1.0	3.0	14.0	55.5	147.9	278.5	465.1
1999	15.3	*	*	*	*	0.7	3.2	13.9	56.6	149.7	285.2	493.9
Hispanic, male												
2006	13.7	*	*	*	*	0.8	3.3	16.2	49.9	135.6	257.7	381.2
2005	14.9	*	*	*	*	0.9	4.5	17.5	58.2	143.9	293.0	404.7
2004	13.4	*	*	*	*	0.9	3.4	16.0	56.9	134.2	279.3	325.0
2003	14.4	*	*	*	*	0.8	3.9	18.1	62.4	146.3	297.2	426.4
2002	13.9	*	*	*	*	0.9	4.1	16.7	58.9	143.7	306.9	425.0
2001	13.6	*	*	*	*	0.8	3.9	18.5	58.7	151.4	285.3	422.8
2000	13.8	*	*	*	*	1.4	3.6	15.5	64.7	160.8	299.4	449.4
1999	13.4	*	*	*	*	0.6	3.3	17.2	62.5	155.8	293.0	468.0
Hispanic, female												
2006	14.7	*	*	*	*	1.0	2.1	10.1	35.8	103.7	208.7	363.7
2005	16.3	*	*	*	*	0.8	2.1	12.0	40.7	112.2	248.9	404.7
2004	15.7	*	*	*	*	0.8	2.6	10.6	40.5	117.5	244.0	386.8
2003	16.6	*	*	*	*	0.9	2.7	13.0	45.8	120.3	254.9	428.9
2002	16.7	*	*	*	*	0.7	2.9	11.4	46.0	127.6	270.9	444.3
2001	17.1	*	*	*	*	0.7	2.2	11.4	48.3	133.2	275.0	522.2
2000	16.5	*	*	*	*	*	2.4	12.6	47.4	137.7	264.5	472.8
1999	17.3	*	*	*	*	0.8	3.1	10.7	51.4	144.9	280.0	506.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
Non-Hispanic 4/, both sexes												
2006	25.9	*	*	0.1	0.4	1.8	5.1	13.1	35.5	78.9	162.6	280.5
2005	26.9	*	*	0.1	0.5	1.7	4.9	13.2	36.1	83.7	171.6	307.2
2004	26.5	*	*	0.1	0.4	1.6	4.8	13.3	36.0	84.2	172.0	303.5
2003	27.0	*	*	0.1	0.4	1.8	4.8	13.6	37.1	87.7	175.8	312.0
2002	26.9	*	*	0.1	0.5	1.8	5.0	13.6	36.4	88.2	177.2	315.0
2001	26.4	*	*	0.1	0.4	1.7	4.5	13.4	36.5	87.7	176.4	314.8
2000	25.9	*	*	0.1	0.5	1.7	4.4	13.0	36.4	86.8	174.8	313.4
1999	25.7	*	*	0.1	0.4	1.6	4.4	12.8	36.8	88.1	173.1	310.0
Non-Hispanic 4/, male												
2006	26.3	*	*	*	0.5	2.1	6.3	16.5	43.1	93.0	189.9	317.0
2005	26.8	*	*	*	0.6	2.1	6.0	16.4	43.7	96.3	196.9	340.9
2004	26.2	*	*	*	0.4	1.9	6.1	16.5	42.7	97.3	197.7	333.5
2003	26.4	*	*	*	0.5	2.1	6.0	16.5	43.5	100.9	201.8	335.9
2002	25.8	*	*	0.1	0.5	2.2	6.2	16.3	43.1	98.2	196.6	344.0
2001	25.0	*	*	0.1	0.4	1.9	5.5	16.1	41.8	98.0	194.3	338.9
2000	24.2	*	*	*	0.5	2.0	5.3	15.2	40.7	95.0	194.6	339.7
1999	24.1	*	*	*	0.4	1.8	5.3	15.3	41.0	96.3	193.8	339.5
Non-Hispanic 4/, female												
2006	25.5	*	*	*	0.4	1.6	4.0	9.9	28.4	66.8	144.0	263.5
2005	27.1	*	*	0.1	0.5	1.3	3.8	10.1	29.1	73.0	154.4	291.9
2004	26.8	*	*	0.1	0.4	1.4	3.6	10.3	29.8	73.1	154.7	290.1
2003	27.6	*	*	*	0.3	1.5	3.6	10.9	31.2	76.7	158.5	301.5
2002	27.9	*	*	*	0.4	1.4	3.8	11.1	30.1	79.8	164.3	302.5
2001	27.8	*	*	*	0.3	1.4	3.6	10.9	31.6	79.1	164.6	304.7
2000	27.6	*	*	*	0.4	1.4	3.5	11.0	32.4	80.1	161.8	302.8
1999	27.3	*	*	*	0.4	1.4	3.6	10.4	32.8	81.4	159.6	298.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
Non-Hispanic white, both sexes												
2006 5/	25.4	*	*	*	0.3	1.4	4.4	11.0	30.3	68.9	149.0	264.2
2005 5/	26.5	*	*	0.1	0.4	1.3	4.1	11.1	30.9	73.1	158.4	289.8
2004 5/	26.1	*	*	*	0.3	1.4	4.1	11.3	30.7	73.7	157.8	284.5
2003 5/	26.6	*	*	*	0.3	1.5	4.1	11.3	31.4	76.8	161.6	293.8
2002	26.4	*	*	*	0.4	1.5	4.3	11.3	30.6	76.8	163.3	296.4
2001	26.0	*	*	*	0.3	1.3	3.9	11.3	30.7	77.0	162.7	297.4
2000	25.4	*	*	*	0.4	1.4	3.9	10.7	30.9	76.1	159.9	295.5
1999	25.0	*	*	*	0.4	1.4	3.8	10.5	30.1	76.8	159.1	290.2
Non-Hispanic white, male												
2006 5/	26.3	*	*	*	0.3	1.6	5.4	14.2	37.6	83.9	179.6	306.8
2005 5/	26.8	*	*	*	0.5	1.5	5.2	14.1	38.1	86.5	186.6	331.5
2004 5/	26.3	*	*	*	0.3	1.6	5.3	14.3	37.3	87.7	187.1	321.7
2003 5/	26.6	*	*	*	0.4	1.7	5.2	14.1	37.9	91.3	191.7	327.3
2002	26.0	*	*	*	0.4	1.9	5.5	13.8	37.5	88.9	187.4	338.4
2001	25.1	*	*	*	0.3	1.5	4.8	13.7	36.4	89.8	183.5	332.1
2000	24.3	*	*	*	0.5	1.7	4.8	13.0	35.4	86.6	184.5	330.4
1999	24.0	*	*	*	0.4	1.5	4.6	13.0	35.0	87.7	184.6	326.0
Non-Hispanic white, female												
2006 5/	24.5	*	*	*	0.3	1.2	3.3	7.9	23.3	55.8	127.8	244.3
2005 5/	26.3	*	*	*	0.4	1.0	3.1	8.1	24.1	61.5	139.1	270.7
2004 5/	25.9	*	*	*	0.3	1.2	3.0	8.3	24.3	61.7	137.8	267.8
2003 5/	26.5	*	*	*	0.3	1.3	2.9	8.5	25.2	64.4	141.3	279.0
2002	26.8	*	*	*	0.4	1.2	3.2	8.8	24.1	66.5	147.0	278.3
2001	26.9	*	*	*	0.2	1.2	2.9	8.8	25.3	66.0	148.9	283.0
2000	26.5	*	*	*	0.3	1.1	3.1	8.6	26.6	67.3	143.8	281.4
1999	26.0	*	*	*	0.3	1.2	3.0	8.1	25.4	67.7	142.4	275.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
Non-Hispanic black, both sexes												
2006 5/	33.6	*	*	*	0.8	4.4	10.6	28.0	79.3	170.0	316.4	475.9
2005 5/	34.4	*	*	*	1.2	4.1	9.8	28.5	81.3	180.6	325.0	513.4
2004 5/	34.4	*	*	*	1.0	3.4	9.8	28.8	81.9	182.6	336.4	536.5
2003 5/	35.0	*	*	*	0.7	4.0	9.9	30.8	87.5	188.9	340.0	532.2
2002	34.8	*	*	*	1.0	3.8	9.9	30.9	85.9	191.5	340.2	544.7
2001	34.1	*	*	*	1.0	3.8	9.1	30.1	86.5	190.8	337.9	533.8
2000	34.0	*	*	*	1.0	3.6	8.2	31.0	83.9	190.6	354.8	530.1
1999	34.1	*	*	*	0.8	3.1	9.1	31.0	92.0	194.0	336.1	530.9
Non-Hispanic black, male												
2006 5/	31.7	*	*	*	1.0	5.3	13.4	34.3	92.6	191.3	334.2	468.2
2005 5/	31.8	*	*	*	1.2	5.0	12.0	34.1	96.2	197.4	346.5	471.0
2004 5/	31.3	*	*	*	1.0	4.0	12.2	34.0	93.4	201.9	344.3	508.1
2003 5/	30.9	*	*	*	0.9	4.9	11.9	34.6	97.8	201.2	342.8	457.1
2002	30.0	*	*	*	1.1	4.6	12.1	35.8	96.6	193.3	332.6	423.4
2001	29.4	*	*	*	0.9	5.0	10.5	35.0	93.8	189.5	345.1	452.7
2000	28.3	*	*	*	1.0	4.0	9.7	34.2	89.7	186.9	339.4	456.8
1999	28.6	*	*	*	*	3.4	10.7	35.3	95.4	189.7	317.7	506.3
Non-Hispanic black, female												
2006 5/	35.3	*	*	*	0.7	3.7	8.1	22.6	68.6	154.7	306.2	479.1
2005 5/	36.7	*	*	*	1.1	3.3	7.8	23.7	69.4	168.6	312.9	530.4
2004 5/	37.2	*	*	*	1.0	2.9	7.8	24.3	72.6	168.7	332.0	547.7
2003 5/	38.7	*	*	*	*	3.2	8.1	27.6	79.2	180.2	338.5	561.3
2002	39.2	*	*	*	0.8	3.0	8.0	26.8	77.3	190.3	344.5	591.1
2001	38.4	*	*	*	1.0	2.8	7.8	25.9	80.6	191.7	333.9	563.8
2000	39.1	*	*	*	1.0	3.3	7.0	28.2	79.1	193.3	363.5	557.3
1999	39.0	*	*	*	0.9	2.7	7.7	27.2	89.2	197.0	346.5	540.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
All origins 3/, both sexes												
2006	0.9	*	*	*	*	*	0.1	0.2	0.6	1.5	5.5	22.7
2005	1.1	*	*	*	*	*	0.1	0.3	0.5	1.9	7.0	30.8
2004	1.0	*	*	*	*	*	0.1	0.3	0.6	1.6	6.9	30.3
2003	1.1	*	*	*	*	*	0.1	0.2	0.6	1.9	7.7	35.5
2002	1.3	*	*	*	*	*	0.1	0.3	0.6	2.1	8.9	41.5
2001	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	8.7	42.9
2000	1.5	*	*	*	*	*	0.1	0.3	0.7	2.5	10.4	50.0
1999	1.5	*	*	*	*	*	0.1	0.3	0.7	2.3	10.2	53.9
All origins 3/, male												
2006	0.6	*	*	*	*	*	0.1	0.2	0.5	1.5	5.6	18.7
2005	0.8	*	*	*	*	*	*	0.3	0.6	1.8	6.5	27.9
2004	0.8	*	*	*	*	*	*	0.2	0.6	1.7	6.8	27.4
2003	0.8	*	*	*	*	*	*	0.2	0.6	2.0	7.8	30.6
2002	0.9	*	*	*	*	*	0.1	0.2	0.7	2.2	8.5	37.0
2001	0.9	*	*	*	*	*	0.1	0.3	0.7	2.1	8.5	35.2
2000	1.0	*	*	*	*	*	*	0.2	0.8	2.7	9.7	43.7
1999	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	10.3	48.4
All origins 3/, female												
2006	1.1	*	*	*	*	*	0.1	0.2	0.6	1.4	5.4	24.5
2005	1.4	*	*	*	*	*	0.1	0.3	0.4	1.9	7.4	32.1
2004	1.3	*	*	*	*	*	0.1	0.3	0.5	1.5	7.0	31.7
2003	1.5	*	*	*	*	*	*	0.3	0.6	1.7	7.7	37.6
2002	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	9.1	43.5
2001	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	8.9	46.1
2000	2.0	*	*	*	*	*	0.2	0.3	0.6	2.4	10.9	52.6
1999	2.0	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	56.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
Hispanic, both sexes												
2006	0.3	*	*	*	*	*	*	*	*	*	5.7	19.2
2005	0.3	*	*	*	*	*	*	*	*	1.5	6.9	23.1
2004	0.3	*	*	*	*	*	*	*	*	*	4.7	19.5
2003	0.3	*	*	*	*	*	*	*	*	*	5.6	28.5
2002	0.4	*	*	*	*	*	*	*	*	1.8	8.2	29.4
2001	0.4	*	*	*	*	*	*	*	*	*	8.1	38.8
2000	0.4	*	*	*	*	*	*	*	*	2.2	8.5	39.8
1999	0.4	*	*	*	*	*	*	*	*	*	6.5	51.2
Hispanic, male												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.3	*	*	*	*	*	*	*	*	*	7.0	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	8.9	43.8
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	51.8
Hispanic, female												
2006	0.4	*	*	*	*	*	*	*	*	*	5.5	22.8
2005	0.4	*	*	*	*	*	*	*	*	*	6.7	22.8
2004	0.4	*	*	*	*	*	*	*	*	*	5.2	22.3
2003	0.4	*	*	*	*	*	*	*	*	*	6.1	28.9
2002	0.4	*	*	*	*	*	*	*	*	*	8.4	29.0
2001	0.5	*	*	*	*	*	*	*	*	*	7.6	36.3
2000	0.5	*	*	*	*	*	*	*	*	*	8.6	41.5
1999	0.5	*	*	*	*	*	*	*	*	*	*	51.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
Non-Hispanic 4/, both sexes												
2006	1.0	*	*	*	*	*	0.1	0.2	0.6	1.5	5.5	22.8
2005	1.2	*	*	*	*	*	0.1	0.3	0.5	1.9	7.0	31.1
2004	1.2	*	*	*	*	*	0.1	0.3	0.6	1.6	7.0	30.8
2003	1.3	*	*	*	*	*	0.1	0.3	0.6	1.9	7.8	35.7
2002	1.5	*	*	*	*	*	0.1	0.3	0.7	2.1	8.9	41.8
2001	1.4	*	*	*	*	*	0.1	0.3	0.6	2.1	8.8	43.0
2000	1.7	*	*	*	*	*	0.1	0.3	0.7	2.5	10.4	50.3
1999	1.7	*	*	*	*	*	0.1	0.3	0.7	2.4	10.3	53.9
Non-Hispanic 4/, male												
2006	0.7	*	*	*	*	*	*	0.2	0.6	1.6	5.6	18.9
2005	0.8	*	*	*	*	*	*	0.3	0.6	1.8	6.4	28.0
2004	0.8	*	*	*	*	*	*	0.3	0.6	1.7	7.0	28.0
2003	0.9	*	*	*	*	*	*	0.2	0.6	2.0	7.9	30.7
2002	1.0	*	*	*	*	*	0.1	0.3	0.8	2.2	8.5	37.1
2001	1.0	*	*	*	*	*	*	0.3	0.7	2.2	8.5	34.8
2000	1.1	*	*	*	*	*	*	0.3	0.9	2.7	9.6	43.9
1999	1.2	*	*	*	*	*	0.1	0.3	0.9	2.7	10.4	48.1
Non-Hispanic 4/, female												
2006	1.2	*	*	*	*	*	0.1	0.3	0.7	1.4	5.4	24.6
2005	1.5	*	*	*	*	*	0.1	0.3	0.5	2.0	7.4	32.5
2004	1.4	*	*	*	*	*	0.1	0.3	0.5	1.6	7.0	32.1
2003	1.6	*	*	*	*	*	*	0.3	0.7	1.8	7.7	37.9
2002	1.8	*	*	*	*	*	0.1	0.3	0.6	2.0	9.1	43.9
2001	1.9	*	*	*	*	*	*	0.3	0.6	2.0	8.9	46.4
2000	2.2	*	*	*	*	*	0.2	0.3	0.6	2.4	10.9	52.8
1999	2.2	*	*	*	*	*	0.2	0.3	0.6	2.1	10.3	56.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
Non-Hispanic white, both sexes												
2006 5/	1.0	*	*	*	*	*	0.1	0.2	0.6	1.4	5.3	22.2
2005 5/	1.3	*	*	*	*	*	0.1	0.3	0.5	1.8	6.5	30.3
2004 5/	1.2	*	*	*	*	*	0.1	0.3	0.5	1.5	6.7	29.6
2003 5/	1.4	*	*	*	*	*	0.1	0.2	0.6	1.7	7.4	34.4
2002	1.5	*	*	*	*	*	0.1	0.3	0.6	1.9	8.5	40.6
2001	1.5	*	*	*	*	*	0.1	0.2	0.6	2.0	8.2	42.2
2000	1.8	*	*	*	*	*	0.1	0.2	0.6	2.3	10.0	49.0
1999	1.8	*	*	*	*	*	0.1	0.3	0.6	2.0	9.6	52.8
Non-Hispanic white, male												
2006 5/	0.7	*	*	*	*	*	*	0.2	0.5	1.4	5.1	18.4
2005 5/	0.9	*	*	*	*	*	*	0.3	0.5	1.7	5.8	27.4
2004 5/	0.9	*	*	*	*	*	*	0.2	0.5	1.5	6.5	26.9
2003 5/	0.9	*	*	*	*	*	*	0.2	0.5	1.8	7.2	29.6
2002	1.1	*	*	*	*	*	*	0.2	0.6	2.0	8.0	36.5
2001	1.0	*	*	*	*	*	*	0.2	0.5	1.9	7.9	33.3
2000	1.2	*	*	*	*	*	*	0.2	0.7	2.3	9.0	43.0
1999	1.2	*	*	*	*	*	*	0.2	0.7	2.2	9.4	46.2
Non-Hispanic white, female												
2006 5/	1.3	*	*	*	*	*	0.1	0.2	0.7	1.4	5.4	24.0
2005 5/	1.6	*	*	*	*	*	*	0.3	0.5	1.8	7.0	31.7
2004 5/	1.5	*	*	*	*	*	*	0.3	0.5	1.5	6.8	30.8
2003 5/	1.8	*	*	*	*	*	*	0.2	0.6	1.7	7.6	36.5
2002	2.0	*	*	*	*	*	*	0.3	0.6	1.8	8.8	42.3
2001	2.0	*	*	*	*	*	*	0.3	0.6	2.0	8.3	45.9
2000	2.3	*	*	*	*	*	0.2	0.3	0.6	2.3	10.6	51.5
1999	2.3	*	*	*	*	*	0.2	0.3	0.5	1.9	9.8	55.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
Non-Hispanic black, both sexes												
2006 5/	0.9	*	*	*	*	*	*	0.5	1.1	3.0	8.8	31.3
2005 5/	1.1	*	*	*	*	*	*	0.6	1.1	3.4	12.8	43.1
2004 5/	1.1	*	*	*	*	*	*	0.7	1.2	3.4	10.7	47.2
2003 5/	1.2	*	*	*	*	*	*	0.7	1.4	3.2	13.2	55.9
2002	1.4	*	*	*	*	*	*	0.4	1.8	4.3	14.7	60.2
2001	1.3	*	*	*	*	*	*	0.8	1.2	3.7	16.1	55.2
2000	1.5	*	*	*	*	*	*	0.6	1.7	4.8	16.3	67.4
1999	1.7	*	*	*	*	*	*	0.7	2.0	5.6	19.6	70.8
Non-Hispanic black, male												
2006 5/	0.9	*	*	*	*	*	*	*	1.4	4.1	12.2	29.3
2005 5/	0.9	*	*	*	*	*	*	*	1.7	3.1	12.6	37.1
2004 5/	1.0	*	*	*	*	*	*	*	1.7	4.6	12.3	45.7
2003 5/	1.0	*	*	*	*	*	*	*	1.8	4.5	17.9	49.0
2002	1.1	*	*	*	*	*	*	*	2.9	4.5	16.4	45.9
2001	1.2	*	*	*	*	*	*	1.2	1.9	5.6	16.5	58.9
2000	1.3	*	*	*	*	*	*	*	2.8	6.6	17.9	64.6
1999	1.5	*	*	*	*	*	*	*	2.9	8.3	23.1	76.1
Non-Hispanic black, female												
2006 5/	1.0	*	*	*	*	*	*	*	*	2.3	6.8	32.1
2005 5/	1.4	*	*	*	*	*	*	*	*	3.6	12.9	45.6
2004 5/	1.3	*	*	*	*	*	*	*	*	2.6	9.8	47.8
2003 5/	1.4	*	*	*	*	*	*	*	*	2.4	10.6	58.6
2002	1.6	*	*	*	*	*	*	*	*	4.1	13.8	65.6
2001	1.4	*	*	*	*	*	*	*	*	2.4	15.8	53.9
2000	1.8	*	*	*	*	*	*	*	*	3.6	15.5	68.5
1999	1.8	*	*	*	*	*	*	*	*	3.7	17.7	68.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
All origins 3/, both sexes												
2006	0.8	*	*	*	*	*	0.1	0.2	0.5	1.4	5.2	20.9
2005	1.0	*	*	*	*	*	0.1	0.3	0.5	1.8	6.6	28.9
2004	1.0	*	*	*	*	*	0.1	0.3	0.5	1.6	6.7	28.3
2003	1.1	*	*	*	*	*	0.1	0.2	0.6	1.7	7.4	33.4
2002	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	8.4	38.4
2001	1.2	*	*	*	*	*	0.1	0.3	0.6	1.9	8.2	40.2
2000	1.4	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.3
1999	1.4	*	*	*	*	*	0.1	0.3	0.6	2.2	9.5	50.0
All origins 3/, male												
2006	0.6	*	*	*	*	*	*	0.2	0.5	1.4	5.2	17.5
2005	0.7	*	*	*	*	*	*	0.3	0.6	1.7	6.2	26.3
2004	0.7	*	*	*	*	*	*	0.2	0.6	1.7	6.6	25.3
2003	0.8	*	*	*	*	*	*	0.2	0.6	1.9	7.4	28.2
2002	0.9	*	*	*	*	*	0.1	0.2	0.7	2.1	8.0	35.1
2001	0.8	*	*	*	*	*	*	0.2	0.6	1.9	8.0	33.6
2000	0.9	*	*	*	*	*	*	0.2	0.7	2.5	8.9	40.7
1999	1.0	*	*	*	*	*	0.1	0.3	0.8	2.5	9.4	44.4
All origins 3/, female												
2006	1.0	*	*	*	*	*	0.1	0.2	0.6	1.4	5.1	22.4
2005	1.3	*	*	*	*	*	0.1	0.3	0.4	1.8	6.9	30.2
2004	1.2	*	*	*	*	*	0.1	0.3	0.5	1.5	6.7	29.6
2003	1.4	*	*	*	*	*	*	0.2	0.6	1.7	7.4	35.6
2002	1.6	*	*	*	*	*	0.1	0.2	0.5	1.8	8.6	39.8
2001	1.6	*	*	*	*	*	0.1	0.3	0.5	1.9	8.2	43.0
2000	1.8	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	48.6
1999	1.8	*	*	*	*	*	0.1	0.3	0.5	2.0	9.5	52.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
Hispanic, both sexes												
2006	0.3	*	*	*	*	*	*	*	*	*	5.6	17.6
2005	0.3	*	*	*	*	*	*	*	*	*	6.6	21.8
2004	0.3	*	*	*	*	*	*	*	*	*	4.4	19.5
2003	0.3	*	*	*	*	*	*	*	*	*	4.7	25.9
2002	0.4	*	*	*	*	*	*	*	*	1.8	8.2	28.3
2001	0.4	*	*	*	*	*	*	*	*	*	7.9	38.1
2000	0.4	*	*	*	*	*	*	*	*	2.0	7.7	38.5
1999	0.4	*	*	*	*	*	*	*	*	*	6.2	47.7
Hispanic, male												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.3	*	*	*	*	*	*	*	*	*	6.7	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	8.9	41.9
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	45.3
Hispanic, female												
2006	0.4	*	*	*	*	*	*	*	*	*	5.5	21.0
2005	0.4	*	*	*	*	*	*	*	*	*	6.5	21.5
2004	0.3	*	*	*	*	*	*	*	*	*	*	22.3
2003	0.3	*	*	*	*	*	*	*	*	*	5.3	26.6
2002	0.4	*	*	*	*	*	*	*	*	*	8.4	28.2
2001	0.4	*	*	*	*	*	*	*	*	*	7.3	36.3
2000	0.5	*	*	*	*	*	*	*	*	*	7.6	41.5
1999	0.5	*	*	*	*	*	*	*	*	*	*	48.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
Non-Hispanic 4/, both sexes												
2006	0.9	*	*	*	*	*	0.1	0.2	0.6	1.4	5.1	21.0
2005	1.1	*	*	*	*	*	0.1	0.3	0.5	1.8	6.6	29.2
2004	1.1	*	*	*	*	*	0.1	0.3	0.5	1.6	6.8	28.6
2003	1.2	*	*	*	*	*	0.1	0.2	0.6	1.8	7.5	33.7
2002	1.3	*	*	*	*	*	0.1	0.2	0.6	1.9	8.3	38.7
2001	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	8.2	40.2
2000	1.5	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.5
1999	1.6	*	*	*	*	*	0.1	0.3	0.7	2.2	9.6	50.0
Non-Hispanic 4/, male												
2006	0.6	*	*	*	*	*	*	0.2	0.5	1.5	5.2	17.9
2005	0.8	*	*	*	*	*	*	0.3	0.6	1.7	6.1	26.5
2004	0.8	*	*	*	*	*	*	0.2	0.6	1.7	6.7	25.8
2003	0.9	*	*	*	*	*	*	0.2	0.6	1.8	7.5	28.4
2002	1.0	*	*	*	*	*	0.1	0.2	0.7	2.1	8.0	35.3
2001	0.9	*	*	*	*	*	*	0.3	0.6	2.0	8.0	33.2
2000	1.0	*	*	*	*	*	*	0.2	0.8	2.5	8.8	40.9
1999	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	9.5	44.3
Non-Hispanic 4/, female												
2006	1.1	*	*	*	*	*	0.1	0.2	0.6	1.4	5.1	22.5
2005	1.4	*	*	*	*	*	0.1	0.3	0.4	1.9	6.9	30.5
2004	1.4	*	*	*	*	*	0.1	0.3	0.5	1.5	6.8	29.9
2003	1.5	*	*	*	*	*	*	0.2	0.6	1.7	7.5	36.0
2002	1.7	*	*	*	*	*	0.1	0.3	0.5	1.8	8.6	40.1
2001	1.7	*	*	*	*	*	*	0.3	0.6	1.9	8.3	43.2
2000	2.0	*	*	*	*	*	0.2	0.3	0.6	2.2	10.3	48.7
1999	2.0	*	*	*	*	*	0.2	0.3	0.5	2.0	9.7	52.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
Non-Hispanic white, both sexes												
2006 5/	0.9	*	*	*	*	*	0.1	0.2	0.5	1.3	4.9	20.3
2005 5/	1.2	*	*	*	*	*	0.1	0.2	0.5	1.6	6.1	28.5
2004 5/	1.1	*	*	*	*	*	0.1	0.2	0.5	1.4	6.4	27.3
2003 5/	1.3	*	*	*	*	*	0.1	0.2	0.5	1.6	7.2	32.3
2002	1.4	*	*	*	*	*	0.1	0.2	0.5	1.8	8.0	37.4
2001	1.4	*	*	*	*	*	*	0.2	0.5	1.8	7.6	39.4
2000	1.6	*	*	*	*	*	0.1	0.2	0.6	2.1	9.3	45.3
1999	1.6	*	*	*	*	*	0.1	0.2	0.5	1.9	8.9	48.8
Non-Hispanic white, male												
2006 5/	0.7	*	*	*	*	*	*	0.2	0.4	1.3	4.7	17.2
2005 5/	0.8	*	*	*	*	*	*	0.2	0.5	1.6	5.6	25.9
2004 5/	0.8	*	*	*	*	*	*	0.2	0.5	1.4	6.2	24.6
2003 5/	0.9	*	*	*	*	*	*	0.2	0.4	1.6	6.9	27.3
2002	1.0	*	*	*	*	*	*	0.2	0.5	1.9	7.5	34.5
2001	0.9	*	*	*	*	*	*	0.2	0.5	1.7	7.4	31.7
2000	1.1	*	*	*	*	*	*	0.2	0.6	2.1	8.3	39.8
1999	1.1	*	*	*	*	*	*	0.2	0.6	2.0	8.6	42.0
Non-Hispanic white, female												
2006 5/	1.2	*	*	*	*	*	0.1	0.2	0.6	1.3	5.1	21.8
2005 5/	1.5	*	*	*	*	*	*	0.3	0.4	1.7	6.5	29.7
2004 5/	1.5	*	*	*	*	*	*	0.3	0.5	1.4	6.6	28.5
2003 5/	1.7	*	*	*	*	*	*	0.2	0.6	1.6	7.4	34.5
2002	1.8	*	*	*	*	*	*	0.3	0.5	1.7	8.3	38.6
2001	1.9	*	*	*	*	*	*	0.3	0.6	1.9	7.7	42.7
2000	2.2	*	*	*	*	*	0.1	0.2	0.6	2.0	10.0	47.4
1999	2.2	*	*	*	*	*	0.2	0.3	0.4	1.8	9.2	51.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
Non-Hispanic black, both sexes												
2006 5/	0.9	*	*	*	*	*	*	0.5	1.0	3.0	8.2	30.0
2005 5/	1.1	*	*	*	*	*	*	0.6	1.1	3.3	12.3	41.0
2004 5/	1.1	*	*	*	*	*	*	0.6	1.1	3.3	10.5	46.3
2003 5/	1.2	*	*	*	*	*	*	0.7	1.3	3.1	13.0	54.1
2002	1.3	*	*	*	*	*	*	*	1.7	3.6	14.2	57.2
2001	1.2	*	*	*	*	*	*	0.6	1.1	3.4	15.0	52.1
2000	1.4	*	*	*	*	*	*	0.6	1.6	4.8	15.1	64.2
1999	1.6	*	*	*	*	*	*	0.6	1.8	5.5	18.1	66.0
Non-Hispanic black, male												
2006 5/	0.8	*	*	*	*	*	*	*	*	4.1	11.1	29.3
2005 5/	0.8	*	*	*	*	*	*	*	1.7	3.1	12.1	34.3
2004 5/	0.9	*	*	*	*	*	*	*	1.6	4.4	12.0	43.7
2003 5/	1.0	*	*	*	*	*	*	*	1.8	4.3	17.3	46.9
2002	1.0	*	*	*	*	*	*	*	2.7	4.0	15.2	44.9
2001	1.1	*	*	*	*	*	*	1.0	*	5.0	15.9	56.6
2000	1.2	*	*	*	*	*	*	*	2.5	6.4	17.3	62.2
1999	1.4	*	*	*	*	*	*	*	2.6	8.0	20.6	74.9
Non-Hispanic black, female												
2006 5/	0.9	*	*	*	*	*	*	*	*	2.3	6.5	30.3
2005 5/	1.3	*	*	*	*	*	*	*	*	3.5	12.4	43.7
2004 5/	1.2	*	*	*	*	*	*	*	*	2.5	9.7	47.4
2003 5/	1.4	*	*	*	*	*	*	*	*	2.3	10.6	57.0
2002	1.5	*	*	*	*	*	*	*	*	3.4	13.6	61.9
2001	1.3	*	*	*	*	*	*	*	*	2.3	14.5	50.5
2000	1.6	*	*	*	*	*	*	*	*	3.6	13.9	65.0
1999	1.7	*	*	*	*	*	*	*	*	3.7	16.6	62.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
All origins 3/, both sexes												
2006	0.1	*	*	*	*	*	*	*	*	*	0.4	1.8
2005	0.1	*	*	*	*	*	*	*	*	0.1	0.4	1.8
2004	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
2003	0.1	*	*	*	*	*	*	*	*	0.1	0.3	2.1
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.5	3.1
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.7
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.7
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
All origins 3/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	0.4	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	1.6
2004	0.0	*	*	*	*	*	*	*	*	*	*	2.1
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2002	0.1	*	*	*	*	*	*	*	*	*	0.6	1.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	1.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	3.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.9	3.9
All origins 3/, female												
2006	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
2005	0.1	*	*	*	*	*	*	*	*	*	0.5	1.9
2004	0.1	*	*	*	*	*	*	*	*	*	*	2.1
2003	0.1	*	*	*	*	*	*	*	*	*	*	2.0
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	3.1
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.7	4.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	3.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
Non-Hispanic 4/, both sexes												
2006	0.1	*	*	*	*	*	*	*	*	*	0.4	1.8
2005	0.1	*	*	*	*	*	*	*	*	*	0.4	1.9
2004	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2003	0.1	*	*	*	*	*	*	*	*	0.1	0.3	2.1
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.5	3.2
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.6	2.7
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.8
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
Non-Hispanic 4/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	0.4	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	1.6
2004	0.0	*	*	*	*	*	*	*	*	*	*	2.2
2003	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2002	0.1	*	*	*	*	*	*	*	*	*	0.6	1.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	1.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	3.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.9	3.8
Non-Hispanic 4/, female												
2006	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
2005	0.1	*	*	*	*	*	*	*	*	*	0.5	2.0
2004	0.1	*	*	*	*	*	*	*	*	*	*	2.1
2003	0.1	*	*	*	*	*	*	*	*	*	*	1.9
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	3.2
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.7	4.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
Non-Hispanic white, both sexes												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	0.4	1.9
2005 5/	0.1	*	*	*	*	*	*	*	*	*	0.4	1.8
2004 5/	0.1	*	*	*	*	*	*	*	*	*	0.3	2.3
2003 5/	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.2
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.8
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.6	3.8
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
Non-Hispanic white, male												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	1.5
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	2.0
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	3.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.8	4.2
Non-Hispanic white, female												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2005 5/	0.1	*	*	*	*	*	*	*	*	*	0.5	2.0
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	2.0
2002	0.2	*	*	*	*	*	*	*	*	*	0.5	3.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	3.3
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.6	4.0
1999	0.2	*	*	*	*	*	*	*	*	*	0.6	3.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
All origins 3/, both sexes												
2006	0.2	1.5	0.1	*	0.1	0.1	0.2	0.2	0.2	0.4	0.7	0.9
2005	0.2	1.4	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.4	0.6	0.9
2004	0.2	1.6	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.7	0.9
2003	0.3	1.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.5	0.7	1.1
2002	0.2	1.8	*	0.1	0.1	0.1	0.1	0.3	0.3	0.6	0.7	1.2
2001	0.3	1.9	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.8	0.9
2000	0.3	2.4	0.2	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999	0.3	3.1	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.5
All origins 3/, male												
2006	0.2	1.4	*	*	*	0.1	0.2	0.3	0.3	0.4	0.7	*
2005	0.2	1.4	0.3	*	*	0.1	0.2	0.3	0.4	0.4	0.6	*
2004	0.2	1.5	*	*	*	0.1	0.2	0.4	0.3	0.4	0.7	*
2003	0.3	1.9	*	*	0.1	0.1	0.2	0.4	0.5	0.5	0.8	*
2002	0.3	2.0	*	*	*	*	0.2	0.4	0.4	0.7	0.8	*
2001	0.3	1.9	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000	0.3	2.5	0.3	*	*	0.1	0.2	0.3	0.5	0.7	1.3	*
1999	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.0	1.7
All origins 3/, female												
2006	0.2	1.6	*	*	*	*	0.2	0.2	0.2	0.4	0.6	0.8
2005	0.2	1.4	*	*	*	*	0.1	0.2	0.3	0.4	0.6	0.9
2004	0.2	1.7	*	*	*	*	0.1	0.2	0.3	0.6	0.6	0.7
2003	0.2	2.0	*	*	*	*	0.1	0.2	0.3	0.5	0.6	1.0
2002	0.2	1.7	*	*	*	*	*	0.2	0.3	0.5	0.6	1.1
2001	0.2	2.0	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.8
2000	0.2	2.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.0
1999	0.3	3.0	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
Hispanic, both sexes												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
Non-Hispanic 4/, both sexes												
2006	0.2	1.6	*	*	*	0.1	0.2	0.3	0.2	0.4	0.7	0.9
2005	0.2	1.5	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.4	0.6	0.9
2004	0.2	1.8	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.7	0.9
2003	0.3	1.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.5	0.7	1.1
2002	0.3	2.1	*	0.1	0.1	0.1	0.1	0.3	0.3	0.6	0.6	1.2
2001	0.3	2.2	0.2	0.1	0.1	0.1	0.2	0.3	0.5	0.5	0.7	0.9
2000	0.3	2.7	0.3	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999	0.3	3.2	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.4
Non-Hispanic 4/, male												
2006	0.2	1.7	*	*	*	*	0.2	0.3	0.3	0.4	0.8	*
2005	0.3	1.5	*	*	*	0.1	0.2	0.3	0.4	0.4	0.6	*
2004	0.3	1.8	*	*	*	*	0.2	0.4	0.3	0.4	0.7	*
2003	0.3	1.6	*	*	*	0.1	0.2	0.4	0.5	0.6	0.8	*
2002	0.3	2.1	*	*	*	*	0.2	0.4	0.4	0.7	0.8	*
2001	0.3	2.1	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000	0.3	2.7	*	*	*	*	0.2	0.3	0.5	0.8	1.2	*
1999	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.1	*
Non-Hispanic 4/, female												
2006	0.2	1.6	*	*	*	*	0.2	0.2	0.2	0.4	0.6	0.8
2005	0.2	1.5	*	*	*	*	0.2	0.2	0.3	0.4	0.6	0.8
2004	0.2	1.9	*	*	*	*	*	0.2	0.3	0.6	0.6	0.7
2003	0.2	2.2	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.0
2002	0.2	2.0	*	*	*	*	*	0.2	0.3	0.5	0.5	1.2
2001	0.2	2.3	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.7
2000	0.3	2.7	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.1
1999	0.3	3.3	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
Non-Hispanic white, both sexes												
2006 5/	0.2	1.5	*	*	*	0.1	0.1	0.3	0.2	0.4	0.6	0.9
2005 5/	0.2	1.1	*	*	*	*	0.1	0.2	0.3	0.4	0.5	0.8
2004 5/	0.2	1.6	*	*	*	*	0.1	0.2	0.3	0.4	0.6	0.8
2003 5/	0.2	1.5	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.0
2002	0.2	1.9	*	0.1	*	*	0.1	0.2	0.2	0.5	0.5	1.2
2001	0.2	1.6	*	*	0.1	0.1	0.1	0.2	0.4	0.5	0.7	0.7
2000	0.2	2.1	*	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.1
1999	0.3	2.2	0.3	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.3
Non-Hispanic white, male												
2006 5/	0.2	*	*	*	*	*	0.1	0.3	0.2	0.4	0.8	*
2005 5/	0.2	*	*	*	*	*	0.2	0.3	0.3	0.4	0.6	*
2004 5/	0.2	1.9	*	*	*	*	0.2	0.3	0.2	0.3	0.7	*
2003 5/	0.3	1.8	*	*	*	*	0.1	0.3	0.4	0.5	0.8	*
2002	0.2	1.9	*	*	*	*	0.1	0.2	0.3	0.7	0.7	*
2001	0.3	*	*	*	*	*	*	0.3	0.4	0.5	0.8	*
2000	0.3	2.3	*	*	*	*	0.1	0.3	0.4	0.7	1.1	*
1999	0.3	2.8	*	*	*	*	0.1	0.3	0.4	0.6	0.9	*
Non-Hispanic white, female												
2006 5/	0.2	*	*	*	*	*	0.1	0.2	0.2	0.4	0.5	0.8
2005 5/	0.2	*	*	*	*	*	*	0.2	0.3	0.3	0.5	0.8
2004 5/	0.2	*	*	*	*	*	*	*	0.3	0.5	0.5	*
2003 5/	0.2	*	*	*	*	*	*	0.2	0.2	0.5	0.6	0.8
2002	0.2	1.8	*	*	*	*	*	0.2	0.2	0.4	0.4	1.3
2001	0.2	1.9	*	*	*	*	*	0.2	0.3	0.4	0.6	*
2000	0.2	1.8	*	*	*	*	*	0.2	0.2	0.3	0.8	1.0
1999	0.3	*	*	*	*	*	0.1	0.2	0.3	0.5	0.8	1.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
Non-Hispanic black, both sexes												
2006 5/	0.3	*	*	*	*	*	0.5	0.4	0.7	*	*	*
2005 5/	0.4	*	*	*	*	*	0.4	0.6	*	*	*	*
2004 5/	0.4	*	*	*	*	*	*	0.8	0.7	*	*	*
2003 5/	0.5	3.5	*	*	*	*	0.4	0.7	1.3	*	*	*
2002	0.5	3.1	*	*	*	*	*	0.9	0.9	1.3	*	*
2001	0.5	4.7	*	*	*	*	0.5	0.7	1.0	*	*	*
2000	0.6	5.9	*	*	*	*	0.5	0.6	0.9	1.7	*	*
1999	0.6	7.0	*	*	*	*	0.4	0.9	*	1.3	2.7	*
Non-Hispanic black, male												
2006 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.4	*	*	*	*	*	*	0.9	*	*	*	*
2003 5/	0.5	*	*	*	*	*	*	*	*	*	*	*
2002	0.6	*	*	*	*	*	*	1.3	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*	*	*
1999	0.7	*	*	*	*	*	*	1.5	*	*	*	*
Non-Hispanic black, female												
2006 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.5	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	9.6	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
All origins 3/, both sexes												
2006	6.5	*	*	*	*	*	*	0.2	1.3	12.2	69.8	144.8
2005	6.6	*	*	*	*	*	*	0.2	1.4	13.0	71.2	143.7
2004	6.1	*	*	*	*	*	*	0.2	1.2	12.0	67.5	135.8
2003	6.2	*	*	*	*	*	*	0.2	1.3	12.7	67.8	138.2
2002	5.9	*	*	*	*	*	*	0.1	1.2	12.2	63.9	135.2
2001	5.8	*	*	*	*	*	*	0.1	1.2	11.7	64.6	134.2
2000	5.6	*	*	*	*	*	*	0.1	1.1	11.5	61.9	131.9
1999	5.2	*	*	*	*	*	*	0.1	1.0	11.0	58.2	124.4
All origins 3/, male												
2006	7.7	*	*	*	*	*	*	0.2	1.7	17.6	104.7	231.7
2005	7.7	*	*	*	*	*	*	0.2	1.9	18.7	107.2	228.6
2004	7.2	*	*	*	*	*	*	0.2	1.6	17.3	101.7	220.0
2003	7.1	*	*	*	*	*	*	0.2	1.6	18.2	100.7	222.4
2002	6.8	*	*	*	*	*	*	0.2	1.5	17.9	94.7	220.1
2001	6.7	*	*	*	*	*	*	0.2	1.8	16.8	97.0	224.4
2000	6.4	*	*	*	*	*	*	0.2	1.5	16.1	92.8	222.3
1999	6.0	*	*	*	*	*	*	0.2	1.4	15.7	89.5	205.6
All origins 3/, female												
2006	5.4	*	*	*	*	*	*	0.1	0.8	7.7	45.9	104.1
2005	5.5	*	*	*	*	*	*	0.1	1.0	8.1	46.8	104.7
2004	5.1	*	*	*	*	*	*	*	0.8	7.5	44.4	97.9
2003	5.3	*	*	*	*	*	*	0.1	0.9	8.1	45.8	101.0
2002	5.0	*	*	*	*	*	*	*	0.9	7.4	43.5	98.4
2001	4.9	*	*	*	*	*	*	*	0.7	7.5	43.3	96.4
2000	4.8	*	*	*	*	*	*	0.1	0.8	7.6	41.7	95.0
1999	4.5	*	*	*	*	*	*	*	0.7	7.1	37.8	91.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
Hispanic, both sexes												
2006	1.6	*	*	*	*	*	*	*	*	8.9	41.5	93.9
2005	1.7	*	*	*	*	*	*	*	0.9	9.0	48.5	95.0
2004	1.4	*	*	*	*	*	*	*	1.0	6.8	44.1	85.1
2003	1.3	*	*	*	*	*	*	*	*	7.9	40.3	77.7
2002	1.3	*	*	*	*	*	*	*	1.1	8.6	37.2	76.5
2001	1.3	*	*	*	*	*	*	*	*	8.6	41.5	86.3
2000	1.2	*	*	*	*	*	*	*	*	7.8	37.7	86.3
1999	1.1	*	*	*	*	*	*	*	*	8.1	38.3	77.2
Hispanic, male												
2006	1.8	*	*	*	*	*	*	*	*	11.7	63.9	151.1
2005	1.9	*	*	*	*	*	*	*	*	13.5	71.2	138.2
2004	1.6	*	*	*	*	*	*	*	*	9.5	66.5	126.1
2003	1.5	*	*	*	*	*	*	*	*	12.4	58.3	129.3
2002	1.4	*	*	*	*	*	*	*	*	13.6	50.1	113.8
2001	1.4	*	*	*	*	*	*	*	*	11.2	63.0	127.6
2000	1.3	*	*	*	*	*	*	*	*	9.7	55.1	143.1
1999	1.3	*	*	*	*	*	*	*	*	11.4	58.9	105.7
Hispanic, female												
2006	1.3	*	*	*	*	*	*	*	*	6.6	25.9	63.5
2005	1.5	*	*	*	*	*	*	*	*	5.3	32.8	72.3
2004	1.2	*	*	*	*	*	*	*	*	4.7	28.6	64.1
2003	1.1	*	*	*	*	*	*	*	*	4.3	27.9	51.6
2002	1.1	*	*	*	*	*	*	*	*	4.7	28.4	58.0
2001	1.2	*	*	*	*	*	*	*	*	6.6	26.9	66.1
2000	1.1	*	*	*	*	*	*	*	*	6.3	26.1	58.4
1999	1.0	*	*	*	*	*	*	*	*	5.5	24.6	63.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
Non-Hispanic 4/, both sexes												
2006	7.4	*	*	*	*	*	*	0.2	1.3	12.5	71.5	147.2
2005	7.4	*	*	*	*	*	*	0.2	1.5	13.2	72.5	145.9
2004	6.9	*	*	*	*	*	*	0.2	1.2	12.4	68.6	137.9
2003	7.0	*	*	*	*	*	*	0.2	1.3	13.0	69.1	140.6
2002	6.6	*	*	*	*	*	*	0.1	1.2	12.4	65.1	137.4
2001	6.5	*	*	*	*	*	*	0.1	1.2	11.9	65.5	135.6
2000	6.2	*	*	*	*	*	*	0.1	1.1	11.7	62.8	133.3
1999	5.8	*	*	*	*	*	*	0.1	1.1	11.1	58.9	125.8
Non-Hispanic 4/, male												
2006	8.7	*	*	*	*	*	*	0.2	1.8	18.0	107.2	235.9
2005	8.7	*	*	*	*	*	*	0.2	2.0	19.1	109.3	233.1
2004	8.1	*	*	*	*	*	*	0.3	1.6	17.8	103.4	224.4
2003	8.0	*	*	*	*	*	*	0.2	1.7	18.5	102.7	226.5
2002	7.6	*	*	*	*	*	*	0.2	1.5	18.2	96.7	224.4
2001	7.5	*	*	*	*	*	*	0.2	1.8	17.1	98.3	227.8
2000	7.1	*	*	*	*	*	*	0.2	1.5	16.5	94.2	225.2
1999	6.7	*	*	*	*	*	*	0.2	1.5	15.9	90.6	209.4
Non-Hispanic 4/, female												
2006	6.1	*	*	*	*	*	*	0.1	0.8	7.8	47.1	106.0
2005	6.1	*	*	*	*	*	*	*	1.0	8.3	47.5	106.1
2004	5.7	*	*	*	*	*	*	*	0.8	7.8	45.3	99.2
2003	5.9	*	*	*	*	*	*	0.2	0.9	8.4	46.6	102.8
2002	5.6	*	*	*	*	*	*	*	0.9	7.6	44.2	99.8
2001	5.4	*	*	*	*	*	*	*	0.7	7.6	44.0	97.3
2000	5.3	*	*	*	*	*	*	*	0.8	7.7	42.3	96.1
1999	4.9	*	*	*	*	*	*	*	0.7	7.1	38.3	92.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
Non-Hispanic white, both sexes												
2006 5/	8.9	*	*	*	*	*	*	0.1	1.4	13.3	76.8	157.0
2005 5/	8.9	*	*	*	*	*	*	0.2	1.6	14.2	77.4	155.5
2004 5/	8.2	*	*	*	*	*	*	0.2	1.3	13.5	73.3	145.5
2003 5/	8.3	*	*	*	*	*	*	0.2	1.3	14.0	73.7	148.8
2002	7.9	*	*	*	*	*	*	0.2	1.3	13.3	69.6	145.8
2001	7.7	*	*	*	*	*	*	0.1	1.3	12.8	69.9	143.5
2000	7.3	*	*	*	*	*	*	0.1	1.2	12.6	66.7	141.2
1999	6.8	*	*	*	*	*	*	0.1	1.1	11.9	62.4	132.8
Non-Hispanic white, male												
2006 5/	10.4	*	*	*	*	*	*	0.2	1.9	19.1	114.0	252.4
2005 5/	10.4	*	*	*	*	*	*	0.2	2.1	20.3	116.1	247.9
2004 5/	9.7	*	*	*	*	*	*	0.3	1.7	19.2	110.0	236.3
2003 5/	9.5	*	*	*	*	*	*	0.2	1.7	19.8	109.0	237.9
2002	9.1	*	*	*	*	*	*	0.2	1.6	19.3	102.9	237.8
2001	9.0	*	*	*	*	*	*	0.2	2.0	18.3	104.5	241.7
2000	8.4	*	*	*	*	*	*	0.2	1.6	17.6	99.2	238.6
1999	7.9	*	*	*	*	*	*	0.2	1.6	17.1	95.7	221.0
Non-Hispanic white, female												
2006 5/	7.4	*	*	*	*	*	*	*	0.9	8.4	51.0	112.5
2005 5/	7.4	*	*	*	*	*	*	*	1.1	9.0	50.7	113.2
2004 5/	6.8	*	*	*	*	*	*	*	0.9	8.5	48.3	104.8
2003 5/	7.1	*	*	*	*	*	*	0.2	1.0	9.0	49.8	109.6
2002	6.7	*	*	*	*	*	*	*	1.0	8.1	47.3	106.0
2001	6.5	*	*	*	*	*	*	*	0.7	8.2	46.9	102.7
2000	6.3	*	*	*	*	*	*	*	0.9	8.3	45.3	101.9
1999	5.8	*	*	*	*	*	*	*	0.7	7.6	40.5	97.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
Non-Hispanic black, both sexes												
2006 5/	1.8	*	*	*	*	*	*	*	1.0	7.5	30.6	51.8
2005 5/	1.9	*	*	*	*	*	*	*	0.8	7.3	34.4	57.3
2004 5/	1.7	*	*	*	*	*	*	*	0.7	5.5	29.8	61.6
2003 5/	1.7	*	*	*	*	*	*	*	1.2	7.2	29.4	53.5
2002	1.5	*	*	*	*	*	*	*	*	6.7	25.7	51.7
2001	1.5	*	*	*	*	*	*	*	0.8	5.6	25.7	59.0
2000	1.5	*	*	*	*	*	*	*	*	5.8	27.3	52.5
1999	1.4	*	*	*	*	*	*	*	0.9	5.9	24.8	51.7
Non-Hispanic black, male												
2006 5/	2.1	*	*	*	*	*	*	*	*	12.1	51.4	71.5
2005 5/	2.3	*	*	*	*	*	*	*	*	12.6	54.9	99.0
2004 5/	1.9	*	*	*	*	*	*	*	*	8.6	44.5	106.7
2003 5/	2.0	*	*	*	*	*	*	*	1.8	10.6	45.6	99.1
2002	1.8	*	*	*	*	*	*	*	*	11.1	40.4	87.5
2001	1.7	*	*	*	*	*	*	*	*	8.8	38.1	99.3
2000	1.8	*	*	*	*	*	*	*	*	9.0	47.5	88.1
1999	1.6	*	*	*	*	*	*	*	*	8.7	38.5	87.9
Non-Hispanic black, female												
2006 5/	1.5	*	*	*	*	*	*	*	*	4.3	18.8	43.7
2005 5/	1.5	*	*	*	*	*	*	*	*	3.6	22.7	40.6
2004 5/	1.5	*	*	*	*	*	*	*	*	3.3	21.6	43.8
2003 5/	1.4	*	*	*	*	*	*	*	*	4.8	20.2	35.9
2002	1.3	*	*	*	*	*	*	*	*	3.6	17.4	38.0
2001	1.4	*	*	*	*	*	*	*	*	3.3	18.7	44.1
2000	1.2	*	*	*	*	*	*	*	*	3.5	16.0	39.2
1999	1.3	*	*	*	*	*	*	*	*	3.9	17.2	38.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
All origins 3/, both sexes												
2006	24.2	*	*	*	*	*	*	0.2	2.1	20.2	175.6	848.3
2005	24.2	*	*	*	*	*	*	0.2	2.1	20.5	177.3	861.6
2004	22.5	*	*	*	*	*	*	0.2	1.9	19.7	168.7	818.8
2003	21.8	*	*	*	*	*	*	0.2	2.0	20.9	164.4	802.4
2002	20.4	*	*	*	*	*	*	0.1	1.9	19.7	158.1	752.3
2001	18.9	*	*	*	*	*	*	0.2	2.1	18.7	147.5	710.3
2000	17.6	*	*	*	*	*	*	0.2	2.0	18.7	139.6	667.7
1999	16.0	*	*	*	*	*	*	0.2	1.9	17.4	129.5	601.3
All origins 3/, male												
2006	14.3	*	*	*	*	*	*	0.2	2.0	19.7	156.6	638.6
2005	14.1	*	*	*	*	*	*	0.1	2.0	19.3	157.7	639.9
2004	13.1	*	*	*	*	*	*	0.2	1.8	18.5	149.0	619.6
2003	12.8	*	*	*	*	*	*	0.2	1.8	20.2	147.6	606.5
2002	12.0	*	*	*	*	*	*	0.1	1.8	18.6	142.2	572.0
2001	11.3	*	*	*	*	*	*	0.2	1.9	17.7	136.3	556.1
2000	10.5	*	*	*	*	*	*	0.2	1.8	18.7	127.6	522.1
1999	9.8	*	*	*	*	*	*	0.2	1.9	17.3	122.6	485.7
All origins 3/, female												
2006	33.8	*	*	*	*	*	*	0.2	2.1	20.5	188.7	946.4
2005	33.9	*	*	*	*	*	*	0.2	2.3	21.4	190.5	963.4
2004	31.5	*	*	*	*	*	*	0.3	1.9	20.8	181.9	908.4
2003	30.5	*	*	*	*	*	*	0.2	2.1	21.5	175.7	889.0
2002	28.5	*	*	*	*	*	*	0.1	2.0	20.6	168.7	830.5
2001	26.3	*	*	*	*	*	*	0.2	2.3	19.5	154.8	774.9
2000	24.5	*	*	*	*	*	*	0.2	2.2	18.6	147.4	727.0
1999	21.9	*	*	*	*	*	*	0.2	1.8	17.5	134.0	648.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
Hispanic, both sexes												
2006	5.4	*	*	*	*	*	*	*	1.9	12.5	110.4	514.5
2005	5.2	*	*	*	*	*	*	*	1.3	13.0	109.8	507.8
2004	4.6	*	*	*	*	*	*	*	1.0	12.0	99.5	490.9
2003	4.6	*	*	*	*	*	*	*	1.3	14.6	103.7	493.4
2002	3.8	*	*	*	*	*	*	*	1.0	10.8	89.2	438.5
2001	3.4	*	*	*	*	*	*	*	*	9.9	83.9	399.5
2000	3.2	*	*	*	*	*	*	*	*	11.9	79.2	380.2
1999	2.9	*	*	*	*	*	*	*	*	9.7	75.4	351.5
Hispanic, male												
2006	3.5	*	*	*	*	*	*	*	2.1	13.1	107.2	404.8
2005	3.5	*	*	*	*	*	*	*	*	13.9	112.6	412.2
2004	2.8	*	*	*	*	*	*	*	*	11.5	93.0	374.1
2003	3.1	*	*	*	*	*	*	*	*	13.8	101.2	426.4
2002	2.3	*	*	*	*	*	*	*	*	10.5	80.4	341.3
2001	2.1	*	*	*	*	*	*	*	*	8.8	83.2	306.6
2000	2.1	*	*	*	*	*	*	*	*	13.3	81.6	284.2
1999	1.8	*	*	*	*	*	*	*	*	11.2	66.7	278.2
Hispanic, female												
2006	7.4	*	*	*	*	*	*	*	1.6	12.0	112.6	572.8
2005	7.0	*	*	*	*	*	*	*	1.6	12.3	107.9	557.9
2004	6.5	*	*	*	*	*	*	*	*	12.5	104.0	551.0
2003	6.2	*	*	*	*	*	*	*	*	15.3	105.3	527.3
2002	5.4	*	*	*	*	*	*	*	*	10.9	95.3	486.6
2001	4.7	*	*	*	*	*	*	*	*	10.7	84.3	444.9
2000	4.3	*	*	*	*	*	*	*	*	10.8	77.6	427.3
1999	4.0	*	*	*	*	*	*	*	*	8.4	81.2	386.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
Non-Hispanic 4/, both sexes												
2006	27.4	*	*	*	*	*	*	0.2	2.1	20.7	179.5	864.3
2005	27.3	*	*	*	*	*	*	0.2	2.2	21.0	181.0	877.5
2004	25.4	*	*	*	*	*	*	0.2	1.9	20.2	172.3	832.6
2003	24.5	*	*	*	*	*	*	0.2	2.0	21.3	167.3	814.0
2002	22.9	*	*	*	*	*	*	0.1	2.0	20.3	161.1	763.2
2001	21.2	*	*	*	*	*	*	0.2	2.2	19.2	150.2	720.5
2000	19.6	*	*	*	*	*	*	0.2	2.1	19.0	142.0	676.7
1999	17.7	*	*	*	*	*	*	0.2	2.0	17.8	131.5	608.9
Non-Hispanic 4/, male												
2006	16.3	*	*	*	*	*	*	0.3	2.0	20.2	159.5	651.1
2005	15.9	*	*	*	*	*	*	0.2	2.0	19.7	160.2	650.9
2004	14.9	*	*	*	*	*	*	0.2	1.9	19.0	152.0	631.2
2003	14.4	*	*	*	*	*	*	0.2	1.9	20.5	149.8	613.6
2002	13.6	*	*	*	*	*	*	0.2	1.9	19.1	145.0	581.4
2001	12.7	*	*	*	*	*	*	0.2	2.0	18.3	138.6	565.2
2000	11.7	*	*	*	*	*	*	0.2	1.8	19.0	129.6	531.0
1999	10.9	*	*	*	*	*	*	0.2	2.0	17.7	124.8	492.8
Non-Hispanic 4/, female												
2006	38.0	*	*	*	*	*	*	0.2	2.2	21.2	193.2	963.4
2005	38.2	*	*	*	*	*	*	0.2	2.3	22.1	195.2	980.9
2004	35.3	*	*	*	*	*	*	0.3	2.0	21.3	186.0	922.7
2003	34.1	*	*	*	*	*	*	0.2	2.2	22.0	179.0	902.1
2002	31.9	*	*	*	*	*	*	0.1	2.1	21.2	171.8	841.6
2001	29.3	*	*	*	*	*	*	0.2	2.4	20.0	157.8	785.1
2000	27.2	*	*	*	*	*	*	0.2	2.3	19.1	150.0	735.6
1999	24.2	*	*	*	*	*	*	0.2	1.9	18.0	135.9	655.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
Non-Hispanic white, both sexes												
2006 5/	32.2	*	*	*	*	*	*	0.3	2.3	21.7	187.1	896.5
2005 5/	31.9	*	*	*	*	*	*	0.2	2.4	21.7	187.5	909.1
2004 5/	29.6	*	*	*	*	*	*	0.3	2.1	21.1	178.1	861.9
2003 5/	28.7	*	*	*	*	*	*	0.2	2.2	22.2	173.9	846.9
2002	26.9	*	*	*	*	*	*	0.2	2.1	21.3	168.2	793.9
2001	24.8	*	*	*	*	*	*	0.2	2.4	20.3	155.6	749.2
2000	22.9	*	*	*	*	*	*	0.2	2.2	19.9	147.4	705.1
1999	20.7	*	*	*	*	*	*	0.2	2.1	18.6	136.7	636.6
Non-Hispanic white, male												
2006 5/	19.1	*	*	*	*	*	*	0.3	2.1	21.0	165.9	676.4
2005 5/	18.5	*	*	*	*	*	*	0.2	2.2	20.1	164.7	673.6
2004 5/	17.3	*	*	*	*	*	*	0.2	2.0	19.6	157.0	653.3
2003 5/	16.8	*	*	*	*	*	*	0.2	2.0	21.2	154.8	638.1
2002	15.8	*	*	*	*	*	*	0.2	2.0	19.9	150.5	605.4
2001	14.7	*	*	*	*	*	*	0.2	2.1	19.1	143.0	587.2
2000	13.6	*	*	*	*	*	*	0.2	2.0	19.8	134.6	551.7
1999	12.7	*	*	*	*	*	*	0.2	2.2	18.3	129.8	515.1
Non-Hispanic white, female												
2006 5/	44.8	*	*	*	*	*	*	0.2	2.4	22.4	201.8	999.2
2005 5/	44.8	*	*	*	*	*	*	0.3	2.5	23.0	203.3	1,016.8
2004 5/	41.4	*	*	*	*	*	*	0.4	2.2	22.3	192.5	955.3
2003 5/	40.2	*	*	*	*	*	*	0.2	2.5	23.0	186.8	938.8
2002	37.6	*	*	*	*	*	*	0.2	2.3	22.5	180.1	875.3
2001	34.4	*	*	*	*	*	*	0.2	2.7	21.3	163.9	816.6
2000	31.9	*	*	*	*	*	*	0.2	2.5	20.0	155.8	767.0
1999	28.4	*	*	*	*	*	*	0.2	2.0	18.8	141.2	685.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
Non-Hispanic black, both sexes												
2006 5/	11.7	*	*	*	*	*	*	*	1.7	19.1	149.3	674.5
2005 5/	12.3	*	*	*	*	*	*	*	1.5	22.7	160.1	698.8
2004 5/	11.4	*	*	*	*	*	*	*	1.3	19.4	155.4	666.9
2003 5/	10.1	*	*	*	*	*	*	*	1.1	19.6	140.3	591.1
2002	9.1	*	*	*	*	*	*	*	1.2	16.8	124.2	548.7
2001	8.7	*	*	*	*	*	*	*	1.1	15.5	126.9	513.2
2000	7.7	*	*	*	*	*	*	*	1.3	16.1	111.6	450.9
1999	6.7	*	*	*	*	*	*	*	1.2	15.5	99.1	374.7
Non-Hispanic black, male												
2006 5/	6.5	*	*	*	*	*	*	*	1.7	19.3	134.9	478.3
2005 5/	7.2	*	*	*	*	*	*	*	*	22.8	152.8	527.2
2004 5/	6.5	*	*	*	*	*	*	*	*	19.4	138.6	507.0
2003 5/	6.2	*	*	*	*	*	*	*	*	20.5	138.8	465.6
2002	5.5	*	*	*	*	*	*	*	*	17.9	122.5	414.6
2001	5.3	*	*	*	*	*	*	*	*	16.3	123.6	420.4
2000	4.6	*	*	*	*	*	*	*	*	16.9	98.7	378.1
1999	4.2	*	*	*	*	*	*	*	*	16.8	90.7	320.9
Non-Hispanic black, female												
2006 5/	16.4	*	*	*	*	*	*	*	1.6	18.9	157.5	754.5
2005 5/	16.9	*	*	*	*	*	*	*	1.7	22.7	164.3	767.9
2004 5/	15.8	*	*	*	*	*	*	*	1.4	19.4	164.9	730.0
2003 5/	13.7	*	*	*	*	*	*	*	*	18.9	141.1	639.7
2002	12.5	*	*	*	*	*	*	*	*	16.1	125.2	599.9
2001	11.8	*	*	*	*	*	*	*	*	15.0	128.7	547.5
2000	10.6	*	*	*	*	*	*	*	*	15.6	118.8	478.0
1999	9.0	*	*	*	*	*	*	*	*	14.5	103.8	394.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
All origins 3/, both sexes												
2006	275.1	12.3	1.3	0.9	3.2	10.1	35.3	108.7	255.5	629.6	1,846.2	5,928.2
2005	288.8	12.2	1.3	0.9	3.3	10.1	36.1	110.5	263.2	667.2	1,963.8	6,398.5
2004	293.3	13.8	1.5	0.8	3.2	10.0	36.6	110.9	268.2	697.7	2,035.6	6,616.6
2003	310.3	13.9	1.5	0.8	3.5	10.3	38.2	113.4	284.9	748.3	2,170.8	7,138.0
2002	318.6	15.9	1.4	0.9	3.2	10.0	37.8	114.3	294.8	788.4	2,257.0	7,389.6
2001	323.9	14.9	1.9	0.9	3.2	10.2	37.1	113.4	301.2	810.4	2,322.2	7,673.7
2000	332.9	16.7	1.5	0.9	3.3	9.5	36.8	115.7	319.2	849.3	2,396.2	8,030.3
1999	340.6	17.0	1.5	0.9	3.4	9.6	37.7	116.2	327.3	887.3	2,474.9	8,197.1
All origins 3/, male												
2006	267.7	13.4	1.4	1.0	4.0	13.4	47.5	153.3	355.1	821.3	2,235.3	6,151.8
2005	277.9	13.4	1.5	1.0	4.2	13.0	48.6	155.8	365.7	864.5	2,368.4	6,615.9
2004	281.2	14.7	1.5	0.8	4.0	12.8	49.0	156.7	371.5	903.9	2,462.3	6,818.1
2003	296.3	15.2	1.4	1.0	4.2	12.9	51.1	160.9	393.3	976.3	2,636.1	7,329.3
2002	303.3	16.6	1.5	0.9	4.1	12.7	50.8	162.3	407.0	1,027.8	2,731.1	7,638.8
2001	306.2	15.3	1.9	0.9	3.9	12.8	49.9	160.4	414.0	1,057.6	2,823.3	7,936.0
2000	315.8	17.5	1.6	1.0	3.9	11.8	49.5	165.1	441.6	1,114.1	2,926.6	8,448.1
1999	323.0	17.9	1.6	1.0	4.0	12.1	51.2	165.8	452.0	1,168.4	3,024.0	8,603.2
All origins 3/, female												
2006	282.4	11.2	1.3	0.8	2.4	6.7	23.1	65.5	162.9	467.4	1,580.1	5,823.6
2005	299.3	11.0	1.1	0.8	2.3	7.0	23.7	66.6	167.9	500.8	1,689.1	6,298.7
2004	304.9	12.9	1.5	0.8	2.3	7.1	24.2	66.6	172.2	524.5	1,748.4	6,526.0
2003	323.9	12.5	1.5	0.7	2.8	7.5	25.3	67.5	184.4	557.8	1,860.0	7,053.5
2002	333.3	15.1	1.3	0.8	2.2	7.2	24.9	68.0	191.0	589.1	1,942.2	7,281.5
2001	340.9	14.5	1.8	0.9	2.4	7.6	24.3	68.1	196.9	605.6	1,992.9	7,563.9
2000	349.5	15.9	1.4	0.8	2.8	7.2	24.3	68.0	206.3	631.2	2,050.4	7,860.1
1999	357.5	16.0	1.4	0.8	2.7	7.2	24.4	68.5	212.5	657.0	2,118.2	8,033.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
Hispanic, both sexes												
2006	86.5	10.2	1.1	0.8	2.7	6.3	19.9	70.4	188.4	496.9	1,410.4	4,096.9
2005	91.4	10.5	1.1	0.8	2.8	5.9	20.9	71.1	197.8	545.3	1,564.7	4,469.4
2004	89.3	11.5	1.2	0.6	2.5	5.6	21.2	71.3	202.2	555.6	1,568.0	4,583.7
2003	93.4	11.0	1.3	0.6	2.5	6.4	20.5	71.9	212.2	594.9	1,683.5	5,176.6
2002	94.3	13.1	1.2	0.7	2.6	6.1	20.4	73.6	225.8	621.8	1,728.4	5,369.7
2001	96.2	12.0	1.5	0.8	2.4	6.4	19.8	75.4	216.4	644.9	1,828.4	6,036.2
2000	96.5	13.7	1.3	0.7	2.6	6.4	21.3	75.0	236.3	660.8	1,852.8	6,168.2
1999	99.2	15.2	1.4	0.7	2.5	6.5	21.4	80.5	242.7	690.0	1,935.5	6,335.9
Hispanic, male												
2006	87.1	11.1	1.3	1.0	3.6	8.5	27.4	97.7	258.6	645.2	1,697.6	4,103.7
2005	92.3	11.3	1.5	0.9	3.3	7.9	28.7	100.6	273.3	712.2	1,895.1	4,556.5
2004	90.6	11.7	1.6	0.5	3.5	7.8	29.2	101.1	282.3	730.6	1,928.2	4,600.7
2003	92.6	12.2	1.5	0.7	3.3	7.8	27.9	102.8	295.6	779.5	2,007.9	5,079.7
2002	94.2	12.2	*	0.6	3.3	8.1	27.4	102.9	316.5	822.6	2,067.7	5,494.9
2001	95.4	12.7	1.4	0.8	3.0	8.3	26.8	107.2	297.2	852.4	2,220.5	6,155.6
2000	96.0	13.7	1.4	0.9	3.1	8.1	27.9	107.7	330.6	870.1	2,272.6	6,304.3
1999	99.2	12.7	1.4	0.9	3.0	8.6	29.7	116.3	333.2	925.3	2,356.2	6,595.2
Hispanic, female												
2006	85.9	9.3	*	0.6	1.8	3.5	11.6	42.4	123.8	376.4	1,209.4	4,093.2
2005	90.4	9.7	*	0.7	2.1	3.5	12.2	41.1	128.8	410.4	1,334.6	4,423.7
2004	87.9	11.3	*	0.7	1.4	2.9	12.2	41.4	129.5	415.2	1,319.3	4,575.0
2003	94.3	9.7	*	*	1.6	4.6	12.3	41.0	137.0	447.8	1,460.7	5,225.8
2002	94.4	14.0	1.3	0.8	1.7	3.7	12.8	44.4	144.4	462.0	1,496.0	5,307.7
2001	97.1	11.2	1.6	0.8	1.8	4.2	12.3	44.1	144.6	481.2	1,563.4	5,977.9
2000	96.9	13.8	*	*	2.0	4.5	14.3	42.9	152.6	496.2	1,570.9	6,101.4
1999	99.1	17.9	1.4	*	2.0	4.0	12.7	45.7	162.6	505.6	1,655.5	6,210.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
Non-Hispanic 4/, both sexes												
2006	307.3	12.9	1.4	0.9	3.3	11.1	38.0	112.8	260.5	638.1	1,870.0	6,013.2
2005	321.4	12.8	1.4	0.9	3.4	11.0	38.6	114.5	267.9	674.9	1,983.7	6,483.0
2004	325.9	14.4	1.5	0.9	3.3	11.0	38.9	114.7	272.6	705.9	2,057.5	6,698.1
2003	343.8	14.5	1.5	0.9	3.7	11.1	40.7	117.2	289.5	756.6	2,190.9	7,206.9
2002	352.2	16.5	1.5	0.9	3.3	10.8	40.2	117.8	299.1	796.7	2,276.3	7,451.4
2001	356.7	15.5	1.9	0.9	3.3	10.9	39.3	116.5	306.3	818.6	2,338.1	7,715.2
2000	365.6	17.3	1.6	0.9	3.5	10.1	38.6	118.8	323.8	857.8	2,412.2	8,076.4
1999	372.7	17.4	1.5	0.9	3.6	10.2	39.5	118.8	331.8	895.4	2,489.2	8,239.8
Non-Hispanic 4/, male												
2006	300.1	14.1	1.4	1.0	4.1	14.8	51.2	159.4	362.1	832.0	2,264.9	6,255.9
2005	310.3	14.0	1.5	1.0	4.4	14.3	52.1	161.6	372.1	873.3	2,391.7	6,716.9
2004	313.4	15.5	1.4	0.9	4.1	14.1	52.3	162.2	377.3	913.1	2,486.9	6,917.7
2003	329.5	15.9	1.4	1.0	4.4	14.1	54.6	166.3	399.0	986.2	2,662.3	7,419.6
2002	336.4	17.6	1.7	1.0	4.2	13.8	54.2	167.5	412.3	1,037.1	2,755.7	7,713.8
2001	338.1	15.8	2.0	0.9	4.1	13.7	53.0	164.7	420.6	1,066.8	2,842.6	7,990.6
2000	347.7	18.3	1.7	1.0	4.1	12.6	52.0	169.4	447.3	1,124.0	2,946.0	8,513.1
1999	354.2	19.1	1.7	1.0	4.2	12.7	53.6	169.3	457.9	1,177.2	3,041.8	8,657.6
Non-Hispanic 4/, female												
2006	314.1	11.7	1.4	0.8	2.5	7.4	25.0	68.0	165.9	473.6	1,600.4	5,900.4
2005	332.0	11.4	1.3	0.9	2.4	7.8	25.4	69.2	170.9	506.9	1,707.0	6,376.3
2004	337.9	13.2	1.6	0.9	2.5	8.0	25.9	69.0	175.3	531.4	1,768.8	6,600.0
2003	357.5	13.1	1.6	0.8	3.0	8.1	27.1	70.0	187.7	564.0	1,876.4	7,113.4
2002	367.2	15.3	1.3	0.8	2.3	7.9	26.5	70.0	194.1	595.9	1,958.5	7,338.2
2001	374.3	15.2	1.8	0.9	2.5	8.2	25.9	70.1	200.4	612.3	2,006.9	7,600.7
2000	382.6	16.3	1.5	0.8	2.9	7.6	25.4	70.0	209.6	638.0	2,064.5	7,899.8
1999	390.2	15.5	1.4	0.8	2.9	7.7	25.7	70.2	215.4	664.0	2,130.7	8,072.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
Non-Hispanic white, both sexes												
2006 5/	333.1	9.5	1.2	0.8	2.9	8.9	32.8	98.7	236.4	602.6	1,851.6	6,119.6
2005 5/	348.4	9.8	1.1	0.8	2.8	8.8	33.3	99.5	242.6	640.3	1,963.3	6,596.0
2004 5/	353.1	12.0	1.2	0.7	2.8	8.8	33.9	99.7	247.1	670.5	2,036.7	6,790.0
2003 5/	372.4	10.9	1.2	0.8	3.0	9.0	34.8	101.4	262.4	717.8	2,169.7	7,304.7
2002	381.9	13.4	1.4	0.8	2.7	8.6	34.9	102.9	272.0	760.6	2,252.2	7,546.1
2001	385.8	12.1	1.5	0.8	2.7	8.9	33.3	101.3	278.5	779.4	2,313.0	7,811.4
2000	395.5	14.6	1.4	0.8	2.8	8.1	32.8	103.9	295.3	820.2	2,384.9	8,179.0
1999	402.7	14.2	1.3	0.8	3.0	8.3	33.3	104.1	302.5	856.8	2,462.9	8,342.9
Non-Hispanic white, male												
2006 5/	322.3	10.2	1.1	0.9	3.7	12.1	45.6	143.1	332.0	790.3	2,252.6	6,398.5
2005 5/	333.6	10.7	1.3	0.8	3.6	11.6	46.2	144.4	341.7	834.9	2,380.1	6,880.5
2004 5/	336.9	13.0	1.1	0.8	3.5	11.6	47.1	145.3	346.5	876.3	2,476.1	7,045.5
2003 5/	354.1	11.3	0.9	0.9	3.6	11.7	48.5	148.4	368.3	945.5	2,648.3	7,561.6
2002	362.2	14.8	1.6	0.9	3.5	11.1	48.7	151.1	381.0	1,001.2	2,744.3	7,845.8
2001	363.0	13.0	1.6	0.8	3.3	11.4	46.7	148.1	389.7	1,027.1	2,827.4	8,116.3
2000	373.6	15.5	1.6	0.9	3.3	10.5	46.1	153.2	415.4	1,088.2	2,930.7	8,650.5
1999	379.8	16.1	1.5	0.8	3.4	10.6	46.9	152.7	424.4	1,139.8	3,030.2	8,776.1
Non-Hispanic white, female												
2006 5/	343.6	8.8	1.3	0.6	2.0	5.6	20.0	55.0	145.3	439.9	1,573.4	5,989.5
2005 5/	362.8	8.9	0.9	0.7	1.9	5.9	20.3	55.3	148.4	472.1	1,676.5	6,465.9
2004 5/	368.7	10.9	1.2	0.7	2.0	6.1	20.6	54.8	152.6	493.5	1,737.3	6,675.5
2003 5/	390.0	10.4	1.4	0.6	2.4	6.2	21.2	55.1	161.9	522.8	1,846.4	7,191.6
2002	400.8	12.0	1.1	0.7	1.9	6.0	21.1	55.4	168.8	555.5	1,922.5	7,416.6
2001	407.6	11.0	1.5	0.8	2.0	6.4	19.9	55.3	173.3	569.4	1,971.8	7,684.6
2000	416.5	13.6	1.1	0.6	2.2	5.7	19.5	55.3	182.0	594.6	2,026.2	7,988.9
1999	424.7	12.1	1.1	0.7	2.5	6.0	19.8	56.2	187.7	620.0	2,091.4	8,170.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
Non-Hispanic black, both sexes												
2006 5/	255.2	26.3	2.2	1.4	5.8	23.8	74.6	225.6	502.9	1,051.1	2,378.1	5,656.4
2005 5/	265.1	23.5	2.4	1.6	6.4	24.0	76.5	233.4	522.3	1,084.7	2,531.9	6,088.3
2004 5/	269.2	22.4	3.0	1.5	6.1	23.8	76.8	236.2	535.6	1,140.7	2,606.3	6,436.5
2003 5/	282.2	26.6	2.9	1.5	7.1	23.7	81.3	244.5	565.5	1,224.4	2,745.6	6,923.9
2002	285.1	28.3	2.1	1.4	6.0	24.3	78.2	242.6	578.0	1,243.1	2,851.9	7,136.9
2001	289.7	30.1	3.7	1.3	6.8	23.5	80.7	243.2	587.2	1,293.5	2,915.8	7,366.6
2000	294.2	28.8	2.5	1.5	7.0	22.3	79.1	246.4	609.6	1,333.1	2,995.1	7,576.5
1999	298.5	30.2	2.6	1.4	7.0	21.6	83.7	246.8	623.7	1,382.8	3,044.9	7,624.4
Non-Hispanic black, male												
2006 5/	259.6	29.6	2.7	1.3	6.6	30.9	94.6	302.6	694.8	1,390.3	2,867.7	5,493.9
2005 5/	264.0	26.8	2.4	1.6	8.4	30.3	97.5	311.4	715.4	1,400.0	2,994.3	5,778.7
2004 5/	266.0	24.8	2.9	1.5	7.8	29.0	95.7	312.4	730.6	1,453.6	3,071.2	6,280.6
2003 5/	277.5	31.7	3.3	1.6	8.7	29.2	101.4	324.7	753.7	1,562.4	3,299.8	6,625.9
2002	277.2	29.1	2.0	1.5	7.3	30.1	97.5	319.3	775.5	1,576.9	3,318.5	6,982.7
2001	281.2	29.7	3.8	1.2	8.3	28.9	100.1	319.1	772.1	1,639.6	3,461.6	7,413.8
2000	284.3	30.0	2.3	1.4	7.8	26.5	97.6	324.0	807.5	1,658.1	3,527.8	7,755.5
1999	291.1	31.5	2.7	1.6	8.7	26.6	105.2	329.7	833.1	1,738.6	3,542.5	7,952.2
Non-Hispanic black, female												
2006 5/	251.3	22.8	1.8	1.4	4.9	17.3	56.9	159.6	348.4	808.9	2,100.3	5,722.7
2005 5/	266.1	20.1	2.4	1.6	4.5	18.1	58.0	166.5	366.9	859.2	2,269.7	6,212.9
2004 5/	272.2	20.0	3.0	1.6	4.3	19.1	60.1	170.8	379.0	917.6	2,343.8	6,498.1
2003 5/	286.5	21.3	2.4	1.3	5.5	18.8	63.5	175.8	414.5	984.0	2,432.6	7,039.3
2002	292.3	27.4	2.2	1.4	4.7	18.9	61.2	176.8	419.5	1,006.4	2,588.0	7,195.9
2001	297.4	30.6	3.6	1.4	5.3	18.6	63.5	178.1	438.5	1,049.3	2,609.4	7,349.1
2000	303.1	27.5	2.9	1.5	6.3	18.6	62.8	179.8	450.3	1,104.6	2,695.3	7,510.0
1999	305.1	28.9	2.4	1.3	5.3	17.2	64.7	176.0	455.9	1,131.8	2,766.3	7,501.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
All origins 3/, both sexes												
2006	211.0	8.4	1.0	0.6	2.5	8.2	28.3	88.0	207.3	490.3	1,383.1	4,480.8
2005	220.0	8.7	0.9	0.6	2.7	8.1	28.9	89.7	214.8	518.9	1,460.8	4,778.4
2004	222.2	10.3	1.2	0.6	2.5	7.9	29.3	90.2	218.8	541.6	1,506.3	4,895.9
2003	235.6	11.0	1.2	0.6	2.7	8.2	30.7	92.5	233.2	585.0	1,611.1	5,278.4
2002	241.7	12.4	1.1	0.6	2.5	7.9	30.5	93.7	241.5	615.9	1,677.2	5,446.8
2001	245.8	11.9	1.5	0.7	2.5	8.0	29.6	92.9	246.9	635.1	1,725.7	5,664.2
2000	252.6	13.0	1.2	0.7	2.6	7.4	29.2	94.2	261.2	665.6	1,780.3	5,926.1
1999	259.9	13.8	1.2	0.7	2.8	7.6	30.2	95.7	269.9	701.7	1,849.9	6,063.0
All origins 3/, male												
2006	214.0	8.8	1.1	0.7	3.3	11.2	39.5	128.9	296.8	660.5	1,743.5	4,819.9
2005	221.1	9.4	1.0	0.6	3.6	10.8	40.7	131.5	306.9	692.3	1,829.4	5,143.4
2004	222.8	10.9	1.1	0.6	3.2	10.5	40.9	132.3	312.8	723.8	1,893.6	5,239.3
2003	235.0	12.1	1.1	0.7	3.4	10.5	42.8	136.2	331.7	785.3	2,030.3	5,621.5
2002	240.7	12.9	1.1	0.7	3.3	10.5	43.1	138.4	343.4	827.1	2,110.1	5,823.5
2001	242.5	11.8	1.5	0.7	3.2	10.3	41.7	136.6	349.8	851.3	2,177.3	6,040.5
2000	249.8	13.3	1.4	0.8	3.2	9.6	41.4	140.2	371.7	898.3	2,248.1	6,430.0
1999	257.0	13.9	1.3	0.8	3.4	9.8	43.0	142.0	383.0	950.6	2,336.4	6,561.6
All origins 3/, female												
2006	208.0	7.9	0.9	0.6	1.8	5.1	17.0	48.5	124.1	346.3	1,136.7	4,322.1
2005	218.9	8.0	0.9	0.6	1.7	5.3	17.1	49.2	129.1	372.7	1,210.5	4,610.8
2004	221.6	9.7	1.2	0.6	1.7	5.2	17.7	49.6	131.5	388.6	1,245.6	4,741.5
2003	236.2	9.8	1.3	0.5	2.1	5.7	18.6	50.2	141.9	417.5	1,331.1	5,126.7
2002	242.7	11.8	1.0	0.6	1.7	5.2	18.0	50.6	147.2	440.1	1,389.7	5,283.3
2001	249.0	12.0	1.4	0.7	1.8	5.6	17.6	50.7	151.8	455.9	1,428.9	5,506.8
2000	255.3	12.5	1.0	0.5	2.1	5.2	17.2	49.8	159.3	474.0	1,475.1	5,720.9
1999	262.6	13.7	1.1	0.6	2.1	5.4	17.5	51.0	165.6	497.9	1,533.9	5,862.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
Hispanic, both sexes												
2006	65.3	6.5	0.8	0.6	1.8	4.6	14.0	51.7	142.2	377.2	1,064.9	3,140.6
2005	69.2	7.6	0.6	0.4	1.9	4.5	15.0	52.3	153.1	420.1	1,176.9	3,407.5
2004	67.2	8.9	1.1	0.4	1.8	4.0	15.2	52.4	155.7	424.2	1,177.6	3,446.8
2003	70.9	8.7	1.1	0.4	1.9	4.7	14.9	53.3	165.4	461.2	1,261.0	3,932.9
2002	71.9	10.2	0.8	0.4	1.9	4.4	14.7	55.3	176.8	484.3	1,297.6	4,121.9
2001	73.3	9.7	1.1	0.5	1.7	4.6	13.8	55.2	170.7	504.5	1,376.3	4,607.5
2000	73.1	11.2	1.0	0.6	1.9	4.3	14.8	55.9	183.3	506.5	1,395.9	4,701.8
1999	76.2	12.7	1.2	0.6	1.8	4.5	15.4	60.9	192.7	536.3	1,467.8	4,888.6
Hispanic, male												
2006	67.7	7.1	*	0.7	2.5	6.5	20.1	75.6	202.3	505.6	1,308.4	3,257.9
2005	72.1	8.0	*	0.5	2.4	6.3	21.7	77.9	219.3	561.5	1,469.2	3,534.2
2004	70.2	8.9	1.4	*	2.7	5.9	22.1	77.6	224.6	572.2	1,489.0	3,496.8
2003	72.2	9.5	*	*	2.7	6.1	21.1	79.6	235.6	625.0	1,543.5	3,874.5
2002	74.0	9.6	*	*	2.6	6.1	20.6	80.5	256.0	657.7	1,599.5	4,301.8
2001	74.6	10.3	*	*	2.1	6.1	20.0	82.9	242.2	683.7	1,702.7	4,784.3
2000	74.7	11.2	1.4	0.7	2.4	5.7	20.7	84.3	264.8	684.8	1,733.2	4,897.5
1999	77.7	10.3	1.4	0.7	2.1	6.5	22.5	92.2	272.1	737.0	1,816.0	5,012.2
Hispanic, female												
2006	62.6	5.9	*	*	0.9	2.3	7.1	27.3	86.9	273.0	894.5	3,078.3
2005	66.2	7.2	*	*	1.4	2.3	7.4	26.2	92.6	305.9	973.4	3,341.3
2004	64.1	9.0	*	*	0.8	1.7	7.5	27.0	93.1	305.5	962.7	3,421.2
2003	69.6	7.8	*	*	1.1	3.1	8.0	27.0	102.1	330.6	1,067.0	3,962.5
2002	69.7	10.8	*	*	1.1	2.4	8.2	30.2	105.7	346.4	1,090.8	4,032.8
2001	71.8	9.0	*	*	1.2	2.8	7.1	27.9	107.2	363.1	1,155.7	4,521.1
2000	71.5	11.2	*	*	1.3	2.7	8.5	28.2	111.2	366.3	1,169.4	4,605.8
1999	74.6	15.2	*	*	1.5	2.3	7.9	30.5	122.5	379.0	1,236.1	4,828.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
Non-Hispanic 4/, both sexes												
2006	235.8	8.9	1.0	0.6	2.7	9.0	30.7	92.0	212.2	497.6	1,400.4	4,542.5
2005	244.9	9.0	1.0	0.7	2.8	8.9	31.2	93.5	219.3	525.1	1,474.7	4,837.9
2004	247.0	10.7	1.2	0.6	2.6	8.8	31.5	93.9	223.1	548.4	1,521.3	4,953.4
2003	261.0	11.5	1.2	0.7	2.9	8.9	32.9	96.1	237.5	591.6	1,625.0	5,324.5
2002	267.1	12.9	1.1	0.7	2.6	8.7	32.7	97.1	245.5	622.2	1,690.6	5,484.9
2001	270.7	12.3	1.5	0.7	2.6	8.7	31.6	96.0	251.6	641.4	1,736.4	5,688.6
2000	277.3	13.3	1.2	0.7	2.8	8.0	30.8	97.1	265.7	672.7	1,791.1	5,953.6
1999	284.3	14.0	1.2	0.7	2.9	8.2	31.8	98.3	274.1	708.5	1,859.2	6,086.3
Non-Hispanic 4/, male												
2006	240.3	9.4	1.1	0.7	3.4	12.4	43.1	134.8	303.8	669.9	1,767.6	4,899.0
2005	247.1	9.8	1.0	0.7	3.8	12.0	44.0	137.2	313.2	699.9	1,847.0	5,222.0
2004	248.5	11.5	1.1	0.6	3.3	11.7	44.0	137.7	318.7	731.9	1,911.8	5,317.0
2003	261.4	12.8	1.1	0.8	3.5	11.6	46.1	141.5	337.4	793.4	2,050.1	5,690.6
2002	267.0	13.7	1.2	0.7	3.4	11.6	46.4	143.6	348.6	834.6	2,128.7	5,874.4
2001	268.0	12.0	1.7	0.7	3.3	11.2	44.6	141.1	356.1	858.7	2,192.5	6,077.8
2000	275.2	13.7	1.4	0.8	3.3	10.3	43.8	144.5	377.5	907.0	2,263.4	6,474.6
1999	282.0	14.8	1.3	0.8	3.6	10.5	45.4	145.7	388.7	958.3	2,349.8	6,602.0
Non-Hispanic 4/, female												
2006	231.4	8.4	1.0	0.6	1.9	5.7	18.6	50.8	127.0	351.4	1,149.7	4,376.7
2005	242.8	8.3	1.0	0.7	1.8	5.9	18.7	51.5	131.9	377.2	1,222.3	4,662.6
2004	245.5	9.8	1.3	0.6	1.9	5.9	19.2	51.8	134.3	393.8	1,258.8	4,790.8
2003	260.5	10.2	1.3	0.6	2.3	6.2	20.1	52.4	144.6	422.4	1,341.5	5,163.6
2002	267.2	12.0	1.1	0.6	1.8	5.8	19.3	52.4	150.0	444.8	1,400.2	5,316.8
2001	273.3	12.7	1.4	0.7	1.9	6.1	18.9	52.6	154.8	460.9	1,437.0	5,526.8
2000	279.3	12.8	1.1	0.6	2.2	5.7	18.1	51.5	162.2	479.3	1,483.4	5,742.9
1999	286.4	13.3	1.1	0.6	2.3	5.9	18.5	52.6	168.2	503.4	1,540.8	5,879.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
Non-Hispanic white, both sexes												
2006 5/	257.4	6.5	0.9	0.5	2.3	7.4	27.6	83.1	196.8	475.5	1,394.5	4,642.7
2005 5/	267.1	7.0	0.8	0.5	2.3	7.2	27.8	84.0	202.4	504.3	1,466.8	4,942.5
2004 5/	269.1	9.1	0.9	0.5	2.1	7.2	28.2	84.2	206.6	526.3	1,514.3	5,038.3
2003 5/	284.0	8.7	0.9	0.6	2.3	7.3	29.2	85.8	219.6	566.7	1,616.7	5,412.2
2002	290.8	10.7	1.0	0.6	2.1	6.9	29.3	87.5	227.2	599.7	1,679.2	5,568.4
2001	294.1	9.8	1.2	0.6	2.1	7.1	27.6	86.4	233.0	616.7	1,724.0	5,776.0
2000	301.3	11.4	1.1	0.5	2.1	6.4	27.2	87.7	246.4	648.1	1,777.4	6,047.9
1999	308.2	11.3	1.0	0.6	2.4	6.7	27.6	88.7	254.0	683.2	1,846.1	6,176.6
Non-Hispanic white, male												
2006 5/	260.3	6.8	0.9	0.6	3.0	10.3	39.8	124.5	284.5	644.3	1,767.4	5,032.8
2005 5/	267.8	7.5	0.8	0.6	3.1	9.9	40.3	126.2	293.0	677.6	1,849.3	5,374.1
2004 5/	269.1	9.8	0.8	0.5	2.7	9.7	40.6	126.9	298.8	709.5	1,915.1	5,430.9
2003 5/	282.9	9.4	0.7	0.7	2.8	9.8	42.2	129.8	317.7	767.3	2,049.9	5,821.0
2002	289.2	11.6	1.1	0.7	2.8	9.4	42.9	133.1	327.6	813.5	2,129.9	5,994.1
2001	289.5	10.3	1.3	0.6	2.6	9.4	40.4	130.7	335.8	834.7	2,190.4	6,195.4
2000	297.5	12.0	1.2	0.7	2.6	8.6	40.1	134.3	356.3	885.1	2,261.9	6,606.6
1999	304.0	12.4	1.1	0.6	2.8	8.7	40.6	135.0	365.6	935.2	2,352.6	6,713.7
Non-Hispanic white, female												
2006 5/	254.7	6.1	0.9	0.5	1.5	4.4	15.5	42.2	113.2	329.1	1,135.8	4,460.8
2005 5/	266.4	6.5	0.7	0.5	1.4	4.5	15.2	42.4	116.1	354.6	1,203.6	4,745.1
2004 5/	269.1	8.3	1.0	0.5	1.5	4.6	15.6	42.2	118.9	368.6	1,241.2	4,862.4
2003 5/	285.1	8.0	1.1	0.5	1.9	4.7	16.2	42.4	126.6	394.8	1,324.0	5,232.2
2002	292.3	9.6	0.8	0.5	1.4	4.4	15.8	42.6	132.0	417.4	1,377.2	5,384.5
2001	298.4	9.2	1.1	0.6	1.5	4.7	14.8	42.7	136.0	431.8	1,414.7	5,601.6
2000	304.9	10.8	0.9	0.4	1.7	4.2	14.3	41.9	142.9	448.5	1,458.9	5,822.7
1999	312.3	10.2	0.8	0.6	1.9	4.6	14.5	43.1	149.0	472.3	1,514.5	5,962.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
Non-Hispanic black, both sexes												
2006 5/	189.3	17.9	1.8	1.0	4.8	19.1	55.8	170.1	382.6	786.1	1,729.7	4,103.9
2005 5/	196.3	16.5	1.8	1.2	5.3	19.0	58.0	175.8	403.4	806.0	1,837.3	4,373.6
2004 5/	198.7	15.6	2.5	1.2	5.0	18.5	58.3	179.9	410.1	852.4	1,866.7	4,610.3
2003 5/	209.4	20.4	2.3	1.2	5.7	18.8	61.4	185.8	434.7	925.6	1,983.5	4,977.8
2002	212.2	21.9	1.7	1.0	5.1	19.2	59.4	184.7	450.2	936.3	2,067.0	5,151.0
2001	215.5	22.9	3.1	1.1	5.6	18.7	61.3	183.6	454.5	977.9	2,123.4	5,307.7
2000	218.6	20.5	2.0	1.2	5.9	17.9	58.7	185.4	473.2	1,014.8	2,177.9	5,424.6
1999	223.8	24.8	2.1	1.2	6.0	17.5	63.6	189.1	488.5	1,058.0	2,223.2	5,514.7
Non-Hispanic black, male												
2006 5/	198.8	18.8	2.1	1.1	5.6	25.4	72.9	235.0	537.9	1,062.7	2,164.2	4,129.6
2005 5/	201.5	19.0	*	1.1	7.2	24.6	76.8	242.9	559.3	1,059.7	2,229.0	4,277.2
2004 5/	203.2	17.0	2.5	1.1	6.5	23.5	75.1	245.1	570.4	1,111.3	2,262.0	4,695.5
2003 5/	212.6	24.5	2.5	1.3	7.4	23.1	79.7	253.9	589.2	1,210.9	2,449.4	4,897.8
2002	212.9	23.0	*	1.0	6.3	25.0	77.6	250.9	615.0	1,207.5	2,470.0	5,161.2
2001	215.7	20.7	3.4	1.0	7.1	23.6	78.9	247.6	610.5	1,259.3	2,586.9	5,495.4
2000	217.6	20.1	2.0	1.2	6.9	22.1	75.6	252.5	639.6	1,281.7	2,615.1	5,677.2
1999	224.9	24.6	2.3	1.4	7.7	22.1	83.8	259.8	668.4	1,355.1	2,631.6	5,877.1
Non-Hispanic black, female												
2006 5/	180.6	17.0	*	0.9	3.9	13.2	40.6	114.4	257.7	588.6	1,483.2	4,093.4
2005 5/	191.6	14.0	2.1	1.3	3.4	13.9	41.3	118.2	278.0	624.6	1,615.3	4,412.4
2004 5/	194.6	14.2	2.4	1.4	3.5	14.0	43.5	124.0	281.4	667.9	1,643.6	4,576.6
2003 5/	206.5	16.1	2.0	1.1	4.0	14.8	45.3	127.4	310.7	722.8	1,720.5	5,008.9
2002	211.6	20.8	1.9	1.0	3.9	14.1	43.3	128.0	317.9	744.0	1,839.1	5,147.0
2001	215.4	25.2	2.9	1.1	4.0	14.2	45.7	128.7	328.9	779.4	1,863.2	5,238.2
2000	219.5	20.8	2.0	1.1	4.9	14.1	43.7	127.8	339.3	827.1	1,931.9	5,330.8
1999	222.8	25.0	1.9	1.0	4.4	13.3	45.8	128.7	344.3	848.5	1,994.5	5,379.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
All origins 3/, both sexes												
2006	1.1	*	*	*	*	0.1	0.2	0.4	1.1	3.2	8.6	16.7
2005	1.1	*	*	*	*	0.1	0.2	0.4	1.2	3.3	8.5	19.0
2004	1.1	*	*	*	*	0.1	0.2	0.5	1.1	3.3	8.8	17.1
2003	1.2	*	*	*	*	0.1	0.2	0.5	1.3	3.8	9.6	20.5
2002	1.2	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.6	17.8
2001	1.2	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.4	17.7
2000	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.5	16.8
1999	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.0	18.0
All origins 3/, male												
2006	0.7	*	*	*	*	*	0.1	0.4	0.9	2.4	6.2	12.0
2005	0.7	*	*	*	*	*	0.2	0.4	1.0	2.5	6.2	14.0
2004	0.7	*	*	*	*	*	0.2	0.4	0.8	2.4	7.3	11.5
2003	0.8	*	*	*	*	*	0.2	0.4	1.1	2.9	7.5	16.2
2002	0.8	*	*	*	*	*	0.2	0.5	1.2	3.0	6.7	11.8
2001	0.7	*	*	*	*	*	0.2	0.5	1.1	2.8	6.6	12.8
2000	0.8	*	*	*	*	*	0.3	0.6	1.4	2.9	6.6	12.2
1999	0.8	*	*	*	*	*	0.2	0.5	1.2	3.5	6.8	10.3
All origins 3/, female												
2006	1.5	*	*	*	*	*	0.2	0.4	1.3	3.9	10.3	18.9
2005	1.5	*	*	*	*	*	0.2	0.5	1.4	4.0	10.1	21.3
2004	1.5	*	*	*	*	*	0.3	0.6	1.4	4.1	9.9	19.5
2003	1.7	*	*	*	*	*	0.2	0.7	1.5	4.5	11.0	22.3
2002	1.7	*	*	*	*	*	0.2	0.7	2.0	4.6	11.6	20.4
2001	1.7	*	*	*	*	*	0.3	0.7	1.7	5.1	11.3	19.8
2000	1.8	*	*	*	*	0.1	0.4	0.8	2.0	5.8	11.4	18.7
1999	1.9	*	*	*	*	0.1	0.3	0.8	2.2	5.6	12.2	21.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Hispanic, both sexes												
2006	0.4	*	*	*	*	*	*	*	1.1	2.5	6.6	10.2
2005	0.4	*	*	*	*	*	*	0.5	1.1	3.0	5.6	8.6
2004	0.4	*	*	*	*	*	*	*	1.3	3.5	6.3	*
2003	0.4	*	*	*	*	*	*	0.5	1.0	3.3	6.7	*
2002	0.5	*	*	*	*	*	*	0.7	1.8	4.6	7.0	*
2001	0.4	*	*	*	*	*	*	*	1.5	4.2	5.4	*
2000	0.4	*	*	*	*	*	*	*	1.8	4.4	5.3	*
1999	0.5	*	*	*	*	*	*	0.8	1.6	4.2	7.3	*
Hispanic, male												
2006	0.3	*	*	*	*	*	*	*	*	*	*	*
2005	0.2	*	*	*	*	*	*	*	*	*	*	*
2004	0.3	*	*	*	*	*	*	*	*	*	*	*
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.5	*	*	*	*	*	*	*	*	2.6	8.9	12.0
2005	0.6	*	*	*	*	*	*	*	*	3.7	8.1	*
2004	0.5	*	*	*	*	*	*	*	2.1	3.7	6.4	*
2003	0.5	*	*	*	*	*	*	*	*	4.0	8.4	*
2002	0.8	*	*	*	*	*	*	*	2.1	5.9	10.7	*
2001	0.6	*	*	*	*	*	*	*	2.0	5.2	*	*
2000	0.6	*	*	*	*	*	*	*	*	5.6	6.9	*
1999	0.8	*	*	*	*	*	*	*	*	5.9	10.4	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Non-Hispanic 4/, both sexes												
2006	1.2	*	*	*	*	0.1	0.2	0.4	1.1	3.3	8.7	17.0
2005	1.3	*	*	*	*	0.1	0.2	0.4	1.2	3.3	8.7	19.5
2004	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.3	9.0	17.7
2003	1.4	*	*	*	*	*	0.2	0.5	1.3	3.8	9.7	20.9
2002	1.4	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.7	18.1
2001	1.3	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.6	18.0
2000	1.4	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.7	17.1
1999	1.4	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.1	18.3
Non-Hispanic 4/, male												
2006	0.8	*	*	*	*	*	0.1	0.4	0.9	2.4	6.3	12.3
2005	0.8	*	*	*	*	*	0.2	0.4	1.0	2.5	6.4	14.6
2004	0.8	*	*	*	*	*	0.2	0.4	0.8	2.3	7.3	11.9
2003	0.9	*	*	*	*	*	0.2	0.4	1.1	2.9	7.6	16.6
2002	0.8	*	*	*	*	*	0.2	0.5	1.2	3.0	7.0	12.0
2001	0.8	*	*	*	*	*	0.3	0.5	1.1	2.8	6.7	13.1
2000	0.9	*	*	*	*	*	0.3	0.6	1.4	2.9	6.7	12.5
1999	0.8	*	*	*	*	*	0.2	0.5	1.2	3.6	6.9	10.7
Non-Hispanic 4/, female												
2006	1.6	*	*	*	*	*	0.2	0.4	1.3	4.0	10.3	19.2
2005	1.7	*	*	*	*	*	0.2	0.5	1.4	4.0	10.3	21.8
2004	1.7	*	*	*	*	*	0.2	0.6	1.4	4.1	10.0	20.2
2003	1.8	*	*	*	*	*	0.3	0.7	1.5	4.6	11.1	22.8
2002	1.8	*	*	*	*	*	0.2	0.7	2.0	4.5	11.6	20.8
2001	1.8	*	*	*	*	*	0.3	0.7	1.7	5.1	11.5	20.0
2000	1.9	*	*	*	*	*	0.4	0.8	2.0	5.8	11.6	19.0
1999	2.0	*	*	*	*	0.1	0.3	0.8	2.3	5.5	12.2	21.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Non-Hispanic white, both sexes												
2006 5/	1.4	*	*	*	*	*	0.1	0.3	1.0	3.3	9.2	18.4
2005 5/	1.4	*	*	*	*	*	0.2	0.4	1.1	3.4	9.2	21.0
2004 5/	1.4	*	*	*	*	*	0.1	0.5	1.0	3.4	9.5	18.9
2003 5/	1.5	*	*	*	*	*	0.2	0.4	1.2	3.9	10.4	22.3
2002	1.5	*	*	*	*	*	0.2	0.5	1.5	3.9	10.2	19.3
2001	1.5	*	*	*	*	*	0.2	0.5	1.3	4.0	10.2	19.0
2000	1.6	*	*	*	*	*	0.3	0.6	1.7	4.6	10.3	18.0
1999	1.6	*	*	*	*	0.1	0.2	0.6	1.7	4.8	10.6	19.2
Non-Hispanic white, male												
2006 5/	0.8	*	*	*	*	*	*	0.3	0.8	2.4	6.7	13.2
2005 5/	0.9	*	*	*	*	*	0.2	0.3	0.9	2.5	6.7	15.4
2004 5/	0.9	*	*	*	*	*	*	0.4	0.7	2.4	7.8	12.8
2003 5/	1.0	*	*	*	*	*	0.2	0.3	0.9	3.0	8.3	17.7
2002	0.9	*	*	*	*	*	0.2	0.4	1.1	3.0	7.3	12.9
2001	0.9	*	*	*	*	*	0.2	0.4	1.1	2.7	7.0	13.7
2000	0.9	*	*	*	*	*	0.3	0.5	1.5	2.9	7.2	13.0
1999	0.9	*	*	*	*	*	0.2	0.5	1.1	3.8	7.2	11.6
Non-Hispanic white, female												
2006 5/	1.9	*	*	*	*	*	*	0.4	1.2	4.1	10.9	20.8
2005 5/	1.9	*	*	*	*	*	0.2	0.4	1.2	4.2	10.9	23.6
2004 5/	1.9	*	*	*	*	*	0.2	0.5	1.3	4.2	10.6	21.6
2003 5/	2.1	*	*	*	*	*	0.2	0.5	1.4	4.6	11.8	24.4
2002	2.1	*	*	*	*	*	0.2	0.6	1.9	4.6	12.2	22.0
2001	2.0	*	*	*	*	*	0.2	0.5	1.5	5.1	12.2	21.2
2000	2.1	*	*	*	*	*	0.3	0.7	1.9	6.0	12.4	20.0
1999	2.2	*	*	*	*	*	0.2	0.7	2.2	5.7	12.9	22.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Non-Hispanic black, both sexes												
2006 5/	0.7	*	*	*	*	*	0.4	0.8	1.7	3.0	5.2	*
2005 5/	0.6	*	*	*	*	*	*	1.0	1.5	2.4	4.5	6.3
2004 5/	0.7	*	*	*	*	*	0.6	1.0	1.6	2.7	4.1	6.0
2003 5/	0.8	*	*	*	*	*	0.5	1.4	2.7	4.0	4.2	7.1
2002	0.7	*	*	*	*	*	0.4	1.0	2.4	2.9	4.4	*
2001	0.8	*	*	*	*	*	0.7	1.4	2.1	4.2	4.3	6.9
2000	0.7	*	*	*	*	*	0.6	1.0	2.0	3.6	4.0	8.6
1999	0.8	*	*	*	*	*	0.5	1.2	2.2	3.4	4.4	8.4
Non-Hispanic black, male												
2006 5/	0.5	*	*	*	*	*	*	*	1.7	*	*	*
2005 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.5	*	*	*	*	*	*	0.9	*	*	*	*
2003 5/	0.6	*	*	*	*	*	*	0.9	2.8	3.0	*	*
2002	0.5	*	*	*	*	*	*	*	1.8	2.9	*	*
2001	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	0.8	*	*	*	*	*	*	0.9	1.8	3.3	6.2	*
2005 5/	0.7	*	*	*	*	*	*	1.1	1.6	2.9	4.6	*
2004 5/	0.8	*	*	*	*	*	*	1.1	1.7	3.4	5.2	*
2003 5/	1.0	*	*	*	*	*	*	1.7	2.6	4.6	5.2	*
2002	0.8	*	*	*	*	*	*	1.1	2.9	2.9	5.1	*
2001	1.2	*	*	*	*	*	0.9	1.7	2.6	5.4	5.4	*
2000	1.0	*	*	*	*	*	0.8	1.4	2.7	4.4	4.9	*
1999	1.0	*	*	*	*	*	*	1.5	2.7	4.1	4.4	11.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
All origins 3/, both sexes												
2006	9.9	*	*	*	0.1	0.7	3.1	8.3	13.1	20.4	47.8	194.0
2005	9.9	*	*	*	0.1	0.8	3.1	7.8	13.0	20.4	48.8	198.6
2004	9.7	*	*	*	0.1	0.8	3.0	7.8	13.4	21.7	49.3	192.8
2003	9.7	*	*	*	0.1	0.8	2.9	7.9	13.1	22.1	51.0	195.0
2002	9.2	*	*	*	0.1	0.7	2.7	7.2	12.5	21.3	49.6	187.0
2001	8.7	*	*	*	0.1	0.6	2.4	6.3	10.9	19.8	48.7	189.5
2000	8.4	*	*	*	0.1	0.5	2.2	5.6	11.0	19.6	48.2	193.1
1999	8.1	*	*	*	0.1	0.5	2.1	5.3	10.7	19.1	47.7	184.9
All origins 3/, male												
2006	9.3	*	*	*	0.2	1.0	4.2	11.7	18.0	25.7	49.7	143.0
2005	9.1	*	*	*	0.1	1.1	4.4	10.9	18.1	25.2	48.7	147.5
2004	8.9	*	*	*	0.1	1.1	4.1	10.7	18.5	27.3	48.0	145.4
2003	8.8	*	*	*	0.2	1.0	4.0	10.8	17.2	27.1	51.0	142.2
2002	8.2	*	*	*	0.1	1.1	3.8	9.8	16.9	25.2	49.4	134.6
2001	7.5	*	*	*	*	0.8	3.3	8.6	14.4	23.6	48.4	134.4
2000	7.2	*	*	*	0.1	0.7	2.9	7.8	14.5	23.7	46.5	139.1
1999	6.8	*	*	*	0.1	0.7	2.9	7.0	13.7	21.9	46.8	133.6
All origins 3/, female												
2006	10.6	*	*	*	*	0.4	2.0	5.0	8.5	15.9	46.5	217.9
2005	10.7	*	*	*	*	0.5	1.9	4.7	8.3	16.4	48.9	222.0
2004	10.5	*	*	*	*	0.4	1.9	5.1	8.6	16.9	50.2	214.1
2003	10.7	*	*	*	*	0.5	1.9	5.2	9.3	17.8	51.0	218.3
2002	10.2	*	*	*	*	0.4	1.6	4.6	8.4	18.1	49.8	209.7
2001	9.8	*	*	*	*	0.4	1.6	4.1	7.8	16.6	48.8	212.6
2000	9.7	*	*	*	*	0.4	1.5	3.5	7.8	16.2	49.4	215.1
1999	9.4	*	*	*	*	0.3	1.3	3.6	7.9	16.8	48.3	205.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
Hispanic, both sexes												
2006	3.6	*	*	*	*	0.4	1.8	4.9	9.5	18.4	43.1	157.7
2005	3.5	*	*	*	*	0.4	1.7	4.9	10.8	16.5	42.8	145.0
2004	3.6	*	*	*	*	0.3	1.7	4.8	11.0	22.2	47.0	150.3
2003	4.0	*	*	*	*	0.4	1.8	5.7	11.8	22.9	54.2	179.3
2002	3.7	*	*	*	*	0.4	1.7	4.9	12.2	22.4	51.4	165.7
2001	3.3	*	*	*	*	0.3	1.7	4.9	10.0	21.0	46.9	151.9
2000	2.8	*	*	*	*	0.3	1.2	3.8	8.2	18.4	46.0	143.3
1999	2.7	*	*	*	*	*	1.2	3.3	8.9	16.8	44.8	152.2
Hispanic, male												
2006	3.7	*	*	*	*	0.6	2.7	7.1	13.1	23.8	44.6	125.2
2005	3.5	*	*	*	*	0.6	2.5	7.0	14.3	19.5	45.8	118.3
2004	3.8	*	*	*	*	*	2.7	6.1	16.0	27.8	55.4	131.7
2003	4.3	*	*	*	*	0.5	2.9	8.3	17.0	30.8	63.7	144.7
2002	3.9	*	*	*	*	0.6	2.6	7.4	17.0	28.5	61.1	137.2
2001	3.4	*	*	*	*	*	2.7	7.0	13.8	26.4	48.3	144.7
2000	2.7	*	*	*	*	*	1.6	5.4	10.7	22.4	50.6	112.9
1999	2.6	*	*	*	*	*	1.8	4.7	12.5	20.9	42.8	107.8
Hispanic, female												
2006	3.6	*	*	*	*	*	0.8	2.6	6.3	14.0	42.0	174.9
2005	3.4	*	*	*	*	*	0.9	2.8	7.6	14.1	40.7	159.0
2004	3.4	*	*	*	*	*	*	3.4	6.4	17.7	41.2	159.9
2003	3.7	*	*	*	*	*	*	3.1	7.2	16.6	47.7	196.9
2002	3.4	*	*	*	*	*	*	2.5	7.9	17.5	44.7	179.9
2001	3.1	*	*	*	*	*	*	2.7	6.5	16.8	45.9	155.5
2000	2.9	*	*	*	*	*	0.8	2.1	6.1	15.3	42.9	158.3
1999	2.9	*	*	*	*	*	*	2.0	5.8	13.6	46.1	173.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
Non-Hispanic 4/, both sexes												
2006	11.0	*	*	*	0.1	0.8	3.3	8.6	13.3	20.5	47.9	195.6
2005	10.9	*	*	*	0.1	0.9	3.3	8.0	13.2	20.6	49.0	200.7
2004	10.7	*	*	*	0.1	0.9	3.2	8.1	13.5	21.5	49.3	194.3
2003	10.6	*	*	*	0.1	0.9	3.1	8.1	13.1	21.9	50.7	195.0
2002	10.0	*	*	*	0.1	0.8	2.8	7.4	12.4	21.0	49.3	187.1
2001	9.4	*	*	*	*	0.7	2.5	6.3	10.9	19.6	48.6	190.3
2000	9.2	*	*	*	0.1	0.6	2.3	5.7	11.2	19.5	48.1	194.3
1999	8.8	*	*	*	0.1	0.6	2.2	5.4	10.7	19.1	47.6	185.3
Non-Hispanic 4/, male												
2006	10.3	*	*	*	0.2	1.1	4.5	12.2	18.3	25.7	49.8	143.8
2005	10.0	*	*	*	0.1	1.2	4.7	11.3	18.3	25.4	48.7	148.7
2004	9.8	*	*	*	0.2	1.3	4.3	11.1	18.5	27.1	47.4	145.6
2003	9.5	*	*	*	0.2	1.1	4.1	10.9	17.1	26.7	50.1	141.7
2002	8.9	*	*	*	0.1	1.2	4.0	10.0	16.8	24.6	48.5	133.7
2001	8.1	*	*	*	*	0.9	3.4	8.6	14.2	23.3	48.3	133.4
2000	7.8	*	*	*	0.1	0.7	3.0	7.9	14.6	23.5	46.0	139.6
1999	7.4	*	*	*	0.1	0.8	3.1	7.2	13.6	21.7	46.7	134.2
Non-Hispanic 4/, female												
2006	11.7	*	*	*	*	0.4	2.2	5.2	8.7	16.1	46.7	219.6
2005	11.8	*	*	*	*	0.6	2.0	4.9	8.3	16.6	49.2	224.5
2004	11.6	*	*	*	*	0.5	2.1	5.2	8.8	16.7	50.6	216.0
2003	11.7	*	*	*	*	0.6	2.1	5.4	9.4	17.8	51.1	218.5
2002	11.1	*	*	*	*	0.5	1.7	4.8	8.4	18.0	49.9	210.2
2001	10.7	*	*	*	*	0.4	1.7	4.2	7.8	16.5	48.8	214.0
2000	10.6	*	*	*	*	0.4	1.6	3.6	8.0	16.2	49.5	216.5
1999	10.2	*	*	*	*	0.3	1.4	3.7	8.0	16.9	48.2	205.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
Non-Hispanic white, both sexes												
2006 5/	10.3	*	*	*	0.1	0.5	2.4	6.3	10.0	15.5	41.9	188.0
2005 5/	10.1	*	*	*	*	0.6	2.3	5.6	10.0	16.1	42.3	190.6
2004 5/	9.8	*	*	*	*	0.5	2.2	5.6	9.8	17.0	42.7	184.6
2003 5/	9.7	*	*	*	*	0.5	2.0	5.3	9.5	16.7	44.3	184.9
2002	9.1	*	*	*	*	0.4	1.9	5.1	8.8	16.1	42.2	177.0
2001	8.6	*	*	*	*	0.4	1.6	4.1	7.6	14.8	42.2	180.4
2000	8.3	*	*	*	*	0.3	1.5	3.6	7.5	14.5	41.2	182.5
1999	7.9	*	*	*	*	0.3	1.3	3.4	7.3	14.1	40.6	175.2
Non-Hispanic white, male												
2006 5/	9.0	*	*	*	0.2	0.7	3.5	9.4	14.0	19.3	42.4	134.7
2005 5/	8.8	*	*	*	*	0.9	3.4	8.3	14.2	19.9	42.0	137.6
2004 5/	8.6	*	*	*	*	0.8	3.2	8.3	13.6	21.6	40.9	135.9
2003 5/	8.2	*	*	*	*	0.8	3.0	7.8	13.0	20.4	43.5	130.8
2002	7.6	*	*	*	*	0.7	2.8	7.3	12.2	19.0	41.2	123.5
2001	6.9	*	*	*	*	0.6	2.3	5.9	10.5	18.1	41.5	122.0
2000	6.6	*	*	*	*	0.4	2.1	5.4	10.3	18.0	38.8	129.5
1999	6.2	*	*	*	*	0.6	2.0	4.9	9.6	16.2	39.4	123.8
Non-Hispanic white, female												
2006 5/	11.4	*	*	*	*	0.2	1.3	3.2	6.2	12.3	41.6	212.9
2005 5/	11.3	*	*	*	*	0.3	1.2	2.9	5.9	12.8	42.5	214.9
2004 5/	11.0	*	*	*	*	*	1.2	3.0	6.2	13.0	43.9	206.5
2003 5/	11.1	*	*	*	*	0.3	1.1	2.8	6.3	13.6	44.9	208.7
2002	10.5	*	*	*	*	0.2	1.0	2.9	5.5	13.6	42.9	200.1
2001	10.2	*	*	*	*	0.2	0.8	2.3	4.9	11.9	42.6	204.7
2000	9.9	*	*	*	*	0.2	0.8	1.9	4.9	11.6	42.7	203.9
1999	9.6	*	*	*	*	*	0.6	2.0	5.0	12.4	41.4	195.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
Non-Hispanic black, both sexes												
2006 5/	17.9	*	*	*	*	2.6	9.2	25.9	42.7	66.2	119.0	304.2
2005 5/	18.3	*	*	*	0.4	2.6	9.8	25.6	42.0	64.0	129.0	341.9
2004 5/	18.2	*	*	*	0.3	2.8	9.5	26.7	45.8	65.4	129.1	318.6
2003 5/	18.5	*	*	*	0.4	2.6	9.8	28.5	45.1	70.3	128.4	334.9
2002	17.7	*	*	*	*	2.9	8.8	24.8	45.1	68.3	134.2	322.1
2001	16.5	*	*	*	*	2.2	8.5	23.2	39.4	66.1	125.6	328.7
2000	16.6	*	*	*	*	2.2	7.5	21.6	43.1	67.6	132.7	344.1
1999	16.0	*	*	*	0.4	1.9	8.1	21.3	40.1	67.4	132.7	308.5
Non-Hispanic black, male												
2006 5/	19.5	*	*	*	*	3.7	11.3	34.3	59.6	90.2	147.1	278.5
2005 5/	19.1	*	*	*	*	3.0	13.2	34.7	59.1	82.6	138.8	304.5
2004 5/	19.0	*	*	*	*	3.7	12.1	34.0	65.4	87.5	133.8	271.3
2003 5/	18.7	*	*	*	*	3.2	11.5	35.6	57.5	93.4	139.1	291.9
2002	18.0	*	*	*	*	4.1	11.7	31.9	61.9	84.0	141.8	264.7
2001	16.3	*	*	*	*	2.7	10.5	30.4	50.1	79.8	137.4	288.7
2000	16.1	*	*	*	*	2.7	9.2	28.4	56.0	81.9	139.5	275.9
1999	15.6	*	*	*	*	2.2	10.3	26.8	51.1	80.6	143.2	266.2
Non-Hispanic black, female												
2006 5/	16.5	*	*	*	*	1.5	7.3	18.7	29.1	49.0	103.0	314.6
2005 5/	17.5	*	*	*	*	2.3	6.9	17.9	28.2	50.7	123.5	356.9
2004 5/	17.5	*	*	*	*	2.0	7.3	20.4	30.1	49.6	126.4	337.3
2003 5/	18.4	*	*	*	*	2.1	8.2	22.4	35.1	53.9	122.4	351.6
2002	17.4	*	*	*	*	1.8	6.2	18.7	31.7	57.1	129.8	344.1
2001	16.7	*	*	*	*	1.7	6.7	17.1	30.8	56.5	118.9	343.5
2000	17.1	*	*	*	*	1.8	5.9	15.8	32.7	57.5	128.9	369.4
1999	16.4	*	*	*	*	1.6	6.1	16.5	31.3	58.1	126.8	324.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
All origins 3/, both sexes												
2006	1.0	*	*	*	*	0.1	0.2	0.4	0.8	1.9	6.2	22.6
2005	1.1	*	*	*	*	0.1	0.2	0.4	0.8	2.3	7.0	25.3
2004	1.0	*	*	*	*	0.1	0.2	0.5	0.9	2.1	6.6	24.8
2003	1.1	*	*	*	*	0.1	0.2	0.5	1.0	2.2	7.0	25.6
2002	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.8	23.5
2001	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.3	6.8	23.3
2000	1.0	*	*	*	*	0.1	0.2	0.4	0.9	2.3	7.0	24.3
1999	1.2	*	*	*	*	0.1	0.3	0.4	1.1	2.6	8.9	29.1
All origins 3/, male												
2006	0.9	*	*	*	*	0.1	0.3	0.5	1.1	2.0	6.9	23.8
2005	0.9	*	*	*	*	0.1	0.2	0.6	1.0	2.5	7.2	27.4
2004	0.9	*	*	*	*	*	0.3	0.6	1.1	2.2	7.2	24.3
2003	0.9	*	*	*	*	*	0.3	0.6	1.2	2.3	7.4	26.7
2002	0.9	*	*	*	*	*	0.2	0.5	1.2	2.4	7.0	23.1
2001	0.8	*	*	*	*	*	0.3	0.5	1.0	2.5	6.8	24.2
2000	0.8	*	*	*	*	*	0.2	0.5	0.9	2.5	7.2	27.8
1999	1.0	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.3
All origins 3/, female												
2006	1.1	*	*	*	*	*	0.1	0.4	0.6	1.8	5.7	22.1
2005	1.2	*	*	*	*	*	0.1	0.3	0.6	2.1	6.8	24.3
2004	1.2	*	*	*	*	*	0.1	0.4	0.7	1.9	6.2	25.0
2003	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.8	25.2
2002	1.2	*	*	*	*	*	0.1	0.3	0.8	2.0	6.7	23.7
2001	1.1	*	*	*	*	*	0.2	0.2	0.9	2.1	6.8	22.9
2000	1.1	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.9
1999	1.4	*	*	*	*	*	0.2	0.3	0.9	2.4	8.5	28.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
Hispanic, both sexes												
2006	0.3	*	*	*	*	*	*	*	0.9	2.3	4.9	12.9
2005	0.5	*	*	*	*	*	*	*	1.2	3.3	7.1	22.2
2004	0.5	*	*	*	*	*	*	*	1.3	1.8	8.9	20.0
2003	0.4	*	*	*	*	*	*	*	*	2.7	5.9	25.4
2002	0.4	*	*	*	*	*	*	*	1.3	2.9	8.5	22.7
2001	0.4	*	*	*	*	*	*	*	1.1	3.0	7.4	26.3
2000	0.4	*	*	*	*	*	*	*	*	2.5	6.3	25.2
1999	0.4	*	*	*	*	*	*	*	*	3.1	8.1	23.2
Hispanic, male												
2006	0.4	*	*	*	*	*	*	*	*	*	*	*
2005	0.4	*	*	*	*	*	*	*	*	*	8.4	*
2004	0.4	*	*	*	*	*	*	*	*	*	10.4	*
2003	0.4	*	*	*	*	*	*	*	*	*	*	35.4
2002	0.4	*	*	*	*	*	*	*	*	*	8.2	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.3	*	*	*	*	*	*	*	*	*	*	12.6
2005	0.5	*	*	*	*	*	*	*	*	3.3	6.3	22.8
2004	0.5	*	*	*	*	*	*	*	*	*	7.9	18.0
2003	0.4	*	*	*	*	*	*	*	*	3.1	*	20.3
2002	0.5	*	*	*	*	*	*	*	*	*	8.7	21.6
2001	0.5	*	*	*	*	*	*	*	*	*	7.3	24.2
2000	0.4	*	*	*	*	*	*	*	*	*	*	26.7
1999	0.5	*	*	*	*	*	*	*	*	*	8.7	20.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
Non-Hispanic 4/, both sexes												
2006	1.1	*	*	*	*	0.1	0.2	0.4	0.8	1.9	6.2	23.0
2005	1.2	*	*	*	*	0.1	0.2	0.5	0.8	2.2	6.9	25.4
2004	1.1	*	*	*	*	*	0.2	0.5	0.9	2.1	6.5	24.9
2003	1.2	*	*	*	*	0.1	0.2	0.5	1.0	2.2	7.1	25.6
2002	1.1	*	*	*	*	0.1	0.2	0.4	1.0	2.1	6.7	23.5
2001	1.1	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.7	23.1
2000	1.1	*	*	*	*	0.1	0.2	0.4	0.9	2.2	7.0	24.2
1999	1.3	*	*	*	*	0.1	0.3	0.4	1.1	2.5	8.9	29.3
Non-Hispanic 4/, male												
2006	1.0	*	*	*	*	0.1	0.3	0.5	1.0	2.0	6.9	24.3
2005	1.0	*	*	*	*	0.1	0.2	0.6	0.9	2.4	7.0	27.7
2004	1.0	*	*	*	*	*	0.3	0.6	1.1	2.2	7.0	24.4
2003	1.0	*	*	*	*	*	0.3	0.7	1.2	2.3	7.4	26.3
2002	0.9	*	*	*	*	*	0.2	0.6	1.2	2.3	6.9	22.9
2001	0.9	*	*	*	*	*	0.3	0.5	1.0	2.4	6.7	23.8
2000	0.9	*	*	*	*	*	0.2	0.5	0.9	2.4	7.0	27.9
1999	1.1	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.4
Non-Hispanic 4/, female												
2006	1.2	*	*	*	*	*	0.1	0.4	0.6	1.7	5.8	22.5
2005	1.3	*	*	*	*	*	0.2	0.3	0.6	2.0	6.8	24.4
2004	1.3	*	*	*	*	*	0.1	0.4	0.7	1.9	6.1	25.2
2003	1.3	*	*	*	*	*	0.1	0.4	0.7	2.1	6.8	25.3
2002	1.2	*	*	*	*	*	0.2	0.3	0.8	2.0	6.5	23.7
2001	1.2	*	*	*	*	*	0.2	0.2	0.9	2.0	6.7	22.8
2000	1.2	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.8
1999	1.5	*	*	*	*	*	0.2	0.3	0.9	2.4	8.4	28.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
Non-Hispanic white, both sexes												
2006 5/	0.9	*	*	*	*	*	*	0.2	0.4	1.1	5.1	21.3
2005 5/	1.0	*	*	*	*	*	*	0.2	0.3	1.3	5.5	24.0
2004 5/	0.9	*	*	*	*	*	*	0.1	0.4	1.3	5.3	23.0
2003 5/	1.0	*	*	*	*	*	*	0.1	0.4	1.3	5.9	23.2
2002	0.9	*	*	*	*	*	*	0.1	0.4	1.2	5.6	20.8
2001	0.9	*	*	*	*	*	*	0.1	0.5	1.4	5.6	20.7
2000	0.9	*	*	*	*	*	0.1	0.1	0.4	1.5	5.6	21.8
1999	1.1	*	*	*	*	*	*	0.2	0.5	1.4	7.3	26.1
Non-Hispanic white, male												
2006 5/	0.8	*	*	*	*	*	*	0.2	0.4	1.1	5.6	22.9
2005 5/	0.8	*	*	*	*	*	*	0.2	0.4	1.4	5.6	25.7
2004 5/	0.8	*	*	*	*	*	*	0.2	0.5	1.5	5.8	22.8
2003 5/	0.8	*	*	*	*	*	*	0.2	0.5	1.6	6.6	23.6
2002	0.7	*	*	*	*	*	*	*	0.5	1.4	5.9	20.2
2001	0.7	*	*	*	*	*	*	0.2	0.4	1.5	5.8	22.2
2000	0.7	*	*	*	*	*	*	0.2	0.4	1.6	6.1	24.3
1999	0.9	*	*	*	*	*	*	0.2	0.5	1.6	8.3	28.7
Non-Hispanic white, female												
2006 5/	1.1	*	*	*	*	*	*	0.1	0.3	1.1	4.7	20.5
2005 5/	1.2	*	*	*	*	*	*	*	0.2	1.2	5.5	23.2
2004 5/	1.1	*	*	*	*	*	*	*	0.3	1.1	4.9	23.1
2003 5/	1.1	*	*	*	*	*	*	*	0.3	1.1	5.5	23.0
2002	1.1	*	*	*	*	*	*	*	0.3	1.1	5.4	21.0
2001	1.1	*	*	*	*	*	*	*	0.5	1.3	5.4	20.1
2000	1.1	*	*	*	*	*	*	*	0.3	1.4	5.2	20.8
1999	1.2	*	*	*	*	*	*	*	0.4	1.3	6.6	25.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
Non-Hispanic black, both sexes												
2006 5/	2.2	*	*	*	*	*	0.9	2.2	4.5	8.7	18.8	42.5
2005 5/	2.4	*	*	*	*	0.5	1.0	2.5	4.6	9.8	21.8	42.9
2004 5/	2.4	*	*	*	*	*	1.2	2.9	5.0	8.7	20.0	49.2
2003 5/	2.5	*	*	*	*	0.4	1.3	3.1	5.5	9.3	19.9	53.8
2002	2.5	*	*	*	*	*	1.0	2.8	6.2	10.0	19.8	57.2
2001	2.3	*	*	*	*	0.4	1.2	2.1	5.1	10.1	20.2	50.6
2000	2.3	*	*	*	*	*	0.8	2.2	4.7	10.0	22.6	50.9
1999	2.8	*	*	*	*	*	1.5	2.3	6.2	12.7	27.3	66.0
Non-Hispanic black, male												
2006 5/	2.2	*	*	*	*	*	1.3	2.7	6.3	10.7	23.4	40.3
2005 5/	2.4	*	*	*	*	*	1.3	3.4	6.0	11.3	23.5	50.4
2004 5/	2.3	*	*	*	*	*	1.9	3.6	6.3	8.8	23.7	42.7
2003 5/	2.4	*	*	*	*	*	1.8	4.1	7.2	9.1	19.0	51.1
2002	2.3	*	*	*	*	*	1.0	3.9	7.7	10.2	19.3	62.4
2001	2.2	*	*	*	*	*	1.6	3.2	6.0	12.0	18.6	45.0
2000	2.1	*	*	*	*	*	1.0	2.9	5.4	11.1	20.1	64.6
1999	2.6	*	*	*	*	*	1.8	3.3	7.8	13.8	26.0	60.6
Non-Hispanic black, female												
2006 5/	2.1	*	*	*	*	*	*	1.7	3.1	7.2	16.3	43.3
2005 5/	2.3	*	*	*	*	*	0.7	1.7	3.4	8.7	20.8	39.8
2004 5/	2.4	*	*	*	*	*	*	2.3	4.0	8.5	17.9	51.8
2003 5/	2.6	*	*	*	*	*	0.8	2.2	4.2	9.4	20.4	54.9
2002	2.6	*	*	*	*	*	0.9	1.8	5.0	9.9	20.1	55.2
2001	2.4	*	*	*	*	*	0.8	1.2	4.3	8.7	21.1	52.6
2000	2.4	*	*	*	*	*	*	1.6	4.2	9.2	24.0	45.8
1999	3.1	*	*	*	*	*	1.2	1.5	4.8	12.0	28.0	68.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
All origins 3/, both sexes												
2006	142.1	0.5	*	*	0.3	2.6	15.3	57.7	147.9	352.5	959.8	2,904.7
2005	150.4	*	*	*	0.4	2.5	15.6	59.6	154.2	376.2	1,029.8	3,158.1
2004	153.7	*	*	*	0.3	2.7	16.2	59.9	158.2	395.9	1,072.6	3,287.4
2003	165.1	0.5	*	0.1	0.4	2.7	17.0	62.2	169.8	430.2	1,160.7	3,610.2
2002	171.4	0.6	*	0.0	0.3	2.5	17.1	63.8	177.7	458.9	1,220.2	3,775.0
2001	176.3	0.6	*	0.0	0.4	2.8	16.7	64.0	184.0	477.6	1,271.3	3,958.2
2000	183.1	0.8	*	*	0.3	2.3	16.7	66.1	196.7	505.1	1,324.2	4,182.5
1999	189.8	0.7	*	0.1	0.4	2.5	17.2	67.7	203.5	536.6	1,384.8	4,318.1
All origins 3/, male												
2006	152.2	*	*	*	0.5	4.0	23.2	89.1	220.7	494.2	1,269.7	3,303.0
2005	159.0	*	*	*	0.6	3.8	23.6	92.0	229.4	521.7	1,351.5	3,576.8
2004	161.6	*	*	*	0.4	4.0	24.4	92.8	234.8	549.1	1,410.3	3,680.3
2003	172.2	*	*	*	0.5	4.0	25.7	97.2	251.5	598.8	1,524.4	4,005.2
2002	178.4	*	*	*	0.4	3.7	26.1	99.4	263.3	640.1	1,598.9	4,205.3
2001	181.7	*	*	*	0.6	4.1	25.5	99.5	271.6	663.7	1,663.5	4,395.8
2000	188.7	*	*	*	0.4	3.5	25.8	103.5	290.2	705.2	1,736.8	4,701.7
1999	195.4	*	*	*	0.5	3.7	26.6	106.2	299.3	749.8	1,813.7	4,837.5
All origins 3/, female												
2006	132.3	*	*	*	0.2	1.1	7.4	27.4	80.1	232.6	747.9	2,718.3
2005	142.0	*	*	*	0.1	1.2	7.6	28.2	84.2	253.4	811.4	2,965.8
2004	146.1	*	*	*	0.1	1.4	8.0	28.0	87.1	267.3	845.3	3,110.6
2003	158.1	*	*	*	0.3	1.4	8.4	28.3	94.0	289.2	917.7	3,435.5
2002	164.7	*	*	*	0.2	1.3	8.2	29.5	98.5	308.1	968.9	3,588.3
2001	171.2	*	*	*	0.2	1.5	8.1	29.7	103.1	323.4	1,013.6	3,775.1
2000	177.6	*	*	*	0.2	1.2	7.6	29.9	110.4	340.3	1,055.2	3,971.0
1999	184.5	*	*	*	0.2	1.4	7.9	30.7	115.1	362.0	1,106.3	4,108.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Hispanic, both sexes												
2006	47.2	*	*	*	0.3	1.3	7.2	34.3	106.8	286.6	813.3	2,296.7
2005	51.0	*	*	*	0.3	1.3	7.9	35.7	115.4	326.7	910.6	2,528.4
2004	49.6	*	*	*	*	1.2	7.7	35.3	117.4	326.5	906.9	2,613.4
2003	52.1	*	*	*	0.3	1.4	7.5	35.4	122.0	355.0	973.0	2,970.0
2002	54.0	*	*	*	*	1.2	7.6	38.0	135.9	382.0	1,010.8	3,182.9
2001	55.9	*	*	*	*	1.6	6.8	38.7	130.2	400.9	1,099.1	3,620.2
2000	55.9	*	*	*	*	1.1	7.6	39.5	143.6	405.3	1,111.3	3,697.2
1999	58.7	*	*	*	*	1.4	7.4	44.8	150.4	433.5	1,178.0	3,876.2
Hispanic, male												
2006	49.9	*	*	*	*	1.9	10.9	52.7	156.5	394.1	1,022.8	2,453.9
2005	54.2	*	*	*	*	1.9	12.0	55.2	171.0	449.6	1,165.5	2,694.8
2004	52.7	*	*	*	*	1.9	11.9	55.2	175.1	455.7	1,175.3	2,695.4
2003	53.7	*	*	*	0.5	2.1	11.0	55.8	179.7	494.6	1,214.9	2,933.9
2002	56.5	*	*	*	*	1.9	11.4	58.0	204.0	530.4	1,271.7	3,374.9
2001	57.9	*	*	*	*	2.4	10.4	60.3	191.9	559.9	1,393.3	3,828.2
2000	58.2	*	*	*	*	1.6	12.1	62.7	214.8	561.5	1,420.0	3,865.6
1999	60.8	*	*	*	*	2.1	12.2	71.3	218.3	608.5	1,495.9	3,987.7
Hispanic, female												
2006	44.4	*	*	*	*	0.6	2.9	15.5	60.9	199.2	666.6	2,213.2
2005	47.6	*	*	*	*	*	3.2	15.9	64.5	227.5	733.1	2,441.3
2004	46.2	*	*	*	*	*	3.1	15.4	65.0	222.8	721.6	2,571.3
2003	50.4	*	*	*	*	0.6	3.7	15.0	69.8	243.7	806.9	2,988.3
2002	51.4	*	*	*	*	*	3.5	18.1	74.6	263.9	832.2	3,087.8
2001	53.7	*	*	*	*	*	2.8	17.4	75.4	275.4	900.3	3,518.6
2000	53.6	*	*	*	*	*	2.8	16.8	80.6	282.5	904.0	3,614.6
1999	56.6	*	*	*	*	*	2.3	19.0	90.5	296.4	966.5	3,822.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Non-Hispanic 4/, both sexes												
2006	158.2	0.6	*	*	0.3	2.9	16.8	60.3	150.9	356.5	967.2	2,931.2
2005	166.7	*	*	*	0.4	2.8	16.9	62.0	156.9	379.0	1,034.7	3,183.8
2004	170.3	*	*	*	0.3	3.1	17.5	62.3	160.9	399.6	1,079.3	3,312.1
2003	182.5	0.6	*	0.1	0.4	3.0	18.4	64.7	172.8	433.9	1,167.1	3,629.3
2002	189.0	*	*	*	0.3	2.8	18.4	66.1	180.1	462.1	1,226.5	3,787.0
2001	193.6	0.7	*	*	0.4	3.1	18.0	66.1	187.3	480.9	1,274.9	3,959.4
2000	200.5	0.8	*	*	0.3	2.6	17.7	68.1	199.5	509.0	1,328.8	4,187.0
1999	207.1	0.8	*	0.1	0.4	2.7	18.3	69.4	206.3	540.3	1,388.1	4,319.6
Non-Hispanic 4/, male												
2006	170.6	*	*	*	0.5	4.6	25.5	93.2	225.4	500.0	1,282.9	3,344.5
2005	177.2	*	*	*	0.6	4.3	25.7	95.9	233.5	525.5	1,359.6	3,618.5
2004	179.9	*	*	*	0.5	4.6	26.4	96.6	238.7	553.4	1,419.8	3,722.6
2003	191.5	*	*	*	0.5	4.5	27.9	101.1	255.9	603.6	1,536.0	4,045.5
2002	197.6	*	*	*	0.5	4.1	28.3	103.1	266.5	644.3	1,609.7	4,228.4
2001	200.4	*	*	*	0.6	4.5	27.5	102.8	276.1	667.7	1,670.1	4,408.5
2000	207.6	*	*	*	0.4	3.9	27.4	106.7	294.0	710.4	1,744.6	4,721.1
1999	214.1	*	*	*	0.5	4.0	28.3	108.8	303.4	754.2	1,820.2	4,854.5
Non-Hispanic 4/, female												
2006	146.4	*	*	*	0.2	1.3	8.2	28.7	81.6	234.7	751.6	2,739.1
2005	156.7	*	*	*	0.1	1.4	8.3	29.4	85.7	254.9	814.5	2,985.4
2004	161.1	*	*	*	0.2	1.6	8.7	29.3	88.6	269.9	850.4	3,128.5
2003	173.9	*	*	*	0.3	1.5	9.0	29.6	95.6	291.5	920.9	3,446.4
2002	180.7	*	*	*	0.2	1.5	8.8	30.5	100.0	309.9	972.5	3,596.5
2001	187.2	*	*	*	0.2	1.6	8.7	30.8	105.0	325.6	1,015.5	3,772.7
2000	193.8	*	*	*	0.2	1.3	8.2	31.0	112.1	342.8	1,057.9	3,971.0
1999	200.6	*	*	*	0.3	1.5	8.5	31.5	116.6	364.6	1,107.8	4,105.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Non-Hispanic white, both sexes												
2006 5/	174.5	*	*	*	0.3	2.8	16.5	57.4	144.0	345.7	966.4	2,992.4
2005 5/	183.9	*	*	*	0.3	2.6	16.4	58.8	149.0	369.5	1,033.7	3,253.9
2004 5/	187.7	*	*	*	0.2	2.8	17.2	59.1	153.5	388.1	1,079.6	3,369.7
2003 5/	200.8	*	*	*	0.4	2.8	17.9	61.1	164.5	421.6	1,164.6	3,691.6
2002	208.0	*	*	*	0.3	2.6	17.9	62.8	171.2	450.8	1,223.1	3,848.2
2001	212.6	*	*	*	0.3	2.9	17.3	62.6	177.8	467.8	1,271.3	4,022.8
2000	220.2	*	*	*	0.3	2.4	16.9	64.6	189.5	496.4	1,324.5	4,261.3
1999	227.2	*	*	*	0.3	2.5	17.3	65.8	196.4	527.7	1,386.5	4,390.2
Non-Hispanic white, male												
2006 5/	187.5	*	*	*	0.5	4.4	25.6	90.2	216.7	488.6	1,288.8	3,441.2
2005 5/	195.1	*	*	*	0.5	4.0	25.4	92.4	224.3	517.2	1,367.9	3,732.9
2004 5/	198.0	*	*	*	0.3	4.3	26.4	93.4	230.4	543.1	1,431.6	3,814.0
2003 5/	210.2	*	*	*	0.4	4.3	27.5	97.2	247.3	591.8	1,541.1	4,149.5
2002	217.2	*	*	*	0.4	4.0	28.0	100.0	256.8	636.0	1,618.3	4,324.5
2001	219.6	*	*	*	0.5	4.3	26.9	99.5	266.3	656.5	1,678.7	4,506.9
2000	227.6	*	*	*	0.4	3.7	26.8	103.5	283.5	701.7	1,752.3	4,833.4
1999	234.2	*	*	*	0.5	3.6	27.2	105.1	292.3	745.2	1,834.9	4,954.1
Non-Hispanic white, female												
2006 5/	161.9	*	*	*	*	1.1	7.3	25.1	74.6	221.9	742.7	2,783.1
2005 5/	173.1	*	*	*	*	1.3	7.4	25.7	77.3	241.8	803.8	3,034.8
2004 5/	177.7	*	*	*	*	1.3	7.9	25.4	80.4	254.7	839.7	3,170.6
2003 5/	191.8	*	*	*	0.3	1.4	8.2	25.5	85.9	275.9	910.4	3,490.0
2002	199.1	*	*	*	0.2	1.3	7.8	26.1	90.2	293.0	958.3	3,642.5
2001	205.8	*	*	*	0.2	1.4	7.6	26.4	94.2	307.7	1,001.1	3,821.4
2000	213.1	*	*	*	0.2	1.0	6.9	26.4	100.9	323.6	1,043.3	4,030.7
1999	220.5	*	*	*	*	1.4	7.4	27.2	106.1	345.6	1,092.9	4,165.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Non-Hispanic black, both sexes												
2006 5/	116.5	*	*	*	0.5	4.3	23.1	92.8	238.7	516.6	1,143.3	2,640.8
2005 5/	121.7	*	*	*	0.6	4.1	24.0	96.3	255.2	530.1	1,226.6	2,830.2
2004 5/	124.8	*	*	*	0.7	4.9	24.0	97.9	259.1	576.3	1,255.5	3,038.5
2003 5/	133.2	*	*	*	0.7	4.4	25.5	101.8	277.3	622.3	1,375.4	3,328.7
2002	137.0	*	*	*	0.5	4.1	25.8	103.6	292.4	643.9	1,433.9	3,481.1
2001	141.2	*	*	*	0.9	4.8	26.4	104.6	302.1	681.4	1,487.0	3,628.7
2000	144.4	*	*	*	0.4	4.4	26.1	107.7	318.9	710.1	1,534.5	3,701.7
1999	147.9	*	*	*	0.8	4.1	28.2	110.8	324.7	741.6	1,559.0	3,817.9
Non-Hispanic black, male												
2006 5/	124.5	*	*	*	*	6.3	32.2	133.8	347.2	717.2	1,476.8	2,685.6
2005 5/	127.0	*	*	*	0.8	6.3	33.6	140.2	364.5	712.6	1,542.6	2,808.9
2004 5/	129.2	*	*	*	1.2	6.7	32.9	141.2	368.2	771.0	1,551.8	3,133.7
2003 5/	137.6	*	*	*	0.9	6.4	36.2	148.3	387.0	836.1	1,758.6	3,309.2
2002	139.5	*	*	*	*	5.3	36.2	146.9	413.2	850.8	1,762.3	3,572.8
2001	143.2	*	*	*	1.2	6.8	37.1	149.0	418.0	903.1	1,834.9	3,791.8
2000	145.7	*	*	*	*	6.1	36.3	152.2	444.2	916.5	1,887.6	3,911.2
1999	150.0	*	*	*	0.9	6.0	40.3	160.0	458.4	967.5	1,873.3	4,082.5
Non-Hispanic black, female												
2006 5/	109.2	*	*	*	*	2.5	15.2	57.7	151.4	373.4	954.0	2,622.6
2005 5/	116.8	*	*	*	*	2.1	15.5	58.7	167.2	399.7	1,047.6	2,838.7
2004 5/	120.8	*	*	*	*	3.2	16.0	60.7	171.5	437.6	1,088.3	3,000.9
2003 5/	129.1	*	*	*	*	2.6	16.0	62.0	189.2	470.4	1,159.1	3,336.2
2002	134.8	*	*	*	*	2.9	16.7	66.5	195.4	497.1	1,248.1	3,446.0
2001	139.4	*	*	*	*	2.9	16.9	66.5	208.9	525.0	1,291.6	3,568.3
2000	143.2	*	*	*	*	2.9	17.1	69.4	218.1	565.0	1,335.7	3,623.9
1999	146.0	*	*	*	0.7	2.4	17.4	68.7	217.5	582.3	1,383.0	3,719.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
All origins 3/, both sexes												
2006	47.2	*	*	*	0.1	1.0	6.1	22.5	58.2	131.1	319.0	825.8
2005	50.9	*	*	*	0.2	1.0	6.2	23.7	61.1	143.1	348.2	923.1
2004	53.4	*	*	*	0.1	1.1	6.4	24.2	64.5	153.4	370.3	992.9
2003	58.7	*	*	*	0.2	1.0	7.0	25.5	70.9	168.6	412.7	1,118.8
2002	62.3	*	*	*	0.2	1.0	7.0	26.5	74.9	186.1	444.4	1,191.3
2001	64.9	*	*	*	0.2	1.2	7.1	27.6	78.4	193.2	471.2	1,265.4
2000	68.5	0.5	*	*	0.1	0.9	6.9	28.9	86.6	209.1	503.3	1,338.5
1999	71.5	*	*	*	0.2	1.1	7.2	29.6	90.0	225.0	529.3	1,379.3
All origins 3/, male												
2006	51.6	*	*	*	0.2	1.5	9.2	34.2	85.6	181.9	411.8	936.3
2005	54.8	*	*	*	0.2	1.5	9.2	36.2	89.9	195.5	445.9	1,045.0
2004	57.4	*	*	*	0.2	1.6	9.6	37.3	95.3	209.5	477.0	1,120.5
2003	62.6	*	*	*	0.2	1.5	10.4	39.8	104.1	231.0	530.6	1,263.5
2002	66.2	*	*	*	0.2	1.4	10.5	40.6	109.2	256.2	569.2	1,366.4
2001	68.6	*	*	*	0.2	1.7	10.6	42.4	114.0	264.7	604.1	1,461.3
2000	72.7	*	*	*	0.1	1.3	10.6	44.7	125.7	286.9	648.3	1,568.3
1999	76.1	*	*	*	0.2	1.5	11.0	46.3	131.8	310.0	684.4	1,622.1
All origins 3/, female												
2006	43.0	*	*	*	*	0.5	3.0	11.2	32.6	88.1	255.6	774.1
2005	47.2	*	*	*	*	0.5	3.2	11.6	34.3	98.9	281.8	867.1
2004	49.6	*	*	*	*	0.6	3.3	11.6	35.9	106.3	298.5	935.5
2003	54.8	*	*	*	0.1	0.6	3.6	11.7	40.0	116.5	333.9	1,054.8
2002	58.4	*	*	*	0.1	0.5	3.6	12.9	43.2	127.7	361.5	1,115.3
2001	61.3	*	*	*	*	0.7	3.5	13.3	45.4	134.0	383.9	1,183.5
2000	64.6	*	*	*	0.1	0.5	3.3	13.6	50.5	145.1	408.7	1,244.8
1999	67.0	*	*	*	0.1	0.6	3.4	13.5	51.5	155.4	428.6	1,281.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Hispanic, both sexes												
2006	15.7	*	*	*	*	0.5	2.7	11.5	37.7	100.5	273.6	692.9
2005	17.6	*	*	*	*	0.5	2.7	12.5	43.2	119.4	318.7	787.2
2004	17.3	*	*	*	*	0.5	2.9	12.9	44.2	120.4	315.7	816.8
2003	18.5	*	*	*	*	0.5	3.0	13.3	44.5	137.7	341.6	952.0
2002	19.7	*	*	*	*	0.5	3.0	14.2	55.1	146.2	364.3	1,048.2
2001	20.0	*	*	*	*	0.7	2.5	14.8	49.5	149.8	394.9	1,185.5
2000	20.1	*	*	*	*	0.5	3.0	15.5	55.0	154.3	404.7	1,177.1
1999	21.1	*	*	*	*	0.5	2.9	17.8	56.1	164.0	433.7	1,227.8
Hispanic, male												
2006	16.5	*	*	*	*	0.7	3.9	16.7	52.7	137.7	338.6	742.0
2005	18.8	*	*	*	*	0.6	3.8	18.9	62.2	160.9	406.2	875.4
2004	18.4	*	*	*	*	0.6	4.3	20.1	65.4	164.1	403.4	840.6
2003	18.9	*	*	*	*	0.7	4.1	20.6	64.1	187.5	409.2	963.6
2002	20.6	*	*	*	*	0.7	4.2	21.0	80.8	199.3	460.0	1,107.7
2001	20.9	*	*	*	*	1.0	3.9	22.7	69.4	206.9	488.7	1,331.3
2000	21.1	*	*	*	*	0.7	4.7	23.4	80.0	207.6	514.8	1,273.8
1999	22.0	*	*	*	*	0.8	4.6	26.7	79.0	225.8	558.9	1,276.8
Hispanic, female												
2006	14.9	*	*	*	*	*	1.4	6.2	23.9	70.4	228.2	666.8
2005	16.4	*	*	*	*	*	1.6	6.0	25.8	85.8	257.7	741.0
2004	16.0	*	*	*	*	*	1.3	5.6	24.9	85.4	255.1	804.5
2003	18.0	*	*	*	*	*	1.7	6.0	26.8	98.1	295.2	946.1
2002	18.7	*	*	*	*	*	1.8	7.4	32.1	103.9	298.8	1,018.8
2001	19.1	*	*	*	*	*	1.1	7.0	31.8	104.7	331.5	1,114.2
2000	19.1	*	*	*	*	*	1.2	7.8	32.9	112.3	330.8	1,129.7
1999	20.1	*	*	*	*	*	1.1	9.1	35.9	115.5	350.4	1,204.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Non-Hispanic 4/, both sexes												
2006	52.6	*	*	*	0.1	1.2	6.7	23.7	59.8	133.1	321.3	831.0
2005	56.5	*	*	*	0.2	1.1	6.8	24.9	62.5	144.7	349.4	928.4
2004	59.2	*	*	*	0.1	1.3	7.0	25.4	66.0	155.4	372.7	999.2
2003	64.9	*	*	*	0.2	1.2	7.6	26.7	72.8	170.4	415.5	1,123.8
2002	68.7	*	*	*	0.2	1.1	7.6	27.7	76.3	188.2	447.3	1,194.0
2001	71.4	*	*	*	0.2	1.3	7.6	28.8	80.3	195.6	473.6	1,265.3
2000	75.3	*	*	*	0.1	1.0	7.4	30.0	88.6	211.9	506.3	1,340.8
1999	78.2	*	*	*	0.2	1.2	7.7	30.5	92.2	227.9	531.9	1,381.0
Non-Hispanic 4/, male												
2006	57.9	*	*	*	0.2	1.8	10.2	36.2	88.2	184.6	415.9	945.2
2005	61.2	*	*	*	0.3	1.7	10.2	38.0	92.0	197.8	447.5	1,052.8
2004	64.0	*	*	*	0.2	1.8	10.5	39.1	97.5	212.1	480.2	1,132.5
2003	69.7	*	*	*	0.2	1.7	11.4	41.8	106.9	233.4	535.7	1,274.5
2002	73.5	*	*	*	0.2	1.6	11.5	42.6	111.3	259.2	573.4	1,374.2
2001	75.9	*	*	*	0.2	1.8	11.6	44.2	117.0	267.7	607.9	1,463.9
2000	80.3	*	*	*	0.2	1.5	11.4	46.5	128.7	290.9	652.6	1,576.2
1999	83.8	*	*	*	0.2	1.7	11.8	47.8	135.1	313.8	687.7	1,631.5
Non-Hispanic 4/, female												
2006	47.6	*	*	*	*	0.6	3.3	11.8	33.4	89.4	256.7	778.0
2005	52.0	*	*	*	*	0.5	3.5	12.2	35.0	99.7	282.9	871.7
2004	54.7	*	*	*	*	0.7	3.6	12.2	36.7	107.7	300.4	939.6
2003	60.3	*	*	*	0.2	0.6	3.8	12.3	41.0	117.5	335.3	1,057.5
2002	64.1	*	*	*	*	0.6	3.8	13.3	44.0	128.9	363.8	1,116.3
2001	67.1	*	*	*	*	0.8	3.8	13.9	46.3	135.7	385.4	1,182.7
2000	70.6	*	*	*	0.1	0.6	3.5	14.1	51.6	146.8	411.0	1,245.6
1999	72.9	*	*	*	*	0.7	3.7	13.9	52.5	157.3	430.7	1,280.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Non-Hispanic white, both sexes												
2006 5/	58.0	*	*	*	0.1	1.1	6.8	23.4	58.8	131.4	321.4	841.4
2005 5/	62.3	*	*	*	0.1	1.0	6.8	24.5	61.3	143.7	349.1	941.9
2004 5/	65.3	*	*	*	0.1	1.2	7.0	24.8	64.9	152.9	374.2	1,010.1
2003 5/	71.5	*	*	*	0.2	1.1	7.5	25.9	71.4	168.3	415.1	1,137.4
2002	75.5	*	*	*	0.1	1.1	7.5	27.0	74.7	186.1	445.9	1,202.8
2001	78.4	*	*	*	0.1	1.2	7.5	28.1	78.7	193.0	473.0	1,276.8
2000	82.7	*	*	*	0.1	0.9	7.3	29.3	86.9	209.3	505.3	1,355.3
1999	86.0	*	*	*	0.1	1.1	7.5	30.0	90.7	226.1	532.4	1,394.2
Non-Hispanic white, male												
2006 5/	64.0	*	*	*	0.2	1.7	10.5	36.3	87.5	184.4	418.6	964.6
2005 5/	67.8	*	*	*	0.3	1.6	10.4	38.2	91.3	198.9	451.0	1,077.9
2004 5/	70.9	*	*	*	*	1.9	10.7	38.8	97.3	210.8	487.4	1,156.5
2003 5/	77.0	*	*	*	0.2	1.6	11.5	41.2	106.5	232.9	537.8	1,300.9
2002	81.2	*	*	*	0.2	1.6	11.5	42.5	110.8	260.2	575.8	1,393.0
2001	83.7	*	*	*	0.2	1.7	11.6	44.3	116.8	267.7	612.5	1,483.2
2000	88.5	*	*	*	0.2	1.4	11.4	46.7	128.0	291.0	656.7	1,600.7
1999	92.4	*	*	*	0.2	1.5	11.8	48.1	134.6	315.0	695.7	1,653.8
Non-Hispanic white, female												
2006 5/	52.2	*	*	*	*	0.5	3.1	10.7	31.6	85.4	254.0	783.9
2005 5/	57.0	*	*	*	*	0.4	3.2	11.1	32.7	95.9	279.1	879.7
2004 5/	59.9	*	*	*	*	0.6	3.3	10.9	34.2	103.1	297.1	944.5
2003 5/	66.2	*	*	*	*	0.6	3.5	10.8	38.0	113.0	332.2	1,065.3
2002	70.1	*	*	*	*	0.5	3.6	11.8	40.6	122.9	358.8	1,120.7
2001	73.4	*	*	*	*	0.8	3.4	12.2	42.7	129.6	380.6	1,190.9
2000	77.2	*	*	*	*	0.5	3.1	12.3	48.2	140.5	405.9	1,256.3
1999	79.8	*	*	*	*	0.7	3.3	12.3	49.3	151.8	425.5	1,290.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Non-Hispanic black, both sexes												
2006 5/	38.7	*	*	*	*	1.7	8.5	32.4	82.5	175.7	374.2	811.7
2005 5/	41.1	*	*	*	*	1.8	9.1	33.6	86.9	182.8	413.4	901.2
2004 5/	43.4	*	*	*	0.3	1.8	9.1	36.2	93.0	208.2	424.3	991.3
2003 5/	47.3	*	*	*	*	1.7	10.4	38.6	100.6	223.1	486.5	1,103.6
2002	50.2	*	*	*	*	1.7	10.0	39.0	109.1	242.6	523.0	1,212.9
2001	51.6	*	*	*	0.4	1.9	10.2	40.2	111.6	256.0	543.7	1,247.0
2000	54.0	*	*	*	*	1.8	10.1	41.9	122.3	273.0	576.6	1,281.8
1999	55.1	*	*	*	*	1.7	10.5	42.0	123.2	282.4	589.9	1,328.6
Non-Hispanic black, male												
2006 5/	40.6	*	*	*	*	2.5	11.9	45.1	117.3	233.0	463.5	824.6
2005 5/	41.8	*	*	*	*	2.6	12.2	46.6	120.2	233.2	499.2	898.3
2004 5/	43.9	*	*	*	*	2.1	12.0	51.6	126.2	269.9	491.7	1,020.2
2003 5/	48.2	*	*	*	*	2.4	14.3	54.7	135.9	287.1	606.9	1,121.9
2002	50.1	*	*	*	*	2.2	13.7	53.4	144.5	305.4	631.3	1,296.3
2001	50.9	*	*	*	*	2.5	13.5	53.4	146.1	320.8	646.6	1,389.4
2000	53.7	*	*	*	*	2.2	13.6	56.3	163.1	341.2	693.3	1,426.6
1999	55.2	*	*	*	*	2.5	14.5	58.1	168.7	357.0	691.4	1,509.4
Non-Hispanic black, female												
2006 5/	37.0	*	*	*	*	1.0	5.5	21.6	54.5	134.8	323.4	806.4
2005 5/	40.5	*	*	*	*	1.0	6.3	22.5	60.2	146.8	364.8	902.3
2004 5/	43.0	*	*	*	*	1.5	6.5	23.0	66.2	164.3	386.3	979.8
2003 5/	46.5	*	*	*	*	1.0	6.9	24.7	72.2	177.6	418.5	1,096.5
2002	50.4	*	*	*	*	1.3	6.7	26.7	80.7	198.0	461.8	1,181.0
2001	52.3	*	*	*	*	1.3	7.4	28.8	83.8	210.3	485.9	1,194.3
2000	54.2	*	*	*	*	1.4	7.0	29.6	89.5	225.0	510.9	1,228.0
1999	54.9	*	*	*	*	1.0	7.0	28.2	86.8	229.7	533.1	1,261.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
All origins 3/, both sexes												
2006	1.3	*	*	*	*	0.1	0.2	0.8	1.9	3.6	8.0	21.5
2005	1.2	*	*	*	*	*	0.2	0.8	1.7	3.3	7.2	20.8
2004	0.8	*	*	*	*	*	0.2	0.5	1.2	2.2	5.2	14.8
2003	1.1	*	*	*	*	*	0.2	0.7	1.4	3.2	6.5	20.0
2002	1.2	*	*	*	*	*	0.2	0.9	1.7	3.5	7.1	21.1
2001	1.2	*	*	*	*	*	0.2	0.8	1.8	3.5	6.9	21.5
2000	1.2	*	*	*	*	*	0.2	0.7	2.0	3.5	7.3	21.9
1999	1.2	*	*	*	*	*	0.2	0.7	1.9	3.7	7.0	21.9
All origins 3/, male												
2006	1.4	*	*	*	*	*	0.2	1.2	2.6	4.6	9.6	21.8
2005	1.2	*	*	*	*	*	0.3	1.2	2.5	4.3	8.5	20.2
2004	0.9	*	*	*	*	*	0.3	0.7	1.8	3.0	6.3	16.3
2003	1.1	*	*	*	*	*	0.3	1.0	2.1	4.3	8.0	19.5
2002	1.2	*	*	*	*	*	0.2	1.3	2.5	4.7	8.7	21.4
2001	1.2	*	*	*	*	*	0.3	1.2	2.6	4.7	8.5	20.7
2000	1.3	*	*	*	*	*	0.3	1.2	2.9	4.8	9.1	20.7
1999	1.2	*	*	*	*	*	0.3	1.0	2.8	4.9	8.8	22.3
All origins 3/, female												
2006	1.3	*	*	*	*	*	0.2	0.5	1.2	2.7	6.8	21.4
2005	1.2	*	*	*	*	*	0.1	0.5	0.9	2.4	6.2	21.1
2004	0.8	*	*	*	*	*	0.1	0.3	0.7	1.4	4.4	14.1
2003	1.0	*	*	*	*	*	0.1	0.4	0.9	2.2	5.5	20.2
2002	1.1	*	*	*	*	*	0.1	0.5	1.0	2.4	5.9	21.0
2001	1.1	*	*	*	*	*	0.1	0.4	1.1	2.5	5.9	21.8
2000	1.1	*	*	*	*	*	0.1	0.3	1.1	2.4	6.1	22.4
1999	1.1	*	*	*	*	*	*	0.4	1.1	2.6	5.8	21.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
Hispanic, both sexes												
2006	0.2	*	*	*	*	*	*	*	*	1.5	3.6	8.6
2005	0.2	*	*	*	*	*	*	*	*	*	3.2	10.7
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.2	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.2	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
Non-Hispanic 4/, both sexes												
2006	1.5	*	*	*	*	*	0.2	0.9	2.0	3.7	8.2	22.1
2005	1.4	*	*	*	*	*	0.2	0.9	1.8	3.4	7.4	21.2
2004	1.0	*	*	*	*	*	0.2	0.6	1.3	2.3	5.4	15.3
2003	1.2	*	*	*	*	*	0.3	0.7	1.5	3.3	6.7	20.6
2002	1.3	*	*	*	*	*	0.2	0.9	1.8	3.6	7.3	21.8
2001	1.3	*	*	*	*	*	0.2	0.8	1.9	3.6	7.1	22.0
2000	1.3	*	*	*	*	*	0.2	0.8	2.1	3.6	7.5	22.2
1999	1.3	*	*	*	*	*	0.2	0.7	2.0	3.8	7.2	22.2
Non-Hispanic 4/, male												
2006	1.6	*	*	*	*	*	0.3	1.3	2.8	4.8	10.0	22.6
2005	1.4	*	*	*	*	*	0.3	1.3	2.7	4.4	8.7	20.9
2004	1.0	*	*	*	*	*	0.3	0.8	1.9	3.2	6.5	17.0
2003	1.3	*	*	*	*	*	0.4	1.1	2.2	4.5	8.4	20.2
2002	1.4	*	*	*	*	*	0.3	1.4	2.6	4.9	9.0	22.1
2001	1.4	*	*	*	*	*	0.3	1.3	2.7	4.8	8.7	20.9
2000	1.4	*	*	*	*	*	0.3	1.3	3.1	4.9	9.3	21.0
1999	1.4	*	*	*	*	*	0.3	1.1	2.9	5.2	9.0	22.9
Non-Hispanic 4/, female												
2006	1.4	*	*	*	*	*	0.2	0.5	1.3	2.8	7.0	21.9
2005	1.3	*	*	*	*	*	0.1	0.5	1.0	2.5	6.5	21.4
2004	0.9	*	*	*	*	*	0.1	0.3	0.8	1.5	4.6	14.6
2003	1.2	*	*	*	*	*	0.1	0.4	0.9	2.3	5.7	20.7
2002	1.3	*	*	*	*	*	0.1	0.5	1.1	2.5	6.1	21.6
2001	1.3	*	*	*	*	*	0.1	0.4	1.1	2.6	6.0	22.4
2000	1.2	*	*	*	*	*	0.1	0.3	1.2	2.5	6.3	22.7
1999	1.2	*	*	*	*	*	*	0.4	1.1	2.7	6.0	21.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
Non-Hispanic white, both sexes												
2006 5/	1.6	*	*	*	*	*	0.2	0.8	1.9	3.5	8.1	22.3
2005 5/	1.4	*	*	*	*	*	0.2	0.7	1.6	3.2	7.1	20.9
2004 5/	1.0	*	*	*	*	*	0.2	0.5	1.2	2.1	5.1	15.6
2003 5/	1.3	*	*	*	*	*	0.2	0.7	1.4	3.1	6.6	20.9
2002	1.4	*	*	*	*	*	0.2	0.8	1.7	3.5	7.2	22.0
2001	1.4	*	*	*	*	*	0.2	0.8	1.7	3.4	7.0	22.0
2000	1.4	*	*	*	*	*	0.2	0.7	1.9	3.4	7.3	22.5
1999	1.4	*	*	*	*	*	0.2	0.6	1.8	3.6	7.0	22.7
Non-Hispanic white, male												
2006 5/	1.6	*	*	*	*	*	0.2	1.2	2.6	4.6	9.6	22.7
2005 5/	1.5	*	*	*	*	*	0.3	1.1	2.4	4.1	8.4	20.3
2004 5/	1.0	*	*	*	*	*	0.3	0.7	1.7	2.8	6.3	16.9
2003 5/	1.3	*	*	*	*	*	0.4	1.0	2.0	4.4	7.9	20.4
2002	1.5	*	*	*	*	*	0.3	1.2	2.5	4.6	9.0	22.7
2001	1.4	*	*	*	*	*	0.3	1.2	2.5	4.5	8.7	20.9
2000	1.5	*	*	*	*	*	0.3	1.2	2.8	4.6	9.0	21.3
1999	1.4	*	*	*	*	*	0.2	0.9	2.6	4.9	8.7	23.6
Non-Hispanic white, female												
2006 5/	1.5	*	*	*	*	*	0.2	0.4	1.2	2.6	7.0	22.2
2005 5/	1.4	*	*	*	*	*	*	0.3	0.8	2.4	6.3	21.2
2004 5/	1.0	*	*	*	*	*	*	0.3	0.7	1.5	4.3	15.0
2003 5/	1.3	*	*	*	*	*	*	0.4	0.8	2.1	5.7	21.1
2002	1.4	*	*	*	*	*	*	0.5	1.0	2.5	6.0	21.8
2001	1.4	*	*	*	*	*	*	0.3	1.0	2.5	5.9	22.4
2000	1.3	*	*	*	*	*	*	0.3	1.0	2.4	6.1	23.0
1999	1.3	*	*	*	*	*	*	0.3	1.1	2.6	5.8	22.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
Non-Hispanic black, both sexes												
2006 5/	1.4	*	*	*	*	*	0.5	1.7	3.5	6.1	11.3	21.0
2005 5/	1.5	*	*	*	*	*	*	2.0	3.9	5.7	11.8	27.9
2004 5/	1.1	*	*	*	*	*	*	1.4	3.0	4.2	9.1	15.5
2003 5/	1.2	*	*	*	*	*	0.5	1.3	3.0	5.5	10.1	20.2
2002	1.2	*	*	*	*	*	*	1.7	3.0	6.1	9.4	21.5
2001	1.3	*	*	*	*	*	0.5	1.6	4.0	5.8	8.5	24.3
2000	1.3	*	*	*	*	*	0.4	1.5	4.3	5.6	10.8	21.9
1999	1.2	*	*	*	*	*	0.5	1.9	3.6	5.8	10.9	17.5
Non-Hispanic black, male												
2006 5/	1.6	*	*	*	*	*	*	2.3	5.2	8.1	15.6	23.8
2005 5/	1.6	*	*	*	*	*	*	2.4	6.0	8.1	15.7	28.5
2004 5/	1.3	*	*	*	*	*	*	1.9	4.7	7.1	12.0	23.4
2003 5/	1.5	*	*	*	*	*	*	2.0	4.9	7.7	16.4	24.5
2002	1.4	*	*	*	*	*	*	2.8	4.4	9.4	11.7	21.9
2001	1.5	*	*	*	*	*	*	2.5	5.6	8.8	10.5	24.3
2000	1.5	*	*	*	*	*	*	2.3	6.3	8.7	14.2	*
1999	1.5	*	*	*	*	*	0.8	2.7	6.1	8.4	12.8	*
Non-Hispanic black, female												
2006 5/	1.2	*	*	*	*	*	*	1.2	2.1	4.7	8.8	19.8
2005 5/	1.3	*	*	*	*	*	*	1.6	2.2	4.0	9.5	27.6
2004 5/	0.8	*	*	*	*	*	*	0.9	1.7	2.0	7.4	12.4
2003 5/	0.9	*	*	*	*	*	*	*	1.5	3.9	6.5	18.6
2002	1.1	*	*	*	*	*	*	0.8	2.0	3.7	8.1	21.3
2001	1.1	*	*	*	*	*	*	*	2.6	3.8	7.4	24.4
2000	1.1	*	*	*	*	*	*	*	2.7	3.5	8.9	22.2
1999	1.0	*	*	*	*	*	*	1.1	1.6	3.9	9.8	16.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
All origins 3/, both sexes												
2006	93.5	*	*	*	0.2	1.5	9.0	34.4	87.8	217.9	632.8	2,057.4
2005	98.2	*	*	*	0.2	1.5	9.2	35.0	91.4	229.8	674.5	2,214.2
2004	99.5	*	*	*	0.2	1.6	9.6	35.1	92.5	240.3	697.1	2,279.7
2003	105.3	*	*	*	0.2	1.7	9.8	35.9	97.5	258.4	741.5	2,471.4
2002	108.0	*	*	*	0.2	1.4	9.9	36.5	101.0	269.4	768.8	2,562.5
2001	110.3	*	*	*	0.2	1.6	9.5	35.6	103.8	280.9	793.1	2,671.3
2000	113.3	*	*	*	0.2	1.4	9.5	36.4	108.1	292.4	813.6	2,822.1
1999	117.2	*	*	*	0.2	1.4	9.8	37.5	111.5	308.0	848.5	2,917.0
All origins 3/, male												
2006	99.2	*	*	*	0.3	2.4	13.8	53.7	132.4	307.7	848.3	2,344.9
2005	102.9	*	*	*	0.3	2.3	14.1	54.7	136.9	321.9	897.1	2,511.6
2004	103.3	*	*	*	0.2	2.4	14.5	54.8	137.7	336.5	926.9	2,543.6
2003	108.5	*	*	*	0.3	2.5	14.9	56.4	145.4	363.5	985.8	2,722.2
2002	110.9	*	*	*	0.2	2.2	15.4	57.5	151.5	379.3	1,021.0	2,817.4
2001	111.9	*	*	*	0.3	2.4	14.6	55.9	154.9	394.3	1,050.9	2,913.8
2000	114.8	*	*	*	0.2	2.1	14.8	57.7	161.6	413.5	1,079.4	3,112.7
1999	118.0	*	*	*	0.3	2.1	15.3	59.0	164.7	434.9	1,120.5	3,193.1
All origins 3/, female												
2006	88.0	*	*	*	*	0.6	4.2	15.7	46.3	141.9	485.5	1,922.9
2005	93.7	*	*	*	*	0.7	4.3	16.0	49.0	152.2	523.3	2,077.6
2004	95.7	*	*	*	*	0.7	4.6	16.2	50.5	159.6	542.4	2,161.0
2003	102.3	*	*	*	0.1	0.8	4.7	16.2	53.1	170.5	578.3	2,360.5
2002	105.2	*	*	*	*	0.7	4.4	16.2	54.3	178.0	601.4	2,451.9
2001	108.8	*	*	*	0.1	0.7	4.4	16.0	56.7	186.9	623.7	2,569.8
2000	111.9	*	*	*	*	0.6	4.3	16.0	58.8	192.8	640.3	2,703.8
1999	116.4	*	*	*	0.1	0.7	4.3	16.7	62.6	204.0	671.8	2,805.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
Hispanic, both sexes												
2006	31.3	*	*	*	*	0.8	4.4	22.6	68.5	184.6	536.0	1,595.2
2005	33.2	*	*	*	*	0.8	5.1	23.0	71.8	205.9	588.8	1,730.5
2004	32.2	*	*	*	*	0.8	4.8	22.4	72.9	205.5	589.6	1,795.7
2003	33.5	*	*	*	*	0.9	4.6	22.0	77.2	216.4	629.5	2,012.8
2002	34.2	*	*	*	*	0.7	4.5	23.6	80.1	234.6	644.0	2,129.7
2001	35.7	*	*	*	*	0.9	4.2	23.8	80.0	249.4	701.7	2,426.6
2000	35.6	*	*	*	*	0.6	4.6	23.9	88.0	249.7	703.2	2,508.8
1999	37.5	*	*	*	*	0.9	4.5	26.8	93.5	268.8	741.8	2,640.7
Hispanic, male												
2006	33.3	*	*	*	*	1.2	7.0	35.7	103.2	254.5	680.7	1,705.1
2005	35.2	*	*	*	*	1.3	8.2	36.0	108.3	287.2	754.3	1,811.9
2004	34.2	*	*	*	*	1.3	7.5	35.0	109.3	291.0	769.7	1,853.5
2003	34.6	*	*	*	*	1.4	6.9	35.1	115.3	306.1	804.2	1,965.7
2002	35.7	*	*	*	*	1.2	7.2	36.6	122.3	329.1	808.8	2,262.2
2001	36.8	*	*	*	*	1.4	6.5	37.5	121.4	350.2	901.5	2,479.8
2000	36.9	*	*	*	*	0.9	7.4	39.1	134.2	351.6	900.8	2,577.7
1999	38.7	*	*	*	*	1.3	7.6	44.3	138.5	381.8	932.8	2,706.7
Hispanic, female												
2006	29.2	*	*	*	*	*	1.4	9.2	36.6	127.8	434.6	1,536.8
2005	31.0	*	*	*	*	*	1.6	9.7	38.5	140.3	473.5	1,687.9
2004	30.1	*	*	*	*	*	1.8	9.8	39.9	136.9	465.3	1,766.0
2003	32.3	*	*	*	*	*	2.0	8.9	42.8	144.9	509.6	2,036.7
2002	32.6	*	*	*	*	*	1.6	10.5	42.2	159.4	531.2	2,064.1
2001	34.5	*	*	*	*	*	1.7	10.3	43.3	169.9	566.7	2,400.7
2000	34.2	*	*	*	*	*	1.7	8.9	47.2	169.6	570.5	2,475.0
1999	36.3	*	*	*	*	*	1.3	9.7	53.8	180.2	614.8	2,608.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
Non-Hispanic 4/, both sexes												
2006	104.1	*	*	*	0.2	1.7	9.8	35.7	89.1	219.7	637.6	2,078.0
2005	108.9	*	*	*	0.2	1.7	9.9	36.3	92.6	230.9	677.9	2,234.1
2004	110.1	*	*	*	0.2	1.8	10.3	36.3	93.6	241.8	701.2	2,297.5
2003	116.3	*	*	*	0.2	1.8	10.6	37.2	98.5	260.2	744.8	2,484.9
2002	118.9	*	*	*	0.2	1.6	10.6	37.5	102.0	270.3	771.9	2,571.2
2001	120.9	*	*	*	0.2	1.7	10.2	36.5	105.1	281.6	794.2	2,672.2
2000	123.9	*	*	*	0.2	1.5	10.1	37.3	108.8	293.5	815.0	2,824.0
1999	127.6	*	*	*	0.2	1.5	10.4	38.1	112.1	308.6	849.1	2,916.3
Non-Hispanic 4/, male												
2006	111.1	*	*	*	0.3	2.7	15.0	55.7	134.4	310.5	857.1	2,376.7
2005	114.6	*	*	*	0.3	2.6	15.2	56.6	138.8	323.3	903.3	2,544.8
2004	114.9	*	*	*	0.2	2.7	15.6	56.7	139.3	338.1	933.0	2,573.1
2003	120.4	*	*	*	0.3	2.8	16.2	58.3	146.7	365.6	991.9	2,750.7
2002	122.7	*	*	*	0.3	2.4	16.6	59.2	152.6	380.2	1,027.3	2,832.2
2001	123.1	*	*	*	0.3	2.6	15.7	57.3	156.4	395.1	1,053.5	2,923.7
2000	126.0	*	*	*	0.2	2.3	15.7	58.9	162.3	414.7	1,082.7	3,123.9
1999	128.9	*	*	*	0.3	2.3	16.2	59.9	165.4	435.2	1,123.5	3,200.1
Non-Hispanic 4/, female												
2006	97.4	*	*	*	*	0.6	4.7	16.4	47.0	142.6	487.8	1,939.2
2005	103.4	*	*	*	*	0.8	4.7	16.7	49.7	152.8	525.1	2,092.3
2004	105.6	*	*	*	*	0.8	5.0	16.7	51.1	160.7	545.5	2,174.3
2003	112.4	*	*	*	0.1	0.9	5.1	16.9	53.7	171.7	580.0	2,368.1
2002	115.4	*	*	*	*	0.8	4.8	16.7	55.0	178.4	602.6	2,458.6
2001	118.8	*	*	*	0.1	0.8	4.8	16.5	57.5	187.3	624.1	2,567.6
2000	122.0	*	*	*	*	0.7	4.6	16.5	59.3	193.5	640.6	2,702.7
1999	126.4	*	*	*	0.1	0.8	4.7	17.3	62.9	204.6	671.1	2,802.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
Non-Hispanic white, both sexes												
2006 5/	114.9	*	*	*	0.2	1.6	9.4	33.2	83.2	210.8	636.9	2,128.7
2005 5/	120.2	*	*	*	0.2	1.6	9.5	33.6	86.1	222.7	677.4	2,291.1
2004 5/	121.4	*	*	*	0.1	1.6	10.0	33.9	87.4	233.1	700.2	2,343.9
2003 5/	128.0	*	*	*	0.2	1.7	10.2	34.6	91.7	250.2	743.0	2,533.4
2002	131.0	*	*	*	0.2	1.5	10.2	34.9	94.8	261.3	770.0	2,623.4
2001	132.7	*	*	*	0.2	1.6	9.6	33.8	97.4	271.4	791.2	2,724.1
2000	136.1	*	*	*	0.2	1.4	9.4	34.6	100.7	283.7	811.9	2,883.6
1999	139.9	*	*	*	0.2	1.4	9.6	35.2	103.9	297.9	847.2	2,973.2
Non-Hispanic white, male												
2006 5/	121.9	*	*	*	0.3	2.6	14.8	52.8	126.6	299.6	860.6	2,453.9
2005 5/	125.9	*	*	*	0.3	2.4	14.8	53.1	130.7	314.2	908.5	2,634.7
2004 5/	126.1	*	*	*	0.2	2.4	15.4	53.9	131.5	329.5	937.9	2,640.6
2003 5/	131.9	*	*	*	0.2	2.6	15.7	55.1	138.9	354.5	995.5	2,828.2
2002	134.5	*	*	*	0.2	2.3	16.2	56.4	143.5	371.2	1,033.6	2,908.8
2001	134.4	*	*	*	0.3	2.5	15.0	54.1	147.0	384.3	1,057.5	3,002.8
2000	137.7	*	*	*	0.2	2.2	15.1	55.7	152.7	406.1	1,086.5	3,211.3
1999	140.4	*	*	*	0.2	2.1	15.2	56.1	155.1	425.3	1,130.5	3,276.7
Non-Hispanic white, female												
2006 5/	108.1	*	*	*	*	0.5	4.1	14.0	41.9	133.8	481.7	1,977.0
2005 5/	114.7	*	*	*	*	0.8	4.2	14.4	43.7	143.6	518.4	2,133.9
2004 5/	116.9	*	*	*	*	0.7	4.5	14.2	45.6	150.1	538.3	2,211.1
2003 5/	124.3	*	*	*	*	0.8	4.6	14.4	47.0	160.9	572.4	2,403.6
2002	127.7	*	*	*	*	0.7	4.2	13.8	48.7	167.6	593.4	2,500.1
2001	131.1	*	*	*	*	0.7	4.1	13.9	50.5	175.6	614.6	2,608.1
2000	134.6	*	*	*	*	0.5	3.8	13.8	51.8	180.6	631.3	2,751.4
1999	139.3	*	*	*	*	0.7	4.0	14.6	55.7	191.3	661.7	2,852.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
Non-Hispanic black, both sexes												
2006 5/	76.4	*	*	*	*	2.5	14.1	58.7	152.7	334.8	757.9	1,808.1
2005 5/	79.1	*	*	*	*	2.3	14.5	60.7	164.4	341.6	801.4	1,901.2
2004 5/	80.3	*	*	*	0.4	3.0	14.6	60.3	163.1	363.9	822.2	2,031.7
2003 5/	84.7	*	*	*	0.4	2.7	14.6	61.9	173.6	393.8	878.9	2,204.8
2002	85.5	*	*	*	*	2.2	15.5	62.9	180.3	395.2	901.5	2,246.7
2001	88.2	*	*	*	0.5	2.9	15.6	62.9	186.6	419.6	934.8	2,357.3
2000	89.1	*	*	*	*	2.5	15.6	64.3	192.3	431.5	947.1	2,398.0
1999	91.6	*	*	*	0.4	2.4	17.2	66.9	197.9	453.5	958.2	2,471.9
Non-Hispanic black, male												
2006 5/	82.3	*	*	*	*	3.7	19.7	86.4	224.7	476.1	997.7	1,837.1
2005 5/	83.6	*	*	*	*	3.5	20.9	91.2	238.4	471.3	1,027.6	1,882.1
2004 5/	83.9	*	*	*	*	4.5	20.4	87.7	237.3	493.9	1,048.2	2,090.1
2003 5/	87.9	*	*	*	*	4.0	21.3	91.6	246.2	541.3	1,135.2	2,162.8
2002	88.0	*	*	*	*	3.0	22.2	90.8	264.4	536.0	1,119.3	2,254.6
2001	90.8	*	*	*	*	4.3	22.9	93.1	266.3	573.5	1,177.8	2,378.1
2000	90.4	*	*	*	*	3.7	22.1	93.6	274.7	566.6	1,180.1	2,463.5
1999	93.3	*	*	*	*	3.5	25.1	99.2	283.6	602.1	1,169.2	2,552.9
Non-Hispanic black, female												
2006 5/	71.0	*	*	*	*	1.4	9.2	34.9	94.8	233.9	621.8	1,796.3
2005 5/	75.0	*	*	*	*	1.1	8.9	34.6	104.8	248.9	673.2	1,908.8
2004 5/	76.9	*	*	*	*	1.6	9.4	36.8	103.6	271.3	694.6	2,008.6
2003 5/	81.7	*	*	*	*	1.5	8.7	36.5	115.4	288.8	734.1	2,221.1
2002	83.3	*	*	*	*	1.6	9.6	39.0	112.8	295.4	778.3	2,243.7
2001	86.0	*	*	*	*	1.6	9.2	36.9	122.4	310.9	798.4	2,349.6
2000	88.0	*	*	*	*	1.5	10.0	39.1	125.9	336.5	815.9	2,373.7
1999	90.1	*	*	*	*	1.3	10.2	39.4	129.2	348.6	840.0	2,441.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
All origins 3/, both sexes												
2006	20.4	*	*	*	0.0	0.5	3.3	13.4	31.3	55.1	114.2	345.5
2005	21.2	*	*	*	*	0.5	3.2	13.9	31.5	55.0	124.8	376.4
2004	21.7	*	*	*	*	0.5	3.3	13.6	32.0	58.7	129.0	400.4
2003	23.1	*	*	*	*	0.5	3.4	13.9	32.7	62.7	142.5	442.9
2002	23.6	*	*	*	*	0.5	3.4	13.4	33.1	64.7	146.2	474.5
2001	23.9	*	*	*	0.1	0.5	3.2	12.9	33.3	65.6	152.7	493.6
2000	24.7	*	*	*	*	0.4	3.0	13.2	34.0	68.3	158.1	534.6
1999	25.8	*	*	*	*	0.4	3.3	13.0	35.3	72.7	166.2	563.4
All origins 3/, male												
2006	23.2	*	*	*	*	0.8	5.1	21.7	48.8	79.1	146.8	365.3
2005	23.6	*	*	*	*	0.7	5.1	22.2	49.0	78.7	158.8	390.7
2004	23.9	*	*	*	*	0.8	5.2	22.0	49.0	83.6	164.9	414.0
2003	25.2	*	*	*	*	0.9	5.2	22.7	49.9	88.9	181.1	462.1
2002	25.4	*	*	*	*	0.7	5.4	21.5	51.2	91.8	185.3	491.8
2001	25.3	*	*	*	*	0.7	5.0	21.0	51.4	92.4	193.9	502.0
2000	26.0	*	*	*	*	0.6	4.9	21.5	52.6	97.2	199.3	550.8
1999	26.8	*	*	*	*	0.6	5.3	20.9	53.7	103.0	207.0	578.6
All origins 3/, female												
2006	17.7	*	*	*	*	0.2	1.5	5.5	15.1	34.8	92.0	336.1
2005	18.8	*	*	*	*	0.2	1.4	5.7	15.3	35.0	101.6	369.9
2004	19.5	*	*	*	*	0.2	1.4	5.5	16.1	37.7	104.8	394.3
2003	21.1	*	*	*	*	0.2	1.5	5.5	16.7	40.8	116.7	434.4
2002	21.9	*	*	*	*	0.2	1.3	5.5	16.2	42.1	120.2	467.1
2001	22.5	*	*	*	*	0.2	1.4	5.1	16.7	43.3	125.5	490.1
2000	23.5	*	*	*	*	0.1	1.2	5.2	16.8	44.6	131.3	528.0
1999	24.8	*	*	*	*	0.2	1.3	5.3	18.3	47.9	139.8	557.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
Hispanic, both sexes												
2006	7.7	*	*	*	*	*	1.7	9.1	28.2	53.9	98.8	260.2
2005	8.2	*	*	*	*	*	1.7	10.9	29.1	55.2	113.8	274.7
2004	7.9	*	*	*	*	*	1.7	9.7	29.3	55.8	118.9	279.7
2003	8.3	*	*	*	*	0.3	1.6	9.8	29.4	57.3	124.6	357.1
2002	8.2	*	*	*	*	0.4	1.9	8.9	29.7	61.1	124.1	358.6
2001	8.5	*	*	*	*	0.4	1.5	10.1	29.7	65.9	132.5	417.0
2000	8.4	*	*	*	*	*	1.6	9.5	30.5	65.7	128.2	452.5
1999	9.0	*	*	*	*	*	1.5	10.4	33.0	72.1	147.5	469.4
Hispanic, male												
2006	9.5	*	*	*	*	*	2.7	15.3	46.4	81.1	126.8	277.4
2005	10.2	*	*	*	*	*	2.8	17.9	47.8	82.8	156.0	308.8
2004	9.8	*	*	*	*	*	2.7	16.6	47.1	83.3	169.5	297.0
2003	10.2	*	*	*	*	0.5	2.6	17.1	48.7	88.6	169.5	397.1
2002	9.7	*	*	*	*	0.6	2.9	14.8	49.3	90.8	153.9	415.0
2001	10.1	*	*	*	*	0.6	2.2	17.1	47.9	100.2	181.1	440.0
2000	9.9	*	*	*	*	*	2.7	16.6	50.5	92.4	180.9	493.8
1999	10.6	*	*	*	*	*	2.5	17.8	50.0	115.8	194.0	485.3
Hispanic, female												
2006	5.8	*	*	*	*	*	*	2.7	11.5	31.9	79.2	251.0
2005	6.0	*	*	*	*	*	*	3.7	11.9	32.8	84.4	256.8
2004	5.9	*	*	*	*	*	*	2.8	13.2	33.8	84.0	270.8
2003	6.2	*	*	*	*	*	*	2.5	12.0	32.4	93.7	336.7
2002	6.5	*	*	*	*	*	*	3.1	12.0	37.6	103.7	330.7
2001	6.8	*	*	*	*	*	*	3.2	13.5	38.7	99.7	405.8
2000	6.8	*	*	*	*	*	*	2.6	12.8	44.6	92.8	432.3
1999	7.4	*	*	*	*	*	*	3.2	18.0	37.9	116.6	461.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
Non-Hispanic 4/, both sexes												
2006	22.5	*	*	*	*	0.6	3.6	13.9	31.4	54.8	114.7	349.3
2005	23.3	*	*	*	*	0.5	3.5	14.1	31.5	54.6	124.8	380.6
2004	23.8	*	*	*	*	0.5	3.6	14.0	32.0	58.3	128.9	404.7
2003	25.3	*	*	*	*	0.6	3.6	14.2	32.6	62.6	142.6	445.3
2002	25.9	*	*	*	*	0.5	3.6	13.7	32.9	64.3	146.4	477.8
2001	26.0	*	*	*	*	0.5	3.5	13.0	33.3	65.0	152.6	494.8
2000	26.9	*	*	*	*	0.4	3.2	13.4	33.8	67.9	158.3	535.9
1999	27.9	*	*	*	*	0.4	3.5	13.1	35.0	72.0	165.7	564.9
Non-Hispanic 4/, male												
2006	25.5	*	*	*	*	1.0	5.6	22.3	48.6	78.2	147.2	369.6
2005	25.8	*	*	*	*	0.8	5.5	22.6	48.7	77.7	158.1	394.0
2004	26.2	*	*	*	*	0.9	5.6	22.5	48.9	82.8	163.7	418.3
2003	27.5	*	*	*	*	1.0	5.6	23.1	49.3	88.2	180.4	463.3
2002	27.8	*	*	*	*	0.7	5.8	22.0	50.7	91.0	185.6	493.3
2001	27.5	*	*	*	*	0.7	5.4	21.2	51.1	91.0	193.2	503.0
2000	28.2	*	*	*	*	0.7	5.1	21.7	52.1	96.4	198.5	551.0
1999	28.9	*	*	*	*	0.6	5.6	21.0	53.3	101.1	205.6	580.2
Non-Hispanic 4/, female												
2006	19.6	*	*	*	*	0.2	1.6	5.8	15.4	34.8	92.5	339.9
2005	20.8	*	*	*	*	0.2	1.5	6.0	15.5	35.0	102.2	374.5
2004	21.5	*	*	*	*	0.2	1.6	5.8	16.3	37.7	105.5	398.6
2003	23.3	*	*	*	*	0.2	1.6	5.7	16.9	41.2	117.4	437.4
2002	24.0	*	*	*	*	0.2	1.4	5.7	16.5	42.0	120.5	471.1
2001	24.6	*	*	*	*	0.3	1.5	5.2	16.8	43.3	126.0	491.4
2000	25.6	*	*	*	*	0.2	1.3	5.4	16.9	44.4	132.2	529.8
1999	27.0	*	*	*	*	0.2	1.4	5.4	18.2	48.2	139.8	558.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
Non-Hispanic white, both sexes												
2006 5/	23.2	*	*	*	*	0.5	3.3	12.5	28.0	49.1	108.5	346.8
2005 5/	24.0	*	*	*	*	0.4	3.3	12.7	27.5	49.1	117.9	377.8
2004 5/	24.6	*	*	*	*	0.5	3.5	12.6	28.4	52.4	121.7	399.9
2003 5/	26.2	*	*	*	*	0.6	3.4	12.8	28.6	55.9	135.1	441.5
2002	26.8	*	*	*	*	0.4	3.3	12.4	28.9	57.2	138.2	473.5
2001	26.8	*	*	*	*	0.4	3.2	11.6	28.4	57.1	144.2	490.7
2000	27.6	*	*	*	*	0.4	2.9	11.9	28.9	60.2	149.0	530.2
1999	28.6	*	*	*	*	0.4	3.1	11.3	29.8	63.6	156.2	558.8
Non-Hispanic white, male												
2006 5/	25.9	*	*	*	*	0.9	5.3	20.4	43.5	69.8	139.0	366.7
2005 5/	26.3	*	*	*	*	0.7	5.3	20.7	43.0	69.8	148.8	396.1
2004 5/	26.7	*	*	*	*	0.7	5.5	20.8	43.4	75.2	154.3	412.6
2003 5/	28.0	*	*	*	*	0.9	5.3	21.0	43.9	78.5	170.6	459.5
2002	28.2	*	*	*	*	0.7	5.4	20.3	44.8	81.3	174.4	489.2
2001	27.8	*	*	*	*	0.6	5.0	19.1	44.3	80.4	182.9	499.4
2000	28.4	*	*	*	*	0.6	4.8	19.6	45.2	86.3	186.1	544.7
1999	29.0	*	*	*	*	0.6	5.2	18.4	46.0	90.0	193.5	572.8
Non-Hispanic white, female												
2006 5/	20.6	*	*	*	*	*	1.4	4.7	13.3	31.1	87.3	337.5
2005 5/	21.9	*	*	*	*	0.2	1.2	4.8	12.8	31.2	96.6	369.4
2004 5/	22.6	*	*	*	*	0.2	1.4	4.5	14.1	32.9	99.4	394.2
2003 5/	24.6	*	*	*	*	0.2	1.5	4.6	14.2	36.5	111.0	433.5
2002	25.3	*	*	*	*	0.2	1.1	4.6	13.8	36.6	113.9	466.8
2001	25.8	*	*	*	*	0.2	1.3	4.1	13.3	37.3	118.6	487.0
2000	26.9	*	*	*	*	*	1.1	4.3	13.4	38.2	124.6	524.4
1999	28.2	*	*	*	*	0.2	1.1	4.2	14.6	41.6	131.8	553.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
Non-Hispanic black, both sexes												
2006 5/	24.4	*	*	*	*	1.1	5.8	25.5	63.7	111.6	201.2	433.7
2005 5/	25.3	*	*	*	*	0.8	5.4	26.1	68.6	110.2	218.2	475.2
2004 5/	25.7	*	*	*	*	0.9	4.9	26.1	67.3	120.8	229.1	509.2
2003 5/	26.7	*	*	*	*	0.9	5.1	26.1	70.1	129.6	244.1	546.7
2002	27.7	*	*	*	*	0.8	6.0	25.7	71.9	137.2	261.1	580.4
2001	28.5	*	*	*	*	0.7	5.8	25.7	78.1	144.8	266.7	595.9
2000	29.3	*	*	*	*	0.7	5.3	27.3	78.9	147.2	280.2	652.2
1999	30.7	*	*	*	*	0.8	6.5	28.1	82.0	158.7	291.0	670.3
Non-Hispanic black, male												
2006 5/	29.5	*	*	*	*	1.5	8.4	38.9	100.8	170.9	275.8	471.0
2005 5/	29.7	*	*	*	*	1.3	7.9	39.8	106.1	168.1	296.8	452.0
2004 5/	29.7	*	*	*	*	1.6	7.1	38.5	105.6	173.1	308.9	540.6
2003 5/	31.2	*	*	*	*	1.6	7.8	41.1	106.0	191.8	334.2	588.1
2002	31.9	*	*	*	*	1.1	8.9	38.8	111.8	198.7	358.4	603.9
2001	32.3	*	*	*	*	1.0	8.8	39.9	118.1	209.4	352.3	609.8
2000	33.2	*	*	*	*	1.1	7.9	42.3	120.8	209.3	371.5	711.6
1999	34.5	*	*	*	*	1.1	9.8	43.6	123.7	228.2	378.4	714.3
Non-Hispanic black, female												
2006 5/	19.8	*	*	*	*	*	3.5	14.0	33.8	69.3	158.9	418.5
2005 5/	21.3	*	*	*	*	*	3.2	14.3	38.4	68.9	173.7	484.5
2004 5/	22.1	*	*	*	*	*	3.1	15.4	36.5	83.4	184.0	496.7
2003 5/	22.7	*	*	*	*	*	2.7	13.3	41.2	85.3	193.3	530.7
2002	24.0	*	*	*	*	*	3.4	14.4	39.8	93.6	206.1	571.5
2001	25.0	*	*	*	*	*	3.2	13.6	45.9	99.3	218.7	590.7
2000	25.8	*	*	*	*	*	2.9	14.4	45.1	103.4	228.9	630.2
1999	27.1	*	*	*	*	*	3.6	14.7	48.5	109.7	242.1	653.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
All origins 3/, both sexes												
2006	73.1	*	*	*	0.1	1.0	5.7	21.0	56.5	162.8	518.6	1,712.0
2005	77.0	*	*	*	0.2	1.1	6.0	21.2	59.8	174.9	549.7	1,837.8
2004	77.8	*	*	*	0.1	1.1	6.2	21.5	60.5	181.7	568.1	1,879.3
2003	82.2	*	*	*	0.2	1.1	6.5	22.0	64.9	195.7	599.0	2,028.5
2002	84.4	*	*	*	0.1	1.0	6.5	23.1	68.0	204.7	622.6	2,088.0
2001	86.4	*	*	*	0.2	1.1	6.2	22.7	70.5	215.3	640.5	2,177.7
2000	88.6	*	*	*	0.1	1.0	6.5	23.2	74.1	224.1	655.5	2,287.5
1999	91.4	*	*	*	0.2	1.0	6.5	24.5	76.3	235.2	682.3	2,353.6
All origins 3/, male												
2006	76.1	*	*	*	0.2	1.6	8.7	32.1	83.7	228.7	701.5	1,979.5
2005	79.3	*	*	*	0.2	1.6	9.0	32.5	88.0	243.3	738.3	2,120.9
2004	79.4	*	*	*	0.2	1.7	9.3	32.8	88.7	252.9	762.0	2,129.6
2003	83.3	*	*	*	0.2	1.6	9.7	33.7	95.5	274.6	804.7	2,260.1
2002	85.5	*	*	*	0.2	1.5	10.0	35.9	100.3	287.5	835.7	2,325.6
2001	86.5	*	*	*	0.2	1.8	9.5	34.9	103.5	301.9	856.9	2,411.8
2000	88.8	*	*	*	0.2	1.5	9.9	36.2	109.0	316.3	880.1	2,561.9
1999	91.2	*	*	*	0.2	1.5	10.0	38.1	111.0	331.9	913.6	2,614.4
All origins 3/, female												
2006	70.3	*	*	*	*	0.4	2.7	10.2	31.2	107.1	393.5	1,586.8
2005	74.8	*	*	*	*	0.5	2.9	10.3	33.7	117.2	421.7	1,707.7
2004	76.2	*	*	*	*	0.5	3.2	10.6	34.3	121.9	437.6	1,766.7
2003	81.1	*	*	*	0.1	0.6	3.2	10.8	36.4	129.7	461.6	1,926.2
2002	83.3	*	*	*	*	0.5	3.1	10.7	38.0	135.9	481.1	1,984.9
2001	86.2	*	*	*	0.1	0.5	3.0	11.0	40.0	143.6	498.2	2,079.8
2000	88.4	*	*	*	*	0.5	3.0	10.7	42.0	148.2	509.0	2,175.8
1999	91.6	*	*	*	0.1	0.5	3.0	11.4	44.3	156.1	532.1	2,248.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Hispanic, both sexes												
2006	23.6	*	*	*	*	0.6	2.7	13.5	40.3	130.6	437.2	1,335.0
2005	25.0	*	*	*	*	0.6	3.4	12.1	42.8	150.8	475.0	1,455.8
2004	24.3	*	*	*	*	0.6	3.2	12.7	43.6	149.7	470.7	1,516.0
2003	25.2	*	*	*	*	0.6	3.0	12.2	47.8	159.1	504.9	1,655.7
2002	26.0	*	*	*	*	0.4	2.7	14.6	50.4	173.4	519.9	1,771.1
2001	27.2	*	*	*	*	0.6	2.7	13.7	50.4	183.6	569.1	2,009.6
2000	27.2	*	*	*	*	0.5	3.0	14.3	57.5	184.0	575.0	2,056.3
1999	28.5	*	*	*	*	0.6	3.0	16.4	60.5	196.7	594.3	2,171.4
Hispanic, male												
2006	23.8	*	*	*	*	0.9	4.3	20.4	56.8	173.4	553.9	1,427.7
2005	25.0	*	*	*	*	0.9	5.5	18.1	60.5	204.4	598.3	1,503.1
2004	24.3	*	*	*	*	0.9	4.8	18.3	62.2	207.8	600.2	1,556.5
2003	24.4	*	*	*	*	0.9	4.3	18.0	66.6	217.6	634.7	1,568.6
2002	25.9	*	*	*	*	0.6	4.3	21.8	73.0	238.3	654.9	1,847.2
2001	26.7	*	*	*	*	0.9	4.3	20.4	73.6	250.0	720.4	2,039.8
2000	27.0	*	*	*	*	0.7	4.7	22.6	83.6	259.1	719.8	2,084.0
1999	28.1	*	*	*	*	1.0	5.1	26.5	88.5	266.0	738.8	2,221.4
Hispanic, female												
2006	23.3	*	*	*	*	*	1.0	6.5	25.1	95.9	355.4	1,285.7
2005	25.0	*	*	*	*	*	1.0	6.0	26.6	107.4	389.1	1,431.1
2004	24.2	*	*	*	*	*	1.3	7.0	26.7	103.2	381.3	1,495.2
2003	26.1	*	*	*	*	*	1.6	6.4	30.7	112.5	415.8	1,700.0
2002	26.1	*	*	*	*	*	0.9	7.4	30.2	121.8	427.5	1,733.3
2001	27.7	*	*	*	*	*	1.0	7.1	29.8	131.2	466.9	1,994.8
2000	27.5	*	*	*	*	*	1.2	6.2	34.4	124.9	477.7	2,042.7
1999	28.9	*	*	*	*	*	0.8	6.5	35.8	142.3	498.2	2,147.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Non-Hispanic 4/, both sexes												
2006	81.6	*	*	*	0.1	1.1	6.2	21.8	57.7	164.9	522.9	1,728.7
2005	85.6	*	*	*	0.2	1.2	6.4	22.2	61.1	176.4	553.1	1,853.4
2004	86.3	*	*	*	0.2	1.2	6.7	22.4	61.6	183.5	572.4	1,892.8
2003	91.0	*	*	*	0.2	1.2	7.0	22.9	65.9	197.6	602.3	2,039.6
2002	93.1	*	*	*	0.1	1.1	7.1	23.8	69.0	206.0	625.4	2,093.4
2001	94.9	*	*	*	0.2	1.2	6.7	23.5	71.8	216.7	641.6	2,177.4
2000	97.0	*	*	*	0.1	1.1	6.9	23.9	75.0	225.6	656.7	2,288.1
1999	99.7	*	*	*	0.2	1.1	6.9	25.1	77.1	236.5	683.4	2,351.4
Non-Hispanic 4/, male												
2006	85.6	*	*	*	0.2	1.8	9.5	33.4	85.8	232.3	709.8	2,007.1
2005	88.8	*	*	*	0.3	1.8	9.7	34.0	90.1	245.6	745.1	2,150.8
2004	88.7	*	*	*	0.2	1.8	10.0	34.2	90.4	255.3	769.4	2,154.8
2003	92.9	*	*	*	0.2	1.8	10.6	35.2	97.4	277.5	811.5	2,287.4
2002	94.9	*	*	*	0.2	1.6	10.8	37.2	102.0	289.3	841.7	2,338.9
2001	95.6	*	*	*	0.2	2.0	10.3	36.1	105.3	304.1	860.3	2,420.7
2000	97.8	*	*	*	0.2	1.7	10.5	37.2	110.2	318.2	884.2	2,572.9
1999	100.0	*	*	*	0.2	1.6	10.6	38.9	112.1	334.1	917.9	2,619.9
Non-Hispanic 4/, female												
2006	77.9	*	*	*	*	0.4	3.0	10.6	31.6	107.7	395.3	1,599.3
2005	82.6	*	*	*	*	0.6	3.2	10.7	34.2	117.8	422.8	1,717.8
2004	84.1	*	*	*	*	0.6	3.4	11.0	34.9	123.0	439.9	1,775.7
2003	89.2	*	*	*	0.1	0.7	3.4	11.2	36.7	130.6	462.6	1,930.7
2002	91.4	*	*	*	*	0.6	3.4	11.0	38.5	136.4	482.1	1,987.5
2001	94.2	*	*	*	*	0.5	3.2	11.3	40.7	143.9	498.0	2,076.2
2000	96.3	*	*	*	*	0.5	3.3	11.1	42.4	149.1	508.5	2,172.9
1999	99.4	*	*	*	*	0.6	3.3	11.8	44.8	156.4	531.2	2,243.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Non-Hispanic white, both sexes												
2006 5/	91.6	*	*	*	0.1	1.1	6.1	20.7	55.2	161.7	528.4	1,781.9
2005 5/	96.2	*	*	*	0.1	1.2	6.2	20.9	58.6	173.6	559.5	1,913.3
2004 5/	96.8	*	*	*	0.1	1.1	6.5	21.3	59.1	180.6	578.5	1,944.1
2003 5/	101.8	*	*	*	0.1	1.1	6.7	21.8	63.1	194.3	607.9	2,091.9
2002	104.3	*	*	*	0.1	1.1	6.9	22.5	65.9	204.1	631.9	2,149.8
2001	106.0	*	*	*	0.1	1.1	6.4	22.2	69.0	214.3	647.0	2,233.4
2000	108.4	*	*	*	0.1	1.0	6.5	22.7	71.9	223.5	662.9	2,353.3
1999	111.3	*	*	*	0.1	1.0	6.5	23.9	74.1	234.2	690.9	2,414.4
Non-Hispanic white, male												
2006 5/	95.9	*	*	*	0.2	1.7	9.5	32.3	83.2	229.7	721.6	2,087.2
2005 5/	99.6	*	*	*	0.2	1.7	9.4	32.5	87.7	244.4	759.7	2,238.6
2004 5/	99.3	*	*	*	0.2	1.7	9.9	33.1	88.1	254.3	783.6	2,228.0
2003 5/	103.9	*	*	*	0.2	1.7	10.4	34.1	94.9	276.0	824.8	2,368.6
2002	106.2	*	*	*	0.2	1.6	10.7	36.1	98.7	289.9	859.2	2,419.7
2001	106.6	*	*	*	0.2	1.8	10.0	34.9	102.7	303.9	874.6	2,503.4
2000	109.2	*	*	*	0.2	1.6	10.3	36.1	107.5	319.8	900.4	2,666.6
1999	111.4	*	*	*	0.2	1.5	10.1	37.7	109.1	335.3	936.9	2,703.9
Non-Hispanic white, female												
2006 5/	87.5	*	*	*	*	0.4	2.7	9.3	28.6	102.8	394.5	1,639.5
2005 5/	92.9	*	*	*	*	0.6	2.9	9.5	30.9	112.4	421.8	1,764.6
2004 5/	94.3	*	*	*	*	0.5	3.1	9.7	31.5	117.3	438.8	1,816.9
2003 5/	99.8	*	*	*	*	0.6	3.1	9.7	32.8	124.3	461.4	1,970.1
2002	102.4	*	*	*	*	0.5	3.1	9.2	34.9	131.0	479.6	2,033.3
2001	105.3	*	*	*	*	0.5	2.9	9.8	37.2	138.2	496.0	2,121.0
2000	107.7	*	*	*	*	0.4	2.7	9.5	38.3	142.4	506.8	2,227.0
1999	111.1	*	*	*	*	0.5	2.8	10.4	41.1	149.7	529.9	2,299.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Non-Hispanic black, both sexes												
2006 5/	51.9	*	*	*	*	1.4	8.4	33.2	89.0	223.2	556.6	1,374.4
2005 5/	53.8	*	*	*	*	1.5	9.1	34.6	95.8	231.4	583.2	1,426.0
2004 5/	54.6	*	*	*	0.4	2.1	9.6	34.2	95.9	243.2	593.1	1,522.6
2003 5/	57.9	*	*	*	*	1.8	9.5	35.8	103.6	264.2	634.8	1,658.1
2002	57.8	*	*	*	*	1.5	9.6	37.2	108.4	258.0	640.3	1,666.3
2001	59.7	*	*	*	0.4	2.1	9.8	37.1	108.4	274.7	668.1	1,761.4
2000	59.8	*	*	*	*	1.8	10.4	37.0	113.4	284.4	666.8	1,745.8
1999	61.0	*	*	*	*	1.6	10.7	38.9	115.9	294.7	667.1	1,801.6
Non-Hispanic black, male												
2006 5/	52.7	*	*	*	*	2.2	11.3	47.5	123.9	305.2	721.9	1,366.1
2005 5/	53.9	*	*	*	*	2.2	13.0	51.4	132.3	303.3	730.8	1,430.2
2004 5/	54.2	*	*	*	*	3.0	13.4	49.2	131.6	320.8	739.3	1,549.6
2003 5/	56.7	*	*	*	*	2.4	13.4	50.5	140.2	349.5	801.1	1,574.7
2002	56.1	*	*	*	*	1.9	13.3	52.0	152.5	337.3	760.8	1,650.8
2001	58.4	*	*	*	*	3.3	14.1	53.2	148.2	364.1	825.5	1,768.3
2000	57.3	*	*	*	*	2.6	14.1	51.3	153.9	357.2	808.7	1,751.9
1999	58.8	*	*	*	*	2.4	15.3	55.5	160.0	374.0	790.8	1,838.6
Non-Hispanic black, female												
2006 5/	51.2	*	*	*	*	0.7	5.7	21.0	61.0	164.6	462.9	1,377.8
2005 5/	53.7	*	*	*	*	0.8	5.7	20.3	66.4	180.0	499.6	1,424.3
2004 5/	54.9	*	*	*	*	1.2	6.3	21.3	67.1	187.8	510.6	1,511.9
2003 5/	59.1	*	*	*	*	1.2	6.1	23.2	74.2	203.5	540.8	1,690.4
2002	59.3	*	*	*	*	1.1	6.2	24.5	73.0	201.8	572.2	1,672.2
2001	60.9	*	*	*	*	1.0	6.0	23.3	76.5	211.7	579.7	1,758.9
2000	62.1	*	*	*	*	1.1	7.1	24.7	80.7	233.1	587.0	1,743.5
1999	63.0	*	*	*	*	0.8	6.6	24.6	80.7	238.9	597.9	1,787.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
All origins 3/, both sexes												
2006	56.9	7.7	0.9	0.6	2.1	4.7	9.5	21.2	44.4	112.3	360.7	1,342.8
2005	57.6	8.3	0.9	0.6	2.2	4.6	9.8	21.5	45.7	116.8	366.7	1,377.4
2004	56.6	10.0	1.1	0.5	2.1	4.3	9.7	21.5	45.2	118.6	368.9	1,373.9
2003	58.5	10.4	1.1	0.6	2.2	4.5	10.3	21.4	48.0	126.8	382.8	1,427.2
2002	58.8	11.7	1.0	0.6	2.1	4.5	10.3	21.7	48.7	129.6	390.9	1,443.5
2001	58.6	11.1	1.4	0.6	2.0	4.4	9.9	21.6	49.5	131.4	389.5	1,475.5
2000	58.8	12.0	1.1	0.6	2.2	4.4	9.9	21.5	50.9	134.2	391.3	1,509.4
1999	59.4	12.9	1.2	0.6	2.3	4.4	10.3	21.6	52.9	138.8	398.5	1,512.9
All origins 3/, male												
2006	51.0	8.0	1.0	0.6	2.5	5.9	11.6	27.2	56.2	136.2	411.1	1,338.1
2005	51.4	8.9	0.9	0.6	2.8	5.8	12.3	27.6	57.5	140.5	415.8	1,377.8
2004	50.6	10.5	1.1	0.5	2.6	5.3	12.0	27.7	57.7	142.8	420.8	1,377.7
2003	52.3	11.4	1.0	0.7	2.6	5.3	12.7	27.2	60.7	154.2	440.0	1,431.3
2002	52.4	12.1	1.0	0.6	2.7	5.6	12.8	28.2	60.8	156.5	448.2	1,448.6
2001	51.8	10.8	1.5	0.6	2.4	5.2	12.4	27.6	61.7	158.7	452.0	1,473.4
2000	52.2	12.2	1.2	0.7	2.6	5.2	12.2	27.8	64.7	164.0	451.0	1,549.1
1999	53.1	13.1	1.3	0.7	2.7	5.3	12.9	27.7	67.5	172.7	459.7	1,548.9
All origins 3/, female												
2006	62.6	7.5	0.9	0.5	1.5	3.5	7.3	15.3	33.5	92.1	326.3	1,345.0
2005	63.5	7.7	0.8	0.6	1.5	3.4	7.3	15.5	34.6	96.7	333.3	1,377.3
2004	62.4	9.4	1.1	0.6	1.5	3.3	7.4	15.5	33.6	98.4	334.0	1,372.1
2003	64.5	9.3	1.2	0.5	1.7	3.7	7.9	15.7	36.3	103.8	344.5	1,425.4
2002	65.0	11.4	1.0	0.5	1.4	3.4	7.8	15.5	37.5	107.2	352.8	1,441.3
2001	65.2	11.5	1.3	0.6	1.5	3.6	7.5	15.9	38.3	108.7	348.4	1,476.4
2000	65.1	11.8	0.9	0.5	1.8	3.5	7.5	15.3	38.2	109.6	352.3	1,493.2
1999	65.5	12.7	1.0	0.6	1.8	3.6	7.7	15.6	39.4	111.1	358.7	1,498.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
Hispanic, both sexes												
2006	13.7	6.3	0.8	0.5	1.4	2.8	4.7	11.9	23.9	67.5	197.0	663.2
2005	13.9	7.5	0.6	0.4	1.5	2.7	5.1	10.9	24.7	70.6	210.8	703.3
2004	13.2	8.5	1.0	0.4	1.5	2.3	5.3	11.4	24.7	70.3	208.5	659.3
2003	14.1	8.3	1.0	0.3	1.4	2.8	5.3	11.3	29.7	77.3	221.2	749.9
2002	13.3	9.7	0.7	0.4	1.5	2.6	5.1	11.4	25.7	72.6	219.8	740.0
2001	13.3	9.2	1.0	0.5	1.3	2.6	4.9	10.9	27.9	75.4	217.5	798.5
2000	13.6	10.2	1.0	0.5	1.7	2.8	5.5	11.6	28.8	75.9	227.0	829.4
1999	13.8	12.2	1.2	0.5	1.6	2.8	6.3	11.6	30.6	78.7	229.6	827.9
Hispanic, male												
2006	13.5	6.9	*	0.6	1.9	3.9	6.1	15.1	30.6	82.9	231.4	658.6
2005	13.7	8.0	*	*	1.9	3.7	6.9	14.9	31.8	87.0	247.5	697.4
2004	13.0	8.4	1.3	*	2.2	3.3	7.1	15.5	31.6	83.6	241.8	641.6
2003	13.6	9.1	*	*	1.8	3.4	7.0	14.7	37.0	95.0	253.5	752.7
2002	12.9	9.1	*	*	2.1	3.4	6.4	14.3	31.7	92.5	256.9	758.0
2001	12.6	9.6	*	*	1.6	3.2	6.5	15.0	34.1	91.1	248.6	775.2
2000	13.2	9.9	1.3	0.7	2.1	3.5	6.7	15.1	37.2	94.1	250.8	890.8
1999	13.7	10.3	1.4	0.6	1.8	3.8	8.2	15.4	38.8	102.4	267.5	888.6
Hispanic, female												
2006	13.9	5.7	*	*	0.8	1.5	3.2	8.5	17.8	54.9	172.8	665.6
2005	14.1	7.0	*	*	1.2	1.5	3.1	6.8	18.3	57.3	185.1	706.4
2004	13.5	8.6	*	*	0.8	1.1	3.3	7.3	18.5	59.6	185.5	668.4
2003	14.5	7.6	*	*	0.9	2.2	3.3	8.0	23.2	63.2	199.1	748.4
2002	13.7	10.3	*	*	0.8	1.6	3.7	8.5	20.3	56.7	194.5	731.1
2001	13.9	8.7	*	*	0.9	1.9	3.2	6.9	22.5	63.0	196.4	809.8
2000	14.0	10.6	*	*	1.1	1.9	4.3	8.1	21.4	61.6	211.0	799.3
1999	13.9	14.1	*	*	1.3	1.6	4.2	8.0	23.4	60.1	204.3	798.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
Non-Hispanic 4/, both sexes												
2006	64.3	8.2	1.0	0.6	2.2	5.2	10.3	22.2	46.1	115.5	370.4	1,375.6
2005	64.8	8.6	1.0	0.6	2.3	5.0	10.6	22.6	47.3	120.1	375.4	1,408.5
2004	63.6	10.3	1.1	0.6	2.1	4.7	10.4	22.5	46.7	122.0	377.3	1,404.5
2003	65.4	10.9	1.1	0.6	2.3	4.9	11.0	22.3	49.3	130.0	390.5	1,453.7
2002	65.7	12.2	1.1	0.6	2.2	4.9	11.0	22.7	50.4	133.2	398.4	1,469.2
2001	65.2	11.6	1.5	0.6	2.1	4.7	10.6	22.6	51.1	134.7	396.5	1,497.8
2000	65.1	12.4	1.1	0.6	2.3	4.7	10.4	22.3	52.4	137.5	397.5	1,530.9
1999	65.6	13.1	1.1	0.6	2.4	4.7	10.8	22.4	54.3	142.0	404.4	1,533.9
Non-Hispanic 4/, male												
2006	57.8	8.3	1.0	0.6	2.7	6.5	12.6	28.6	58.3	139.8	421.7	1,374.1
2005	58.0	9.2	1.0	0.6	3.0	6.3	13.2	29.0	59.5	144.2	425.3	1,412.5
2004	57.0	11.0	1.0	0.6	2.7	5.8	12.8	28.9	59.6	146.8	430.2	1,412.5
2003	58.6	12.0	1.0	0.7	2.8	5.8	13.5	28.4	62.2	158.0	448.9	1,460.5
2002	58.7	12.8	1.1	0.7	2.8	6.2	13.7	29.5	62.9	160.4	456.6	1,477.3
2001	57.8	11.0	1.6	0.6	2.6	5.6	13.2	28.7	63.6	162.5	460.6	1,499.0
2000	58.0	12.7	1.2	0.7	2.7	5.5	12.9	28.9	66.4	167.8	458.9	1,573.6
1999	58.6	13.8	1.3	0.7	2.9	5.5	13.5	28.7	69.3	176.2	466.4	1,571.2
Non-Hispanic 4/, female												
2006	70.4	8.0	0.9	0.5	1.7	3.9	8.0	16.1	34.8	94.9	335.4	1,376.4
2005	71.2	7.9	1.0	0.6	1.6	3.8	8.0	16.4	36.0	99.6	341.5	1,406.6
2004	69.9	9.6	1.2	0.6	1.6	3.7	8.0	16.4	34.8	101.1	341.7	1,400.9
2003	71.8	9.7	1.2	0.5	1.9	4.0	8.5	16.4	37.3	106.5	351.5	1,450.7
2002	72.3	11.6	1.0	0.5	1.5	3.7	8.4	16.1	38.7	110.4	359.7	1,465.7
2001	72.3	12.2	1.3	0.6	1.6	3.9	8.0	16.7	39.4	111.6	354.4	1,497.3
2000	71.8	12.0	1.0	0.5	1.9	3.8	7.9	15.9	39.4	112.5	357.5	1,513.7
1999	72.2	12.3	1.0	0.6	1.9	3.9	8.1	16.2	40.5	113.9	364.2	1,519.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
Non-Hispanic white, both sexes												
2006 5/	70.4	5.8	0.8	0.5	1.9	4.1	8.6	18.9	41.4	109.8	372.0	1,422.6
2005 5/	70.6	6.7	0.7	0.5	1.9	3.9	8.8	19.1	42.0	114.0	376.1	1,453.0
2004 5/	69.3	8.7	0.8	0.5	1.8	3.8	8.6	19.0	41.9	116.5	377.3	1,442.1
2003 5/	71.0	8.1	0.8	0.5	1.9	3.8	9.1	18.8	44.1	123.1	391.4	1,490.1
2002	71.3	10.0	0.9	0.5	1.8	3.8	9.3	19.0	45.3	127.6	398.1	1,503.1
2001	70.6	9.1	1.1	0.5	1.7	3.7	8.5	19.0	45.8	128.7	394.8	1,533.2
2000	70.3	10.5	0.9	0.5	1.7	3.7	8.4	18.7	47.4	131.2	395.8	1,564.2
1999	70.4	10.5	0.9	0.5	2.0	3.7	8.7	18.7	48.3	135.2	401.1	1,565.9
Non-Hispanic white, male												
2006 5/	62.1	5.7	0.8	0.6	2.3	5.1	10.5	24.4	52.5	132.9	424.0	1,420.8
2005 5/	62.2	6.9	0.8	0.5	2.5	4.9	11.2	24.9	53.1	136.5	427.1	1,462.5
2004 5/	60.9	9.3	0.7	0.5	2.3	4.6	10.8	24.7	53.6	140.9	428.9	1,445.4
2003 5/	62.6	8.7	0.7	0.6	2.3	4.6	11.4	24.3	56.0	150.6	450.4	1,499.3
2002	62.7	10.7	1.1	0.6	2.3	4.7	11.8	25.2	57.1	154.1	457.1	1,512.9
2001	61.4	9.4	1.2	0.6	2.0	4.4	10.9	24.8	57.5	155.8	457.4	1,530.7
2000	61.6	10.9	1.1	0.6	2.0	4.4	10.7	24.6	60.7	161.0	457.6	1,606.4
1999	61.7	11.6	1.1	0.5	2.3	4.4	11.2	24.5	62.1	168.5	462.9	1,595.5
Non-Hispanic white, female												
2006 5/	78.4	5.8	0.8	0.5	1.4	3.0	6.7	13.5	30.9	89.9	335.9	1,423.5
2005 5/	78.8	6.4	0.6	0.4	1.3	2.8	6.4	13.3	31.4	94.6	340.9	1,448.6
2004 5/	77.3	8.2	0.9	0.4	1.3	2.9	6.4	13.3	30.7	95.6	342.1	1,440.6
2003 5/	79.1	7.5	1.0	0.4	1.6	3.0	6.8	13.5	32.7	99.6	351.5	1,486.1
2002	79.6	9.3	0.8	0.5	1.2	2.8	6.8	13.0	34.1	105.0	358.5	1,498.8
2001	79.4	8.8	1.0	0.5	1.3	3.1	6.1	13.4	34.8	105.7	353.3	1,534.2
2000	78.7	10.1	0.8	0.4	1.4	2.9	6.2	12.9	34.8	106.0	355.3	1,547.2
1999	78.8	9.4	0.8	0.5	1.7	3.0	6.2	13.1	35.3	107.3	360.6	1,554.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
Non-Hispanic black, both sexes												
2006 5/	52.0	16.8	1.7	0.9	3.9	11.8	22.2	48.4	95.0	191.6	443.4	1,111.9
2005 5/	53.4	15.6	1.8	1.2	4.3	11.7	22.9	50.4	100.2	199.7	455.4	1,152.4
2004 5/	52.7	15.3	2.4	1.1	3.8	10.4	23.0	51.4	98.5	199.4	458.1	1,197.9
2003 5/	54.4	19.8	2.0	1.0	4.5	11.3	24.4	51.1	104.2	219.8	455.6	1,253.2
2002	54.3	20.9	1.6	0.9	4.2	11.8	23.4	52.6	104.1	211.2	474.7	1,285.5
2001	54.7	21.3	3.0	1.0	4.4	11.0	24.5	52.2	105.7	216.1	486.4	1,292.9
2000	54.6	19.6	2.0	1.1	4.9	10.8	23.7	52.8	104.5	223.6	484.3	1,319.4
1999	56.2	23.2	2.0	1.0	4.7	11.0	25.3	53.5	115.3	232.9	499.9	1,314.0
Non-Hispanic black, male												
2006 5/	52.1	17.9	2.0	1.0	4.5	14.8	28.0	63.5	123.2	242.0	513.3	1,122.4
2005 5/	52.6	18.4	*	1.1	5.8	14.6	28.4	63.9	128.3	251.6	520.0	1,107.6
2004 5/	52.3	16.4	2.5	1.0	4.8	12.6	27.7	65.3	128.9	242.3	550.5	1,244.7
2003 5/	53.4	23.6	2.2	1.2	5.8	13.0	29.6	65.1	134.7	269.3	530.4	1,239.1
2002	52.6	21.8	*	0.9	5.1	14.8	28.2	67.5	130.4	259.4	543.4	1,258.0
2001	53.5	18.8	3.3	0.9	5.5	13.4	29.3	63.9	134.9	261.9	593.5	1,362.9
2000	53.3	19.1	2.0	1.2	5.7	12.7	28.7	68.3	132.9	269.8	565.5	1,416.1
1999	56.2	22.9	2.3	1.2	6.0	13.4	30.9	68.7	149.5	291.0	584.7	1,466.6
Non-Hispanic black, female												
2006 5/	51.9	15.8	*	0.8	3.3	8.9	17.0	35.4	72.3	155.6	403.7	1,107.6
2005 5/	54.2	12.7	2.1	1.2	2.8	9.0	18.0	38.9	77.6	162.6	418.8	1,170.4
2004 5/	53.1	14.2	2.3	1.2	2.8	8.4	19.0	39.4	74.0	168.8	405.9	1,179.4
2003 5/	55.4	15.8	1.8	0.9	3.3	9.7	19.8	39.2	79.6	184.6	413.4	1,258.7
2002	55.9	19.9	1.9	0.8	3.2	9.0	19.2	39.9	83.0	177.0	435.9	1,296.0
2001	55.8	23.8	2.7	1.0	3.2	8.9	20.3	42.2	82.3	183.7	426.2	1,267.0
2000	55.8	20.1	1.9	1.0	4.2	9.0	19.2	39.6	81.6	191.1	438.5	1,283.4
1999	56.3	23.6	1.8	0.9	3.4	8.9	20.4	40.5	87.9	192.0	452.4	1,257.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
All origins 3/, both sexes												
2006	0.4	*	*	*	*	0.1	0.2	0.5	0.7	1.2	2.4	2.2
2005	0.4	*	*	*	*	0.1	0.2	0.4	0.7	1.3	2.5	3.0
2004	0.4	*	*	*	*	0.1	0.2	0.5	0.6	1.3	2.5	3.2
2003	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.5	3.0
2002	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4	2.6
2001	0.4	*	*	*	*	0.1	0.3	0.5	0.6	1.3	2.5	2.5
2000	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.4	2.8
1999	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9
All origins 3/, male												
2006	0.4	*	*	*	*	*	0.2	0.7	0.9	1.4	2.7	2.3
2005	0.5	*	*	*	*	0.1	0.2	0.5	0.9	1.5	3.3	5.1
2004	0.5	*	*	*	*	0.1	0.2	0.6	0.7	1.4	3.1	4.8
2003	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.2	4.1
2002	0.5	*	*	*	*	0.2	0.3	0.6	0.7	1.6	3.0	3.5
2001	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.5	3.8
2000	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.5	3.5	4.2
1999	0.5	*	*	*	*	0.1	0.3	0.6	0.9	1.4	3.2	4.7
All origins 3/, female												
2006	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	2.2	2.1
2005	0.3	*	*	*	*	*	0.1	0.3	0.5	1.1	2.0	2.1
2004	0.4	*	*	*	*	*	0.1	0.3	0.4	1.1	2.0	2.5
2003	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	2.0	2.5
2002	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	2.0	2.2
2001	0.4	*	*	*	*	*	0.2	0.3	0.5	1.1	1.8	2.0
2000	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	1.8	2.3
1999	0.3	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
Hispanic, both sexes												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.2	*	*	*	*	*	*	*	*	*	*	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.3	*	*	*	*	*	*	*	*	*	*	*
2004	0.3	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
Non-Hispanic 4/, both sexes												
2006	0.4	*	*	*	*	0.1	0.2	0.5	0.8	1.2	2.5	2.1
2005	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.5	3.1
2004	0.4	*	*	*	*	0.1	0.2	0.5	0.5	1.3	2.5	3.2
2003	0.5	*	*	*	*	0.1	0.3	0.4	0.8	1.3	2.5	3.0
2002	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4	2.7
2001	0.5	*	*	*	*	0.1	0.3	0.5	0.6	1.4	2.5	2.6
2000	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.5	2.9
1999	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9
Non-Hispanic 4/, male												
2006	0.5	*	*	*	*	*	0.2	0.7	1.0	1.4	2.8	2.3
2005	0.5	*	*	*	*	*	0.2	0.6	0.8	1.4	3.3	5.3
2004	0.5	*	*	*	*	*	0.2	0.6	0.7	1.4	3.1	4.8
2003	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.3	4.1
2002	0.5	*	*	*	*	0.2	0.3	0.6	0.7	1.6	3.0	3.6
2001	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.6	3.9
2000	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.4	3.6	4.2
1999	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.2	4.7
Non-Hispanic 4/, female												
2006	0.4	*	*	*	*	*	0.2	0.3	0.6	1.0	2.2	2.1
2005	0.4	*	*	*	*	*	0.1	0.3	0.5	1.1	2.0	2.2
2004	0.4	*	*	*	*	*	0.1	0.3	0.4	1.2	2.0	2.6
2003	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	1.9	2.5
2002	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	2.0	2.3
2001	0.4	*	*	*	*	*	0.3	0.3	0.5	1.2	1.8	2.0
2000	0.4	*	*	*	*	*	0.2	0.3	0.6	1.0	1.8	2.3
1999	0.4	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
Non-Hispanic white, both sexes												
2006 5/	0.4	*	*	*	*	*	0.2	0.4	0.6	1.2	2.4	2.1
2005 5/	0.4	*	*	*	*	*	0.1	0.3	0.5	1.2	2.5	3.2
2004 5/	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.4	3.3
2003 5/	0.5	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.5	3.2
2002	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.5	2.7
2001	0.4	*	*	*	*	*	0.2	0.4	0.5	1.2	2.5	2.7
2000	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.1	2.4	2.8
1999	0.4	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	3.0
Non-Hispanic white, male												
2006 5/	0.5	*	*	*	*	*	0.2	0.5	0.8	1.4	2.7	2.5
2005 5/	0.5	*	*	*	*	*	0.2	0.4	0.6	1.4	3.4	5.6
2004 5/	0.5	*	*	*	*	*	0.2	0.4	0.6	1.4	3.1	4.9
2003 5/	0.5	*	*	*	*	*	0.3	0.4	0.8	1.3	3.4	4.5
2002	0.5	*	*	*	*	0.2	0.3	0.5	0.6	1.5	3.2	3.7
2001	0.5	*	*	*	*	*	0.2	0.5	0.7	1.4	3.6	4.0
2000	0.5	*	*	*	*	*	0.3	0.5	0.8	1.3	3.6	4.3
1999	0.5	*	*	*	*	*	0.2	0.4	0.7	1.3	3.3	5.1
Non-Hispanic white, female												
2006 5/	0.4	*	*	*	*	*	0.2	0.3	0.4	1.0	2.2	1.9
2005 5/	0.4	*	*	*	*	*	*	0.2	0.4	1.1	1.9	2.1
2004 5/	0.4	*	*	*	*	*	*	0.2	0.4	1.1	1.9	2.5
2003 5/	0.4	*	*	*	*	*	0.2	0.2	0.5	1.0	1.9	2.7
2002	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.0	2.3
2001	0.4	*	*	*	*	*	0.2	0.2	0.4	1.1	1.7	2.1
2000	0.4	*	*	*	*	*	0.1	0.2	0.4	1.0	1.7	2.2
1999	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8	2.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
Non-Hispanic black, both sexes												
2006 5/	0.6	*	*	*	*	*	0.4	1.1	2.0	1.7	3.5	*
2005 5/	0.6	*	*	*	*	*	0.5	1.1	2.0	1.8	3.7	*
2004 5/	0.5	*	*	*	*	*	*	1.4	1.2	2.1	2.8	*
2003 5/	0.6	*	*	*	*	*	0.6	1.3	2.0	2.2	3.0	*
2002	0.5	*	*	*	*	*	0.4	1.3	1.6	2.3	*	*
2001	0.7	*	*	*	*	*	0.7	1.3	1.7	2.6	4.0	*
2000	0.6	*	*	*	*	*	0.6	1.4	1.6	1.9	2.9	*
1999	0.6	*	*	*	*	*	0.7	1.2	2.0	2.6	*	*
Non-Hispanic black, male												
2006 5/	0.7	*	*	*	*	*	*	1.5	2.7	*	*	*
2005 5/	0.7	*	*	*	*	*	*	1.5	2.8	*	*	*
2004 5/	0.6	*	*	*	*	*	*	1.9	*	*	*	*
2003 5/	0.7	*	*	*	*	*	*	2.0	2.3	*	*	*
2002	0.6	*	*	*	*	*	*	1.6	1.9	*	*	*
2001	0.8	*	*	*	*	*	0.8	1.9	2.2	3.9	*	*
2000	0.7	*	*	*	*	*	0.8	1.8	*	*	*	*
1999	0.7	*	*	*	*	*	1.0	1.9	2.5	*	*	*
Non-Hispanic black, female												
2006 5/	0.5	*	*	*	*	*	*	*	1.4	*	3.2	*
2005 5/	0.5	*	*	*	*	*	*	0.8	1.3	*	4.0	*
2004 5/	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2003 5/	0.5	*	*	*	*	*	*	*	1.7	2.1	*	*
2002	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2001	0.5	*	*	*	*	*	0.7	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	1.0	1.6	*	*	*
1999	0.5	*	*	*	*	*	*	*	1.6	2.9	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
All origins 3/, both sexes												
2006	0.3	*	0.1	0.1	0.1	0.1	0.2	0.2	0.4	0.6	1.2	1.8
2005	0.3	0.5	*	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2004	0.3	0.5	0.2	*	0.1	0.1	0.2	0.3	0.4	0.6	1.2	1.8
2003	0.3	0.8	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	1.6
2002	0.3	0.6	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	2.1
2001	0.3	0.8	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000	0.3	0.5	0.2	0.0	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.8
1999	0.3	0.7	0.2	0.1	0.1	0.1	0.3	0.3	0.5	0.8	1.3	2.0
All origins 3/, male												
2006	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.3	1.8
2005	0.3	*	*	*	0.1	0.2	0.2	0.4	0.4	0.8	1.5	1.9
2004	0.3	*	*	*	*	*	0.2	0.3	0.5	0.7	1.3	2.3
2003	0.3	*	*	*	0.2	0.1	0.2	0.4	0.4	1.0	1.3	1.5
2002	0.3	*	*	*	0.1	0.2	0.2	0.4	0.5	0.9	1.2	2.4
2001	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000	0.3	*	*	*	0.1	0.1	0.3	0.2	0.4	0.8	1.5	2.1
1999	0.3	*	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.4	2.5
All origins 3/, female												
2006	0.3	*	*	*	*	*	0.2	0.2	0.3	0.5	1.0	1.8
2005	0.3	*	*	0.1	*	0.1	0.2	0.3	0.3	0.6	1.0	1.9
2004	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.5	1.2	1.6
2003	0.3	*	*	*	0.1	*	0.2	0.2	0.3	0.7	1.1	1.7
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.3	2.0
2001	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	2.0
2000	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.0	1.7
1999	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	1.3	1.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Hispanic, both sexes												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.2	*	*	*	*	*	*	*	*	*	*	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.2	*	*	*	*	*	*	*	*	*	*	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.2	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Non-Hispanic 4/, both sexes												
2006	0.3	*	*	*	0.1	0.1	0.2	0.3	0.4	0.6	1.2	1.8
2005	0.3	*	*	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2004	0.3	*	*	*	0.1	0.1	0.2	0.3	0.4	0.6	1.2	1.8
2003	0.3	0.9	*	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	1.7
2002	0.3	0.7	0.2	*	0.1	0.1	0.2	0.3	0.4	0.8	1.3	2.1
2001	0.3	0.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.1	1.8
1999	0.3	0.8	0.2	*	0.1	0.1	0.3	0.3	0.6	0.8	1.3	2.1
Non-Hispanic 4/, male												
2006	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.4	1.8
2005	0.3	*	*	*	*	0.1	0.2	0.4	0.4	0.8	1.5	2.0
2004	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	1.3	2.4
2003	0.3	*	*	*	0.2	0.2	0.3	0.4	0.4	1.0	1.3	1.6
2002	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.9	1.2	2.3
2001	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000	0.3	*	*	*	0.1	0.2	0.3	0.3	0.4	0.9	1.4	2.1
1999	0.3	*	*	*	*	0.2	0.3	0.3	0.6	0.9	1.4	2.6
Non-Hispanic 4/, female												
2006	0.3	*	*	*	*	*	0.2	0.2	0.3	0.5	1.1	1.9
2005	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.0	1.9
2004	0.3	*	*	*	*	*	0.1	0.2	0.3	0.5	1.2	1.6
2003	0.3	*	*	*	*	*	0.2	0.2	0.3	0.7	1.1	1.7
2002	0.3	*	*	*	*	*	0.1	0.2	0.3	0.7	1.3	2.0
2001	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	1.9
2000	0.3	*	*	*	*	*	0.1	0.2	0.3	0.6	1.0	1.7
1999	0.3	*	*	*	*	*	0.2	0.2	0.5	0.7	1.2	1.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Non-Hispanic white, both sexes												
2006 5/	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.6	1.3	1.9
2005 5/	0.3	*	*	*	0.1	0.1	0.2	0.3	0.3	0.7	1.1	1.9
2004 5/	0.3	*	*	*	*	0.1	0.1	0.2	0.4	0.6	1.2	1.8
2003 5/	0.3	*	*	*	0.1	*	0.2	0.2	0.4	0.9	1.2	1.7
2002	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.8	1.2	2.1
2001	0.3	*	*	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.1	1.8
1999	0.3	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.3	2.0
Non-Hispanic white, male												
2006 5/	0.3	*	*	*	*	*	0.2	0.3	0.5	0.8	1.5	1.9
2005 5/	0.3	*	*	*	*	*	0.2	0.3	0.3	0.8	1.4	2.0
2004 5/	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.2	2.3
2003 5/	0.3	*	*	*	0.2	*	0.2	0.3	0.4	1.0	1.4	1.8
2002	0.3	*	*	*	*	0.2	0.2	0.3	0.4	0.9	1.2	2.3
2001	0.3	*	*	*	*	*	0.2	0.3	0.5	0.7	1.2	2.0
2000	0.3	*	*	*	*	*	0.2	0.2	0.4	0.8	1.4	2.1
1999	0.3	*	*	*	*	*	0.2	0.3	0.7	0.8	1.4	2.6
Non-Hispanic white, female												
2006 5/	0.3	*	*	*	*	*	0.2	0.2	0.3	0.5	1.1	1.9
2005 5/	0.3	*	*	*	*	*	0.2	0.2	0.3	0.6	0.9	1.8
2004 5/	0.3	*	*	*	*	*	*	0.2	0.3	0.6	1.1	1.6
2003 5/	0.3	*	*	*	*	*	*	0.2	0.3	0.7	1.1	1.7
2002	0.3	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.1
2001	0.3	*	*	*	*	*	0.2	0.2	0.3	0.6	1.3	1.8
2000	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.0	1.7
1999	0.3	*	*	*	*	*	0.2	0.2	0.5	0.7	1.2	1.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Non-Hispanic black, both sexes												
2006 5/	0.3	*	*	*	*	*	*	0.4	*	*	*	*
2005 5/	0.4	*	*	*	*	*	*	0.6	0.8	*	2.1	*
2004 5/	0.4	*	*	*	*	*	0.5	0.5	*	*	2.3	*
2003 5/	0.5	*	*	*	*	0.4	0.5	0.5	*	*	*	*
2002	0.4	*	*	*	*	*	*	0.6	0.9	*	2.1	*
2001	0.4	*	*	*	*	*	0.4	0.7	*	*	*	*
2000	0.4	*	*	*	*	*	0.4	0.5	*	*	*	*
1999	0.5	*	*	*	*	*	0.5	0.5	*	1.2	*	*
Non-Hispanic black, male												
2006 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.5	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.5	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.5	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.5	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
All origins 3/, both sexes												
2006	20.2	*	*	*	0.1	0.2	0.8	2.8	8.9	31.5	132.7	614.4
2005	19.9	0.7	*	*	0.1	0.2	0.7	2.6	8.6	31.6	132.8	618.6
2004	19.5	0.7	*	*	0.1	0.2	0.7	2.5	8.4	32.2	131.2	621.2
2003	19.8	0.9	*	*	0.1	0.2	0.7	2.4	8.8	33.5	133.2	641.1
2002	19.6	0.6	*	*	0.1	0.3	0.6	2.5	8.8	33.8	132.8	645.0
2001	20.0	0.6	*	*	0.1	0.2	0.7	2.4	9.4	34.3	136.3	674.6
2000	19.8	0.8	*	0.1	0.1	0.2	0.6	2.2	9.1	34.8	136.1	684.9
1999	19.7	*	*	*	0.1	0.2	0.6	2.1	8.9	35.4	137.2	681.9
All origins 3/, male												
2006	16.2	*	*	*	0.1	0.3	0.9	3.7	10.8	36.9	148.6	599.0
2005	15.8	*	*	*	0.1	0.3	0.9	3.1	10.3	36.0	147.8	603.1
2004	15.4	*	*	*	*	0.3	0.8	3.0	10.1	37.5	147.0	605.5
2003	15.7	1.4	*	*	0.1	0.3	0.9	2.9	10.8	39.7	148.6	629.8
2002	15.3	*	*	*	*	0.3	0.8	3.1	10.5	39.2	149.0	621.4
2001	15.5	*	*	*	*	0.3	0.8	3.1	11.1	40.3	153.8	646.4
2000	15.3	1.1	*	*	0.1	0.2	0.7	2.7	10.9	41.1	153.7	672.2
1999	15.2	*	*	*	0.1	0.2	0.7	2.7	10.8	42.5	152.0	667.0
All origins 3/, female												
2006	24.0	*	*	*	*	0.2	0.6	2.0	7.0	26.9	121.8	621.6
2005	23.9	*	*	*	*	0.2	0.5	2.0	7.0	27.9	122.5	625.6
2004	23.4	*	*	*	*	0.2	0.6	2.0	6.8	27.8	120.5	628.2
2003	23.7	*	*	*	*	0.2	0.5	1.9	7.0	28.3	122.9	646.1
2002	23.7	*	*	*	*	0.2	0.5	1.8	7.2	29.2	122.0	655.3
2001	24.3	*	*	*	*	0.2	0.6	1.8	7.8	29.3	124.7	686.4
2000	24.1	*	*	*	*	0.1	0.5	1.6	7.5	29.6	124.6	690.0
1999	24.0	*	*	*	*	0.2	0.5	1.6	7.1	29.6	127.6	687.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
Hispanic, both sexes												
2006	4.1	*	*	*	*	*	*	1.5	4.4	18.8	74.5	304.8
2005	4.0	*	*	*	*	*	*	1.1	4.5	18.5	75.6	316.6
2004	3.7	*	*	*	*	*	0.4	1.3	4.7	19.2	72.9	287.3
2003	4.0	*	*	*	*	*	*	1.3	5.7	22.5	77.1	330.1
2002	3.6	*	*	*	*	*	*	1.2	4.6	19.9	75.2	316.0
2001	3.7	*	*	*	*	*	0.4	1.3	5.6	18.8	72.9	349.5
2000	3.6	*	*	*	*	*	*	0.7	5.8	18.9	72.1	369.6
1999	3.7	*	*	*	*	*	*	1.0	5.3	19.5	73.5	377.4
Hispanic, male												
2006	3.5	*	*	*	*	*	*	2.1	4.9	22.9	89.8	276.3
2005	3.3	*	*	*	*	*	*	1.4	4.2	21.5	90.2	295.1
2004	3.1	*	*	*	*	*	*	1.7	6.0	22.1	84.1	263.4
2003	3.5	*	*	*	*	*	*	1.3	7.8	26.0	92.8	312.5
2002	3.0	*	*	*	*	*	*	1.1	5.8	23.1	85.8	296.2
2001	3.0	*	*	*	*	*	*	1.8	6.0	21.8	75.6	333.3
2000	2.9	*	*	*	*	*	*	*	7.7	23.0	74.2	380.9
1999	3.0	*	*	*	*	*	*	*	6.1	24.6	82.9	358.0
Hispanic, female												
2006	4.8	*	*	*	*	*	*	0.9	4.0	15.5	63.7	319.9
2005	4.8	*	*	*	*	*	*	*	4.7	16.0	65.4	327.8
2004	4.4	*	*	*	*	*	*	1.0	3.4	16.9	65.2	299.6
2003	4.6	*	*	*	*	*	*	1.2	3.7	19.7	66.4	339.1
2002	4.4	*	*	*	*	*	*	1.3	3.6	17.3	68.0	325.8
2001	4.5	*	*	*	*	*	*	*	5.2	16.4	71.0	357.4
2000	4.3	*	*	*	*	*	*	*	4.2	15.6	70.7	364.0
1999	4.4	*	*	*	*	*	*	*	4.6	15.5	67.3	386.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
Non-Hispanic 4/, both sexes												
2006	22.9	*	*	*	0.1	0.2	0.8	3.0	9.2	32.4	136.1	629.3
2005	22.5	0.7	*	*	0.1	0.3	0.8	2.7	8.9	32.5	136.0	632.5
2004	22.0	0.8	*	*	0.1	0.3	0.8	2.6	8.7	33.1	134.2	635.3
2003	22.2	1.0	*	*	0.1	0.3	0.8	2.5	9.1	34.1	135.9	653.2
2002	22.0	*	*	*	0.1	0.3	0.7	2.6	9.1	34.6	135.3	656.8
2001	22.4	*	*	*	0.1	0.3	0.7	2.5	9.6	35.2	138.8	685.6
2000	22.1	0.8	*	*	0.1	0.2	0.6	2.3	9.3	35.7	138.5	694.8
1999	21.8	*	*	*	0.1	0.2	0.6	2.2	9.1	36.3	139.4	691.0
Non-Hispanic 4/, male												
2006	18.5	*	*	*	0.1	0.3	1.0	3.9	11.3	37.9	152.1	616.3
2005	18.0	*	*	*	0.1	0.3	1.0	3.3	10.8	37.0	151.0	618.8
2004	17.5	*	*	*	*	0.3	0.9	3.1	10.4	38.6	150.3	621.3
2003	17.7	1.4	*	*	0.1	0.3	1.0	3.1	11.0	40.5	151.2	643.2
2002	17.3	*	*	*	*	0.4	0.9	3.3	10.9	40.2	151.8	634.7
2001	17.4	*	*	*	*	0.3	0.9	3.2	11.4	41.4	157.2	658.0
2000	17.2	*	*	*	0.1	0.2	0.8	2.9	11.0	42.1	156.9	683.3
1999	16.9	*	*	*	0.1	0.2	0.8	2.8	11.0	43.5	154.5	677.1
Non-Hispanic 4/, female												
2006	27.1	*	*	*	*	0.2	0.7	2.2	7.3	27.8	125.2	635.3
2005	26.9	*	*	*	*	0.2	0.6	2.1	7.2	28.8	125.7	638.7
2004	26.3	*	*	*	*	0.2	0.6	2.1	7.1	28.6	123.3	641.6
2003	26.5	*	*	*	*	0.2	0.6	2.0	7.3	28.8	125.6	657.6
2002	26.5	*	*	*	*	0.2	0.5	1.9	7.5	29.9	124.3	666.4
2001	27.1	*	*	*	*	0.2	0.6	1.9	8.0	30.1	126.8	697.0
2000	26.7	*	*	*	*	0.2	0.5	1.7	7.7	30.5	126.5	699.5
1999	26.5	*	*	*	*	0.2	0.5	1.6	7.2	30.4	129.7	696.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
Non-Hispanic white, both sexes												
2006 5/	26.0	*	*	*	*	0.1	0.6	2.2	8.2	31.2	138.3	653.7
2005 5/	25.4	*	*	*	0.1	0.2	0.5	2.1	7.8	31.1	136.7	655.9
2004 5/	24.8	*	*	*	*	0.2	0.6	2.0	7.6	32.3	135.7	657.1
2003 5/	25.0	*	*	*	*	0.2	0.5	1.9	8.0	32.8	137.1	673.5
2002	24.7	*	*	*	*	0.2	0.5	2.0	8.2	33.3	135.9	677.3
2001	25.1	*	*	*	*	0.2	0.5	2.1	8.7	33.9	139.7	706.8
2000	24.7	*	*	*	*	0.1	0.5	1.9	8.4	34.3	139.2	714.3
1999	24.3	*	*	*	*	0.1	0.5	1.7	8.1	34.9	139.1	709.1
Non-Hispanic white, male												
2006 5/	20.7	*	*	*	*	0.2	0.7	2.8	10.2	36.4	154.9	641.8
2005 5/	20.0	*	*	*	*	0.2	0.7	2.5	9.4	35.2	152.3	646.8
2004 5/	19.5	*	*	*	*	0.2	0.7	2.4	9.0	37.8	151.7	643.0
2003 5/	19.7	*	*	*	*	0.3	0.7	2.4	9.7	39.3	153.2	665.3
2002	19.2	*	*	*	*	0.2	0.7	2.6	9.8	38.6	153.2	656.2
2001	19.3	*	*	*	*	0.2	0.7	2.7	10.5	40.0	158.1	679.1
2000	19.0	*	*	*	*	0.2	0.6	2.4	10.1	41.0	157.9	701.6
1999	18.6	*	*	*	*	*	0.6	2.2	10.1	42.2	154.7	690.8
Non-Hispanic white, female												
2006 5/	31.0	*	*	*	*	*	0.5	1.6	6.3	26.8	126.8	659.3
2005 5/	30.5	*	*	*	*	*	0.4	1.6	6.2	27.5	125.9	660.0
2004 5/	30.0	*	*	*	*	*	0.4	1.5	6.2	27.6	124.7	663.4
2003 5/	30.1	*	*	*	*	*	0.4	1.5	6.3	27.2	126.3	677.2
2002	30.0	*	*	*	*	0.2	0.4	1.4	6.7	28.8	124.2	686.4
2001	30.7	*	*	*	*	0.2	0.4	1.4	6.9	28.8	127.5	718.4
2000	30.2	*	*	*	*	*	0.3	1.3	6.8	28.7	126.9	719.4
1999	29.9	*	*	*	*	*	0.3	1.2	6.3	28.8	128.8	716.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
Non-Hispanic black, both sexes												
2006 5/	14.6	*	*	*	*	0.8	2.3	8.7	20.0	52.5	149.9	492.6
2005 5/	14.9	*	*	*	*	0.8	2.4	7.5	20.7	54.8	163.9	497.8
2004 5/	14.1	*	*	*	*	0.7	2.0	7.7	19.9	49.2	153.0	513.2
2003 5/	14.5	*	*	*	*	0.7	2.4	7.1	21.0	55.3	152.7	538.7
2002	14.2	*	*	*	*	0.9	1.9	7.4	19.0	54.1	158.3	534.5
2001	14.3	*	*	*	*	0.6	2.0	6.6	20.0	55.5	159.1	551.2
2000	14.1	*	*	*	*	0.5	1.7	5.9	19.0	56.2	156.3	564.8
1999	14.3	*	*	*	*	0.6	1.7	6.5	19.2	57.3	164.8	557.4
Non-Hispanic black, male												
2006 5/	12.9	*	*	*	*	1.0	3.1	11.5	24.1	64.6	168.5	473.7
2005 5/	12.7	*	*	*	*	1.1	3.5	9.4	26.0	65.0	182.8	431.0
2004 5/	12.2	*	*	*	*	1.0	2.3	9.5	25.6	56.6	178.5	510.1
2003 5/	12.3	*	*	*	*	0.8	2.8	8.5	26.8	64.3	165.9	521.0
2002	12.0	*	*	*	*	1.1	2.3	9.2	23.3	65.9	176.4	489.0
2001	12.0	*	*	*	*	0.9	2.3	7.6	22.6	66.3	185.4	547.5
2000	11.6	*	*	*	*	*	1.8	7.3	21.8	62.9	176.5	589.4
1999	12.2	*	*	*	*	*	1.9	8.6	23.0	66.4	180.4	609.7
Non-Hispanic black, female												
2006 5/	16.2	*	*	*	*	*	1.7	6.2	16.6	43.8	139.3	500.4
2005 5/	16.9	*	*	*	*	*	1.4	5.8	16.5	47.5	153.2	524.7
2004 5/	15.7	*	*	*	*	*	1.8	6.1	15.4	43.9	138.6	514.4
2003 5/	16.5	*	*	*	*	*	2.0	5.8	16.3	48.9	145.2	545.6
2002	16.3	*	*	*	*	*	1.5	5.7	15.5	45.8	148.1	551.8
2001	16.4	*	*	*	*	*	1.7	5.6	18.0	47.9	144.4	552.6
2000	16.3	*	*	*	*	*	1.6	4.7	16.8	51.5	145.0	555.6
1999	16.2	*	*	*	*	*	1.5	4.8	16.1	50.9	156.0	537.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
All origins 3/, both sexes												
2006	36.0	6.9	0.7	0.5	1.9	4.3	8.3	17.6	34.4	79.0	224.5	724.4
2005	37.0	7.1	0.7	0.5	2.0	4.2	8.7	18.2	36.0	83.2	230.2	754.0
2004	36.5	8.7	0.9	0.5	1.9	3.9	8.7	18.3	35.8	84.5	234.1	747.7
2003	38.0	8.6	0.9	0.5	1.9	4.1	9.1	18.2	38.1	91.2	246.0	781.5
2002	38.5	10.5	0.8	0.5	1.9	4.0	9.3	18.6	38.8	93.7	254.4	793.7
2001	37.9	9.7	1.1	0.5	1.8	4.0	8.8	18.5	39.1	95.1	249.5	796.5
2000	38.3	10.6	0.9	0.5	2.0	4.0	8.8	18.6	40.8	97.4	251.6	820.0
1999	39.0	11.9	0.9	0.5	2.1	4.1	9.2	18.7	42.8	101.3	257.6	826.1
All origins 3/, male												
2006	34.0	6.9	0.8	0.6	2.3	5.4	10.3	22.6	43.9	97.1	258.4	735.0
2005	34.9	7.7	0.8	0.5	2.6	5.2	10.9	23.6	46.0	102.3	263.3	767.7
2004	34.5	9.2	0.9	0.5	2.4	4.8	10.7	23.8	46.3	103.1	269.4	765.2
2003	35.8	9.2	0.9	0.6	2.3	4.8	11.3	23.3	48.5	112.1	287.0	795.9
2002	36.3	10.9	0.8	0.5	2.5	5.0	11.5	24.1	49.0	114.8	294.9	821.3
2001	35.6	9.1	1.2	0.5	2.2	4.7	11.1	23.6	49.4	116.1	293.4	821.3
2000	36.1	10.5	1.0	0.6	2.3	4.8	10.9	24.3	52.6	120.7	292.4	870.7
1999	37.1	12.1	1.0	0.6	2.5	4.8	11.6	24.2	55.3	127.9	303.1	874.7
All origins 3/, female												
2006	38.0	6.8	0.7	0.4	1.4	3.1	6.4	12.8	25.6	63.7	201.3	719.5
2005	39.0	6.4	0.7	0.4	1.3	3.1	6.5	13.0	26.8	67.1	207.8	747.6
2004	38.5	8.2	0.8	0.5	1.3	2.9	6.6	13.0	26.0	68.9	210.3	739.9
2003	40.1	7.9	1.0	0.4	1.5	3.4	7.0	13.3	28.4	73.7	218.6	775.1
2002	40.6	10.0	0.8	0.4	1.3	3.1	7.1	13.2	29.3	76.2	227.6	781.7
2001	40.2	10.4	1.0	0.5	1.3	3.2	6.5	13.5	29.6	77.7	220.7	786.1
2000	40.4	10.7	0.7	0.4	1.6	3.2	6.8	13.1	29.9	78.3	225.0	799.3
1999	40.9	11.6	0.8	0.5	1.7	3.3	6.9	13.5	31.3	79.6	228.0	806.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Hispanic, both sexes												
2006	9.2	5.3	0.6	0.4	1.2	2.6	4.2	9.8	18.6	47.4	120.2	354.5
2005	9.5	6.4	*	0.3	1.4	2.5	4.5	9.2	19.3	50.3	132.5	384.2
2004	9.1	7.2	0.7	0.3	1.4	2.1	4.5	9.5	18.7	49.6	131.9	369.1
2003	9.7	6.8	0.9	0.3	1.2	2.6	4.8	9.5	23.2	53.0	141.3	418.2
2002	9.3	8.6	0.6	0.3	1.4	2.4	4.7	9.6	19.7	50.9	141.6	419.6
2001	9.3	8.0	0.8	0.4	1.2	2.2	4.3	9.2	21.6	55.3	141.6	444.6
2000	9.6	8.7	0.7	0.5	1.6	2.5	4.9	10.2	22.0	54.9	152.1	457.2
1999	9.8	11.5	0.9	0.4	1.3	2.5	5.5	10.1	24.6	57.2	151.9	447.6
Hispanic, male												
2006	9.6	5.8	*	0.5	1.7	3.6	5.5	12.2	24.6	58.1	139.8	378.9
2005	10.0	6.9	*	*	1.7	3.4	6.1	12.7	26.3	62.3	153.7	401.0
2004	9.4	6.9	*	*	2.0	3.1	6.2	12.8	23.5	59.9	153.8	374.1
2003	9.8	7.0	*	*	1.6	3.0	6.4	12.7	28.3	66.2	159.1	437.2
2002	9.6	8.0	*	*	2.0	3.1	6.0	12.2	24.2	67.1	168.2	455.1
2001	9.3	7.9	*	*	1.5	2.8	5.7	12.6	27.3	67.7	170.4	438.1
2000	9.9	8.1	*	0.7	2.1	3.3	6.0	13.7	28.6	68.8	172.6	505.9
1999	10.3	9.8	*	*	1.5	3.4	7.2	13.7	31.9	75.6	179.4	526.2
Hispanic, female												
2006	8.8	4.6	*	*	0.7	1.3	2.8	7.4	13.2	38.7	106.4	341.5
2005	9.0	5.9	*	*	1.0	1.4	2.7	5.6	12.9	40.5	117.7	375.4
2004	8.8	7.4	*	*	0.7	1.0	2.6	6.1	14.3	41.3	116.8	366.6
2003	9.6	6.6	*	*	0.8	2.0	3.1	6.3	18.6	42.4	129.0	408.6
2002	9.0	9.3	*	*	0.8	1.5	3.3	7.0	15.8	38.0	123.4	402.0
2001	9.2	8.0	*	*	0.9	1.6	2.8	5.8	16.5	45.5	122.1	447.7
2000	9.3	9.3	*	*	1.0	1.7	3.8	6.8	16.1	44.0	138.3	433.3
1999	9.2	13.3	*	*	1.1	1.5	3.7	6.6	18.1	42.9	133.6	409.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Non-Hispanic 4/, both sexes												
2006	40.6	7.3	0.8	0.5	2.0	4.7	9.0	18.5	35.7	81.3	230.6	742.4
2005	41.5	7.2	0.8	0.5	2.1	4.6	9.4	19.1	37.4	85.5	235.7	770.9
2004	40.9	9.1	0.9	0.5	2.0	4.3	9.3	19.2	37.1	87.0	239.4	764.1
2003	42.4	9.0	0.9	0.5	2.1	4.4	9.7	19.1	39.1	93.7	250.9	795.8
2002	42.9	10.9	0.8	0.5	2.0	4.4	9.9	19.4	40.2	96.5	259.4	807.6
2001	42.1	10.1	1.2	0.5	1.9	4.3	9.4	19.3	40.4	97.5	253.9	807.8
2000	42.3	11.0	0.9	0.5	2.0	4.3	9.3	19.3	42.0	99.8	255.4	831.4
1999	43.0	11.9	0.9	0.5	2.2	4.3	9.6	19.5	44.0	103.6	261.3	837.9
Non-Hispanic 4/, male												
2006	38.5	7.2	0.8	0.6	2.5	5.9	11.1	23.7	45.5	99.8	265.4	753.7
2005	39.2	7.9	0.8	0.5	2.8	5.7	11.7	24.7	47.4	105.0	269.4	786.5
2004	38.8	9.7	0.9	0.5	2.4	5.2	11.4	24.9	48.1	106.0	275.4	784.1
2003	40.0	9.9	0.8	0.6	2.5	5.2	12.0	24.3	49.8	115.1	293.1	811.6
2002	40.6	11.6	0.8	0.6	2.6	5.5	12.3	25.3	50.8	117.7	300.6	836.7
2001	39.6	9.3	1.4	0.5	2.4	5.1	11.8	24.5	50.9	118.8	298.5	835.3
2000	40.0	10.9	1.0	0.6	2.4	5.1	11.5	25.1	54.1	123.4	297.1	883.9
1999	40.9	12.7	1.0	0.6	2.6	5.1	12.1	25.0	56.7	130.5	307.4	886.8
Non-Hispanic 4/, female												
2006	42.7	7.4	0.8	0.4	1.5	3.6	7.0	13.4	26.6	65.5	206.9	737.2
2005	43.7	6.5	0.8	0.5	1.4	3.4	7.1	13.7	28.0	69.1	212.8	763.9
2004	43.0	8.3	0.9	0.5	1.4	3.4	7.2	13.7	27.0	70.9	215.2	755.1
2003	44.6	8.1	1.0	0.4	1.6	3.6	7.5	14.0	29.1	75.8	222.8	788.8
2002	45.1	10.1	0.8	0.4	1.4	3.3	7.6	13.8	30.3	78.7	232.2	795.0
2001	44.5	11.0	1.0	0.5	1.4	3.5	6.9	14.2	30.6	79.7	224.6	796.3
2000	44.5	11.0	0.8	0.4	1.7	3.4	7.1	13.6	30.9	80.4	228.3	810.2
1999	44.9	11.1	0.9	0.5	1.8	3.6	7.2	14.1	32.2	81.5	231.4	818.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Non-Hispanic white, both sexes												
2006 5/	43.7	5.4	0.6	0.4	1.7	3.7	7.6	16.1	32.2	76.8	230.0	765.0
2005 5/	44.5	5.9	0.7	0.4	1.7	3.6	8.0	16.4	33.4	81.0	235.8	792.0
2004 5/	43.7	7.9	0.7	0.4	1.7	3.4	7.8	16.5	33.5	82.4	238.0	779.9
2003 5/	45.2	7.1	0.7	0.5	1.7	3.5	8.2	16.3	35.1	88.3	250.6	811.6
2002	45.8	9.2	0.7	0.5	1.6	3.4	8.4	16.5	36.2	92.3	258.6	820.9
2001	44.7	8.1	0.9	0.4	1.5	3.4	7.6	16.3	36.2	92.9	251.4	821.8
2000	44.9	9.6	0.8	0.4	1.6	3.4	7.6	16.3	38.0	95.0	253.1	845.3
1999	45.4	9.8	0.8	0.5	1.9	3.4	7.8	16.4	39.0	98.3	258.4	851.7
Non-Hispanic white, male												
2006 5/	40.5	5.2	0.6	0.5	2.2	4.7	9.4	20.7	40.9	94.4	264.8	774.7
2005 5/	41.4	6.2	0.7	0.5	2.3	4.5	10.2	21.6	42.7	99.1	270.0	808.1
2004 5/	40.6	8.4	0.7	0.4	2.0	4.2	9.7	21.7	43.6	101.0	272.8	795.1
2003 5/	42.0	7.5	0.6	0.6	2.0	4.2	10.2	21.0	45.1	108.9	292.5	827.7
2002	42.7	9.8	0.8	0.5	2.1	4.2	10.7	21.9	46.2	113.1	299.5	850.7
2001	41.3	8.1	1.0	0.5	1.9	4.0	9.9	21.3	45.8	113.8	294.4	845.6
2000	41.7	9.6	0.8	0.5	1.8	4.1	9.6	21.5	49.3	117.9	294.7	898.3
1999	42.4	10.8	0.9	0.5	2.1	4.1	10.1	21.5	50.6	124.1	303.5	897.0
Non-Hispanic white, female												
2006 5/	46.7	5.6	0.6	0.4	1.2	2.8	5.8	11.5	23.9	61.6	205.8	760.4
2005 5/	47.6	5.5	0.6	0.3	1.2	2.6	5.8	11.2	24.5	65.4	212.2	784.6
2004 5/	46.7	7.3	0.7	0.4	1.2	2.6	5.8	11.3	23.8	66.3	214.4	773.1
2003 5/	48.3	6.7	0.8	0.3	1.4	2.7	6.1	11.6	25.6	70.6	222.3	804.5
2002	48.9	8.5	0.6	0.4	1.1	2.5	6.2	11.2	26.6	74.6	231.1	808.1
2001	47.9	8.1	0.7	0.4	1.1	2.7	5.3	11.5	27.2	75.2	222.8	811.9
2000	47.9	9.6	0.7	0.3	1.3	2.7	5.6	11.1	27.4	75.7	225.7	824.0
1999	48.2	8.7	0.6	0.5	1.6	2.8	5.5	11.4	28.1	76.7	228.9	833.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Non-Hispanic black, both sexes												
2006 5/	36.5	14.3	1.5	0.8	3.5	10.6	19.2	38.2	72.5	136.9	288.9	614.7
2005 5/	37.4	12.7	1.2	0.9	3.9	10.5	19.7	41.3	76.7	142.3	285.8	649.7
2004 5/	37.7	12.8	1.9	0.9	3.5	9.4	20.3	41.7	76.7	147.1	300.0	678.7
2003 5/	38.8	15.2	1.6	0.8	4.0	10.1	20.9	42.3	80.8	161.3	298.6	711.8
2002	39.1	17.8	1.2	0.7	3.9	10.5	20.9	43.4	82.7	153.6	312.3	747.1
2001	39.3	18.2	2.4	0.9	3.8	10.0	21.4	43.7	83.3	157.1	322.1	737.9
2000	39.5	16.1	1.5	0.9	4.5	9.8	21.0	45.1	83.1	164.4	323.9	748.9
1999	40.9	20.9	1.5	0.9	4.4	10.1	22.3	45.2	93.3	171.8	331.4	752.7
Non-Hispanic black, male												
2006 5/	38.2	14.8	1.7	0.9	4.0	13.4	24.1	49.9	95.9	174.7	340.1	647.8
2005 5/	38.8	15.2	*	1.0	5.4	12.9	23.9	52.5	98.6	184.1	331.0	672.7
2004 5/	39.0	13.9	1.9	0.7	4.5	11.2	24.6	53.0	101.2	181.6	366.5	728.6
2003 5/	39.9	18.3	1.9	0.9	5.1	11.4	25.6	53.9	105.1	201.8	359.8	718.1
2002	39.6	19.7	*	0.8	4.7	13.3	25.0	55.8	104.0	189.7	363.1	763.6
2001	40.3	16.0	2.7	0.9	5.0	11.9	25.9	53.5	109.1	190.6	401.2	813.1
2000	40.5	15.0	1.7	1.0	5.1	11.7	25.4	58.6	108.9	202.9	384.1	821.9
1999	42.8	20.8	*	1.1	5.6	12.1	27.4	57.6	123.1	220.6	401.5	855.7
Non-Hispanic black, female												
2006 5/	34.9	13.8	*	0.7	3.0	8.0	14.9	28.3	53.7	109.8	259.8	601.3
2005 5/	36.2	10.1	*	0.8	2.5	8.2	15.9	31.8	59.0	112.3	260.1	640.4
2004 5/	36.5	11.6	1.8	1.1	2.5	7.7	16.5	32.0	57.0	122.6	262.4	659.0
2003 5/	37.8	11.9	*	0.7	2.8	8.8	16.7	32.3	61.2	132.5	264.0	709.4
2002	38.7	15.8	*	*	3.0	8.0	17.2	32.8	65.5	127.9	283.5	740.8
2001	38.5	20.5	2.1	0.9	2.7	8.3	17.3	35.3	62.5	133.5	277.6	710.1
2000	38.6	17.3	*	0.8	3.8	8.1	17.0	33.4	62.4	137.3	290.0	721.7
1999	39.2	21.1	*	*	3.2	8.2	17.9	34.7	69.5	137.3	292.2	714.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential hypertension and hypertensive renal disease (I10,I12,I15)												
All origins 3/, both sexes												
2006	8.0	*	*	*	0.0	0.3	0.9	3.0	6.9	16.8	51.0	189.4
2005	8.4	*	*	*	0.1	0.2	0.9	2.7	6.4	17.7	55.6	210.0
2004	7.9	*	*	*	0.1	0.3	0.8	2.7	6.3	17.1	52.6	198.5
2003	7.5	*	*	*	0.1	0.2	0.8	2.5	6.3	16.9	51.7	188.9
2002	7.0	*	*	*	0.1	0.2	0.8	2.3	5.7	16.0	48.2	180.4
2001	6.8	*	*	*	0.1	0.3	0.7	2.4	5.8	15.5	47.7	171.9
2000	6.4	*	*	*	*	0.2	0.8	2.3	5.9	15.1	45.5	162.9
1999	6.1	*	*	*	*	0.2	0.7	2.2	5.5	15.2	43.6	152.1
All origins 3/, male												
2006	6.4	*	*	*	*	0.3	1.1	3.8	8.4	18.5	51.2	159.2
2005	6.5	*	*	*	*	0.2	1.1	3.5	7.9	19.2	55.3	169.3
2004	6.1	*	*	*	*	0.3	1.0	3.5	7.4	18.2	53.3	166.0
2003	5.7	*	*	*	*	0.3	1.0	3.3	7.4	18.4	48.9	153.5
2002	5.4	*	*	*	*	0.2	1.0	2.8	6.6	17.1	47.3	153.4
2001	5.2	*	*	*	*	0.3	0.9	3.0	6.5	16.6	45.8	150.8
2000	5.0	*	*	*	*	0.2	1.0	3.0	7.0	15.8	45.2	142.4
1999	4.7	*	*	*	*	0.2	0.9	2.7	6.4	16.5	42.9	132.4
All origins 3/, female												
2006	9.5	*	*	*	*	0.2	0.7	2.2	5.5	15.3	50.8	203.5
2005	10.3	*	*	*	*	0.2	0.7	2.0	5.1	16.5	55.8	228.7
2004	9.5	*	*	*	*	0.2	0.6	2.0	5.3	16.1	52.1	213.2
2003	9.3	*	*	*	*	0.2	0.7	1.8	5.3	15.7	53.5	204.6
2002	8.6	*	*	*	*	0.2	0.6	1.8	4.9	15.1	48.8	192.1
2001	8.3	*	*	*	*	0.2	0.6	1.8	5.2	14.5	48.9	180.8
2000	7.8	*	*	*	*	0.2	0.6	1.7	4.9	14.6	45.7	171.3
1999	7.4	*	*	*	*	0.2	0.5	1.7	4.7	14.1	43.9	160.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential hypertension and hypertensive renal disease (I10,I12,I15)												
Hispanic, both sexes												
2006	2.8	*	*	*	*	*	0.5	2.3	7.1	16.9	40.1	147.1
2005	3.1	*	*	*	*	*	0.6	1.9	6.3	18.6	52.9	169.4
2004	2.7	*	*	*	*	*	0.4	2.2	5.4	16.1	44.3	172.2
2003	2.6	*	*	*	*	*	0.4	2.0	5.8	16.7	44.5	171.0
2002	2.4	*	*	*	*	*	0.5	1.1	6.2	15.5	47.2	148.6
2001	2.4	*	*	*	*	*	*	1.8	4.5	16.5	48.9	160.1
2000	2.3	*	*	*	*	*	0.5	1.8	6.4	15.0	45.8	148.0
1999	2.2	*	*	*	*	*	*	2.2	5.0	15.2	48.3	129.8
Hispanic, male												
2006	2.4	*	*	*	*	*	0.6	2.8	8.4	19.0	38.9	131.9
2005	2.5	*	*	*	*	*	0.8	2.1	6.1	20.5	54.1	140.7
2004	2.3	*	*	*	*	*	*	2.9	6.4	18.4	48.6	147.1
2003	2.1	*	*	*	*	*	*	2.8	7.4	15.5	44.5	133.9
2002	1.9	*	*	*	*	*	*	1.2	7.1	15.3	46.4	132.2
2001	1.9	*	*	*	*	*	*	2.3	3.9	16.8	51.0	129.5
2000	2.1	*	*	*	*	*	0.8	2.8	8.1	15.2	52.6	141.1
1999	1.9	*	*	*	*	*	*	2.4	5.2	18.5	51.6	131.6
Hispanic, female												
2006	3.3	*	*	*	*	*	*	1.9	6.0	15.3	40.9	155.2
2005	3.7	*	*	*	*	*	*	1.7	6.5	17.1	52.1	184.4
2004	3.1	*	*	*	*	*	*	1.5	4.6	14.3	41.2	185.1
2003	3.2	*	*	*	*	*	*	1.2	4.4	17.6	44.5	189.8
2002	3.0	*	*	*	*	*	*	*	5.5	15.7	47.8	156.7
2001	2.9	*	*	*	*	*	*	1.2	5.0	16.2	47.4	175.0
2000	2.5	*	*	*	*	*	*	*	4.9	14.9	41.3	151.3
1999	2.5	*	*	*	*	*	*	1.9	4.8	12.7	46.1	128.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential hypertension and hypertensive renal disease (I10,I12,I15)												
Non-Hispanic 4/, both sexes												
2006	8.8	*	*	*	*	0.3	1.0	3.1	6.8	16.7	51.6	191.2
2005	9.3	*	*	*	*	0.2	1.0	2.8	6.4	17.6	55.6	211.8
2004	8.7	*	*	*	*	0.3	0.9	2.8	6.4	17.1	52.9	199.4
2003	8.3	*	*	*	*	0.3	0.9	2.6	6.3	16.8	51.9	189.4
2002	7.7	*	*	*	*	0.2	0.8	2.4	5.7	15.9	48.1	181.1
2001	7.4	*	*	*	0.1	0.3	0.8	2.4	5.8	15.3	47.5	172.0
2000	7.0	*	*	*	*	0.2	0.8	2.4	5.8	15.1	45.3	162.9
1999	6.6	*	*	*	*	0.2	0.7	2.2	5.6	15.1	43.2	152.3
Non-Hispanic 4/, male												
2006	7.1	*	*	*	*	0.4	1.2	3.9	8.3	18.4	51.9	160.5
2005	7.2	*	*	*	*	0.3	1.2	3.7	8.0	19.0	55.3	170.7
2004	6.8	*	*	*	*	0.3	1.1	3.5	7.4	18.1	53.5	166.7
2003	6.3	*	*	*	*	0.3	1.0	3.3	7.3	18.5	49.1	154.4
2002	5.9	*	*	*	*	0.3	1.0	3.0	6.6	17.1	47.2	153.9
2001	5.7	*	*	*	*	0.3	1.0	3.0	6.6	16.5	45.3	151.1
2000	5.4	*	*	*	*	0.2	1.0	3.0	6.9	15.7	44.7	142.0
1999	5.1	*	*	*	*	0.2	0.9	2.7	6.5	16.4	42.4	131.9
Non-Hispanic 4/, female												
2006	10.5	*	*	*	*	0.2	0.7	2.2	5.5	15.3	51.4	205.5
2005	11.3	*	*	*	*	0.2	0.8	2.0	5.0	16.4	55.9	230.6
2004	10.5	*	*	*	*	0.3	0.6	2.0	5.4	16.2	52.6	214.0
2003	10.2	*	*	*	*	0.2	0.8	1.8	5.4	15.5	53.8	204.7
2002	9.4	*	*	*	*	0.2	0.6	1.9	4.8	15.0	48.7	192.8
2001	9.0	*	*	*	*	0.3	0.6	1.9	5.1	14.4	48.9	180.7
2000	8.5	*	*	*	*	0.2	0.6	1.8	4.9	14.6	45.8	171.3
1999	8.0	*	*	*	*	0.2	0.6	1.7	4.7	14.1	43.7	160.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential hypertension and hypertensive renal disease (I10,I12,I15)												
Non-Hispanic white, both sexes												
2006 5/	8.5	*	*	*	*	0.1	0.5	1.9	4.9	13.1	45.8	180.8
2005 5/	9.0	*	*	*	*	*	0.5	1.7	4.5	13.4	49.6	201.5
2004 5/	8.3	*	*	*	*	0.1	0.5	1.7	4.2	13.1	46.7	188.3
2003 5/	8.0	*	*	*	*	0.1	0.5	1.6	4.2	12.6	46.2	181.0
2002	7.4	*	*	*	*	0.1	0.4	1.5	3.9	12.5	42.2	170.6
2001	7.0	*	*	*	*	0.1	0.4	1.4	3.9	11.7	41.8	162.8
2000	6.5	*	*	*	*	0.1	0.4	1.3	3.8	11.6	39.9	152.7
1999	6.1	*	*	*	*	0.1	0.3	1.2	3.6	11.8	37.3	141.7
Non-Hispanic white, male												
2006 5/	6.5	*	*	*	*	0.2	0.6	2.5	6.1	14.2	45.9	152.0
2005 5/	6.6	*	*	*	*	*	0.7	2.3	5.7	14.3	48.6	160.8
2004 5/	6.3	*	*	*	*	0.2	0.7	2.2	5.2	14.2	46.9	159.6
2003 5/	5.8	*	*	*	*	*	0.7	2.1	4.9	14.4	43.6	145.4
2002	5.5	*	*	*	*	*	0.5	2.0	4.7	13.5	41.0	145.4
2001	5.1	*	*	*	*	0.2	0.5	1.7	4.5	12.8	39.6	142.1
2000	4.9	*	*	*	*	*	0.6	1.8	4.7	12.4	39.5	131.8
1999	4.6	*	*	*	*	0.2	0.5	1.6	4.5	13.0	36.5	119.8
Non-Hispanic white, female												
2006 5/	10.3	*	*	*	*	*	0.3	1.2	3.6	12.1	45.7	194.2
2005 5/	11.2	*	*	*	*	*	0.4	1.1	3.3	12.6	50.3	220.1
2004 5/	10.2	*	*	*	*	*	0.3	1.1	3.3	12.3	46.6	201.2
2003 5/	10.0	*	*	*	*	*	0.4	1.1	3.5	11.2	47.9	196.6
2002	9.2	*	*	*	*	*	0.3	1.1	3.1	11.5	42.9	181.4
2001	8.8	*	*	*	*	*	0.3	1.0	3.3	10.8	43.2	171.5
2000	8.1	*	*	*	*	*	0.3	0.9	3.1	10.9	40.1	161.1
1999	7.6	*	*	*	*	*	0.2	0.9	2.8	10.8	37.8	150.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential hypertension and hypertensive renal disease (I10,I12,I15)												
Non-Hispanic black, both sexes												
2006 5/	12.8	*	*	*	*	1.2	3.8	11.3	23.4	50.1	118.9	322.8
2005 5/	13.1	*	*	*	*	1.0	3.6	10.8	23.0	55.2	124.3	345.2
2004 5/	12.8	*	*	*	*	1.0	3.2	10.4	24.3	52.8	124.9	339.6
2003 5/	12.2	*	*	*	*	1.0	3.2	9.5	24.7	55.1	121.0	304.3
2002	11.5	*	*	*	*	0.9	3.4	9.2	21.4	48.3	120.0	312.4
2001	11.3	*	*	*	*	1.1	3.2	10.1	22.4	48.4	113.9	291.5
2000	11.0	*	*	*	*	0.7	3.2	9.9	23.3	49.5	108.9	285.9
1999	10.7	*	*	*	*	0.7	3.2	9.4	21.8	47.5	112.9	272.2
Non-Hispanic black, male												
2006 5/	11.7	*	*	*	*	1.4	4.9	14.2	28.7	60.6	129.0	274.9
2005 5/	11.8	*	*	*	*	1.1	4.0	13.6	29.8	65.0	139.0	294.0
2004 5/	11.1	*	*	*	*	1.2	4.0	13.2	28.8	57.6	139.2	268.3
2003 5/	10.8	*	*	*	*	1.3	3.5	13.0	30.0	61.3	120.9	280.2
2002	9.9	*	*	*	*	1.0	4.3	11.3	24.0	52.6	129.5	276.8
2001	10.0	*	*	*	*	1.2	3.9	13.1	25.8	55.5	119.7	277.2
2000	9.5	*	*	*	*	*	3.8	12.1	28.4	51.5	110.5	268.9
1999	9.4	*	*	*	*	*	3.8	12.0	24.8	52.3	119.1	268.6
Non-Hispanic black, female												
2006 5/	13.9	*	*	*	*	1.0	2.8	8.8	19.1	42.7	113.1	342.3
2005 5/	14.3	*	*	*	*	0.8	3.1	8.3	17.6	48.2	116.0	365.8
2004 5/	14.3	*	*	*	*	1.0	2.5	8.0	20.8	49.4	116.9	367.8
2003 5/	13.4	*	*	*	*	*	2.8	6.5	20.5	50.8	121.1	313.6
2002	13.0	*	*	*	*	0.7	2.6	7.4	19.4	45.3	114.6	326.1
2001	12.4	*	*	*	*	1.0	2.5	7.5	19.7	43.4	110.7	296.9
2000	12.4	*	*	*	*	*	2.6	8.0	19.3	48.1	108.0	292.2
1999	11.8	*	*	*	*	0.7	2.7	7.2	19.4	44.1	109.4	273.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
All origins 3/, both sexes												
2006	45.8	3.4	0.3	0.2	0.5	1.3	5.1	14.7	33.3	96.3	335.1	1,039.6
2005	48.4	3.1	0.4	0.2	0.5	1.4	5.2	15.0	33.0	101.1	359.0	1,141.8
2004	51.1	3.1	0.3	0.2	0.5	1.4	5.4	14.9	34.3	107.8	386.2	1,245.9
2003	54.2	2.5	0.3	0.2	0.5	1.5	5.5	15.0	35.6	112.9	410.7	1,370.1
2002	56.4	2.9	0.3	0.2	0.4	1.4	5.4	15.1	37.2	120.3	431.0	1,445.9
2001	57.4	2.7	0.4	0.2	0.5	1.5	5.5	15.1	38.0	123.4	443.9	1,500.2
2000	59.6	3.3	0.3	0.2	0.5	1.5	5.8	16.0	41.0	128.6	461.3	1,589.2
1999	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8
All origins 3/, male												
2006	37.0	3.9	0.3	0.3	0.5	1.4	5.3	16.4	38.7	108.0	345.5	932.4
2005	38.8	3.5	0.5	0.3	0.4	1.5	5.2	16.5	38.5	113.6	372.9	1,023.3
2004	40.7	3.4	0.3	0.2	0.5	1.4	5.6	16.7	39.5	121.1	402.9	1,118.1
2003	42.9	2.8	0.3	0.2	0.5	1.6	5.8	16.7	40.8	127.8	431.4	1,236.0
2002	44.2	3.2	0.4	0.2	0.5	1.4	5.3	16.7	42.7	135.0	445.9	1,317.9
2001	45.2	3.1	0.3	0.2	0.5	1.6	5.7	16.7	43.4	140.4	467.3	1,380.2
2000	46.9	3.8	*	0.2	0.5	1.5	5.8	17.5	47.2	145.0	490.8	1,484.3
1999	47.1	3.3	0.3	0.2	0.5	1.5	5.8	16.7	46.5	147.3	500.4	1,512.8
All origins 3/, female												
2006	54.4	2.9	0.4	0.2	0.5	1.2	4.8	13.0	28.2	86.5	328.0	1,089.8
2005	57.8	2.6	0.3	0.2	0.5	1.2	5.1	13.6	27.9	90.5	349.5	1,196.1
2004	61.2	2.8	*	0.2	0.5	1.4	5.1	13.1	29.5	96.6	374.9	1,303.4
2003	65.1	2.2	0.3	0.1	0.5	1.4	5.3	13.4	30.9	100.5	396.8	1,429.4
2002	68.2	2.5	0.3	0.2	0.3	1.4	5.5	13.6	32.1	108.1	421.2	1,501.5
2001	69.2	2.3	0.4	0.2	0.5	1.5	5.4	13.6	32.9	109.3	428.6	1,550.4
2000	71.8	2.7	0.4	0.2	0.5	1.5	5.7	14.5	35.3	115.1	442.1	1,632.0
1999	72.3	2.1	0.3	0.2	0.5	1.4	5.6	13.8	35.1	117.2	449.8	1,656.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
Hispanic, both sexes												
2006	15.8	3.4	*	*	0.7	1.2	4.6	14.4	34.2	87.3	262.1	687.0
2005	16.0	2.6	*	*	0.6	1.0	4.6	15.0	33.4	89.4	275.5	738.4
2004	16.4	2.2	*	*	0.5	1.2	4.9	14.6	34.9	97.0	293.1	813.0
2003	16.7	*	*	*	0.4	1.2	4.5	14.9	35.3	98.4	316.8	880.5
2002	16.6	2.4	*	0.3	0.4	1.3	4.4	15.3	35.9	102.9	319.9	898.0
2001	17.4	*	*	*	0.6	1.4	5.0	15.9	35.4	106.2	341.2	1,042.9
2000	17.5	*	*	*	0.5	1.7	5.2	15.2	39.8	113.2	344.1	1,082.9
1999	17.4	*	*	*	0.5	1.6	4.9	15.7	37.7	116.9	348.3	1,077.6
Hispanic, male												
2006	14.3	*	*	*	0.8	1.4	5.5	17.0	41.1	100.1	292.8	581.9
2005	14.4	*	*	*	0.7	1.2	5.2	17.8	40.3	106.2	294.0	692.4
2004	15.0	*	*	*	0.5	1.5	5.9	17.5	42.9	114.4	323.3	778.9
2003	14.9	*	*	*	*	1.3	5.2	18.1	43.5	113.9	337.1	837.4
2002	15.0	*	*	*	*	1.5	5.1	18.6	45.0	124.6	338.5	856.7
2001	15.7	*	*	*	0.7	1.6	5.3	18.7	43.5	127.2	386.3	1,005.6
2000	15.8	*	*	*	*	1.9	5.5	18.1	48.8	136.1	392.9	1,029.9
1999	16.1	*	*	*	0.6	1.8	5.7	19.5	46.1	140.8	395.2	1,136.6
Hispanic, female												
2006	17.5	*	*	*	0.6	0.9	3.7	11.8	27.8	76.9	240.6	742.9
2005	17.7	*	*	*	*	0.9	3.9	12.1	27.1	75.8	262.6	762.5
2004	17.9	*	*	*	*	0.9	3.9	11.8	27.7	83.0	272.2	830.4
2003	18.6	*	*	*	*	1.1	3.7	11.7	27.8	86.0	302.8	902.3
2002	18.4	*	*	*	*	1.0	3.7	12.0	27.6	85.6	307.2	918.5
2001	19.1	*	*	*	*	1.2	4.7	13.1	28.2	89.6	310.7	1,061.2
2000	19.4	*	*	*	*	1.4	4.9	12.4	31.9	95.2	311.3	1,108.9
1999	18.8	*	*	*	*	1.4	4.0	11.9	30.3	98.1	317.1	1,049.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
Non-Hispanic 4/, both sexes												
2006	50.9	3.5	0.4	0.2	0.4	1.3	5.2	14.6	33.1	96.8	339.1	1,056.5
2005	53.8	3.2	0.4	0.2	0.4	1.4	5.2	15.0	33.0	101.7	363.3	1,159.8
2004	56.7	3.3	0.3	0.2	0.5	1.5	5.4	14.8	34.1	108.4	390.8	1,263.8
2003	60.1	2.6	0.3	0.2	0.6	1.5	5.7	15.0	35.6	113.7	415.0	1,388.4
2002	62.4	3.0	0.3	0.2	0.4	1.5	5.5	15.0	37.2	121.3	435.5	1,465.0
2001	63.2	2.8	0.3	0.2	0.5	1.5	5.6	14.9	38.0	124.3	447.7	1,514.0
2000	65.4	3.6	0.3	0.2	0.5	1.5	5.8	15.9	40.9	129.2	465.0	1,604.5
1999	65.7	2.9	0.3	0.2	0.5	1.4	5.8	15.1	40.6	131.2	473.8	1,630.4
Non-Hispanic 4/, male												
2006	41.1	4.1	0.3	0.3	0.5	1.4	5.3	16.2	38.4	108.3	348.2	950.4
2005	43.0	3.6	0.5	0.3	0.4	1.6	5.2	16.3	38.3	113.9	377.0	1,039.8
2004	45.0	3.7	0.3	0.2	0.5	1.4	5.6	16.5	39.0	121.4	406.7	1,133.7
2003	47.6	2.8	*	0.2	0.6	1.6	5.9	16.5	40.4	128.4	435.7	1,252.9
2002	48.9	3.5	0.4	0.2	0.5	1.4	5.3	16.4	42.3	135.4	450.1	1,336.0
2001	49.7	3.3	0.3	0.2	0.5	1.5	5.7	16.4	43.2	140.9	469.8	1,392.9
2000	51.5	4.2	*	0.2	0.5	1.5	5.8	17.3	46.8	145.1	493.7	1,499.9
1999	51.5	3.7	0.3	0.2	0.5	1.4	5.8	16.4	46.3	147.3	503.6	1,524.8
Non-Hispanic 4/, female												
2006	60.3	2.8	0.4	0.2	0.4	1.2	5.0	13.1	28.2	87.0	332.8	1,105.8
2005	64.1	2.8	*	0.2	0.5	1.3	5.2	13.7	28.0	91.4	354.1	1,214.5
2004	67.8	3.0	*	0.2	0.5	1.5	5.2	13.2	29.5	97.4	380.0	1,321.9
2003	72.0	2.4	0.3	0.1	0.5	1.4	5.5	13.5	31.0	101.4	401.1	1,448.0
2002	75.3	2.5	*	0.2	0.3	1.5	5.8	13.7	32.4	109.5	425.8	1,520.7
2001	76.1	2.4	0.4	0.2	0.5	1.5	5.5	13.5	33.2	110.4	433.2	1,564.4
2000	78.7	2.8	0.4	0.2	0.5	1.5	5.8	14.6	35.5	116.1	446.4	1,646.8
1999	79.2	2.0	*	0.2	0.5	1.4	5.8	13.9	35.4	118.1	454.4	1,672.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
Non-Hispanic white, both sexes												
2006 5/	54.2	2.4	0.4	0.2	0.4	1.0	3.8	11.1	26.8	87.1	331.9	1,068.3
2005 5/	57.4	2.4	0.3	0.2	0.3	1.1	3.9	11.1	26.9	92.3	355.8	1,171.8
2004 5/	60.6	2.6	0.3	0.2	0.4	1.1	4.2	10.9	27.8	98.7	381.8	1,276.4
2003 5/	64.2	2.0	0.2	0.1	0.5	1.2	4.1	10.9	28.7	103.6	406.6	1,401.1
2002	67.0	2.2	0.3	0.2	0.4	1.2	4.1	11.1	30.8	110.8	427.5	1,479.2
2001	67.6	1.9	0.3	0.2	0.4	1.3	4.2	10.8	31.2	113.0	439.5	1,525.0
2000	70.0	2.6	0.3	0.2	0.5	1.2	4.3	12.0	34.2	119.3	455.7	1,616.8
1999	70.4	2.4	0.3	0.2	0.4	1.1	4.3	11.2	33.4	120.7	465.1	1,648.0
Non-Hispanic white, male												
2006 5/	42.6	2.6	*	0.3	0.4	1.1	3.9	12.1	30.3	96.5	340.5	960.2
2005 5/	44.8	2.8	0.4	0.2	0.3	1.2	4.0	12.1	30.7	102.4	368.2	1,050.5
2004 5/	47.0	2.9	*	0.2	0.5	1.1	4.4	12.1	31.1	110.0	396.9	1,145.3
2003 5/	49.7	1.9	*	0.2	0.5	1.3	4.2	12.1	32.1	116.9	426.0	1,264.2
2002	51.3	2.7	*	0.2	0.5	1.1	4.0	12.1	34.5	123.2	441.1	1,345.9
2001	52.0	2.2	*	0.2	0.4	1.3	4.3	11.9	35.1	128.0	460.5	1,399.0
2000	53.9	3.3	*	0.2	0.5	1.2	4.3	13.0	38.7	133.1	482.3	1,505.9
1999	53.9	3.1	*	0.2	0.4	1.1	4.4	12.0	38.0	135.0	492.8	1,535.1
Non-Hispanic white, female												
2006 5/	65.5	2.2	0.4	*	0.4	1.0	3.7	10.1	23.5	79.0	325.9	1,118.7
2005 5/	69.6	2.0	*	*	0.4	1.1	3.9	10.2	23.3	83.6	347.2	1,227.3
2004 5/	73.7	2.2	*	0.2	0.4	1.2	4.0	9.8	24.7	89.0	371.6	1,335.1
2003 5/	78.2	2.1	*	*	0.4	1.1	4.0	9.7	25.5	92.1	393.6	1,461.3
2002	82.1	*	*	0.2	0.3	1.2	4.3	10.1	27.2	100.2	418.4	1,536.7
2001	82.6	*	*	*	0.3	1.3	4.1	9.8	27.4	100.3	425.6	1,577.4
2000	85.5	1.9	*	0.2	0.4	1.2	4.3	10.9	29.9	107.6	438.3	1,661.6
1999	86.1	*	*	*	0.4	1.1	4.3	10.5	29.1	108.8	447.0	1,693.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
Non-Hispanic black, both sexes												
2006 5/	44.7	7.7	*	0.3	0.7	2.8	12.7	37.9	83.0	181.3	442.7	1,023.0
2005 5/	46.5	6.2	*	0.4	0.8	3.1	12.5	40.5	81.5	186.0	475.7	1,122.3
2004 5/	48.7	6.2	*	*	0.8	3.3	12.7	40.0	86.0	199.4	519.1	1,226.1
2003 5/	51.1	4.9	*	*	1.0	3.2	14.2	42.0	91.3	204.9	539.1	1,366.0
2002	51.7	5.6	*	0.4	0.6	3.2	13.3	41.6	89.7	218.7	558.8	1,398.0
2001	52.9	6.9	*	*	0.9	3.0	13.6	43.5	92.8	226.0	567.2	1,465.5
2000	54.3	8.0	*	*	0.7	3.0	14.9	44.3	95.2	226.3	593.8	1,545.8
1999	53.9	4.7	*	*	0.7	2.9	14.7	42.2	97.7	235.7	596.4	1,501.5
Non-Hispanic black, male												
2006 5/	40.8	10.2	*	*	0.6	2.9	13.3	44.6	108.6	222.7	479.4	894.3
2005 5/	41.8	6.8	*	*	0.8	3.5	12.8	45.9	106.6	228.5	510.4	998.2
2004 5/	43.1	7.1	*	*	1.0	3.1	13.3	46.0	110.2	240.1	558.7	1,077.1
2003 5/	44.9	*	*	*	0.8	3.5	14.9	48.1	114.1	241.8	597.6	1,200.7
2002	45.1	*	*	*	0.7	2.9	12.7	47.5	112.7	267.3	595.2	1,266.8
2001	46.2	8.3	*	*	0.7	3.1	13.5	49.9	113.9	273.5	620.5	1,359.4
2000	47.7	9.2	*	*	*	2.8	15.1	50.4	116.9	272.1	664.9	1,479.5
1999	47.5	*	*	*	*	2.9	15.0	49.6	119.5	279.1	656.7	1,455.9
Non-Hispanic black, female												
2006 5/	48.3	*	*	*	0.8	2.7	12.2	32.1	62.5	151.8	421.9	1,075.5
2005 5/	50.8	*	*	*	0.9	2.7	12.2	35.9	61.3	155.6	456.0	1,172.3
2004 5/	53.8	*	*	*	*	3.5	12.2	34.8	66.6	170.4	496.7	1,285.0
2003 5/	56.7	*	*	*	1.1	3.0	13.7	36.8	73.0	178.6	506.0	1,430.0
2002	57.7	*	*	*	*	3.5	13.8	36.6	71.3	184.2	538.2	1,448.1
2001	59.0	*	*	*	1.1	2.9	13.6	38.1	75.9	192.5	537.2	1,504.8
2000	60.3	*	*	*	0.8	3.2	14.8	39.2	77.8	194.1	553.8	1,570.4
1999	59.8	*	*	*	0.8	2.8	14.4	35.9	80.2	205.1	562.7	1,518.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
All origins 3/, both sexes												
2006	2.9	*	*	*	*	*	0.1	0.3	1.2	4.5	18.9	89.9
2005	4.0	*	*	*	*	*	0.1	0.5	1.4	5.4	25.9	132.9
2004	4.0	*	*	*	*	*	0.1	0.4	1.6	6.1	26.3	137.0
2003	4.5	*	*	*	*	*	0.1	0.5	1.7	6.6	29.8	154.6
2002	4.8	*	*	*	*	*	0.1	0.5	1.8	7.1	31.9	168.1
2001	4.9	*	*	*	*	*	0.1	0.5	1.9	7.3	33.1	179.1
2000	5.1	*	*	*	*	*	0.1	0.5	1.9	7.7	34.6	188.8
1999	5.4	*	*	*	*	*	0.1	0.5	2.0	8.0	35.9	202.0
All origins 3/, male												
2006	2.3	*	*	*	*	*	0.1	0.5	1.6	5.3	20.5	85.3
2005	3.1	*	*	*	*	*	0.1	0.6	1.8	6.6	29.0	121.7
2004	3.1	*	*	*	*	*	0.1	0.5	2.0	7.3	29.7	128.2
2003	3.4	*	*	*	*	*	0.2	0.7	2.2	7.8	33.6	138.6
2002	3.7	*	*	*	*	*	0.1	0.6	2.4	8.8	35.5	156.6
2001	3.7	*	*	*	*	*	0.1	0.6	2.4	9.0	35.9	167.2
2000	3.9	*	*	*	*	*	0.2	0.6	2.5	9.6	39.7	176.5
1999	4.0	*	*	*	*	*	0.1	0.7	2.7	9.6	40.0	189.0
All origins 3/, female												
2006	3.5	*	*	*	*	*	*	0.2	0.9	3.8	17.9	92.1
2005	4.9	*	*	*	*	*	*	0.3	1.1	4.4	23.9	138.1
2004	4.9	*	*	*	*	*	*	0.2	1.2	5.1	24.0	141.0
2003	5.5	*	*	*	*	*	*	0.3	1.2	5.6	27.3	161.7
2002	5.9	*	*	*	*	*	0.1	0.3	1.3	5.7	29.5	173.1
2001	6.2	*	*	*	*	*	*	0.3	1.4	5.9	31.2	184.1
2000	6.3	*	*	*	*	*	*	0.3	1.2	6.2	31.3	193.8
1999	6.7	*	*	*	*	*	*	0.3	1.4	6.7	33.2	207.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
Hispanic, both sexes												
2006	0.6	*	*	*	*	*	*	*	*	2.6	11.5	50.5
2005	0.9	*	*	*	*	*	*	*	0.9	3.9	17.7	79.6
2004	0.9	*	*	*	*	*	*	*	1.0	4.3	18.6	74.7
2003	0.9	*	*	*	*	*	*	*	1.1	4.6	18.9	86.0
2002	1.1	*	*	*	*	*	*	*	1.0	4.2	22.7	108.6
2001	1.1	*	*	*	*	*	*	*	*	4.0	23.1	130.7
2000	1.2	*	*	*	*	*	*	*	*	5.9	24.5	132.7
1999	1.2	*	*	*	*	*	*	*	1.5	5.2	27.1	129.1
Hispanic, male												
2006	0.5	*	*	*	*	*	*	*	*	3.4	12.3	45.1
2005	0.8	*	*	*	*	*	*	*	*	5.2	20.4	83.4
2004	0.8	*	*	*	*	*	*	*	*	5.1	22.9	88.3
2003	0.9	*	*	*	*	*	*	*	*	6.1	24.2	95.4
2002	0.9	*	*	*	*	*	*	*	*	5.2	26.7	102.1
2001	0.9	*	*	*	*	*	*	*	*	4.8	27.3	127.6
2000	1.0	*	*	*	*	*	*	*	*	8.2	32.0	104.8
1999	1.1	*	*	*	*	*	*	*	*	6.8	31.3	151.0
Hispanic, female												
2006	0.8	*	*	*	*	*	*	*	*	*	10.8	53.3
2005	1.1	*	*	*	*	*	*	*	*	2.7	15.8	77.6
2004	1.0	*	*	*	*	*	*	*	*	3.7	15.6	67.7
2003	1.0	*	*	*	*	*	*	*	*	3.4	15.3	81.3
2002	1.3	*	*	*	*	*	*	*	*	3.5	20.0	111.9
2001	1.4	*	*	*	*	*	*	*	*	3.5	20.2	132.2
2000	1.4	*	*	*	*	*	*	*	*	4.0	19.5	146.4
1999	1.4	*	*	*	*	*	*	*	*	4.0	24.3	118.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
Non-Hispanic 4/, both sexes												
2006	3.3	*	*	*	*	*	0.1	0.3	1.3	4.6	19.4	91.8
2005	4.5	*	*	*	*	*	0.1	0.5	1.5	5.5	26.4	135.3
2004	4.5	*	*	*	*	*	0.1	0.4	1.6	6.3	26.7	139.7
2003	5.0	*	*	*	*	*	0.1	0.5	1.7	6.7	30.2	157.1
2002	5.4	*	*	*	*	*	0.1	0.5	1.9	7.3	32.2	170.2
2001	5.5	*	*	*	*	*	0.1	0.5	1.9	7.5	33.4	180.5
2000	5.7	*	*	*	*	*	0.1	0.5	1.9	7.8	35.0	190.3
1999	5.9	*	*	*	*	*	0.1	0.5	2.1	8.1	36.1	203.9
Non-Hispanic 4/, male												
2006	2.6	*	*	*	*	*	*	0.5	1.7	5.5	20.9	87.5
2005	3.5	*	*	*	*	*	0.1	0.7	1.9	6.7	29.5	123.7
2004	3.5	*	*	*	*	*	0.1	0.5	2.0	7.5	30.0	130.1
2003	3.8	*	*	*	*	*	0.2	0.7	2.2	7.9	34.0	140.3
2002	4.1	*	*	*	*	*	0.1	0.6	2.5	9.0	35.8	158.7
2001	4.1	*	*	*	*	*	0.1	0.6	2.4	9.2	36.1	168.2
2000	4.3	*	*	*	*	*	0.2	0.7	2.6	9.7	39.9	178.7
1999	4.4	*	*	*	*	*	0.2	0.7	2.7	9.7	40.3	189.9
Non-Hispanic 4/, female												
2006	3.9	*	*	*	*	*	*	0.2	0.9	3.9	18.3	93.9
2005	5.5	*	*	*	*	*	*	0.3	1.1	4.6	24.3	140.6
2004	5.5	*	*	*	*	*	*	0.2	1.2	5.2	24.4	144.0
2003	6.2	*	*	*	*	*	*	0.3	1.2	5.8	27.7	164.5
2002	6.6	*	*	*	*	*	*	0.3	1.4	5.9	29.8	175.1
2001	6.8	*	*	*	*	*	*	0.3	1.5	6.0	31.6	185.5
2000	7.0	*	*	*	*	*	*	0.3	1.3	6.3	31.8	195.1
1999	7.4	*	*	*	*	*	*	0.3	1.5	6.9	33.4	209.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
Non-Hispanic white, both sexes												
2006 5/	3.8	*	*	*	*	*	*	0.3	1.2	4.6	19.6	95.2
2005 5/	5.2	*	*	*	*	*	0.1	0.4	1.5	5.4	27.1	141.3
2004 5/	5.3	*	*	*	*	*	0.1	0.4	1.6	6.2	27.6	145.0
2003 5/	5.8	*	*	*	*	*	0.1	0.5	1.6	6.6	30.8	162.2
2002	6.1	*	*	*	*	*	0.1	0.4	1.8	7.3	32.5	177.0
2001	6.3	*	*	*	*	*	0.1	0.5	1.8	7.5	34.1	187.4
2000	6.5	*	*	*	*	*	0.1	0.4	1.8	7.8	35.5	196.8
1999	6.7	*	*	*	*	*	0.1	0.5	2.0	8.2	36.7	209.9
Non-Hispanic white, male												
2006 5/	2.9	*	*	*	*	*	*	0.5	1.6	5.3	20.9	90.3
2005 5/	4.0	*	*	*	*	*	*	0.7	1.9	6.5	30.3	130.6
2004 5/	4.1	*	*	*	*	*	*	0.6	2.0	7.6	30.8	136.8
2003 5/	4.3	*	*	*	*	*	0.2	0.7	2.2	7.7	34.1	145.2
2002	4.6	*	*	*	*	*	0.1	0.5	2.3	9.0	35.9	165.7
2001	4.7	*	*	*	*	*	0.1	0.6	2.3	9.2	36.6	175.2
2000	4.9	*	*	*	*	*	0.2	0.6	2.6	9.8	40.6	183.6
1999	4.9	*	*	*	*	*	0.1	0.7	2.6	9.6	40.4	194.6
Non-Hispanic white, female												
2006 5/	4.6	*	*	*	*	*	*	0.2	0.9	3.9	18.7	97.6
2005 5/	6.4	*	*	*	*	*	*	0.2	1.1	4.5	24.9	146.2
2004 5/	6.4	*	*	*	*	*	*	0.2	1.2	5.1	25.4	148.7
2003 5/	7.2	*	*	*	*	*	*	0.3	1.1	5.8	28.5	169.6
2002	7.6	*	*	*	*	*	*	0.3	1.3	5.9	30.2	181.9
2001	7.9	*	*	*	*	*	*	0.3	1.3	6.1	32.5	192.6
2000	8.0	*	*	*	*	*	*	0.2	1.1	6.2	32.1	202.1
1999	8.5	*	*	*	*	*	*	0.3	1.4	6.9	34.3	215.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
Non-Hispanic black, both sexes												
2006 5/	1.9	*	*	*	*	*	*	0.6	2.1	6.6	20.9	72.7
2005 5/	2.2	*	*	*	*	*	*	0.8	2.1	7.6	24.0	99.7
2004 5/	2.2	*	*	*	*	*	*	0.5	2.2	7.5	24.3	103.6
2003 5/	2.7	*	*	*	*	*	*	0.8	2.7	9.1	30.8	122.9
2002	2.8	*	*	*	*	*	*	1.0	3.0	8.1	37.0	118.0
2001	2.7	*	*	*	*	*	*	0.7	3.4	8.9	30.7	126.1
2000	3.0	*	*	*	*	*	*	1.1	3.3	9.1	35.6	140.2
1999	3.1	*	*	*	*	*	*	0.7	3.3	9.3	35.0	155.8
Non-Hispanic black, male												
2006 5/	1.7	*	*	*	*	*	*	*	3.1	8.9	26.1	76.0
2005 5/	1.7	*	*	*	*	*	*	0.9	2.8	9.8	26.6	77.1
2004 5/	1.7	*	*	*	*	*	*	*	3.0	8.6	30.2	78.2
2003 5/	2.3	*	*	*	*	*	*	1.2	3.7	12.0	40.4	105.5
2002	2.4	*	*	*	*	*	*	1.4	4.3	10.2	44.8	109.4
2001	2.2	*	*	*	*	*	*	1.0	3.9	11.0	37.2	113.2
2000	2.5	*	*	*	*	*	*	1.5	4.1	11.1	40.4	150.3
1999	2.6	*	*	*	*	*	*	*	4.5	12.0	44.4	166.4
Non-Hispanic black, female												
2006 5/	2.0	*	*	*	*	*	*	*	1.4	4.9	18.0	71.4
2005 5/	2.7	*	*	*	*	*	*	*	1.5	6.0	22.6	108.8
2004 5/	2.7	*	*	*	*	*	*	*	1.7	6.7	20.9	113.6
2003 5/	3.0	*	*	*	*	*	*	*	1.9	7.1	25.4	129.6
2002	3.2	*	*	*	*	*	*	*	2.0	6.6	32.5	121.2
2001	3.2	*	*	*	*	*	*	*	3.0	7.4	27.1	130.9
2000	3.5	*	*	*	*	*	*	*	2.7	7.7	32.8	136.5
1999	3.5	*	*	*	*	*	*	*	2.3	7.4	29.8	151.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
All origins 3/, both sexes												
2006	7.5	*	*	*	0.2	0.4	1.0	2.7	6.8	21.7	58.0	128.5
2005	8.0	*	*	*	0.1	0.4	1.1	2.6	7.5	24.1	62.6	135.5
2004	8.1	*	*	*	0.1	0.4	1.0	2.7	7.2	25.1	64.3	139.3
2003	8.5	*	*	*	0.2	0.4	1.0	2.9	8.1	26.9	67.6	146.0
2002	8.6	*	*	*	0.2	0.4	1.0	2.7	8.6	29.1	68.7	148.4
2001	8.9	*	*	*	0.2	0.4	1.1	2.6	8.7	29.2	71.8	158.2
2000	9.3	*	*	*	0.2	0.4	1.0	2.8	9.2	32.2	74.6	163.2
1999	9.2	*	*	*	0.1	0.4	1.1	2.7	9.3	31.5	75.8	165.2
All origins 3/, male												
2006	8.0	*	*	*	0.2	0.5	1.5	3.8	9.6	29.0	74.5	155.0
2005	8.5	*	*	*	0.2	0.5	1.5	3.6	10.5	32.8	81.8	158.2
2004	8.5	*	*	*	0.2	0.6	1.4	3.8	9.9	33.4	82.8	166.5
2003	9.2	*	*	*	0.2	0.5	1.4	4.1	11.3	37.0	91.8	179.7
2002	9.4	*	*	*	0.2	0.5	1.3	3.7	11.9	39.8	92.3	187.4
2001	9.6	*	*	*	0.2	0.6	1.5	3.6	12.0	40.3	97.0	197.3
2000	10.2	*	*	*	0.2	0.5	1.2	3.8	13.1	45.4	102.9	214.9
1999	10.1	*	*	*	0.2	0.5	1.3	3.7	13.3	44.4	104.2	207.5
All origins 3/, female												
2006	7.0	*	*	*	0.1	0.2	0.6	1.6	4.2	15.5	46.7	116.1
2005	7.4	*	*	*	*	0.3	0.7	1.6	4.7	16.7	49.5	125.0
2004	7.6	*	*	*	*	0.3	0.7	1.6	4.7	18.1	51.8	127.0
2003	7.8	*	*	*	0.1	0.2	0.7	1.8	5.2	18.4	51.4	131.1
2002	7.9	*	*	*	0.1	0.3	0.7	1.7	5.5	20.1	53.0	131.5
2001	8.2	*	*	*	0.1	0.3	0.7	1.7	5.6	19.9	55.3	141.8
2000	8.3	*	*	*	0.1	0.3	0.8	1.8	5.6	21.4	56.2	142.2
1999	8.4	*	*	*	*	0.2	0.8	1.7	5.6	21.0	57.3	148.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
Hispanic, both sexes												
2006	2.0	*	*	*	*	0.4	0.8	1.7	4.2	12.8	31.9	71.6
2005	2.1	*	*	*	*	*	0.8	1.8	4.1	13.3	41.7	74.4
2004	2.0	*	*	*	*	0.3	0.7	2.0	5.2	13.9	34.5	77.1
2003	2.2	*	*	*	*	0.3	0.7	1.6	4.6	14.1	42.3	106.2
2002	2.2	*	*	*	*	0.3	0.8	1.7	5.9	14.8	41.0	92.6
2001	2.0	*	*	*	*	*	0.7	2.1	4.8	13.7	39.0	95.0
2000	2.3	*	*	*	*	0.4	0.7	1.8	5.8	20.2	42.5	102.8
1999	2.1	*	*	*	*	*	0.7	1.3	5.7	16.4	44.0	110.8
Hispanic, male												
2006	2.3	*	*	*	*	0.5	1.1	2.1	6.0	17.2	45.2	86.8
2005	2.5	*	*	*	*	*	1.0	2.6	6.5	18.8	57.5	105.9
2004	2.3	*	*	*	*	*	0.9	3.0	7.3	20.5	44.4	89.7
2003	2.5	*	*	*	*	*	0.9	2.2	7.4	19.0	58.7	138.5
2002	2.3	*	*	*	*	*	1.0	2.2	7.1	19.9	56.6	102.1
2001	2.2	*	*	*	*	*	0.9	2.7	6.3	20.0	53.2	108.6
2000	2.5	*	*	*	*	*	0.8	2.1	8.0	25.7	62.0	131.0
1999	2.4	*	*	*	*	*	0.9	1.7	7.7	22.2	62.0	163.9
Hispanic, female												
2006	1.8	*	*	*	*	*	*	1.3	2.6	9.2	22.6	63.5
2005	1.7	*	*	*	*	*	*	0.9	1.9	8.9	30.7	58.0
2004	1.8	*	*	*	*	*	*	1.0	3.3	8.7	27.7	70.6
2003	1.9	*	*	*	*	*	*	*	2.1	10.1	31.1	89.8
2002	2.0	*	*	*	*	*	*	1.1	4.9	10.8	30.4	87.9
2001	1.8	*	*	*	*	*	*	1.6	3.4	8.8	29.3	88.4
2000	2.1	*	*	*	*	*	*	1.5	3.9	15.8	29.4	89.0
1999	1.9	*	*	*	*	*	*	*	4.0	11.9	31.9	85.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
Non-Hispanic 4/, both sexes												
2006	8.4	*	*	*	0.2	0.4	1.0	2.8	7.0	22.3	59.5	131.2
2005	8.9	*	*	*	0.1	0.4	1.1	2.6	7.7	24.8	63.7	138.2
2004	9.0	*	*	*	0.1	0.5	1.1	2.8	7.4	25.9	65.8	141.8
2003	9.5	*	*	*	0.2	0.4	1.1	3.1	8.4	27.7	68.7	147.5
2002	9.6	*	*	*	0.2	0.4	1.0	2.8	8.8	30.0	69.8	150.3
2001	9.9	*	*	*	0.2	0.5	1.2	2.6	8.9	30.1	73.2	160.1
2000	10.2	*	*	*	0.2	0.4	1.0	2.8	9.4	32.9	75.8	165.0
1999	10.2	*	*	*	0.1	0.4	1.1	2.8	9.5	32.3	76.9	166.8
Non-Hispanic 4/, male												
2006	9.1	*	*	*	0.2	0.5	1.5	4.0	9.9	29.8	76.2	158.5
2005	9.5	*	*	*	0.2	0.5	1.6	3.7	10.8	33.7	83.0	160.6
2004	9.6	*	*	*	0.2	0.6	1.5	3.9	10.1	34.3	84.9	170.1
2003	10.4	*	*	*	0.2	0.5	1.5	4.3	11.6	38.2	93.4	181.5
2002	10.5	*	*	*	0.2	0.5	1.3	3.9	12.3	41.1	93.9	190.7
2001	10.7	*	*	*	0.2	0.6	1.6	3.6	12.3	41.5	98.8	200.5
2000	11.3	*	*	*	0.2	0.5	1.3	3.9	13.5	46.5	104.4	217.9
1999	11.2	*	*	*	0.2	0.6	1.4	3.8	13.7	45.5	105.8	209.0
Non-Hispanic 4/, female												
2006	7.9	*	*	*	*	0.2	0.6	1.6	4.4	16.0	48.2	118.5
2005	8.3	*	*	*	*	0.3	0.7	1.6	4.9	17.3	50.5	128.0
2004	8.5	*	*	*	*	0.3	0.7	1.7	4.9	18.8	53.1	129.2
2003	8.6	*	*	*	*	0.3	0.7	1.9	5.5	19.0	52.3	132.6
2002	8.8	*	*	*	*	0.3	0.7	1.7	5.5	20.7	53.9	132.8
2001	9.1	*	*	*	*	0.3	0.8	1.7	5.8	20.6	56.3	143.3
2000	9.2	*	*	*	0.1	0.2	0.8	1.8	5.7	21.7	57.1	143.6
1999	9.3	*	*	*	*	0.2	0.8	1.8	5.7	21.5	58.2	149.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
Non-Hispanic white, both sexes												
2006 5/	9.3	*	*	*	0.2	0.3	0.9	2.4	6.7	22.4	59.8	132.6
2005 5/	9.8	*	*	*	0.1	0.3	0.9	2.2	7.4	24.9	64.0	138.9
2004 5/	9.9	*	*	*	0.2	0.4	0.9	2.4	6.9	26.2	66.3	141.9
2003 5/	10.4	*	*	*	0.2	0.3	0.9	2.6	8.2	28.2	69.5	148.3
2002	10.6	*	*	*	0.1	0.3	0.9	2.3	8.4	30.4	70.9	150.9
2001	10.8	*	*	*	0.2	0.4	1.0	2.3	8.5	30.5	73.5	160.1
2000	11.2	*	*	*	0.1	0.3	0.8	2.4	9.0	33.5	76.3	164.8
1999	11.3	*	*	*	0.1	0.4	1.0	2.5	9.4	32.9	77.7	166.7
Non-Hispanic white, male												
2006 5/	10.0	*	*	*	0.2	0.5	1.3	3.5	9.5	30.1	77.8	163.3
2005 5/	10.4	*	*	*	0.2	0.4	1.2	3.1	10.4	34.2	83.7	164.5
2004 5/	10.5	*	*	*	0.2	0.5	1.3	3.5	9.4	35.0	86.4	173.0
2003 5/	11.5	*	*	*	0.2	0.4	1.2	3.8	11.3	39.2	94.7	185.8
2002	11.6	*	*	*	0.2	0.4	1.2	3.3	11.8	42.0	96.3	194.6
2001	11.8	*	*	*	0.2	0.6	1.3	3.1	12.0	42.4	100.3	204.7
2000	12.6	*	*	*	0.2	0.5	1.0	3.5	13.2	47.9	106.3	222.7
1999	12.5	*	*	*	0.2	0.5	1.3	3.4	13.8	47.0	107.9	212.8
Non-Hispanic white, female												
2006 5/	8.6	*	*	*	*	0.2	0.5	1.3	4.0	15.7	47.3	118.3
2005 5/	9.1	*	*	*	*	0.3	0.6	1.4	4.6	16.9	50.4	127.2
2004 5/	9.3	*	*	*	*	0.2	0.6	1.4	4.5	18.6	52.6	128.0
2003 5/	9.4	*	*	*	*	0.2	0.5	1.5	5.2	18.8	52.4	131.8
2002	9.6	*	*	*	*	0.3	0.6	1.4	5.2	20.5	53.9	132.1
2001	9.9	*	*	*	*	0.3	0.6	1.5	5.3	20.4	55.7	141.6
2000	9.9	*	*	*	*	0.2	0.6	1.4	5.2	21.3	56.6	141.5
1999	10.1	*	*	*	*	0.2	0.7	1.5	5.3	21.1	57.8	148.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
Non-Hispanic black, both sexes												
2006 5/	6.5	*	*	*	*	0.7	2.1	5.8	11.7	27.0	65.8	134.0
2005 5/	6.9	*	*	*	*	0.9	2.4	5.5	12.3	29.9	70.5	147.5
2004 5/	6.8	*	*	*	*	0.9	2.3	5.3	12.9	28.6	71.2	156.9
2003 5/	6.8	*	*	*	*	0.7	2.4	6.4	12.1	29.6	71.1	152.9
2002	6.9	*	*	*	*	0.9	2.0	6.0	13.6	31.7	69.1	157.6
2001	7.3	*	*	*	*	0.8	2.5	5.2	14.1	32.3	80.5	175.7
2000	7.3	*	*	*	*	0.7	2.2	5.7	14.5	33.4	78.9	180.0
1999	7.0	*	*	*	*	0.6	2.0	5.4	12.5	32.1	77.3	180.1
Non-Hispanic black, male												
2006 5/	6.5	*	*	*	*	1.0	3.2	8.0	16.6	35.4	69.0	119.1
2005 5/	7.1	*	*	*	*	1.1	3.7	8.0	16.9	37.0	89.1	132.3
2004 5/	6.9	*	*	*	*	1.2	3.1	7.5	18.3	36.1	81.0	161.6
2003 5/	6.9	*	*	*	*	1.2	3.1	8.4	16.7	36.4	91.5	141.7
2002	7.0	*	*	*	*	1.2	2.7	8.2	19.5	39.2	78.8	168.5
2001	7.2	*	*	*	*	1.0	3.5	7.4	17.9	40.3	97.2	168.6
2000	7.2	*	*	*	*	0.9	2.8	7.6	18.7	41.8	96.9	179.7
1999	6.7	*	*	*	*	0.9	2.4	7.4	16.0	40.0	90.7	184.2
Non-Hispanic black, female												
2006 5/	6.5	*	*	*	*	*	1.2	3.9	7.7	20.9	64.1	140.1
2005 5/	6.7	*	*	*	*	*	1.2	3.4	8.5	24.9	59.9	153.6
2004 5/	6.8	*	*	*	*	*	1.6	3.5	8.6	23.3	65.7	155.0
2003 5/	6.8	*	*	*	*	*	1.7	4.6	8.4	24.7	59.6	157.2
2002	6.9	*	*	*	*	*	1.4	4.2	8.9	26.3	63.7	153.4
2001	7.4	*	*	*	*	*	1.6	3.3	11.0	26.6	71.1	178.4
2000	7.5	*	*	*	*	*	1.6	4.0	11.2	27.5	68.8	180.1
1999	7.2	*	*	*	*	*	1.7	3.6	9.7	26.6	69.9	178.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
All origins 3/, both sexes												
2006	4.4	*	*	*	0.1	0.3	0.8	1.8	4.3	13.9	35.4	62.7
2005	4.7	*	*	*	0.1	0.3	0.8	1.8	4.8	15.4	38.1	64.9
2004	4.7	*	*	*	0.1	0.3	0.7	1.8	4.5	15.9	39.6	64.8
2003	5.1	*	*	*	0.1	0.3	0.7	2.0	5.3	17.8	42.5	70.0
2002	5.1	*	*	*	0.1	0.3	0.7	1.8	5.5	19.1	43.3	69.4
2001	5.3	*	*	*	0.1	0.3	0.7	1.7	5.5	19.2	45.7	77.5
2000	5.6	*	*	*	0.1	0.3	0.7	1.8	6.2	21.8	47.2	79.2
1999	5.7	*	*	*	0.1	0.2	0.7	1.7	6.0	21.7	48.6	80.3
All origins 3/, male												
2006	5.2	*	*	*	0.1	0.4	1.2	2.8	6.6	19.5	48.4	87.6
2005	5.6	*	*	*	0.1	0.4	1.2	2.8	7.3	22.4	52.6	91.0
2004	5.6	*	*	*	0.2	0.5	1.1	2.7	6.9	23.1	54.8	95.0
2003	6.3	*	*	*	0.1	0.4	1.1	3.1	8.2	26.0	61.9	105.5
2002	6.4	*	*	*	0.2	0.4	1.0	2.7	8.5	28.5	62.7	106.6
2001	6.5	*	*	*	0.2	0.4	1.2	2.6	8.5	28.7	66.4	114.8
2000	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.9	70.6	127.3
1999	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.7	72.7	125.3
All origins 3/, female												
2006	3.6	*	*	*	*	0.1	0.3	0.9	2.2	9.3	26.5	51.0
2005	3.8	*	*	*	*	0.2	0.3	0.8	2.5	9.5	28.2	52.9
2004	3.8	*	*	*	*	0.2	0.3	0.8	2.3	9.8	29.3	51.2
2003	3.9	*	*	*	*	0.1	0.3	1.0	2.6	11.0	29.5	54.2
2002	4.0	*	*	*	*	0.1	0.3	0.8	2.7	11.2	30.5	53.3
2001	4.2	*	*	*	*	0.1	0.3	0.8	2.8	11.3	32.1	61.9
2000	4.3	*	*	*	*	0.1	0.4	0.9	3.0	12.7	32.0	59.6
1999	4.4	*	*	*	*	*	0.4	0.8	2.8	12.7	32.9	62.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
Hispanic, both sexes												
2006	1.1	*	*	*	*	*	0.6	1.0	2.4	7.1	16.3	31.7
2005	1.1	*	*	*	*	*	0.5	1.1	2.6	7.3	19.9	29.1
2004	1.1	*	*	*	*	*	0.4	1.3	3.3	7.9	18.0	30.4
2003	1.3	*	*	*	*	*	0.4	0.9	3.1	8.3	26.7	49.2
2002	1.1	*	*	*	*	*	0.4	1.0	3.8	7.5	21.2	31.0
2001	1.1	*	*	*	*	*	0.4	1.2	2.5	8.2	22.9	31.9
2000	1.3	*	*	*	*	*	0.4	1.3	3.4	11.2	27.1	41.8
1999	1.1	*	*	*	*	*	0.5	0.9	3.4	9.2	24.0	40.7
Hispanic, male												
2006	1.4	*	*	*	*	*	1.0	1.5	3.9	10.9	24.0	47.4
2005	1.5	*	*	*	*	*	0.8	1.9	4.8	10.7	30.7	43.6
2004	1.5	*	*	*	*	*	0.7	2.2	5.5	12.6	25.4	40.6
2003	1.6	*	*	*	*	*	*	1.4	5.1	12.5	38.3	80.0
2002	1.4	*	*	*	*	*	0.7	1.6	5.5	12.0	32.0	43.5
2001	1.5	*	*	*	*	*	0.7	1.8	3.8	14.2	34.9	53.3
2000	1.6	*	*	*	*	*	*	1.8	5.2	16.0	42.3	64.5
1999	1.5	*	*	*	*	*	*	1.5	5.3	14.7	33.9	79.8
Hispanic, female												
2006	0.7	*	*	*	*	*	*	*	*	4.1	10.8	23.4
2005	0.7	*	*	*	*	*	*	*	*	4.5	12.3	21.5
2004	0.8	*	*	*	*	*	*	*	*	4.1	12.8	25.2
2003	0.9	*	*	*	*	*	*	*	*	5.0	18.7	33.6
2002	0.8	*	*	*	*	*	*	*	2.3	4.0	13.8	24.9
2001	0.7	*	*	*	*	*	*	*	*	3.5	14.8	21.4
2000	1.0	*	*	*	*	*	*	*	*	7.5	16.8	30.7
1999	0.8	*	*	*	*	*	*	*	*	4.8	17.3	21.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
Non-Hispanic 4/, both sexes												
2006	5.0	*	*	*	0.1	0.3	0.8	1.9	4.5	14.4	36.6	64.2
2005	5.3	*	*	*	0.1	0.3	0.8	1.9	5.0	16.0	39.1	66.5
2004	5.3	*	*	*	0.1	0.3	0.8	1.8	4.6	16.4	40.7	66.2
2003	5.7	*	*	*	0.1	0.3	0.8	2.1	5.5	18.5	43.2	70.7
2002	5.7	*	*	*	0.1	0.3	0.7	1.8	5.6	19.8	44.3	70.8
2001	6.0	*	*	*	0.1	0.3	0.8	1.7	5.7	19.8	46.7	79.0
2000	6.2	*	*	*	0.1	0.3	0.7	1.8	6.3	22.4	47.9	80.3
1999	6.3	*	*	*	0.1	0.3	0.8	1.8	6.2	22.4	49.5	81.7
Non-Hispanic 4/, male												
2006	5.9	*	*	*	0.2	0.4	1.3	3.0	6.8	20.1	49.9	89.8
2005	6.3	*	*	*	0.1	0.4	1.3	2.9	7.4	23.2	53.8	93.3
2004	6.3	*	*	*	0.2	0.5	1.2	2.7	7.0	23.8	56.4	97.5
2003	7.0	*	*	*	0.2	0.4	1.2	3.2	8.5	26.9	63.0	106.6
2002	7.2	*	*	*	0.2	0.4	1.1	2.9	8.7	29.6	64.1	109.1
2001	7.3	*	*	*	0.2	0.5	1.2	2.7	8.8	29.5	67.7	117.0
2000	7.8	*	*	*	*	0.4	1.0	2.7	9.8	33.8	71.6	129.6
1999	7.8	*	*	*	*	0.4	1.1	2.8	9.8	33.6	74.3	127.0
Non-Hispanic 4/, female												
2006	4.1	*	*	*	*	*	0.4	0.9	2.3	9.6	27.5	52.3
2005	4.3	*	*	*	*	0.2	0.3	0.9	2.7	9.9	29.1	54.2
2004	4.2	*	*	*	*	0.2	0.4	0.9	2.4	10.2	30.2	52.3
2003	4.4	*	*	*	*	0.2	0.3	1.0	2.7	11.4	30.0	55.0
2002	4.4	*	*	*	*	0.1	0.4	0.9	2.7	11.6	31.2	54.3
2001	4.7	*	*	*	*	0.2	0.3	0.8	2.9	11.8	32.9	63.2
2000	4.7	*	*	*	*	0.1	0.4	0.9	3.1	13.0	32.5	60.4
1999	4.8	*	*	*	*	*	0.4	0.9	2.9	13.2	33.4	63.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
Non-Hispanic white, both sexes												
2006 5/	5.5	*	*	*	0.1	0.2	0.7	1.6	4.3	14.7	37.7	66.0
2005 5/	5.8	*	*	*	0.1	0.3	0.6	1.5	4.9	16.5	40.2	67.9
2004 5/	5.9	*	*	*	0.1	0.3	0.6	1.6	4.3	17.2	41.9	68.4
2003 5/	6.4	*	*	*	0.1	0.2	0.6	1.8	5.4	19.3	44.8	73.2
2002	6.5	*	*	*	0.1	0.2	0.6	1.5	5.5	20.7	46.0	72.8
2001	6.7	*	*	*	0.1	0.3	0.6	1.5	5.6	20.7	47.9	81.3
2000	7.0	*	*	*	*	0.2	0.6	1.6	6.3	23.4	49.3	82.0
1999	7.1	*	*	*	0.1	0.2	0.7	1.6	6.3	23.4	51.1	83.7
Non-Hispanic white, male												
2006 5/	6.6	*	*	*	0.2	0.4	1.1	2.6	6.7	20.6	52.0	92.8
2005 5/	7.0	*	*	*	*	0.4	1.0	2.4	7.4	24.2	55.6	96.7
2004 5/	7.0	*	*	*	0.2	0.4	1.0	2.4	6.6	24.8	58.3	101.4
2003 5/	7.9	*	*	*	0.2	0.3	1.0	2.8	8.4	28.3	65.5	111.2
2002	8.0	*	*	*	*	0.3	0.9	2.4	8.6	31.0	67.1	114.7
2001	8.2	*	*	*	0.2	0.5	1.0	2.3	8.7	30.9	70.0	122.9
2000	8.8	*	*	*	*	0.3	0.8	2.5	9.8	35.6	74.2	135.3
1999	8.9	*	*	*	*	0.4	1.0	2.4	10.1	35.4	77.5	131.8
Non-Hispanic white, female												
2006 5/	4.5	*	*	*	*	*	0.3	0.7	2.1	9.6	27.8	53.4
2005 5/	4.7	*	*	*	*	0.2	0.3	0.7	2.5	9.9	29.7	54.8
2004 5/	4.7	*	*	*	*	*	0.3	0.7	2.2	10.6	30.7	53.6
2003 5/	4.9	*	*	*	*	*	0.2	0.8	2.5	11.6	30.9	56.5
2002	4.9	*	*	*	*	*	0.3	0.7	2.6	11.9	31.9	54.7
2001	5.2	*	*	*	*	*	0.3	0.6	2.7	12.0	33.3	64.0
2000	5.2	*	*	*	*	*	0.3	0.7	2.9	13.1	32.9	60.5
1999	5.3	*	*	*	*	*	0.4	0.7	2.7	13.3	33.9	64.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
Non-Hispanic black, both sexes												
2006 5/	3.3	*	*	*	*	0.5	1.5	4.1	6.7	14.6	28.3	47.8
2005 5/	3.4	*	*	*	*	0.5	1.8	4.1	7.0	14.8	31.1	51.3
2004 5/	3.3	*	*	*	*	0.6	1.7	3.6	7.6	13.3	32.4	44.6
2003 5/	3.3	*	*	*	*	0.6	1.7	4.4	7.3	14.5	30.4	47.3
2002	3.3	*	*	*	*	0.6	1.4	4.3	7.5	14.8	30.2	50.5
2001	3.5	*	*	*	*	0.4	1.6	3.6	7.7	14.7	38.0	55.9
2000	3.6	*	*	*	*	0.5	1.5	3.3	8.2	17.3	36.8	63.0
1999	3.4	*	*	*	*	0.4	1.5	3.9	6.5	16.8	34.0	61.1
Non-Hispanic black, male												
2006 5/	3.7	*	*	*	*	0.9	2.4	5.9	9.8	19.3	30.3	52.2
2005 5/	4.0	*	*	*	*	0.8	3.2	6.2	10.1	18.4	39.2	52.3
2004 5/	3.9	*	*	*	*	1.0	2.6	5.5	11.8	18.5	37.4	57.9
2003 5/	3.9	*	*	*	*	1.1	2.5	6.5	11.0	17.9	39.8	52.2
2002	3.9	*	*	*	*	1.0	2.3	6.6	12.1	19.3	34.9	49.2
2001	3.9	*	*	*	*	*	2.6	5.4	11.3	19.9	47.4	49.7
2000	4.0	*	*	*	*	*	2.3	4.7	11.6	22.6	47.2	70.5
1999	3.7	*	*	*	*	*	2.0	6.0	9.3	22.1	38.8	77.3
Non-Hispanic black, female												
2006 5/	2.9	*	*	*	*	*	0.8	2.4	4.3	11.2	27.1	46.0
2005 5/	3.0	*	*	*	*	*	*	2.3	4.4	12.2	26.5	50.9
2004 5/	2.7	*	*	*	*	*	0.9	2.0	4.3	9.5	29.6	39.4
2003 5/	2.8	*	*	*	*	*	1.0	2.5	4.3	12.1	25.1	45.4
2002	2.8	*	*	*	*	*	*	2.2	3.8	11.6	27.5	51.0
2001	3.1	*	*	*	*	*	0.7	2.0	4.8	11.1	32.7	58.2
2000	3.3	*	*	*	*	*	0.9	2.1	5.5	13.5	30.9	60.2
1999	3.1	*	*	*	*	*	1.0	2.0	4.3	13.1	31.3	55.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
All origins 3/, both sexes												
2006	3.1	*	*	*	0.1	0.1	0.2	0.8	2.5	7.8	22.6	65.8
2005	3.3	*	*	*	*	0.1	0.3	0.8	2.7	8.7	24.5	70.6
2004	3.4	*	*	*	*	0.1	0.3	1.0	2.7	9.2	24.7	74.5
2003	3.4	*	*	*	0.1	0.1	0.3	0.9	2.8	9.1	25.0	76.0
2002	3.5	*	*	*	0.1	0.1	0.3	0.9	3.1	10.0	25.3	79.0
2001	3.5	*	*	*	*	0.1	0.4	0.9	3.2	10.0	26.1	80.7
2000	3.6	*	*	*	0.1	0.1	0.3	1.0	3.1	10.4	27.4	84.0
1999	3.6	*	*	*	*	0.1	0.3	0.9	3.3	9.8	27.2	84.9
All origins 3/, male												
2006	2.8	*	*	*	*	*	0.2	1.0	3.0	9.5	26.1	67.3
2005	2.9	*	*	*	*	*	0.3	0.8	3.2	10.4	29.2	67.2
2004	2.9	*	*	*	*	0.1	0.3	1.1	3.0	10.3	28.0	71.6
2003	3.0	*	*	*	*	0.1	0.3	1.0	3.1	11.0	29.9	74.2
2002	3.0	*	*	*	*	*	0.3	1.0	3.4	11.3	29.6	80.8
2001	3.1	*	*	*	*	0.1	0.3	0.9	3.5	11.6	30.6	82.5
2000	3.2	*	*	*	*	0.1	0.3	1.1	3.6	12.5	32.3	87.6
1999	3.1	*	*	*	*	0.1	0.3	1.0	3.8	11.7	31.5	82.2
All origins 3/, female												
2006	3.4	*	*	*	*	0.1	0.2	0.7	2.0	6.3	20.2	65.0
2005	3.7	*	*	*	*	0.1	0.4	0.7	2.2	7.2	21.3	72.2
2004	3.9	*	*	*	*	0.1	0.4	0.8	2.4	8.3	22.5	75.8
2003	3.8	*	*	*	*	*	0.3	0.9	2.6	7.5	21.8	76.9
2002	4.0	*	*	*	*	0.2	0.3	0.8	2.8	8.9	22.5	78.2
2001	4.0	*	*	*	*	0.1	0.4	0.9	2.9	8.6	23.1	79.9
2000	4.0	*	*	*	*	0.1	0.4	0.9	2.6	8.7	24.2	82.6
1999	4.1	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.5	86.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Hispanic, both sexes												
2006	0.9	*	*	*	*	*	*	0.7	1.8	5.7	15.6	39.9
2005	1.0	*	*	*	*	*	*	0.6	1.4	6.1	21.8	45.3
2004	0.9	*	*	*	*	*	*	0.7	1.8	6.1	16.5	46.6
2003	0.9	*	*	*	*	*	*	0.7	1.5	5.7	15.6	57.0
2002	1.1	*	*	*	*	*	0.3	0.7	2.1	7.3	19.8	61.5
2001	0.9	*	*	*	*	*	*	0.9	2.2	5.5	16.0	63.2
2000	1.0	*	*	*	*	*	*	*	2.4	8.9	15.4	61.0
1999	1.0	*	*	*	*	*	*	*	2.4	7.2	20.0	70.2
Hispanic, male												
2006	0.9	*	*	*	*	*	*	*	2.1	6.3	21.2	39.5
2005	1.0	*	*	*	*	*	*	*	1.8	8.1	26.7	62.3
2004	0.8	*	*	*	*	*	*	*	*	8.0	19.0	49.0
2003	0.9	*	*	*	*	*	*	*	2.2	6.5	20.3	58.5
2002	0.9	*	*	*	*	*	*	*	*	7.8	24.6	58.6
2001	0.8	*	*	*	*	*	*	*	2.5	5.8	18.3	55.2
2000	0.9	*	*	*	*	*	*	*	2.7	9.7	19.7	66.5
1999	0.9	*	*	*	*	*	*	*	*	7.5	28.2	84.1
Hispanic, female												
2006	1.0	*	*	*	*	*	*	*	*	5.1	11.7	40.1
2005	1.0	*	*	*	*	*	*	*	*	4.4	18.4	36.5
2004	1.0	*	*	*	*	*	*	*	1.9	4.5	14.8	45.4
2003	1.0	*	*	*	*	*	*	*	*	5.1	12.4	56.3
2002	1.3	*	*	*	*	*	*	*	2.6	6.8	16.6	63.0
2001	1.1	*	*	*	*	*	*	*	*	5.4	14.5	67.0
2000	1.1	*	*	*	*	*	*	*	*	8.3	12.5	58.4
1999	1.1	*	*	*	*	*	*	*	2.4	7.1	14.6	63.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Non-Hispanic 4/, both sexes												
2006	3.5	*	*	*	*	0.1	0.2	0.9	2.6	7.9	23.0	67.0
2005	3.7	*	*	*	*	0.1	0.3	0.8	2.8	8.8	24.6	71.7
2004	3.8	*	*	*	*	0.1	0.3	1.0	2.8	9.4	25.1	75.6
2003	3.8	*	*	*	0.1	0.1	0.3	1.0	2.9	9.3	25.5	76.7
2002	3.9	*	*	*	0.1	0.1	0.3	0.9	3.2	10.2	25.6	79.5
2001	3.9	*	*	*	*	0.1	0.4	0.9	3.2	10.3	26.5	81.1
2000	4.0	*	*	*	0.1	0.1	0.3	1.0	3.1	10.5	27.9	84.7
1999	3.9	*	*	*	*	0.1	0.3	1.0	3.3	9.9	27.5	85.1
Non-Hispanic 4/, male												
2006	3.1	*	*	*	*	*	0.2	1.0	3.1	9.7	26.3	68.7
2005	3.2	*	*	*	*	*	0.3	0.8	3.3	10.5	29.2	67.3
2004	3.2	*	*	*	*	*	0.3	1.1	3.1	10.5	28.5	72.6
2003	3.3	*	*	*	*	*	0.3	1.0	3.1	11.3	30.3	74.9
2002	3.4	*	*	*	*	*	0.3	1.0	3.5	11.5	29.8	81.7
2001	3.4	*	*	*	*	0.1	0.3	0.9	3.5	12.0	31.1	83.5
2000	3.5	*	*	*	*	0.1	0.2	1.2	3.6	12.7	32.8	88.2
1999	3.4	*	*	*	*	0.1	0.3	1.0	3.9	11.9	31.5	82.0
Non-Hispanic 4/, female												
2006	3.8	*	*	*	*	*	0.2	0.7	2.0	6.4	20.7	66.2
2005	4.1	*	*	*	*	0.1	0.4	0.7	2.3	7.4	21.4	73.7
2004	4.3	*	*	*	*	*	0.4	0.8	2.5	8.5	22.9	76.9
2003	4.2	*	*	*	*	*	0.4	0.9	2.8	7.6	22.3	77.6
2002	4.3	*	*	*	*	0.2	0.3	0.8	2.8	9.0	22.8	78.5
2001	4.4	*	*	*	*	0.1	0.4	0.9	2.9	8.8	23.5	80.1
2000	4.4	*	*	*	*	*	0.4	0.9	2.6	8.7	24.6	83.3
1999	4.5	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.8	86.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Non-Hispanic white, both sexes												
2006 5/	3.7	*	*	*	*	*	0.2	0.8	2.3	7.7	22.1	66.6
2005 5/	3.9	*	*	*	*	0.1	0.3	0.7	2.6	8.4	23.7	70.9
2004 5/	4.0	*	*	*	*	0.1	0.3	0.9	2.5	9.0	24.4	73.6
2003 5/	4.0	*	*	*	*	0.1	0.3	0.8	2.8	8.9	24.6	75.1
2002	4.1	*	*	*	*	0.1	0.3	0.8	2.9	9.7	24.9	78.2
2001	4.2	*	*	*	*	0.1	0.3	0.8	2.9	9.8	25.6	78.8
2000	4.3	*	*	*	*	0.1	0.3	0.9	2.8	10.1	27.1	82.8
1999	4.2	*	*	*	*	0.1	0.3	0.9	3.1	9.6	26.5	83.0
Non-Hispanic white, male												
2006 5/	3.4	*	*	*	*	*	0.2	0.9	2.8	9.5	25.8	70.5
2005 5/	3.4	*	*	*	*	*	0.2	0.7	3.1	10.0	28.1	67.8
2004 5/	3.5	*	*	*	*	*	0.3	1.1	2.8	10.2	28.1	71.6
2003 5/	3.5	*	*	*	*	*	0.2	0.9	2.9	10.8	29.2	74.6
2002	3.6	*	*	*	*	*	0.2	0.9	3.2	11.0	29.3	79.9
2001	3.6	*	*	*	*	*	0.2	0.8	3.3	11.5	30.3	81.8
2000	3.8	*	*	*	*	*	0.2	1.0	3.3	12.3	32.1	87.4
1999	3.6	*	*	*	*	*	0.3	1.0	3.6	11.6	30.4	81.0
Non-Hispanic white, female												
2006 5/	4.0	*	*	*	*	*	0.2	0.6	1.9	6.1	19.5	64.9
2005 5/	4.4	*	*	*	*	*	0.4	0.7	2.1	7.0	20.8	72.4
2004 5/	4.6	*	*	*	*	*	0.3	0.7	2.3	8.0	21.9	74.4
2003 5/	4.5	*	*	*	*	*	0.3	0.7	2.7	7.3	21.6	75.4
2002	4.7	*	*	*	*	*	0.3	0.7	2.6	8.6	21.9	77.4
2001	4.7	*	*	*	*	*	0.4	0.9	2.6	8.3	22.4	77.6
2000	4.7	*	*	*	*	*	0.3	0.7	2.2	8.2	23.8	81.0
1999	4.8	*	*	*	*	*	0.4	0.8	2.5	7.9	24.0	83.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Non-Hispanic black, both sexes												
2006 5/	3.2	*	*	*	*	*	0.6	1.7	5.0	12.4	37.5	86.3
2005 5/	3.4	*	*	*	*	*	0.6	1.4	5.3	15.1	39.3	96.1
2004 5/	3.5	*	*	*	*	*	0.6	1.7	5.3	15.3	38.8	112.2
2003 5/	3.5	*	*	*	*	*	0.7	2.0	4.8	15.0	40.7	105.6
2002	3.6	*	*	*	*	*	0.6	1.8	6.2	16.8	39.0	107.1
2001	3.8	*	*	*	*	*	0.9	1.7	6.4	17.5	42.5	119.9
2000	3.7	*	*	*	*	*	0.6	2.4	6.3	16.1	42.1	117.0
1999	3.6	*	*	*	*	*	0.6	1.5	5.9	15.3	43.3	119.0
Non-Hispanic black, male												
2006 5/	2.8	*	*	*	*	*	*	2.1	6.8	16.0	38.7	66.9
2005 5/	3.1	*	*	*	*	*	*	1.9	6.8	18.6	49.9	79.9
2004 5/	3.0	*	*	*	*	*	*	2.0	6.5	17.5	43.6	103.6
2003 5/	3.0	*	*	*	*	*	*	1.9	5.7	18.5	51.7	89.5
2002	3.1	*	*	*	*	*	*	1.6	7.4	19.9	44.0	119.2
2001	3.3	*	*	*	*	*	0.9	2.0	6.6	20.5	49.8	119.0
2000	3.2	*	*	*	*	*	*	2.9	7.1	19.2	49.7	109.2
1999	3.0	*	*	*	*	*	*	1.4	6.7	17.9	51.9	107.0
Non-Hispanic black, female												
2006 5/	3.6	*	*	*	*	*	*	1.4	3.5	9.8	36.9	94.2
2005 5/	3.7	*	*	*	*	*	*	1.1	4.1	12.7	33.4	102.6
2004 5/	4.1	*	*	*	*	*	0.8	1.5	4.4	13.7	36.1	115.7
2003 5/	4.0	*	*	*	*	*	0.8	2.1	4.1	12.5	34.5	111.8
2002	4.0	*	*	*	*	*	0.8	1.9	5.1	14.6	36.1	102.4
2001	4.4	*	*	*	*	*	0.9	1.4	6.3	15.5	38.4	120.2
2000	4.2	*	*	*	*	*	0.7	1.9	5.7	14.0	37.9	119.9
1999	4.2	*	*	*	*	*	0.8	1.6	5.4	13.5	38.5	123.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
All origins 3/, both sexes												
2006	1.3	0.8	*	*	0.1	0.3	0.6	1.2	1.7	2.9	7.0	18.4
2005	1.6	0.6	*	*	0.1	0.3	0.8	1.5	2.1	3.8	8.6	23.0
2004	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.1	3.6	9.0	23.3
2003	1.6	0.9	*	*	0.1	0.3	0.8	1.4	2.0	4.1	9.0	24.0
2002	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.2	3.8	9.8	24.4
2001	1.6	0.5	*	*	0.1	0.3	0.6	1.2	2.2	4.6	9.4	26.1
2000	1.6	0.7	*	*	0.1	0.3	0.7	1.4	2.1	4.3	9.4	25.6
1999	1.4	0.6	*	*	0.1	0.2	0.6	1.0	1.9	3.8	9.1	23.4
All origins 3/, male												
2006	1.2	*	*	*	*	0.3	0.7	1.5	2.1	2.9	7.0	16.2
2005	1.5	*	*	*	*	0.3	0.7	1.8	2.6	4.4	8.4	19.2
2004	1.4	*	*	*	*	0.4	0.7	1.6	2.5	3.9	9.7	19.6
2003	1.4	1.0	*	*	0.1	0.3	0.8	1.6	2.4	4.2	8.5	19.1
2002	1.4	*	*	*	0.1	0.3	0.7	1.4	2.4	3.9	9.5	20.5
2001	1.4	*	*	*	*	0.3	0.7	1.4	2.5	4.9	9.8	22.0
2000	1.4	*	*	*	*	0.3	0.8	1.6	2.3	4.7	9.7	22.2
1999	1.3	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.6	20.8
All origins 3/, female												
2006	1.5	*	*	*	0.1	0.3	0.6	1.0	1.4	3.0	7.0	19.5
2005	1.8	*	*	*	0.1	0.4	0.8	1.1	1.7	3.3	8.8	24.8
2004	1.7	*	*	*	*	0.3	0.7	1.1	1.8	3.4	8.5	25.0
2003	1.9	*	*	*	0.1	0.4	0.8	1.2	1.7	4.0	9.4	26.2
2002	1.9	*	*	*	*	0.4	0.7	1.2	2.1	3.8	10.1	26.2
2001	1.9	*	*	*	0.1	0.4	0.6	1.0	2.0	4.4	9.2	27.8
2000	1.8	*	*	*	0.1	0.4	0.6	1.1	2.0	4.0	9.2	26.9
1999	1.6	*	*	*	*	0.2	0.5	1.0	1.8	3.4	8.8	24.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
Hispanic, both sexes												
2006	0.5	*	*	*	*	*	*	0.9	1.2	2.3	4.7	11.3
2005	0.7	*	*	*	*	*	0.6	1.0	1.5	3.5	8.1	15.0
2004	0.5	*	*	*	*	*	0.3	1.2	1.6	2.0	4.8	17.6
2003	0.6	*	*	*	*	*	0.4	1.1	1.4	2.6	6.4	17.6
2002	0.6	*	*	*	*	*	0.4	1.0	1.4	3.1	6.3	18.3
2001	0.6	*	*	*	*	*	*	0.9	*	4.2	7.4	32.5
2000	0.6	*	*	*	*	*	0.6	1.1	1.3	3.3	8.7	24.6
1999	0.5	*	*	*	*	*	*	1.1	1.5	4.1	5.4	24.6
Hispanic, male												
2006	0.4	*	*	*	*	*	*	1.2	*	*	*	*
2005	0.6	*	*	*	*	*	*	1.2	2.0	3.9	*	*
2004	0.5	*	*	*	*	*	*	1.4	2.2	*	*	*
2003	0.6	*	*	*	*	*	*	1.4	*	*	*	*
2002	0.5	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	1.3	*	4.0	8.9	*
2000	0.6	*	*	*	*	*	0.8	*	*	*	10.8	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.5	*	*	*	*	*	*	*	*	2.6	4.9	*
2005	0.8	*	*	*	*	*	0.7	*	*	3.1	10.7	13.7
2004	0.6	*	*	*	*	*	*	*	*	*	4.9	23.8
2003	0.6	*	*	*	*	*	*	*	*	*	6.6	19.5
2002	0.7	*	*	*	*	*	*	*	*	3.2	6.2	23.2
2001	0.6	*	*	*	*	*	*	*	*	4.4	6.3	33.5
2000	0.7	*	*	*	*	*	*	*	*	*	7.3	23.7
1999	0.6	*	*	*	*	*	*	*	*	4.3	*	23.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
Non-Hispanic 4/, both sexes												
2006	1.5	0.8	*	*	0.1	0.3	0.7	1.3	1.7	3.0	7.1	18.8
2005	1.8	0.8	*	*	0.1	0.4	0.8	1.5	2.1	3.8	8.6	23.4
2004	1.8	0.8	*	*	0.1	0.4	0.8	1.4	2.2	3.7	9.2	23.5
2003	1.8	0.8	*	*	0.1	0.4	0.8	1.4	2.1	4.1	9.1	24.3
2002	1.8	0.6	*	*	0.1	0.4	0.7	1.4	2.3	3.9	10.0	24.7
2001	1.8	*	*	*	0.1	0.4	0.7	1.2	2.3	4.6	9.5	25.8
2000	1.8	0.7	*	*	0.1	0.4	0.7	1.4	2.2	4.4	9.4	25.6
1999	1.6	0.7	*	*	0.1	0.3	0.6	1.0	1.9	3.7	9.3	23.3
Non-Hispanic 4/, male												
2006	1.3	*	*	*	*	0.3	0.8	1.5	2.1	2.9	7.1	16.5
2005	1.6	*	*	*	*	0.3	0.8	1.9	2.6	4.5	8.7	19.3
2004	1.6	*	*	*	*	0.4	0.8	1.6	2.5	4.0	9.9	20.2
2003	1.5	*	*	*	*	0.3	0.8	1.6	2.4	4.2	8.6	19.3
2002	1.5	*	*	*	*	0.4	0.7	1.5	2.5	3.9	9.6	21.1
2001	1.5	*	*	*	*	0.3	0.7	1.4	2.6	4.9	9.8	21.6
2000	1.5	*	*	*	*	0.3	0.8	1.6	2.3	4.8	9.6	22.1
1999	1.4	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.7	20.5
Non-Hispanic 4/, female												
2006	1.6	*	*	*	0.1	0.3	0.6	1.0	1.4	3.0	7.1	19.8
2005	1.9	*	*	*	*	0.4	0.8	1.1	1.7	3.3	8.6	25.3
2004	1.9	*	*	*	*	0.4	0.7	1.1	1.9	3.5	8.7	25.0
2003	2.1	*	*	*	0.1	0.4	0.9	1.2	1.7	4.1	9.5	26.5
2002	2.1	*	*	*	*	0.4	0.7	1.2	2.1	3.8	10.2	26.2
2001	2.0	*	*	*	0.1	0.4	0.6	1.0	2.1	4.3	9.2	27.6
2000	2.0	*	*	*	*	0.4	0.6	1.2	2.1	4.1	9.3	27.0
1999	1.8	*	*	*	*	0.3	0.5	0.9	1.8	3.3	9.0	24.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
Non-Hispanic white, both sexes												
2006 5/	1.5	0.9	*	*	0.1	0.3	0.6	1.1	1.6	2.8	6.9	18.5
2005 5/	1.8	*	*	*	*	0.3	0.6	1.3	1.9	3.6	8.3	23.9
2004 5/	1.8	*	*	*	*	0.3	0.6	1.2	2.0	3.5	9.0	23.4
2003 5/	1.8	*	*	*	0.1	0.3	0.7	1.3	1.9	3.9	8.7	23.8
2002	1.8	*	*	*	*	0.3	0.6	1.2	2.1	3.6	9.6	24.5
2001	1.8	*	*	*	*	0.3	0.6	1.0	2.1	4.3	9.2	25.7
2000	1.8	*	*	*	*	0.3	0.6	1.2	1.9	4.0	9.2	25.5
1999	1.6	*	*	*	*	0.2	0.5	0.9	1.8	3.5	9.0	23.0
Non-Hispanic white, male												
2006 5/	1.3	*	*	*	*	0.2	0.7	1.3	1.9	2.9	6.8	16.3
2005 5/	1.6	*	*	*	*	0.3	0.7	1.7	2.4	4.1	8.3	19.4
2004 5/	1.6	*	*	*	*	0.3	0.7	1.4	2.4	3.8	9.6	20.3
2003 5/	1.5	*	*	*	*	0.3	0.7	1.3	2.2	3.9	8.0	18.7
2002	1.5	*	*	*	*	0.3	0.6	1.3	2.2	3.7	9.1	20.5
2001	1.5	*	*	*	*	0.2	0.6	1.2	2.3	4.6	9.4	21.2
2000	1.5	*	*	*	*	0.2	0.7	1.5	2.0	4.5	9.4	21.8
1999	1.4	*	*	*	*	0.2	0.6	0.9	2.0	3.9	9.3	20.3
Non-Hispanic white, female												
2006 5/	1.7	*	*	*	*	0.3	0.5	0.8	1.3	2.8	7.0	19.5
2005 5/	2.0	*	*	*	*	0.3	0.6	1.0	1.5	3.2	8.4	26.0
2004 5/	2.0	*	*	*	*	0.3	0.6	0.9	1.6	3.3	8.5	24.8
2003 5/	2.1	*	*	*	*	0.3	0.6	1.2	1.6	3.9	9.3	26.1
2002	2.2	*	*	*	*	0.3	0.6	1.1	1.9	3.6	9.9	26.3
2001	2.1	*	*	*	*	0.3	0.5	0.8	1.9	4.1	9.0	27.6
2000	2.0	*	*	*	*	0.3	0.5	1.0	1.9	3.6	9.0	27.0
1999	1.8	*	*	*	*	0.2	0.5	0.8	1.7	3.1	8.8	24.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
Non-Hispanic black, both sexes												
2006 5/	1.9	*	*	*	*	0.8	1.7	2.9	3.5	4.9	11.1	26.0
2005 5/	2.1	*	*	*	*	0.8	1.8	3.0	4.6	6.5	14.0	24.3
2004 5/	2.1	*	*	*	*	1.0	2.0	3.0	4.2	6.8	13.9	29.1
2003 5/	2.2	*	*	*	*	0.9	2.0	2.7	4.4	7.4	15.5	33.6
2002	2.2	*	*	*	*	1.0	1.7	2.7	5.0	6.9	16.8	29.9
2001	2.0	*	*	*	*	0.8	1.6	2.3	4.9	7.9	14.3	29.7
2000	2.1	*	*	*	*	0.9	1.5	2.8	4.6	9.0	14.4	29.9
1999	1.8	*	*	*	*	0.5	1.5	2.4	3.0	6.8	14.0	28.5
Non-Hispanic black, male												
2006 5/	1.8	*	*	*	*	0.8	1.5	3.5	4.2	5.1	12.0	22.9
2005 5/	2.1	*	*	*	*	*	1.7	3.8	5.7	8.9	16.0	23.8
2004 5/	2.0	*	*	*	*	1.1	2.0	3.5	4.5	7.3	15.4	25.4
2003 5/	2.1	*	*	*	*	0.9	1.4	3.5	5.4	8.4	17.9	33.0
2002	2.1	*	*	*	*	1.1	1.8	3.0	5.6	7.8	18.5	31.7
2001	2.0	*	*	*	*	*	1.8	2.4	6.5	9.2	15.9	28.9
2000	2.0	*	*	*	*	0.9	1.8	3.5	5.3	9.1	13.3	30.5
1999	1.7	*	*	*	*	*	1.9	2.9	2.7	8.4	17.2	*
Non-Hispanic black, female												
2006 5/	1.9	*	*	*	*	0.8	1.8	2.3	2.9	4.8	10.6	27.3
2005 5/	2.1	*	*	*	*	0.9	1.9	2.4	3.6	4.8	12.9	24.5
2004 5/	2.3	*	*	*	*	0.8	1.9	2.6	3.9	6.5	13.0	30.5
2003 5/	2.3	*	*	*	*	1.0	2.6	1.9	3.6	6.8	14.2	33.8
2002	2.3	*	*	*	*	1.0	1.6	2.4	4.5	6.2	15.9	29.3
2001	2.1	*	*	*	*	0.9	1.4	2.1	3.7	7.0	13.5	29.9
2000	2.3	*	*	*	*	1.0	1.3	2.3	4.0	8.8	15.1	29.7
1999	1.8	*	*	*	*	*	1.1	2.0	3.2	5.6	12.3	30.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
All origins 3/, both sexes												
2006	18.8	6.4	0.8	0.2	0.4	0.8	1.9	4.6	10.0	32.0	127.8	502.5
2005	21.3	6.5	0.7	0.3	0.4	0.9	2.1	5.1	11.3	35.5	142.2	593.9
2004	20.3	6.7	0.7	0.2	0.4	0.8	2.0	4.6	10.8	34.6	139.3	582.6
2003	22.4	8.0	1.0	0.4	0.5	0.9	2.2	5.2	11.2	37.3	151.1	666.1
2002	22.8	6.5	0.7	0.2	0.4	0.9	2.2	4.8	11.2	37.5	156.9	696.6
2001	21.8	7.4	0.7	0.2	0.5	0.9	2.2	4.6	10.7	36.3	148.5	685.6
2000	23.2	7.6	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.1	160.3	744.1
1999	22.8	8.4	0.8	0.2	0.5	0.8	2.4	4.6	11.0	37.2	157.0	751.8
All origins 3/, male												
2006	17.4	7.7	0.8	0.2	0.5	0.9	2.1	5.6	12.3	38.9	156.9	586.4
2005	19.2	7.0	0.7	0.3	0.5	1.0	2.5	6.2	13.6	42.8	174.3	673.5
2004	18.6	8.0	0.6	0.2	0.6	1.0	2.2	5.6	13.0	42.8	172.7	677.2
2003	20.1	9.4	1.1	0.3	0.6	1.1	2.5	6.4	13.2	46.5	186.8	757.8
2002	20.4	7.8	0.8	0.2	0.5	0.9	2.4	5.8	13.6	46.4	194.7	805.8
2001	19.6	8.1	0.7	0.2	0.5	1.0	2.5	5.7	13.1	45.0	186.4	808.4
2000	20.8	8.4	0.7	0.2	0.5	1.0	2.8	5.8	14.4	48.3	199.8	892.0
1999	20.3	8.8	0.8	0.3	0.5	1.0	2.6	5.9	12.8	46.9	196.5	894.7
All origins 3/, female												
2006	20.2	5.0	0.8	0.2	0.4	0.8	1.7	3.7	7.8	26.3	107.8	463.3
2005	23.2	5.8	0.7	0.3	0.3	0.8	1.8	4.1	9.1	29.4	120.4	557.4
2004	22.0	5.4	0.9	0.2	0.3	0.6	1.8	3.6	8.8	27.7	116.8	540.0
2003	24.6	6.6	0.9	0.4	0.5	0.7	1.9	4.1	9.4	29.5	127.2	625.6
2002	25.1	5.1	0.6	0.3	0.4	0.8	1.9	3.8	9.0	30.1	131.8	649.2
2001	23.9	6.7	0.7	0.2	0.4	0.8	1.9	3.5	8.5	29.1	123.7	634.2
2000	25.6	6.8	0.7	0.2	0.5	0.8	1.9	3.7	9.5	31.5	134.6	683.9
1999	25.3	8.0	0.9	0.2	0.5	0.7	2.1	3.4	9.4	29.3	131.3	694.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
Hispanic, both sexes												
2006	6.7	6.2	0.7	*	0.3	0.6	1.2	3.7	9.8	30.8	114.2	390.9
2005	7.2	7.1	0.9	*	*	0.5	1.5	4.0	10.4	34.4	122.3	456.9
2004	7.0	5.8	0.7	*	0.5	0.6	1.3	3.8	9.6	33.2	125.2	482.3
2003	7.4	7.6	1.1	0.3	0.5	0.7	1.3	4.0	10.7	35.0	134.3	516.2
2002	7.3	5.8	*	*	*	0.6	1.3	3.4	10.5	33.8	142.6	566.0
2001	7.4	7.0	1.1	*	*	0.3	1.3	3.6	9.1	35.4	137.2	660.3
2000	7.4	7.8	*	0.3	0.3	0.7	1.6	3.8	11.2	37.6	136.3	643.6
1999	6.6	8.5	*	*	0.4	0.4	1.5	3.5	9.5	31.7	118.5	614.6
Hispanic, male												
2006	6.3	7.3	*	*	*	0.7	1.4	4.5	12.6	36.7	131.5	440.9
2005	6.8	7.8	*	*	*	0.6	1.7	5.1	12.9	42.3	149.7	496.9
2004	6.6	8.0	*	*	0.6	0.7	1.5	4.9	11.5	38.9	152.0	528.2
2003	6.9	9.1	1.4	*	0.5	0.9	1.3	5.2	11.6	43.4	165.7	577.2
2002	7.0	7.7	*	*	*	0.7	1.7	4.7	13.9	43.8	173.2	655.9
2001	6.9	7.4	*	*	*	*	1.7	5.0	11.4	40.7	173.5	765.6
2000	7.2	7.6	*	*	*	0.8	2.1	5.3	14.8	46.0	180.4	763.9
1999	6.3	10.3	*	*	*	*	2.1	4.9	10.4	41.1	148.1	720.3
Hispanic, female												
2006	7.1	5.1	*	*	*	*	0.9	2.8	7.3	26.0	102.0	364.3
2005	7.7	6.4	*	*	*	*	1.1	2.9	8.1	28.1	103.3	436.0
2004	7.6	*	*	*	*	*	1.2	2.6	7.9	28.7	106.7	458.8
2003	7.9	6.1	*	*	*	*	1.2	2.7	9.9	28.4	112.7	485.2
2002	7.6	*	*	*	*	*	0.9	2.1	7.4	25.9	121.7	521.4
2001	7.9	6.5	*	*	*	*	1.0	2.1	7.1	31.2	112.7	608.8
2000	7.7	8.0	*	*	*	0.7	1.0	2.4	7.9	31.0	106.6	584.6
1999	7.0	6.5	*	*	*	*	0.8	2.2	8.6	24.3	98.9	563.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
Non-Hispanic 4/, both sexes												
2006	20.9	6.3	0.8	0.2	0.5	0.9	2.1	4.7	10.0	32.0	128.4	507.5
2005	23.6	6.2	0.6	0.3	0.5	1.0	2.2	5.2	11.3	35.5	143.1	599.9
2004	22.4	6.9	0.8	0.2	0.4	0.8	2.1	4.6	10.9	34.5	139.7	586.0
2003	24.7	8.1	1.0	0.4	0.5	1.0	2.4	5.3	11.2	37.2	151.6	671.1
2002	25.1	6.7	0.7	0.2	0.4	0.9	2.3	4.9	11.3	37.6	157.1	700.2
2001	23.8	7.5	0.6	0.2	0.5	1.0	2.3	4.7	10.7	36.1	148.6	684.8
2000	25.4	7.5	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.0	160.7	745.7
1999	25.0	8.4	0.9	0.2	0.5	0.9	2.4	4.7	11.0	37.4	158.0	754.0
Non-Hispanic 4/, male												
2006	19.4	7.8	0.8	0.2	0.5	0.9	2.2	5.7	12.3	38.9	158.2	593.8
2005	21.4	6.7	0.6	0.3	0.6	1.0	2.6	6.3	13.6	42.6	175.4	682.2
2004	20.6	8.0	0.6	0.2	0.5	1.0	2.3	5.6	13.0	42.7	173.5	683.2
2003	22.3	9.4	1.1	0.3	0.6	1.2	2.7	6.4	13.2	46.4	187.6	764.6
2002	22.5	7.8	0.8	0.2	0.5	1.0	2.5	5.8	13.5	46.3	195.1	810.7
2001	21.5	8.2	0.6	0.2	0.5	1.1	2.6	5.7	13.1	44.9	186.5	808.2
2000	22.7	8.5	0.7	0.2	0.6	1.1	2.9	5.7	14.3	48.2	199.7	894.9
1999	22.2	8.4	0.9	0.3	0.5	1.1	2.7	5.9	12.9	47.0	197.7	897.9
Non-Hispanic 4/, female												
2006	22.3	4.8	0.8	0.2	0.4	0.9	1.9	3.8	7.8	26.2	108.0	467.4
2005	25.7	5.7	0.6	0.3	0.3	0.9	1.9	4.2	9.2	29.5	121.2	562.3
2004	24.2	5.9	0.9	0.2	0.3	0.6	1.9	3.7	8.9	27.5	117.0	542.5
2003	27.1	6.8	1.0	0.4	0.5	0.8	2.1	4.3	9.3	29.5	127.7	630.0
2002	27.5	5.4	0.7	0.3	0.4	0.8	2.0	4.0	9.2	30.3	131.8	652.6
2001	26.1	6.7	0.6	0.2	0.4	0.8	2.0	3.6	8.5	28.9	123.8	633.5
2000	27.9	6.5	0.6	0.2	0.5	0.8	2.0	3.8	9.6	31.4	135.3	685.4
1999	27.6	8.4	0.9	0.2	0.5	0.8	2.2	3.5	9.4	29.5	132.2	696.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
Non-Hispanic white, both sexes												
2006 5/	23.1	4.5	0.6	0.2	0.4	0.8	1.9	4.2	9.5	30.9	128.0	514.4
2005 5/	26.2	4.1	0.5	0.3	0.5	0.9	2.0	4.7	10.4	33.9	143.3	612.6
2004 5/	24.8	4.8	0.6	0.2	0.4	0.8	1.8	4.0	10.1	33.2	138.8	593.8
2003 5/	27.4	6.2	0.9	0.3	0.5	0.8	2.1	4.7	10.1	35.7	151.9	684.5
2002	27.9	4.9	0.5	0.2	0.4	0.8	2.0	4.4	10.3	36.7	156.9	712.7
2001	26.2	4.9	0.5	0.2	0.4	0.8	1.9	4.0	9.9	35.0	147.2	693.9
2000	27.9	5.0	0.5	0.2	0.5	0.8	2.1	4.1	10.7	36.8	160.2	759.8
1999	27.5	5.3	0.6	0.2	0.4	0.8	2.0	4.0	10.0	36.1	158.8	766.9
Non-Hispanic white, male												
2006 5/	21.2	5.8	0.6	0.2	0.5	0.8	2.1	4.9	11.5	37.1	156.9	598.8
2005 5/	23.3	5.1	0.5	0.2	0.6	1.0	2.3	5.5	12.2	40.1	174.6	693.4
2004 5/	22.4	5.5	0.5	0.2	0.5	0.9	2.0	4.9	12.0	40.3	170.7	691.1
2003 5/	24.1	7.4	0.9	0.3	0.6	1.1	2.3	5.6	11.4	43.6	185.5	774.1
2002	24.6	6.1	0.5	0.2	0.5	0.9	2.2	5.2	12.1	44.9	193.2	820.6
2001	23.1	5.8	0.6	0.2	0.5	0.9	2.1	4.8	11.7	42.7	182.5	813.9
2000	24.5	5.8	0.5	0.2	0.5	0.9	2.4	4.7	12.6	44.7	197.5	908.1
1999	23.9	5.5	0.5	0.3	0.4	0.9	2.1	4.8	11.5	44.6	197.3	909.3
Non-Hispanic white, female												
2006 5/	25.0	3.0	0.6	*	0.4	0.8	1.8	3.5	7.5	25.6	107.9	475.0
2005 5/	28.9	3.0	0.5	0.3	0.3	0.9	1.8	3.8	8.8	28.6	121.7	575.7
2004 5/	27.1	4.2	0.7	0.2	0.3	0.6	1.7	3.2	8.3	27.0	117.0	550.2
2003 5/	30.6	5.0	0.8	0.4	0.5	0.6	1.8	3.8	8.8	28.9	129.3	645.1
2002	31.1	3.6	0.5	0.3	0.3	0.8	1.8	3.7	8.6	29.7	132.5	666.1
2001	29.2	3.9	0.5	0.2	0.4	0.8	1.7	3.3	8.1	28.5	123.7	644.0
2000	31.3	4.2	0.6	0.2	0.4	0.8	1.8	3.4	9.0	30.1	135.7	700.0
1999	31.0	5.1	0.8	0.2	0.4	0.6	1.9	3.1	8.7	29.0	133.5	710.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
Non-Hispanic black, both sexes												
2006 5/	13.9	13.7	1.4	*	0.6	1.4	3.4	9.0	16.2	45.5	141.2	439.6
2005 5/	15.3	15.1	1.1	0.4	0.5	1.5	3.9	10.1	20.3	53.2	151.8	475.4
2004 5/	15.2	14.7	1.4	0.3	0.7	1.2	4.1	9.3	19.1	50.4	158.4	511.7
2003 5/	15.9	14.4	1.1	0.4	0.7	1.9	4.5	10.5	21.1	55.0	157.7	528.3
2002	16.1	13.3	1.7	*	0.6	1.6	4.5	8.9	21.6	50.9	167.7	579.8
2001	16.0	16.9	1.0	*	0.8	1.7	4.9	9.9	19.1	48.8	169.5	583.1
2000	16.9	17.5	1.2	*	0.8	1.8	4.6	10.3	22.3	60.8	172.2	604.5
1999	16.7	20.0	1.7	*	0.9	2.0	5.5	10.8	20.9	54.7	158.3	634.7
Non-Hispanic black, male												
2006 5/	13.6	16.3	*	*	0.7	1.4	3.7	11.7	21.8	59.5	182.1	512.2
2005 5/	15.1	14.0	*	*	0.7	1.7	5.0	13.2	28.1	71.2	191.5	537.6
2004 5/	15.5	17.0	*	*	*	1.6	4.9	12.1	24.8	73.2	221.0	607.6
2003 5/	16.5	16.1	*	*	0.8	2.3	5.5	13.3	29.8	79.1	220.8	642.5
2002	15.9	14.2	1.8	*	0.7	1.8	5.2	11.5	29.6	66.5	231.2	695.8
2001	16.3	16.6	*	*	0.8	2.1	5.9	13.5	26.6	70.0	242.5	718.4
2000	17.3	18.8	1.7	*	0.8	2.4	5.8	14.3	30.3	84.1	235.4	763.2
1999	16.9	20.1	2.1	*	0.8	2.3	6.7	15.3	27.3	77.9	215.1	803.4
Non-Hispanic black, female												
2006 5/	14.1	10.9	*	*	*	1.4	3.0	6.6	11.7	35.4	118.0	409.9
2005 5/	15.5	16.2	*	*	*	1.2	3.0	7.4	14.0	40.3	129.3	450.4
2004 5/	15.0	12.2	1.8	*	0.7	0.8	3.3	6.9	14.5	34.1	123.0	473.8
2003 5/	15.4	12.6	*	*	*	1.6	3.7	8.1	14.0	37.8	122.1	484.1
2002	16.2	12.3	*	*	*	1.4	3.9	6.6	15.2	39.8	131.8	535.5
2001	15.6	17.1	*	*	0.8	1.3	4.0	6.9	13.0	33.9	128.5	533.0
2000	16.5	16.2	*	*	0.8	1.2	3.5	6.9	15.9	44.4	136.7	545.6
1999	16.5	20.0	*	*	0.9	1.8	4.5	7.0	15.8	38.4	126.4	571.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
All origins 3/, both sexes												
2006	0.3	*	*	0.1	*	*	*	0.1	0.1	0.4	1.8	7.0
2005	0.6	*	*	0.0	*	*	*	0.1	0.2	0.7	3.7	19.3
2004	0.4	*	*	*	*	*	*	0.1	0.2	0.5	2.2	11.4
2003	0.6	0.8	0.4	0.1	0.1	*	0.1	0.1	0.3	1.0	3.6	17.0
2002	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	8.8
2001	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.2
2000	0.6	*	*	*	*	*	0.0	0.1	0.4	1.1	4.0	20.2
1999	0.6	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.2
All origins 3/, male												
2006	0.2	*	*	*	*	*	*	*	0.1	0.5	2.0	8.1
2005	0.5	*	*	*	*	*	*	0.1	0.3	0.7	3.6	18.2
2004	0.3	*	*	*	*	*	*	*	0.2	0.7	2.4	11.1
2003	0.5	*	0.4	*	*	*	*	0.2	0.3	1.1	3.7	17.0
2002	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	8.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.2
2000	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.0	20.7
1999	0.4	*	*	*	*	*	*	*	0.3	0.9	4.2	19.9
All origins 3/, female												
2006	0.3	*	*	*	*	*	*	*	0.1	0.4	1.7	6.5
2005	0.8	*	*	*	*	*	*	*	0.2	0.8	3.7	19.8
2004	0.5	*	*	*	*	*	*	*	0.2	0.4	2.1	11.5
2003	0.7	*	0.3	0.1	*	*	*	0.1	0.3	0.9	3.5	17.1
2002	0.3	*	*	*	*	*	*	*	*	0.2	1.3	8.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000	0.8	*	*	*	*	*	*	0.1	0.4	0.9	4.0	19.9
1999	0.7	*	*	*	*	*	*	*	0.3	0.9	3.4	21.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
Hispanic, both sexes												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
Non-Hispanic 4/, both sexes												
2006	0.3	*	*	*	*	*	*	0.1	0.1	0.5	1.9	7.3
2005	0.7	*	*	*	*	*	*	0.1	0.2	0.8	3.8	19.8
2004	0.4	*	*	*	*	*	*	0.1	0.2	0.6	2.3	11.8
2003	0.7	0.7	0.4	0.1	0.1	*	0.1	0.2	0.3	1.1	3.6	17.5
2002	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	9.0
2001	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.3
2000	0.7	*	*	*	*	*	0.0	0.1	0.4	1.1	4.1	20.5
1999	0.7	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.6
Non-Hispanic 4/, male												
2006	0.3	*	*	*	*	*	*	*	*	0.5	2.1	8.4
2005	0.5	*	*	*	*	*	*	0.1	0.3	0.7	3.8	18.7
2004	0.3	*	*	*	*	*	*	*	0.2	0.7	2.5	11.6
2003	0.5	*	0.4	*	*	*	*	0.2	0.3	1.1	3.7	17.5
2002	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	9.2
2001	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2000	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.1	21.1
1999	0.5	*	*	*	*	*	*	*	0.3	0.8	4.3	20.7
Non-Hispanic 4/, female												
2006	0.4	*	*	*	*	*	*	*	0.1	0.4	1.8	6.7
2005	0.9	*	*	*	*	*	*	*	0.2	0.8	3.8	20.3
2004	0.5	*	*	*	*	*	*	*	0.2	0.4	2.2	11.8
2003	0.8	*	0.4	0.1	*	*	*	0.1	0.3	1.0	3.6	17.6
2002	0.3	*	*	*	*	*	*	*	*	0.2	1.3	8.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000	0.9	*	*	*	*	*	*	0.1	0.4	1.0	4.1	20.3
1999	0.8	*	*	*	*	*	*	*	0.3	0.9	3.4	22.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
Non-Hispanic white, both sexes												
2006 5/	0.4	*	*	*	*	*	*	0.1	0.2	0.5	2.0	7.8
2005 5/	0.8	*	*	*	*	*	*	0.1	0.2	0.8	4.1	21.5
2004 5/	0.5	*	*	*	*	*	*	*	0.2	0.6	2.5	12.7
2003 5/	0.8	*	0.3	0.1	*	*	*	0.2	0.3	1.1	3.9	18.8
2002	0.3	*	*	*	*	*	*	*	0.1	0.3	1.6	9.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2000	0.8	*	*	*	*	*	*	0.1	0.4	1.2	4.3	21.7
1999	0.8	*	*	*	*	*	*	0.1	0.3	0.9	3.9	23.3
Non-Hispanic white, male												
2006 5/	0.3	*	*	*	*	*	*	*	*	0.6	2.2	8.9
2005 5/	0.6	*	*	*	*	*	*	*	0.3	0.8	4.1	19.9
2004 5/	0.4	*	*	*	*	*	*	*	0.2	0.8	2.5	12.7
2003 5/	0.6	*	*	*	*	*	*	0.2	0.3	1.2	4.0	18.9
2002	0.3	*	*	*	*	*	*	*	*	0.4	1.7	10.0
2001	0.1	*	*	*	*	*	*	*	*	*	*	2.4
2000	0.6	*	*	*	*	*	*	*	0.4	1.4	4.3	22.0
1999	0.6	*	*	*	*	*	*	*	0.3	0.9	4.6	22.5
Non-Hispanic white, female												
2006 5/	0.4	*	*	*	*	*	*	*	0.2	0.4	2.0	7.3
2005 5/	1.0	*	*	*	*	*	*	*	0.2	0.9	4.2	22.2
2004 5/	0.6	*	*	*	*	*	*	*	0.2	0.5	2.4	12.8
2003 5/	1.0	*	*	*	*	*	*	0.1	0.3	1.1	3.9	18.7
2002	0.4	*	*	*	*	*	*	*	*	*	1.4	9.5
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2000	1.0	*	*	*	*	*	*	0.1	0.5	1.0	4.3	21.6
1999	1.0	*	*	*	*	*	*	*	0.3	1.0	3.5	23.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
Non-Hispanic black, both sexes												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.3	*	*	*	*	*	*	*	*	*	*	5.9
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	9.5
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
All origins 3/, both sexes												
2006	18.5	5.9	0.7	0.1	0.4	0.8	1.9	4.6	9.8	31.6	125.9	495.5
2005	20.6	6.0	0.6	0.2	0.4	0.9	2.1	5.0	11.1	34.8	138.5	574.7
2004	19.9	6.3	0.6	0.2	0.4	0.7	2.0	4.5	10.7	34.0	137.0	571.2
2003	21.8	7.3	0.7	0.2	0.5	0.9	2.2	5.1	10.9	36.3	147.5	649.1
2002	22.5	6.3	0.7	0.2	0.4	0.9	2.1	4.7	11.1	37.2	155.5	687.8
2001	21.7	7.2	0.7	0.2	0.4	0.8	2.2	4.6	10.6	36.2	148.1	683.4
2000	22.6	7.4	0.6	0.2	0.4	0.9	2.3	4.6	11.5	38.0	156.4	723.9
1999	22.2	8.1	0.8	0.2	0.4	0.8	2.3	4.6	10.7	36.4	153.3	730.6
All origins 3/, male												
2006	17.1	7.2	0.6	0.1	0.4	0.8	2.1	5.5	12.2	38.4	155.0	578.3
2005	18.8	6.6	0.6	0.2	0.5	0.9	2.4	6.1	13.3	42.1	170.7	655.3
2004	18.3	7.5	0.5	0.2	0.5	0.9	2.2	5.5	12.8	42.1	170.3	666.1
2003	19.6	8.5	0.7	0.2	0.5	1.1	2.5	6.2	12.9	45.4	183.2	740.9
2002	20.2	7.6	0.8	0.2	0.4	0.9	2.4	5.7	13.4	46.0	193.1	796.9
2001	19.5	7.8	0.7	0.2	0.5	0.9	2.5	5.7	13.0	44.9	186.0	806.2
2000	20.3	8.1	0.6	0.2	0.5	1.0	2.7	5.6	14.0	47.1	195.8	871.3
1999	19.8	8.5	0.7	0.2	0.4	0.9	2.6	5.8	12.5	46.0	192.3	874.7
All origins 3/, female												
2006	19.9	4.6	0.7	*	0.4	0.8	1.7	3.7	7.7	25.9	106.1	456.7
2005	22.5	5.3	0.6	0.2	0.2	0.8	1.8	4.0	8.9	28.7	116.7	537.6
2004	21.5	5.0	0.8	0.2	0.3	0.5	1.8	3.5	8.6	27.3	114.6	528.5
2003	23.9	6.0	0.6	0.3	0.4	0.7	1.9	4.0	9.1	28.6	123.7	608.5
2002	24.8	5.0	0.6	0.2	0.3	0.8	1.9	3.8	9.0	29.8	130.6	640.5
2001	23.8	6.7	0.7	0.2	0.4	0.7	1.9	3.5	8.4	29.0	123.2	632.0
2000	24.8	6.6	0.6	0.2	0.4	0.8	1.9	3.6	9.1	30.6	130.6	663.9
1999	24.6	7.7	0.8	0.2	0.4	0.7	2.0	3.4	9.1	28.4	128.0	672.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
Hispanic, both sexes												
2006	6.6	5.7	0.5	*	0.3	0.6	1.2	3.6	9.7	30.7	113.8	388.9
2005	7.1	6.2	0.7	*	*	0.5	1.5	3.9	10.3	34.1	120.9	450.5
2004	7.0	5.2	0.6	*	0.4	0.6	1.3	3.7	9.6	32.9	124.9	479.5
2003	7.2	6.5	0.7	*	0.5	0.7	1.2	3.9	10.4	34.8	132.4	511.0
2002	7.2	5.5	*	*	*	0.6	1.3	3.4	10.5	33.6	142.1	563.2
2001	7.3	7.0	1.0	*	*	0.3	1.3	3.5	9.1	35.3	136.3	660.3
2000	7.3	7.5	*	*	0.3	0.7	1.6	3.8	11.0	37.2	134.3	633.0
1999	6.5	7.9	*	*	0.4	0.4	1.5	3.4	9.2	30.9	116.7	607.6
Hispanic, male												
2006	6.2	6.4	*	*	*	0.7	1.4	4.4	12.5	36.5	131.5	438.7
2005	6.7	6.7	*	*	*	0.6	1.7	4.9	12.8	42.0	149.0	488.2
2004	6.5	7.3	*	*	0.6	0.7	1.5	4.9	11.5	38.5	151.3	526.8
2003	6.7	7.5	*	*	*	0.8	1.3	5.2	11.4	42.8	163.0	571.1
2002	6.9	7.3	*	*	*	0.6	1.7	4.7	13.9	43.6	172.8	654.2
2001	6.8	7.4	*	*	*	*	1.7	5.0	11.4	40.7	172.6	765.6
2000	7.1	7.6	*	*	*	0.7	2.1	5.2	14.6	45.6	178.0	751.8
1999	6.1	10.3	*	*	*	*	2.0	4.7	9.9	39.5	147.0	718.2
Hispanic, female												
2006	7.0	4.8	*	*	*	*	0.9	2.8	7.1	26.0	101.4	362.5
2005	7.5	5.7	*	*	*	*	1.1	2.9	8.0	27.8	101.4	430.8
2004	7.5	*	*	*	*	*	1.2	2.5	7.9	28.4	106.7	455.2
2003	7.7	5.4	*	*	*	*	1.1	2.6	9.6	28.4	111.4	480.5
2002	7.5	*	*	*	*	*	0.9	2.1	7.4	25.7	121.1	518.1
2001	7.8	6.5	*	*	*	*	0.9	2.1	7.1	31.0	111.8	608.8
2000	7.6	7.4	*	*	*	*	1.0	2.3	7.9	30.7	105.0	574.7
1999	6.8	5.4	*	*	*	*	0.8	2.2	8.5	24.1	96.4	554.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
Non-Hispanic 4/, both sexes												
2006	20.6	5.9	0.7	0.1	0.4	0.9	2.0	4.7	9.9	31.6	126.5	500.2
2005	22.9	5.9	0.5	0.2	0.4	0.9	2.2	5.1	11.1	34.7	139.2	580.1
2004	22.0	6.6	0.6	0.2	0.4	0.8	2.1	4.6	10.7	33.9	137.4	574.2
2003	24.0	7.4	0.6	0.3	0.5	0.9	2.3	5.2	10.9	36.2	148.0	653.6
2002	24.8	6.5	0.7	0.2	0.4	0.9	2.3	4.9	11.1	37.3	155.6	691.2
2001	23.8	7.2	0.6	0.2	0.5	0.9	2.3	4.6	10.6	36.0	148.2	682.5
2000	24.7	7.3	0.6	0.2	0.5	0.9	2.4	4.6	11.4	37.9	156.6	725.2
1999	24.3	8.1	0.8	0.2	0.4	0.9	2.4	4.6	10.7	36.5	154.2	732.4
Non-Hispanic 4/, male												
2006	19.1	7.4	0.7	0.1	0.5	0.9	2.2	5.6	12.2	38.4	156.1	585.5
2005	20.9	6.5	0.5	0.2	0.6	1.0	2.5	6.2	13.3	41.9	171.5	663.5
2004	20.3	7.5	0.5	0.2	0.5	1.0	2.3	5.6	12.8	42.0	171.0	671.6
2003	21.7	8.7	0.7	0.2	0.5	1.1	2.7	6.3	12.9	45.3	183.9	747.1
2002	22.3	7.6	0.8	0.2	0.5	1.0	2.5	5.8	13.4	45.9	193.5	801.5
2001	21.4	7.8	0.6	0.2	0.5	1.0	2.6	5.7	13.0	44.8	186.1	805.9
2000	22.2	8.2	0.7	0.2	0.5	1.0	2.8	5.6	13.9	46.8	195.7	873.8
1999	21.7	8.0	0.8	0.2	0.5	1.1	2.6	5.8	12.5	46.2	193.4	877.2
Non-Hispanic 4/, female												
2006	21.9	4.4	0.7	*	0.4	0.9	1.9	3.8	7.7	25.8	106.2	460.6
2005	24.8	5.2	0.5	0.3	0.3	0.8	1.9	4.1	9.0	28.7	117.3	542.0
2004	23.7	5.6	0.8	0.2	0.3	0.6	1.8	3.6	8.7	27.1	114.8	530.7
2003	26.3	6.1	0.6	0.3	0.4	0.7	2.0	4.1	9.0	28.5	124.1	612.4
2002	27.2	5.3	0.6	0.2	0.4	0.8	2.0	4.0	9.1	30.1	130.6	643.7
2001	26.0	6.6	0.6	0.2	0.4	0.8	2.0	3.6	8.5	28.8	123.3	631.3
2000	27.1	6.4	0.5	0.1	0.4	0.8	2.0	3.7	9.2	30.4	131.2	665.2
1999	26.8	8.2	0.9	0.2	0.4	0.8	2.2	3.5	9.1	28.5	128.8	674.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
Non-Hispanic white, both sexes												
2006 5/	22.7	4.1	0.6	0.1	0.4	0.8	1.9	4.2	9.3	30.4	125.9	506.5
2005 5/	25.3	3.8	0.4	0.2	0.4	0.9	2.0	4.6	10.2	33.1	139.2	591.2
2004 5/	24.3	4.6	0.5	0.1	0.4	0.7	1.8	4.0	9.9	32.6	136.3	581.1
2003 5/	26.6	5.7	0.6	0.3	0.5	0.8	2.0	4.5	9.8	34.6	148.0	665.7
2002	27.6	4.7	0.5	0.2	0.4	0.8	2.0	4.4	10.2	36.4	155.3	703.1
2001	26.1	4.6	0.5	0.2	0.4	0.8	1.9	4.0	9.8	34.9	146.7	691.5
2000	27.1	4.8	0.5	0.2	0.4	0.8	2.1	3.9	10.3	35.6	155.9	738.1
1999	26.8	5.2	0.6	0.2	0.4	0.7	1.9	3.9	9.7	35.2	154.8	743.7
Non-Hispanic white, male												
2006 5/	20.9	5.4	0.6	*	0.5	0.8	2.0	4.9	11.4	36.5	154.7	589.9
2005 5/	22.7	4.7	*	0.2	0.6	0.9	2.2	5.4	11.9	39.3	170.5	673.6
2004 5/	22.0	5.3	*	*	0.5	0.9	2.0	4.8	11.8	39.5	168.2	678.4
2003 5/	23.5	6.8	0.6	0.2	0.5	1.0	2.3	5.4	11.1	42.5	181.5	755.2
2002	24.3	6.0	0.5	0.2	0.5	0.9	2.2	5.1	11.9	44.5	191.5	810.6
2001	23.0	5.3	0.6	0.2	0.5	0.9	2.1	4.8	11.6	42.6	182.0	811.5
2000	23.8	5.4	0.4	0.2	0.5	0.8	2.4	4.6	12.2	43.3	193.2	886.1
1999	23.4	5.4	*	0.2	0.4	0.9	2.0	4.7	11.1	43.8	192.7	886.8
Non-Hispanic white, female												
2006 5/	24.6	2.8	0.6	*	0.4	0.8	1.7	3.4	7.4	25.2	105.9	467.7
2005 5/	27.9	2.8	0.4	0.3	0.3	0.8	1.7	3.7	8.5	27.7	117.6	553.5
2004 5/	26.5	3.8	0.6	*	0.2	0.5	1.6	3.2	8.1	26.6	114.6	537.5
2003 5/	29.6	4.5	0.5	0.3	0.4	0.6	1.8	3.7	8.5	27.9	125.4	626.4
2002	30.7	3.4	0.5	0.2	0.3	0.8	1.8	3.7	8.5	29.5	131.1	656.6
2001	29.1	3.8	0.4	0.2	0.3	0.7	1.7	3.2	8.0	28.3	123.3	641.7
2000	30.3	4.2	0.5	*	0.3	0.8	1.7	3.3	8.5	29.1	131.4	678.4
1999	30.0	4.9	0.7	0.2	0.3	0.6	1.8	3.0	8.4	28.0	130.0	686.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
Non-Hispanic black, both sexes												
2006 5/	13.8	12.9	1.2	*	0.6	1.4	3.3	8.9	16.2	45.3	140.2	437.7
2005 5/	15.1	14.3	1.0	*	0.5	1.4	3.8	9.9	20.1	52.9	150.4	471.3
2004 5/	15.1	14.0	1.2	*	0.6	1.2	4.0	9.2	18.9	50.0	156.7	509.2
2003 5/	15.6	13.3	0.9	*	0.6	1.9	4.5	10.2	20.8	54.4	156.4	522.4
2002	16.0	13.1	1.5	*	0.6	1.6	4.5	8.9	21.4	50.5	167.3	577.4
2001	15.9	16.9	1.0	*	0.8	1.7	4.9	9.9	19.0	48.7	169.1	581.5
2000	16.6	17.2	1.2	*	0.8	1.8	4.5	10.1	22.1	59.9	170.8	595.0
1999	16.5	19.3	1.7	*	0.9	2.0	5.4	10.7	20.7	54.0	156.6	628.9
Non-Hispanic black, male												
2006 5/	13.5	15.7	*	*	0.7	1.4	3.7	11.6	21.8	59.3	180.5	509.4
2005 5/	15.0	14.0	*	*	0.7	1.7	4.8	13.0	27.7	71.0	190.6	530.0
2004 5/	15.4	15.8	*	*	*	1.6	4.9	12.0	24.7	72.7	218.2	606.6
2003 5/	16.2	15.2	*	*	0.7	2.3	5.4	13.0	29.4	78.5	219.9	639.3
2002	15.8	13.9	1.6	*	*	1.8	5.2	11.5	29.3	66.0	230.6	693.6
2001	16.3	16.6	*	*	0.8	2.0	5.9	13.5	26.4	69.7	242.5	716.1
2000	17.1	18.8	1.7	*	0.7	2.3	5.7	14.0	30.1	83.4	233.9	750.3
1999	16.8	19.1	2.0	*	0.8	2.3	6.6	15.1	27.2	77.0	213.3	799.9
Non-Hispanic black, female												
2006 5/	14.0	10.0	*	*	*	1.4	3.0	6.6	11.7	35.2	117.4	408.4
2005 5/	15.3	14.6	*	*	*	1.2	2.9	7.3	14.0	40.0	127.6	447.7
2004 5/	14.8	12.2	1.7	*	0.7	0.8	3.3	6.9	14.3	33.8	122.0	470.6
2003 5/	15.0	11.3	*	*	*	1.5	3.7	7.8	13.9	37.2	120.6	477.1
2002	16.1	12.3	*	*	*	1.4	3.9	6.6	15.1	39.5	131.5	533.0
2001	15.6	17.1	*	*	0.8	1.3	4.0	6.9	13.0	33.9	127.9	531.7
2000	16.3	15.5	*	*	0.8	1.2	3.5	6.8	15.6	43.3	135.3	537.3
1999	16.3	19.6	*	*	0.9	1.7	4.4	6.9	15.5	37.7	124.8	565.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
All origins 3/, both sexes												
2006	0.1	1.3	*	*	*	*	*	*	*	0.1	0.5	2.0
2005	0.1	1.2	*	*	*	*	*	*	*	0.1	0.6	3.3
2004	0.1	1.5	*	*	*	*	*	*	*	0.1	0.4	2.6
2003	0.1	1.3	0.2	*	*	*	*	*	0.1	0.2	0.6	3.5
2002	0.1	1.3	0.1	*	*	*	*	*	*	0.2	0.5	3.3
2001	0.1	1.2	*	*	*	*	*	*	*	0.2	0.5	3.3
2000	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.0
All origins 3/, male												
2006	0.1	1.4	*	*	*	*	*	*	*	*	0.4	1.8
2005	0.1	1.4	*	*	*	*	*	*	*	*	0.8	3.1
2004	0.1	1.6	*	*	*	*	*	*	*	*	*	2.5
2003	0.1	1.7	*	*	*	*	*	*	*	*	0.7	2.4
2002	0.1	1.9	*	*	*	*	*	*	*	*	0.5	2.8
2001	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2000	0.1	1.3	*	*	*	*	*	*	*	0.3	0.7	3.8
1999	0.1	1.9	*	*	*	*	*	*	*	*	1.1	4.0
All origins 3/, female												
2006	0.1	1.1	*	*	*	*	*	*	*	*	0.5	2.1
2005	0.2	1.0	*	*	*	*	*	*	*	*	0.5	3.5
2004	0.1	1.3	*	*	*	*	*	*	*	*	0.5	2.7
2003	0.2	*	*	*	*	*	*	*	*	*	0.6	3.9
2002	0.2	*	*	*	*	*	*	*	*	0.2	0.5	3.5
2001	0.2	1.2	*	*	*	*	*	*	*	*	0.5	3.7
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.8
1999	0.2	1.5	*	*	*	*	*	*	*	*	0.8	6.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
Non-Hispanic 4/, both sexes												
2006	0.1	1.4	*	*	*	*	*	*	*	0.1	0.5	2.1
2005	0.2	1.4	*	*	*	*	*	*	*	0.1	0.6	3.4
2004	0.1	1.4	*	*	*	*	*	*	*	*	0.4	2.6
2003	0.2	1.2	0.2	*	*	*	*	*	*	0.2	0.6	3.6
2002	0.1	1.3	0.2	*	*	*	*	*	*	0.2	0.5	3.4
2001	0.1	1.3	*	*	*	*	*	*	*	0.2	0.4	3.3
2000	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.1
Non-Hispanic 4/, male												
2006	0.1	1.5	*	*	*	*	*	*	*	*	0.4	1.9
2005	0.1	1.7	*	*	*	*	*	*	*	*	0.8	3.2
2004	0.1	1.6	*	*	*	*	*	*	*	*	*	2.5
2003	0.1	1.4	*	*	*	*	*	*	*	*	0.7	2.4
2002	0.1	2.1	*	*	*	*	*	*	*	*	0.6	2.9
2001	0.1	1.2	*	*	*	*	*	*	*	*	*	2.4
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.7	3.7
1999	0.2	2.0	*	*	*	*	*	*	*	*	1.1	4.1
Non-Hispanic 4/, female												
2006	0.1	1.4	*	*	*	*	*	*	*	*	0.5	2.2
2005	0.2	*	*	*	*	*	*	*	*	*	0.5	3.6
2004	0.1	*	*	*	*	*	*	*	*	*	0.5	2.7
2003	0.2	*	*	*	*	*	*	*	*	*	0.6	4.1
2002	0.2	*	*	*	*	*	*	*	*	0.2	0.5	3.6
2001	0.2	1.4	*	*	*	*	*	*	*	*	0.5	3.7
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.7	4.8
1999	0.3	1.5	*	*	*	*	*	*	*	*	0.8	6.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
Non-Hispanic white, both sexes												
2006 5/	0.1	1.2	*	*	*	*	*	*	*	*	0.5	2.3
2005 5/	0.2	1.0	*	*	*	*	*	*	*	0.1	0.7	3.6
2004 5/	0.1	1.2	*	*	*	*	*	*	*	*	0.4	2.8
2003 5/	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	3.8
2002	0.2	1.1	*	*	*	*	*	*	*	0.2	0.5	3.5
2001	0.1	1.0	*	*	*	*	*	*	*	0.1	0.5	3.5
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.7	4.7
1999	0.2	1.1	*	*	*	*	*	*	*	0.2	0.9	6.4
Non-Hispanic white, male												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	2.0
2005 5/	0.1	*	*	*	*	*	*	*	*	*	0.9	3.4
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	2.6
2003 5/	0.1	*	*	*	*	*	*	*	*	*	0.7	2.7
2002	0.1	1.8	*	*	*	*	*	*	*	*	0.6	3.0
2001	0.1	*	*	*	*	*	*	*	*	*	*	2.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.7	3.7
1999	0.2	*	*	*	*	*	*	*	*	*	1.1	4.1
Non-Hispanic white, female												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	0.6	2.4
2005 5/	0.2	*	*	*	*	*	*	*	*	*	0.5	3.7
2004 5/	0.1	*	*	*	*	*	*	*	*	*	0.5	2.9
2003 5/	0.2	*	*	*	*	*	*	*	*	*	0.6	4.3
2002	0.2	*	*	*	*	*	*	*	*	*	0.5	3.7
2001	0.2	*	*	*	*	*	*	*	*	*	0.5	3.9
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.7	5.1
1999	0.3	*	*	*	*	*	*	*	*	*	0.8	7.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
Non-Hispanic black, both sexes												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	4.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
All origins 3/, both sexes												
2006	0.1	1.3	*	*	*	*	*	*	*	*	0.3	1.1
2005	0.1	1.2	*	*	*	*	*	*	*	*	0.4	1.9
2004	0.1	1.4	*	*	*	*	*	*	*	*	0.2	1.4
2003	0.1	1.3	0.2	*	*	*	*	*	*	0.1	0.3	2.1
2002	0.1	1.2	0.1	*	*	*	*	*	*	0.1	0.3	2.0
2001	0.1	1.2	*	*	*	*	*	*	*	0.1	0.3	2.1
2000	0.1	1.1	*	*	*	*	*	*	*	0.2	0.4	2.6
1999	0.1	1.6	0.2	*	*	*	*	*	*	0.2	0.6	3.6
All origins 3/, male												
2006	0.1	1.4	*	*	*	*	*	*	*	*	*	1.2
2005	0.1	1.4	*	*	*	*	*	*	*	*	0.5	1.7
2004	0.1	1.6	*	*	*	*	*	*	*	*	*	1.8
2003	0.1	1.7	*	*	*	*	*	*	*	*	*	*
2002	0.1	1.8	*	*	*	*	*	*	*	*	*	1.7
2001	0.1	1.2	*	*	*	*	*	*	*	*	*	*
2000	0.1	1.2	*	*	*	*	*	*	*	*	0.6	2.4
1999	0.1	1.9	*	*	*	*	*	*	*	*	0.7	2.6
All origins 3/, female												
2006	0.1	1.1	*	*	*	*	*	*	*	*	0.3	1.1
2005	0.1	1.0	*	*	*	*	*	*	*	*	0.3	2.0
2004	0.1	1.3	*	*	*	*	*	*	*	*	0.3	1.3
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2001	0.1	1.2	*	*	*	*	*	*	*	*	0.4	2.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.7
1999	0.2	1.4	*	*	*	*	*	*	*	*	0.5	4.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
Non-Hispanic 4/, both sexes												
2006	0.1	1.4	*	*	*	*	*	*	*	*	0.3	1.2
2005	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.0
2004	0.1	1.4	*	*	*	*	*	*	*	*	0.3	1.4
2003	0.1	1.1	0.2	*	*	*	*	*	*	0.1	0.4	2.2
2002	0.1	1.3	0.2	*	*	*	*	*	*	0.1	0.3	2.1
2001	0.1	1.3	*	*	*	*	*	*	*	0.1	0.3	2.1
2000	0.1	1.1	*	*	*	*	*	*	*	0.2	0.4	2.6
1999	0.1	1.7	0.2	*	*	*	*	*	*	0.2	0.6	3.6
Non-Hispanic 4/, male												
2006	0.1	1.5	*	*	*	*	*	*	*	*	*	1.3
2005	0.1	1.6	*	*	*	*	*	*	*	*	0.5	1.7
2004	0.1	1.6	*	*	*	*	*	*	*	*	*	1.7
2003	0.1	1.4	*	*	*	*	*	*	*	*	*	*
2002	0.1	2.0	*	*	*	*	*	*	*	*	*	1.7
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
1999	0.1	2.0	*	*	*	*	*	*	*	*	0.7	2.6
Non-Hispanic 4/, female												
2006	0.1	1.4	*	*	*	*	*	*	*	*	0.3	1.1
2005	0.1	*	*	*	*	*	*	*	*	*	0.4	2.1
2004	0.1	*	*	*	*	*	*	*	*	*	0.3	1.3
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	2.6
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2001	0.1	1.4	*	*	*	*	*	*	*	*	0.4	2.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	2.7
1999	0.2	1.4	*	*	*	*	*	*	*	*	0.5	4.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
Non-Hispanic white, both sexes												
2006 5/	0.1	1.2	*	*	*	*	*	*	*	*	0.3	1.2
2005 5/	0.1	1.0	*	*	*	*	*	*	*	*	0.5	2.0
2004 5/	0.1	1.2	*	*	*	*	*	*	*	*	0.2	1.5
2003 5/	0.1	1.0	*	*	*	*	*	*	*	*	0.4	2.3
2002	0.1	1.0	*	*	*	*	*	*	*	0.1	0.3	2.1
2001	0.1	1.0	*	*	*	*	*	*	*	*	0.3	2.2
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.4	2.7
1999	0.2	1.1	*	*	*	*	*	*	*	0.1	0.6	3.9
Non-Hispanic white, male												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	0.6	1.8
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	1.8
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	1.7	*	*	*	*	*	*	*	*	*	1.7
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.6	2.3
1999	0.1	*	*	*	*	*	*	*	*	*	0.8	2.6
Non-Hispanic white, female												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2005 5/	0.1	*	*	*	*	*	*	*	*	*	0.4	2.2
2004 5/	0.1	*	*	*	*	*	*	*	*	*	0.3	1.4
2003 5/	0.1	*	*	*	*	*	*	*	*	*	0.4	2.8
2002	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	2.8
1999	0.2	*	*	*	*	*	*	*	*	*	0.5	4.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
Non-Hispanic black, both sexes												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	4.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
All origins 3/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	0.2	0.9
2005	0.0	*	*	*	*	*	*	*	*	*	0.2	1.4
2004	0.0	*	*	*	*	*	*	*	*	*	0.2	1.2
2003	0.0	*	*	*	*	*	*	*	*	*	0.3	1.3
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	1.3
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	1.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.4
All origins 3/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	*	1.1
2005	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2004	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2003	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.3
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
Non-Hispanic 4/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	0.2	1.0
2005	0.0	*	*	*	*	*	*	*	*	*	0.2	1.5
2004	0.0	*	*	*	*	*	*	*	*	*	0.2	1.2
2003	0.0	*	*	*	*	*	*	*	*	*	0.3	1.4
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	1.3
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.2	1.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.4
Non-Hispanic 4/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	*	1.1
2005	0.1	*	*	*	*	*	*	*	*	*	*	1.5
2004	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2003	0.1	*	*	*	*	*	*	*	*	*	*	1.5
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.3
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
Non-Hispanic white, both sexes												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	0.2	1.1
2005 5/	0.1	*	*	*	*	*	*	*	*	*	0.2	1.6
2004 5/	0.0	*	*	*	*	*	*	*	*	*	0.2	1.3
2003 5/	0.1	*	*	*	*	*	*	*	*	*	0.3	1.5
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	1.4
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.5
Non-Hispanic white, male												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	1.6
2004 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	1.3
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	1.6
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	1.6
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	1.6
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2001	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	3.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
All origins 3/, both sexes												
2006	41.6	0.7	0.3	0.3	0.4	0.6	1.9	9.1	39.2	149.3	363.4	589.1
2005	44.2	0.8	0.3	0.3	0.4	0.6	2.0	9.4	42.0	160.5	385.6	637.2
2004	41.5	0.9	0.3	0.3	0.4	0.6	2.0	8.4	40.4	153.8	366.7	601.7
2003	43.5	0.8	0.3	0.3	0.5	0.7	2.1	8.7	43.3	163.2	383.0	635.1
2002	43.3	1.0	0.4	0.3	0.5	0.8	2.2	8.7	42.4	163.0	386.7	637.6
2001	43.2	1.0	0.3	0.3	0.4	0.7	2.2	8.5	44.1	167.9	379.8	644.7
2000	43.4	0.9	0.3	0.3	0.5	0.7	2.1	8.6	44.2	169.4	386.1	648.6
1999	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0
All origins 3/, male												
2006	40.2	0.9	0.3	0.4	0.4	0.7	1.8	9.2	41.9	163.2	437.3	760.8
2005	42.8	*	0.4	0.3	0.4	0.6	1.9	9.7	45.0	180.7	463.7	819.9
2004	40.6	1.2	0.4	0.4	0.5	0.7	2.0	8.8	43.1	172.1	445.6	811.1
2003	42.4	1.1	0.5	0.4	0.5	0.8	1.9	9.1	46.5	183.6	464.9	865.9
2002	42.9	1.1	0.6	0.4	0.6	0.8	2.2	9.1	45.2	184.8	480.8	894.8
2001	42.7	1.1	0.4	0.3	0.5	0.7	2.0	8.8	46.9	191.3	475.1	916.9
2000	43.5	1.2	0.4	0.4	0.6	0.8	1.9	9.0	47.8	195.2	488.5	967.9
1999	45.6	*	0.4	0.4	0.6	0.8	1.8	8.6	52.3	210.7	513.2	996.7
All origins 3/, female												
2006	43.0	*	*	0.2	0.3	0.6	2.0	8.9	36.6	137.5	312.8	508.7
2005	45.5	*	0.3	0.2	0.3	0.6	2.1	9.0	39.2	143.4	332.5	553.3
2004	42.5	*	*	0.2	0.3	0.5	2.1	8.1	38.0	138.4	313.6	507.4
2003	44.4	*	*	0.2	0.4	0.6	2.3	8.2	40.3	146.0	328.3	533.0
2002	43.7	*	0.3	0.3	0.4	0.7	2.3	8.2	39.8	144.9	324.1	526.0
2001	43.7	*	*	0.2	0.4	0.7	2.3	8.1	41.5	148.5	317.3	530.8
2000	43.2	*	0.3	0.3	0.4	0.7	2.2	8.3	41.0	148.2	319.2	518.5
1999	43.4	*	0.3	0.2	0.5	0.9	2.1	8.4	43.1	149.8	322.9	504.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
Hispanic, both sexes												
2006	7.5	*	*	*	*	0.3	0.7	2.6	12.0	45.5	153.5	374.4
2005	8.1	*	*	*	*	0.3	0.8	3.5	12.5	50.6	171.9	414.6
2004	7.5	*	*	*	*	0.4	1.1	3.1	12.2	51.7	158.7	391.5
2003	8.0	*	*	*	0.3	0.4	1.0	3.5	12.7	53.3	171.0	460.2
2002	7.9	*	*	*	*	0.4	1.2	3.5	13.9	53.8	176.0	457.3
2001	7.7	*	*	*	*	0.3	1.0	3.6	14.0	55.8	169.5	480.2
2000	7.6	*	*	*	*	0.3	1.1	3.3	14.3	53.1	181.9	481.1
1999	8.4	*	*	*	*	0.5	1.6	3.9	14.6	62.6	194.8	536.0
Hispanic, male												
2006	7.5	*	*	*	*	*	0.7	2.8	14.0	56.6	196.3	484.9
2005	8.3	*	*	*	*	*	1.0	3.6	15.8	64.5	234.2	526.8
2004	7.6	*	*	*	*	0.5	1.4	2.8	13.7	68.4	206.4	518.4
2003	8.2	*	*	*	*	*	1.0	3.2	16.6	68.1	231.2	646.5
2002	8.1	*	*	*	*	*	1.0	3.8	17.5	69.2	243.3	602.4
2001	7.8	*	*	*	*	*	0.7	3.2	16.1	75.5	224.0	676.1
2000	8.0	*	*	*	*	*	0.9	3.4	18.2	72.4	250.3	671.1
1999	8.9	*	*	*	*	*	1.4	3.6	17.6	81.9	272.7	836.8
Hispanic, female												
2006	7.5	*	*	*	*	*	0.7	2.5	10.1	36.5	123.5	315.7
2005	7.8	*	*	*	*	*	*	3.5	9.5	39.3	128.6	355.8
2004	7.4	*	*	*	*	*	0.7	3.4	10.9	38.3	125.7	326.3
2003	7.7	*	*	*	*	*	1.0	3.8	9.3	41.5	129.6	365.6
2002	7.6	*	*	*	*	*	1.4	3.1	10.6	41.5	129.8	385.5
2001	7.5	*	*	*	*	*	1.2	4.1	12.1	40.3	132.7	384.4
2000	7.2	*	*	*	*	*	1.3	3.3	10.8	38.0	136.0	387.8
1999	8.0	*	*	*	*	*	1.8	4.2	12.0	47.5	142.9	391.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
Non-Hispanic 4/, both sexes												
2006	47.5	0.8	0.3	0.3	0.4	0.7	2.1	9.8	41.4	157.1	375.9	599.3
2005	50.2	0.8	0.4	0.3	0.4	0.7	2.2	10.0	44.4	168.5	397.6	647.1
2004	47.0	0.8	0.3	0.3	0.5	0.7	2.2	9.0	42.7	160.9	377.8	610.3
2003	49.0	0.8	0.3	0.3	0.5	0.8	2.3	9.1	45.6	170.6	393.3	641.2
2002	48.6	1.1	0.4	0.4	0.5	0.8	2.4	9.2	44.6	170.1	396.0	643.5
2001	48.4	1.1	0.3	0.3	0.5	0.8	2.3	8.9	46.3	174.8	388.6	649.4
2000	48.3	1.0	0.3	0.4	0.5	0.8	2.2	9.0	46.3	176.1	393.8	653.2
1999	49.4	1.0	0.4	0.4	0.6	0.9	2.0	8.9	49.8	183.5	405.1	648.0
Non-Hispanic 4/, male												
2006	46.1	*	0.3	0.4	0.5	0.8	2.0	9.9	44.1	171.0	451.8	775.2
2005	48.8	*	0.5	0.3	0.5	0.7	2.1	10.4	47.3	188.9	476.6	833.9
2004	46.2	*	0.4	0.4	0.6	0.8	2.1	9.5	45.3	178.9	458.4	824.6
2003	48.1	*	0.5	0.4	0.6	0.9	2.1	9.7	48.7	191.1	476.2	874.6
2002	48.4	1.2	0.6	0.4	0.6	0.9	2.4	9.6	47.3	191.9	491.2	906.0
2001	48.0	*	0.4	0.3	0.6	0.8	2.2	9.3	49.0	198.1	485.7	924.8
2000	48.7	*	0.4	0.4	0.6	0.9	2.1	9.4	49.7	201.9	497.6	978.0
1999	50.8	*	0.4	0.5	0.7	0.8	1.9	9.0	54.5	217.4	521.9	1,000.2
Non-Hispanic 4/, female												
2006	48.8	*	*	0.2	0.3	0.6	2.2	9.7	38.8	145.3	324.0	517.5
2005	51.5	*	0.3	0.2	0.3	0.7	2.3	9.6	41.6	151.3	344.0	561.8
2004	47.8	*	*	0.2	0.4	0.6	2.3	8.5	40.2	145.6	323.6	514.5
2003	49.9	*	*	0.2	0.4	0.7	2.5	8.6	42.7	153.3	338.0	538.6
2002	48.9	*	*	0.3	0.4	0.8	2.4	8.7	42.0	151.8	332.9	530.3
2001	48.7	*	*	0.2	0.4	0.8	2.4	8.5	43.8	155.4	324.9	534.8
2000	48.0	*	*	0.3	0.4	0.7	2.3	8.7	43.2	154.8	326.2	521.9
1999	48.0	*	*	0.2	0.5	0.9	2.1	8.7	45.4	155.7	329.3	507.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
Non-Hispanic white, both sexes												
2006 5/	55.6	*	*	0.1	0.2	0.5	1.9	9.9	43.7	169.7	398.4	629.8
2005 5/	58.6	*	0.2	0.1	0.2	0.5	2.0	9.8	46.9	181.1	420.9	679.0
2004 5/	54.9	*	*	0.2	0.3	0.5	2.0	8.8	44.8	173.3	399.6	640.0
2003 5/	57.0	*	*	0.2	0.3	0.6	2.0	9.0	47.8	182.6	414.9	671.6
2002	56.4	0.9	0.2	0.2	0.4	0.6	2.0	9.0	46.7	181.6	416.3	673.1
2001	55.9	*	*	0.2	0.3	0.5	1.9	8.7	48.6	185.9	408.6	676.9
2000	55.7	*	*	0.2	0.4	0.6	1.8	8.6	48.2	186.5	413.2	680.8
1999	56.6	*	0.2	0.2	0.4	0.6	1.6	8.5	51.8	193.6	422.4	675.6
Non-Hispanic white, male												
2006 5/	53.0	*	*	0.2	0.2	0.5	1.8	9.8	45.4	180.5	469.7	804.2
2005 5/	56.0	*	*	*	0.3	0.5	1.9	10.1	48.9	198.7	493.9	861.9
2004 5/	53.0	*	*	0.2	0.3	0.5	1.8	9.2	46.3	188.5	476.6	852.5
2003 5/	54.9	*	*	0.2	0.4	0.5	1.8	9.5	50.0	200.2	491.0	903.6
2002	55.1	*	*	0.2	0.4	0.6	2.0	9.3	48.3	200.4	506.7	935.4
2001	54.6	*	*	0.2	0.3	0.5	1.9	9.1	50.5	206.1	500.9	951.5
2000	55.1	*	*	0.2	0.4	0.5	1.7	8.9	50.8	208.8	513.6	1,008.6
1999	57.3	*	*	0.3	0.5	0.5	1.5	8.5	55.6	224.9	534.9	1,033.5
Non-Hispanic white, female												
2006 5/	58.0	*	*	*	0.2	0.5	2.0	9.9	42.1	160.4	348.9	548.5
2005 5/	61.1	*	*	*	0.2	0.6	2.2	9.5	44.9	165.8	370.7	595.3
2004 5/	56.7	*	*	*	0.2	0.5	2.1	8.4	43.5	160.1	347.2	544.8
2003 5/	59.0	*	*	*	0.3	0.6	2.2	8.5	45.7	167.6	363.5	569.5
2002	57.7	*	*	*	0.4	0.6	2.1	8.6	45.1	165.5	355.7	559.8
2001	57.2	*	*	*	0.3	0.6	2.0	8.3	46.8	168.8	347.3	562.7
2000	56.2	*	*	0.2	0.3	0.6	1.9	8.3	45.8	167.6	347.2	548.7
1999	55.9	*	*	*	0.4	0.6	1.8	8.5	48.3	167.4	348.8	532.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
Non-Hispanic black, both sexes												
2006 5/	20.3	*	*	1.1	1.2	2.0	4.1	12.3	38.1	102.9	229.9	337.4
2005 5/	21.8	*	1.2	0.8	1.2	1.8	4.0	14.2	40.3	116.4	244.8	369.2
2004 5/	19.9	*	0.8	1.0	1.3	1.9	4.0	13.0	39.1	103.9	225.9	335.9
2003 5/	20.9	*	1.0	1.0	1.3	2.0	4.4	12.9	41.3	115.3	238.5	354.9
2002	21.4	*	1.2	1.1	1.3	2.0	5.2	13.1	41.6	115.0	250.5	370.8
2001	21.1	*	0.9	0.8	1.3	2.5	4.9	12.7	41.3	120.4	236.3	379.3
2000	21.5	*	0.9	1.1	1.4	2.2	4.9	14.5	44.7	123.2	236.5	376.2
1999	22.6	*	*	1.0	1.4	2.7	4.7	14.1	46.7	129.9	265.3	377.0
Non-Hispanic black, male												
2006 5/	22.7	*	*	1.3	1.4	2.4	4.0	13.6	48.8	139.7	345.1	522.3
2005 5/	24.7	*	1.6	1.0	1.5	2.0	4.0	15.5	49.9	160.1	381.2	596.6
2004 5/	22.8	*	*	1.4	1.6	2.6	4.2	14.4	51.4	142.5	342.3	569.0
2003 5/	24.2	*	*	1.3	1.6	2.9	4.1	13.7	51.5	158.0	386.7	608.4
2002	24.9	*	1.6	1.2	2.0	2.3	5.7	14.6	53.1	160.0	395.7	653.1
2001	24.5	*	*	1.0	1.7	2.8	4.7	13.5	50.4	170.1	385.6	682.6
2000	25.1	*	*	1.3	1.8	2.8	4.9	15.2	55.3	178.9	372.4	696.3
1999	27.0	*	*	1.4	1.9	2.7	4.7	15.6	60.4	186.1	438.4	707.2
Non-Hispanic black, female												
2006 5/	18.1	*	*	0.8	0.9	1.6	4.2	11.1	29.5	76.6	164.6	262.0
2005 5/	19.1	*	*	*	0.9	1.6	3.9	13.1	32.5	85.1	167.5	277.7
2004 5/	17.2	*	*	0.6	1.0	1.3	3.9	11.9	29.3	76.3	160.2	243.8
2003 5/	17.9	*	*	*	1.1	1.2	4.7	12.2	33.2	85.0	154.8	256.7
2002	18.2	*	*	1.1	0.7	1.7	4.8	11.8	32.3	83.0	168.5	263.0
2001	18.0	*	*	0.7	1.0	2.2	5.0	11.9	34.0	85.3	152.5	266.9
2000	18.2	*	*	0.8	1.0	1.6	4.9	13.8	36.1	84.1	159.9	257.3
1999	18.6	*	*	0.6	0.9	2.7	4.7	12.8	35.8	90.3	168.4	253.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
All origins 3/, both sexes												
2006	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.5	6.0
2005	0.3	0.6	*	*	*	*	*	0.1	0.2	0.6	1.6	7.4
2004	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.5	6.5
2003	0.3	0.5	*	*	*	*	*	0.1	0.2	0.6	1.9	7.3
2002	0.3	0.6	*	*	*	*	*	0.1	0.2	0.7	2.2	8.1
2001	0.3	0.5	*	*	*	*	0.0	0.1	0.2	0.7	2.1	9.1
2000	0.4	*	*	*	*	*	0.0	0.1	0.3	0.9	2.7	11.8
1999	0.4	0.5	*	*	*	*	*	0.1	0.4	0.9	2.8	10.7
All origins 3/, male												
2006	0.2	*	*	*	*	*	*	*	0.2	0.5	1.5	5.9
2005	0.2	*	*	*	*	*	*	*	0.2	0.6	1.9	7.3
2004	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.6	7.0
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.6	2.2	7.0
2002	0.3	*	*	*	*	*	*	0.1	0.2	0.9	2.5	8.8
2001	0.3	*	*	*	*	*	*	0.1	0.3	0.9	2.3	9.2
2000	0.4	*	*	*	*	*	*	*	0.3	1.0	3.6	12.9
1999	0.4	*	*	*	*	*	*	*	0.4	1.1	3.3	11.9
All origins 3/, female												
2006	0.3	*	*	*	*	*	*	*	0.2	0.5	1.5	6.0
2005	0.3	*	*	*	*	*	*	*	0.2	0.7	1.4	7.4
2004	0.3	*	*	*	*	*	*	*	0.2	0.4	1.5	6.3
2003	0.3	*	*	*	*	*	*	*	0.1	0.6	1.7	7.5
2002	0.4	*	*	*	*	*	*	*	0.2	0.6	2.1	7.8
2001	0.4	*	*	*	*	*	*	0.1	0.2	0.6	1.9	9.1
2000	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.1	11.4
1999	0.5	*	*	*	*	*	*	*	0.4	0.8	2.5	10.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
Hispanic, both sexes												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	11.1
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
Non-Hispanic 4/, both sexes												
2006	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.5	6.0
2005	0.3	0.6	*	*	*	*	*	0.1	0.3	0.7	1.6	7.3
2004	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.5	6.6
2003	0.3	*	*	*	*	*	*	0.1	0.2	0.6	1.9	7.3
2002	0.4	0.6	*	*	*	*	*	0.1	0.2	0.8	2.2	8.0
2001	0.4	*	*	*	*	*	*	0.1	0.2	0.8	2.1	9.1
2000	0.5	*	*	*	*	*	*	0.1	0.3	0.9	2.7	11.8
1999	0.5	*	*	*	*	*	*	0.1	0.4	1.0	2.8	10.7
Non-Hispanic 4/, male												
2006	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	6.1
2005	0.3	*	*	*	*	*	*	*	0.3	0.6	1.9	7.4
2004	0.3	*	*	*	*	*	*	0.1	0.3	0.7	1.6	7.0
2003	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.2	6.9
2002	0.3	*	*	*	*	*	*	0.1	0.2	0.9	2.5	8.8
2001	0.3	*	*	*	*	*	*	0.1	0.3	1.0	2.3	9.3
2000	0.4	*	*	*	*	*	*	*	0.3	0.9	3.6	12.8
1999	0.4	*	*	*	*	*	*	*	0.5	1.1	3.3	11.9
Non-Hispanic 4/, female												
2006	0.3	*	*	*	*	*	*	*	0.2	0.5	1.5	6.0
2005	0.4	*	*	*	*	*	*	*	0.3	0.7	1.3	7.2
2004	0.3	*	*	*	*	*	*	*	0.2	0.4	1.5	6.4
2003	0.4	*	*	*	*	*	*	*	0.1	0.6	1.7	7.5
2002	0.4	*	*	*	*	*	*	*	0.2	0.6	2.0	7.6
2001	0.4	*	*	*	*	*	*	*	0.2	0.6	2.0	9.1
2000	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.1	11.4
1999	0.5	*	*	*	*	*	*	*	0.4	0.8	2.4	10.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
Non-Hispanic white, both sexes												
2006 5/	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.6	6.3
2005 5/	0.4	*	*	*	*	*	*	0.1	0.2	0.7	1.6	7.8
2004 5/	0.3	*	*	*	*	*	*	0.1	0.2	0.6	1.6	6.7
2003 5/	0.4	*	*	*	*	*	*	0.1	0.2	0.6	2.0	7.5
2002	0.4	*	*	*	*	*	*	0.1	0.2	0.8	2.3	8.4
2001	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.2	9.7
2000	0.5	*	*	*	*	*	*	0.1	0.3	0.9	2.8	12.6
1999	0.5	*	*	*	*	*	*	0.1	0.5	1.0	2.8	11.2
Non-Hispanic white, male												
2006 5/	0.2	*	*	*	*	*	*	*	*	0.5	1.5	6.5
2005 5/	0.3	*	*	*	*	*	*	*	0.3	0.6	1.9	7.5
2004 5/	0.3	*	*	*	*	*	*	*	0.3	0.7	1.7	7.2
2003 5/	0.3	*	*	*	*	*	*	*	0.2	0.6	2.2	7.1
2002	0.4	*	*	*	*	*	*	*	0.2	1.0	2.6	9.3
2001	0.3	*	*	*	*	*	*	*	0.3	1.0	2.3	9.6
2000	0.4	*	*	*	*	*	*	*	0.3	0.9	3.8	13.8
1999	0.4	*	*	*	*	*	*	*	0.5	1.2	3.2	12.4
Non-Hispanic white, female												
2006 5/	0.4	*	*	*	*	*	*	*	0.2	0.5	1.6	6.2
2005 5/	0.4	*	*	*	*	*	*	*	0.2	0.8	1.4	7.9
2004 5/	0.4	*	*	*	*	*	*	*	0.2	0.5	1.6	6.5
2003 5/	0.4	*	*	*	*	*	*	*	*	0.7	1.8	7.6
2002	0.5	*	*	*	*	*	*	*	0.2	0.7	2.1	8.0
2001	0.5	*	*	*	*	*	*	*	0.2	0.6	2.1	9.7
2000	0.6	*	*	*	*	*	*	*	0.3	0.8	2.1	12.2
1999	0.6	*	*	*	*	*	*	*	0.4	0.8	2.5	10.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
Non-Hispanic black, both sexes												
2006 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	2.2	*
1999	0.3	*	*	*	*	*	*	*	*	*	2.9	*
Non-Hispanic black, male												
2006 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
All origins 3/, both sexes												
2006	4.2	*	*	*	*	*	0.2	1.0	4.8	17.3	36.8	45.3
2005	4.7	*	*	*	*	*	0.2	1.0	5.5	20.1	41.4	51.9
2004	4.6	*	*	*	*	*	0.2	1.0	5.4	20.5	41.0	49.9
2003	5.1	*	*	*	*	*	0.2	1.1	6.3	22.2	46.0	54.6
2002	5.4	*	*	*	*	*	0.2	1.1	6.4	24.0	48.0	59.3
2001	5.7	*	*	*	*	*	0.2	1.1	7.2	26.2	50.4	62.1
2000	5.9	*	*	*	*	*	0.2	1.3	7.4	27.8	52.7	63.3
1999	6.4	*	*	*	*	*	0.2	1.3	8.5	30.7	56.9	62.1
All origins 3/, male												
2006	4.3	*	*	*	*	*	0.2	1.3	5.3	19.6	45.4	63.7
2005	5.0	*	*	*	*	*	0.2	1.3	6.7	23.9	52.4	71.5
2004	4.9	*	*	*	*	*	0.2	1.2	6.2	24.3	52.2	74.4
2003	5.3	*	*	*	*	*	0.2	1.4	6.9	25.6	58.1	79.2
2002	5.7	*	*	*	*	*	0.3	1.4	7.2	28.1	63.5	90.0
2001	6.0	*	*	*	*	*	0.2	1.4	7.9	30.8	65.3	101.2
2000	6.3	*	*	*	*	*	0.2	1.5	8.1	33.0	69.1	104.2
1999	6.9	*	*	*	*	*	0.2	1.5	9.5	37.8	76.4	105.4
All origins 3/, female												
2006	4.1	*	*	*	*	*	0.2	0.9	4.4	15.4	30.9	36.7
2005	4.5	*	*	*	*	*	0.1	0.8	4.5	16.9	33.9	42.8
2004	4.4	*	*	*	*	*	0.1	0.8	4.7	17.4	33.5	38.9
2003	4.9	*	*	*	*	*	0.2	0.8	5.7	19.2	37.9	43.7
2002	5.0	*	*	*	*	*	0.1	0.9	5.6	20.6	37.6	46.0
2001	5.4	*	*	*	*	*	0.1	0.9	6.6	22.5	40.6	45.8
2000	5.6	*	*	*	*	*	0.2	1.0	6.8	23.6	41.9	46.7
1999	5.8	*	*	*	*	*	0.2	1.1	7.5	24.9	44.2	44.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
Hispanic, both sexes												
2006	0.6	*	*	*	*	*	*	*	1.7	4.0	15.6	20.7
2005	0.7	*	*	*	*	*	*	*	1.3	5.4	16.6	24.8
2004	0.7	*	*	*	*	*	*	*	0.9	5.0	16.1	30.0
2003	0.8	*	*	*	*	*	*	*	1.4	6.0	20.0	34.7
2002	0.8	*	*	*	*	*	*	*	1.5	6.7	18.8	36.6
2001	0.8	*	*	*	*	*	*	*	1.4	6.6	20.4	39.4
2000	0.8	*	*	*	*	*	*	*	1.5	6.0	22.3	31.2
1999	0.9	*	*	*	*	*	*	*	1.3	7.1	23.5	44.2
Hispanic, male												
2006	0.8	*	*	*	*	*	*	*	2.2	6.2	21.8	31.6
2005	0.8	*	*	*	*	*	*	*	1.8	8.6	22.7	38.6
2004	0.7	*	*	*	*	*	*	*	*	6.5	20.4	47.6
2003	0.9	*	*	*	*	*	*	*	2.1	7.6	29.1	53.9
2002	0.8	*	*	*	*	*	*	*	*	9.2	26.7	53.5
2001	0.9	*	*	*	*	*	*	*	*	9.2	34.0	60.9
2000	0.9	*	*	*	*	*	*	*	*	9.1	32.5	58.4
1999	1.0	*	*	*	*	*	*	*	*	11.2	36.0	77.6
Hispanic, female												
2006	0.5	*	*	*	*	*	*	*	*	*	11.3	15.0
2005	0.6	*	*	*	*	*	*	*	*	2.9	12.3	17.6
2004	0.6	*	*	*	*	*	*	*	*	3.7	13.1	20.9
2003	0.7	*	*	*	*	*	*	*	*	4.7	13.7	25.0
2002	0.7	*	*	*	*	*	*	*	*	4.7	13.5	28.2
2001	0.6	*	*	*	*	*	*	*	*	4.6	11.2	28.9
2000	0.6	*	*	*	*	*	*	*	*	3.6	15.5	*
1999	0.7	*	*	*	*	*	*	*	*	4.0	15.3	28.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
Non-Hispanic 4/, both sexes												
2006	4.8	*	*	*	*	*	0.2	1.1	5.1	18.3	38.1	46.5
2005	5.4	*	*	*	*	*	0.2	1.1	5.9	21.2	42.8	53.1
2004	5.3	*	*	*	*	*	0.2	1.1	5.8	21.7	42.4	50.7
2003	5.8	*	*	*	*	*	0.2	1.2	6.6	23.3	47.3	55.4
2002	6.1	*	*	*	*	*	0.2	1.2	6.8	25.1	49.3	60.1
2001	6.4	*	*	*	*	*	0.2	1.2	7.6	27.5	51.7	62.8
2000	6.7	*	*	*	*	*	0.2	1.3	7.9	29.1	53.8	64.3
1999	7.1	*	*	*	*	*	0.2	1.3	9.0	32.0	58.1	62.5
Non-Hispanic 4/, male												
2006	4.9	*	*	*	*	*	0.3	1.3	5.6	20.6	46.9	65.3
2005	5.7	*	*	*	*	*	0.3	1.4	7.0	25.0	54.1	73.1
2004	5.6	*	*	*	*	*	0.2	1.3	6.6	25.6	54.0	75.5
2003	6.0	*	*	*	*	*	0.2	1.5	7.3	26.8	59.5	80.4
2002	6.5	*	*	*	*	*	0.3	1.5	7.7	29.3	65.2	91.3
2001	6.8	*	*	*	*	*	0.3	1.5	8.4	32.1	66.6	102.7
2000	7.1	*	*	*	*	*	0.2	1.6	8.5	34.3	70.6	105.9
1999	7.8	*	*	*	*	*	0.2	1.5	10.0	39.2	77.8	106.1
Non-Hispanic 4/, female												
2006	4.7	*	*	*	*	*	0.2	0.9	4.6	16.4	32.0	37.8
2005	5.1	*	*	*	*	*	0.1	0.8	4.8	18.0	35.1	44.0
2004	5.0	*	*	*	*	*	0.2	0.9	5.1	18.4	34.6	39.6
2003	5.6	*	*	*	*	*	0.2	0.9	6.0	20.3	39.1	44.5
2002	5.7	*	*	*	*	*	0.1	1.0	6.0	21.7	38.8	46.6
2001	6.1	*	*	*	*	*	0.1	1.0	7.0	23.6	41.9	46.3
2000	6.3	*	*	*	*	*	0.2	1.1	7.3	24.8	42.9	47.5
1999	6.5	*	*	*	*	*	0.2	1.1	8.1	26.1	45.3	45.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
Non-Hispanic white, both sexes												
2006 5/	5.7	*	*	*	*	*	0.2	1.2	5.5	20.0	40.5	49.2
2005 5/	6.3	*	*	*	*	*	0.2	1.1	6.3	23.0	45.2	55.9
2004 5/	6.2	*	*	*	*	*	0.2	1.1	6.2	23.7	45.2	53.4
2003 5/	6.8	*	*	*	*	*	0.2	1.2	7.1	25.2	49.9	57.9
2002	7.1	*	*	*	*	*	0.2	1.3	7.3	27.0	52.3	63.1
2001	7.5	*	*	*	*	*	0.2	1.3	8.2	29.6	54.9	65.9
2000	7.8	*	*	*	*	*	0.2	1.3	8.4	31.2	57.0	66.9
1999	8.3	*	*	*	*	*	0.2	1.3	9.7	34.5	61.2	65.2
Non-Hispanic white, male												
2006 5/	5.7	*	*	*	*	*	0.3	1.4	5.7	22.0	48.7	68.4
2005 5/	6.5	*	*	*	*	*	0.2	1.4	7.3	26.4	55.8	75.5
2004 5/	6.5	*	*	*	*	*	0.2	1.2	6.8	27.2	56.5	78.2
2003 5/	6.9	*	*	*	*	*	0.2	1.5	7.6	28.5	61.3	81.6
2002	7.4	*	*	*	*	*	0.3	1.5	8.0	30.6	67.7	94.3
2001	7.8	*	*	*	*	*	0.3	1.5	8.7	33.8	69.6	106.6
2000	8.1	*	*	*	*	*	0.2	1.4	8.9	35.8	73.9	107.3
1999	8.9	*	*	*	*	*	0.2	1.4	10.4	41.1	80.6	110.0
Non-Hispanic white, female												
2006 5/	5.7	*	*	*	*	*	0.2	1.1	5.2	18.2	34.8	40.3
2005 5/	6.1	*	*	*	*	*	*	0.9	5.4	20.0	37.8	47.0
2004 5/	6.0	*	*	*	*	*	0.2	0.9	5.6	20.6	37.6	42.3
2003 5/	6.7	*	*	*	*	*	0.2	1.0	6.7	22.5	42.3	47.4
2002	6.8	*	*	*	*	*	*	1.1	6.7	24.0	41.9	49.7
2001	7.2	*	*	*	*	*	0.1	1.0	7.7	26.1	45.2	48.9
2000	7.4	*	*	*	*	*	0.2	1.1	8.0	27.3	45.8	50.6
1999	7.7	*	*	*	*	*	0.2	1.2	9.0	29.0	48.5	47.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
Non-Hispanic black, both sexes												
2006 5/	1.9	*	*	*	*	*	0.4	1.0	4.3	10.3	22.6	24.2
2005 5/	2.2	*	*	*	*	*	*	1.4	4.6	12.9	26.7	27.3
2004 5/	1.9	*	*	*	*	*	*	1.3	4.7	11.0	22.5	27.9
2003 5/	2.2	*	*	*	*	*	*	1.2	5.3	13.2	27.7	31.2
2002	2.3	*	*	*	*	*	0.5	1.2	4.6	16.2	28.2	32.1
2001	2.2	*	*	*	*	*	*	1.4	5.6	15.6	23.6	35.9
2000	2.5	*	*	*	*	*	*	2.0	6.1	17.1	27.0	37.5
1999	2.7	*	*	*	*	*	0.4	1.9	6.3	17.3	31.8	37.8
Non-Hispanic black, male												
2006 5/	2.4	*	*	*	*	*	*	1.6	6.8	14.3	36.7	40.3
2005 5/	2.8	*	*	*	*	*	*	2.1	7.5	19.4	42.9	50.4
2004 5/	2.6	*	*	*	*	*	*	2.0	7.1	17.4	37.1	57.9
2003 5/	2.8	*	*	*	*	*	*	1.7	7.5	18.6	46.7	71.4
2002	3.1	*	*	*	*	*	*	1.8	6.7	25.3	49.5	65.6
2001	2.8	*	*	*	*	*	*	1.6	7.9	23.0	40.2	69.3
2000	3.3	*	*	*	*	*	*	3.0	8.2	27.4	41.3	89.2
1999	3.6	*	*	*	*	*	*	3.0	9.1	29.3	55.7	73.7
Non-Hispanic black, female												
2006 5/	1.4	*	*	*	*	*	*	*	2.3	7.5	14.7	17.6
2005 5/	1.6	*	*	*	*	*	*	0.8	2.3	8.2	17.5	18.0
2004 5/	1.3	*	*	*	*	*	*	*	2.7	6.4	14.3	16.1
2003 5/	1.6	*	*	*	*	*	*	*	3.5	9.4	16.9	15.7
2002	1.6	*	*	*	*	*	*	*	2.8	9.8	16.1	19.2
2001	1.7	*	*	*	*	*	*	1.1	3.7	10.4	14.3	23.5
2000	1.8	*	*	*	*	*	*	1.3	4.5	9.8	18.9	18.3
1999	1.8	*	*	*	*	*	*	1.0	4.0	8.9	18.4	24.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
All origins 3/, both sexes												
2006	1.2	*	0.2	0.2	0.3	0.5	0.9	1.3	1.6	2.3	4.8	12.3
2005	1.3	*	0.2	0.2	0.3	0.5	0.8	1.4	1.7	2.5	5.4	14.5
2004	1.3	*	0.2	0.3	0.4	0.5	0.9	1.4	1.8	2.7	5.2	14.1
2003	1.4	*	0.2	0.3	0.4	0.6	0.9	1.5	2.0	2.9	5.8	14.2
2002	1.5	*	0.3	0.3	0.4	0.6	1.1	1.5	2.0	3.2	6.4	14.7
2001	1.5	*	0.2	0.2	0.4	0.6	1.0	1.5	2.2	3.5	6.4	16.1
2000	1.6	*	0.2	0.3	0.4	0.6	1.0	1.6	2.2	4.0	6.9	16.7
1999	1.7	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.2	7.6	17.3
All origins 3/, male												
2006	0.9	*	*	0.3	0.4	0.5	0.7	1.0	1.1	1.7	3.5	9.1
2005	0.9	*	0.3	0.3	0.4	0.5	0.7	1.0	1.2	1.8	4.1	8.7
2004	1.0	*	*	0.3	0.5	0.6	0.7	1.0	1.3	2.1	3.5	11.1
2003	1.0	*	0.4	0.3	0.4	0.6	0.7	1.2	1.6	2.1	4.5	10.7
2002	1.1	*	0.4	0.3	0.5	0.6	0.9	1.2	1.6	2.4	5.0	10.9
2001	1.1	*	0.3	0.3	0.4	0.6	0.9	1.1	1.5	2.7	4.5	12.1
2000	1.2	*	0.3	0.4	0.5	0.7	0.9	1.3	1.7	3.0	5.0	12.9
1999	1.2	*	*	0.4	0.5	0.6	0.8	1.1	1.8	3.2	5.7	12.7
All origins 3/, female												
2006	1.5	*	*	0.1	0.3	0.4	1.0	1.6	2.0	2.9	5.7	13.9
2005	1.7	*	*	0.2	0.2	0.5	1.0	1.8	2.2	3.2	6.3	17.2
2004	1.6	*	*	0.2	0.3	0.4	1.0	1.6	2.2	3.2	6.3	15.4
2003	1.8	*	*	0.2	0.3	0.5	1.1	1.9	2.4	3.6	6.7	15.8
2002	1.8	*	*	0.3	0.3	0.5	1.2	1.8	2.4	3.9	7.3	16.3
2001	1.9	*	*	0.2	0.3	0.6	1.1	1.9	2.8	4.1	7.6	17.8
2000	2.0	*	*	0.3	0.3	0.5	1.2	2.0	2.7	4.8	8.1	18.2
1999	2.1	*	*	0.2	0.4	0.7	1.2	2.2	3.0	5.0	8.8	19.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
Hispanic, both sexes												
2006	0.6	*	*	*	*	0.3	0.5	0.6	1.4	2.4	4.4	16.4
2005	0.6	*	*	*	*	*	0.5	0.8	1.6	2.9	4.8	13.7
2004	0.7	*	*	*	*	0.3	0.5	1.1	1.9	3.3	5.3	11.4
2003	0.7	*	*	*	*	0.3	0.5	1.6	2.2	3.4	6.1	14.5
2002	0.7	*	*	*	*	0.3	0.7	1.2	1.9	3.8	7.0	12.2
2001	0.7	*	*	*	*	*	0.6	1.4	2.4	3.8	7.9	13.8
2000	0.8	*	*	*	*	*	0.6	1.3	2.8	5.6	7.9	16.6
1999	0.9	*	*	*	*	0.3	0.9	1.6	2.4	5.8	10.4	17.5
Hispanic, male												
2006	0.4	*	*	*	*	*	*	*	*	*	*	*
2005	0.5	*	*	*	*	*	*	*	*	*	*	*
2004	0.5	*	*	*	*	*	*	*	*	*	*	*
2003	0.6	*	*	*	*	*	*	1.4	2.2	*	*	*
2002	0.6	*	*	*	*	*	*	1.2	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000	0.7	*	*	*	*	*	*	1.3	*	5.3	*	*
1999	0.7	*	*	*	*	*	0.9	1.4	*	*	*	*
Hispanic, female												
2006	0.8	*	*	*	*	*	*	*	1.7	3.3	5.3	19.8
2005	0.7	*	*	*	*	*	*	*	2.1	3.4	5.6	13.0
2004	0.9	*	*	*	*	*	*	1.5	2.8	3.8	6.9	*
2003	0.9	*	*	*	*	*	*	1.8	2.1	4.1	6.3	17.2
2002	0.9	*	*	*	*	*	1.0	1.2	2.2	5.3	7.6	*
2001	1.0	*	*	*	*	*	*	2.0	2.9	3.9	9.4	18.6
2000	1.0	*	*	*	*	*	*	1.3	3.2	5.8	9.2	*
1999	1.2	*	*	*	*	*	1.0	1.9	3.1	7.1	13.2	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
Non-Hispanic 4/, both sexes												
2006	1.3	*	0.2	0.3	0.3	0.5	0.9	1.4	1.6	2.3	4.8	12.1
2005	1.4	*	0.3	0.3	0.4	0.6	0.9	1.5	1.7	2.5	5.5	14.5
2004	1.4	*	0.2	0.3	0.4	0.5	0.9	1.4	1.8	2.7	5.2	14.2
2003	1.5	*	0.2	0.3	0.4	0.6	1.0	1.5	2.0	2.9	5.8	14.1
2002	1.6	*	0.3	0.3	0.5	0.6	1.1	1.5	2.0	3.1	6.3	14.7
2001	1.6	*	0.2	0.3	0.4	0.7	1.1	1.5	2.2	3.4	6.3	16.1
2000	1.7	*	0.2	0.3	0.5	0.7	1.1	1.6	2.2	3.9	6.8	16.6
1999	1.8	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.1	7.4	17.3
Non-Hispanic 4/, male												
2006	1.0	*	*	0.4	0.4	0.6	0.8	1.1	1.1	1.7	3.5	9.0
2005	1.0	*	0.4	0.3	0.4	0.6	0.7	1.1	1.2	1.7	4.1	8.4
2004	1.0	*	*	0.3	0.5	0.6	0.7	1.1	1.4	2.1	3.5	11.1
2003	1.1	*	0.4	0.4	0.5	0.7	0.8	1.1	1.5	2.0	4.5	10.7
2002	1.2	*	0.4	0.3	0.6	0.7	0.9	1.2	1.6	2.4	4.9	11.1
2001	1.1	*	*	0.3	0.4	0.6	0.9	1.2	1.5	2.6	4.4	12.2
2000	1.2	*	*	0.4	0.6	0.8	0.9	1.3	1.6	2.9	4.9	12.7
1999	1.3	*	*	0.4	0.6	0.7	0.8	1.1	1.8	3.1	5.6	12.6
Non-Hispanic 4/, female												
2006	1.6	*	*	0.2	0.3	0.5	1.1	1.7	2.0	2.8	5.7	13.6
2005	1.9	*	*	0.2	0.3	0.6	1.1	1.9	2.2	3.2	6.3	17.3
2004	1.7	*	*	0.2	0.3	0.5	1.1	1.7	2.2	3.2	6.3	15.5
2003	1.9	*	*	0.2	0.4	0.5	1.2	1.9	2.4	3.6	6.7	15.6
2002	2.0	*	*	0.3	0.3	0.6	1.2	1.9	2.4	3.8	7.2	16.3
2001	2.1	*	*	0.2	0.4	0.7	1.2	1.9	2.8	4.1	7.5	17.7
2000	2.1	*	*	0.3	0.4	0.6	1.2	2.0	2.6	4.7	8.0	18.2
1999	2.2	*	*	0.2	0.4	0.7	1.2	2.2	3.0	4.9	8.6	19.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
Non-Hispanic white, both sexes												
2006 5/	1.1	*	*	0.1	0.2	0.3	0.7	1.0	1.1	1.9	4.5	11.8
2005 5/	1.2	*	*	0.1	0.2	0.4	0.7	1.1	1.3	1.9	4.9	14.5
2004 5/	1.2	*	*	0.1	0.2	0.3	0.7	0.9	1.3	2.3	4.6	13.8
2003 5/	1.3	*	*	0.1	0.2	0.4	0.7	1.0	1.5	2.4	5.3	14.1
2002	1.4	*	*	0.1	0.3	0.5	0.7	1.1	1.5	2.6	6.1	14.7
2001	1.4	*	*	0.1	0.2	0.4	0.7	1.1	1.5	2.9	5.8	15.7
2000	1.4	*	*	0.2	0.3	0.4	0.7	1.1	1.5	3.2	6.4	15.8
1999	1.5	*	*	0.2	0.4	0.4	0.7	1.2	1.9	3.4	7.1	16.5
Non-Hispanic white, male												
2006 5/	0.7	*	*	*	0.2	0.3	0.5	0.7	0.7	1.3	3.2	9.0
2005 5/	0.8	*	*	*	0.2	0.4	0.5	0.8	0.9	1.3	3.5	8.2
2004 5/	0.8	*	*	*	0.3	0.3	0.5	0.7	0.9	1.7	2.8	10.4
2003 5/	0.9	*	*	*	0.3	0.4	0.5	0.8	1.1	1.7	3.8	10.6
2002	0.9	*	*	0.2	0.3	0.5	0.5	0.8	1.1	1.8	4.5	10.8
2001	0.9	*	*	0.2	0.2	0.3	0.6	0.8	1.0	2.1	3.9	11.4
2000	1.0	*	*	0.2	0.3	0.5	0.6	0.9	1.0	2.3	4.6	11.1
1999	1.0	*	*	0.2	0.4	0.4	0.5	0.7	1.4	2.4	5.2	10.6
Non-Hispanic white, female												
2006 5/	1.5	*	*	*	*	0.3	0.8	1.3	1.6	2.4	5.4	13.2
2005 5/	1.7	*	*	*	0.2	0.5	0.8	1.3	1.7	2.5	5.9	17.4
2004 5/	1.6	*	*	*	0.2	0.4	0.8	1.1	1.7	2.9	5.9	15.3
2003 5/	1.7	*	*	*	0.2	0.5	0.9	1.3	2.0	2.9	6.3	15.7
2002	1.8	*	*	*	0.3	0.5	0.9	1.3	1.8	3.2	7.1	16.4
2001	1.9	*	*	*	0.2	0.4	0.8	1.4	2.1	3.6	7.1	17.5
2000	1.9	*	*	0.2	0.3	0.4	0.9	1.3	2.0	4.0	7.6	17.7
1999	2.1	*	*	*	0.3	0.5	0.9	1.6	2.3	4.2	8.3	18.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
Non-Hispanic black, both sexes												
2006 5/	2.5	*	*	1.0	1.1	1.7	2.5	4.0	5.2	5.5	6.1	12.7
2005 5/	2.7	*	1.1	0.8	1.1	1.6	2.4	4.3	5.5	7.2	9.9	12.0
2004 5/	2.7	*	*	0.9	1.2	1.7	2.5	4.6	5.8	5.8	9.4	15.3
2003 5/	2.8	*	0.8	0.9	1.2	1.7	2.8	4.9	5.2	7.2	9.3	11.9
2002	3.0	*	0.9	1.1	1.3	1.6	3.5	4.9	6.6	8.0	7.9	11.8
2001	3.1	*	*	0.8	1.2	2.2	3.3	4.7	6.9	7.7	10.7	16.5
2000	3.3	*	*	1.1	1.3	2.0	3.2	5.5	6.9	9.4	10.4	22.3
1999	3.3	*	*	1.0	1.3	2.4	3.0	5.4	7.0	9.8	10.6	20.7
Non-Hispanic black, male												
2006 5/	2.3	*	*	1.2	1.4	2.0	2.4	3.5	4.7	4.8	*	*
2005 5/	2.3	*	1.6	1.0	1.4	1.7	2.2	3.0	4.1	5.3	10.7	*
2004 5/	2.6	*	*	1.2	1.5	2.4	2.3	4.0	5.3	5.7	8.8	*
2003 5/	2.5	*	*	1.3	1.4	2.5	2.4	4.0	4.4	5.5	8.9	*
2002	2.8	*	*	1.1	1.9	2.0	3.7	4.0	5.9	7.5	6.7	*
2001	2.7	*	*	0.9	1.4	2.5	3.0	4.1	4.7	6.8	8.7	*
2000	2.8	*	*	1.3	1.6	2.5	3.0	3.7	6.1	7.9	6.5	24.7
1999	2.9	*	*	1.4	1.7	2.7	2.8	4.0	5.5	9.2	8.4	*
Non-Hispanic black, female												
2006 5/	2.7	*	*	0.7	0.9	1.3	2.6	4.4	5.5	6.1	7.1	15.3
2005 5/	3.1	*	*	*	0.8	1.5	2.6	5.4	6.6	8.6	9.5	13.8
2004 5/	2.8	*	*	*	1.0	1.1	2.7	5.1	6.2	5.9	9.7	16.9
2003 5/	3.0	*	*	*	1.1	1.0	3.2	5.7	5.9	8.4	9.5	13.6
2002	3.1	*	*	1.0	*	1.3	3.4	5.7	7.2	8.4	8.6	12.5
2001	3.4	*	*	0.7	0.9	2.0	3.6	5.1	8.6	8.4	11.8	17.1
2000	3.7	*	*	0.8	1.0	1.5	3.4	7.0	7.6	10.4	12.7	21.4
1999	3.6	*	*	0.6	0.9	2.2	3.2	6.5	8.1	10.2	11.7	21.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
All origins 3/, both sexes												
2006	36.0	*	*	*	*	0.1	0.9	6.6	32.6	129.1	320.2	525.4
2005	37.8	*	*	*	*	0.1	1.0	6.9	34.5	137.2	337.2	563.5
2004	35.3	*	*	*	*	0.1	0.9	6.0	33.0	130.0	319.0	531.1
2003	36.6	*	*	*	0.1	0.1	1.0	5.9	34.8	137.5	329.2	558.9
2002	36.1	*	*	*	*	0.1	0.9	5.9	33.8	135.1	330.1	555.5
2001	35.7	*	*	*	0.1	0.1	0.9	5.7	34.5	137.5	321.0	557.4
2000	35.4	*	*	*	*	0.1	0.8	5.6	34.3	136.7	323.8	556.7
1999	36.0	*	*	*	*	0.1	0.7	5.5	36.2	141.4	330.6	555.8
All origins 3/, male												
2006	34.8	*	*	*	*	0.1	0.8	6.9	35.3	141.5	386.8	682.2
2005	36.6	*	*	*	*	0.1	1.0	7.3	37.0	154.4	405.2	732.4
2004	34.5	*	*	*	*	0.1	1.0	6.5	35.2	145.0	388.4	718.6
2003	35.9	*	*	*	*	0.1	1.0	6.4	37.8	155.4	400.1	769.0
2002	35.7	*	*	*	*	0.1	1.0	6.4	36.2	153.4	409.8	785.1
2001	35.3	*	*	*	*	0.1	0.9	6.2	37.2	157.0	403.0	794.5
2000	35.6	*	*	*	*	*	0.8	6.1	37.7	158.3	410.9	838.0
1999	37.1	*	*	*	*	*	0.8	5.9	40.5	168.6	427.9	866.7
All origins 3/, female												
2006	37.1	*	*	*	*	0.1	0.9	6.4	30.1	118.7	274.8	452.1
2005	39.0	*	*	*	*	*	1.0	6.4	32.2	122.6	291.0	485.9
2004	36.2	*	*	*	*	*	0.9	5.5	30.8	117.3	272.2	446.8
2003	37.4	*	*	*	*	*	1.0	5.5	32.1	122.6	281.9	466.0
2002	36.5	*	*	*	*	0.1	0.9	5.4	31.5	119.8	277.2	455.8
2001	36.0	*	*	*	*	0.1	1.0	5.2	32.0	121.3	267.1	458.2
2000	35.2	*	*	*	*	*	0.8	5.2	31.2	119.0	267.1	442.2
1999	35.0	*	*	*	*	0.2	0.7	5.0	32.2	119.1	267.4	430.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
Hispanic, both sexes												
2006	6.2	*	*	*	*	*	*	1.6	8.9	39.0	132.1	331.8
2005	6.7	*	*	*	*	*	*	2.5	9.7	42.0	148.9	367.9
2004	6.1	*	*	*	*	*	0.5	1.7	9.3	43.2	135.4	345.4
2003	6.3	*	*	*	*	*	0.4	1.7	9.1	43.7	143.5	403.7
2002	6.3	*	*	*	*	*	0.4	2.1	10.4	43.1	147.8	397.4
2001	6.0	*	*	*	*	*	*	1.9	10.0	45.2	139.6	418.9
2000	5.9	*	*	*	*	*	0.4	1.7	9.9	40.7	149.1	422.7
1999	6.5	*	*	*	*	*	0.4	1.9	10.8	49.3	158.1	462.3
Hispanic, male												
2006	6.2	*	*	*	*	*	*	1.8	10.7	49.0	170.1	439.8
2005	6.9	*	*	*	*	*	*	2.6	13.0	53.5	205.8	468.2
2004	6.3	*	*	*	*	*	0.7	1.7	11.2	59.2	181.0	455.3
2003	6.6	*	*	*	*	*	*	1.5	12.2	57.8	194.4	574.2
2002	6.6	*	*	*	*	*	*	2.2	14.0	57.9	208.9	535.4
2001	6.3	*	*	*	*	*	*	2.1	12.2	62.5	181.5	605.7
2000	6.2	*	*	*	*	*	*	1.6	13.6	56.6	209.0	584.5
1999	7.0	*	*	*	*	*	*	1.6	13.9	65.9	228.9	735.4
Hispanic, female												
2006	6.1	*	*	*	*	*	*	1.4	7.3	30.8	105.6	274.4
2005	6.5	*	*	*	*	*	*	2.3	6.6	32.7	109.3	315.4
2004	5.9	*	*	*	*	*	*	1.7	7.6	30.4	104.0	288.8
2003	6.0	*	*	*	*	*	*	1.9	6.3	32.5	108.5	317.2
2002	5.9	*	*	*	*	*	*	1.9	7.1	31.3	106.0	329.1
2001	5.8	*	*	*	*	*	*	1.8	8.1	31.5	111.2	327.7
2000	5.6	*	*	*	*	*	*	1.7	6.6	28.2	109.0	343.3
1999	5.9	*	*	*	*	*	*	2.1	8.1	36.3	111.0	330.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
Non-Hispanic 4/, both sexes												
2006	41.1	*	*	*	*	0.1	1.0	7.2	34.6	136.0	331.4	534.6
2005	43.0	*	*	*	*	0.1	1.1	7.3	36.5	144.2	347.8	572.1
2004	40.0	*	*	*	*	0.1	1.0	6.4	34.8	136.0	328.7	538.8
2003	41.4	*	*	*	*	0.1	1.1	6.3	36.7	143.8	338.3	564.3
2002	40.6	*	*	*	*	0.1	1.0	6.3	35.6	141.0	338.1	560.8
2001	40.0	*	*	*	0.1	0.1	1.0	6.0	36.2	143.2	328.6	561.3
2000	39.5	*	*	*	*	0.1	0.9	6.0	36.0	142.3	330.5	560.5
1999	40.0	*	*	*	*	0.1	0.8	5.8	37.9	146.4	336.8	557.6
Non-Hispanic 4/, male												
2006	40.0	*	*	*	*	0.1	0.9	7.5	37.3	148.2	399.8	694.8
2005	41.8	*	*	*	*	*	1.1	7.8	38.9	161.6	416.5	745.0
2004	39.3	*	*	*	*	0.2	1.0	7.0	37.1	150.6	399.4	730.9
2003	40.7	*	*	*	*	0.1	1.0	6.9	39.6	161.7	410.1	776.7
2002	40.4	*	*	*	*	0.1	1.1	6.9	37.8	159.2	418.5	794.9
2001	39.8	*	*	*	*	0.1	1.0	6.6	38.9	162.6	412.4	800.6
2000	40.0	*	*	*	*	*	0.9	6.5	39.3	163.8	418.4	846.6
1999	41.4	*	*	*	*	*	0.8	6.3	42.2	173.9	435.1	869.6
Non-Hispanic 4/, female												
2006	42.1	*	*	*	*	0.1	1.0	7.0	32.0	125.6	284.8	460.1
2005	44.1	*	*	*	*	*	1.1	6.9	34.3	129.4	301.2	493.2
2004	40.8	*	*	*	*	*	1.0	5.9	32.7	123.7	281.2	453.0
2003	42.0	*	*	*	*	*	1.1	5.8	34.1	128.9	290.3	471.0
2002	40.8	*	*	*	*	0.1	1.0	5.8	33.4	125.8	284.9	459.8
2001	40.1	*	*	*	*	*	1.0	5.5	33.8	127.1	273.6	461.7
2000	39.1	*	*	*	*	*	0.9	5.5	32.9	124.5	273.2	444.8
1999	38.7	*	*	*	*	0.2	0.7	5.2	33.9	123.9	272.9	432.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
Non-Hispanic white, both sexes												
2006 5/	48.5	*	*	*	*	0.1	1.0	7.6	36.9	147.3	351.8	562.5
2005 5/	50.6	*	*	*	*	0.1	1.1	7.6	39.0	155.5	369.2	600.8
2004 5/	47.1	*	*	*	*	0.1	1.1	6.7	37.1	146.7	348.1	566.1
2003 5/	48.6	*	*	*	*	0.1	1.1	6.6	39.0	154.4	357.6	592.2
2002	47.5	*	*	*	*	0.1	1.0	6.5	37.7	151.2	355.7	586.9
2001	46.6	*	*	*	*	0.1	1.0	6.2	38.6	152.6	345.7	585.7
2000	46.0	*	*	*	*	*	0.8	6.1	37.9	151.2	347.1	585.5
1999	46.3	*	*	*	*	0.1	0.7	5.9	39.9	154.8	351.4	582.8
Non-Hispanic white, male												
2006 5/	46.4	*	*	*	*	*	0.9	7.8	38.8	156.8	416.3	720.3
2005 5/	48.4	*	*	*	*	*	1.1	7.8	40.5	170.4	432.7	770.7
2004 5/	45.5	*	*	*	*	0.2	1.1	7.2	38.4	158.9	415.6	756.7
2003 5/	46.9	*	*	*	*	*	1.0	7.1	41.1	169.5	423.7	804.2
2002	46.4	*	*	*	*	*	1.1	6.9	39.0	167.0	431.9	821.0
2001	45.6	*	*	*	*	*	1.0	6.7	40.5	169.3	425.2	823.9
2000	45.7	*	*	*	*	*	0.9	6.4	40.5	169.8	431.4	876.5
1999	47.1	*	*	*	*	*	0.8	6.2	43.4	180.3	445.9	900.5
Non-Hispanic white, female												
2006 5/	50.5	*	*	*	*	*	1.0	7.5	35.1	139.2	307.0	488.9
2005 5/	52.8	*	*	*	*	*	1.2	7.3	37.6	142.5	325.5	523.1
2004 5/	48.8	*	*	*	*	*	1.1	6.3	35.9	136.2	302.2	480.7
2003 5/	50.2	*	*	*	*	*	1.1	6.1	37.0	141.5	313.1	498.8
2002	48.6	*	*	*	*	*	1.0	6.2	36.4	137.7	304.6	485.7
2001	47.6	*	*	*	*	*	1.0	5.7	36.8	138.5	292.9	486.6
2000	46.3	*	*	*	*	*	0.8	5.8	35.5	135.5	291.6	468.2
1999	45.5	*	*	*	*	*	0.7	5.5	36.6	133.4	289.5	456.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
Non-Hispanic black, both sexes												
2006 5/	15.8	*	*	*	*	*	1.2	7.2	28.4	86.5	199.5	296.8
2005 5/	16.8	*	*	*	*	*	1.2	8.5	29.8	95.7	207.2	328.2
2004 5/	15.0	*	*	*	*	*	1.1	6.9	28.4	86.8	193.2	288.1
2003 5/	15.7	*	*	*	*	*	1.3	6.5	30.5	94.6	200.3	306.1
2002	15.9	*	*	*	*	*	1.2	6.8	30.2	90.3	213.1	322.7
2001	15.6	*	*	*	*	*	1.3	6.5	28.7	96.1	200.8	323.4
2000	15.4	*	*	*	*	*	1.3	6.8	31.4	95.8	196.8	312.6
1999	16.4	*	*	*	*	*	1.2	6.6	33.2	101.9	220.1	314.0
Non-Hispanic black, male												
2006 5/	17.9	*	*	*	*	*	1.1	8.3	36.9	120.1	301.7	473.7
2005 5/	19.5	*	*	*	*	*	1.2	10.3	38.0	134.5	325.7	534.8
2004 5/	17.4	*	*	*	*	*	1.3	8.0	38.8	119.0	295.8	495.9
2003 5/	18.6	*	*	*	*	*	1.4	7.7	39.3	133.6	329.0	526.3
2002	18.7	*	*	*	*	*	1.3	8.5	40.3	126.6	337.9	572.1
2001	18.8	*	*	*	*	*	1.4	7.5	37.6	139.1	334.9	592.5
2000	18.7	*	*	*	*	*	1.3	8.5	40.7	142.7	322.4	578.9
1999	20.3	*	*	*	*	*	1.5	8.3	45.4	146.5	369.3	610.9
Non-Hispanic black, female												
2006 5/	13.8	*	*	*	*	*	1.3	6.2	21.6	62.5	141.5	224.6
2005 5/	14.4	*	*	*	*	*	1.2	6.8	23.2	68.0	140.0	245.1
2004 5/	12.9	*	*	*	*	*	1.0	5.9	20.1	63.8	135.2	206.0
2003 5/	13.0	*	*	*	*	*	1.2	5.5	23.5	66.8	127.6	220.8
2002	13.3	*	*	*	*	*	1.1	5.3	22.1	64.6	142.4	227.4
2001	12.7	*	*	*	*	*	1.3	5.7	21.5	65.8	125.5	223.7
2000	12.4	*	*	*	*	*	1.3	5.3	24.0	62.8	126.1	213.7
1999	12.9	*	*	*	*	*	1.0	5.0	23.3	70.4	136.6	203.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
All origins 3/, both sexes												
2006	0.3	*	*	*	*	*	*	0.0	0.2	1.0	3.0	5.1
2005	0.3	*	*	*	*	*	*	0.1	0.2	0.9	3.4	5.7
2004	0.4	*	*	*	*	*	*	*	0.2	1.1	3.4	6.0
2003	0.4	*	*	*	*	*	*	*	0.3	1.3	3.8	6.4
2002	0.4	*	*	*	*	*	*	*	0.3	1.2	3.8	6.8
2001	0.4	*	*	*	*	*	*	*	0.3	1.5	4.1	6.2
2000	0.4	*	*	*	*	*	*	*	0.2	1.5	3.9	7.5
1999	0.4	*	*	*	*	*	*	*	0.2	1.5	4.1	6.1
All origins 3/, male												
2006	0.6	*	*	*	*	*	*	*	0.3	2.0	7.2	15.3
2005	0.7	*	*	*	*	*	*	*	0.4	2.0	8.1	17.1
2004	0.7	*	*	*	*	*	*	*	0.5	2.4	8.1	18.4
2003	0.7	*	*	*	*	*	*	*	0.5	2.7	9.1	19.8
2002	0.7	*	*	*	*	*	*	*	0.5	2.6	9.0	21.6
2001	0.8	*	*	*	*	*	*	*	0.5	3.1	9.9	20.2
2000	0.8	*	*	*	*	*	*	*	0.5	3.2	9.6	24.9
1999	0.8	*	*	*	*	*	*	*	0.5	3.2	10.1	20.4
All origins 3/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
Non-Hispanic 4/, both sexes												
2006	0.4	*	*	*	*	*	*	*	0.2	1.0	3.2	5.3
2005	0.4	*	*	*	*	*	*	0.1	0.2	1.0	3.6	5.8
2004	0.4	*	*	*	*	*	*	*	0.3	1.2	3.5	6.1
2003	0.4	*	*	*	*	*	*	*	0.3	1.3	3.9	6.6
2002	0.4	*	*	*	*	*	*	*	0.3	1.2	3.9	6.9
2001	0.4	*	*	*	*	*	*	*	0.3	1.5	4.2	6.2
2000	0.5	*	*	*	*	*	*	*	0.2	1.6	4.0	7.6
1999	0.4	*	*	*	*	*	*	*	0.3	1.5	4.3	6.2
Non-Hispanic 4/, male												
2006	0.7	*	*	*	*	*	*	*	0.3	2.1	7.6	16.0
2005	0.8	*	*	*	*	*	*	*	0.5	2.1	8.5	17.8
2004	0.8	*	*	*	*	*	*	*	0.5	2.5	8.4	18.9
2003	0.9	*	*	*	*	*	*	*	0.5	2.8	9.3	20.7
2002	0.8	*	*	*	*	*	*	*	0.5	2.6	9.3	21.9
2001	0.9	*	*	*	*	*	*	*	0.5	3.3	10.2	20.5
2000	0.9	*	*	*	*	*	*	*	0.5	3.3	9.7	25.7
1999	0.9	*	*	*	*	*	*	*	0.5	3.4	10.5	21.1
Non-Hispanic 4/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
Non-Hispanic white, both sexes												
2006 5/	0.4	*	*	*	*	*	*	*	0.2	1.1	3.4	5.7
2005 5/	0.5	*	*	*	*	*	*	*	0.3	1.1	3.8	6.2
2004 5/	0.5	*	*	*	*	*	*	*	0.3	1.3	3.8	6.3
2003 5/	0.5	*	*	*	*	*	*	*	0.3	1.5	4.1	7.1
2002	0.5	*	*	*	*	*	*	*	0.3	1.4	4.1	7.3
2001	0.5	*	*	*	*	*	*	*	0.3	1.6	4.5	6.6
2000	0.5	*	*	*	*	*	*	*	0.3	1.7	4.1	8.1
1999	0.5	*	*	*	*	*	*	*	0.2	1.6	4.6	6.4
Non-Hispanic white, male												
2006 5/	0.8	*	*	*	*	*	*	*	0.3	2.3	8.1	17.1
2005 5/	0.9	*	*	*	*	*	*	*	0.5	2.3	9.0	18.7
2004 5/	0.9	*	*	*	*	*	*	*	0.6	2.7	9.0	19.6
2003 5/	1.0	*	*	*	*	*	*	*	0.6	3.0	9.8	22.3
2002	1.0	*	*	*	*	*	*	*	0.6	2.8	9.8	23.2
2001	1.0	*	*	*	*	*	*	*	0.5	3.5	10.8	21.6
2000	1.1	*	*	*	*	*	*	*	0.5	3.6	10.1	27.2
1999	1.0	*	*	*	*	*	*	*	0.5	3.6	11.1	21.8
Non-Hispanic white, female												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
Non-Hispanic black, both sexes												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.1	*	*	*	*	*	*	*	*	*	2.4	*
2002	0.2	*	*	*	*	*	*	*	*	*	2.3	*
2001	0.2	*	*	*	*	*	*	*	*	*	2.2	*
2000	0.2	*	*	*	*	*	*	*	*	*	2.6	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.3	*	*	*	*	*	*	*	*	*	6.3	*
2002	0.3	*	*	*	*	*	*	*	*	*	5.9	*
2001	0.3	*	*	*	*	*	*	*	*	*	6.0	*
2000	0.3	*	*	*	*	*	*	*	*	*	6.5	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
All origins 3/, both sexes												
2006	5.6	*	*	*	0.1	0.2	0.4	1.0	2.7	9.3	40.5	155.8
2005	5.8	*	*	*	0.1	0.2	0.4	1.1	2.7	9.3	42.6	163.9
2004	5.7	*	*	*	0.1	0.2	0.4	0.9	2.6	9.6	42.9	164.9
2003	6.0	*	*	*	0.1	0.2	0.4	1.0	2.8	9.6	45.0	176.2
2002	6.1	*	*	*	0.1	0.2	0.4	0.9	2.5	9.8	46.3	186.0
2001	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.8	189.4
2000	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6
1999	5.5	*	*	*	0.1	0.2	0.4	0.8	2.5	9.5	41.1	175.6
All origins 3/, male												
2006	6.0	*	*	*	*	0.3	0.5	1.3	3.5	12.6	57.2	224.8
2005	6.0	*	*	*	0.1	0.2	0.4	1.3	3.5	12.5	57.9	229.3
2004	5.9	*	*	*	0.1	0.2	0.4	1.2	3.2	12.9	58.6	237.2
2003	6.1	*	*	*	0.1	0.2	0.4	1.2	3.5	12.8	61.7	246.2
2002	6.3	*	*	*	0.1	0.2	0.5	1.1	3.1	13.0	63.8	268.5
2001	6.1	*	*	*	0.1	0.2	0.5	1.2	3.2	13.3	63.1	270.3
2000	6.1	*	*	*	0.1	0.2	0.5	1.2	3.0	14.4	62.3	276.0
1999	5.7	*	*	*	0.1	0.2	0.4	0.9	3.2	13.1	59.7	258.4
All origins 3/, female												
2006	5.2	*	*	*	*	0.1	0.3	0.8	1.9	6.4	29.1	123.5
2005	5.7	*	*	*	0.1	0.1	0.4	0.9	2.1	6.7	32.2	133.9
2004	5.5	*	*	*	*	0.1	0.3	0.7	1.9	6.8	32.3	132.4
2003	5.8	*	*	*	*	0.1	0.3	0.7	2.1	6.9	33.8	145.3
2002	5.9	*	*	*	*	0.2	0.3	0.6	2.0	7.1	34.7	150.2
2001	6.0	*	*	*	*	0.2	0.3	0.7	2.1	7.2	34.5	155.6
2000	5.8	*	*	*	*	0.1	0.3	0.7	1.9	7.0	33.0	151.6
1999	5.3	*	*	*	*	0.2	0.3	0.6	1.8	6.6	29.1	142.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
Hispanic, both sexes												
2006	1.3	*	*	*	*	*	*	0.5	1.3	5.7	25.9	77.5
2005	1.4	*	*	*	*	*	*	0.7	1.4	5.3	24.3	109.9
2004	1.3	*	*	*	*	*	*	*	1.6	5.1	26.3	101.3
2003	1.3	*	*	*	*	*	*	*	1.7	6.5	22.8	113.0
2002	1.4	*	*	*	*	*	*	*	1.8	5.3	29.0	131.4
2001	1.5	*	*	*	*	*	*	*	1.8	6.1	27.4	161.3
2000	1.4	*	*	*	*	*	*	0.6	1.9	5.9	27.9	137.4
1999	1.2	*	*	*	*	*	*	*	1.6	5.5	24.6	130.5
Hispanic, male												
2006	1.3	*	*	*	*	*	*	*	2.1	7.3	31.9	86.8
2005	1.4	*	*	*	*	*	*	*	2.2	6.1	32.1	154.4
2004	1.3	*	*	*	*	*	*	*	2.4	7.4	31.8	121.9
2003	1.3	*	*	*	*	*	*	*	2.0	8.7	32.2	123.1
2002	1.4	*	*	*	*	*	*	*	2.6	6.1	43.1	167.3
2001	1.5	*	*	*	*	*	*	*	2.5	8.4	36.2	220.9
2000	1.4	*	*	*	*	*	*	*	2.5	7.4	42.8	169.3
1999	1.1	*	*	*	*	*	*	*	*	6.8	33.9	159.6
Hispanic, female												
2006	1.3	*	*	*	*	*	*	*	*	4.3	21.7	72.5
2005	1.3	*	*	*	*	*	*	*	*	4.7	18.8	86.7
2004	1.3	*	*	*	*	*	*	*	*	3.3	22.5	90.8
2003	1.4	*	*	*	*	*	*	*	*	4.9	16.3	107.8
2002	1.4	*	*	*	*	*	*	*	*	4.6	19.4	113.6
2001	1.5	*	*	*	*	*	*	*	*	4.3	21.5	132.2
2000	1.4	*	*	*	*	*	*	*	*	4.8	17.8	121.7
1999	1.3	*	*	*	*	*	*	*	*	4.5	18.4	116.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
Non-Hispanic 4/, both sexes												
2006	6.4	*	*	*	0.1	0.2	0.4	1.1	2.8	9.5	41.4	159.6
2005	6.6	*	*	*	0.1	0.2	0.4	1.2	2.8	9.6	43.6	166.3
2004	6.4	*	*	*	0.1	0.2	0.4	1.0	2.6	9.9	43.7	167.4
2003	6.7	*	*	*	0.1	0.2	0.4	1.0	2.9	9.7	46.0	178.5
2002	6.8	*	*	*	0.1	0.2	0.5	0.9	2.6	10.1	47.1	187.8
2001	6.7	*	*	*	0.1	0.2	0.4	1.0	2.7	10.2	46.5	190.0
2000	6.5	*	*	*	0.1	0.2	0.4	1.0	2.5	10.6	45.1	189.0
1999	6.0	*	*	*	0.1	0.2	0.4	0.8	2.5	9.7	41.7	176.8
Non-Hispanic 4/, male												
2006	6.9	*	*	*	*	0.3	0.5	1.3	3.6	13.0	58.8	232.3
2005	6.8	*	*	*	0.1	0.2	0.5	1.4	3.6	12.9	59.3	232.8
2004	6.7	*	*	*	*	0.2	0.5	1.3	3.3	13.2	60.1	242.7
2003	6.9	*	*	*	0.1	0.2	0.4	1.2	3.6	13.0	63.1	251.5
2002	7.0	*	*	*	0.1	0.2	0.6	1.2	3.1	13.4	64.7	272.1
2001	6.8	*	*	*	0.1	0.2	0.6	1.3	3.2	13.6	64.2	271.6
2000	6.8	*	*	*	0.1	0.2	0.5	1.2	3.1	14.8	62.8	279.7
1999	6.3	*	*	*	0.1	0.3	0.5	1.0	3.3	13.3	60.6	261.6
Non-Hispanic 4/, female												
2006	5.9	*	*	*	*	0.1	0.3	0.8	2.1	6.6	29.5	125.8
2005	6.3	*	*	*	0.1	0.1	0.4	1.0	2.2	6.8	32.9	136.0
2004	6.1	*	*	*	*	0.2	0.3	0.7	2.0	7.1	32.8	133.8
2003	6.5	*	*	*	*	0.2	0.4	0.8	2.1	7.0	34.6	146.5
2002	6.6	*	*	*	*	0.2	0.3	0.6	2.1	7.3	35.4	151.4
2001	6.6	*	*	*	*	0.2	0.3	0.8	2.2	7.4	34.9	156.0
2000	6.3	*	*	*	*	0.1	0.3	0.8	2.0	7.1	33.6	152.3
1999	5.8	*	*	*	*	0.2	0.3	0.7	1.8	6.8	29.4	142.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
Non-Hispanic white, both sexes												
2006 5/	7.3	*	*	*	*	0.2	0.4	1.1	2.8	9.2	42.0	164.1
2005 5/	7.4	*	*	*	0.1	0.2	0.5	1.1	2.7	9.4	43.5	169.0
2004 5/	7.2	*	*	*	0.1	0.2	0.4	1.0	2.4	9.7	44.1	170.4
2003 5/	7.5	*	*	*	0.1	0.2	0.4	1.0	2.6	9.5	45.7	181.0
2002	7.7	*	*	*	0.1	0.2	0.4	0.9	2.5	9.7	47.3	191.3
2001	7.5	*	*	*	0.1	0.2	0.4	0.9	2.6	9.9	46.6	191.2
2000	7.3	*	*	*	0.1	0.2	0.4	0.9	2.3	10.0	45.0	192.5
1999	6.7	*	*	*	0.1	0.2	0.4	0.7	2.4	9.2	41.2	179.3
Non-Hispanic white, male												
2006 5/	7.8	*	*	*	*	0.3	0.5	1.3	3.6	12.5	59.2	239.9
2005 5/	7.6	*	*	*	*	0.2	0.5	1.3	3.4	12.4	58.7	237.8
2004 5/	7.5	*	*	*	*	0.2	0.5	1.3	2.9	12.7	60.2	247.4
2003 5/	7.6	*	*	*	*	0.2	0.4	1.2	3.3	12.6	62.4	256.5
2002	7.9	*	*	*	*	0.3	0.6	1.1	3.0	12.9	64.9	278.4
2001	7.6	*	*	*	*	0.3	0.6	1.2	3.1	13.2	64.4	276.9
2000	7.5	*	*	*	*	0.2	0.5	1.1	2.9	13.7	62.7	288.1
1999	6.9	*	*	*	*	0.2	0.4	0.9	3.0	12.4	59.6	267.4
Non-Hispanic white, female												
2006 5/	6.7	*	*	*	*	*	0.3	0.8	2.1	6.4	30.1	128.7
2005 5/	7.2	*	*	*	*	0.2	0.4	1.0	2.1	6.8	33.1	137.5
2004 5/	6.9	*	*	*	*	*	0.3	0.7	2.0	7.1	33.1	135.9
2003 5/	7.3	*	*	*	*	0.2	0.3	0.8	2.0	6.8	34.4	147.8
2002	7.4	*	*	*	*	0.2	0.3	0.6	2.0	7.1	35.6	153.7
2001	7.4	*	*	*	*	0.2	0.3	0.7	2.1	7.0	34.7	155.5
2000	7.1	*	*	*	*	*	0.3	0.7	1.8	7.0	33.4	153.9
1999	6.5	*	*	*	*	0.2	0.3	0.6	1.7	6.5	29.3	144.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
Non-Hispanic black, both sexes												
2006 5/	3.8	*	*	*	*	0.4	0.7	1.4	3.8	14.1	42.9	130.9
2005 5/	4.3	*	*	*	*	*	0.5	1.8	4.6	13.5	53.2	155.4
2004 5/	4.0	*	*	*	*	*	0.6	1.4	5.0	14.2	47.6	147.4
2003 5/	4.4	*	*	*	*	*	0.7	1.3	5.3	13.9	55.5	165.7
2002	4.2	*	*	*	*	*	0.7	1.4	3.8	15.2	52.8	167.3
2001	4.4	*	*	*	*	*	0.6	1.6	4.1	15.4	53.6	187.3
2000	4.3	*	*	*	*	*	0.7	1.9	4.3	17.9	52.0	160.6
1999	4.3	*	*	*	*	*	0.7	1.5	4.5	16.4	52.6	168.5
Non-Hispanic black, male												
2006 5/	4.2	*	*	*	*	*	0.8	1.7	5.3	21.2	68.4	177.8
2005 5/	4.6	*	*	*	*	*	*	2.5	6.3	20.0	81.0	205.5
2004 5/	4.3	*	*	*	*	*	0.8	1.8	7.3	21.5	68.7	207.3
2003 5/	4.5	*	*	*	*	*	*	1.7	7.2	19.8	79.6	215.2
2002	4.4	*	*	*	*	*	0.9	2.0	4.9	22.3	74.1	237.4
2001	4.3	*	*	*	*	*	0.8	1.9	5.5	20.8	75.6	243.7
2000	4.5	*	*	*	*	*	0.8	2.7	5.2	28.4	75.0	210.2
1999	4.7	*	*	*	*	*	0.9	1.8	6.6	25.3	81.6	229.4
Non-Hispanic black, female												
2006 5/	3.4	*	*	*	*	*	*	1.2	2.5	9.0	28.4	111.7
2005 5/	4.0	*	*	*	*	*	*	1.1	3.2	8.9	37.5	135.2
2004 5/	3.8	*	*	*	*	*	*	1.0	3.1	8.9	35.7	123.7
2003 5/	4.3	*	*	*	*	*	0.8	1.0	3.7	9.7	41.9	146.5
2002	4.1	*	*	*	*	*	*	0.9	3.0	10.2	40.8	140.5
2001	4.6	*	*	*	*	*	*	1.4	2.9	11.6	41.3	166.4
2000	4.1	*	*	*	*	*	*	1.2	3.6	10.4	39.1	142.2
1999	4.0	*	*	*	*	*	*	1.2	2.9	10.2	36.4	145.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)												
All origins 3/, both sexes												
2006	9.2	8.1	0.6	0.2	0.2	0.5	1.2	3.1	8.9	28.2	73.4	136.6
2005	9.1	7.4	0.6	0.2	0.3	0.5	1.1	3.2	8.8	28.8	72.5	136.5
2004	9.3	6.5	0.7	0.2	0.3	0.4	1.2	3.2	9.5	29.9	74.2	139.4
2003	8.8	7.0	0.5	0.1	0.3	0.5	1.1	3.3	8.8	28.1	68.5	138.1
2002	8.7	6.9	0.5	0.2	0.3	0.5	1.2	3.2	9.2	28.7	68.0	133.5
2001	9.2	7.3	0.6	0.2	0.4	0.5	1.3	3.5	9.6	30.1	73.2	142.7
2000	9.1	7.2	0.6	0.2	0.3	0.6	1.4	3.5	9.9	29.6	72.0	141.7
1999	9.0	6.6	0.6	0.2	0.3	0.6	1.3	3.3	9.5	29.4	70.7	147.6
All origins 3/, male												
2006	9.3	9.2	0.7	0.2	0.3	0.5	1.2	3.3	10.3	35.0	93.1	163.7
2005	9.0	8.9	0.7	0.2	0.3	0.4	1.0	3.5	10.2	34.9	90.7	157.7
2004	9.3	6.6	0.7	0.2	0.3	0.4	1.2	3.5	10.8	36.9	93.9	168.8
2003	8.7	8.0	0.6	0.2	0.4	0.6	1.1	3.5	10.2	34.1	85.3	165.0
2002	8.5	7.9	0.7	0.2	0.3	0.4	1.3	3.3	10.5	35.3	83.9	156.6
2001	8.9	9.0	0.6	0.2	0.4	0.5	1.4	3.8	10.9	36.5	89.3	169.1
2000	9.0	7.9	0.6	0.2	0.4	0.6	1.5	3.8	11.4	35.7	90.5	179.7
1999	8.9	7.5	0.7	0.2	0.3	0.6	1.4	3.5	10.6	36.2	89.9	182.3
All origins 3/, female												
2006	9.2	6.9	0.5	0.2	0.2	0.5	1.2	3.0	7.5	22.6	59.9	124.0
2005	9.2	5.8	0.5	0.1	0.2	0.5	1.1	2.9	7.6	23.7	60.2	126.7
2004	9.3	6.4	0.6	0.1	0.3	0.4	1.2	3.0	8.4	24.1	61.0	126.1
2003	8.9	5.9	0.5	0.1	0.3	0.5	1.1	3.0	7.4	23.1	57.2	126.2
2002	8.8	5.8	0.4	0.2	0.3	0.5	1.1	3.1	8.0	23.1	57.5	123.4
2001	9.4	5.5	0.6	0.2	0.3	0.6	1.2	3.1	8.4	24.8	62.6	131.6
2000	9.2	6.4	0.6	0.2	0.3	0.6	1.2	3.2	8.6	24.7	59.9	126.2
1999	9.1	5.8	0.5	0.2	0.2	0.7	1.1	3.1	8.5	23.7	58.3	133.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)												
Hispanic, both sexes												
2006	3.6	6.3	*	*	0.3	0.3	0.7	2.6	7.6	25.4	59.5	106.8
2005	3.7	5.8	*	*	0.3	0.4	0.7	2.6	8.8	26.9	67.7	104.0
2004	3.6	5.8	*	*	*	*	0.9	2.4	7.8	27.2	71.0	112.7
2003	3.4	6.6	*	*	*	0.4	0.9	2.5	8.9	25.5	58.2	118.7
2002	3.3	5.2	*	*	*	0.4	0.7	2.7	8.3	26.1	58.0	112.0
2001	3.4	5.0	*	*	*	*	0.7	3.1	8.9	25.3	65.5	134.4
2000	3.5	5.6	*	*	*	0.5	1.1	2.6	9.8	25.4	64.2	135.4
1999	3.4	5.0	*	*	*	0.4	0.9	3.2	9.6	26.0	64.0	134.0
Hispanic, male												
2006	3.4	7.1	*	*	*	*	0.9	2.7	10.1	28.9	67.3	112.8
2005	3.5	6.3	*	*	*	0.5	0.7	3.0	9.9	29.1	81.8	114.6
2004	3.4	5.6	*	*	*	*	0.8	2.9	9.1	31.1	82.3	112.1
2003	3.3	6.6	*	*	*	*	1.1	3.1	11.3	30.8	63.3	127.8
2002	3.2	4.9	*	*	*	*	0.8	2.6	9.8	34.6	64.0	108.8
2001	3.4	5.0	*	*	*	*	0.9	3.4	10.6	30.2	78.3	167.6
2000	3.3	5.8	*	*	*	*	1.3	3.0	11.9	29.3	81.1	133.0
1999	3.3	5.2	*	*	*	*	1.1	3.1	11.6	30.5	82.4	161.8
Hispanic, female												
2006	3.7	5.5	*	*	*	*	*	2.6	5.3	22.7	54.0	103.6
2005	3.9	5.3	*	*	*	*	0.8	2.2	7.8	25.2	57.9	98.4
2004	3.9	6.1	*	*	*	*	0.9	1.9	6.6	24.1	63.2	113.1
2003	3.6	6.6	*	*	*	*	*	1.8	6.7	21.2	54.8	114.1
2002	3.4	5.4	*	*	*	*	*	2.8	6.9	19.3	54.0	113.6
2001	3.5	5.0	*	*	*	*	*	2.8	7.5	21.4	56.8	118.2
2000	3.6	5.3	*	*	*	*	0.9	2.2	7.8	22.2	52.8	136.5
1999	3.5	*	*	*	*	*	*	3.3	7.8	22.4	51.7	120.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)												
Non-Hispanic 4/, both sexes												
2006	10.2	8.6	0.7	0.2	0.2	0.6	1.3	3.2	9.0	28.4	74.1	138.1
2005	10.0	7.8	0.6	0.2	0.3	0.5	1.1	3.3	8.8	28.9	72.7	138.0
2004	10.2	6.7	0.7	0.2	0.3	0.5	1.3	3.3	9.6	30.1	74.3	140.5
2003	9.6	7.0	0.6	0.1	0.4	0.6	1.2	3.3	8.7	28.2	68.9	138.7
2002	9.5	7.3	0.5	0.2	0.3	0.5	1.2	3.2	9.2	28.8	68.4	134.1
2001	10.0	7.8	0.6	0.2	0.4	0.6	1.3	3.5	9.7	30.3	73.3	142.7
2000	9.8	7.5	0.6	0.2	0.4	0.7	1.4	3.6	9.9	29.8	72.1	141.6
1999	9.7	6.9	0.6	0.2	0.3	0.7	1.3	3.3	9.4	29.5	70.8	147.7
Non-Hispanic 4/, male												
2006	10.4	9.8	0.8	0.2	0.3	0.5	1.3	3.3	10.3	35.4	94.7	166.4
2005	10.0	9.6	0.8	0.2	0.3	0.4	1.1	3.6	10.2	35.3	91.0	159.9
2004	10.3	6.9	0.8	0.2	0.3	0.5	1.3	3.6	10.9	37.2	94.4	171.6
2003	9.5	8.3	0.6	0.2	0.4	0.6	1.1	3.5	10.1	34.2	86.3	166.5
2002	9.4	8.7	0.7	0.2	0.3	0.4	1.3	3.4	10.5	35.3	84.7	158.5
2001	9.8	10.0	0.6	0.2	0.4	0.6	1.4	3.9	10.9	36.7	89.5	169.1
2000	9.8	8.4	0.6	0.2	0.4	0.6	1.5	3.9	11.3	36.0	90.7	181.3
1999	9.6	7.9	0.8	0.2	0.4	0.7	1.4	3.6	10.5	36.4	89.9	182.7
Non-Hispanic 4/, female												
2006	10.0	7.3	0.5	0.2	0.2	0.6	1.3	3.0	7.7	22.5	60.1	124.9
2005	10.1	5.9	0.4	0.2	0.2	0.5	1.2	3.0	7.5	23.6	60.3	128.0
2004	10.1	6.5	0.6	0.1	0.3	0.5	1.3	3.1	8.5	24.1	60.8	126.6
2003	9.7	5.7	0.5	*	0.3	0.5	1.2	3.2	7.5	23.2	57.3	126.4
2002	9.6	5.9	0.4	0.2	0.3	0.5	1.2	3.1	8.0	23.3	57.5	123.6
2001	10.2	5.5	0.6	0.2	0.3	0.6	1.3	3.1	8.5	25.0	62.7	131.8
2000	9.9	6.6	0.5	0.2	0.3	0.7	1.3	3.3	8.6	24.7	60.1	125.5
1999	9.8	5.9	0.5	0.2	0.2	0.7	1.2	3.1	8.5	23.8	58.4	133.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)												
Non-Hispanic white, both sexes												
2006 5/	11.4	5.4	0.6	0.2	0.2	0.5	1.2	2.8	8.6	29.2	77.0	142.9
2005 5/	11.1	6.0	0.5	0.2	0.2	0.4	1.0	2.9	8.5	29.4	74.8	141.8
2004 5/	11.4	4.5	0.6	0.1	0.3	0.4	1.1	2.9	9.4	30.9	76.9	145.4
2003 5/	10.7	4.9	0.4	0.1	0.3	0.5	0.9	3.0	8.5	29.1	71.3	143.7
2002	10.5	4.7	0.4	0.2	0.3	0.4	1.1	2.9	8.8	29.3	70.3	138.6
2001	11.0	4.8	0.5	0.2	0.3	0.5	1.1	3.0	9.3	30.9	75.6	145.9
2000	10.8	5.4	0.5	0.2	0.3	0.6	1.2	3.2	9.6	30.2	73.8	145.7
1999	10.6	4.9	0.6	0.2	0.3	0.6	1.1	3.0	9.1	29.7	72.3	150.7
Non-Hispanic white, male												
2006 5/	11.6	5.6	0.7	0.2	0.2	0.5	1.2	3.1	10.1	36.3	98.2	171.5
2005 5/	11.0	7.3	0.7	0.2	0.2	0.4	1.0	3.2	9.7	36.1	92.9	162.2
2004 5/	11.4	4.6	0.6	*	0.3	0.4	1.0	3.2	10.6	38.1	97.3	177.3
2003 5/	10.6	5.5	0.4	0.2	0.3	0.6	1.0	3.2	9.7	35.2	88.8	172.0
2002	10.3	5.6	0.5	*	0.3	0.4	1.1	3.0	10.1	36.0	87.0	164.1
2001	10.8	6.2	0.4	0.2	0.4	0.4	1.3	3.4	10.5	37.7	92.4	173.4
2000	10.7	6.6	0.5	0.2	0.3	0.6	1.3	3.5	11.0	36.4	92.6	185.9
1999	10.5	5.6	0.6	0.2	0.4	0.7	1.2	3.2	10.0	37.0	92.0	186.1
Non-Hispanic white, female												
2006 5/	11.2	5.2	*	*	*	0.6	1.1	2.6	7.2	23.1	62.3	129.6
2005 5/	11.2	4.8	*	0.2	0.2	0.4	1.0	2.6	7.4	23.6	62.4	132.5
2004 5/	11.3	4.4	0.5	*	0.3	0.4	1.1	2.7	8.3	24.7	63.0	131.0
2003 5/	10.8	4.4	*	*	0.3	0.4	0.9	2.7	7.3	23.9	59.4	131.2
2002	10.7	3.7	*	0.2	0.2	0.5	1.0	2.7	7.7	23.6	59.1	127.6
2001	11.3	3.4	0.5	0.2	0.3	0.6	0.9	2.6	8.2	25.1	64.4	134.5
2000	11.0	4.1	0.5	*	0.3	0.6	1.1	3.0	8.4	25.0	61.5	129.5
1999	10.7	4.0	0.5	0.2	0.2	0.5	0.9	2.7	8.2	23.6	59.4	136.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)												
Non-Hispanic black, both sexes												
2006 5/	6.4	20.3	0.9	*	0.5	0.8	2.2	6.0	13.4	26.0	51.9	96.6
2005 5/	6.6	15.7	0.9	*	0.4	0.9	2.2	6.3	12.3	29.6	57.1	104.0
2004 5/	6.4	15.3	1.1	*	0.4	0.9	2.7	6.4	13.6	27.9	53.9	94.4
2003 5/	6.0	15.3	1.1	*	0.7	1.0	2.5	6.6	12.3	24.6	51.3	91.0
2002	6.2	17.3	1.2	0.4	0.7	0.9	2.4	6.2	13.8	27.1	52.3	90.7
2001	6.9	19.0	1.1	*	0.7	1.3	2.9	7.5	13.9	29.6	56.9	110.8
2000	6.7	17.7	0.8	*	0.7	1.1	2.7	6.9	13.7	30.6	58.6	99.8
1999	6.7	15.8	1.1	*	0.4	1.4	2.9	6.6	13.6	31.3	58.7	112.8
Non-Hispanic black, male												
2006 5/	6.4	24.7	*	*	0.6	0.8	2.3	5.8	14.4	33.4	67.6	119.1
2005 5/	6.7	20.5	*	*	*	0.8	2.0	6.9	15.4	34.4	76.5	134.2
2004 5/	6.5	16.4	*	*	*	1.1	3.2	6.9	15.5	34.8	69.9	117.9
2003 5/	6.2	18.6	*	*	0.9	1.1	2.3	6.4	15.3	31.4	69.0	116.1
2002	6.3	20.3	*	*	*	0.9	2.8	6.5	16.4	32.5	67.1	102.8
2001	6.6	24.3	*	*	*	1.4	2.6	7.8	15.7	32.9	69.0	131.7
2000	6.8	17.1	*	*	0.8	1.2	3.2	8.0	15.6	36.2	74.0	135.0
1999	6.6	18.0	*	*	*	1.1	2.9	7.3	15.9	35.6	73.5	146.2
Non-Hispanic black, female												
2006 5/	6.4	15.8	*	*	*	0.8	2.1	6.2	12.5	20.7	43.1	87.4
2005 5/	6.5	10.7	*	*	*	1.0	2.4	5.9	9.8	26.2	46.1	91.9
2004 5/	6.3	14.2	*	*	*	0.8	2.3	6.0	12.2	23.0	44.9	85.1
2003 5/	5.9	11.9	*	*	*	1.0	2.7	6.6	9.8	19.8	41.4	81.3
2002	6.2	14.2	*	*	0.7	0.9	2.0	5.9	11.7	23.3	43.9	86.1
2001	7.2	13.4	*	*	*	1.2	3.2	7.2	12.4	27.2	50.0	103.1
2000	6.6	18.3	*	*	*	1.0	2.3	6.0	12.1	26.6	49.8	86.8
1999	6.9	13.6	*	*	*	1.7	2.9	5.9	11.9	28.2	50.4	100.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
All origins 3/, both sexes												
2006	1.1	*	*	*	*	*	0.2	0.7	1.2	2.7	7.2	20.7
2005	1.2	*	*	*	*	0.1	0.2	0.7	1.2	2.9	8.0	21.4
2004	1.3	*	*	*	*	*	0.2	0.6	1.3	3.2	8.9	24.4
2003	1.3	*	*	*	*	0.1	0.2	0.7	1.4	3.2	9.9	26.4
2002	1.4	*	*	*	*	0.1	0.3	0.7	1.5	3.6	10.1	28.3
2001	1.6	*	*	*	*	0.1	0.3	0.7	1.7	3.9	11.4	33.4
2000	1.6	*	*	*	*	*	0.3	0.8	1.8	4.1	11.4	35.6
1999	1.6	*	*	*	*	0.1	0.3	0.7	1.8	4.5	12.3	34.6
All origins 3/, male												
2006	1.1	*	*	*	*	*	0.2	0.9	1.6	3.2	8.2	22.0
2005	1.2	*	*	*	*	*	0.3	1.0	1.7	3.6	9.7	22.4
2004	1.2	*	*	*	*	*	0.3	0.9	1.6	3.6	10.2	24.6
2003	1.3	*	*	*	*	*	0.3	1.0	1.8	4.1	11.8	28.9
2002	1.4	*	*	*	*	*	0.3	0.9	1.9	4.6	12.2	31.6
2001	1.5	*	*	*	*	0.1	0.4	1.0	2.2	4.8	13.0	34.9
2000	1.6	*	*	*	*	*	0.4	1.2	2.4	5.3	13.7	40.7
1999	1.6	*	*	*	*	*	0.3	0.9	2.3	6.1	14.8	37.5
All origins 3/, female												
2006	1.1	*	*	*	*	*	0.1	0.4	0.8	2.3	6.5	20.1
2005	1.2	*	*	*	*	*	0.2	0.4	0.8	2.4	6.8	21.0
2004	1.3	*	*	*	*	*	0.1	0.4	1.0	2.8	8.0	24.3
2003	1.4	*	*	*	*	*	0.2	0.4	1.0	2.4	8.7	25.3
2002	1.5	*	*	*	*	*	0.2	0.5	1.2	2.9	8.7	26.9
2001	1.7	*	*	*	*	*	0.1	0.5	1.1	3.2	10.4	32.7
2000	1.6	*	*	*	*	*	0.1	0.5	1.3	3.0	10.0	33.5
1999	1.7	*	*	*	*	*	0.2	0.5	1.3	3.3	10.6	33.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
Hispanic, both sexes												
2006	0.4	*	*	*	*	*	*	0.5	0.8	2.5	5.6	19.6
2005	0.4	*	*	*	*	*	*	0.6	0.9	2.2	7.0	16.7
2004	0.4	*	*	*	*	*	*	*	1.4	2.4	7.2	16.6
2003	0.5	*	*	*	*	*	*	0.8	1.2	3.4	8.7	17.1
2002	0.5	*	*	*	*	*	*	0.5	1.6	3.3	8.8	20.5
2001	0.4	*	*	*	*	*	*	*	1.2	2.9	9.7	23.8
2000	0.5	*	*	*	*	*	*	*	1.6	3.3	9.9	30.5
1999	0.5	*	*	*	*	*	*	*	1.5	3.6	9.2	26.7
Hispanic, male												
2006	0.4	*	*	*	*	*	*	*	*	3.7	7.0	*
2005	0.4	*	*	*	*	*	*	*	*	*	9.4	*
2004	0.4	*	*	*	*	*	*	*	*	*	9.7	*
2003	0.6	*	*	*	*	*	*	1.1	*	5.5	11.5	*
2002	0.6	*	*	*	*	*	*	*	2.5	5.4	11.1	*
2001	0.5	*	*	*	*	*	*	*	*	*	16.5	*
2000	0.5	*	*	*	*	*	*	*	2.9	4.4	*	*
1999	0.6	*	*	*	*	*	*	*	*	4.8	14.1	*
Hispanic, female												
2006	0.4	*	*	*	*	*	*	*	*	*	4.6	21.0
2005	0.4	*	*	*	*	*	*	*	*	*	5.3	18.9
2004	0.4	*	*	*	*	*	*	*	*	*	5.4	15.8
2003	0.4	*	*	*	*	*	*	*	*	*	6.8	20.3
2002	0.4	*	*	*	*	*	*	*	*	*	7.3	17.4
2001	0.4	*	*	*	*	*	*	*	*	*	*	25.1
2000	0.5	*	*	*	*	*	*	*	*	*	10.9	29.7
1999	0.4	*	*	*	*	*	*	*	*	*	*	22.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
Non-Hispanic 4/, both sexes												
2006	1.2	*	*	*	*	*	0.2	0.7	1.2	2.8	7.3	20.7
2005	1.3	*	*	*	*	0.1	0.3	0.7	1.3	3.0	8.0	21.6
2004	1.4	*	*	*	*	*	0.2	0.7	1.3	3.2	8.9	24.6
2003	1.5	*	*	*	*	0.1	0.2	0.7	1.4	3.2	9.9	26.7
2002	1.6	*	*	*	*	0.1	0.3	0.7	1.5	3.7	10.2	28.6
2001	1.7	*	*	*	*	0.1	0.3	0.7	1.7	4.0	11.5	33.6
2000	1.8	*	*	*	*	*	0.3	0.8	1.8	4.1	11.5	35.7
1999	1.8	*	*	*	*	*	0.3	0.7	1.8	4.6	12.4	34.8
Non-Hispanic 4/, male												
2006	1.2	*	*	*	*	*	0.2	0.9	1.7	3.2	8.3	22.2
2005	1.3	*	*	*	*	*	0.4	1.0	1.7	3.7	9.7	23.0
2004	1.3	*	*	*	*	*	0.3	0.9	1.6	3.7	10.2	24.8
2003	1.4	*	*	*	*	*	0.3	0.9	1.8	4.0	11.7	29.6
2002	1.5	*	*	*	*	*	0.4	0.9	1.9	4.5	12.3	31.7
2001	1.6	*	*	*	*	0.1	0.4	1.0	2.2	4.8	12.8	35.5
2000	1.8	*	*	*	*	*	0.4	1.2	2.4	5.4	13.9	41.0
1999	1.7	*	*	*	*	*	0.4	1.0	2.3	6.1	14.8	37.7
Non-Hispanic 4/, female												
2006	1.3	*	*	*	*	*	0.2	0.4	0.9	2.4	6.6	20.0
2005	1.3	*	*	*	*	*	0.2	0.4	0.9	2.4	6.9	21.0
2004	1.5	*	*	*	*	*	0.2	0.5	1.0	2.8	8.1	24.6
2003	1.5	*	*	*	*	*	0.2	0.4	1.0	2.5	8.7	25.4
2002	1.6	*	*	*	*	*	0.2	0.5	1.2	3.0	8.8	27.3
2001	1.8	*	*	*	*	*	0.2	0.5	1.2	3.3	10.6	32.8
2000	1.8	*	*	*	*	*	0.1	0.5	1.3	3.1	9.9	33.5
1999	1.9	*	*	*	*	*	0.2	0.5	1.3	3.3	10.8	33.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
Non-Hispanic white, both sexes												
2006 5/	1.3	*	*	*	*	*	0.1	0.6	1.2	2.6	7.3	21.2
2005 5/	1.4	*	*	*	*	*	0.2	0.6	1.2	2.9	8.0	21.8
2004 5/	1.5	*	*	*	*	*	0.2	0.6	1.2	3.1	9.0	25.4
2003 5/	1.6	*	*	*	*	*	0.2	0.6	1.3	3.2	9.8	27.3
2002	1.7	*	*	*	*	*	0.2	0.6	1.4	3.6	10.4	29.4
2001	1.9	*	*	*	*	*	0.2	0.7	1.5	3.9	11.6	34.7
2000	1.9	*	*	*	*	*	0.2	0.7	1.8	4.0	11.6	36.3
1999	2.0	*	*	*	*	*	0.2	0.6	1.7	4.5	12.4	35.6
Non-Hispanic white, male												
2006 5/	1.2	*	*	*	*	*	0.1	0.7	1.6	2.9	8.2	22.2
2005 5/	1.4	*	*	*	*	*	0.3	0.8	1.6	3.5	9.7	23.1
2004 5/	1.4	*	*	*	*	*	0.2	0.8	1.5	3.6	10.0	25.7
2003 5/	1.5	*	*	*	*	*	0.2	0.8	1.6	4.0	11.3	29.8
2002	1.6	*	*	*	*	*	0.3	0.8	1.7	4.4	12.4	32.6
2001	1.7	*	*	*	*	*	0.3	0.8	2.0	4.5	12.7	36.3
2000	1.9	*	*	*	*	*	0.3	1.1	2.3	5.2	13.7	41.0
1999	1.8	*	*	*	*	*	0.3	0.8	2.1	5.9	14.9	38.6
Non-Hispanic white, female												
2006 5/	1.4	*	*	*	*	*	0.2	0.4	0.8	2.4	6.7	20.7
2005 5/	1.4	*	*	*	*	*	0.2	0.4	0.9	2.4	6.8	21.2
2004 5/	1.7	*	*	*	*	*	0.1	0.4	1.0	2.7	8.3	25.3
2003 5/	1.7	*	*	*	*	*	0.2	0.4	1.0	2.6	8.8	26.2
2002	1.8	*	*	*	*	*	0.2	0.5	1.1	3.0	9.0	27.9
2001	2.1	*	*	*	*	*	0.1	0.5	1.1	3.4	10.9	34.0
2000	2.0	*	*	*	*	*	0.1	0.4	1.3	3.1	10.2	34.4
1999	2.1	*	*	*	*	*	0.2	0.4	1.3	3.2	10.8	34.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
Non-Hispanic black, both sexes												
2006 5/	0.9	*	*	*	*	*	0.4	1.4	2.1	3.8	7.3	14.9
2005 5/	1.0	*	*	*	*	*	0.5	1.3	1.7	4.0	7.3	17.5
2004 5/	1.0	*	*	*	*	*	0.5	1.3	2.1	4.3	7.8	16.1
2003 5/	1.0	*	*	*	*	*	0.4	1.5	2.2	3.2	11.1	17.3
2002	1.0	*	*	*	*	*	0.6	1.2	2.6	4.2	7.9	18.8
2001	1.1	*	*	*	*	*	0.6	1.4	2.8	4.5	9.0	20.9
2000	1.2	*	*	*	*	*	0.5	1.7	2.6	4.9	10.1	26.7
1999	1.3	*	*	*	*	*	0.8	1.5	2.8	5.7	11.0	25.5
Non-Hispanic black, male												
2006 5/	1.2	*	*	*	*	*	*	2.3	2.9	5.8	9.5	*
2005 5/	1.2	*	*	*	*	*	0.8	2.2	2.7	5.9	8.4	20.0
2004 5/	1.1	*	*	*	*	*	*	1.8	3.0	4.7	11.1	*
2003 5/	1.3	*	*	*	*	*	*	2.3	3.4	5.2	16.4	*
2002	1.2	*	*	*	*	*	0.8	2.0	3.9	6.0	11.1	*
2001	1.4	*	*	*	*	*	1.0	2.3	4.3	7.0	11.7	24.3
2000	1.4	*	*	*	*	*	0.8	2.4	3.8	7.2	13.9	31.7
1999	1.5	*	*	*	*	*	1.2	2.4	4.5	8.3	12.2	28.5
Non-Hispanic black, female												
2006 5/	0.8	*	*	*	*	*	*	*	1.5	2.3	6.2	14.2
2005 5/	0.8	*	*	*	*	*	*	*	*	2.7	6.7	16.5
2004 5/	0.9	*	*	*	*	*	*	0.9	1.3	4.1	6.0	16.9
2003 5/	0.8	*	*	*	*	*	*	*	*	*	8.1	16.1
2002	0.9	*	*	*	*	*	*	*	1.6	2.9	6.1	18.4
2001	0.9	*	*	*	*	*	*	*	1.5	2.7	7.4	19.7
2000	1.0	*	*	*	*	*	*	1.1	1.7	3.3	8.0	24.9
1999	1.1	*	*	*	*	*	*	*	*	3.9	10.3	24.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
All origins 3/, both sexes												
2006	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	0.8	1.7
2005	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.8	2.1
2004	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.7
2003	0.2	*	*	*	*	*	0.1	0.1	0.2	0.3	1.0	1.7
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.9	2.0
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.1
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.6
1999	0.1	*	*	0.1	*	*	0.1	0.1	0.2	0.4	0.8	1.5
All origins 3/, male												
2006	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.2	2.3
2005	0.2	*	*	*	*	*	*	*	0.3	0.6	1.1	2.8
2004	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.3	2.5
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.4
2002	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.3	2.2
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.8
2000	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.0	2.9
All origins 3/, female												
2006	0.1	*	*	*	*	*	*	*	0.2	0.3	0.4	1.5
2005	0.1	*	*	*	*	*	*	*	0.1	0.2	0.7	1.8
2004	0.1	*	*	*	*	*	*	*	0.2	0.4	0.9	1.4
2003	0.1	*	*	*	*	*	*	0.1	*	*	0.7	1.3
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.7	2.0
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
Hispanic, both sexes												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
Non-Hispanic 4/, both sexes												
2006	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	1.7
2005	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.2
2004	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8
2003	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.6
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	2.1
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.2
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.7
1999	0.1	*	*	*	*	*	0.0	0.1	0.2	0.4	0.8	1.5
Non-Hispanic 4/, male												
2006	0.2	*	*	*	*	*	*	*	0.2	0.7	1.2	2.3
2005	0.2	*	*	*	*	*	*	*	0.3	0.7	1.1	3.0
2004	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.5
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.2
2002	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.3	2.3
2001	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.0	3.0
2000	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999	0.2	*	*	*	*	*	*	0.2	0.3	0.4	1.0	3.0
Non-Hispanic 4/, female												
2006	0.1	*	*	*	*	*	*	*	0.2	0.3	0.4	1.5
2005	0.1	*	*	*	*	*	*	*	*	0.2	0.7	1.8
2004	0.1	*	*	*	*	*	*	*	0.2	0.4	0.9	1.5
2003	0.1	*	*	*	*	*	*	*	*	*	0.8	1.4
2002	0.2	*	*	*	*	*	*	*	*	0.3	0.7	2.0
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
Non-Hispanic white, both sexes												
2006 5/	0.2	*	*	*	*	*	*	*	0.1	0.5	0.7	1.8
2005 5/	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.3
2004 5/	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8
2003 5/	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.0	1.7
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.0	2.1
2001	0.2	*	*	*	*	*	*	0.1	0.1	0.5	0.7	2.1
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.8
1999	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.8	1.6
Non-Hispanic white, male												
2006 5/	0.2	*	*	*	*	*	*	*	0.2	0.7	1.1	2.4
2005 5/	0.2	*	*	*	*	*	*	*	0.3	0.7	1.1	3.1
2004 5/	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	2.3
2003 5/	0.2	*	*	*	*	*	*	*	0.2	0.5	1.5	2.4
2002	0.2	*	*	*	*	*	*	0.2	0.3	0.6	1.4	2.1
2001	0.2	*	*	*	*	*	*	*	0.2	0.7	1.0	2.6
2000	0.2	*	*	*	*	*	*	*	0.3	0.4	1.2	2.1
1999	0.2	*	*	*	*	*	*	0.2	0.2	0.4	0.9	3.1
Non-Hispanic white, female												
2006 5/	0.1	*	*	*	*	*	*	*	*	0.3	0.4	1.5
2005 5/	0.1	*	*	*	*	*	*	*	*	*	0.7	1.9
2004 5/	0.2	*	*	*	*	*	*	*	*	0.4	0.8	1.6
2003 5/	0.1	*	*	*	*	*	*	*	*	*	0.8	1.3
2002	0.2	*	*	*	*	*	*	*	*	0.3	0.7	2.1
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.4	2.0
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.7	1.6
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
Non-Hispanic black, both sexes												
2006 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
All origins 3/, both sexes												
2006	0.6	0.8	*	*	*	*	0.1	0.2	0.6	1.2	3.9	12.1
2005	0.6	1.0	*	*	*	*	0.1	0.2	0.6	1.1	3.5	12.1
2004	0.5	1.0	*	*	*	*	0.1	0.2	0.4	1.3	3.9	11.0
2003	0.6	0.9	*	*	*	*	0.1	0.3	0.6	1.3	3.6	11.9
2002	0.6	0.7	*	*	*	*	0.1	0.2	0.5	1.2	3.8	12.8
2001	0.5	1.1	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.6
2000	0.5	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.1
1999	0.5	1.2	*	*	*	*	0.1	0.2	0.4	1.2	3.7	12.7
All origins 3/, male												
2006	0.5	0.9	*	*	*	*	0.1	0.2	0.5	1.3	4.1	13.5
2005	0.4	1.3	*	*	*	*	*	0.2	0.6	1.0	3.3	12.7
2004	0.5	1.0	*	*	*	*	0.1	0.2	0.4	1.4	3.8	11.3
2003	0.5	1.0	*	*	*	*	*	0.3	0.5	1.3	3.9	12.3
2002	0.4	*	*	*	*	*	*	0.2	0.4	1.2	3.7	11.7
2001	0.4	1.4	*	*	*	*	*	0.2	0.5	1.1	3.5	12.0
2000	0.4	1.4	*	*	*	*	*	0.2	0.4	1.1	4.0	12.6
1999	0.4	1.3	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.2
All origins 3/, female												
2006	0.7	*	*	*	*	*	*	0.2	0.6	1.2	3.8	11.4
2005	0.7	*	*	*	*	*	0.1	0.2	0.6	1.2	3.6	11.8
2004	0.6	1.0	*	*	*	*	*	0.2	0.5	1.2	4.0	10.8
2003	0.6	*	*	*	*	*	*	0.2	0.6	1.3	3.4	11.8
2002	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.9	13.3
2001	0.6	*	*	*	*	*	*	0.2	0.5	1.2	3.6	12.8
2000	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.3
1999	0.6	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
Hispanic, both sexes												
2006	0.2	*	*	*	*	*	*	*	*	*	3.5	*
2005	0.2	*	*	*	*	*	*	*	*	*	*	11.1
2004	0.2	*	*	*	*	*	*	*	*	*	3.5	*
2003	0.2	*	*	*	*	*	*	*	*	*	3.6	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	4.3	12.5
2000	0.2	*	*	*	*	*	*	*	*	*	4.5	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.3	*	*	*	*	*	*	*	*	*	*	*
2004	0.3	*	*	*	*	*	*	*	*	*	*	*
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
Non-Hispanic 4/, both sexes												
2006	0.6	0.9	*	*	*	*	0.1	0.2	0.6	1.2	3.9	12.3
2005	0.6	1.1	*	*	*	*	0.1	0.2	0.6	1.1	3.5	12.1
2004	0.6	0.9	*	*	*	*	0.1	0.2	0.5	1.3	3.9	11.2
2003	0.6	1.1	*	*	*	*	0.1	0.2	0.6	1.3	3.6	12.1
2002	0.6	0.8	*	*	*	*	0.1	0.2	0.5	1.2	3.8	13.0
2001	0.6	1.0	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.5
2000	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.7	13.1
1999	0.6	1.3	*	*	*	*	0.1	0.1	0.5	1.1	3.7	12.7
Non-Hispanic 4/, male												
2006	0.6	*	*	*	*	*	*	0.2	0.6	1.3	4.1	13.9
2005	0.5	1.5	*	*	*	*	*	0.2	0.6	1.0	3.4	12.8
2004	0.5	*	*	*	*	*	0.1	0.2	0.5	1.4	3.9	11.6
2003	0.5	*	*	*	*	*	*	0.3	0.5	1.3	3.9	12.5
2002	0.5	*	*	*	*	*	*	0.2	0.4	1.3	3.7	11.9
2001	0.5	1.2	*	*	*	*	*	0.2	0.5	1.2	3.5	12.1
2000	0.5	1.4	*	*	*	*	*	0.2	0.4	1.1	3.9	12.8
1999	0.5	1.5	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.3
Non-Hispanic 4/, female												
2006	0.7	*	*	*	*	*	*	0.2	0.6	1.2	3.9	11.6
2005	0.7	*	*	*	*	*	0.1	0.2	0.6	1.2	3.6	11.8
2004	0.7	*	*	*	*	*	*	0.2	0.4	1.2	4.0	11.0
2003	0.7	*	*	*	*	*	*	0.2	0.6	1.3	3.3	12.0
2002	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.9	13.4
2001	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.5	12.7
2000	0.7	*	*	*	*	*	*	0.2	0.5	1.1	3.6	13.2
1999	0.7	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
Non-Hispanic white, both sexes												
2006 5/	0.7	0.9	*	*	*	*	0.1	0.2	0.6	1.2	3.9	12.8
2005 5/	0.7	1.1	*	*	*	*	0.1	0.2	0.6	1.1	3.6	12.5
2004 5/	0.7	1.0	*	*	*	*	0.1	0.2	0.4	1.3	4.1	11.4
2003 5/	0.7	1.1	*	*	*	*	*	0.2	0.5	1.3	3.6	12.4
2002	0.7	0.9	*	*	*	*	0.1	0.2	0.4	1.2	4.0	13.6
2001	0.6	0.9	*	*	*	*	*	0.2	0.5	1.2	3.5	12.8
2000	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.7	13.3
1999	0.6	1.4	*	*	*	*	0.1	0.1	0.5	1.1	3.7	13.0
Non-Hispanic white, male												
2006 5/	0.6	*	*	*	*	*	*	0.2	0.6	1.3	4.0	14.5
2005 5/	0.5	*	*	*	*	*	*	0.2	0.6	1.0	3.4	12.9
2004 5/	0.6	*	*	*	*	*	*	0.2	0.5	1.4	4.0	11.7
2003 5/	0.6	*	*	*	*	*	*	0.3	0.5	1.3	3.9	12.6
2002	0.5	*	*	*	*	*	*	0.2	0.4	1.2	3.8	12.2
2001	0.5	*	*	*	*	*	*	0.2	0.5	1.2	3.5	11.9
2000	0.5	*	*	*	*	*	*	0.2	0.4	1.1	4.0	13.0
1999	0.5	*	*	*	*	*	*	0.2	0.6	1.1	4.1	13.0
Non-Hispanic white, female												
2006 5/	0.8	*	*	*	*	*	*	0.2	0.6	1.2	3.9	12.0
2005 5/	0.8	*	*	*	*	*	*	0.2	0.6	1.2	3.8	12.3
2004 5/	0.8	*	*	*	*	*	*	0.2	0.4	1.2	4.1	11.3
2003 5/	0.8	*	*	*	*	*	*	0.2	0.5	1.3	3.3	12.4
2002	0.8	*	*	*	*	*	*	0.2	0.4	1.2	4.1	14.2
2001	0.8	*	*	*	*	*	*	0.2	0.5	1.1	3.5	13.1
2000	0.8	*	*	*	*	*	*	0.2	0.5	1.1	3.6	13.4
1999	0.7	*	*	*	*	*	*	*	0.4	1.1	3.4	12.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
Non-Hispanic black, both sexes												
2006 5/	0.5	*	*	*	*	*	*	*	0.7	1.8	4.6	9.6
2005 5/	0.4	*	*	*	*	*	*	*	0.9	1.5	3.2	9.3
2004 5/	0.4	*	*	*	*	*	*	*	0.7	1.7	3.2	9.5
2003 5/	0.4	*	*	*	*	*	*	*	1.0	1.4	4.0	9.2
2002	0.4	*	*	*	*	*	*	*	0.8	1.6	3.2	7.9
2001	0.4	*	*	*	*	*	*	*	0.9	1.5	3.6	11.2
2000	0.4	*	*	*	*	*	*	*	*	1.3	4.0	12.7
1999	0.4	*	*	*	*	*	*	*	*	1.5	4.2	10.0
Non-Hispanic black, male												
2006 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	0.5	*	*	*	*	*	*	*	*	*	4.3	8.6
2005 5/	0.4	*	*	*	*	*	*	*	*	*	*	8.0
2004 5/	0.4	*	*	*	*	*	*	*	*	*	3.2	8.8
2003 5/	0.5	*	*	*	*	*	*	*	*	*	3.9	8.3
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	9.0
2000	0.5	*	*	*	*	*	*	*	*	*	4.3	12.6
1999	0.5	*	*	*	*	*	*	*	*	*	4.2	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
All origins 3/, both sexes												
2006	9.2	*	*	*	0.1	0.8	5.8	17.8	22.8	26.0	29.0	19.4
2005	9.3	*	*	*	0.1	0.8	6.1	17.7	23.5	27.2	29.0	19.7
2004	9.2	*	*	*	*	0.8	6.3	18.0	22.6	27.7	28.8	19.7
2003	9.5	*	*	*	*	0.9	6.8	18.3	23.0	29.5	30.0	20.1
2002	9.5	*	*	*	0.1	0.9	7.0	18.0	22.9	29.4	31.4	21.4
2001	9.5	*	*	*	0.1	1.0	7.4	18.5	22.7	30.0	30.2	22.2
2000	9.4	*	*	*	0.1	1.0	7.5	17.7	23.8	29.8	31.0	23.1
1999	9.4	*	*	*	0.1	1.0	7.3	17.4	23.7	30.6	31.9	23.2
All origins 3/, male												
2006	12.1	*	*	*	*	1.0	7.6	25.9	33.5	34.7	35.9	26.1
2005	12.3	*	*	*	*	1.0	8.1	26.0	34.8	35.8	36.4	27.6
2004	12.1	*	*	*	*	1.1	8.7	26.3	32.8	37.0	36.7	25.5
2003	12.5	*	*	*	*	1.2	9.2	27.5	33.8	39.1	36.9	27.5
2002	12.3	*	*	*	*	1.2	9.3	26.9	33.1	38.5	38.4	28.4
2001	12.4	*	*	*	*	1.3	9.8	27.5	33.1	40.2	36.8	32.0
2000	12.5	*	*	*	*	1.4	10.3	26.9	34.4	39.9	38.4	34.3
1999	12.5	*	*	*	*	1.3	10.3	26.9	34.6	41.6	39.6	33.1
All origins 3/, female												
2006	6.4	*	*	*	*	0.5	4.1	10.0	13.0	18.6	24.2	16.2
2005	6.4	*	*	*	*	0.6	4.1	9.7	12.9	19.9	23.9	16.0
2004	6.3	*	*	*	*	0.4	4.0	10.0	13.1	19.9	23.4	17.1
2003	6.5	*	*	*	*	0.5	4.4	9.4	13.1	21.4	25.4	16.8
2002	6.7	*	*	*	*	0.7	4.8	9.5	13.5	21.9	26.7	18.4
2001	6.7	*	*	*	*	0.7	5.0	9.8	13.1	21.5	25.9	18.1
2000	6.5	*	*	*	*	0.7	4.7	8.8	14.0	21.5	26.1	18.5
1999	6.4	*	*	*	*	0.8	4.4	8.3	13.7	21.6	26.9	19.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
Hispanic, both sexes												
2006	8.1	*	*	*	*	1.1	6.7	23.7	34.2	43.0	54.4	40.3
2005	8.3	*	*	*	*	1.0	7.3	23.5	36.9	48.8	54.0	34.7
2004	8.2	*	*	*	*	1.0	7.6	24.8	34.6	47.3	60.2	26.2
2003	8.5	*	*	*	*	1.4	7.9	24.3	38.4	51.1	62.0	31.1
2002	8.8	*	*	*	*	1.1	8.7	26.9	38.4	56.2	53.5	45.5
2001	8.9	*	*	*	*	1.2	9.7	28.0	39.0	54.4	60.4	36.9
2000	9.0	*	*	*	*	1.4	9.8	27.2	41.4	58.3	63.8	48.4
1999	8.8	*	*	*	*	1.3	10.5	26.6	39.0	56.3	59.4	51.2
Hispanic, male												
2006	11.0	*	*	*	*	1.4	9.9	37.4	54.1	56.9	61.0	51.9
2005	11.6	*	*	*	*	1.4	10.9	37.8	58.4	69.1	68.8	36.1
2004	11.1	*	*	*	*	1.5	11.6	40.8	52.4	59.4	70.8	29.4
2003	11.6	*	*	*	*	2.0	12.3	40.2	62.2	61.3	71.3	35.4
2002	12.2	*	*	*	*	1.6	13.5	44.2	60.0	77.8	64.4	45.2
2001	12.7	*	*	*	*	1.8	15.0	46.4	63.9	77.1	76.0	41.9
2000	12.7	*	*	*	*	2.3	15.0	45.6	66.1	83.4	80.1	60.5
1999	12.3	*	*	*	*	1.8	16.5	44.3	62.6	79.1	70.9	66.9
Hispanic, female												
2006	5.0	*	*	*	*	0.6	3.0	9.7	16.0	31.6	49.8	34.2
2005	4.8	*	*	*	*	*	3.2	8.9	17.1	32.4	43.7	33.9
2004	5.0	*	*	*	*	*	3.1	8.6	18.5	37.6	52.9	24.5
2003	5.2	*	*	*	*	0.6	3.2	8.3	17.0	42.9	55.6	28.9
2002	5.2	*	*	*	*	*	3.5	9.7	18.9	38.9	46.1	45.6
2001	5.0	*	*	*	*	*	4.0	9.9	17.0	36.5	49.9	34.4
2000	5.1	*	*	*	*	*	4.1	9.3	19.5	38.7	52.8	42.5
1999	5.0	*	*	*	*	*	4.1	9.3	18.1	38.4	51.7	43.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
Non-Hispanic 4/, both sexes												
2006	9.4	*	*	*	0.1	0.7	5.7	17.1	21.8	24.6	27.3	18.3
2005	9.4	*	*	*	0.1	0.7	5.9	17.0	22.3	25.5	27.4	18.9
2004	9.3	*	*	*	*	0.7	6.1	17.2	21.5	26.2	27.0	19.4
2003	9.6	*	*	*	*	0.8	6.6	17.6	21.7	27.9	28.2	19.6
2002	9.5	*	*	*	0.1	0.9	6.7	17.0	21.5	27.5	30.2	20.4
2001	9.5	*	*	*	0.1	0.9	7.1	17.5	21.3	28.2	28.7	21.6
2000	9.4	*	*	*	0.1	1.0	7.1	16.7	22.3	27.9	29.5	22.1
1999	9.5	*	*	*	*	1.0	6.9	16.5	22.5	29.0	30.7	22.1
Non-Hispanic 4/, male												
2006	12.3	*	*	*	*	0.9	7.1	24.5	31.5	32.9	34.2	24.6
2005	12.3	*	*	*	*	0.8	7.6	24.4	32.7	33.2	34.4	27.1
2004	12.3	*	*	*	*	1.0	8.2	24.6	31.0	35.3	34.8	25.3
2003	12.6	*	*	*	*	1.1	8.6	26.0	31.3	37.5	35.0	27.1
2002	12.2	*	*	*	*	1.1	8.6	24.9	30.7	35.7	37.0	27.7
2001	12.3	*	*	*	*	1.2	9.1	25.6	30.6	37.6	34.9	31.4
2000	12.4	*	*	*	*	1.2	9.6	25.0	31.8	37.1	36.5	33.0
1999	12.5	*	*	*	*	1.2	9.5	25.2	32.4	39.3	38.2	31.6
Non-Hispanic 4/, female												
2006	6.6	*	*	*	*	0.5	4.2	10.0	12.7	17.6	22.6	15.3
2005	6.6	*	*	*	*	0.6	4.2	9.8	12.5	18.9	22.7	15.2
2004	6.5	*	*	*	*	0.5	4.1	10.1	12.7	18.5	21.8	16.7
2003	6.7	*	*	*	*	0.5	4.6	9.5	12.7	19.8	23.7	16.3
2002	6.9	*	*	*	*	0.7	4.9	9.4	13.0	20.6	25.8	17.3
2001	6.9	*	*	*	*	0.7	5.1	9.8	12.7	20.4	24.7	17.5
2000	6.7	*	*	*	*	0.7	4.7	8.7	13.6	20.3	24.9	17.7
1999	6.6	*	*	*	*	0.8	4.4	8.1	13.3	20.6	25.8	18.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
Non-Hispanic white, both sexes												
2006 5/	10.3	*	*	*	*	0.7	5.9	17.6	22.1	25.7	28.5	19.0
2005 5/	10.2	*	*	*	*	0.6	6.2	17.1	22.3	26.5	28.2	19.5
2004 5/	10.1	*	*	*	*	0.7	6.2	17.4	21.3	27.1	28.3	19.9
2003 5/	10.3	*	*	*	*	0.8	6.8	17.5	21.5	28.9	29.7	20.3
2002	10.2	*	*	*	*	0.8	6.9	16.8	21.4	28.5	31.5	21.2
2001	10.1	*	*	*	*	0.9	7.0	17.0	21.0	28.9	29.9	22.5
2000	10.0	*	*	*	*	0.8	7.0	16.2	22.1	28.4	30.6	22.8
1999	9.9	*	*	*	*	0.9	6.6	15.8	21.8	29.5	31.7	22.6
Non-Hispanic white, male												
2006 5/	13.4	*	*	*	*	0.9	7.4	25.1	31.9	33.9	35.4	25.6
2005 5/	13.3	*	*	*	*	0.8	7.9	24.6	32.5	34.2	34.9	28.1
2004 5/	13.2	*	*	*	*	1.0	8.3	24.7	30.6	36.1	36.3	26.4
2003 5/	13.6	*	*	*	*	1.0	9.0	25.9	30.8	38.7	36.5	27.9
2002	13.2	*	*	*	*	1.0	8.8	24.7	30.5	36.6	38.7	28.7
2001	13.0	*	*	*	*	1.1	8.9	24.8	29.7	38.1	36.3	32.9
2000	13.1	*	*	*	*	1.1	9.4	24.4	31.5	37.7	37.8	34.1
1999	13.1	*	*	*	*	1.1	9.1	24.2	31.5	39.7	39.2	32.7
Non-Hispanic white, female												
2006 5/	7.2	*	*	*	*	0.5	4.3	10.2	12.9	18.5	23.6	15.9
2005 5/	7.2	*	*	*	*	0.5	4.4	9.7	12.6	19.9	23.6	15.6
2004 5/	7.1	*	*	*	*	0.5	4.2	10.1	12.5	19.4	22.8	17.0
2003 5/	7.2	*	*	*	*	0.5	4.6	9.3	12.6	20.5	25.2	17.0
2002	7.5	*	*	*	*	0.6	5.0	9.0	12.9	21.5	26.8	18.0
2001	7.4	*	*	*	*	0.7	5.1	9.3	12.8	21.1	25.7	18.2
2000	7.0	*	*	*	*	0.6	4.5	8.0	13.2	20.6	25.9	18.2
1999	6.9	*	*	*	*	0.7	4.1	7.5	12.7	21.0	26.7	18.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
Non-Hispanic black, both sexes												
2006 5/	6.1	*	*	*	*	0.4	4.6	15.8	22.4	19.0	16.3	10.9
2005 5/	6.7	*	*	*	*	0.6	4.3	18.0	25.3	20.5	19.3	12.0
2004 5/	6.8	*	*	*	*	0.4	5.6	18.7	25.9	22.0	14.4	12.1
2003 5/	7.1	*	*	*	*	0.8	5.8	20.2	26.1	24.1	14.1	12.5
2002	7.0	*	*	*	*	0.9	5.6	21.2	25.3	21.8	18.3	12.1
2001	7.7	*	*	*	*	0.9	7.5	22.9	26.3	25.3	18.2	10.6
2000	7.7	*	*	*	*	1.1	8.0	22.6	26.1	26.5	18.1	14.0
1999	8.0	*	*	*	*	1.0	8.2	23.0	29.7	26.9	21.7	16.8
Non-Hispanic black, male												
2006 5/	8.3	*	*	*	*	*	5.9	22.9	33.9	29.3	24.5	*
2005 5/	9.2	*	*	*	*	*	5.5	26.5	39.9	31.4	29.4	*
2004 5/	9.4	*	*	*	*	*	7.6	27.6	40.0	34.5	20.5	*
2003 5/	9.8	*	*	*	*	1.0	7.3	31.0	41.2	33.7	20.2	22.4
2002	9.3	*	*	*	*	1.1	6.9	30.7	38.2	32.4	22.0	*
2001	10.7	*	*	*	*	1.0	10.4	35.1	42.0	39.7	22.5	*
2000	10.3	*	*	*	*	1.1	10.7	33.4	38.5	38.2	25.0	*
1999	11.0	*	*	*	*	1.1	11.3	35.8	43.8	41.0	30.0	*
Non-Hispanic black, female												
2006 5/	4.2	*	*	*	*	*	3.5	9.7	13.2	11.6	11.7	9.0
2005 5/	4.4	*	*	*	*	*	3.2	10.8	13.6	12.7	13.5	10.7
2004 5/	4.4	*	*	*	*	*	3.8	11.0	14.5	13.1	10.9	11.2
2003 5/	4.6	*	*	*	*	*	4.6	10.9	14.0	17.3	10.6	8.7
2002	5.0	*	*	*	*	0.8	4.5	13.2	15.0	14.2	16.2	10.5
2001	4.9	*	*	*	*	0.7	5.0	12.5	13.7	15.1	15.8	9.8
2000	5.4	*	*	*	*	1.1	5.7	13.4	16.2	18.3	14.2	11.8
1999	5.4	*	*	*	*	1.0	5.5	12.1	18.5	16.9	17.0	15.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
All origins 3/, both sexes												
2006	4.4	*	*	*	*	0.6	4.0	10.5	12.1	9.4	5.9	2.2
2005	4.4	*	*	*	*	0.5	4.1	10.6	12.3	9.9	5.7	1.8
2004	4.3	*	*	*	*	0.6	4.2	10.6	11.7	10.0	5.5	2.1
2003	4.3	*	*	*	*	0.6	4.4	10.5	11.6	9.8	5.7	1.8
2002	4.2	*	*	*	*	0.6	4.4	10.3	11.6	10.1	5.7	2.1
2001	4.3	*	*	*	*	0.7	4.8	10.7	11.5	10.1	5.7	2.4
2000	4.3	*	*	*	*	0.7	4.9	10.3	12.2	10.3	6.1	2.2
1999	4.3	*	*	*	*	0.6	4.8	10.4	12.0	10.5	6.5	2.8
All origins 3/, male												
2006	6.4	*	*	*	*	0.9	5.1	15.5	18.9	15.7	10.2	4.7
2005	6.5	*	*	*	*	0.7	5.4	15.7	19.5	15.9	10.2	4.2
2004	6.3	*	*	*	*	0.8	5.8	15.5	18.2	16.5	10.1	4.6
2003	6.4	*	*	*	*	0.9	6.0	16.1	18.3	16.1	10.1	3.6
2002	6.2	*	*	*	*	0.8	5.8	15.6	18.3	16.5	9.7	4.7
2001	6.4	*	*	*	*	0.9	6.5	16.1	18.4	16.6	10.1	5.5
2000	6.5	*	*	*	*	1.0	6.8	16.0	19.0	17.3	11.5	5.3
1999	6.5	*	*	*	*	0.8	6.8	16.3	19.2	18.1	11.2	6.1
All origins 3/, female												
2006	2.4	*	*	*	*	0.4	2.8	5.8	5.8	4.2	3.0	1.1
2005	2.3	*	*	*	*	0.4	2.8	5.6	5.5	4.9	2.7	0.7
2004	2.3	*	*	*	*	0.3	2.7	5.7	5.6	4.5	2.3	1.0
2003	2.2	*	*	*	*	0.3	2.9	5.1	5.4	4.5	2.7	1.0
2002	2.3	*	*	*	*	0.4	3.0	5.2	5.4	4.8	3.1	1.0
2001	2.3	*	*	*	*	0.4	3.1	5.5	5.1	4.7	2.8	1.0
2000	2.2	*	*	*	*	0.4	3.1	4.8	6.0	4.5	2.5	1.0
1999	2.1	*	*	*	*	0.5	2.9	4.7	5.4	4.3	3.4	1.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
Hispanic, both sexes												
2006	4.3	*	*	*	*	0.9	4.6	15.1	19.3	16.9	14.5	*
2005	4.2	*	*	*	*	0.8	4.9	14.4	18.9	20.2	11.4	*
2004	4.1	*	*	*	*	0.7	5.3	15.2	17.2	16.1	14.6	*
2003	4.3	*	*	*	*	0.9	5.6	14.4	21.4	17.7	14.1	*
2002	4.3	*	*	*	*	0.7	5.5	16.1	19.9	20.8	10.5	*
2001	4.7	*	*	*	*	0.9	6.3	17.1	21.4	21.1	17.0	*
2000	4.8	*	*	*	*	1.0	6.9	16.8	23.4	22.9	15.4	*
1999	4.6	*	*	*	*	0.8	6.9	16.8	21.7	20.6	14.4	*
Hispanic, male												
2006	6.9	*	*	*	*	1.3	6.9	24.7	33.8	32.1	27.2	*
2005	6.8	*	*	*	*	1.2	7.6	23.9	33.4	38.1	23.7	*
2004	6.6	*	*	*	*	1.2	8.2	26.6	29.7	28.6	28.3	*
2003	6.9	*	*	*	*	1.3	8.8	25.5	38.4	29.0	26.5	*
2002	7.1	*	*	*	*	1.2	9.0	27.0	36.3	40.3	18.1	*
2001	7.7	*	*	*	*	1.5	10.0	29.3	38.9	41.9	30.4	*
2000	7.9	*	*	*	*	1.8	10.6	29.3	42.8	46.6	33.9	*
1999	7.4	*	*	*	*	1.2	10.8	29.1	40.0	39.5	27.1	*
Hispanic, female												
2006	1.6	*	*	*	*	*	2.1	5.3	6.0	4.6	5.5	*
2005	1.4	*	*	*	*	*	2.0	4.8	5.6	5.7	*	*
2004	1.4	*	*	*	*	*	2.0	3.6	5.9	6.1	5.2	*
2003	1.5	*	*	*	*	*	2.1	3.4	6.1	8.7	5.5	*
2002	1.4	*	*	*	*	*	1.8	5.1	5.1	5.3	*	*
2001	1.5	*	*	*	*	*	2.2	5.1	5.9	4.7	7.9	*
2000	1.4	*	*	*	*	*	2.9	4.5	6.2	4.3	*	*
1999	1.5	*	*	*	*	*	2.8	4.9	5.6	5.7	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
Non-Hispanic 4/, both sexes												
2006	4.4	*	*	*	*	0.5	3.8	10.0	11.4	8.8	5.4	1.9
2005	4.4	*	*	*	*	0.5	3.9	10.1	11.6	9.0	5.4	1.7
2004	4.3	*	*	*	*	0.5	4.1	10.0	11.2	9.5	4.9	2.1
2003	4.2	*	*	*	*	0.5	4.2	10.1	10.8	9.1	5.2	1.7
2002	4.2	*	*	*	*	0.6	4.2	9.7	10.8	9.3	5.5	1.9
2001	4.2	*	*	*	*	0.6	4.6	10.1	10.7	9.3	5.2	2.2
2000	4.2	*	*	*	*	0.6	4.7	9.7	11.3	9.5	5.7	2.2
1999	4.2	*	*	*	*	0.6	4.6	9.8	11.3	9.9	6.1	2.5
Non-Hispanic 4/, male												
2006	6.3	*	*	*	*	0.7	4.8	14.3	17.5	14.3	9.1	3.9
2005	6.4	*	*	*	*	0.6	5.0	14.6	18.2	14.1	9.4	3.8
2004	6.3	*	*	*	*	0.7	5.3	14.2	17.2	15.6	9.1	4.5
2003	6.2	*	*	*	*	0.8	5.5	15.0	16.6	15.1	9.2	3.4
2002	6.0	*	*	*	*	0.7	5.3	14.4	16.7	14.8	9.2	4.3
2001	6.2	*	*	*	*	0.8	5.9	14.8	16.7	14.9	9.2	5.1
2000	6.2	*	*	*	*	0.8	6.3	14.7	17.1	15.4	10.5	5.0
1999	6.4	*	*	*	*	0.7	6.2	15.1	17.7	16.7	10.5	5.5
Non-Hispanic 4/, female												
2006	2.5	*	*	*	*	0.3	2.9	5.8	5.8	4.1	2.9	1.0
2005	2.5	*	*	*	*	0.4	2.9	5.7	5.5	4.8	2.7	0.8
2004	2.4	*	*	*	*	0.3	2.8	5.9	5.6	4.3	2.2	1.0
2003	2.3	*	*	*	*	0.3	3.0	5.3	5.4	4.1	2.6	1.0
2002	2.4	*	*	*	*	0.5	3.1	5.2	5.4	4.8	3.0	0.9
2001	2.4	*	*	*	*	0.5	3.2	5.5	5.0	4.7	2.5	1.0
2000	2.3	*	*	*	*	0.4	3.1	4.8	5.9	4.5	2.5	1.0
1999	2.2	*	*	*	*	0.5	2.9	4.7	5.3	4.2	3.3	1.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
Non-Hispanic white, both sexes												
2006 5/	4.7	*	*	*	*	0.5	4.0	10.4	11.8	9.0	5.6	2.0
2005 5/	4.7	*	*	*	*	0.4	4.1	10.4	11.8	9.2	5.5	1.8
2004 5/	4.6	*	*	*	*	0.5	4.2	10.3	11.1	9.7	5.1	2.2
2003 5/	4.5	*	*	*	*	0.5	4.4	10.1	10.7	9.3	5.4	1.7
2002	4.4	*	*	*	*	0.5	4.3	9.7	10.8	9.5	5.5	1.9
2001	4.4	*	*	*	*	0.6	4.5	9.9	10.5	9.3	5.3	2.1
2000	4.3	*	*	*	*	0.6	4.5	9.4	11.1	9.5	5.6	2.2
1999	4.3	*	*	*	*	0.6	4.4	9.5	10.8	9.7	6.0	2.4
Non-Hispanic white, male												
2006 5/	6.9	*	*	*	*	0.7	5.1	14.9	18.0	14.5	9.3	4.0
2005 5/	6.8	*	*	*	*	0.5	5.2	14.9	18.4	14.2	9.4	3.9
2004 5/	6.7	*	*	*	*	0.7	5.5	14.6	17.0	15.7	9.2	4.7
2003 5/	6.6	*	*	*	*	0.8	5.6	15.1	16.4	15.3	9.4	3.5
2002	6.4	*	*	*	*	0.7	5.4	14.4	16.7	14.9	9.4	4.1
2001	6.4	*	*	*	*	0.8	5.8	14.7	16.4	14.6	9.3	4.8
2000	6.5	*	*	*	*	0.8	6.1	14.5	17.0	15.4	10.6	5.1
1999	6.5	*	*	*	*	0.7	6.0	14.7	17.1	16.3	10.3	5.1
Non-Hispanic white, female												
2006 5/	2.7	*	*	*	*	0.3	3.0	6.0	6.0	4.2	3.0	1.1
2005 5/	2.6	*	*	*	*	0.3	3.0	5.9	5.5	4.9	2.8	0.8
2004 5/	2.5	*	*	*	*	0.3	3.0	6.0	5.5	4.5	2.3	1.0
2003 5/	2.4	*	*	*	*	0.3	3.1	5.3	5.2	4.1	2.8	1.0
2002	2.4	*	*	*	*	0.4	3.2	5.1	5.2	4.9	3.0	1.0
2001	2.4	*	*	*	*	0.4	3.2	5.3	5.0	4.8	2.5	1.0
2000	2.2	*	*	*	*	0.4	3.0	4.4	5.5	4.6	2.4	1.0
1999	2.2	*	*	*	*	0.5	2.8	4.3	4.9	4.2	3.2	1.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
Non-Hispanic black, both sexes												
2006 5/	2.7	*	*	*	*	*	2.6	7.7	9.5	7.8	4.1	*
2005 5/	3.1	*	*	*	*	*	2.7	9.1	11.5	8.8	4.6	*
2004 5/	3.1	*	*	*	*	*	3.0	9.2	12.8	9.1	3.2	*
2003 5/	3.3	*	*	*	*	*	3.4	10.0	12.7	9.4	3.4	*
2002	3.2	*	*	*	*	0.5	3.0	10.2	12.3	8.7	5.1	*
2001	3.7	*	*	*	*	0.5	4.4	11.7	13.1	10.8	4.6	*
2000	3.9	*	*	*	*	0.7	5.1	12.3	13.5	10.4	5.6	*
1999	4.2	*	*	*	*	0.5	5.1	12.9	15.4	11.9	7.2	*
Non-Hispanic black, male												
2006 5/	3.8	*	*	*	*	*	3.1	11.3	15.0	13.7	7.5	*
2005 5/	4.4	*	*	*	*	*	3.2	13.7	18.6	15.4	9.0	*
2004 5/	4.6	*	*	*	*	*	4.1	13.2	20.7	17.2	*	*
2003 5/	4.9	*	*	*	*	*	4.2	16.0	20.3	16.2	6.9	*
2002	4.6	*	*	*	*	*	3.9	15.1	18.6	15.1	8.8	*
2001	5.5	*	*	*	*	*	6.1	17.4	21.7	20.0	7.8	*
2000	5.5	*	*	*	*	*	7.1	17.9	19.6	18.4	10.2	*
1999	6.1	*	*	*	*	*	6.9	19.7	24.6	21.9	12.8	*
Non-Hispanic black, female												
2006 5/	1.7	*	*	*	*	*	2.0	4.6	5.1	3.5	*	*
2005 5/	1.9	*	*	*	*	*	2.3	5.2	5.8	4.1	*	*
2004 5/	1.8	*	*	*	*	*	2.0	5.7	6.4	3.4	*	*
2003 5/	1.9	*	*	*	*	*	2.6	4.8	6.6	4.6	*	*
2002	2.1	*	*	*	*	*	2.2	6.1	7.2	4.2	*	*
2001	2.2	*	*	*	*	*	2.8	6.8	6.2	4.3	*	*
2000	2.5	*	*	*	*	*	3.4	7.5	8.6	4.8	*	*
1999	2.4	*	*	*	*	*	3.5	7.0	8.1	4.8	4.0	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
All origins 3/, both sexes												
2006	4.8	*	*	*	*	0.2	1.9	7.3	10.7	16.6	23.0	17.2
2005	4.9	*	*	*	*	0.2	2.0	7.1	11.2	17.3	23.2	17.8
2004	4.9	*	*	*	*	0.2	2.1	7.5	10.9	17.8	23.3	17.6
2003	5.2	*	*	*	*	0.3	2.4	7.8	11.4	19.7	24.3	18.2
2002	5.2	*	*	*	0.1	0.3	2.6	7.7	11.3	19.3	25.7	19.3
2001	5.2	*	*	*	*	0.3	2.6	7.8	11.2	19.9	24.5	19.9
2000	5.1	*	*	*	0.1	0.3	2.5	7.3	11.6	19.5	24.9	20.9
1999	5.1	*	*	*	*	0.4	2.5	7.0	11.7	20.1	25.5	20.4
All origins 3/, male												
2006	5.7	*	*	*	*	0.2	2.5	10.4	14.6	19.0	25.7	21.4
2005	5.8	*	*	*	*	0.2	2.7	10.3	15.3	19.9	26.2	23.3
2004	5.8	*	*	*	*	0.3	2.9	10.8	14.6	20.5	26.6	21.0
2003	6.2	*	*	*	*	0.3	3.2	11.4	15.5	23.0	26.8	23.9
2002	6.1	*	*	*	*	0.4	3.5	11.2	14.8	22.0	28.7	23.7
2001	6.1	*	*	*	*	0.4	3.4	11.4	14.7	23.6	26.6	26.5
2000	6.0	*	*	*	*	0.4	3.5	10.9	15.4	22.7	26.9	29.0
1999	6.0	*	*	*	*	0.5	3.5	10.6	15.3	23.6	28.4	27.0
All origins 3/, female												
2006	4.0	*	*	*	*	0.2	1.3	4.2	7.1	14.5	21.2	15.2
2005	4.0	*	*	*	*	0.2	1.3	4.1	7.4	15.1	21.2	15.3
2004	4.1	*	*	*	*	0.1	1.3	4.2	7.5	15.4	21.1	16.1
2003	4.3	*	*	*	*	0.2	1.5	4.3	7.6	17.0	22.6	15.8
2002	4.5	*	*	*	*	0.3	1.8	4.3	8.1	17.1	23.7	17.4
2001	4.4	*	*	*	*	0.2	1.9	4.3	8.0	16.8	23.1	17.1
2000	4.3	*	*	*	*	0.3	1.6	4.0	8.0	16.9	23.6	17.5
1999	4.3	*	*	*	*	0.3	1.4	3.6	8.3	17.3	23.5	17.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
Hispanic, both sexes												
2006	3.8	*	*	*	*	*	2.0	8.6	14.9	26.0	40.0	32.9
2005	4.1	*	*	*	*	*	2.4	9.1	18.0	28.6	42.6	30.4
2004	4.1	*	*	*	*	*	2.3	9.6	17.4	31.2	45.6	24.3
2003	4.2	*	*	*	*	0.4	2.3	9.8	17.0	33.4	47.9	26.9
2002	4.5	*	*	*	*	0.4	3.2	10.8	18.5	35.3	43.0	38.8
2001	4.3	*	*	*	*	*	3.5	10.9	17.6	33.3	43.5	30.6
2000	4.3	*	*	*	*	0.4	2.9	10.5	18.0	35.4	48.4	44.5
1999	4.2	*	*	*	*	0.5	3.6	9.7	17.2	35.7	45.0	40.0
Hispanic, male												
2006	4.1	*	*	*	*	*	3.0	12.7	20.3	24.8	33.8	33.8
2005	4.8	*	*	*	*	*	3.4	13.9	25.0	31.0	45.1	*
2004	4.5	*	*	*	*	*	3.4	14.1	22.7	30.8	42.6	*
2003	4.7	*	*	*	*	0.6	3.5	14.7	23.8	32.3	44.9	*
2002	5.1	*	*	*	*	*	4.6	17.1	23.8	37.5	46.4	*
2001	5.0	*	*	*	*	*	5.1	17.1	25.0	35.1	45.6	*
2000	4.8	*	*	*	*	0.6	4.4	16.2	23.3	36.7	46.2	50.4
1999	4.9	*	*	*	*	0.6	5.7	15.1	22.6	39.5	43.8	45.3
Hispanic, female												
2006	3.4	*	*	*	*	*	0.9	4.4	10.0	27.0	44.3	32.4
2005	3.4	*	*	*	*	*	1.2	4.2	11.5	26.7	40.9	33.9
2004	3.7	*	*	*	*	*	1.1	5.0	12.5	31.5	47.7	24.5
2003	3.7	*	*	*	*	*	1.1	5.0	10.8	34.3	50.0	26.6
2002	3.8	*	*	*	*	*	1.7	4.6	13.8	33.6	40.8	42.3
2001	3.4	*	*	*	*	*	1.8	4.7	11.0	31.8	42.0	33.5
2000	3.7	*	*	*	*	*	1.3	4.8	13.3	34.3	49.9	41.5
1999	3.5	*	*	*	*	*	1.3	4.5	12.4	32.7	45.8	37.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
Non-Hispanic 4/, both sexes												
2006	5.0	*	*	*	*	0.2	1.8	7.1	10.3	15.8	21.9	16.4
2005	5.1	*	*	*	*	0.2	1.9	6.9	10.6	16.4	22.0	17.2
2004	5.0	*	*	*	*	0.2	2.1	7.2	10.4	16.7	22.0	17.3
2003	5.4	*	*	*	*	0.2	2.4	7.5	10.9	18.7	23.0	17.9
2002	5.4	*	*	*	*	0.3	2.5	7.3	10.7	18.2	24.8	18.5
2001	5.3	*	*	*	*	0.3	2.5	7.5	10.7	18.9	23.6	19.4
2000	5.2	*	*	*	0.1	0.3	2.5	7.0	11.0	18.4	23.8	19.9
1999	5.2	*	*	*	*	0.3	2.3	6.7	11.2	19.2	24.6	19.6
Non-Hispanic 4/, male												
2006	6.0	*	*	*	*	0.2	2.4	10.1	14.1	18.6	25.1	20.8
2005	6.0	*	*	*	*	0.2	2.6	9.8	14.5	19.1	25.0	23.3
2004	6.0	*	*	*	*	0.3	2.8	10.4	13.9	19.7	25.7	20.7
2003	6.4	*	*	*	*	0.3	3.2	11.0	14.7	22.3	25.9	23.7
2002	6.2	*	*	*	*	0.4	3.3	10.6	14.0	20.9	27.7	23.4
2001	6.2	*	*	*	*	0.4	3.1	10.8	13.8	22.8	25.7	26.4
2000	6.1	*	*	*	*	0.3	3.3	10.3	14.7	21.7	26.0	28.0
1999	6.1	*	*	*	*	0.4	3.3	10.1	14.7	22.6	27.7	26.2
Non-Hispanic 4/, female												
2006	4.1	*	*	*	*	0.1	1.3	4.2	6.9	13.5	19.7	14.3
2005	4.1	*	*	*	*	0.2	1.3	4.1	7.0	14.1	20.0	14.4
2004	4.1	*	*	*	*	0.1	1.3	4.1	7.1	14.2	19.6	15.7
2003	4.4	*	*	*	*	0.2	1.6	4.2	7.3	15.7	21.1	15.3
2002	4.6	*	*	*	*	0.3	1.8	4.2	7.6	15.8	22.8	16.4
2001	4.5	*	*	*	*	0.2	1.9	4.3	7.7	15.7	22.2	16.5
2000	4.4	*	*	*	*	0.3	1.6	3.9	7.6	15.8	22.4	16.7
1999	4.4	*	*	*	*	0.3	1.4	3.5	8.0	16.3	22.6	17.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
Non-Hispanic white, both sexes												
2006 5/	5.5	*	*	*	*	0.2	1.9	7.2	10.3	16.6	22.9	16.9
2005 5/	5.5	*	*	*	*	0.2	2.1	6.7	10.5	17.3	22.7	17.7
2004 5/	5.5	*	*	*	*	0.2	2.0	7.1	10.2	17.4	23.2	17.7
2003 5/	5.9	*	*	*	*	0.2	2.4	7.4	10.8	19.6	24.3	18.6
2002	5.9	*	*	*	*	0.3	2.6	7.1	10.7	19.0	26.0	19.3
2001	5.8	*	*	*	*	0.3	2.5	7.1	10.5	19.6	24.7	20.4
2000	5.7	*	*	*	*	0.3	2.4	6.7	11.0	18.9	25.0	20.6
1999	5.6	*	*	*	*	0.3	2.3	6.4	11.0	19.8	25.6	20.2
Non-Hispanic white, male												
2006 5/	6.5	*	*	*	*	0.2	2.3	10.2	13.9	19.4	26.1	21.5
2005 5/	6.5	*	*	*	*	0.2	2.7	9.7	14.1	20.0	25.4	24.2
2004 5/	6.5	*	*	*	*	0.2	2.8	10.2	13.6	20.4	27.1	21.7
2003 5/	6.9	*	*	*	*	0.2	3.3	10.9	14.4	23.4	27.1	24.5
2002	6.7	*	*	*	*	0.3	3.4	10.3	13.7	21.7	29.3	24.7
2001	6.6	*	*	*	*	0.4	3.0	10.2	13.3	23.5	26.9	28.1
2000	6.6	*	*	*	*	0.3	3.4	9.9	14.6	22.3	27.2	29.0
1999	6.6	*	*	*	*	0.4	3.2	9.5	14.4	23.3	28.9	27.5
Non-Hispanic white, female												
2006 5/	4.6	*	*	*	*	*	1.4	4.2	6.9	14.2	20.6	14.8
2005 5/	4.6	*	*	*	*	0.2	1.4	3.8	7.1	15.0	20.8	14.8
2004 5/	4.5	*	*	*	*	*	1.2	4.1	7.0	14.9	20.5	16.0
2003 5/	4.8	*	*	*	*	0.2	1.5	4.0	7.4	16.4	22.4	16.0
2002	5.0	*	*	*	*	0.2	1.8	3.9	7.7	16.6	23.8	17.0
2001	5.0	*	*	*	*	0.3	1.9	4.1	7.9	16.3	23.2	17.2
2000	4.8	*	*	*	*	0.3	1.5	3.6	7.6	16.0	23.5	17.2
1999	4.8	*	*	*	*	0.2	1.4	3.2	7.7	16.8	23.5	17.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
Non-Hispanic black, both sexes												
2006 5/	3.4	*	*	*	*	*	2.1	8.1	12.9	11.2	12.2	10.1
2005 5/	3.6	*	*	*	*	*	1.6	9.0	13.8	11.7	14.7	10.7
2004 5/	3.6	*	*	*	*	*	2.6	9.5	13.1	12.9	11.2	11.2
2003 5/	3.8	*	*	*	*	0.4	2.5	10.1	13.4	14.7	10.6	11.0
2002	3.8	*	*	*	*	0.4	2.6	11.0	13.0	13.1	13.2	9.7
2001	3.9	*	*	*	*	*	3.2	11.2	13.2	14.5	13.6	7.8
2000	3.8	*	*	*	*	0.4	2.9	10.3	12.7	16.1	12.6	12.4
1999	3.9	*	*	*	*	0.5	3.1	10.1	14.3	15.0	14.5	12.3
Non-Hispanic black, male												
2006 5/	4.4	*	*	*	*	*	2.8	11.6	18.9	15.6	17.0	*
2005 5/	4.7	*	*	*	*	*	2.3	12.8	21.3	16.1	20.5	*
2004 5/	4.8	*	*	*	*	*	3.5	14.5	19.2	17.2	15.1	*
2003 5/	4.9	*	*	*	*	*	3.0	14.9	20.9	17.5	13.3	*
2002	4.7	*	*	*	*	*	2.9	15.5	19.6	17.3	13.2	*
2001	5.3	*	*	*	*	*	4.3	17.6	20.3	19.7	14.7	*
2000	4.8	*	*	*	*	*	3.6	15.5	18.9	19.8	14.8	*
1999	4.9	*	*	*	*	*	4.5	16.0	19.3	19.1	17.2	*
Non-Hispanic black, female												
2006 5/	2.5	*	*	*	*	*	1.4	5.1	8.1	8.1	9.5	9.0
2005 5/	2.5	*	*	*	*	*	0.9	5.7	7.7	8.6	11.4	10.0
2004 5/	2.6	*	*	*	*	*	1.8	5.3	8.1	9.8	9.0	10.8
2003 5/	2.8	*	*	*	*	*	2.0	6.0	7.4	12.6	9.1	8.3
2002	3.0	*	*	*	*	*	2.3	7.1	7.7	10.1	13.3	9.6
2001	2.7	*	*	*	*	*	2.1	5.7	7.5	10.8	13.0	*
2000	2.9	*	*	*	*	*	2.3	5.9	7.7	13.5	11.3	11.3
1999	2.9	*	*	*	*	*	1.9	5.0	10.4	12.1	13.0	12.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
All origins 3/, both sexes												
2006	1.0	*	*	*	*	0.0	0.1	0.3	0.9	2.5	7.8	21.8
2005	1.0	*	*	*	*	*	0.1	0.3	0.7	2.4	8.1	22.9
2004	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	7.8	23.8
2003	1.0	*	*	*	*	0.1	0.1	0.3	0.7	2.7	8.2	21.8
2002	1.0	*	*	*	*	0.1	0.1	0.3	0.8	2.8	7.8	22.9
2001	1.0	*	*	*	*	*	0.1	0.3	1.0	2.9	7.5	23.7
2000	1.0	*	*	*	*	*	0.1	0.3	0.8	2.4	7.5	24.8
1999	1.0	*	*	*	*	*	0.1	0.3	0.9	2.6	7.8	23.9
All origins 3/, male												
2006	0.9	*	*	*	*	*	0.1	0.3	0.9	3.1	9.6	21.7
2005	0.9	*	*	*	*	*	*	0.3	0.8	3.2	9.4	24.3
2004	0.9	*	*	*	*	*	0.1	0.4	1.0	2.9	9.3	25.5
2003	0.9	*	*	*	*	*	0.1	0.3	0.8	3.2	10.1	22.7
2002	0.9	*	*	*	*	*	0.1	0.4	1.0	3.1	9.8	24.1
2001	1.0	*	*	*	*	*	0.1	0.4	1.2	3.2	9.3	26.9
2000	0.9	*	*	*	*	*	0.1	0.3	0.9	2.9	9.4	27.6
1999	0.9	*	*	*	*	*	0.1	0.3	1.1	3.0	9.3	27.7
All origins 3/, female												
2006	1.1	*	*	*	*	*	0.1	0.3	0.8	2.0	6.6	21.9
2005	1.1	*	*	*	*	*	0.1	0.3	0.6	1.8	7.2	22.3
2004	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.8	23.0
2003	1.1	*	*	*	*	*	*	0.3	0.5	2.2	6.9	21.4
2002	1.1	*	*	*	*	*	0.1	0.2	0.7	2.5	6.4	22.4
2001	1.1	*	*	*	*	*	*	0.3	0.8	2.6	6.3	22.4
2000	1.1	*	*	*	*	*	0.1	0.3	0.6	2.1	6.2	23.7
1999	1.1	*	*	*	*	*	0.1	0.4	0.7	2.2	6.8	22.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Hispanic, both sexes												
2006	0.4	*	*	*	*	*	*	*	0.9	2.5	8.6	20.0
2005	0.5	*	*	*	*	*	*	*	1.0	3.4	7.7	29.9
2004	0.5	*	*	*	*	*	*	*	1.0	2.8	9.6	27.1
2003	0.4	*	*	*	*	*	*	*	*	3.6	9.5	18.7
2002	0.5	*	*	*	*	*	*	*	1.3	3.6	9.2	24.4
2001	0.5	*	*	*	*	*	*	*	1.2	4.5	8.5	31.3
2000	0.5	*	*	*	*	*	*	*	*	3.8	10.1	29.2
1999	0.5	*	*	*	*	*	*	*	1.2	3.3	11.9	32.3
Hispanic, male												
2006	0.3	*	*	*	*	*	*	*	*	*	9.5	*
2005	0.5	*	*	*	*	*	*	*	*	4.1	10.4	28.6
2004	0.5	*	*	*	*	*	*	*	*	*	12.2	*
2003	0.5	*	*	*	*	*	*	*	*	4.4	13.8	*
2002	0.5	*	*	*	*	*	*	*	*	4.0	9.4	*
2001	0.5	*	*	*	*	*	*	*	*	5.0	11.2	*
2000	0.5	*	*	*	*	*	*	*	*	5.3	15.7	*
1999	0.5	*	*	*	*	*	*	*	*	*	15.1	*
Hispanic, female												
2006	0.5	*	*	*	*	*	*	*	*	*	8.0	22.8
2005	0.6	*	*	*	*	*	*	*	*	2.9	5.8	30.6
2004	0.6	*	*	*	*	*	*	*	*	2.8	7.9	31.0
2003	0.4	*	*	*	*	*	*	*	*	2.9	6.6	19.5
2002	0.5	*	*	*	*	*	*	*	*	3.2	9.0	23.2
2001	0.6	*	*	*	*	*	*	*	*	4.1	6.6	30.7
2000	0.5	*	*	*	*	*	*	*	*	*	*	31.7
1999	0.6	*	*	*	*	*	*	*	*	*	9.7	32.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Non-Hispanic 4/, both sexes												
2006	1.1	*	*	*	*	*	0.1	0.3	0.9	2.5	7.8	21.9
2005	1.1	*	*	*	*	*	0.1	0.3	0.7	2.3	8.1	22.6
2004	1.1	*	*	*	*	*	0.1	0.3	0.8	2.4	7.7	23.6
2003	1.1	*	*	*	*	0.1	0.1	0.3	0.7	2.6	8.1	21.9
2002	1.1	*	*	*	*	*	0.1	0.3	0.8	2.7	7.7	22.8
2001	1.1	*	*	*	*	*	0.1	0.3	0.9	2.8	7.5	23.4
2000	1.1	*	*	*	*	*	0.1	0.3	0.8	2.3	7.4	24.6
1999	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	7.6	23.6
Non-Hispanic 4/, male												
2006	1.0	*	*	*	*	*	*	0.3	0.9	3.1	9.6	22.0
2005	1.0	*	*	*	*	*	*	0.3	0.8	3.1	9.3	24.0
2004	1.0	*	*	*	*	*	0.1	0.3	0.9	2.9	9.1	25.8
2003	1.0	*	*	*	*	*	*	0.3	0.8	3.1	9.9	23.0
2002	1.0	*	*	*	*	*	0.1	0.4	0.9	3.0	9.8	23.9
2001	1.0	*	*	*	*	*	0.1	0.4	1.1	3.1	9.2	26.6
2000	1.0	*	*	*	*	*	0.1	0.3	0.9	2.7	9.2	27.7
1999	1.0	*	*	*	*	*	0.1	0.3	1.0	3.0	9.0	27.5
Non-Hispanic 4/, female												
2006	1.2	*	*	*	*	*	*	0.3	0.8	1.9	6.5	21.9
2005	1.2	*	*	*	*	*	0.1	0.2	0.6	1.7	7.3	21.9
2004	1.2	*	*	*	*	*	0.1	0.3	0.7	2.0	6.7	22.6
2003	1.2	*	*	*	*	*	*	0.3	0.5	2.2	6.9	21.4
2002	1.2	*	*	*	*	*	0.1	0.2	0.6	2.4	6.3	22.4
2001	1.2	*	*	*	*	*	*	0.3	0.8	2.5	6.3	22.1
2000	1.2	*	*	*	*	*	0.1	0.3	0.6	2.0	6.2	23.4
1999	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.6	22.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Non-Hispanic white, both sexes												
2006 5/	1.2	*	*	*	*	*	0.1	0.3	0.8	2.2	7.7	22.4
2005 5/	1.2	*	*	*	*	*	*	0.2	0.6	2.1	8.1	22.7
2004 5/	1.2	*	*	*	*	*	0.1	0.3	0.7	2.3	7.5	24.0
2003 5/	1.2	*	*	*	*	*	0.1	0.2	0.6	2.6	8.1	22.0
2002	1.2	*	*	*	*	*	0.1	0.3	0.7	2.6	7.5	23.4
2001	1.2	*	*	*	*	*	0.1	0.3	0.8	2.6	7.3	23.6
2000	1.2	*	*	*	*	*	0.1	0.2	0.7	2.2	7.3	25.3
1999	1.2	*	*	*	*	*	0.1	0.3	0.7	2.4	7.6	24.0
Non-Hispanic white, male												
2006 5/	1.1	*	*	*	*	*	*	0.3	0.8	2.8	9.6	23.2
2005 5/	1.1	*	*	*	*	*	*	0.2	0.7	2.9	9.5	24.5
2004 5/	1.1	*	*	*	*	*	*	0.3	0.9	2.9	9.1	27.0
2003 5/	1.1	*	*	*	*	*	*	0.2	0.7	3.1	9.8	23.3
2002	1.1	*	*	*	*	*	*	0.3	0.9	3.0	9.8	24.5
2001	1.1	*	*	*	*	*	*	0.4	1.0	2.9	9.2	26.4
2000	1.0	*	*	*	*	*	*	0.2	0.8	2.6	9.3	28.9
1999	1.1	*	*	*	*	*	*	0.3	0.9	2.9	9.3	27.8
Non-Hispanic white, female												
2006 5/	1.3	*	*	*	*	*	*	0.2	0.7	1.6	6.4	22.0
2005 5/	1.3	*	*	*	*	*	*	0.2	0.5	1.4	7.1	21.8
2004 5/	1.3	*	*	*	*	*	*	0.2	0.6	1.8	6.5	22.7
2003 5/	1.3	*	*	*	*	*	*	0.2	0.5	2.1	6.9	21.4
2002	1.3	*	*	*	*	*	*	0.2	0.6	2.3	6.1	22.9
2001	1.3	*	*	*	*	*	*	0.2	0.7	2.2	6.1	22.4
2000	1.3	*	*	*	*	*	*	0.2	0.6	1.8	6.0	23.9
1999	1.3	*	*	*	*	*	*	0.3	0.6	1.9	6.5	22.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Non-Hispanic black, both sexes												
2006 5/	0.9	*	*	*	*	*	*	0.6	1.4	4.1	9.3	17.5
2005 5/	0.9	*	*	*	*	*	*	0.6	1.2	4.2	9.4	21.0
2004 5/	0.8	*	*	*	*	*	*	0.8	1.6	3.6	9.3	19.6
2003 5/	0.8	*	*	*	*	*	*	0.7	1.2	3.0	8.1	21.1
2002	0.7	*	*	*	*	*	*	0.6	1.1	3.3	9.6	15.1
2001	0.9	*	*	*	*	*	*	0.7	1.7	4.8	8.3	20.0
2000	0.8	*	*	*	*	*	*	0.7	1.2	3.8	8.3	18.1
1999	0.7	*	*	*	*	*	0.4	0.5	1.5	3.7	7.3	15.5
Non-Hispanic black, male												
2006 5/	0.6	*	*	*	*	*	*	*	1.6	3.8	9.7	*
2005 5/	0.8	*	*	*	*	*	*	*	*	5.3	10.4	20.0
2004 5/	0.7	*	*	*	*	*	*	1.0	1.6	3.7	9.1	*
2003 5/	0.7	*	*	*	*	*	*	*	*	3.3	10.7	21.3
2002	0.6	*	*	*	*	*	*	*	*	*	10.8	*
2001	0.8	*	*	*	*	*	*	*	2.2	5.0	6.3	27.7
2000	0.6	*	*	*	*	*	*	*	*	3.4	8.0	*
1999	0.6	*	*	*	*	*	*	*	*	3.8	7.2	*
Non-Hispanic black, female												
2006 5/	1.0	*	*	*	*	*	*	*	1.3	4.2	9.0	20.2
2005 5/	1.0	*	*	*	*	*	*	*	*	3.4	8.9	21.4
2004 5/	1.0	*	*	*	*	*	*	*	1.6	3.6	9.3	21.7
2003 5/	0.8	*	*	*	*	*	*	0.8	*	2.8	6.7	21.0
2002	0.8	*	*	*	*	*	*	*	*	3.8	9.0	16.3
2001	1.0	*	*	*	*	*	*	*	*	4.6	9.4	17.1
2000	1.0	*	*	*	*	*	*	*	*	4.0	8.5	19.6
1999	0.8	*	*	*	*	*	*	*	*	3.6	7.4	15.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
All origins 3/, both sexes												
2006	15.1	3.9	*	*	0.2	0.7	1.8	5.2	13.8	39.4	111.4	290.5
2005	14.8	3.9	*	0.1	0.2	0.7	1.7	4.8	13.6	39.3	110.3	288.3
2004	14.5	4.3	*	0.1	0.2	0.6	1.8	5.0	13.6	38.6	108.4	286.6
2003	14.6	4.5	*	0.1	0.2	0.7	1.8	4.9	13.6	40.1	109.5	293.1
2002	14.2	4.3	*	0.1	0.2	0.7	1.7	4.7	13.0	39.2	109.1	288.6
2001	13.9	3.3	*	*	0.2	0.6	1.7	4.6	13.0	40.2	104.2	287.7
2000	13.2	4.3	*	0.1	0.2	0.6	1.6	4.4	12.8	38.0	100.8	277.8
1999	12.7	4.4	*	0.1	0.2	0.6	1.6	4.0	12.0	37.1	97.6	268.9
All origins 3/, male												
2006	15.0	4.8	*	*	0.3	0.8	2.1	6.3	16.0	45.1	136.8	377.1
2005	14.6	4.2	*	*	0.2	0.7	1.9	5.9	16.1	43.8	135.9	376.5
2004	14.1	5.3	*	*	0.2	0.6	2.0	5.9	15.1	43.6	134.3	372.2
2003	14.3	5.2	*	*	0.3	0.7	2.1	6.0	15.4	45.3	136.8	387.6
2002	13.9	5.3	*	*	0.2	0.6	1.8	5.5	14.3	45.8	136.2	385.5
2001	13.5	3.7	*	*	0.2	0.6	1.8	5.5	14.1	46.1	130.0	393.7
2000	12.9	5.3	*	*	0.2	0.6	1.9	5.3	14.2	42.9	127.3	380.7
1999	12.4	4.6	*	*	0.2	0.6	1.8	4.7	13.4	42.3	125.3	371.1
All origins 3/, female												
2006	15.3	3.0	*	*	0.2	0.6	1.5	4.1	11.8	34.6	94.1	250.0
2005	15.0	3.5	*	*	0.2	0.7	1.5	3.7	11.4	35.5	92.9	247.8
2004	14.8	3.2	*	*	0.2	0.7	1.6	4.1	12.2	34.3	90.9	248.1
2003	14.9	3.8	*	*	0.2	0.7	1.5	3.9	12.0	35.6	91.3	251.4
2002	14.5	3.2	*	*	0.2	0.7	1.5	4.0	11.8	33.7	91.1	246.6
2001	14.2	2.9	*	*	0.2	0.6	1.6	3.9	11.9	35.2	87.1	243.3
2000	13.6	3.2	*	*	0.2	0.6	1.4	3.5	11.5	34.0	83.5	235.9
1999	13.0	4.1	*	*	0.2	0.6	1.3	3.4	10.8	33.0	79.6	227.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
Hispanic, both sexes												
2006	5.8	3.7	*	*	*	0.5	1.0	4.9	14.4	40.3	99.5	206.2
2005	5.5	3.6	*	*	*	0.4	1.1	3.9	14.8	38.3	95.8	193.4
2004	5.3	4.1	*	*	*	0.3	1.2	4.5	14.4	41.2	91.9	183.6
2003	5.4	4.0	*	*	*	0.4	1.3	4.6	15.2	39.6	90.2	233.2
2002	4.9	4.2	*	*	*	0.5	1.1	4.3	13.5	36.2	88.7	201.2
2001	5.0	*	*	*	*	0.4	1.4	3.9	12.8	41.4	94.5	222.0
2000	4.7	3.2	*	*	*	0.4	1.0	3.5	13.6	37.5	84.7	229.6
1999	4.7	2.6	*	*	*	0.3	1.3	3.6	14.0	35.6	88.7	232.2
Hispanic, male												
2006	5.6	*	*	*	*	*	1.2	6.2	15.8	48.2	113.8	241.3
2005	5.4	*	*	*	*	*	1.1	5.2	17.7	41.3	112.9	259.0
2004	5.2	5.4	*	*	*	*	1.4	5.3	17.0	44.4	109.4	231.2
2003	5.2	*	*	*	*	*	1.6	5.6	16.2	42.4	107.4	297.1
2002	4.7	4.9	*	*	*	*	1.3	4.8	15.1	40.9	112.0	235.9
2001	4.7	*	*	*	*	*	1.6	3.8	15.2	43.3	112.2	281.9
2000	4.4	*	*	*	*	*	1.4	4.4	14.3	40.9	101.3	292.2
1999	4.7	*	*	*	*	*	1.2	4.4	17.8	39.1	110.5	336.4
Hispanic, female												
2006	6.1	*	*	*	*	*	0.8	3.6	13.2	33.9	89.4	187.5
2005	5.5	*	*	*	*	*	1.0	2.5	12.1	35.9	84.0	159.0
2004	5.5	*	*	*	*	*	0.9	3.7	11.9	38.6	79.8	159.2
2003	5.7	*	*	*	*	*	1.0	3.5	14.4	37.4	78.5	200.8
2002	5.1	*	*	*	*	0.6	1.0	3.8	12.1	32.4	72.8	184.0
2001	5.4	*	*	*	*	*	1.1	4.0	10.7	39.8	82.5	192.7
2000	4.9	*	*	*	*	*	*	2.5	12.9	34.8	73.6	198.8
1999	4.7	*	*	*	*	*	1.3	2.9	10.7	32.9	74.2	182.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
Non-Hispanic 4/, both sexes												
2006	16.7	4.0	*	*	0.2	0.7	1.9	5.2	13.7	39.2	112.0	294.4
2005	16.4	3.9	*	*	0.2	0.8	1.8	4.9	13.5	39.3	111.1	292.5
2004	15.9	4.3	*	*	0.2	0.7	1.9	5.0	13.5	38.3	109.0	290.9
2003	16.0	4.6	*	*	0.2	0.8	1.9	4.9	13.5	40.0	110.2	294.9
2002	15.6	4.3	*	*	0.2	0.7	1.7	4.7	12.9	39.3	109.8	291.2
2001	15.1	3.5	*	*	0.2	0.6	1.7	4.7	12.9	40.0	104.3	289.6
2000	14.4	4.4	*	0.1	0.2	0.7	1.7	4.4	12.6	37.9	101.2	279.0
1999	13.8	4.8	*	0.1	0.2	0.7	1.6	4.1	11.8	37.1	97.7	269.2
Non-Hispanic 4/, male												
2006	16.7	5.1	*	*	0.3	0.9	2.3	6.3	16.0	44.8	138.0	384.0
2005	16.2	4.2	*	*	0.2	0.8	2.1	6.0	15.9	43.9	137.2	382.0
2004	15.6	5.2	*	*	0.2	0.6	2.2	5.9	15.0	43.4	135.3	378.6
2003	15.8	5.4	*	*	0.3	0.8	2.1	6.0	15.2	45.4	138.0	390.8
2002	15.4	5.4	*	*	0.2	0.7	1.9	5.6	14.2	45.9	136.9	391.0
2001	14.8	3.7	*	*	0.2	0.6	1.9	5.6	14.0	46.1	130.6	397.6
2000	14.1	5.4	*	*	0.2	0.7	2.0	5.4	14.1	42.9	128.2	384.0
1999	13.5	4.8	*	*	0.2	0.7	1.9	4.7	13.0	42.3	125.7	370.9
Non-Hispanic 4/, female												
2006	16.8	2.9	*	*	0.2	0.6	1.6	4.2	11.6	34.6	94.2	252.7
2005	16.5	3.5	*	*	0.2	0.7	1.5	3.8	11.3	35.4	93.4	251.6
2004	16.2	3.4	*	*	0.2	0.7	1.7	4.1	12.2	33.9	91.3	251.6
2003	16.2	3.6	*	*	0.2	0.7	1.6	3.9	11.8	35.5	91.7	252.7
2002	15.8	3.2	*	*	0.2	0.7	1.6	3.9	11.7	33.7	91.8	248.2
2001	15.4	3.3	*	*	0.2	0.6	1.6	3.8	12.0	34.8	87.1	244.6
2000	14.7	3.4	*	*	0.2	0.6	1.5	3.6	11.3	33.8	83.6	236.6
1999	14.1	4.8	*	*	0.2	0.7	1.3	3.5	10.7	32.9	79.5	228.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
Non-Hispanic white, both sexes												
2006 5/	16.5	3.4	*	*	0.2	0.5	1.2	3.5	10.3	32.1	104.2	284.8
2005 5/	16.2	2.7	*	*	0.2	0.5	1.1	3.4	9.8	32.7	103.5	284.0
2004 5/	15.8	2.6	*	*	0.2	0.4	1.3	3.3	9.8	32.2	101.8	282.1
2003 5/	15.8	3.5	*	*	0.1	0.5	1.1	3.2	10.0	33.2	102.7	285.9
2002	15.4	3.3	*	*	0.1	0.5	1.1	3.1	9.5	33.0	102.5	281.1
2001	14.9	2.4	*	*	0.2	0.4	1.1	3.1	9.3	33.5	96.6	279.7
2000	14.2	2.7	*	*	0.2	0.4	1.1	2.9	9.3	31.4	94.1	269.4
1999	13.4	3.5	*	*	0.1	0.4	1.0	2.6	8.7	31.1	89.9	255.3
Non-Hispanic white, male												
2006 5/	16.7	4.6	*	*	0.3	0.6	1.5	4.3	12.3	37.7	130.5	380.7
2005 5/	16.2	2.7	*	*	0.2	0.5	1.4	4.2	11.7	38.2	130.4	379.1
2004 5/	15.6	3.1	*	*	0.2	0.4	1.5	3.8	11.1	37.2	129.4	373.0
2003 5/	15.7	4.4	*	*	0.2	0.5	1.3	3.8	11.4	38.5	131.0	386.3
2002	15.3	4.1	*	*	*	0.5	1.2	3.6	10.3	39.7	130.8	382.6
2001	14.8	2.5	*	*	0.2	0.4	1.3	3.7	10.5	39.9	123.7	392.6
2000	14.1	3.5	*	*	0.2	0.5	1.3	3.5	10.6	37.3	121.7	378.9
1999	13.3	3.9	*	*	*	0.5	1.2	2.9	9.9	36.6	117.3	359.6
Non-Hispanic white, female												
2006 5/	16.4	2.1	*	*	*	0.4	1.0	2.7	8.4	27.3	86.0	240.1
2005 5/	16.2	2.7	*	*	0.2	0.5	0.9	2.6	8.0	27.9	85.0	240.5
2004 5/	16.0	2.1	*	*	0.2	0.5	1.1	2.7	8.6	27.8	83.0	241.4
2003 5/	15.9	2.5	*	*	*	0.5	0.9	2.6	8.7	28.7	83.5	241.7
2002	15.6	2.5	*	*	*	0.4	1.0	2.6	8.7	27.3	83.6	237.3
2001	15.0	2.3	*	*	0.2	0.4	0.9	2.4	8.3	28.0	78.6	232.7
2000	14.2	1.8	*	*	0.2	0.4	1.0	2.3	8.0	26.5	76.0	225.2
1999	13.5	3.1	*	*	*	0.4	0.8	2.2	7.5	26.4	71.9	213.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
Non-Hispanic black, both sexes												
2006 5/	22.1	6.8	*	*	0.5	2.0	6.1	16.7	42.7	104.5	213.8	449.4
2005 5/	21.5	9.1	*	*	0.5	2.3	5.7	15.2	45.1	101.6	211.9	429.8
2004 5/	21.0	10.4	*	*	0.5	2.0	5.8	17.1	45.6	96.6	205.1	437.2
2003 5/	21.4	9.5	*	*	0.6	2.3	6.4	16.8	43.7	105.6	209.9	446.5
2002	20.5	8.5	*	*	0.6	2.1	5.5	16.5	41.8	101.5	205.5	444.0
2001	20.2	8.5	*	*	0.5	1.8	5.5	16.2	42.5	104.2	204.0	432.9
2000	19.5	12.0	*	*	0.4	1.9	5.5	16.2	41.2	101.2	197.3	410.2
1999	19.2	10.2	*	*	0.5	2.0	5.6	15.6	37.8	96.6	196.4	436.8
Non-Hispanic black, male												
2006 5/	21.0	7.4	*	*	*	2.2	7.1	20.6	51.0	117.5	256.1	496.6
2005 5/	20.3	11.2	*	*	*	2.4	6.4	19.4	56.3	104.8	249.8	487.2
2004 5/	19.9	12.4	*	*	*	2.0	6.4	21.5	51.1	109.9	235.9	520.3
2003 5/	20.8	9.9	*	*	0.7	2.6	7.4	21.8	52.5	123.3	248.7	526.3
2002	19.7	10.6	*	*	*	2.0	6.2	20.6	50.1	116.3	237.1	568.9
2001	18.6	8.9	*	*	*	2.2	5.3	19.9	46.3	116.0	235.3	517.4
2000	18.3	13.6	*	*	*	2.0	6.7	20.8	45.6	105.7	234.2	496.7
1999	18.1	8.7	*	*	*	1.9	6.7	19.4	41.2	103.9	251.1	532.4
Non-Hispanic black, female												
2006 5/	23.0	*	*	*	*	1.8	5.3	13.3	36.1	95.3	189.8	430.1
2005 5/	22.5	6.8	*	*	*	2.2	5.1	11.6	36.1	99.3	190.5	406.7
2004 5/	22.0	8.4	*	*	*	1.9	5.2	13.2	41.2	87.2	187.7	404.4
2003 5/	21.9	9.0	*	*	*	2.0	5.6	12.5	36.6	93.0	187.9	415.6
2002	21.2	6.3	*	*	0.7	2.2	4.9	13.1	35.1	91.0	187.7	396.3
2001	21.7	8.1	*	*	*	1.4	5.6	13.0	39.4	95.8	186.5	401.7
2000	20.7	10.2	*	*	*	1.8	4.4	12.2	37.6	98.1	176.6	378.1
1999	20.2	11.8	*	*	*	2.0	4.7	12.3	35.1	91.5	165.8	401.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
All origins 3/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	0.3	0.6
2005	0.0	*	*	*	*	*	*	*	*	*	0.3	0.8
2004	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.8
2003	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.6
2002	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.2
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
All origins 3/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2003	0.0	*	*	*	*	*	*	*	*	*	0.4	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	1.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.7	*
All origins 3/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2005	0.0	*	*	*	*	*	*	*	*	*	0.3	0.7
2004	0.1	*	*	*	*	*	*	*	*	*	0.3	0.8
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	0.6
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	0.9
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
Non-Hispanic 4/, both sexes												
2006	0.1	*	*	*	*	*	*	*	*	*	0.3	0.7
2005	0.0	*	*	*	*	*	*	*	*	*	0.3	0.7
2004	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.8
2003	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.5
2002	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.8
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.2
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
Non-Hispanic 4/, male												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	1.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.7	*
Non-Hispanic 4/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2005	0.1	*	*	*	*	*	*	*	*	*	0.3	0.7
2004	0.1	*	*	*	*	*	*	*	*	*	0.3	0.8
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	0.6
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	0.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
Non-Hispanic white, both sexes												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	0.3	0.7
2005 5/	0.1	*	*	*	*	*	*	*	*	*	0.3	0.6
2004 5/	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.7
2003 5/	0.1	*	*	*	*	*	*	*	*	0.2	0.4	*
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	0.8
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	0.8
2000	0.1	*	*	*	*	*	*	*	*	0.1	0.4	1.2
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
Non-Hispanic white, male												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	*
Non-Hispanic white, female												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	0.3	0.8
2003 5/	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	0.8
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	1.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
Non-Hispanic black, both sexes												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)												
All origins 3/, both sexes												
2006	0.6	*	*	*	*	0.1	0.1	0.2	0.5	1.4	4.4	13.4
2005	0.3	*	*	*	*	0.1	0.1	0.1	0.3	0.7	1.9	5.4
2004	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.5	1.3	2.8
2003	0.2	*	*	*	*	*	0.0	0.1	0.2	0.6	1.3	2.8
2002	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.5	1.4	2.8
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.2	2.8
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	2.8
1999	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.6	3.2
All origins 3/, male												
2006	0.6	*	*	*	*	*	*	0.2	0.6	1.5	5.7	17.2
2005	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	2.4	7.5
2004	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.6	4.0
2003	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.6	4.1
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.8	3.8
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.7	3.7
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	4.5
1999	0.2	*	*	*	*	*	*	0.1	0.3	0.5	2.2	5.0
All origins 3/, female												
2006	0.6	*	*	*	*	*	*	0.2	0.4	1.2	3.5	11.6
2005	0.3	*	*	*	*	*	*	*	0.3	0.6	1.6	4.4
2004	0.2	*	*	*	*	*	*	*	0.2	0.4	1.1	2.2
2003	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	2.2
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	2.3
2001	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.1
1999	0.2	*	*	*	*	*	*	*	0.2	0.6	1.3	2.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)												
Hispanic, both sexes												
2006	0.2	*	*	*	*	*	*	*	*	1.5	4.2	8.6
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.2	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)												
Non-Hispanic 4/, both sexes												
2006	0.7	*	*	*	*	*	0.1	0.2	0.5	1.4	4.4	13.6
2005	0.3	*	*	*	*	*	0.1	0.1	0.3	0.7	1.9	5.5
2004	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.5	1.4	2.7
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.3	2.8
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.4	2.8
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.2	2.8
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	2.8
1999	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.7	3.2
Non-Hispanic 4/, male												
2006	0.7	*	*	*	*	*	*	0.2	0.6	1.5	5.7	17.4
2005	0.3	*	*	*	*	*	0.1	0.1	0.4	0.8	2.5	7.5
2004	0.2	*	*	*	*	*	*	*	0.3	0.5	1.7	3.8
2003	0.2	*	*	*	*	*	*	0.2	*	0.6	1.6	4.3
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.8	3.8
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.7	3.7
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	4.2
1999	0.2	*	*	*	*	*	*	0.1	0.3	0.5	2.3	5.1
Non-Hispanic 4/, female												
2006	0.7	*	*	*	*	*	*	0.2	0.4	1.2	3.5	11.9
2005	0.3	*	*	*	*	*	*	*	0.3	0.6	1.5	4.5
2004	0.2	*	*	*	*	*	*	*	0.2	0.4	1.1	2.2
2003	0.2	*	*	*	*	*	*	*	0.2	0.6	1.1	2.2
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	2.3
2001	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.2
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)												
Non-Hispanic white, both sexes												
2006 5/	0.7	*	*	*	*	*	*	0.1	0.3	1.1	4.0	12.7
2005 5/	0.3	*	*	*	*	*	*	0.1	0.3	0.6	1.8	5.2
2004 5/	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	2.7
2003 5/	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	2.7
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.6
2001	0.2	*	*	*	*	*	*	*	0.1	0.6	1.1	2.8
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.4
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.6	2.9
Non-Hispanic white, male												
2006 5/	0.7	*	*	*	*	*	*	0.1	0.4	1.3	5.2	16.2
2005 5/	0.3	*	*	*	*	*	*	*	0.3	0.7	2.4	7.3
2004 5/	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	3.6
2003 5/	0.2	*	*	*	*	*	*	0.1	*	0.5	1.4	4.0
2002	0.2	*	*	*	*	*	*	*	*	0.7	1.7	3.8
2001	0.2	*	*	*	*	*	*	*	*	0.8	1.7	4.0
2000	0.2	*	*	*	*	*	*	*	0.2	0.6	1.7	3.7
1999	0.2	*	*	*	*	*	*	*	*	0.5	2.2	4.6
Non-Hispanic white, female												
2006 5/	0.7	*	*	*	*	*	*	*	0.3	0.9	3.1	11.1
2005 5/	0.3	*	*	*	*	*	*	*	0.2	0.5	1.5	4.3
2004 5/	0.2	*	*	*	*	*	*	*	*	0.4	1.1	2.3
2003 5/	0.2	*	*	*	*	*	*	*	*	0.5	1.0	2.1
2002	0.2	*	*	*	*	*	*	*	*	0.4	1.1	2.0
2001	0.2	*	*	*	*	*	*	*	*	0.4	0.8	2.4
2000	0.2	*	*	*	*	*	*	*	*	0.5	1.1	1.9
1999	0.2	*	*	*	*	*	*	*	*	0.5	1.2	2.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)												
Non-Hispanic black, both sexes												
2006 5/	0.9	*	*	*	*	*	*	0.6	1.6	3.9	8.4	23.9
2005 5/	0.4	*	*	*	*	*	*	*	0.9	1.6	3.4	9.3
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	2.1	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.9	*	*	*	*	*	*	*	2.4	4.4	10.6	33.0
2005 5/	0.5	*	*	*	*	*	*	*	*	*	5.9	*
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	0.9	*	*	*	*	*	*	*	*	3.6	7.1	20.2
2005 5/	0.4	*	*	*	*	*	*	*	*	*	*	8.0
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
All origins 3/, both sexes												
2006	14.5	3.7	*	*	0.2	0.6	1.7	5.0	13.3	37.9	106.7	276.4
2005	14.5	3.7	*	*	0.2	0.7	1.6	4.6	13.3	38.4	108.0	282.1
2004	14.2	4.0	*	*	0.2	0.6	1.7	4.8	13.4	37.9	106.6	282.9
2003	14.4	4.2	*	0.1	0.2	0.7	1.7	4.8	13.4	39.3	107.8	289.7
2002	13.9	4.1	*	*	0.2	0.6	1.6	4.6	12.7	38.6	107.2	284.8
2001	13.6	3.1	*	*	0.2	0.5	1.6	4.5	12.7	39.4	102.6	284.0
2000	13.0	4.0	*	*	0.2	0.6	1.6	4.2	12.5	37.2	98.9	273.7
1999	12.4	4.3	*	0.1	0.2	0.6	1.5	3.9	11.7	36.4	95.3	264.8
All origins 3/, male												
2006	14.3	4.4	*	*	0.2	0.7	2.0	6.0	15.4	43.4	130.7	358.8
2005	14.2	4.1	*	*	0.2	0.7	1.8	5.7	15.6	42.8	133.1	368.0
2004	13.8	5.1	*	*	0.2	0.5	2.0	5.7	14.8	42.9	132.1	367.4
2003	14.1	4.8	*	*	0.2	0.7	2.0	5.7	15.2	44.5	134.8	383.1
2002	13.7	5.1	*	*	0.2	0.6	1.8	5.4	14.0	45.0	134.0	380.7
2001	13.2	3.5	*	*	0.2	0.5	1.8	5.3	13.8	45.2	128.0	388.7
2000	12.6	5.1	*	*	0.2	0.6	1.8	5.1	13.8	42.1	125.1	374.2
1999	12.1	4.5	*	*	0.1	0.5	1.7	4.5	13.1	41.5	122.3	364.6
All origins 3/, female												
2006	14.6	3.0	*	*	0.2	0.6	1.4	3.9	11.4	33.2	90.2	237.8
2005	14.7	3.2	*	*	0.2	0.6	1.4	3.6	11.0	34.8	91.0	242.6
2004	14.6	3.0	*	*	0.2	0.6	1.5	3.9	12.0	33.7	89.5	244.9
2003	14.6	3.7	*	*	0.2	0.6	1.5	3.8	11.8	34.9	89.7	248.4
2002	14.2	3.0	*	*	0.2	0.7	1.4	3.8	11.6	33.2	89.5	243.2
2001	14.0	2.7	*	*	0.2	0.5	1.5	3.7	11.8	34.5	85.9	240.1
2000	13.3	2.8	*	*	0.2	0.5	1.3	3.4	11.2	33.3	81.8	232.8
1999	12.7	4.1	*	*	0.2	0.6	1.3	3.3	10.4	32.2	77.8	224.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
Hispanic, both sexes												
2006	5.6	3.7	*	*	*	0.4	0.9	4.7	13.8	38.7	95.0	197.2
2005	5.3	3.6	*	*	*	0.3	1.0	3.7	14.5	37.5	93.5	187.4
2004	5.2	3.9	*	*	*	0.3	1.2	4.4	14.0	40.5	91.0	179.3
2003	5.3	3.8	*	*	*	0.4	1.3	4.4	14.8	39.0	88.5	230.6
2002	4.8	4.2	*	*	*	0.4	1.1	4.2	13.2	35.7	86.9	198.4
2001	4.9	*	*	*	*	0.3	1.2	3.7	12.5	40.5	93.2	217.6
2000	4.5	2.9	*	*	*	0.4	1.0	3.3	13.3	36.9	83.0	225.6
1999	4.5	*	*	*	*	*	1.2	3.5	13.6	34.4	86.9	228.0
Hispanic, male												
2006	5.4	*	*	*	*	*	1.1	6.0	15.1	46.6	108.8	227.8
2005	5.2	*	*	*	*	*	1.0	5.0	17.2	40.3	111.6	249.1
2004	5.0	5.2	*	*	*	*	1.4	5.2	16.4	43.5	108.4	224.2
2003	5.1	*	*	*	*	*	1.5	5.3	15.8	41.9	105.1	295.5
2002	4.6	4.9	*	*	*	*	1.3	4.8	14.4	40.7	109.2	232.6
2001	4.6	*	*	*	*	*	1.4	3.7	14.6	42.5	110.4	278.1
2000	4.3	*	*	*	*	*	1.3	4.2	13.8	40.5	99.3	280.1
1999	4.5	*	*	*	*	*	1.1	4.3	17.2	38.2	109.0	330.0
Hispanic, female												
2006	5.8	*	*	*	*	*	0.7	3.5	12.6	32.4	85.4	180.9
2005	5.3	*	*	*	*	*	0.9	2.5	11.9	35.2	80.9	155.1
2004	5.4	*	*	*	*	*	0.9	3.6	11.9	38.0	79.0	156.3
2003	5.6	*	*	*	*	*	0.9	3.5	13.9	36.6	77.2	197.7
2002	5.0	*	*	*	*	0.6	0.8	3.6	12.0	31.8	71.7	181.5
2001	5.2	*	*	*	*	*	1.0	3.7	10.5	38.9	81.6	188.0
2000	4.8	*	*	*	*	*	*	2.4	12.8	34.0	72.0	198.8
1999	4.5	*	*	*	*	*	1.3	2.7	10.4	31.3	72.2	178.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
Non-Hispanic 4/, both sexes												
2006	16.0	3.7	*	*	0.2	0.7	1.8	5.0	13.2	37.8	107.2	280.0
2005	16.0	3.7	*	*	0.2	0.7	1.7	4.7	13.1	38.5	108.8	286.2
2004	15.7	4.1	*	*	0.2	0.6	1.8	4.8	13.3	37.6	107.2	287.2
2003	15.7	4.3	*	*	0.2	0.7	1.8	4.8	13.2	39.2	108.5	291.4
2002	15.3	4.1	*	*	0.2	0.7	1.7	4.6	12.6	38.6	107.9	287.4
2001	14.9	3.3	*	*	0.2	0.6	1.7	4.6	12.7	39.2	102.7	285.9
2000	14.1	4.2	*	*	0.2	0.6	1.6	4.3	12.3	37.1	99.3	274.9
1999	13.5	4.7	*	*	0.2	0.6	1.5	3.9	11.5	36.4	95.4	265.0
Non-Hispanic 4/, male												
2006	15.9	4.6	*	*	0.3	0.8	2.2	6.0	15.4	43.1	131.8	365.5
2005	15.8	4.1	*	*	0.2	0.8	1.9	5.8	15.5	42.9	134.3	373.6
2004	15.3	5.1	*	*	0.2	0.6	2.1	5.8	14.6	42.7	133.1	374.0
2003	15.5	5.0	*	*	0.2	0.8	2.1	5.8	15.0	44.6	136.0	386.2
2002	15.1	5.1	*	*	0.2	0.6	1.8	5.4	13.9	45.1	134.7	386.2
2001	14.6	3.5	*	*	0.2	0.6	1.8	5.4	13.7	45.3	128.5	392.7
2000	13.8	5.2	*	*	0.2	0.6	1.9	5.1	13.7	42.0	125.9	377.7
1999	13.2	4.6	*	*	0.1	0.6	1.8	4.5	12.7	41.5	122.6	364.4
Non-Hispanic 4/, female												
2006	16.1	2.8	*	*	0.2	0.6	1.5	4.0	11.2	33.2	90.4	240.2
2005	16.2	3.2	*	*	0.2	0.7	1.5	3.7	11.0	34.7	91.5	246.4
2004	16.0	3.1	*	*	0.2	0.6	1.6	4.0	12.0	33.3	89.8	248.5
2003	15.9	3.5	*	*	0.2	0.7	1.5	3.8	11.6	34.7	90.1	249.7
2002	15.5	2.9	*	*	0.2	0.7	1.5	3.8	11.4	33.2	90.2	244.8
2001	15.2	3.1	*	*	0.2	0.5	1.6	3.7	11.8	34.1	85.8	241.5
2000	14.4	3.1	*	*	0.2	0.5	1.4	3.5	11.0	33.2	82.0	233.4
1999	13.8	4.8	*	*	0.2	0.6	1.3	3.3	10.4	32.1	77.8	225.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
Non-Hispanic white, both sexes												
2006 5/	15.8	3.1	*	*	0.2	0.5	1.2	3.4	9.9	30.9	99.8	271.3
2005 5/	15.8	2.5	*	*	0.2	0.5	1.1	3.3	9.5	32.0	101.3	278.1
2004 5/	15.5	2.5	*	*	0.1	0.4	1.3	3.2	9.6	31.5	100.0	278.6
2003 5/	15.5	3.2	*	*	0.1	0.5	1.1	3.1	9.8	32.5	101.1	282.6
2002	15.2	3.1	*	*	0.1	0.4	1.1	3.0	9.2	32.3	100.8	277.6
2001	14.6	2.2	*	*	0.2	0.4	1.1	3.0	9.2	32.7	95.1	276.1
2000	13.9	2.5	*	*	0.2	0.4	1.1	2.8	9.0	30.7	92.3	265.7
1999	13.1	3.5	*	*	0.1	0.4	0.9	2.4	8.4	30.4	87.8	251.6
Non-Hispanic white, male												
2006 5/	15.9	4.1	*	*	0.2	0.6	1.4	4.2	11.9	36.4	124.8	363.4
2005 5/	15.8	2.6	*	*	0.2	0.5	1.3	4.1	11.3	37.3	127.6	370.9
2004 5/	15.3	3.0	*	*	0.2	0.3	1.4	3.7	10.9	36.5	127.2	368.8
2003 5/	15.4	4.0	*	*	0.2	0.5	1.3	3.7	11.2	37.8	129.3	381.9
2002	15.0	3.9	*	*	*	0.4	1.2	3.5	10.0	38.8	128.8	377.8
2001	14.5	2.3	*	*	*	0.3	1.3	3.6	10.3	39.1	121.7	387.6
2000	13.7	3.3	*	*	0.2	0.4	1.2	3.3	10.3	36.5	119.4	373.4
1999	13.0	3.8	*	*	*	0.4	1.1	2.8	9.6	35.9	114.4	353.8
Non-Hispanic white, female												
2006 5/	15.7	2.1	*	*	*	0.4	0.9	2.6	8.1	26.2	82.5	228.4
2005 5/	15.9	2.3	*	*	*	0.5	0.9	2.5	7.7	27.4	83.2	235.7
2004 5/	15.7	2.0	*	*	*	0.4	1.1	2.6	8.4	27.2	81.5	238.2
2003 5/	15.6	2.3	*	*	*	0.5	0.9	2.5	8.5	28.0	82.0	238.9
2002	15.3	2.2	*	*	*	0.4	1.0	2.5	8.5	26.7	82.0	234.3
2001	14.7	2.1	*	*	0.2	0.4	0.9	2.4	8.1	27.3	77.5	229.7
2000	14.0	*	*	*	*	0.3	0.9	2.3	7.7	25.8	74.4	222.2
1999	13.2	3.1	*	*	*	0.4	0.8	2.1	7.3	25.8	70.3	210.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
Non-Hispanic black, both sexes												
2006 5/	21.1	6.6	*	*	0.4	1.8	5.9	15.9	41.1	100.4	205.1	425.2
2005 5/	21.0	8.7	*	*	0.4	2.1	5.5	14.8	44.2	99.8	208.0	418.9
2004 5/	20.7	10.1	*	*	0.5	1.8	5.5	16.7	45.0	95.4	202.5	432.6
2003 5/	21.1	9.3	*	*	0.5	2.2	6.3	16.5	43.3	104.3	207.1	442.3
2002	20.2	8.0	*	*	0.6	2.0	5.4	16.0	41.4	100.9	202.7	436.5
2001	20.0	8.5	*	*	0.4	1.7	5.3	15.8	42.1	102.9	201.9	428.9
2000	19.2	11.4	*	*	0.4	1.8	5.3	15.6	40.7	99.9	194.6	404.2
1999	18.8	10.2	*	*	0.5	1.9	5.4	15.2	37.0	95.3	193.0	428.7
Non-Hispanic black, male												
2006 5/	20.0	7.1	*	*	*	1.9	6.9	19.7	48.7	112.9	245.0	462.7
2005 5/	19.8	11.2	*	*	*	2.1	6.0	18.9	55.1	103.4	243.9	473.9
2004 5/	19.6	12.1	*	*	*	2.0	6.2	21.2	50.2	108.6	233.0	514.2
2003 5/	20.5	9.6	*	*	*	2.5	7.3	21.3	52.1	121.6	244.4	521.0
2002	19.4	10.0	*	*	*	1.8	6.1	20.1	49.8	115.4	233.3	562.3
2001	18.4	8.9	*	*	*	2.1	5.1	19.3	45.7	114.3	233.5	514.0
2000	17.9	13.6	*	*	*	1.9	6.5	19.9	45.0	104.2	231.7	487.3
1999	17.7	8.7	*	*	*	1.8	6.4	19.0	40.5	102.5	246.7	519.4
Non-Hispanic black, female												
2006 5/	22.1	*	*	*	*	1.7	5.0	12.7	35.1	91.4	182.5	409.9
2005 5/	22.1	*	*	*	*	2.1	5.0	11.3	35.4	97.2	187.6	396.8
2004 5/	21.7	8.0	*	*	*	1.6	4.9	12.9	40.9	86.0	185.3	400.4
2003 5/	21.7	9.0	*	*	*	1.9	5.5	12.4	36.3	92.0	186.1	411.9
2002	20.9	*	*	*	*	2.1	4.8	12.5	34.7	90.6	185.4	388.4
2001	21.4	8.1	*	*	*	1.4	5.5	12.8	39.2	94.9	184.1	397.4
2000	20.3	9.2	*	*	*	1.7	4.1	11.8	37.3	96.9	173.7	373.3
1999	19.8	11.8	*	*	*	2.0	4.5	12.0	34.3	90.2	163.0	394.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
All origins 3/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
Non-Hispanic 4/, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
Non-Hispanic white, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)												
All origins 3/, both sexes												
2006	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	4.7
2005	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	1.8	5.2
2004	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	5.2
2003	0.3	*	*	*	*	*	0.1	0.2	0.2	0.6	1.9	5.9
2002	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.3
2001	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	2.0	6.0
2000	0.3	*	*	*	*	*	0.0	0.1	0.2	0.6	2.0	7.5
1999	0.3	*	*	*	*	*	0.1	0.2	0.2	0.7	2.0	7.5
All origins 3/, male												
2006	0.1	*	*	*	*	*	*	*	0.1	0.5	0.8	3.7
2005	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	3.6
2004	0.2	*	*	*	*	*	*	*	0.2	0.4	1.5	4.5
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.4	5.4
2002	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	5.3
2001	0.2	*	*	*	*	*	*	*	*	0.5	1.5	4.3
2000	0.2	*	*	*	*	*	*	*	*	0.3	1.5	5.6
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.7	6.8
All origins 3/, female												
2006	0.3	*	*	*	*	*	*	0.2	0.2	0.5	1.6	5.2
2005	0.4	*	*	*	*	*	*	0.2	0.2	0.6	2.0	5.9
2004	0.4	*	*	*	*	*	0.1	0.2	0.2	0.7	2.1	5.5
2003	0.4	*	*	*	*	*	0.1	0.2	0.3	0.7	2.2	6.1
2002	0.4	*	*	*	*	*	*	0.1	0.3	0.6	2.2	6.7
2001	0.4	*	*	*	*	*	0.1	0.2	0.3	0.5	2.3	6.7
2000	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.3	8.3
1999	0.4	*	*	*	*	*	0.1	0.2	0.3	0.9	2.1	7.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)												
Hispanic, both sexes												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.2	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)												
Non-Hispanic 4/, both sexes												
2006	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	4.8
2005	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	1.8	5.2
2004	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	1.8	5.3
2003	0.3	*	*	*	*	*	0.1	0.2	0.2	0.5	1.9	6.0
2002	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	6.4
2001	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	2.0	6.1
2000	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.0	7.6
1999	0.3	*	*	*	*	*	0.0	0.1	0.2	0.7	1.9	7.5
Non-Hispanic 4/, male												
2006	0.2	*	*	*	*	*	*	*	*	0.4	0.8	3.9
2005	0.2	*	*	*	*	*	*	*	*	0.4	1.5	3.5
2004	0.2	*	*	*	*	*	*	*	0.2	0.4	1.4	4.6
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.4	5.4
2002	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	5.3
2001	0.2	*	*	*	*	*	*	*	*	0.5	1.6	4.3
2000	0.2	*	*	*	*	*	*	*	*	0.3	1.5	5.8
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	6.9
Non-Hispanic 4/, female												
2006	0.3	*	*	*	*	*	*	0.2	0.2	0.5	1.6	5.2
2005	0.4	*	*	*	*	*	*	0.2	0.2	0.6	2.0	5.9
2004	0.4	*	*	*	*	*	0.1	0.2	0.2	0.7	2.1	5.6
2003	0.4	*	*	*	*	*	0.1	0.2	0.3	0.7	2.2	6.2
2002	0.4	*	*	*	*	*	*	0.1	0.3	0.6	2.2	6.8
2001	0.4	*	*	*	*	*	0.1	0.2	0.3	0.4	2.3	6.9
2000	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.3	8.3
1999	0.4	*	*	*	*	*	*	0.2	0.2	0.9	2.1	7.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)												
Non-Hispanic white, both sexes												
2006 5/	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.3	5.0
2005 5/	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.8	5.2
2004 5/	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	5.3
2003 5/	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	6.0
2002	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.5
2001	0.3	*	*	*	*	*	0.1	0.1	0.2	0.4	2.0	6.2
2000	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.9	7.6
1999	0.3	*	*	*	*	*	*	0.1	0.2	0.7	1.9	7.7
Non-Hispanic white, male												
2006 5/	0.2	*	*	*	*	*	*	*	*	0.5	0.8	4.0
2005 5/	0.2	*	*	*	*	*	*	*	*	0.4	1.6	3.4
2004 5/	0.2	*	*	*	*	*	*	*	0.2	0.4	1.4	4.8
2003 5/	0.2	*	*	*	*	*	*	*	*	0.4	1.3	5.8
2002	0.2	*	*	*	*	*	*	*	*	0.5	1.3	5.4
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.5	4.2
2000	0.2	*	*	*	*	*	*	*	*	*	1.5	6.0
1999	0.2	*	*	*	*	*	*	*	*	0.4	1.7	7.3
Non-Hispanic white, female												
2006 5/	0.4	*	*	*	*	*	*	0.2	0.2	0.5	1.6	5.5
2005 5/	0.4	*	*	*	*	*	*	0.1	0.2	0.6	2.0	6.0
2004 5/	0.4	*	*	*	*	*	*	0.1	0.2	0.7	2.0	5.6
2003 5/	0.4	*	*	*	*	*	*	0.2	0.2	0.7	2.2	6.1
2002	0.4	*	*	*	*	*	*	*	0.3	0.7	2.2	6.9
2001	0.4	*	*	*	*	*	*	0.2	0.3	0.5	2.3	7.1
2000	0.5	*	*	*	*	*	*	*	0.3	0.7	2.1	8.2
1999	0.5	*	*	*	*	*	*	0.2	0.2	0.9	2.1	7.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)												
Non-Hispanic black, both sexes												
2006 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.2	*	*	*	*	*	*	*	*	*	2.6	*
2003 5/	0.3	*	*	*	*	*	*	*	*	*	2.5	5.9
2002	0.2	*	*	*	*	*	*	*	*	*	*	6.4
2001	0.2	*	*	*	*	*	*	*	*	*	2.4	*
2000	0.3	*	*	*	*	*	*	*	*	*	3.1	7.9
1999	0.2	*	*	*	*	*	*	*	*	1.2	*	*
Non-Hispanic black, male												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.3	*	*	*	*	*	*	*	*	*	3.2	*
2003 5/	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	3.6	9.2
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
All origins 3/, both sexes												
2006	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.5
2005	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.7
2004	0.2	*	*	*	*	*	*	*	*	0.2	1.4	5.5
2003	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.3
2002	0.2	*	*	*	*	*	*	*	*	0.2	1.3	4.8
2001	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.5
2000	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.0
All origins 3/, male												
2006	0.3	*	*	*	*	*	*	*	*	0.5	3.2	17.2
2005	0.4	*	*	*	*	*	*	*	*	0.6	3.2	18.2
2004	0.4	*	*	*	*	*	*	*	*	0.5	3.4	17.8
2003	0.3	*	*	*	*	*	*	*	*	0.6	3.2	17.2
2002	0.3	*	*	*	*	*	*	*	*	0.5	3.2	15.8
2001	0.3	*	*	*	*	*	*	*	*	0.5	2.8	18.6
2000	0.3	*	*	*	*	*	*	*	*	0.5	3.3	18.1
1999	0.3	*	*	*	*	*	*	*	*	0.6	3.3	17.5
All origins 3/, female												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
Hispanic, both sexes												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
Non-Hispanic 4/, both sexes												
2006	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.5
2005	0.2	*	*	*	*	*	*	*	*	0.3	1.4	5.8
2004	0.2	*	*	*	*	*	*	*	*	0.2	1.4	5.5
2003	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.3
2002	0.2	*	*	*	*	*	*	*	*	0.2	1.3	4.9
2001	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.7
2000	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.1
Non-Hispanic 4/, male												
2006	0.4	*	*	*	*	*	*	*	*	0.5	3.1	17.4
2005	0.4	*	*	*	*	*	*	*	*	0.6	3.4	18.5
2004	0.4	*	*	*	*	*	*	*	*	0.5	3.5	18.0
2003	0.4	*	*	*	*	*	*	*	*	0.5	3.1	17.5
2002	0.3	*	*	*	*	*	*	*	*	0.5	3.2	16.2
2001	0.4	*	*	*	*	*	*	*	*	0.5	2.8	19.3
2000	0.3	*	*	*	*	*	*	*	*	0.4	3.3	18.0
1999	0.3	*	*	*	*	*	*	*	*	0.5	3.4	18.0
Non-Hispanic 4/, female												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
Non-Hispanic white, both sexes												
2006 5/	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.7
2005 5/	0.2	*	*	*	*	*	*	*	*	0.3	1.4	6.0
2004 5/	0.2	*	*	*	*	*	*	*	*	0.2	1.4	5.7
2003 5/	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.6
2002	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.0
2001	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.9
2000	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999	0.2	*	*	*	*	*	*	*	*	0.2	1.4	5.2
Non-Hispanic white, male												
2006 5/	0.4	*	*	*	*	*	*	*	*	0.5	3.2	17.8
2005 5/	0.5	*	*	*	*	*	*	*	*	0.6	3.5	19.1
2004 5/	0.4	*	*	*	*	*	*	*	*	0.5	3.4	18.5
2003 5/	0.4	*	*	*	*	*	*	*	*	0.5	3.1	18.5
2002	0.4	*	*	*	*	*	*	*	*	0.4	3.2	16.6
2001	0.4	*	*	*	*	*	*	*	*	0.5	2.8	20.1
2000	0.4	*	*	*	*	*	*	*	*	0.4	3.4	18.0
1999	0.4	*	*	*	*	*	*	*	*	0.5	3.4	18.4
Non-Hispanic white, female												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
Non-Hispanic black, both sexes												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
All origins 3/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	0.3	0.5
2005	0.0	*	*	*	*	*	*	*	*	0.1	0.3	0.6
2004	0.0	*	*	*	*	*	*	*	*	*	0.2	0.7
2003	0.0	*	*	*	*	*	*	*	*	*	0.2	0.7
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	0.9
2000	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	*
All origins 3/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2006	0.1	*	*	*	*	*	*	*	*	*	0.5	0.7
2005	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.8
2004	0.1	*	*	*	*	*	*	*	*	*	0.3	1.0
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.1
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
Non-Hispanic 4/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	0.3	0.5
2005	0.0	*	*	*	*	*	*	*	*	0.1	0.3	0.6
2004	0.0	*	*	*	*	*	*	*	*	*	*	0.7
2003	0.0	*	*	*	*	*	*	*	*	*	0.2	0.7
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	0.9
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	*
Non-Hispanic 4/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, female												
2006	0.1	*	*	*	*	*	*	*	*	*	0.5	0.8
2005	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.9
2004	0.1	*	*	*	*	*	*	*	*	*	*	1.0
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	1.0
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
Non-Hispanic white, both sexes												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	0.3	0.5
2005 5/	0.0	*	*	*	*	*	*	*	*	*	0.3	0.6
2004 5/	0.0	*	*	*	*	*	*	*	*	*	*	0.7
2003 5/	0.0	*	*	*	*	*	*	*	*	*	0.2	0.7
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	0.9
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Non-Hispanic white, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	0.5	0.8
2005 5/	0.1	*	*	*	*	*	*	*	*	*	0.5	0.8
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	1.0
2003 5/	0.1	*	*	*	*	*	*	*	*	*	0.4	1.0
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
All origins 3/, both sexes												
2006	0.3	...	...	*	0.4	0.8	0.4	0.1	*	*	*	*
2005	0.3	...	...	*	0.4	0.8	0.5	0.1	*	*	*	*
2004	0.2	...	...	*	0.4	0.7	0.4	0.2	*	*	*	*
2003	0.2	...	...	*	0.3	0.6	0.4	0.1	*	*	*	*
2002	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001	0.1	...	...	*	0.3	0.5	0.2	*	*	*	*	*
2000	0.1	...	...	*	0.2	0.5	0.3	*	*	*	*	*
1999	0.1	...	...	*	0.3	0.5	0.3	*	*	*	*	*
All origins 3/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2006	0.5	...	...	*	0.9	1.6	0.9	0.3	*	*	*	*
2005	0.5	...	...	*	0.9	1.6	0.9	0.3	*	*	*	*
2004	0.5	...	...	*	0.8	1.5	0.7	0.4	*	*	*	*
2003	0.4	...	...	*	0.6	1.2	0.7	0.1	*	*	*	*
2002	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
2001	0.3	...	...	*	0.6	1.0	0.5	*	*	*	*	*
2000	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
1999	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
Hispanic, both sexes												
2006	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
2005	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*
2004	0.3	...	...	*	0.4	0.6	0.4	*	*	*	*	*
2003	0.3	...	...	*	0.3	0.6	0.5	*	*	*	*	*
2002	0.2	...	...	*	*	0.4	0.4	*	*	*	*	*
2001	0.2	...	...	*	0.4	0.6	*	*	*	*	*	*
2000	0.2	...	...	*	0.4	0.6	0.4	*	*	*	*	*
1999	0.2	...	...	*	*	0.4	0.5	*	*	*	*	*
Hispanic, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2006	0.7	...	...	*	1.1	1.9	1.1	*	*	*	*	*
2005	0.6	...	...	*	1.0	1.7	1.1	*	*	*	*	*
2004	0.6	...	...	*	0.9	1.4	0.9	*	*	*	*	*
2003	0.5	...	...	*	0.7	1.4	1.0	*	*	*	*	*
2002	0.4	...	...	*	*	0.8	0.8	*	*	*	*	*
2001	0.5	...	...	*	0.8	1.2	*	*	*	*	*	*
2000	0.5	...	...	*	0.8	1.2	0.9	*	*	*	*	*
1999	0.4	...	...	*	*	0.9	1.0	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
Non-Hispanic 4/, both sexes												
2006	0.2	...	...	*	0.4	0.8	0.4	0.1	*	*	*	*
2005	0.2	...	...	*	0.4	0.8	0.5	0.1	*	*	*	*
2004	0.2	...	...	*	0.4	0.7	0.3	0.2	*	*	*	*
2003	0.2	...	...	*	0.3	0.6	0.4	0.1	*	*	*	*
2002	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001	0.1	...	...	*	0.3	0.5	0.2	*	*	*	*	*
2000	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
1999	0.1	...	...	*	0.2	0.5	0.2	*	*	*	*	*
Non-Hispanic 4/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, female												
2006	0.5	...	...	*	0.8	1.6	0.8	0.3	*	*	*	*
2005	0.5	...	...	*	0.9	1.6	0.9	0.3	*	*	*	*
2004	0.4	...	...	*	0.8	1.5	0.7	0.4	*	*	*	*
2003	0.3	...	...	*	0.5	1.1	0.7	0.1	*	*	*	*
2002	0.2	...	...	*	0.4	0.9	0.4	*	*	*	*	*
2001	0.3	...	...	*	0.5	0.9	0.4	*	*	*	*	*
2000	0.3	...	...	*	0.4	0.9	0.5	*	*	*	*	*
1999	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
Non-Hispanic white, both sexes												
2006 5/	0.1	...	...	*	0.3	0.5	0.3	0.1	*	*	*	*
2005 5/	0.2	...	...	*	0.3	0.5	0.3	0.1	*	*	*	*
2004 5/	0.2	...	...	*	0.2	0.5	0.2	0.1	*	*	*	*
2003 5/	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2002	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
2001	0.1	...	...	*	0.2	0.3	0.1	*	*	*	*	*
2000	0.1	...	...	*	0.1	0.3	0.2	*	*	*	*	*
1999	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2006 5/	0.3	...	...	*	0.6	0.9	0.5	0.2	*	*	*	*
2005 5/	0.3	...	...	*	0.6	1.0	0.6	0.2	*	*	*	*
2004 5/	0.3	...	...	*	0.5	1.0	0.5	0.3	*	*	*	*
2003 5/	0.2	...	...	*	0.3	0.7	0.4	*	*	*	*	*
2002	0.1	...	...	*	0.2	0.6	0.3	*	*	*	*	*
2001	0.2	...	...	*	0.3	0.6	0.2	*	*	*	*	*
2000	0.2	...	...	*	0.3	0.6	0.3	*	*	*	*	*
1999	0.2	...	...	*	0.2	0.6	0.3	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
Non-Hispanic black, both sexes												
2006 5/	0.7	...	...	*	0.9	2.2	1.1	0.4	*	*	*	*
2005 5/	0.7	...	...	*	1.1	2.2	1.2	*	*	*	*	*
2004 5/	0.7	...	...	*	1.2	2.0	0.8	0.5	*	*	*	*
2003 5/	0.5	...	...	*	0.7	1.6	1.1	*	*	*	*	*
2002	0.4	...	...	*	0.8	1.2	0.7	*	*	*	*	*
2001	0.4	...	...	*	0.8	1.3	0.7	*	*	*	*	*
2000	0.4	...	...	*	0.6	1.3	0.7	*	*	*	*	*
1999	0.5	...	...	*	0.8	1.3	0.7	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2006 5/	1.3	...	...	*	1.8	4.1	2.1	0.8	*	*	*	*
2005 5/	1.4	...	...	*	2.2	4.2	2.2	*	*	*	*	*
2004 5/	1.3	...	...	*	2.4	3.9	1.5	0.9	*	*	*	*
2003 5/	1.0	...	...	*	1.4	3.0	2.0	*	*	*	*	*
2002	0.8	...	...	*	1.5	2.4	1.4	*	*	*	*	*
2001	0.8	...	...	*	1.6	2.4	1.4	*	*	*	*	*
2000	0.8	...	...	*	1.2	2.4	1.3	*	*	*	*	*
1999	0.9	...	...	*	1.6	2.5	1.4	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
All origins 3/, both sexes												
2006	0.0	...	...	*	*	*	*	*	*	*	*	*
2005	0.0	...	...	*	*	*	*	*	*	*	*	*
2004	0.0	...	...	*	*	*	*	*	*	*	*	*
2003	0.0	...	...	*	*	*	*	*	*	*	*	*
2002	0.0	...	...	*	*	*	*	*	*	*	*	*
2001	0.0	...	...	*	*	*	*	*	*	*	*	*
2000	0.0	...	...	*	*	*	*	*	*	*	*	*
1999	0.0	...	...	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2006	0.0	...	...	*	*	*	*	*	*	*	*	*
2005	0.0	...	...	*	*	*	*	*	*	*	*	*
2004	0.0	...	...	*	*	*	*	*	*	*	*	*
2003	0.0	...	...	*	*	*	*	*	*	*	*	*
2002	0.0	...	...	*	*	*	*	*	*	*	*	*
2001	0.0	...	...	*	*	*	*	*	*	*	*	*
2000	0.0	...	...	*	*	*	*	*	*	*	*	*
1999	0.0	...	...	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
Hispanic, both sexes												
2006	*	...	...	*	*	*	*	*	*	*	*	*
2005	*	...	...	*	*	*	*	*	*	*	*	*
2004	*	...	...	*	*	*	*	*	*	*	*	*
2003	*	...	...	*	*	*	*	*	*	*	*	*
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2006	*	...	...	*	*	*	*	*	*	*	*	*
2005	*	...	...	*	*	*	*	*	*	*	*	*
2004	*	...	...	*	*	*	*	*	*	*	*	*
2003	*	...	...	*	*	*	*	*	*	*	*	*
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
Non-Hispanic 4/, both sexes												
2006	0.0	...	...	*	*	*	*	*	*	*	*	*
2005	0.0	...	...	*	*	*	*	*	*	*	*	*
2004	0.0	...	...	*	*	*	*	*	*	*	*	*
2003	0.0	...	...	*	*	*	*	*	*	*	*	*
2002	0.0	...	...	*	*	*	*	*	*	*	*	*
2001	0.0	...	...	*	*	*	*	*	*	*	*	*
2000	0.0	...	...	*	*	*	*	*	*	*	*	*
1999	0.0	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, female												
2006	0.0	...	...	*	*	*	*	*	*	*	*	*
2005	0.0	...	...	*	*	*	*	*	*	*	*	*
2004	0.0	...	...	*	*	*	*	*	*	*	*	*
2003	0.0	...	...	*	*	*	*	*	*	*	*	*
2002	0.0	...	...	*	*	*	*	*	*	*	*	*
2001	0.0	...	...	*	*	*	*	*	*	*	*	*
2000	0.0	...	...	*	*	*	*	*	*	*	*	*
1999	0.0	...	...	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
Non-Hispanic white, both sexes												
2006 5/	*	...	...	*	*	*	*	*	*	*	*	*
2005 5/	*	...	...	*	*	*	*	*	*	*	*	*
2004 5/	*	...	...	*	*	*	*	*	*	*	*	*
2003 5/	*	...	...	*	*	*	*	*	*	*	*	*
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2006 5/	*	...	...	*	*	*	*	*	*	*	*	*
2005 5/	*	...	...	*	*	*	*	*	*	*	*	*
2004 5/	*	...	...	*	*	*	*	*	*	*	*	*
2003 5/	*	...	...	*	*	*	*	*	*	*	*	*
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
Non-Hispanic black, both sexes												
2006 5/	*	...	...	*	*	*	*	*	*	*	*	*
2005 5/	*	...	...	*	*	*	*	*	*	*	*	*
2004 5/	*	...	...	*	*	*	*	*	*	*	*	*
2003 5/	*	...	...	*	*	*	*	*	*	*	*	*
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2006 5/	*	...	...	*	*	*	*	*	*	*	*	*
2005 5/	*	...	...	*	*	*	*	*	*	*	*	*
2004 5/	*	...	...	*	*	*	*	*	*	*	*	*
2003 5/	*	...	...	*	*	*	*	*	*	*	*	*
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
All origins 3/, both sexes												
2006	0.2	...	...	*	0.4	0.8	0.4	0.1	*	*	*	*
2005	0.2	...	...	*	0.4	0.7	0.4	0.1	*	*	*	*
2004	0.2	...	...	*	0.4	0.7	0.3	0.2	*	*	*	*
2003	0.2	...	...	*	0.3	0.5	0.4	0.1	*	*	*	*
2002	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2000	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
1999	0.1	...	...	*	0.2	0.4	0.3	*	*	*	*	*
All origins 3/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2006	0.5	...	...	*	0.8	1.6	0.9	0.3	*	*	*	*
2005	0.5	...	...	*	0.9	1.5	0.9	0.2	*	*	*	*
2004	0.4	...	...	*	0.8	1.4	0.7	0.4	*	*	*	*
2003	0.3	...	...	*	0.5	1.1	0.7	0.1	*	*	*	*
2002	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
2001	0.3	...	...	*	0.5	0.9	0.4	*	*	*	*	*
2000	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*
1999	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
Hispanic, both sexes												
2006	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*
2005	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*
2004	0.3	...	...	*	0.4	0.6	0.4	*	*	*	*	*
2003	0.2	...	...	*	0.3	0.5	0.5	*	*	*	*	*
2002	0.2	...	...	*	*	0.4	0.4	*	*	*	*	*
2001	0.2	...	...	*	0.4	0.6	*	*	*	*	*	*
2000	0.2	...	...	*	0.3	0.5	0.4	*	*	*	*	*
1999	0.2	...	...	*	*	0.4	0.5	*	*	*	*	*
Hispanic, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2006	0.7	...	...	*	1.1	1.9	1.1	*	*	*	*	*
2005	0.6	...	...	*	1.0	1.7	1.1	*	*	*	*	*
2004	0.6	...	...	*	0.9	1.4	0.9	*	*	*	*	*
2003	0.5	...	...	*	0.7	1.2	1.0	*	*	*	*	*
2002	0.4	...	...	*	*	0.8	0.8	*	*	*	*	*
2001	0.5	...	...	*	0.8	1.2	*	*	*	*	*	*
2000	0.5	...	...	*	0.7	1.1	0.9	*	*	*	*	*
1999	0.4	...	...	*	*	0.9	1.0	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
Non-Hispanic 4/, both sexes												
2006	0.2	...	...	*	0.4	0.7	0.4	0.1	*	*	*	*
2005	0.2	...	...	*	0.4	0.7	0.4	0.1	*	*	*	*
2004	0.2	...	...	*	0.4	0.7	0.3	0.2	*	*	*	*
2003	0.2	...	...	*	0.2	0.5	0.3	0.1	*	*	*	*
2002	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2000	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
1999	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
Non-Hispanic 4/, male												
2006	...	...	...	...	...	...	...	...	...	...	...	...
2005	...	...	...	...	...	...	...	...	...	...	...	...
2004	...	...	...	...	...	...	...	...	...	...	...	...
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, female												
2006	0.4	...	...	*	0.8	1.5	0.8	0.3	*	*	*	*
2005	0.5	...	...	*	0.8	1.5	0.9	0.3	*	*	*	*
2004	0.4	...	...	*	0.7	1.4	0.6	0.3	*	*	*	*
2003	0.3	...	...	*	0.5	1.0	0.7	0.1	*	*	*	*
2002	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
2001	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
2000	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
1999	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
Non-Hispanic white, both sexes												
2006 5/	0.1	...	...	*	0.3	0.5	0.3	0.1	*	*	*	*
2005 5/	0.2	...	...	*	0.3	0.5	0.3	0.1	*	*	*	*
2004 5/	0.1	...	...	*	0.2	0.5	0.2	0.1	*	*	*	*
2003 5/	0.1	...	...	*	0.2	0.3	0.2	*	*	*	*	*
2002	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
2001	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
2000	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
1999	0.1	...	...	*	0.1	0.3	0.1	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2006 5/	0.3	...	...	*	0.6	0.9	0.5	0.2	*	*	*	*
2005 5/	0.3	...	...	*	0.6	1.0	0.6	0.2	*	*	*	*
2004 5/	0.3	...	...	*	0.4	1.0	0.5	0.3	*	*	*	*
2003 5/	0.2	...	...	*	0.3	0.7	0.4	*	*	*	*	*
2002	0.1	...	...	*	0.2	0.5	0.2	*	*	*	*	*
2001	0.1	...	...	*	0.3	0.6	0.2	*	*	*	*	*
2000	0.1	...	...	*	0.3	0.5	0.3	*	*	*	*	*
1999	0.1	...	...	*	0.2	0.5	0.3	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
Non-Hispanic black, both sexes												
2006 5/	0.6	...	...	*	0.9	2.0	1.1	0.4	*	*	*	*
2005 5/	0.7	...	...	*	1.1	2.0	1.1	*	*	*	*	*
2004 5/	0.6	...	...	*	1.1	1.9	0.7	0.5	*	*	*	*
2003 5/	0.5	...	...	*	0.6	1.5	1.0	*	*	*	*	*
2002	0.4	...	...	*	0.7	1.2	0.7	*	*	*	*	*
2001	0.4	...	...	*	0.7	1.2	0.6	*	*	*	*	*
2000	0.3	...	...	*	0.5	1.1	0.6	*	*	*	*	*
1999	0.4	...	...	*	0.8	1.2	0.7	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	...	...	...	...	...	...	...	...	...	...	...	...
2005 5/	...	...	...	...	...	...	...	...	...	...	...	...
2004 5/	...	...	...	...	...	...	...	...	...	...	...	...
2003 5/	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2006 5/	1.2	...	...	*	1.8	3.9	2.0	0.8	*	*	*	*
2005 5/	1.3	...	...	*	2.2	3.9	2.1	*	*	*	*	*
2004 5/	1.2	...	...	*	2.1	3.7	1.3	0.9	*	*	*	*
2003 5/	0.9	...	...	*	1.1	2.8	1.8	*	*	*	*	*
2002	0.7	...	...	*	1.5	2.2	1.2	*	*	*	*	*
2001	0.7	...	...	*	1.3	2.2	1.2	*	*	*	*	*
2000	0.7	...	...	*	1.1	2.1	1.1	*	*	*	*	*
1999	0.8	...	...	*	1.5	2.3	1.3	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
All origins 3/, both sexes												
2006	4.8	346.7	0.4	0.1	*	*	*	*	*	*	*	*
2005	4.9	351.2	0.4	0.1	*	*	*	*	*	*	*	*
2004	4.8	345.3	0.4	0.1	*	*	*	*	*	*	*	*
2003	4.9	356.0	0.5	*	*	*	*	*	*	*	*	*
2002	4.9	349.7	0.4	0.1	0.1	*	*	*	*	*	*	*
2001	4.9	340.5	0.5	0.1	*	*	*	*	*	*	*	*
2000	5.0	365.7	0.5	0.1	*	*	*	*	*	*	*	*
1999	5.1	371.0	0.6	0.1	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	5.5	379.8	0.4	*	*	*	*	*	*	*	*	*
2005	5.6	386.1	0.4	*	*	*	*	*	*	*	*	*
2004	5.6	384.2	0.5	0.1	*	*	*	*	*	*	*	*
2003	5.7	397.4	0.5	*	*	*	*	*	*	*	*	*
2002	5.7	385.7	0.5	*	*	*	*	*	*	*	*	*
2001	5.6	376.6	0.6	0.1	*	*	*	*	*	*	*	*
2000	5.8	403.9	0.7	0.1	*	*	*	*	*	*	*	*
1999	5.9	408.6	0.7	0.1	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	4.2	312.1	0.4	*	*	*	*	*	*	*	*	*
2005	4.2	314.6	0.3	*	*	*	*	*	*	*	*	*
2004	4.1	304.6	0.3	*	*	*	*	*	*	*	*	*
2003	4.2	312.8	0.5	*	*	*	*	*	*	*	*	*
2002	4.2	312.0	0.3	*	*	*	*	*	*	*	*	*
2001	4.2	302.6	0.4	*	*	*	*	*	*	*	*	*
2000	4.3	325.6	0.4	*	*	*	*	*	*	*	*	*
1999	4.4	331.7	0.5	0.1	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
Hispanic, both sexes												
2006	6.3	286.4	*	*	*	*	*	*	*	*	*	*
2005	6.6	300.6	*	*	*	*	*	*	*	*	*	*
2004	6.5	293.4	*	*	*	*	*	*	*	*	*	*
2003	6.6	300.3	*	*	*	*	*	*	*	*	*	*
2002	6.2	284.8	*	*	*	*	*	*	*	*	*	*
2001	6.0	269.6	*	*	*	*	*	*	*	*	*	*
2000	6.1	274.9	*	*	*	*	*	*	*	*	*	*
1999	6.3	281.4	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	6.8	312.6	*	*	*	*	*	*	*	*	*	*
2005	7.3	336.8	*	*	*	*	*	*	*	*	*	*
2004	7.2	329.7	*	*	*	*	*	*	*	*	*	*
2003	7.1	326.4	*	*	*	*	*	*	*	*	*	*
2002	6.8	313.1	*	*	*	*	*	*	*	*	*	*
2001	6.7	302.0	*	*	*	*	*	*	*	*	*	*
2000	6.6	299.0	*	*	*	*	*	*	*	*	*	*
1999	6.8	305.4	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	5.8	259.0	*	*	*	*	*	*	*	*	*	*
2005	5.8	262.8	*	*	*	*	*	*	*	*	*	*
2004	5.7	255.5	*	*	*	*	*	*	*	*	*	*
2003	6.1	273.0	*	*	*	*	*	*	*	*	*	*
2002	5.6	255.2	*	*	*	*	*	*	*	*	*	*
2001	5.3	235.9	*	*	*	*	*	*	*	*	*	*
2000	5.5	249.7	*	*	*	*	*	*	*	*	*	*
1999	5.8	256.4	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
Non-Hispanic 4/, both sexes												
2006	4.5	360.8	0.4	*	*	*	*	*	*	*	*	*
2005	4.6	361.7	0.4	0.1	*	*	*	*	*	*	*	*
2004	4.5	355.1	0.4	0.1	*	*	*	*	*	*	*	*
2003	4.6	365.3	0.5	*	*	*	*	*	*	*	*	*
2002	4.7	361.1	0.4	0.1	*	*	*	*	*	*	*	*
2001	4.6	353.5	0.5	0.1	*	*	*	*	*	*	*	*
2000	4.7	380.5	0.5	0.1	*	*	*	*	*	*	*	*
1999	4.8	383.8	0.6	0.1	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	5.2	395.8	0.5	*	*	*	*	*	*	*	*	*
2005	5.2	396.2	0.5	*	*	*	*	*	*	*	*	*
2004	5.3	394.5	0.5	*	*	*	*	*	*	*	*	*
2003	5.4	409.7	0.5	*	*	*	*	*	*	*	*	*
2002	5.4	398.6	0.5	*	*	*	*	*	*	*	*	*
2001	5.4	390.1	0.6	0.1	*	*	*	*	*	*	*	*
2000	5.5	421.7	0.7	*	*	*	*	*	*	*	*	*
1999	5.6	423.2	0.8	0.1	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	3.9	324.1	0.4	*	*	*	*	*	*	*	*	*
2005	3.9	325.6	*	*	*	*	*	*	*	*	*	*
2004	3.8	313.8	*	*	*	*	*	*	*	*	*	*
2003	3.8	318.9	0.5	*	*	*	*	*	*	*	*	*
2002	4.0	321.7	0.3	*	*	*	*	*	*	*	*	*
2001	3.9	315.1	0.3	*	*	*	*	*	*	*	*	*
2000	4.0	337.2	0.4	*	*	*	*	*	*	*	*	*
1999	4.1	342.4	0.4	0.1	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
Non-Hispanic white, both sexes												
2006 5/	3.0	261.5	0.3	*	*	*	*	*	*	*	*	*
2005 5/	3.1	265.1	0.3	*	*	*	*	*	*	*	*	*
2004 5/	3.1	261.7	0.3	*	*	*	*	*	*	*	*	*
2003 5/	3.2	274.8	0.3	*	*	*	*	*	*	*	*	*
2002	3.2	270.0	0.2	*	*	*	*	*	*	*	*	*
2001	3.2	260.3	0.4	*	*	*	*	*	*	*	*	*
2000	3.2	275.5	0.4	0.1	*	*	*	*	*	*	*	*
1999	3.2	275.1	0.4	0.1	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	3.5	289.6	0.4	*	*	*	*	*	*	*	*	*
2005 5/	3.6	290.8	*	*	*	*	*	*	*	*	*	*
2004 5/	3.6	294.5	*	*	*	*	*	*	*	*	*	*
2003 5/	3.8	310.6	*	*	*	*	*	*	*	*	*	*
2002	3.8	303.0	*	*	*	*	*	*	*	*	*	*
2001	3.7	286.6	0.5	*	*	*	*	*	*	*	*	*
2000	3.7	305.5	0.5	*	*	*	*	*	*	*	*	*
1999	3.8	304.4	0.5	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	2.5	231.9	*	*	*	*	*	*	*	*	*	*
2005 5/	2.7	238.1	*	*	*	*	*	*	*	*	*	*
2004 5/	2.5	227.3	*	*	*	*	*	*	*	*	*	*
2003 5/	2.6	237.3	*	*	*	*	*	*	*	*	*	*
2002	2.7	235.5	*	*	*	*	*	*	*	*	*	*
2001	2.7	232.8	*	*	*	*	*	*	*	*	*	*
2000	2.7	243.9	*	*	*	*	*	*	*	*	*	*
1999	2.7	244.2	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
Non-Hispanic black, both sexes												
2006 5/	13.0	767.0	0.9	*	*	*	*	*	*	*	*	*
2005 5/	13.0	763.4	0.9	*	*	*	*	*	*	*	*	*
2004 5/	12.8	742.1	0.9	*	*	*	*	*	*	*	*	*
2003 5/	12.9	740.3	1.2	*	*	*	*	*	*	*	*	*
2002	13.2	731.8	1.1	*	*	*	*	*	*	*	*	*
2001	13.2	765.0	0.9	*	*	*	*	*	*	*	*	*
2000	13.9	838.3	1.1	*	*	*	*	*	*	*	*	*
1999	14.4	872.7	1.2	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	15.0	830.0	*	*	*	*	*	*	*	*	*	*
2005 5/	15.2	837.9	*	*	*	*	*	*	*	*	*	*
2004 5/	15.1	814.9	*	*	*	*	*	*	*	*	*	*
2003 5/	15.5	827.9	*	*	*	*	*	*	*	*	*	*
2002	15.4	791.0	*	*	*	*	*	*	*	*	*	*
2001	15.8	846.6	*	*	*	*	*	*	*	*	*	*
2000	16.6	932.9	*	*	*	*	*	*	*	*	*	*
1999	17.1	968.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	11.1	701.2	*	*	*	*	*	*	*	*	*	*
2005 5/	10.9	685.6	*	*	*	*	*	*	*	*	*	*
2004 5/	10.8	666.4	*	*	*	*	*	*	*	*	*	*
2003 5/	10.6	649.2	*	*	*	*	*	*	*	*	*	*
2002	11.3	670.1	*	*	*	*	*	*	*	*	*	*
2001	10.9	679.1	*	*	*	*	*	*	*	*	*	*
2000	11.5	740.7	*	*	*	*	*	*	*	*	*	*
1999	12.0	774.4	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
All origins 3/, both sexes												
2006	3.5	140.9	3.2	0.9	1.1	1.1	1.1	1.5	2.3	1.9	3.0	5.6
2005	3.5	135.2	3.2	1.0	1.2	1.1	1.2	1.6	2.4	1.8	3.1	6.4
2004	3.6	137.9	3.6	1.0	1.2	1.0	1.1	1.7	2.2	2.3	3.6	6.1
2003	3.6	140.4	3.4	0.9	1.1	1.1	1.3	1.6	2.4	2.3	3.6	6.8
2002	3.7	139.4	3.4	1.0	1.2	1.2	1.3	1.8	2.4	2.4	3.8	6.6
2001	3.7	136.7	3.6	0.9	1.3	1.2	1.3	1.6	2.3	2.4	3.7	7.2
2000	3.8	150.9	3.2	1.0	1.1	1.2	1.3	1.6	2.3	2.5	3.9	7.5
1999	3.7	144.2	3.6	1.0	1.1	1.2	1.2	1.8	2.3	2.5	4.0	7.8
All origins 3/, male												
2006	3.8	146.6	2.9	0.9	1.3	1.3	1.2	1.8	2.6	2.1	3.6	6.8
2005	3.7	141.1	3.2	0.9	1.5	1.2	1.2	1.7	2.5	1.8	3.6	6.9
2004	3.8	139.0	3.9	1.0	1.3	1.2	1.3	1.8	2.3	2.5	3.7	6.8
2003	3.8	147.6	3.4	0.9	1.3	1.2	1.4	1.7	2.5	2.4	3.5	6.7
2002	3.9	142.6	3.5	1.0	1.4	1.4	1.3	1.9	2.5	2.6	4.0	7.0
2001	3.9	143.1	3.6	1.0	1.5	1.2	1.4	1.7	2.3	2.7	4.1	7.7
2000	4.0	155.4	3.3	1.0	1.3	1.4	1.4	1.9	2.3	2.7	4.0	10.6
1999	3.9	147.9	4.0	1.0	1.3	1.2	1.3	1.9	2.5	2.5	4.2	8.5
All origins 3/, female												
2006	3.2	134.9	3.4	0.8	0.9	0.9	0.9	1.3	2.0	1.8	2.6	5.1
2005	3.3	129.0	3.3	1.0	0.9	1.0	1.1	1.6	2.2	1.8	2.8	6.2
2004	3.4	136.8	3.2	0.9	1.0	0.9	1.0	1.5	2.1	2.2	3.6	5.8
2003	3.4	132.8	3.5	0.9	0.9	1.0	1.2	1.5	2.3	2.2	3.7	6.8
2002	3.5	136.1	3.3	1.0	1.1	1.0	1.2	1.6	2.3	2.3	3.8	6.5
2001	3.4	129.9	3.7	0.8	1.0	1.1	1.2	1.6	2.3	2.2	3.5	7.0
2000	3.5	146.2	3.1	1.0	0.9	1.0	1.1	1.4	2.4	2.4	3.9	6.3
1999	3.5	140.3	3.2	1.1	0.9	1.1	1.1	1.6	2.2	2.6	3.9	7.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Hispanic, both sexes												
2006	4.4	151.0	3.3	0.8	1.0	0.7	0.6	1.1	1.3	1.7	*	*
2005	4.4	148.3	3.3	1.1	1.0	0.7	0.7	1.2	1.7	*	3.2	*
2004	4.3	141.0	4.0	0.9	0.9	0.7	0.6	1.1	1.4	1.6	3.4	*
2003	4.5	153.0	3.7	0.9	0.8	0.8	0.7	1.0	1.7	*	*	*
2002	4.4	150.7	3.5	1.0	1.0	0.7	0.7	1.1	1.3	*	3.8	*
2001	4.4	146.2	3.4	0.8	1.1	0.7	0.8	1.1	1.3	1.9	4.3	*
2000	4.6	153.3	3.8	0.8	0.8	0.7	0.8	1.1	1.8	2.0	4.7	*
1999	4.4	141.8	3.5	0.9	0.9	0.9	0.7	1.4	1.6	2.9	*	*
Hispanic, male												
2006	4.6	153.4	3.6	0.8	1.2	0.9	0.7	1.0	*	*	*	*
2005	4.5	151.2	3.5	1.2	1.1	0.7	0.8	1.1	2.0	*	*	*
2004	4.2	134.3	4.6	1.3	1.1	0.6	0.7	1.3	*	*	*	*
2003	4.5	156.9	3.7	0.7	0.8	0.9	0.8	1.2	*	*	*	*
2002	4.5	153.1	3.8	0.9	1.1	0.9	*	1.3	*	*	*	*
2001	4.7	153.4	3.5	1.0	1.1	0.7	1.0	1.5	*	*	*	*
2000	4.7	153.3	4.0	0.8	1.1	0.7	1.0	1.4	*	*	*	*
1999	4.3	138.1	4.3	0.9	0.9	0.7	*	1.6	*	*	*	*
Hispanic, female												
2006	4.2	148.5	3.1	0.7	0.7	*	*	1.3	*	*	*	*
2005	4.4	145.2	3.1	1.0	0.9	0.6	0.7	1.3	*	*	*	*
2004	4.3	147.9	3.3	0.6	0.6	0.8	*	*	*	*	*	*
2003	4.4	148.9	3.7	1.0	0.8	0.6	*	*	*	*	*	*
2002	4.3	148.2	3.3	1.0	0.9	*	0.8	*	*	*	*	*
2001	4.1	138.8	3.3	0.6	1.1	*	*	*	*	*	*	*
2000	4.4	153.3	3.5	0.7	*	0.7	*	*	*	*	*	*
1999	4.5	145.8	2.7	0.9	0.9	1.0	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Non-Hispanic 4/, both sexes												
2006	3.3	136.8	3.1	0.9	1.1	1.2	1.2	1.6	2.4	1.9	3.1	5.8
2005	3.3	130.4	3.2	1.0	1.2	1.2	1.3	1.7	2.4	1.9	3.1	6.6
2004	3.4	135.9	3.4	1.0	1.2	1.1	1.2	1.7	2.2	2.4	3.7	6.2
2003	3.5	136.0	3.4	0.9	1.1	1.1	1.3	1.7	2.4	2.3	3.7	6.7
2002	3.6	134.9	3.3	1.0	1.2	1.3	1.4	1.8	2.5	2.5	3.8	6.7
2001	3.5	132.8	3.7	0.9	1.3	1.3	1.4	1.7	2.4	2.5	3.7	7.2
2000	3.6	148.8	3.1	1.0	1.2	1.3	1.3	1.7	2.4	2.6	3.9	7.5
1999	3.6	143.1	3.6	1.1	1.2	1.2	1.3	1.8	2.4	2.5	4.0	7.9
Non-Hispanic 4/, male												
2006	3.6	143.6	2.7	0.9	1.3	1.4	1.3	1.9	2.7	2.1	3.6	6.8
2005	3.6	137.1	3.1	0.9	1.6	1.3	1.3	1.8	2.6	1.9	3.6	7.2
2004	3.6	139.1	3.7	0.9	1.4	1.3	1.4	1.9	2.3	2.5	3.7	7.0
2003	3.7	144.2	3.3	1.0	1.4	1.2	1.5	1.8	2.5	2.4	3.5	6.7
2002	3.8	138.1	3.5	1.0	1.4	1.5	1.4	2.0	2.6	2.6	3.9	7.0
2001	3.7	139.4	3.6	1.0	1.6	1.4	1.5	1.7	2.3	2.7	4.0	7.6
2000	3.9	154.4	3.1	1.0	1.4	1.5	1.5	1.9	2.2	2.7	3.9	10.4
1999	3.8	148.4	3.9	1.1	1.4	1.3	1.4	1.9	2.6	2.4	4.2	8.5
Non-Hispanic 4/, female												
2006	3.0	129.7	3.5	0.9	0.9	1.0	1.0	1.3	2.1	1.8	2.7	5.3
2005	3.1	123.3	3.3	1.0	0.9	1.1	1.2	1.6	2.3	1.9	2.8	6.3
2004	3.2	132.6	3.2	1.0	1.1	1.0	1.0	1.6	2.1	2.2	3.6	5.9
2003	3.3	127.6	3.4	0.9	0.9	1.0	1.2	1.6	2.3	2.2	3.8	6.8
2002	3.4	131.5	3.2	1.0	1.1	1.1	1.3	1.7	2.4	2.3	3.8	6.6
2001	3.3	125.8	3.7	0.9	1.0	1.2	1.3	1.7	2.4	2.2	3.5	7.0
2000	3.4	142.8	3.0	1.0	1.0	1.1	1.2	1.4	2.5	2.4	3.9	6.4
1999	3.4	137.4	3.3	1.1	0.9	1.1	1.2	1.6	2.2	2.6	3.9	7.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Non-Hispanic white, both sexes												
2006 5/	3.2	129.5	2.8	0.8	1.0	1.2	1.2	1.6	2.5	2.0	3.2	5.9
2005 5/	3.2	123.0	2.9	0.9	1.2	1.2	1.3	1.8	2.6	2.0	3.3	6.7
2004 5/	3.3	130.5	2.9	0.9	1.1	1.2	1.2	1.8	2.4	2.5	3.7	6.5
2003 5/	3.3	129.8	3.0	0.9	1.1	1.2	1.4	1.7	2.6	2.3	3.8	6.9
2002	3.4	131.2	3.0	1.0	1.2	1.3	1.4	1.9	2.5	2.5	3.9	6.9
2001	3.4	127.4	3.1	0.9	1.2	1.3	1.4	1.7	2.5	2.5	3.8	7.4
2000	3.4	141.2	2.6	0.9	1.1	1.3	1.3	1.7	2.4	2.6	4.0	7.8
1999	3.4	136.8	3.0	0.9	1.1	1.3	1.3	1.8	2.5	2.5	4.0	8.0
Non-Hispanic white, male												
2006 5/	3.4	135.2	2.6	0.9	1.2	1.4	1.4	1.8	2.8	2.1	3.7	6.8
2005 5/	3.4	129.9	3.0	0.8	1.6	1.3	1.3	1.9	2.7	2.0	3.8	7.2
2004 5/	3.4	132.8	3.1	0.9	1.2	1.4	1.4	2.0	2.4	2.7	3.7	7.4
2003 5/	3.4	136.1	2.9	0.9	1.3	1.2	1.5	1.8	2.6	2.3	3.4	6.8
2002	3.6	137.2	3.2	1.0	1.4	1.6	1.4	2.0	2.6	2.7	3.8	7.3
2001	3.6	135.5	3.1	1.0	1.4	1.5	1.5	1.7	2.4	2.9	4.1	7.4
2000	3.6	147.9	2.6	0.9	1.3	1.6	1.4	1.9	2.3	2.6	4.1	10.8
1999	3.6	144.4	3.2	0.9	1.4	1.3	1.4	2.0	2.7	2.4	4.1	8.6
Non-Hispanic white, female												
2006 5/	2.9	123.5	2.9	0.7	0.9	1.0	1.0	1.4	2.2	1.9	2.8	5.5
2005 5/	3.0	115.8	2.9	0.9	0.8	1.0	1.3	1.7	2.4	2.0	2.9	6.5
2004 5/	3.2	128.0	2.8	0.9	1.0	1.1	1.1	1.6	2.3	2.4	3.7	6.1
2003 5/	3.2	123.2	3.0	0.9	0.9	1.1	1.3	1.7	2.5	2.3	4.0	6.9
2002	3.3	125.0	2.9	1.0	1.1	1.1	1.3	1.8	2.5	2.4	3.9	6.7
2001	3.2	118.8	3.2	0.7	1.0	1.2	1.3	1.7	2.6	2.3	3.6	7.4
2000	3.2	134.2	2.6	1.0	1.0	1.1	1.2	1.6	2.5	2.6	4.0	6.6
1999	3.3	128.9	2.9	1.0	0.8	1.2	1.2	1.7	2.3	2.6	4.0	7.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Non-Hispanic black, both sexes												
2006 5/	4.6	174.3	4.2	1.1	1.4	1.3	1.4	1.9	2.0	1.6	3.0	5.6
2005 5/	4.4	163.4	4.1	1.3	1.6	1.5	1.4	1.4	2.0	*	2.1	6.0
2004 5/	4.6	164.3	5.1	1.3	1.8	1.1	1.3	1.7	1.7	1.6	4.0	*
2003 5/	4.6	160.5	4.8	1.1	1.5	1.4	1.2	1.7	2.2	2.4	3.1	5.9
2002	4.7	156.9	4.7	1.3	1.7	1.5	1.6	1.8	2.7	2.4	3.4	*
2001	4.7	163.5	5.4	1.1	2.0	1.3	1.6	1.8	1.9	2.1	3.0	*
2000	5.0	184.1	5.2	1.4	1.6	1.4	1.6	1.6	2.4	2.7	2.8	*
1999	4.8	174.8	5.8	1.6	1.5	1.1	1.4	1.9	2.0	2.9	4.3	7.4
Non-Hispanic black, male												
2006 5/	5.1	182.2	3.2	0.9	1.6	1.7	1.5	2.4	2.4	*	*	*
2005 5/	4.8	167.1	3.6	1.2	2.0	1.7	1.7	1.5	2.3	*	*	*
2004 5/	5.2	170.5	5.9	1.2	2.4	1.4	1.6	1.8	2.0	*	*	*
2003 5/	5.2	173.1	5.0	1.2	1.8	1.5	1.2	2.0	2.8	3.6	*	*
2002	5.2	153.3	4.8	1.4	2.0	1.8	1.7	2.3	3.5	3.1	*	*
2001	5.1	167.5	5.5	1.0	2.5	1.2	1.8	1.7	2.1	*	*	*
2000	5.5	186.0	5.6	1.5	2.1	1.7	1.9	2.2	*	3.9	*	*
1999	5.3	179.5	6.6	1.6	1.8	1.3	1.6	2.3	2.1	*	*	*
Non-Hispanic black, female												
2006 5/	4.2	166.0	5.3	1.2	1.1	1.0	1.2	1.5	1.7	*	*	*
2005 5/	4.1	159.6	4.7	1.4	1.2	1.4	1.1	1.4	1.7	*	*	*
2004 5/	4.1	157.8	4.3	1.4	1.2	0.8	0.9	1.7	1.4	*	3.4	*
2003 5/	4.0	147.3	4.7	1.1	1.2	1.4	1.3	1.5	1.8	*	*	*
2002	4.4	160.6	4.7	1.1	1.3	1.2	1.6	1.5	2.0	*	*	*
2001	4.4	159.2	5.4	1.3	1.4	1.4	1.5	1.8	1.8	2.4	*	*
2000	4.5	182.2	4.8	1.2	1.2	1.1	1.4	1.2	3.1	*	*	*
1999	4.4	170.0	5.0	1.6	1.1	1.0	1.2	1.5	1.9	3.2	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
All origins 3/, both sexes												
2006	10.6	83.8	1.5	0.3	1.6	2.5	3.7	5.2	6.5	12.2	41.7	236.1
2005	10.8	87.4	1.4	0.4	1.8	2.9	4.7	6.7	7.2	12.8	39.0	224.4
2004	10.3	82.4	1.2	0.3	1.6	2.7	4.3	5.8	7.0	12.6	39.8	221.5
2003	10.8	82.9	1.6	0.4	2.0	3.3	5.4	7.0	7.4	13.8	38.9	224.1
2002	10.4	85.7	1.6	0.3	1.8	3.3	5.3	6.7	7.5	13.1	38.8	208.0
2001	11.4	86.5	1.8	0.4	2.4	3.9	6.9	8.9	9.1	15.1	39.8	206.4
2000	11.3	95.6	1.6	0.4	2.2	3.9	6.6	8.3	9.3	16.5	40.7	208.1
1999	9.6	93.2	1.5	0.3	1.6	2.7	4.6	5.9	7.8	14.6	36.1	187.7
All origins 3/, male												
2006	9.5	95.2	1.6	0.4	2.0	3.5	4.8	6.7	8.3	14.8	44.2	191.1
2005	10.0	101.8	1.6	0.5	2.5	4.0	5.8	8.7	9.3	15.5	40.7	183.9
2004	9.5	96.9	1.2	0.3	2.1	3.6	5.6	7.6	9.1	15.0	41.8	183.6
2003	10.2	96.5	1.8	0.4	2.7	4.5	6.9	9.2	9.6	17.1	41.9	177.9
2002	10.1	98.5	1.8	0.4	2.4	4.5	6.9	8.8	10.0	15.9	43.8	175.5
2001	11.3	98.9	2.1	0.5	3.2	5.2	9.0	11.9	12.0	18.0	43.8	175.0
2000	11.5	110.6	1.7	0.5	2.9	5.3	8.9	11.5	12.3	20.6	46.0	182.7
1999	9.4	106.6	1.7	0.4	2.1	3.6	6.0	8.1	10.4	18.2	40.1	166.3
All origins 3/, female												
2006	11.7	71.9	1.3	0.3	1.1	1.5	2.7	3.8	4.8	10.0	39.9	257.2
2005	11.5	72.3	1.3	0.3	1.1	1.9	3.6	4.7	5.3	10.6	37.8	243.0
2004	11.0	67.3	1.1	0.3	1.0	1.8	3.0	4.0	5.1	10.6	38.4	238.5
2003	11.4	68.6	1.5	0.4	1.4	2.1	3.8	5.0	5.3	10.9	36.9	244.6
2002	10.7	72.3	1.4	0.3	1.2	2.1	3.8	4.7	5.3	10.8	35.4	222.1
2001	11.4	73.6	1.5	0.4	1.5	2.5	4.9	6.1	6.3	12.7	37.2	219.5
2000	11.2	79.9	1.5	0.4	1.5	2.5	4.4	5.3	6.5	13.2	37.2	218.5
1999	9.7	79.2	1.3	0.3	1.0	1.9	3.2	3.8	5.5	11.7	33.5	196.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Hispanic, both sexes												
2006	4.0	52.5	1.2	*	1.3	1.9	2.5	3.0	3.8	6.9	18.7	117.0
2005	4.4	59.5	1.2	0.3	1.5	2.4	3.1	3.6	3.6	7.8	18.6	105.2
2004	3.9	50.4	1.1	*	1.6	2.1	3.1	3.6	3.9	6.2	16.4	99.4
2003	4.1	56.8	0.9	0.3	1.4	2.3	3.1	3.7	3.5	6.1	18.4	110.9
2002	4.0	56.6	1.3	*	1.3	2.0	3.0	3.9	4.8	7.0	18.7	91.5
2001	5.9	60.7	1.6	0.5	2.8	4.2	6.0	8.3	6.9	9.9	19.3	108.2
2000	5.9	65.6	1.7	0.5	2.1	4.0	6.7	8.0	6.7	9.9	20.5	102.2
1999	3.8	54.9	1.0	*	1.3	2.2	3.0	3.1	3.8	7.0	19.4	87.0
Hispanic, male												
2006	4.5	62.1	1.1	*	1.9	2.9	3.9	4.2	5.4	9.7	17.4	90.2
2005	5.1	68.9	1.3	*	2.0	3.6	4.3	5.5	5.1	10.2	21.0	99.6
2004	4.5	60.0	1.2	*	2.3	2.8	4.5	4.9	5.8	7.1	15.7	86.9
2003	4.8	69.9	*	*	1.7	3.4	4.4	5.7	5.6	8.1	18.8	95.4
2002	4.6	66.4	1.6	*	1.7	2.9	4.0	4.7	6.3	9.6	23.4	88.7
2001	7.4	67.2	1.8	*	3.9	6.0	8.5	12.6	10.6	14.2	25.0	87.6
2000	7.5	78.6	2.1	*	2.9	5.8	9.9	12.4	9.0	14.3	25.6	106.8
1999	4.4	61.3	*	*	1.9	2.9	4.3	4.7	5.5	8.3	24.0	77.6
Hispanic, female												
2006	3.4	42.5	1.3	*	0.7	0.7	1.0	1.7	2.3	4.6	19.7	131.2
2005	3.5	49.8	1.2	*	0.9	0.9	1.8	1.6	2.3	5.9	17.0	108.2
2004	3.3	40.4	*	*	0.8	1.3	1.6	2.4	2.2	5.5	16.8	105.9
2003	3.2	43.2	*	*	1.0	1.0	1.6	1.7	*	4.6	18.2	118.8
2002	3.3	46.4	*	*	0.9	1.0	2.0	3.2	3.5	5.0	15.5	92.8
2001	4.3	53.9	*	0.6	1.5	2.1	3.4	4.1	3.6	6.5	15.4	118.2
2000	4.1	52.1	1.4	*	1.3	2.0	3.2	3.7	4.6	6.5	17.2	99.9
1999	3.1	48.2	*	*	*	1.3	1.6	1.7	2.4	6.0	16.3	91.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Non-Hispanic 4/, both sexes												
2006	11.7	92.7	1.5	0.3	1.6	2.6	3.9	5.4	6.6	12.5	43.0	241.9
2005	11.8	94.5	1.5	0.4	1.9	3.1	4.9	6.9	7.5	13.1	39.9	229.6
2004	11.2	90.9	1.2	0.3	1.6	2.8	4.5	5.9	7.2	13.0	40.9	226.7
2003	11.8	89.4	1.8	0.5	2.1	3.5	5.7	7.3	7.7	14.3	39.9	228.4
2002	11.3	92.8	1.7	0.4	1.9	3.6	5.6	6.9	7.7	13.5	39.6	211.9
2001	12.1	92.6	1.8	0.4	2.3	3.7	7.0	8.9	9.2	15.4	40.7	209.6
2000	12.0	102.4	1.6	0.4	2.2	3.8	6.5	8.2	9.3	16.8	41.4	211.6
1999	10.3	101.9	1.6	0.4	1.6	2.8	4.7	6.0	8.0	15.0	36.7	190.9
Non-Hispanic 4/, male												
2006	10.3	104.3	1.7	0.4	2.0	3.6	4.9	7.0	8.4	15.1	45.9	196.3
2005	10.8	110.2	1.7	0.5	2.6	4.0	5.9	8.9	9.6	15.8	41.6	187.5
2004	10.2	106.6	1.2	0.3	2.0	3.8	5.7	7.8	9.2	15.5	43.1	188.0
2003	11.0	102.9	2.0	0.5	2.8	4.7	7.2	9.4	9.9	17.7	43.0	181.1
2002	10.9	106.2	1.9	0.4	2.5	4.8	7.3	9.1	10.1	16.3	44.6	178.6
2001	11.8	106.2	2.1	0.5	3.0	4.9	9.0	11.7	12.0	18.2	44.6	178.2
2000	11.9	118.0	1.7	0.5	2.9	5.1	8.6	11.2	12.3	20.7	46.6	185.4
1999	10.0	116.7	1.9	0.4	2.2	3.7	6.1	8.2	10.6	18.7	40.6	169.5
Non-Hispanic 4/, female												
2006	13.0	80.5	1.3	0.3	1.2	1.7	3.0	4.0	5.0	10.4	41.1	263.1
2005	12.7	78.0	1.3	0.3	1.1	2.1	3.9	4.9	5.5	10.8	38.8	248.9
2004	12.1	74.3	1.2	0.3	1.0	1.9	3.2	4.2	5.3	10.9	39.4	244.0
2003	12.6	75.3	1.6	0.5	1.4	2.4	4.1	5.3	5.6	11.4	37.8	249.2
2002	11.7	78.7	1.4	0.3	1.3	2.3	4.0	4.8	5.4	11.2	36.3	226.3
2001	12.3	78.3	1.5	0.4	1.5	2.6	5.1	6.2	6.6	13.1	38.0	222.7
2000	12.1	86.1	1.5	0.4	1.5	2.6	4.5	5.4	6.6	13.6	38.0	222.3
1999	10.5	86.3	1.4	0.3	1.0	2.0	3.3	3.9	5.7	11.9	34.2	199.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Non-Hispanic white, both sexes												
2006 5/	12.4	77.1	1.2	0.3	1.6	2.6	3.8	5.2	6.3	12.1	42.8	247.4
2005 5/	12.3	77.9	1.2	0.3	1.8	3.0	4.8	6.5	6.9	12.5	39.8	234.9
2004 5/	11.7	71.6	1.0	0.2	1.5	2.8	4.3	5.6	6.7	12.5	40.7	230.0
2003 5/	12.3	72.6	1.5	0.4	2.2	3.4	5.5	6.7	7.1	13.7	40.1	232.9
2002	11.6	76.0	1.3	0.3	1.8	3.4	5.4	6.1	7.1	13.1	39.4	215.8
2001	12.4	73.1	1.5	0.3	2.2	3.6	6.9	8.5	8.6	14.7	40.4	213.3
2000	12.2	79.7	1.4	0.3	2.1	3.6	6.3	7.6	8.6	15.9	40.9	215.8
1999	10.2	79.3	1.4	0.3	1.4	2.5	4.1	5.2	7.3	14.3	35.7	193.8
Non-Hispanic white, male												
2006 5/	10.5	89.1	1.4	0.4	2.0	3.5	4.7	6.5	8.0	14.2	45.1	199.4
2005 5/	10.9	94.6	1.4	0.3	2.5	4.0	5.9	8.4	8.8	14.9	41.0	189.8
2004 5/	10.3	83.1	1.0	0.2	2.1	3.7	5.6	7.2	8.4	14.9	42.8	191.0
2003 5/	11.0	87.4	1.6	0.4	2.9	4.6	6.9	8.6	9.0	16.8	42.8	183.3
2002	10.7	88.2	1.4	0.3	2.5	4.7	6.9	7.9	9.3	15.7	43.6	181.5
2001	11.8	85.2	1.7	0.4	2.9	4.9	8.8	11.1	10.9	17.3	44.7	180.4
2000	11.8	91.8	1.7	0.4	2.8	4.8	8.5	10.3	11.1	19.3	46.0	189.6
1999	9.6	92.3	1.7	0.3	1.9	3.3	5.5	6.9	9.5	17.7	39.2	171.1
Non-Hispanic white, female												
2006 5/	14.2	64.5	1.1	0.2	1.1	1.7	2.9	3.9	4.7	10.3	41.2	269.7
2005 5/	13.7	60.4	1.0	0.2	1.1	2.0	3.7	4.6	5.2	10.4	39.0	255.5
2004 5/	13.0	59.4	1.0	0.2	1.0	1.8	3.0	4.0	5.0	10.4	39.3	247.5
2003 5/	13.5	57.1	1.4	0.4	1.4	2.2	4.0	4.8	5.4	11.0	38.3	254.7
2002	12.5	63.2	1.3	0.2	1.1	2.2	3.8	4.3	5.1	10.9	36.6	230.6
2001	13.1	60.4	1.3	0.3	1.5	2.4	5.0	5.9	6.4	12.5	37.6	227.0
2000	12.6	66.9	1.2	0.3	1.3	2.3	4.1	4.9	6.2	13.0	37.5	226.4
1999	10.9	65.5	1.0	0.3	0.9	1.8	2.8	3.5	5.2	11.4	33.4	202.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Non-Hispanic black, both sexes												
2006 5/	11.4	161.4	2.8	0.6	1.9	3.5	5.8	8.9	10.9	19.6	56.9	226.9
2005 5/	12.4	170.4	2.5	0.9	2.5	4.2	7.0	11.7	13.9	22.1	51.8	213.3
2004 5/	11.9	174.5	2.0	0.6	1.8	4.1	6.6	9.8	13.4	20.8	53.1	223.3
2003 5/	12.7	161.6	3.3	0.8	2.5	5.1	8.6	13.6	14.3	24.0	45.7	214.5
2002	12.6	165.4	2.7	0.8	2.6	5.3	8.8	13.9	13.9	20.4	49.9	198.1
2001	12.9	176.9	3.0	0.6	2.7	5.2	9.3	14.0	15.1	24.6	50.3	194.2
2000	13.9	205.6	2.3	0.7	3.2	5.9	9.3	14.8	17.3	29.2	56.1	189.2
1999	13.0	202.0	2.9	0.5	2.7	5.0	9.1	13.2	16.0	24.6	54.8	182.0
Non-Hispanic black, male												
2006 5/	12.0	172.0	3.1	0.7	2.3	4.7	7.3	12.5	14.0	28.0	70.4	196.1
2005 5/	13.3	187.0	2.7	0.9	3.4	5.4	8.1	15.2	19.6	29.5	62.5	196.0
2004 5/	12.8	208.3	2.2	*	2.3	5.3	8.1	13.8	18.9	25.4	59.3	182.9
2003 5/	13.8	171.9	3.6	0.8	3.0	7.0	11.2	17.9	20.7	32.2	52.8	197.1
2002	14.4	186.3	3.6	0.9	3.1	7.1	11.7	19.6	20.0	26.4	68.6	182.7
2001	14.2	195.9	3.7	0.8	3.8	6.1	11.4	18.2	23.8	30.7	52.2	186.0
2000	15.7	236.8	2.2	0.9	3.8	7.8	10.9	20.6	25.1	40.0	66.0	163.2
1999	14.9	228.0	2.9	0.6	3.6	6.9	11.6	19.4	23.0	32.8	66.9	180.7
Non-Hispanic black, female												
2006 5/	10.8	150.2	2.4	*	1.4	2.3	4.4	5.8	8.4	13.6	49.2	239.5
2005 5/	11.5	153.1	2.3	0.8	1.5	3.1	5.9	8.6	9.4	16.9	45.8	220.2
2004 5/	11.0	139.4	1.7	*	1.3	3.1	5.3	6.3	8.9	17.6	49.6	239.3
2003 5/	11.6	150.9	3.0	0.8	1.9	3.5	6.3	10.0	9.3	18.2	41.7	221.2
2002	10.9	143.6	1.9	0.7	2.0	3.7	6.2	9.0	8.9	16.1	39.3	204.0
2001	11.7	156.8	2.2	*	1.7	4.3	7.4	10.3	8.1	20.3	49.2	197.2
2000	12.4	173.4	2.4	*	2.6	4.3	7.8	9.9	11.0	21.6	50.5	198.9
1999	11.2	175.3	2.8	*	1.9	3.2	7.0	7.9	10.3	18.8	48.0	182.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
All origins 3/, both sexes												
2006	79.3	28.4	3.4	2.0	5.0	8.3	18.4	40.1	71.6	151.6	470.5	1,724.2
2005	73.4	29.2	3.8	2.1	4.9	7.9	18.5	40.3	69.5	146.1	434.3	1,559.1
2004	68.8	29.2	3.8	2.2	4.7	8.1	18.3	38.9	67.0	141.4	409.6	1,461.6
2003	69.3	29.7	4.0	2.3	4.9	8.4	18.9	38.9	67.1	144.4	411.5	1,508.4
2002	67.5	28.4	4.2	2.1	4.6	7.9	18.9	37.8	65.3	140.6	405.3	1,493.0
2001	65.5	28.0	4.2	2.1	4.5	7.8	18.0	36.2	63.4	140.9	395.3	1,476.7
2000	63.4	29.0	4.0	2.1	4.5	7.8	18.0	35.4	62.4	136.0	388.5	1,445.7
1999	61.5	30.2	4.2	2.3	4.6	7.6	17.7	33.9	60.5	136.6	373.0	1,410.7
All origins 3/, male												
2006	65.2	32.5	3.6	2.1	5.8	9.5	21.4	48.3	82.7	164.8	474.4	1,455.0
2005	60.9	34.8	4.3	2.3	5.8	8.9	22.0	49.1	80.9	156.1	437.0	1,305.3
2004	57.2	33.8	4.0	2.3	5.4	9.2	21.5	47.8	77.1	151.9	410.5	1,236.6
2003	57.3	34.8	4.2	2.6	5.6	9.3	22.3	47.2	76.5	155.6	415.1	1,258.1
2002	55.8	33.4	4.6	2.3	5.1	9.1	22.4	46.3	74.5	152.6	406.7	1,262.9
2001	53.8	31.4	4.6	2.2	5.2	8.6	21.3	43.7	71.2	153.4	397.4	1,271.6
2000	52.3	32.3	4.3	2.3	5.3	9.0	21.4	42.8	70.7	148.4	391.5	1,266.2
1999	51.0	35.0	4.5	2.3	5.3	8.8	21.6	40.5	69.0	148.7	384.2	1,225.4
All origins 3/, female												
2006	93.0	24.1	3.1	1.8	4.2	7.1	15.3	32.2	61.3	140.5	467.9	1,850.2
2005	85.6	23.2	3.3	1.9	4.0	6.8	15.0	31.8	59.0	137.7	432.5	1,675.6
2004	79.9	24.5	3.6	2.0	3.9	7.0	15.1	30.4	57.6	132.5	408.9	1,562.8
2003	81.0	24.5	3.8	1.9	4.2	7.4	15.6	31.0	58.4	135.0	409.1	1,619.1
2002	78.8	23.0	3.9	1.9	4.0	6.6	15.5	29.6	56.8	130.6	404.4	1,592.8
2001	76.7	24.5	3.8	2.0	3.8	6.9	14.6	29.1	56.2	130.5	394.0	1,562.6
2000	74.0	25.6	3.6	2.0	3.7	6.5	14.6	28.2	54.8	125.8	386.5	1,518.8
1999	71.6	25.3	3.9	2.2	3.8	6.4	13.9	27.4	52.7	126.7	365.7	1,485.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
Hispanic, both sexes												
2006	26.4	24.6	3.3	1.8	4.4	6.1	13.8	28.9	52.9	114.4	326.2	1,067.7
2005	24.0	29.1	4.1	1.6	4.0	6.1	13.3	28.3	52.5	107.5	290.7	890.7
2004	22.3	26.1	3.3	1.9	3.7	5.4	13.1	27.8	49.1	101.0	276.7	880.5
2003	23.1	27.1	4.3	2.1	4.1	6.1	14.0	30.3	53.5	106.9	284.5	887.2
2002	21.6	25.1	3.8	2.1	3.7	5.6	13.0	27.5	48.5	111.2	272.2	870.8
2001	21.2	22.9	3.7	1.7	3.7	5.2	12.3	26.3	49.7	107.2	277.5	991.0
2000	20.7	25.4	3.7	1.7	3.3	5.9	13.8	27.5	47.5	103.4	284.0	913.0
1999	21.5	26.5	3.7	1.6	3.5	6.5	14.0	28.2	51.4	109.3	263.5	1,039.7
Hispanic, male												
2006	24.9	28.0	3.3	1.9	5.3	7.4	17.6	37.3	64.2	127.6	334.2	897.6
2005	23.2	34.6	4.2	1.6	4.8	7.2	16.6	36.7	64.9	115.5	300.0	789.5
2004	21.3	28.7	3.5	2.1	4.5	6.2	17.2	37.5	57.6	109.6	264.7	770.5
2003	22.2	32.6	4.5	2.3	4.4	7.3	16.7	38.4	64.6	116.4	301.8	757.3
2002	21.1	27.2	4.1	2.4	4.0	6.6	17.1	35.5	56.4	126.0	290.1	808.2
2001	20.2	25.4	4.3	1.8	4.1	6.0	15.9	34.7	59.1	117.2	273.7	939.0
2000	20.2	25.3	3.6	1.8	3.9	7.0	18.2	35.6	57.7	119.9	273.9	870.7
1999	21.0	32.1	3.4	1.6	3.8	8.1	18.8	37.0	62.6	123.9	276.3	923.1
Hispanic, female												
2006	28.1	21.1	3.4	1.8	3.5	4.6	9.5	20.4	42.4	103.7	320.7	1,158.1
2005	24.9	23.3	3.9	1.6	3.2	4.8	9.6	19.8	41.1	101.0	284.2	943.7
2004	23.4	23.5	3.1	1.8	2.8	4.4	8.6	17.9	41.3	94.1	285.0	937.0
2003	24.1	21.5	4.0	2.0	3.8	4.7	11.0	22.1	43.5	99.3	272.6	953.1
2002	22.1	22.8	3.5	1.7	3.2	4.4	8.6	19.6	41.4	99.5	260.0	901.9
2001	22.3	20.2	3.1	1.7	3.2	4.2	8.4	18.0	41.4	99.4	280.2	1,016.5
2000	21.4	25.5	3.8	1.5	2.5	4.5	9.0	19.6	38.5	90.4	290.9	933.8
1999	22.0	20.6	4.0	1.7	3.1	4.6	8.8	19.7	41.6	97.9	255.0	1,096.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
Non-Hispanic 4/, both sexes												
2006	88.3	29.5	3.4	2.0	5.1	8.8	19.1	41.2	73.1	154.2	478.9	1,755.2
2005	81.6	28.9	3.7	2.2	5.1	8.3	19.3	41.5	70.8	148.8	442.3	1,589.3
2004	76.2	30.0	3.9	2.2	4.9	8.7	19.1	40.0	68.3	144.0	416.3	1,485.2
2003	76.5	30.1	4.0	2.3	5.0	8.8	19.6	39.6	68.0	146.7	417.4	1,531.7
2002	74.4	29.0	4.3	2.1	4.7	8.3	19.7	38.6	66.4	142.2	410.7	1,514.9
2001	71.8	29.0	4.3	2.2	4.6	8.3	18.6	37.0	64.2	142.7	399.8	1,491.3
2000	69.3	29.6	4.0	2.2	4.7	8.1	18.4	35.8	63.3	137.5	391.9	1,462.0
1999	66.8	30.8	4.3	2.4	4.8	7.8	18.1	34.1	60.9	137.8	376.6	1,420.1
Non-Hispanic 4/, male												
2006	72.4	33.8	3.7	2.2	5.9	10.0	22.1	49.4	84.1	167.2	482.6	1,484.0
2005	67.4	34.5	4.2	2.4	5.9	9.4	23.0	50.3	81.9	158.7	444.6	1,330.8
2004	63.3	35.0	4.1	2.4	5.6	9.9	22.2	48.6	78.4	154.6	418.1	1,257.5
2003	63.0	35.0	4.2	2.6	5.8	9.7	23.1	47.9	77.2	157.9	420.3	1,278.3
2002	61.2	34.9	4.6	2.3	5.4	9.7	23.1	47.1	75.6	153.8	411.5	1,280.6
2001	58.8	32.7	4.6	2.2	5.4	9.2	21.9	44.2	71.7	155.2	402.2	1,282.3
2000	56.9	33.6	4.4	2.3	5.5	9.4	21.7	43.0	71.3	149.4	395.5	1,279.8
1999	55.2	35.1	4.7	2.4	5.6	9.0	21.7	40.5	69.1	149.6	387.8	1,234.0
Non-Hispanic 4/, female												
2006	103.5	25.0	3.0	1.8	4.3	7.7	16.3	33.4	62.8	143.2	476.5	1,881.3
2005	95.1	23.0	3.1	2.0	4.1	7.2	15.8	33.1	60.5	140.4	440.7	1,707.2
2004	88.5	24.7	3.7	2.1	4.2	7.5	16.0	31.6	58.9	135.1	415.1	1,587.0
2003	89.4	25.1	3.7	1.9	4.2	7.9	16.2	31.7	59.5	137.3	415.5	1,643.1
2002	86.8	22.7	4.0	1.9	4.1	7.0	16.4	30.4	57.9	132.5	410.1	1,615.9
2001	84.2	25.2	4.0	2.1	3.9	7.4	15.4	30.0	57.3	132.2	398.3	1,578.3
2000	81.0	25.4	3.5	2.1	3.9	6.8	15.2	28.9	55.9	127.6	389.5	1,535.7
1999	77.9	26.2	3.9	2.3	4.0	6.7	14.4	28.0	53.3	128.2	369.3	1,494.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
Non-Hispanic white, both sexes												
2006 5/	96.9	23.3	3.0	1.9	4.8	8.2	17.6	38.5	69.8	151.7	482.4	1,790.9
2005 5/	88.9	23.2	3.3	2.0	4.8	7.5	17.8	38.5	67.6	145.9	443.5	1,618.2
2004 5/	82.7	24.8	3.5	2.1	4.5	7.6	17.4	36.9	64.8	140.9	417.7	1,511.9
2003 5/	83.0	24.6	3.5	2.1	4.8	7.8	17.8	36.2	64.3	144.1	417.9	1,558.9
2002	80.5	24.7	3.8	2.0	4.3	7.4	17.8	35.3	62.7	139.1	410.5	1,540.5
2001	77.3	23.2	3.9	2.1	4.3	7.2	16.4	33.2	60.1	139.7	399.5	1,512.9
2000	74.3	24.0	3.4	2.1	4.5	7.0	16.2	32.0	59.0	133.9	390.1	1,481.3
1999	71.5	24.2	4.0	2.2	4.5	7.0	15.5	30.1	56.9	133.5	375.1	1,443.8
Non-Hispanic white, male												
2006 5/	78.2	26.8	3.3	2.0	5.9	9.9	20.8	46.9	80.0	163.8	482.2	1,512.3
2005 5/	72.3	27.7	3.8	2.2	5.9	8.9	22.0	47.5	78.1	154.6	442.1	1,354.9
2004 5/	67.4	29.6	3.7	2.3	5.4	9.0	21.1	45.6	74.3	150.0	415.3	1,273.4
2003 5/	66.9	30.4	3.6	2.5	6.0	9.0	21.5	44.3	72.6	153.3	416.3	1,300.3
2002	64.8	31.4	4.0	2.2	5.2	8.9	21.8	43.5	71.1	148.8	408.2	1,293.4
2001	61.9	25.3	4.4	2.2	5.3	8.2	19.8	40.0	66.9	150.8	399.4	1,293.1
2000	59.6	26.7	3.8	2.3	5.4	8.5	19.4	39.2	66.0	144.1	390.9	1,288.3
1999	57.6	27.4	4.5	2.2	5.4	8.4	19.4	36.2	64.5	144.2	382.8	1,239.8
Non-Hispanic white, female												
2006 5/	114.9	19.5	2.7	1.7	3.7	6.5	14.4	30.2	60.1	141.3	482.5	1,920.7
2005 5/	105.0	18.4	2.8	1.7	3.7	6.1	13.7	29.6	57.7	138.4	444.5	1,738.7
2004 5/	97.5	19.8	3.3	1.8	3.6	6.1	13.8	28.4	55.8	133.1	419.4	1,618.7
2003 5/	98.4	18.6	3.5	1.7	3.6	6.5	14.0	28.3	56.3	136.2	419.1	1,672.7
2002	95.5	17.7	3.6	1.8	3.4	5.8	13.9	27.2	54.8	130.8	412.1	1,647.2
2001	92.1	21.0	3.4	2.1	3.3	6.2	13.1	26.4	53.7	130.2	399.5	1,604.4
2000	88.3	21.2	2.9	1.8	3.5	5.5	12.9	25.0	52.5	125.2	389.6	1,559.1
1999	84.9	20.9	3.5	2.2	3.5	5.6	11.7	24.1	49.8	124.5	370.2	1,525.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
Non-Hispanic black, both sexes												
2006 5/	68.7	53.8	4.6	2.6	6.7	14.1	31.1	68.4	116.8	209.8	546.6	1,660.0
2005 5/	65.9	51.0	4.9	3.4	6.7	13.8	31.8	69.9	113.4	204.5	523.9	1,525.4
2004 5/	63.1	50.5	5.4	2.8	7.0	16.0	31.9	69.4	113.2	204.2	489.7	1,425.6
2003 5/	63.0	52.6	5.5	2.8	6.7	15.7	34.0	71.3	113.6	200.3	487.6	1,451.6
2002	61.7	46.1	6.1	2.7	7.0	15.0	34.7	69.9	110.2	197.3	482.8	1,443.4
2001	61.1	53.8	5.6	2.5	6.7	14.7	34.8	71.7	109.1	198.5	476.3	1,431.5
2000	60.1	51.3	6.1	2.9	6.6	15.6	35.1	70.6	109.6	195.0	472.0	1,403.3
1999	58.0	58.4	5.5	3.1	6.9	13.5	36.2	69.3	104.9	202.1	444.8	1,295.3
Non-Hispanic black, male												
2006 5/	61.5	61.0	5.0	3.0	6.8	13.4	33.3	78.7	140.9	243.3	603.7	1,437.6
2005 5/	58.9	63.8	5.8	3.7	7.0	13.9	33.4	80.6	135.0	232.6	573.0	1,261.7
2004 5/	58.1	56.9	5.8	2.7	7.1	16.8	33.1	81.5	135.3	238.7	548.8	1,316.9
2003 5/	57.6	56.8	6.3	3.0	6.0	15.6	36.7	84.1	134.7	237.5	552.6	1,243.4
2002	56.6	50.6	6.6	2.9	6.9	15.8	36.2	83.4	131.4	232.9	526.1	1,331.3
2001	56.4	61.3	5.2	2.5	6.8	15.6	39.3	85.6	128.3	232.9	530.2	1,329.4
2000	56.1	61.4	6.4	2.6	7.1	16.3	39.6	82.4	133.0	231.8	527.9	1,340.9
1999	54.4	66.5	5.6	3.6	7.3	13.9	40.3	81.9	122.9	228.2	509.7	1,303.8
Non-Hispanic black, female												
2006 5/	75.2	46.3	4.1	2.2	6.6	14.7	29.2	59.6	97.4	186.0	514.2	1,750.7
2005 5/	72.3	37.6	4.0	3.1	6.3	13.8	30.4	60.7	96.1	184.5	496.1	1,631.5
2004 5/	67.6	43.8	5.0	3.0	6.9	15.2	30.8	59.0	95.4	179.6	456.3	1,468.5
2003 5/	67.9	48.4	4.6	2.6	7.3	15.7	31.7	60.3	96.6	173.9	450.9	1,532.3
2002	66.3	41.3	5.6	2.4	7.2	14.3	33.3	58.3	93.2	172.0	458.3	1,486.2
2001	65.2	46.0	6.0	2.6	6.6	13.9	30.8	59.7	93.6	174.3	446.1	1,469.3
2000	63.7	40.9	5.8	3.2	6.2	15.0	31.0	60.5	90.8	169.1	440.6	1,426.5
1999	61.3	50.0	5.4	2.6	6.5	13.1	32.7	58.5	90.5	183.7	408.5	1,292.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
All origins 3/, both sexes												
2006	40.6	27.8	9.9	5.6	38.2	37.0	40.2	45.5	36.2	44.5	105.1	274.9
2005	39.7	26.4	10.3	6.0	37.4	34.9	38.6	43.2	35.8	46.3	106.1	279.5
2004	38.1	25.8	10.3	6.5	37.0	32.6	37.3	40.7	33.2	44.0	103.7	276.7
2003	37.6	23.6	10.9	6.4	37.1	31.5	37.8	38.8	32.9	44.1	101.9	278.9
2002	37.0	23.5	10.5	6.6	38.0	31.5	37.2	36.6	31.4	44.2	101.3	275.4
2001	35.7	24.2	11.2	6.9	36.1	29.9	35.4	34.1	30.3	42.8	100.9	276.4
2000	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9	41.9	95.1	273.5
1999	35.1	22.3	12.4	7.6	35.3	29.6	33.8	31.8	30.6	44.6	100.5	282.4
All origins 3/, male												
2006	53.5	31.6	11.1	6.7	56.0	55.8	56.9	65.0	51.6	60.1	135.8	337.1
2005	52.3	27.5	12.4	7.2	54.6	52.2	54.8	62.2	51.3	62.6	139.0	344.2
2004	49.8	29.0	11.8	8.0	53.3	48.5	53.0	58.6	46.9	59.2	134.7	345.5
2003	49.3	26.4	13.0	7.9	53.3	47.1	54.2	55.9	47.1	58.9	132.9	352.5
2002	48.9	26.8	12.9	8.2	54.9	47.7	53.7	53.3	45.2	60.6	131.6	346.6
2001	47.2	26.2	13.3	8.6	52.9	45.4	51.8	50.6	43.0	57.3	133.4	358.5
2000	46.2	25.7	14.0	8.8	52.1	45.0	51.0	49.0	44.1	55.9	124.3	359.7
1999	46.4	25.4	14.4	9.4	50.7	45.0	50.2	47.9	44.0	60.2	132.4	375.4
All origins 3/, female												
2006	28.1	23.8	8.6	4.4	19.4	17.6	23.4	26.6	22.0	31.3	84.1	245.8
2005	27.5	25.2	8.0	4.7	19.2	16.9	22.3	24.7	21.3	32.5	83.8	249.8
2004	26.8	22.5	8.7	5.1	19.9	16.0	21.7	23.4	20.5	31.1	82.9	245.8
2003	26.2	20.7	8.7	4.8	19.9	15.3	21.4	22.4	19.7	31.6	81.1	246.4
2002	25.6	20.0	8.1	5.0	20.1	14.9	20.8	20.5	18.6	30.6	81.2	244.6
2001	24.5	22.1	8.9	5.2	18.4	14.0	19.2	18.0	18.5	30.8	79.6	242.0
2000	23.8	20.5	9.7	5.6	19.1	13.7	17.5	16.8	18.8	30.3	76.1	238.4
1999	24.1	19.0	10.3	5.6	19.2	14.0	17.6	16.3	18.3	31.8	79.7	244.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Hispanic, both sexes												
2006	27.2	19.5	9.7	5.1	36.1	29.7	31.6	35.6	33.2	37.6	74.9	142.8
2005	26.9	18.5	10.6	4.7	37.1	29.4	30.0	34.1	30.3	39.4	79.7	153.6
2004	25.2	17.5	9.0	5.1	33.4	27.4	29.6	31.0	30.5	37.2	74.2	160.8
2003	26.1	15.1	9.9	5.8	33.8	29.4	31.7	32.8	30.6	38.1	70.9	151.8
2002	26.1	15.2	10.9	5.4	35.5	27.6	31.7	32.0	28.7	39.5	77.4	153.0
2001	25.8	18.8	10.3	5.4	32.6	28.2	32.3	31.7	30.5	40.0	73.8	166.3
2000	25.0	15.3	10.4	5.6	31.8	27.3	31.5	30.4	28.8	38.8	72.1	184.5
1999	25.5	17.7	11.9	5.6	30.8	28.4	32.6	31.2	28.5	42.5	69.6	182.4
Hispanic, male												
2006	39.7	20.8	11.3	6.3	55.5	44.9	48.0	54.0	51.2	54.2	95.8	174.8
2005	39.0	18.3	13.3	5.6	55.8	44.9	44.8	52.9	46.6	58.4	106.2	168.1
2004	36.5	18.8	10.5	6.2	50.8	41.6	44.9	47.9	44.8	52.0	99.4	193.3
2003	38.3	17.4	11.9	7.5	50.7	45.5	48.0	50.6	45.3	53.3	102.4	195.5
2002	38.5	19.5	13.5	6.6	53.5	43.0	48.5	48.7	43.5	58.7	105.9	224.2
2001	37.6	18.0	12.7	6.2	49.2	43.8	50.1	48.8	45.0	54.9	105.1	198.1
2000	36.9	17.0	11.6	6.1	48.2	42.9	49.6	48.4	43.2	57.2	93.9	251.9
1999	37.7	20.7	13.5	6.8	46.2	44.1	51.3	50.2	43.5	63.3	94.9	258.8
Hispanic, female												
2006	13.8	18.1	8.0	3.9	14.2	11.0	13.1	16.7	16.7	24.1	60.2	125.8
2005	13.8	18.6	7.7	3.9	15.9	10.5	13.5	14.8	15.3	23.9	61.2	146.0
2004	13.1	16.2	7.5	4.0	13.5	10.2	12.6	14.0	17.5	25.4	56.8	144.0
2003	13.1	12.7	7.8	3.9	14.3	10.1	13.8	15.1	17.3	26.0	49.2	129.7
2002	12.8	10.8	8.2	4.1	14.5	9.5	13.4	15.3	15.3	24.2	57.9	117.7
2001	13.2	19.7	7.8	4.5	13.3	10.1	13.0	14.9	17.7	28.2	52.6	150.8
2000	12.4	13.5	9.2	5.2	12.5	9.2	12.2	12.8	16.0	24.4	57.4	151.3
1999	12.6	14.6	10.3	4.2	12.8	10.1	12.6	12.8	15.2	26.2	52.7	145.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Non-Hispanic 4/, both sexes												
2006	42.8	30.2	9.8	5.7	38.6	38.7	41.5	46.4	36.4	44.9	106.7	281.3
2005	41.8	28.5	10.2	6.3	37.4	36.0	39.9	44.0	36.1	46.7	107.5	285.1
2004	40.1	28.1	10.6	6.8	37.7	33.6	38.5	41.5	33.3	44.3	105.2	281.5
2003	39.2	25.9	11.1	6.5	37.6	31.8	38.5	39.2	32.9	44.3	103.2	283.7
2002	38.5	25.5	10.4	6.9	38.3	32.2	37.8	36.8	31.4	44.4	102.1	279.8
2001	36.9	25.4	11.3	7.2	36.6	30.1	35.6	34.0	30.0	42.8	101.9	279.8
2000	36.0	25.0	12.2	7.5	36.7	29.8	34.2	32.5	30.9	41.9	95.8	276.1
1999	36.2	23.4	12.4	7.9	36.0	29.6	33.7	31.6	30.5	44.5	101.3	285.0
Non-Hispanic 4/, male												
2006	55.8	34.7	10.9	6.8	55.9	58.6	58.3	66.0	51.3	60.4	137.8	345.9
2005	54.5	30.0	12.1	7.6	54.3	53.9	56.4	62.9	51.5	62.7	140.7	352.9
2004	51.9	31.8	12.1	8.4	53.7	50.1	54.2	59.4	46.8	59.5	136.4	352.5
2003	50.9	28.7	13.2	8.0	53.7	47.3	54.9	56.0	47.0	59.0	134.1	358.9
2002	50.3	28.6	12.7	8.4	55.0	48.6	54.2	53.4	44.9	60.3	132.3	351.2
2001	48.5	28.0	13.4	9.0	53.5	45.6	51.7	50.4	42.5	57.1	134.3	364.5
2000	47.3	27.7	14.5	9.4	52.7	45.2	50.8	48.6	43.8	55.5	125.2	363.1
1999	47.4	26.6	14.5	9.9	51.4	44.8	49.7	47.3	43.5	59.7	133.2	378.7
Non-Hispanic 4/, female												
2006	30.4	25.5	8.7	4.5	20.4	19.0	25.0	27.5	22.4	31.8	85.4	251.3
2005	29.6	26.9	8.1	4.9	19.8	18.2	23.6	25.7	21.7	33.1	85.0	254.1
2004	28.8	24.2	9.0	5.3	21.0	17.2	23.1	24.3	20.7	31.5	84.2	249.8
2003	28.1	22.9	8.9	5.0	20.9	16.3	22.4	23.0	19.8	32.0	82.6	250.6
2002	27.3	22.3	8.1	5.2	21.1	15.9	21.8	20.9	18.8	31.0	82.2	249.0
2001	26.0	22.6	9.2	5.3	19.3	14.7	19.9	18.3	18.5	30.8	80.6	244.6
2000	25.2	22.2	9.7	5.6	20.3	14.5	18.1	17.0	18.9	30.6	76.7	240.9
1999	25.5	20.0	10.2	5.9	20.2	14.6	18.0	16.5	18.5	32.0	80.6	247.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Non-Hispanic white, both sexes												
2006 5/	45.7	24.3	8.9	5.3	42.5	41.8	43.6	46.5	35.4	45.1	110.2	295.9
2005 5/	44.4	24.5	9.8	5.9	40.7	38.6	41.7	43.9	35.2	46.5	109.4	297.2
2004 5/	42.8	24.1	10.2	6.4	41.5	35.7	40.3	41.5	32.5	44.3	107.9	295.2
2003 5/	41.6	21.6	10.7	6.2	41.1	33.5	39.8	38.7	32.2	43.9	105.5	296.5
2002	40.6	19.8	9.4	6.3	41.6	33.7	38.7	36.3	30.6	43.9	104.4	290.9
2001	38.5	20.9	10.9	6.6	39.0	31.1	35.9	32.8	29.0	42.4	103.7	289.3
2000	37.3	20.1	11.1	7.2	39.2	30.0	34.2	31.0	30.0	41.0	97.2	285.6
1999	37.2	19.0	11.1	7.4	38.1	29.9	33.5	30.0	28.9	43.3	101.7	293.3
Non-Hispanic white, male												
2006 5/	58.6	27.6	9.8	6.2	61.3	62.5	60.4	64.8	49.1	59.9	141.2	362.3
2005 5/	57.0	25.6	11.8	7.1	58.9	57.1	58.4	61.9	49.1	61.6	141.9	366.3
2004 5/	54.6	28.3	11.8	7.9	58.9	52.5	56.2	58.7	44.8	59.0	138.8	366.8
2003 5/	53.1	24.1	12.9	7.5	58.2	49.5	56.3	54.7	45.4	58.2	135.7	372.9
2002	52.0	23.3	11.1	7.7	59.2	50.3	55.0	52.1	43.0	58.8	133.5	361.6
2001	49.7	22.9	13.2	8.1	56.6	46.7	51.8	48.0	40.3	56.1	135.2	374.7
2000	48.1	22.9	13.4	8.9	55.9	45.2	50.4	45.8	41.6	53.6	125.2	373.8
1999	47.8	22.0	12.9	9.1	53.6	44.9	49.0	44.2	40.7	57.3	132.7	387.1
Non-Hispanic white, female												
2006 5/	33.3	20.8	8.0	4.3	22.7	20.9	26.7	28.3	22.4	32.2	88.7	264.9
2005 5/	32.3	23.4	7.6	4.6	21.6	19.8	25.0	26.1	22.0	33.4	87.1	265.6
2004 5/	31.4	19.7	8.5	4.9	23.2	18.6	24.2	24.6	20.9	31.6	86.8	263.1
2003 5/	30.5	19.0	8.4	4.8	23.2	17.2	23.2	22.9	19.6	31.7	85.1	262.8
2002	29.6	16.0	7.5	4.8	23.2	16.8	22.4	20.6	18.8	31.2	84.9	260.3
2001	27.8	18.7	8.5	5.0	20.7	15.2	19.9	17.9	18.3	30.7	82.8	253.8
2000	26.9	17.1	8.7	5.3	21.9	14.6	18.0	16.3	19.1	30.4	78.8	250.0
1999	27.0	15.9	9.3	5.6	22.0	14.7	17.9	15.9	17.7	31.5	81.5	255.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Non-Hispanic black, both sexes												
2006 5/	36.2	54.8	14.2	7.4	26.8	35.6	42.4	56.2	49.5	48.7	82.0	153.2
2005 5/	35.9	48.6	12.6	7.9	27.7	33.1	39.6	54.0	48.5	53.6	97.8	176.7
2004 5/	33.7	47.0	12.8	8.8	26.3	32.5	37.8	50.7	44.5	48.5	88.2	155.1
2003 5/	33.3	44.7	13.7	8.0	26.4	30.9	39.7	49.3	42.9	51.1	83.1	161.8
2002	34.0	50.1	14.6	9.4	28.4	32.5	39.9	47.6	41.9	49.7	84.6	175.4
2001	34.4	44.5	14.2	9.7	29.6	31.1	40.9	48.5	42.4	51.3	89.6	183.9
2000	34.4	46.5	16.8	9.6	29.6	33.5	40.6	47.8	41.3	53.0	85.3	183.2
1999	36.1	44.1	18.2	10.9	30.5	33.2	40.7	48.3	45.1	58.6	103.8	208.9
Non-Hispanic black, male												
2006 5/	52.5	64.1	16.3	9.1	40.2	57.6	64.1	88.1	79.8	73.9	117.3	202.5
2005 5/	51.3	52.3	14.1	9.8	41.2	52.4	59.3	83.8	80.9	82.3	145.2	227.4
2004 5/	47.9	49.2	14.4	10.6	38.5	52.0	55.6	77.9	71.9	74.1	133.2	218.5
2003 5/	47.4	49.0	15.9	10.1	39.8	48.4	59.4	75.8	69.0	74.9	123.8	228.0
2002	49.0	52.1	18.9	11.9	42.6	52.2	59.7	73.3	69.2	77.2	131.3	262.5
2001	49.5	50.4	15.8	13.1	44.5	48.8	61.3	78.6	68.2	74.9	138.3	256.4
2000	50.3	49.5	19.4	12.3	44.2	53.5	63.3	76.7	69.4	80.4	133.3	277.1
1999	52.3	48.5	21.8	14.3	46.4	53.0	63.0	79.4	71.8	89.6	151.0	318.5
Non-Hispanic black, female												
2006 5/	21.5	45.0	12.0	5.6	13.1	15.2	23.2	28.9	25.1	30.7	62.0	133.0
2005 5/	21.8	44.8	11.1	6.0	14.0	15.3	22.1	28.5	22.5	33.2	70.9	156.3
2004 5/	20.8	44.8	11.1	6.8	13.9	14.7	21.9	27.5	22.4	30.3	62.8	130.1
2003 5/	20.5	40.3	11.4	5.9	12.9	14.9	22.4	26.6	22.0	34.1	60.1	136.2
2002	20.3	48.0	10.1	6.8	14.2	14.7	22.5	25.6	20.0	30.2	58.2	142.1
2001	20.6	38.3	12.4	6.2	14.8	15.1	22.7	22.7	21.6	34.6	62.2	157.0
2000	20.1	43.3	14.2	6.7	15.1	15.5	20.5	23.0	18.7	33.7	58.4	148.3
1999	21.4	39.6	14.5	7.4	14.8	15.4	21.1	21.8	23.7	36.8	77.4	167.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
All origins 3/, both sexes												
2006	16.2	3.4	3.8	3.6	26.8	19.2	16.6	16.8	15.8	16.8	23.6	24.4
2005	16.3	3.6	4.0	3.8	26.8	19.0	16.6	16.7	16.2	18.0	24.3	26.1
2004	16.4	3.6	4.1	4.3	27.3	18.8	16.4	16.7	15.5	17.5	25.5	27.2
2003	16.5	3.7	4.1	4.3	27.7	18.3	17.1	16.6	15.9	17.5	26.2	29.7
2002	16.8	3.1	4.1	4.2	29.1	18.7	17.1	16.4	15.7	18.3	27.0	28.8
2001	16.6	3.7	4.4	4.4	27.9	18.8	17.4	16.0	15.3	17.9	27.9	28.9
2000	16.6	4.6	4.5	4.7	28.1	18.6	17.0	15.8	16.0	18.0	27.0	31.5
1999	16.6	5.0	4.6	4.8	27.6	18.5	16.8	15.4	16.0	19.4	28.4	32.0
All origins 3/, male												
2006	23.1	3.3	4.0	4.1	37.9	28.9	23.9	25.1	22.9	23.4	33.2	42.9
2005	23.3	3.5	4.5	4.4	37.9	28.6	24.2	24.9	23.5	25.3	34.8	46.0
2004	23.0	3.9	4.4	5.0	37.8	27.7	24.0	24.7	22.2	23.7	36.5	46.5
2003	23.3	4.0	4.5	5.2	38.6	27.3	24.8	24.3	23.0	23.7	36.6	52.0
2002	23.7	3.5	4.8	4.9	40.7	28.1	24.6	24.0	22.5	25.3	37.7	53.0
2001	23.5	3.3	5.0	5.3	39.6	28.3	25.1	23.3	21.6	24.3	39.8	52.7
2000	23.3	4.8	5.0	5.4	39.3	27.7	24.8	23.0	22.5	24.3	38.1	59.8
1999	23.1	5.2	5.1	5.8	38.1	27.2	24.0	22.4	22.5	26.1	40.7	63.5
All origins 3/, female												
2006	9.4	3.6	3.6	3.1	15.1	9.1	9.2	8.7	9.2	11.2	17.1	15.7
2005	9.5	3.6	3.5	3.2	15.1	9.1	9.1	8.8	9.4	11.8	17.1	17.0
2004	9.9	3.2	3.8	3.6	16.1	9.5	8.8	9.0	9.3	12.3	18.1	18.5
2003	10.0	3.4	3.7	3.4	16.3	9.0	9.4	9.1	9.4	12.4	19.3	19.9
2002	10.1	2.8	3.4	3.5	16.9	9.2	9.7	9.0	9.3	12.5	19.8	18.3
2001	10.0	4.2	3.8	3.5	15.6	9.2	9.8	8.9	9.4	12.7	20.1	18.9
2000	10.2	4.3	4.0	3.9	16.4	9.3	9.3	8.9	10.0	12.8	19.8	19.9
1999	10.4	4.8	4.2	3.9	16.6	9.6	9.6	8.7	10.1	14.0	20.4	19.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Hispanic, both sexes												
2006	14.8	4.6	4.7	3.6	27.8	17.8	15.3	15.7	15.4	17.4	20.7	11.0
2005	14.9	4.7	5.3	3.6	28.4	18.3	14.9	14.9	14.3	16.3	24.1	15.4
2004	14.5	4.4	4.7	3.9	26.3	18.1	14.9	14.3	15.9	17.0	21.9	20.0
2003	15.2	4.5	4.2	4.1	26.2	19.1	16.4	15.1	15.8	18.6	23.6	24.9
2002	15.3	3.1	5.8	3.8	27.6	18.3	16.0	15.2	15.2	18.4	26.2	25.5
2001	15.6	5.3	5.3	3.8	26.0	18.8	17.3	16.4	17.9	20.2	30.1	23.8
2000	14.8	5.4	5.2	4.1	24.6	17.3	16.2	16.2	15.7	20.4	23.9	25.2
1999	14.4	5.3	5.5	3.8	23.1	17.8	15.4	15.0	15.5	22.0	25.0	25.3
Hispanic, male												
2006	21.7	4.6	5.3	4.3	42.0	26.5	22.4	23.0	22.7	24.2	29.4	*
2005	21.7	4.4	6.5	4.2	41.5	27.4	21.2	22.2	20.9	25.7	35.4	*
2004	21.3	*	5.4	4.6	39.3	27.0	22.2	21.6	21.6	23.9	34.7	28.0
2003	22.2	5.7	4.6	5.5	38.2	29.3	23.9	22.4	21.4	26.2	35.3	46.2
2002	22.4	*	7.1	4.6	40.4	27.5	23.9	21.6	22.1	28.5	37.3	45.2
2001	22.8	*	6.0	4.3	38.4	28.8	25.9	23.2	24.6	28.2	51.0	43.8
2000	21.6	6.1	5.6	4.5	36.1	26.4	24.1	24.1	23.5	29.5	36.4	44.3
1999	20.8	5.7	5.8	4.5	33.3	26.7	23.2	22.6	21.0	31.6	40.1	*
Hispanic, female												
2006	7.4	4.6	4.0	2.9	11.9	7.1	7.4	8.1	8.8	11.9	14.6	*
2005	7.6	5.0	4.1	3.1	13.6	7.2	7.7	7.5	8.3	8.6	16.3	*
2004	7.3	5.0	4.1	3.2	11.3	7.5	6.7	6.9	10.7	11.5	13.1	15.8
2003	7.6	*	3.8	2.7	12.3	7.0	8.3	7.9	10.7	12.5	15.5	*
2002	7.7	*	4.5	2.9	12.6	7.3	7.5	8.7	8.9	10.3	18.5	*
2001	8.0	6.7	4.6	3.2	11.6	7.1	8.0	9.7	12.0	13.9	16.0	*
2000	7.6	*	4.8	3.7	11.0	6.8	7.8	8.5	8.7	13.3	15.5	*
1999	7.6	*	5.2	3.1	11.0	7.6	7.0	7.7	10.6	14.5	14.9	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Non-Hispanic 4/, both sexes												
2006	16.4	3.1	3.5	3.6	26.6	19.5	16.8	16.8	15.8	16.7	23.7	25.0
2005	16.5	3.2	3.6	3.9	26.4	19.1	16.9	16.9	16.3	18.1	24.2	26.5
2004	16.6	3.3	3.9	4.4	27.4	18.9	16.6	16.9	15.4	17.5	25.6	27.5
2003	16.7	3.5	4.0	4.3	28.0	18.0	17.2	16.6	15.8	17.3	26.3	29.8
2002	16.9	3.2	3.7	4.3	29.3	18.8	17.2	16.4	15.6	18.3	26.9	28.9
2001	16.7	3.4	4.2	4.6	28.2	18.8	17.3	15.8	15.0	17.7	27.7	29.0
2000	16.8	4.3	4.3	4.8	28.8	18.7	17.0	15.7	15.9	17.8	27.1	31.7
1999	16.9	4.9	4.4	5.0	28.4	18.5	16.9	15.4	16.0	19.2	28.3	32.2
Non-Hispanic 4/, male												
2006	23.3	2.9	3.5	4.1	36.9	29.5	24.1	25.3	22.8	23.2	33.3	44.2
2005	23.6	3.3	3.9	4.4	37.1	28.8	24.6	25.1	23.6	25.2	34.6	47.1
2004	23.3	3.9	4.1	5.0	37.5	27.8	24.3	24.9	22.2	23.6	36.5	47.3
2003	23.4	3.5	4.5	5.1	38.5	26.7	24.9	24.4	23.0	23.3	36.6	52.0
2002	23.8	3.4	4.2	5.0	40.6	28.1	24.6	24.1	22.4	24.9	37.5	53.2
2001	23.5	3.2	4.7	5.5	39.7	28.1	24.8	23.2	21.3	24.0	39.2	52.9
2000	23.4	4.5	4.8	5.6	39.8	27.8	24.7	22.8	22.2	23.9	38.0	60.4
1999	23.3	5.1	4.9	6.0	39.0	27.2	24.0	22.2	22.4	25.6	40.4	64.3
Non-Hispanic 4/, female												
2006	9.8	3.2	3.5	3.1	15.7	9.5	9.5	8.8	9.2	11.1	17.2	16.1
2005	9.8	3.2	3.3	3.3	15.3	9.5	9.3	8.9	9.5	12.1	17.2	17.1
2004	10.3	2.6	3.8	3.7	17.0	9.9	9.1	9.2	9.1	12.3	18.4	18.6
2003	10.3	3.5	3.6	3.6	17.0	9.4	9.6	9.2	9.2	12.3	19.5	20.1
2002	10.4	2.9	3.1	3.6	17.7	9.5	10.0	9.0	9.3	12.7	19.8	18.4
2001	10.2	3.6	3.6	3.6	16.3	9.5	10.0	8.8	9.1	12.5	20.2	19.1
2000	10.5	4.2	3.8	3.9	17.4	9.7	9.4	8.9	10.1	12.8	20.0	20.1
1999	10.7	4.7	3.9	4.0	17.5	10.0	9.9	8.8	10.0	13.9	20.5	19.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Non-Hispanic white, both sexes												
2006 5/	16.9	2.6	3.0	3.5	28.8	19.8	17.2	17.0	15.5	16.9	24.0	26.0
2005 5/	17.1	2.5	3.3	3.8	28.4	19.6	17.3	17.0	16.4	18.2	24.6	27.5
2004 5/	17.2	2.9	3.7	4.4	29.9	19.1	17.0	16.8	15.4	17.6	26.0	28.8
2003 5/	17.2	3.2	3.8	4.4	30.2	18.2	17.3	16.6	15.6	17.3	26.6	30.6
2002	17.5	2.5	3.3	4.3	31.8	19.0	17.3	16.4	15.5	18.3	27.4	29.7
2001	17.0	3.1	3.8	4.4	29.9	19.0	17.4	15.6	14.7	17.6	28.3	29.9
2000	17.1	3.5	3.8	4.8	30.9	18.5	17.0	15.4	15.8	17.6	27.6	32.4
1999	17.1	4.3	3.7	5.0	30.3	18.6	16.7	15.0	15.6	19.1	28.6	32.5
Non-Hispanic white, male												
2006 5/	23.8	2.6	2.9	3.9	39.7	29.6	24.5	25.2	22.4	23.2	33.5	45.4
2005 5/	24.2	2.8	3.5	4.4	39.6	29.0	24.9	25.3	23.5	25.0	34.8	48.7
2004 5/	23.9	3.3	3.8	5.2	40.6	27.6	24.6	24.5	21.8	23.5	36.6	49.6
2003 5/	23.9	3.3	4.4	5.1	41.1	26.8	24.9	24.1	22.6	23.3	36.6	52.8
2002	24.3	2.9	3.6	5.0	43.6	28.1	24.6	24.0	22.0	24.5	37.5	54.0
2001	23.8	2.9	4.4	5.2	41.9	28.2	24.6	22.7	20.8	23.7	39.6	54.3
2000	23.7	3.8	4.3	5.7	42.5	27.1	24.5	22.2	21.6	23.2	38.1	62.3
1999	23.5	4.9	4.0	5.9	41.2	26.9	23.6	21.6	21.7	25.2	40.3	64.6
Non-Hispanic white, female												
2006 5/	10.2	2.6	3.2	3.1	17.2	9.8	9.9	8.9	9.0	11.3	17.5	17.0
2005 5/	10.3	2.3	3.2	3.2	16.6	10.1	9.7	8.9	9.6	12.3	17.5	17.8
2004 5/	10.7	2.4	3.6	3.6	18.6	10.4	9.3	9.2	9.3	12.5	18.8	19.5
2003 5/	10.7	3.0	3.2	3.7	18.8	9.5	9.7	9.2	8.9	12.1	19.8	20.8
2002	10.9	2.1	2.9	3.5	19.4	9.8	10.1	8.9	9.4	13.0	20.7	19.2
2001	10.6	3.3	3.2	3.5	17.4	9.7	10.2	8.6	9.1	12.5	20.8	19.8
2000	10.9	3.2	3.4	4.0	18.9	9.7	9.5	8.6	10.4	12.8	20.7	20.4
1999	11.0	3.6	3.4	4.0	19.1	10.2	9.9	8.6	9.9	13.9	20.9	19.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Non-Hispanic black, both sexes												
2006 5/	15.6	4.9	5.2	4.0	19.6	22.4	17.9	18.2	19.1	15.6	20.1	15.9
2005 5/	15.4	5.7	4.7	4.1	20.7	20.4	17.8	18.2	17.3	18.3	20.0	16.9
2004 5/	15.6	5.0	5.2	4.5	20.0	21.6	17.4	19.9	17.6	16.4	21.7	15.0
2003 5/	15.6	4.4	5.1	4.3	20.3	20.0	18.8	18.6	18.7	18.2	22.6	21.4
2002	15.7	5.6	5.0	4.7	21.6	21.1	18.5	18.0	17.3	17.3	21.8	18.8
2001	16.3	*	5.4	5.5	22.8	20.5	19.2	19.1	18.4	18.8	21.4	17.8
2000	16.4	6.9	5.9	4.8	22.4	22.2	19.1	19.7	17.5	19.9	19.9	21.9
1999	16.7	7.7	7.1	5.6	22.7	20.6	19.5	19.2	18.5	19.5	26.0	27.8
Non-Hispanic black, male												
2006 5/	23.6	*	5.8	4.9	28.4	36.4	28.3	29.3	30.0	25.6	31.7	37.6
2005 5/	23.4	*	5.1	4.8	30.2	33.5	28.5	28.4	28.0	30.2	34.2	34.3
2004 5/	23.4	6.2	5.4	4.9	28.6	35.1	27.3	31.5	28.2	26.4	38.5	29.5
2003 5/	23.3	*	5.1	5.3	30.3	31.8	29.1	29.7	29.6	26.5	38.1	43.7
2002	23.6	*	6.5	5.4	31.0	34.1	28.0	28.0	28.6	28.7	40.7	47.0
2001	24.5	*	5.8	7.1	33.4	32.6	29.7	30.7	30.2	27.4	36.9	35.8
2000	24.8	7.2	6.4	5.8	32.5	35.6	30.5	30.1	29.6	32.6	36.4	45.8
1999	24.9	*	8.0	7.0	32.9	33.0	29.8	30.7	29.3	29.5	43.5	68.9
Non-Hispanic black, female												
2006 5/	8.2	*	4.6	3.1	10.5	9.4	8.7	8.7	10.3	8.5	13.6	*
2005 5/	8.1	6.5	4.3	3.4	11.1	8.2	8.3	9.4	8.8	9.9	11.9	10.0
2004 5/	8.5	*	5.0	4.0	11.3	9.3	8.7	9.9	9.2	9.2	12.2	9.2
2003 5/	8.5	*	5.2	3.3	10.1	9.3	9.7	9.0	9.9	12.3	13.9	12.8
2002	8.5	*	3.6	3.9	12.1	9.3	10.1	9.4	8.2	9.1	11.1	*
2001	8.9	*	5.0	3.8	12.3	9.6	9.9	9.2	9.0	12.8	12.6	11.1
2000	8.8	*	5.4	3.7	12.5	10.1	8.9	10.8	7.8	11.0	10.6	13.1
1999	9.4	9.3	6.2	4.1	12.6	9.4	10.4	9.5	9.8	12.5	16.3	12.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
All origins 3/, both sexes												
2006	15.1	3.4	3.6	3.3	26.0	18.2	15.3	15.3	14.3	15.4	22.3	23.4
2005	15.3	3.6	3.8	3.6	25.9	18.0	15.4	15.1	14.7	16.7	22.9	25.1
2004	15.3	3.5	4.0	4.1	26.3	17.6	15.1	15.1	14.1	16.1	24.5	26.1
2003	15.4	3.6	3.9	4.0	26.6	17.1	15.7	14.9	14.2	16.2	24.9	28.8
2002	15.7	3.0	3.9	3.9	28.2	17.8	15.8	14.8	14.1	17.0	25.7	28.0
2001	15.4	3.6	4.1	4.1	26.8	17.5	15.7	14.2	13.7	16.3	26.3	27.8
2000	15.4	4.4	4.2	4.3	26.9	17.3	15.3	14.2	14.4	16.5	25.7	30.4
1999	15.2	4.8	4.2	4.3	26.2	16.9	14.9	13.6	14.2	17.8	26.5	30.4
All origins 3/, male												
2006	21.4	3.3	3.8	3.7	36.6	27.4	21.8	22.6	20.5	21.1	30.8	41.0
2005	21.7	3.5	4.2	4.1	36.5	26.9	22.2	22.2	20.9	23.2	32.4	44.1
2004	21.3	3.8	4.2	4.7	36.4	25.8	21.9	21.9	19.8	21.3	34.7	44.2
2003	21.4	3.9	4.4	4.8	36.9	25.5	22.5	21.5	20.0	21.3	34.3	50.0
2002	21.9	3.3	4.5	4.6	39.3	26.5	22.3	21.3	19.9	22.7	35.3	51.7
2001	21.5	3.2	4.5	4.9	38.0	26.2	22.4	20.5	19.0	21.5	37.3	51.1
2000	21.3	4.6	4.7	5.0	37.4	25.5	22.0	20.4	19.8	21.7	35.6	57.5
1999	20.9	5.0	4.7	5.1	36.0	24.7	21.1	19.4	19.3	23.3	38.0	59.8
All origins 3/, female												
2006	9.0	3.5	3.4	2.9	14.7	8.7	8.8	8.2	8.5	10.6	16.5	15.2
2005	9.1	3.6	3.4	3.0	14.7	8.8	8.6	8.2	8.9	11.2	16.5	16.4
2004	9.5	3.2	3.7	3.4	15.7	9.1	8.3	8.5	8.8	11.7	17.7	17.9
2003	9.5	3.3	3.5	3.2	15.8	8.5	8.9	8.5	8.8	11.9	18.7	19.5
2002	9.8	2.8	3.3	3.3	16.6	8.8	9.3	8.6	8.9	12.3	19.3	17.7
2001	9.5	4.0	3.6	3.3	15.1	8.7	9.1	8.1	8.8	12.0	19.2	18.0
2000	9.7	4.2	3.8	3.6	15.9	8.8	8.8	8.2	9.5	12.3	19.2	19.3
1999	9.7	4.7	3.8	3.6	15.9	8.9	8.8	8.0	9.5	13.3	19.1	18.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
Hispanic, both sexes												
2006	14.2	4.6	4.6	3.4	27.2	17.2	14.3	14.6	14.7	16.5	19.4	10.6
2005	14.3	4.7	5.2	3.5	27.7	17.7	14.1	13.8	13.5	15.5	22.5	15.0
2004	13.9	4.4	4.6	3.8	25.6	17.2	13.9	13.2	14.9	16.1	21.3	19.5
2003	14.4	4.5	4.1	3.9	25.5	18.1	15.4	14.2	14.6	17.4	23.1	24.4
2002	14.6	3.1	5.6	3.6	26.6	17.6	15.2	13.9	14.3	17.9	24.4	24.9
2001	14.3	4.9	4.7	3.5	25.1	17.4	15.3	13.8	15.1	17.7	26.3	20.0
2000	13.9	5.2	4.9	3.8	23.7	16.4	14.7	15.1	14.4	18.4	22.3	22.6
1999	13.3	5.3	5.2	3.6	21.9	16.4	13.9	13.7	14.5	19.7	23.5	23.2
Hispanic, male												
2006	20.7	4.6	5.3	4.0	41.1	25.6	20.6	21.3	21.5	22.9	26.9	*
2005	20.7	4.4	6.3	4.0	40.3	26.3	20.2	20.4	19.4	24.6	32.1	*
2004	20.2	*	5.3	4.4	38.3	25.4	20.8	19.6	19.7	22.3	33.3	28.0
2003	21.1	5.7	4.3	5.2	37.2	27.6	22.1	20.8	19.8	24.7	34.1	46.2
2002	21.3	*	6.8	4.4	38.9	26.4	22.6	19.6	20.5	27.5	34.5	43.5
2001	21.0	*	5.3	4.0	36.9	26.8	23.1	20.2	21.2	24.6	44.3	*
2000	20.1	6.1	5.2	4.1	34.7	24.9	21.6	22.0	21.2	26.0	33.4	*
1999	19.1	5.7	5.6	4.1	31.6	24.4	20.8	20.1	19.0	27.2	37.0	*
Hispanic, female												
2006	7.2	4.6	3.9	2.9	11.6	6.8	7.2	7.8	8.5	11.3	14.2	*
2005	7.4	5.0	4.1	2.9	13.4	7.2	7.4	7.0	8.2	8.2	15.8	*
2004	7.1	5.0	3.9	3.2	11.1	7.3	6.4	6.7	10.5	11.2	13.1	15.1
2003	7.3	*	3.8	2.7	11.9	6.6	8.0	7.5	9.8	11.6	15.5	*
2002	7.4	*	4.4	2.7	12.4	7.2	7.3	8.2	8.7	10.2	17.4	*
2001	7.2	6.2	4.1	3.0	11.2	6.5	6.9	7.6	9.7	12.3	14.2	*
2000	7.2	*	4.6	3.6	10.6	6.5	7.3	8.3	8.4	12.4	14.9	*
1999	7.2	*	4.7	3.0	10.6	7.1	6.5	7.4	10.6	13.8	14.6	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
Non-Hispanic 4/, both sexes												
2006	15.3	3.0	3.3	3.3	25.6	18.4	15.4	15.3	14.2	15.3	22.4	24.1
2005	15.4	3.2	3.4	3.6	25.5	18.0	15.5	15.1	14.7	16.7	22.9	25.5
2004	15.5	3.2	3.8	4.1	26.4	17.6	15.2	15.2	14.0	16.1	24.6	26.3
2003	15.5	3.4	3.9	4.0	26.8	16.9	15.7	14.9	14.1	16.0	25.0	28.9
2002	15.9	3.0	3.5	4.0	28.5	17.7	15.8	14.8	14.1	16.9	25.7	28.1
2001	15.5	3.2	3.9	4.2	27.1	17.5	15.7	14.2	13.5	16.2	26.3	28.0
2000	15.6	4.2	4.1	4.4	27.5	17.4	15.4	14.1	14.4	16.3	25.8	30.6
1999	15.4	4.7	4.0	4.5	26.9	16.9	15.0	13.5	14.0	17.6	26.5	30.5
Non-Hispanic 4/, male												
2006	21.5	2.9	3.3	3.6	35.5	27.8	21.9	22.7	20.3	20.9	30.9	42.3
2005	21.8	3.3	3.6	4.1	35.6	27.0	22.5	22.3	20.9	23.0	32.3	45.2
2004	21.5	3.8	3.9	4.7	36.0	25.8	22.0	22.1	19.7	21.2	34.6	45.0
2003	21.4	3.4	4.3	4.7	36.7	24.9	22.5	21.4	19.9	20.9	34.2	49.9
2002	21.9	3.2	4.0	4.6	39.2	26.4	22.2	21.3	19.7	22.3	35.1	51.9
2001	21.5	3.1	4.3	5.1	38.1	25.9	22.2	20.4	18.8	21.3	36.9	51.6
2000	21.4	4.2	4.5	5.1	37.9	25.5	22.0	20.2	19.6	21.3	35.7	58.3
1999	21.0	4.8	4.4	5.2	36.7	24.6	21.1	19.2	19.1	22.9	37.7	60.7
Non-Hispanic 4/, female												
2006	9.3	3.1	3.3	2.9	15.3	9.1	9.0	8.2	8.5	10.5	16.6	15.6
2005	9.4	3.2	3.2	3.1	14.9	9.1	8.7	8.3	8.9	11.4	16.5	16.5
2004	9.8	2.6	3.7	3.4	16.5	9.4	8.6	8.6	8.6	11.7	17.9	18.0
2003	9.8	3.3	3.4	3.3	16.5	8.9	9.0	8.6	8.7	11.9	18.8	19.6
2002	10.1	2.9	3.0	3.4	17.4	9.1	9.5	8.6	8.9	12.4	19.4	17.8
2001	9.8	3.4	3.5	3.3	15.8	9.1	9.4	8.2	8.7	12.0	19.3	18.2
2000	10.0	4.2	3.6	3.6	16.9	9.2	8.9	8.2	9.6	12.2	19.3	19.4
1999	10.0	4.7	3.6	3.7	16.8	9.2	9.1	8.1	9.3	13.2	19.2	18.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
Non-Hispanic white, both sexes												
2006 5/	15.7	2.5	2.8	3.2	27.7	18.6	15.8	15.4	13.9	15.4	22.6	25.1
2005 5/	16.0	2.5	3.1	3.6	27.4	18.5	15.9	15.4	14.7	16.8	23.2	26.6
2004 5/	16.0	2.8	3.5	4.1	28.8	17.7	15.5	15.1	13.9	16.1	25.0	27.6
2003 5/	15.9	3.1	3.6	4.0	28.9	17.0	15.7	14.8	13.7	15.9	25.2	29.6
2002	16.3	2.4	3.0	4.0	30.8	17.8	15.8	14.8	14.0	16.9	26.2	28.9
2001	15.8	3.0	3.5	4.1	28.7	17.6	15.7	13.9	13.3	16.1	26.9	29.0
2000	15.8	3.3	3.7	4.4	29.5	17.0	15.3	13.7	14.2	16.1	26.2	31.3
1999	15.6	4.1	3.4	4.4	28.7	16.8	14.8	13.1	13.7	17.5	26.8	30.9
Non-Hispanic white, male												
2006 5/	22.0	2.6	2.7	3.4	38.1	27.8	22.2	22.6	19.8	20.8	30.9	43.6
2005 5/	22.4	2.8	3.2	4.1	38.2	27.1	22.7	22.5	20.7	22.7	32.5	46.7
2004 5/	22.0	3.3	3.6	4.8	39.1	25.5	22.2	21.7	19.3	21.0	34.8	47.1
2003 5/	21.8	3.2	4.2	4.7	39.1	24.9	22.4	21.1	19.4	20.9	34.3	50.7
2002	22.3	2.7	3.3	4.6	42.1	26.1	22.0	21.2	19.4	21.8	35.2	52.6
2001	21.7	2.9	3.9	4.8	40.1	25.9	21.9	19.9	18.2	21.0	37.3	53.1
2000	21.5	3.4	4.1	5.1	40.3	24.7	21.6	19.6	18.9	20.7	35.6	60.0
1999	21.1	4.7	3.6	5.1	38.8	24.2	20.6	18.5	18.5	22.6	37.6	61.2
Non-Hispanic white, female												
2006 5/	9.7	2.5	3.0	2.9	16.7	9.4	9.3	8.3	8.3	10.7	16.9	16.5
2005 5/	9.8	2.2	3.1	3.0	16.1	9.7	9.1	8.4	9.0	11.7	16.9	17.3
2004 5/	10.2	2.3	3.5	3.3	18.0	9.8	8.7	8.6	8.7	11.9	18.4	18.9
2003 5/	10.2	2.9	3.0	3.3	18.3	9.0	9.1	8.5	8.4	11.6	19.1	20.3
2002	10.5	2.1	2.8	3.3	19.0	9.4	9.7	8.4	8.8	12.7	20.2	18.6
2001	10.1	3.1	3.1	3.3	17.0	9.3	9.6	8.1	8.7	11.9	20.0	19.0
2000	10.3	3.2	3.2	3.6	18.4	9.3	9.0	7.9	9.8	12.3	20.0	19.8
1999	10.3	3.5	3.1	3.7	18.3	9.4	9.0	7.9	9.2	13.2	19.7	18.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
Non-Hispanic black, both sexes												
2006 5/	14.7	4.7	5.0	3.8	19.1	21.3	16.7	16.7	17.8	14.8	19.0	14.3
2005 5/	14.4	5.7	4.5	3.9	19.9	19.4	16.7	16.1	16.3	17.2	18.9	16.1
2004 5/	14.6	4.9	5.0	4.3	19.1	20.5	16.3	18.1	16.1	15.4	20.4	15.0
2003 5/	14.7	4.4	5.0	4.1	19.4	19.1	17.6	16.9	17.3	17.1	21.1	21.1
2002	14.8	5.4	4.9	4.3	21.0	20.3	17.2	16.3	15.7	16.0	20.2	18.5
2001	15.2	*	5.3	5.0	22.2	19.6	17.6	17.2	16.6	17.4	19.9	16.9
2000	15.5	6.9	5.6	4.5	21.6	21.1	17.6	18.3	16.3	18.4	18.8	20.7
1999	15.5	7.6	6.5	5.1	21.7	19.2	17.9	17.4	16.7	18.1	24.0	25.5
Non-Hispanic black, male												
2006 5/	22.2	*	5.7	4.6	27.7	34.5	26.2	26.4	27.6	24.0	30.3	33.9
2005 5/	21.8	*	4.9	4.5	28.7	32.0	26.6	24.9	26.1	27.9	32.2	33.3
2004 5/	21.7	*	5.2	4.7	27.0	33.0	25.2	28.5	25.3	24.7	36.2	29.5
2003 5/	21.7	*	5.0	4.9	28.8	30.2	27.1	26.5	26.7	24.1	35.5	42.6
2002	22.0	*	6.3	5.0	30.1	32.7	25.7	24.8	25.4	25.6	36.6	45.9
2001	22.7	*	5.6	6.5	32.4	31.0	26.9	27.5	26.8	24.2	34.8	35.8
2000	23.1	7.2	6.0	5.4	31.1	33.6	27.9	27.6	27.2	29.8	34.6	44.6
1999	22.8	*	7.0	6.2	31.5	30.7	26.9	27.5	25.8	26.8	41.3	61.8
Non-Hispanic black, female												
2006 5/	7.9	*	4.3	3.1	10.3	9.1	8.3	8.4	10.0	8.3	12.6	*
2005 5/	7.7	6.5	4.1	3.4	10.9	7.9	7.9	8.5	8.4	9.6	11.3	9.2
2004 5/	8.1	*	4.9	3.8	11.1	9.1	8.3	9.2	8.6	8.8	11.4	9.2
2003 5/	8.3	*	4.9	3.2	9.9	9.0	9.3	8.6	9.7	12.1	13.0	12.8
2002	8.3	*	3.5	3.6	11.9	9.1	9.7	9.0	8.0	9.1	10.9	*
2001	8.5	*	4.9	3.4	12.1	9.2	9.4	8.4	8.4	12.6	11.5	9.8
2000	8.5	*	5.2	3.6	12.3	9.8	8.5	10.2	7.5	10.4	9.9	11.8
1999	8.9	9.3	5.9	4.0	12.1	8.9	9.9	8.8	9.3	12.0	14.4	12.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)												
All origins 3/, both sexes												
2006	0.4	*	*	0.1	0.4	0.3	0.5	0.6	0.5	0.5	0.6	0.5
2005	0.4	*	0.2	0.1	0.5	0.4	0.5	0.6	0.5	0.4	0.6	0.5
2004	0.4	*	*	0.1	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6
2003	0.5	*	*	0.1	0.6	0.5	0.5	0.6	0.5	0.4	0.6	0.5
2002	0.4	*	*	0.2	0.4	0.3	0.5	0.5	0.4	0.4	0.5	*
2001	0.5	*	0.2	0.2	0.5	0.4	0.6	0.6	0.5	0.6	0.8	0.7
2000	0.5	*	0.2	0.2	0.6	0.5	0.7	0.6	0.5	0.5	0.7	0.8
1999	0.8	*	0.3	0.3	0.9	0.8	0.9	0.8	0.8	0.8	1.3	1.3
All origins 3/, male												
2006	0.6	*	*	0.2	0.6	0.5	0.8	0.9	0.7	0.8	1.0	*
2005	0.7	*	*	0.2	0.7	0.7	0.7	1.0	0.8	0.6	1.1	*
2004	0.6	*	*	0.2	0.7	0.7	0.8	0.9	0.7	0.7	0.9	*
2003	0.7	*	*	0.2	1.0	0.8	0.8	1.0	0.8	0.8	1.0	*
2002	0.6	*	*	0.2	0.6	0.5	0.8	0.9	0.7	0.7	1.0	*
2001	0.8	*	0.3	0.2	0.8	0.8	1.0	1.0	0.9	1.0	1.1	*
2000	0.8	*	*	0.3	1.0	0.8	1.1	0.9	0.8	0.9	1.2	*
1999	1.1	*	0.3	0.4	1.4	1.2	1.4	1.3	1.2	1.2	1.8	2.9
All origins 3/, female												
2006	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.3	*
2005	0.2	*	*	*	0.2	0.1	0.2	0.3	0.2	0.3	0.3	*
2004	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.3	*	*
2003	0.2	*	*	0.1	0.3	0.2	0.2	0.2	0.2	*	0.4	*
2002	0.1	*	*	0.1	0.1	0.2	0.2	0.1	*	*	*	*
2001	0.2	*	*	0.2	0.2	0.1	0.2	0.3	0.2	0.3	0.6	*
2000	0.3	*	*	0.2	0.3	0.2	0.3	0.4	0.2	0.2	0.5	*
1999	0.4	*	0.3	0.2	0.4	0.4	0.4	0.4	0.4	0.5	1.0	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
Hispanic, both sexes												
2006	0.3	*	*	*	0.4	0.3	0.6	0.7	*	*	*	*
2005	0.4	*	*	*	0.5	0.4	0.4	0.5	*	*	*	*
2004	0.4	*	*	*	0.4	0.5	0.6	0.6	*	*	*	*
2003	0.4	*	*	*	0.5	0.7	0.6	*	*	*	*	*
2002	0.4	*	*	*	0.6	0.3	0.5	0.6	*	*	*	*
2001	0.5	*	*	*	0.6	0.6	0.8	0.9	*	*	*	*
2000	0.6	*	*	*	0.5	0.6	0.9	0.8	*	*	*	*
1999	0.8	*	*	*	0.9	1.0	1.2	1.0	*	2.0	*	*
Hispanic, male												
2006	0.6	*	*	*	0.6	*	1.0	1.1	*	*	*	*
2005	0.6	*	*	*	0.9	0.7	0.7	*	*	*	*	*
2004	0.7	*	*	*	0.5	0.9	0.9	1.1	*	*	*	*
2003	0.7	*	*	*	0.6	1.1	1.0	*	*	*	*	*
2002	0.6	*	*	*	1.0	*	0.8	*	*	*	*	*
2001	0.9	*	*	*	1.0	0.9	1.4	1.3	*	*	*	*
2000	0.9	*	*	*	0.9	0.9	1.4	1.4	*	*	*	*
1999	1.3	*	*	*	1.3	1.6	1.8	1.8	*	*	*	*
Hispanic, female												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)												
Non-Hispanic 4/, both sexes												
2006	0.4	*	*	0.1	0.4	0.3	0.5	0.5	0.5	0.5	0.6	0.5
2005	0.4	*	0.2	0.1	0.5	0.4	0.5	0.6	0.5	0.4	0.6	0.5
2004	0.4	*	*	0.1	0.5	0.4	0.5	0.5	0.4	0.5	0.5	0.6
2003	0.5	*	*	0.1	0.7	0.5	0.5	0.6	0.5	0.4	0.6	0.5
2002	0.4	*	*	0.2	0.3	0.4	0.5	0.5	0.4	0.4	0.5	*
2001	0.5	*	*	0.2	0.5	0.4	0.6	0.6	0.5	0.7	0.7	0.6
2000	0.5	*	*	0.2	0.6	0.5	0.7	0.6	0.5	0.5	0.7	0.8
1999	0.7	*	0.3	0.3	0.9	0.7	0.9	0.8	0.8	0.7	1.3	1.3
Non-Hispanic 4/, male												
2006	0.6	*	*	0.2	0.6	0.5	0.8	0.9	0.7	0.9	1.0	*
2005	0.7	*	*	0.2	0.7	0.7	0.7	1.0	0.7	0.6	1.0	*
2004	0.6	*	*	0.2	0.7	0.7	0.7	0.8	0.7	0.7	0.9	*
2003	0.7	*	*	0.2	1.0	0.8	0.8	1.0	0.8	0.7	1.0	*
2002	0.6	*	*	0.2	0.6	0.6	0.8	0.9	0.7	0.7	0.9	*
2001	0.7	*	*	0.3	0.7	0.7	1.0	0.9	0.8	1.0	1.0	*
2000	0.8	*	*	0.3	1.0	0.8	1.1	0.8	0.8	0.8	1.1	*
1999	1.1	*	0.4	0.4	1.4	1.1	1.4	1.2	1.2	1.0	1.8	3.0
Non-Hispanic 4/, female												
2006	0.2	*	*	*	0.2	0.2	0.2	0.2	0.3	0.2	0.3	*
2005	0.2	*	*	*	0.2	0.2	0.2	0.3	0.2	0.3	0.3	*
2004	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.3	*	*
2003	0.2	*	*	0.1	0.3	0.2	0.3	0.2	0.2	*	0.4	*
2002	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2001	0.2	*	*	0.2	0.2	0.1	0.2	0.3	0.2	0.3	0.5	*
2000	0.3	*	*	0.2	0.3	0.2	0.3	0.4	0.2	0.2	0.4	*
1999	0.4	*	*	0.2	0.5	0.4	0.4	0.4	0.4	0.5	1.0	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)												
Non-Hispanic white, both sexes												
2006 5/	0.4	*	*	0.1	0.4	0.3	0.5	0.5	0.5	0.5	0.6	0.5
2005 5/	0.4	*	*	0.1	0.4	0.4	0.5	0.6	0.4	0.4	0.6	0.5
2004 5/	0.4	*	*	0.2	0.5	0.4	0.5	0.4	0.4	0.5	0.5	0.6
2003 5/	0.5	*	*	0.2	0.8	0.5	0.5	0.5	0.5	0.4	0.6	0.5
2002	0.4	*	*	0.2	0.4	0.4	0.5	0.5	0.4	0.4	0.4	*
2001	0.5	*	*	0.2	0.5	0.4	0.5	0.6	0.4	0.6	0.7	0.5
2000	0.5	*	*	0.3	0.7	0.5	0.6	0.6	0.4	0.5	0.7	0.8
1999	0.8	*	0.3	0.3	1.0	0.7	0.9	0.8	0.8	0.7	1.2	1.3
Non-Hispanic white, male												
2006 5/	0.6	*	*	0.2	0.7	0.5	0.8	0.8	0.7	0.9	1.1	*
2005 5/	0.6	*	*	0.2	0.6	0.6	0.7	0.9	0.7	0.5	1.0	*
2004 5/	0.6	*	*	0.2	0.7	0.6	0.7	0.7	0.7	0.7	0.8	*
2003 5/	0.7	*	*	0.2	1.2	0.7	0.8	0.9	0.8	0.7	1.0	*
2002	0.6	*	*	0.2	0.6	0.6	0.7	0.8	0.6	0.6	0.8	*
2001	0.7	*	*	0.2	0.8	0.7	0.8	0.9	0.8	0.9	0.9	*
2000	0.8	*	*	0.3	1.0	0.8	1.0	0.8	0.7	0.7	1.2	*
1999	1.1	*	*	0.4	1.4	1.1	1.3	1.2	1.2	1.0	1.7	2.8
Non-Hispanic white, female												
2006 5/	0.2	*	*	*	0.2	*	0.1	0.2	0.3	*	*	*
2005 5/	0.2	*	*	*	0.2	*	0.3	0.2	0.2	0.3	0.3	*
2004 5/	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.3	*	*
2003 5/	0.2	*	*	*	0.3	0.2	0.2	0.2	0.2	*	0.4	*
2002	0.1	*	*	*	*	0.2	0.2	*	*	*	*	*
2001	0.2	*	*	*	0.2	*	0.2	0.2	*	0.4	0.5	*
2000	0.3	*	*	0.2	0.3	0.2	0.2	0.4	*	*	0.4	*
1999	0.4	*	*	0.2	0.5	0.4	0.4	0.4	0.4	0.4	0.9	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)												
Non-Hispanic black, both sexes												
2006 5/	0.4	*	*	*	*	0.4	0.6	0.8	*	*	*	*
2005 5/	0.5	*	*	*	0.5	0.6	0.6	1.1	*	*	*	*
2004 5/	0.5	*	*	*	0.7	0.6	0.5	0.9	0.8	*	*	*
2003 5/	0.5	*	*	*	0.4	0.6	0.7	1.0	*	*	*	*
2002	0.5	*	*	*	*	*	0.7	0.9	*	*	*	*
2001	0.6	*	*	0.4	0.4	0.5	1.1	0.9	0.9	*	*	*
2000	0.6	*	*	*	0.5	0.6	1.0	0.8	*	*	*	*
1999	0.7	*	*	0.3	0.6	0.8	0.9	1.0	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.7	*	*	*	*	*	0.9	1.6	*	*	*	*
2005 5/	0.9	*	*	*	0.9	1.0	1.0	2.0	*	*	*	*
2004 5/	0.9	*	*	*	1.2	1.2	1.0	1.5	*	*	*	*
2003 5/	0.9	*	*	*	0.7	1.0	1.1	2.0	*	*	*	*
2002	0.8	*	*	*	*	*	1.3	1.9	*	*	*	*
2001	0.9	*	*	*	*	*	1.8	1.4	*	*	*	*
2000	0.9	*	*	*	0.9	1.0	1.6	1.4	*	*	*	*
1999	1.2	*	*	*	1.0	1.4	1.6	1.8	*	*	*	*
Non-Hispanic black, female												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
All origins 3/, both sexes												
2006	0.6	*	*	0.2	0.5	0.7	0.8	0.9	1.0	0.8	0.8	0.4
2005	0.6	*	*	0.1	0.4	0.6	0.8	1.0	1.1	0.9	0.7	0.5
2004	0.7	*	*	0.1	0.5	0.8	0.8	1.1	1.0	0.9	0.5	0.5
2003	0.7	*	*	0.2	0.5	0.6	0.9	1.1	1.2	0.9	0.7	*
2002	0.7	*	*	0.1	0.5	0.6	0.9	1.0	1.1	0.9	0.8	0.4
2001	0.7	*	0.2	0.1	0.5	0.9	1.0	1.1	1.0	1.0	0.8	0.5
2000	0.7	*	*	0.1	0.6	0.8	0.9	1.0	1.1	1.0	0.6	*
1999	0.7	*	*	0.2	0.5	0.8	0.9	1.0	1.1	0.9	0.5	*
All origins 3/, male												
2006	1.0	*	*	0.2	0.7	1.1	1.3	1.6	1.7	1.4	1.4	*
2005	1.0	*	*	0.1	0.7	1.0	1.2	1.7	1.8	1.5	1.3	*
2004	1.1	*	*	0.1	0.7	1.2	1.4	1.8	1.7	1.7	0.9	*
2003	1.1	*	*	0.2	0.8	0.9	1.5	1.8	2.2	1.6	1.3	*
2002	1.1	*	*	0.1	0.8	1.1	1.5	1.8	1.9	1.9	1.5	*
2001	1.2	*	*	0.2	0.9	1.4	1.6	1.9	1.8	1.7	1.4	*
2000	1.2	*	*	0.2	0.9	1.3	1.6	1.7	1.8	1.7	1.3	*
1999	1.1	*	*	0.3	0.8	1.4	1.4	1.7	2.0	1.6	0.9	*
All origins 3/, female												
2006	0.3	*	*	0.1	0.2	0.3	0.3	0.3	0.4	0.4	0.3	*
2005	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.4	0.3	0.3	*
2004	0.3	*	*	*	0.3	0.3	0.3	0.4	0.3	0.2	*	*
2003	0.2	*	*	0.1	0.2	0.3	0.3	0.4	0.3	0.3	0.3	*
2002	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.3	*	0.3	*
2001	0.3	*	*	0.1	0.2	0.3	0.4	0.4	0.3	0.3	0.4	*
2000	0.2	*	*	*	0.2	0.2	0.2	0.3	0.3	0.3	*	*
1999	0.3	*	*	0.1	0.2	0.3	0.4	0.3	0.3	0.2	0.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
Hispanic, both sexes												
2006	0.3	*	*	*	*	0.3	0.5	*	*	*	*	*
2005	0.2	*	*	*	*	*	*	0.6	*	*	*	*
2004	0.3	*	*	*	0.3	0.4	0.4	*	*	*	*	*
2003	0.3	*	*	*	0.3	0.4	0.5	*	*	*	*	*
2002	0.3	*	*	*	0.3	0.3	*	0.6	*	*	*	*
2001	0.8	*	*	*	0.4	0.8	1.2	1.7	2.1	2.0	*	*
2000	0.3	*	*	*	0.4	0.4	0.6	*	*	*	*	*
1999	0.3	*	*	*	*	0.5	*	*	*	*	*	*
Hispanic, male												
2006	0.4	*	*	*	*	0.5	0.8	*	*	*	*	*
2005	0.4	*	*	*	*	*	*	1.1	*	*	*	*
2004	0.4	*	*	*	*	0.6	*	*	*	*	*	*
2003	0.5	*	*	*	*	0.6	0.7	*	*	*	*	*
2002	0.5	*	*	*	0.5	0.6	*	*	*	*	*	*
2001	0.9	*	*	*	*	1.1	1.4	1.7	*	*	*	*
2000	0.5	*	*	*	*	0.6	1.1	*	*	*	*	*
1999	0.4	*	*	*	*	0.7	*	*	*	*	*	*
Hispanic, female												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.7	*	*	*	*	*	1.0	1.8	2.0	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
Non-Hispanic 4/, both sexes												
2006	0.7	*	*	0.2	0.5	0.7	0.9	1.0	1.1	0.9	0.8	0.5
2005	0.7	*	*	0.1	0.5	0.7	0.8	1.1	1.1	0.9	0.7	0.5
2004	0.7	*	*	0.1	0.5	0.8	0.9	1.2	1.0	0.9	0.5	0.6
2003	0.7	*	*	0.2	0.5	0.6	0.9	1.2	1.2	0.9	0.7	*
2002	0.7	*	*	0.1	0.5	0.7	0.9	1.1	1.1	0.9	0.8	*
2001	0.7	*	*	0.1	0.6	0.9	1.0	1.1	1.0	0.9	0.7	*
2000	0.7	*	*	0.1	0.6	0.8	1.0	1.0	1.1	1.0	0.6	*
1999	0.7	*	*	0.2	0.6	0.9	1.0	1.1	1.1	0.9	0.6	*
Non-Hispanic 4/, male												
2006	1.1	*	*	0.2	0.8	1.2	1.4	1.7	1.7	1.5	1.4	*
2005	1.1	*	*	0.1	0.8	1.1	1.4	1.8	1.9	1.5	1.3	*
2004	1.2	*	*	0.2	0.7	1.3	1.5	1.9	1.8	1.8	1.0	*
2003	1.2	*	*	0.2	0.8	1.0	1.6	2.0	2.3	1.7	1.3	*
2002	1.2	*	*	0.1	0.8	1.2	1.6	1.9	2.0	2.0	1.5	*
2001	1.2	*	*	0.2	0.9	1.4	1.6	1.9	1.8	1.7	1.3	*
2000	1.2	*	*	0.2	1.0	1.5	1.7	1.8	1.9	1.7	1.3	*
1999	1.2	*	*	0.3	0.8	1.5	1.6	1.8	2.1	1.7	1.0	*
Non-Hispanic 4/, female												
2006	0.3	*	*	*	0.2	0.3	0.4	0.3	0.4	0.4	0.4	*
2005	0.3	*	*	0.1	0.2	0.2	0.3	0.3	0.4	0.4	0.3	*
2004	0.3	*	*	*	0.3	0.3	0.3	0.4	0.3	0.2	*	*
2003	0.3	*	*	0.2	0.2	0.3	0.3	0.4	0.3	0.3	0.3	*
2002	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.4	*	0.3	*
2001	0.2	*	*	*	0.2	0.3	0.3	0.3	0.2	0.2	0.4	*
2000	0.2	*	*	*	0.2	0.2	0.3	0.3	0.4	0.3	*	*
1999	0.3	*	*	0.1	0.3	0.3	0.4	0.4	0.3	0.3	0.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
Non-Hispanic white, both sexes												
2006 5/	0.8	*	*	0.2	0.6	0.8	1.0	1.1	1.1	0.9	0.8	*
2005 5/	0.8	*	*	0.2	0.5	0.8	0.9	1.1	1.2	1.0	0.7	0.5
2004 5/	0.8	*	*	0.2	0.6	1.0	1.0	1.3	1.1	1.0	0.6	0.6
2003 5/	0.8	*	*	0.2	0.5	0.7	1.1	1.2	1.3	0.9	0.7	*
2002	0.8	*	*	0.1	0.6	0.8	1.0	1.2	1.2	1.0	0.8	*
2001	0.8	*	*	0.1	0.7	1.0	1.1	1.1	1.0	0.9	0.7	*
2000	0.8	*	*	0.2	0.7	1.0	1.1	1.1	1.2	1.0	0.7	*
1999	0.8	*	*	0.2	0.6	1.0	1.1	1.1	1.1	0.9	0.5	*
Non-Hispanic white, male												
2006 5/	1.3	*	*	0.3	1.0	1.3	1.5	1.8	1.9	1.6	1.5	*
2005 5/	1.3	*	*	0.2	0.8	1.3	1.5	1.9	2.1	1.7	1.3	*
2004 5/	1.3	*	*	0.2	0.8	1.5	1.6	2.1	1.9	1.9	1.0	*
2003 5/	1.3	*	*	0.2	0.8	1.2	1.7	2.1	2.4	1.7	1.3	*
2002	1.3	*	*	*	0.9	1.3	1.8	2.0	2.1	2.0	1.4	*
2001	1.3	*	*	0.2	1.1	1.6	1.8	1.9	1.8	1.7	1.4	*
2000	1.4	*	*	0.2	1.1	1.7	1.8	1.9	2.0	1.8	1.3	*
1999	1.3	*	*	0.3	1.0	1.6	1.7	1.9	2.0	1.7	1.0	*
Non-Hispanic white, female												
2006 5/	0.3	*	*	*	0.3	0.3	0.4	0.4	0.4	0.4	0.3	*
2005 5/	0.3	*	*	*	0.3	0.2	0.3	0.3	0.4	0.4	0.4	*
2004 5/	0.3	*	*	*	0.4	0.4	0.4	0.4	0.3	*	*	*
2003 5/	0.3	*	*	0.2	0.2	0.3	0.4	0.4	0.3	0.3	*	*
2002	0.3	*	*	*	0.3	0.2	0.3	0.3	0.4	*	0.3	*
2001	0.3	*	*	*	0.3	0.3	0.4	0.3	0.2	0.2	0.3	*
2000	0.3	*	*	*	0.3	0.3	0.3	0.3	0.4	0.3	*	*
1999	0.3	*	*	*	0.3	0.4	0.5	0.4	0.3	0.3	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
Non-Hispanic black, both sexes												
2006 5/	0.5	*	*	*	*	0.6	0.6	0.7	0.7	*	*	*
2005 5/	0.4	*	*	*	*	*	0.5	1.0	*	*	*	*
2004 5/	0.4	*	*	*	*	0.5	0.7	0.8	0.8	*	*	*
2003 5/	0.4	*	*	*	0.4	*	0.5	0.7	*	*	*	*
2002	0.4	*	*	*	*	0.4	0.6	0.8	0.9	*	*	*
2001	0.5	*	*	*	*	0.5	0.5	1.0	0.9	*	*	*
2000	0.4	*	*	*	*	0.5	0.5	0.6	*	*	*	*
1999	0.5	*	*	*	0.4	0.5	0.7	0.8	1.1	*	*	*
Non-Hispanic black, male												
2006 5/	0.8	*	*	*	*	1.2	1.2	1.4	*	*	*	*
2005 5/	0.6	*	*	*	*	*	0.9	1.5	*	*	*	*
2004 5/	0.7	*	*	*	*	0.9	1.2	1.5	1.6	*	*	*
2003 5/	0.7	*	*	*	0.8	*	1.0	1.2	*	*	*	*
2002	0.8	*	*	*	*	*	1.0	1.3	1.8	*	*	*
2001	0.8	*	*	*	*	0.9	1.0	1.8	2.1	*	*	*
2000	0.7	*	*	*	*	0.9	1.0	1.1	*	*	*	*
1999	0.9	*	*	*	*	0.9	1.2	1.4	2.3	*	*	*
Non-Hispanic black, female												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
All origins 3/, both sexes												
2006	24.4	24.3	6.1	2.0	11.4	17.8	23.6	28.7	20.5	27.8	81.4	250.5
2005	23.4	22.8	6.3	2.1	10.6	15.8	21.9	26.5	19.6	28.3	81.9	253.4
2004	21.8	22.2	6.1	2.2	9.8	13.8	20.9	24.0	17.7	26.5	78.2	249.5
2003	21.0	19.9	6.8	2.1	9.3	13.2	20.7	22.3	16.9	26.6	75.6	249.2
2002	20.2	20.3	6.4	2.4	8.9	12.7	20.1	20.2	15.7	25.9	74.4	246.7
2001	19.0	20.5	6.8	2.5	8.2	11.1	18.0	18.1	15.0	24.9	73.0	247.5
2000	18.2	18.6	7.4	2.6	7.9	11.0	17.2	16.8	14.9	23.8	68.1	242.0
1999	18.4	17.3	7.7	2.7	7.7	11.1	17.0	16.4	14.6	25.1	72.1	250.3
All origins 3/, male												
2006	30.4	28.3	7.2	2.6	18.1	26.8	32.9	39.9	28.7	36.7	102.6	294.3
2005	29.0	24.0	7.9	2.8	16.7	23.6	30.6	37.3	27.8	37.4	104.2	298.3
2004	26.8	25.1	7.4	3.0	15.4	20.8	28.9	34.0	24.6	35.5	98.2	299.0
2003	26.0	22.4	8.5	2.8	14.7	19.9	29.4	31.6	24.1	35.3	96.3	300.5
2002	25.2	23.3	8.1	3.2	14.2	19.6	29.1	29.3	22.7	35.3	93.9	293.6
2001	23.8	22.9	8.3	3.3	13.3	17.1	26.8	27.3	21.4	33.0	93.6	305.8
2000	22.9	20.9	9.0	3.4	12.8	17.3	26.2	26.0	21.6	31.5	86.2	299.8
1999	23.3	20.2	9.3	3.7	12.5	17.7	26.2	25.6	21.5	34.1	91.7	311.9
All origins 3/, female												
2006	18.6	20.2	5.0	1.4	4.3	8.5	14.1	17.8	12.8	20.1	67.0	230.1
2005	18.0	21.5	4.5	1.4	4.1	7.8	13.2	16.0	11.8	20.7	66.7	232.8
2004	16.9	19.3	4.8	1.5	3.8	6.5	12.9	14.4	11.2	18.9	64.8	227.3
2003	16.2	17.3	5.0	1.4	3.6	6.3	12.0	13.3	10.3	19.3	61.8	226.5
2002	15.4	17.2	4.7	1.5	3.2	5.7	11.1	11.5	9.2	18.1	61.4	226.3
2001	14.5	17.9	5.1	1.6	2.8	4.9	9.4	9.2	9.1	18.1	59.5	223.1
2000	13.6	16.2	5.7	1.7	2.7	4.5	8.2	7.9	8.8	17.5	56.3	218.4
1999	13.7	14.2	6.1	1.7	2.6	4.4	8.0	7.6	8.2	17.8	59.4	225.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
Hispanic, both sexes												
2006	12.4	14.8	5.0	1.5	8.3	11.9	16.2	20.0	17.8	20.2	54.2	131.9
2005	12.0	13.7	5.2	1.1	8.7	11.1	15.2	19.2	15.9	23.1	55.5	138.2
2004	10.6	13.1	4.3	1.2	7.1	9.3	14.7	16.7	14.6	20.2	52.3	140.8
2003	11.0	10.6	5.6	1.6	7.6	10.2	15.3	17.7	14.8	19.6	47.3	127.0
2002	10.8	12.1	5.1	1.5	7.9	9.4	15.7	16.8	13.5	21.1	51.2	127.5
2001	10.1	13.6	5.0	1.6	6.6	9.4	15.0	15.3	12.6	19.8	43.6	142.6
2000	10.2	9.9	5.2	1.5	7.3	10.0	15.3	14.2	13.1	18.4	48.2	159.2
1999	11.1	12.4	6.4	1.7	7.7	10.5	17.2	16.2	13.0	20.5	44.6	157.2
Hispanic, male												
2006	18.0	16.1	6.0	2.0	13.5	18.4	25.5	31.0	28.5	30.0	66.4	155.6
2005	17.3	13.9	6.8	1.4	14.3	17.5	23.6	30.7	25.7	32.7	70.8	146.9
2004	15.2	14.9	5.1	1.5	11.4	14.7	22.6	26.3	23.2	28.1	64.7	165.3
2003	16.1	11.8	7.3	2.1	12.5	16.2	24.1	28.2	23.9	27.1	67.1	149.3
2002	16.1	15.7	6.4	1.9	13.1	15.5	24.6	27.1	21.4	30.2	68.5	179.0
2001	14.8	14.2	6.7	1.9	10.8	14.9	24.3	25.6	20.4	26.8	54.1	154.3
2000	15.3	10.9	5.9	1.6	12.1	16.5	25.6	24.3	19.7	27.6	57.5	207.6
1999	16.9	15.0	7.7	2.3	12.9	17.4	28.2	27.6	22.5	31.6	54.7	217.8
Hispanic, female												
2006	6.4	13.5	4.0	1.0	2.3	3.9	5.7	8.6	7.9	12.3	45.6	119.2
2005	6.2	13.6	3.6	0.8	2.3	3.3	5.8	7.4	7.0	15.3	44.9	133.6
2004	5.8	11.3	3.4	0.8	2.2	2.8	6.0	7.1	6.9	13.9	43.7	128.2
2003	5.4	9.4	3.9	1.2	2.0	3.1	5.5	7.2	6.7	13.5	33.7	115.6
2002	5.2	8.3	3.7	1.2	1.9	2.1	5.9	6.6	6.3	13.8	39.3	102.0
2001	5.2	13.0	3.3	1.3	1.7	2.9	5.0	5.2	5.7	14.3	36.6	136.8
2000	4.9	8.8	4.4	1.5	1.6	2.4	4.4	4.3	7.3	11.1	41.9	135.5
1999	5.0	9.8	5.2	1.1	1.7	2.6	5.5	5.1	4.6	11.7	37.8	127.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
Non-Hispanic 4/, both sexes												
2006	26.4	27.1	6.3	2.1	12.0	19.2	24.8	29.5	20.6	28.3	82.9	256.3
2005	25.2	25.3	6.6	2.4	11.0	16.9	23.0	27.1	19.7	28.6	83.3	258.6
2004	23.5	24.8	6.6	2.5	10.3	14.8	21.8	24.6	17.8	26.8	79.5	254.1
2003	22.5	22.4	7.1	2.2	9.6	13.8	21.4	22.5	17.0	27.0	76.9	253.9
2002	21.6	22.3	6.7	2.6	9.0	13.4	20.6	20.4	15.8	26.1	75.2	250.9
2001	20.2	22.0	7.2	2.6	8.5	11.3	18.3	18.2	15.0	25.1	74.2	250.8
2000	19.2	20.7	7.9	2.8	8.0	11.1	17.3	16.8	14.9	24.1	68.7	244.4
1999	19.3	18.5	8.0	2.9	7.6	11.1	16.8	16.2	14.5	25.3	72.9	252.8
Non-Hispanic 4/, male												
2006	32.6	31.8	7.4	2.7	19.0	29.0	34.2	40.7	28.5	37.2	104.6	301.7
2005	30.9	26.8	8.3	3.2	17.2	25.1	31.8	37.8	27.8	37.5	106.0	305.9
2004	28.7	27.9	8.0	3.3	16.3	22.3	29.9	34.6	24.6	35.9	100.0	305.1
2003	27.5	25.2	8.7	2.9	15.2	20.7	30.0	31.7	24.0	35.7	97.6	306.9
2002	26.5	25.2	8.5	3.5	14.4	20.5	29.6	29.3	22.6	35.4	94.7	298.0
2001	25.0	24.9	8.7	3.5	13.7	17.5	26.9	27.2	21.2	33.2	95.1	311.6
2000	23.9	23.2	9.7	3.8	12.9	17.4	26.1	25.8	21.5	31.6	87.2	302.7
1999	24.0	21.5	9.6	3.9	12.4	17.6	25.7	25.0	21.1	34.1	92.8	314.3
Non-Hispanic 4/, female												
2006	20.6	22.2	5.2	1.5	4.7	9.5	15.5	18.8	13.2	20.7	68.2	235.2
2005	19.8	23.7	4.8	1.6	4.5	8.7	14.3	16.8	12.2	21.1	67.9	237.0
2004	18.6	21.6	5.2	1.6	4.1	7.3	13.9	15.1	11.5	19.2	65.8	231.2
2003	17.8	19.5	5.3	1.4	3.9	6.9	12.9	13.8	10.6	19.7	63.1	230.6
2002	16.9	19.4	4.9	1.6	3.4	6.4	11.8	11.9	9.5	18.4	62.3	230.6
2001	15.7	19.1	5.6	1.7	3.0	5.2	9.9	9.5	9.4	18.3	60.4	225.5
2000	14.7	18.0	6.0	1.7	2.9	4.8	8.6	8.1	8.8	17.8	56.7	220.8
1999	14.8	15.2	6.3	1.9	2.7	4.6	8.2	7.7	8.4	18.1	60.1	228.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
Non-Hispanic white, both sexes												
2006 5/	28.8	21.7	5.9	1.8	13.7	22.1	26.4	29.5	19.9	28.2	86.2	269.8
2005 5/	27.3	21.9	6.4	2.0	12.3	19.0	24.4	26.9	18.8	28.3	84.9	269.7
2004 5/	25.6	21.2	6.5	2.0	11.6	16.6	23.3	24.7	17.1	26.7	81.8	266.4
2003 5/	24.4	18.4	6.9	1.8	10.9	15.3	22.4	22.1	16.6	26.6	78.9	265.9
2002	23.1	17.2	6.1	2.0	9.8	14.7	21.4	19.9	15.1	25.6	77.0	261.2
2001	21.5	17.8	7.1	2.2	9.1	12.1	18.5	17.2	14.2	24.7	75.4	259.4
2000	20.1	16.6	7.3	2.3	8.3	11.5	17.2	15.6	14.1	23.4	69.6	253.2
1999	20.0	14.7	7.4	2.4	7.8	11.3	16.7	14.9	13.3	24.2	73.2	260.8
Non-Hispanic white, male												
2006 5/	34.8	25.0	6.9	2.4	21.6	32.9	36.0	39.6	26.7	36.7	107.8	316.9
2005 5/	32.7	22.8	8.4	2.7	19.3	28.1	33.5	36.7	25.6	36.7	107.1	317.6
2004 5/	30.7	25.0	8.0	2.7	18.3	24.9	31.7	34.2	22.9	35.5	102.2	317.3
2003 5/	29.2	20.8	8.5	2.4	17.1	22.7	31.4	30.6	22.8	34.8	99.1	320.0
2002	27.8	20.4	7.6	2.7	15.6	22.2	30.5	28.1	21.0	34.3	96.0	307.6
2001	25.9	20.0	8.8	2.9	14.7	18.6	27.2	25.3	19.6	32.5	95.6	320.4
2000	24.5	19.1	9.1	3.2	13.5	18.1	26.0	23.6	19.9	30.4	87.2	311.5
1999	24.3	17.1	8.9	3.2	12.5	17.9	25.4	22.7	19.1	32.0	92.4	322.6
Non-Hispanic white, female												
2006 5/	23.1	18.2	4.8	1.2	5.5	11.1	16.8	19.5	13.4	20.9	71.2	247.9
2005 5/	22.0	21.1	4.4	1.3	4.9	9.8	15.3	17.2	12.4	21.1	69.6	247.8
2004 5/	20.7	17.3	4.9	1.3	4.6	8.3	14.9	15.4	11.6	19.1	68.0	243.7
2003 5/	19.7	16.0	5.1	1.1	4.4	7.7	13.5	13.8	10.7	19.6	65.3	242.0
2002	18.6	13.9	4.6	1.2	3.8	7.1	12.3	11.8	9.4	18.3	64.2	241.1
2001	17.2	15.5	5.3	1.5	3.3	5.5	9.8	9.3	9.2	18.1	62.0	234.0
2000	16.0	14.0	5.4	1.3	3.0	4.9	8.4	7.7	8.7	17.6	58.1	229.7
1999	16.0	12.3	5.9	1.5	2.9	4.6	8.0	7.3	7.9	17.6	60.6	236.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
Non-Hispanic black, both sexes												
2006 5/	20.7	49.9	9.0	3.4	7.2	13.3	24.5	38.0	30.3	33.1	61.9	137.2
2005 5/	20.5	42.9	7.9	3.8	7.0	12.7	21.8	35.8	31.2	35.3	77.8	159.8
2004 5/	18.2	42.0	7.6	4.3	6.3	10.9	20.3	30.9	26.8	32.1	66.5	140.2
2003 5/	17.7	40.3	8.5	3.8	6.1	10.9	20.9	30.7	24.3	32.9	60.5	140.4
2002	18.3	44.5	9.5	4.7	6.9	11.4	21.4	29.6	24.6	32.4	62.8	156.7
2001	18.1	41.6	8.7	4.3	6.8	10.5	21.7	29.4	23.9	32.5	68.2	166.1
2000	18.0	39.5	10.9	4.8	7.1	11.3	21.5	28.1	23.8	33.1	65.4	161.2
1999	19.4	36.4	11.1	5.3	7.8	12.7	21.2	29.1	26.6	39.1	77.8	181.1
Non-Hispanic black, male												
2006 5/	28.8	59.5	10.5	4.2	11.7	21.3	35.7	58.8	49.7	48.2	85.6	164.9
2005 5/	27.9	47.3	9.0	5.0	11.0	18.9	30.8	55.4	52.9	52.1	111.0	193.2
2004 5/	24.6	43.0	9.1	5.7	9.9	16.9	28.3	46.4	43.7	47.7	94.7	189.0
2003 5/	24.1	45.3	10.8	4.8	9.5	16.6	30.2	46.1	39.3	48.4	85.7	184.3
2002	25.5	46.7	12.5	6.5	11.6	18.0	31.7	45.3	40.6	48.5	90.6	215.5
2001	25.0	48.2	10.0	6.0	11.0	16.2	31.6	47.9	38.0	47.5	101.4	220.6
2000	25.5	42.3	12.9	6.5	11.7	17.9	32.7	46.6	39.8	47.8	96.9	231.3
1999	27.5	42.3	13.7	7.3	13.5	20.0	33.3	48.7	42.5	60.1	107.6	249.6
Non-Hispanic black, female												
2006 5/	13.3	39.9	7.5	2.5	2.6	5.8	14.5	20.2	14.7	22.2	48.4	125.9
2005 5/	13.7	38.3	6.8	2.6	2.9	7.1	13.8	19.0	13.7	23.3	58.9	146.3
2004 5/	12.3	40.9	6.1	2.8	2.6	5.4	13.3	17.6	13.3	21.0	50.6	120.9
2003 5/	12.0	35.1	6.2	2.7	2.7	5.6	12.7	17.6	12.2	21.8	46.3	123.4
2002	11.8	42.3	6.5	2.9	2.1	5.4	12.3	16.2	11.7	21.1	47.1	134.2
2001	11.8	34.6	7.4	2.4	2.5	5.4	12.9	13.5	12.6	21.9	49.5	145.9
2000	11.3	36.6	8.8	3.0	2.6	5.4	11.6	12.2	10.9	22.8	47.8	135.2
1999	12.0	30.3	8.3	3.3	2.1	6.1	10.7	12.3	13.8	24.3	61.1	155.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
All origins 3/, both sexes												
2006	7.0	0.6	0.2	0.1	0.6	0.8	1.4	3.0	5.0	12.6	47.7	151.6
2005	6.6	*	0.2	0.1	0.6	0.7	1.4	2.8	4.8	12.4	45.6	147.7
2004	6.4	0.6	0.3	0.1	0.6	0.8	1.5	2.8	4.8	12.2	43.8	143.3
2003	5.9	*	0.3	0.1	0.6	0.7	1.4	2.6	4.4	11.2	40.8	135.9
2002	5.6	*	0.2	0.1	0.6	0.8	1.5	2.5	4.1	10.8	38.3	130.4
2001	5.3	0.6	0.2	0.2	0.6	0.9	1.4	2.6	4.0	10.0	35.3	121.5
2000	4.7	*	0.2	0.1	0.6	0.8	1.3	2.3	3.9	9.0	31.1	112.6
1999	4.7	*	0.4	0.1	0.6	0.9	1.4	2.3	3.7	9.0	31.5	110.2
All origins 3/, male												
2006	7.3	*	0.3	0.1	1.0	1.3	2.3	4.7	7.3	16.5	59.4	181.1
2005	7.0	*	0.2	0.1	0.9	1.3	2.2	4.3	7.0	16.2	57.8	174.8
2004	6.8	*	0.4	0.1	1.0	1.3	2.4	4.5	7.1	16.5	54.8	174.4
2003	6.2	*	0.5	0.1	0.9	1.2	2.4	4.0	6.2	14.5	51.6	165.1
2002	6.0	*	0.4	0.1	1.0	1.3	2.3	3.9	6.0	14.5	48.1	158.9
2001	5.8	*	0.3	0.2	1.1	1.5	2.4	4.3	5.9	13.1	45.7	155.7
2000	5.2	*	0.3	0.1	1.0	1.4	2.2	3.7	5.8	11.9	39.9	145.4
1999	5.2	*	0.5	0.2	1.1	1.5	2.4	3.7	5.7	12.4	40.5	142.5
All origins 3/, female												
2006	6.6	*	*	*	0.1	0.2	0.5	1.3	3.0	9.4	39.8	137.8
2005	6.3	*	*	*	0.2	0.2	0.5	1.3	2.7	9.2	37.4	135.2
2004	6.0	*	*	*	0.2	0.2	0.6	1.3	2.7	8.6	36.4	129.2
2003	5.6	*	*	*	0.2	0.2	0.5	1.2	2.7	8.4	33.5	122.9
2002	5.3	*	*	*	0.2	0.2	0.6	1.2	2.3	7.7	31.8	118.1
2001	4.8	*	*	0.1	0.1	0.2	0.5	1.0	2.2	7.4	28.5	107.2
2000	4.3	*	*	*	0.2	0.2	0.5	0.9	2.1	6.6	25.3	99.2
1999	4.3	*	*	*	0.2	0.2	0.5	0.9	1.9	6.3	25.7	97.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
Hispanic, both sexes												
2006	2.7	*	*	*	0.7	1.2	1.8	3.2	4.6	10.7	34.5	90.4
2005	2.5	*	*	*	0.7	0.9	1.3	2.7	4.7	11.8	34.8	91.6
2004	2.4	*	*	*	0.6	1.1	1.8	2.3	4.9	10.2	30.8	87.1
2003	2.1	*	*	*	0.5	0.8	1.4	2.6	5.0	9.2	25.4	80.8
2002	2.1	*	*	*	0.7	0.9	1.6	2.5	4.4	10.0	29.2	71.5
2001	2.0	*	*	*	0.7	1.1	1.8	2.6	4.2	9.1	23.6	70.0
2000	2.0	*	*	*	0.7	1.1	1.3	2.5	4.3	8.0	26.5	81.0
1999	1.9	*	*	*	0.7	1.1	1.4	2.2	3.8	8.1	24.2	86.3
Hispanic, male												
2006	3.5	*	*	*	1.3	2.1	3.1	5.5	7.7	15.1	41.4	102.6
2005	3.1	*	*	*	1.2	1.6	2.2	4.8	7.5	16.4	46.8	87.2
2004	3.1	*	*	*	1.1	1.8	3.0	4.2	8.1	14.0	38.3	106.5
2003	2.7	*	*	*	0.9	1.4	2.6	4.3	7.5	11.8	36.0	90.8
2002	2.8	*	*	*	1.3	1.6	2.7	3.8	7.0	13.8	36.9	97.0
2001	2.8	*	*	*	1.2	1.9	3.3	4.5	7.1	11.4	27.7	72.4
2000	2.7	*	*	*	1.2	2.0	2.3	4.4	6.6	12.7	30.5	104.8
1999	2.7	*	*	*	1.3	1.9	2.4	3.8	7.2	11.4	31.3	120.8
Hispanic, female												
2006	1.8	*	*	*	*	*	*	*	1.7	7.1	29.7	83.9
2005	1.8	*	*	*	*	*	*	*	2.2	8.1	26.5	93.8
2004	1.6	*	*	*	*	*	*	*	2.0	7.2	25.7	77.1
2003	1.5	*	*	*	*	*	*	*	2.8	7.2	18.2	75.8
2002	1.4	*	*	*	*	*	*	1.1	2.1	7.0	23.9	58.9
2001	1.3	*	*	*	*	*	*	*	*	7.2	20.9	68.9
2000	1.2	*	*	*	*	*	*	*	2.3	4.3	23.8	69.2
1999	1.1	*	*	*	*	*	*	*	*	5.5	19.4	69.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
Non-Hispanic 4/, both sexes												
2006	7.7	*	0.2	0.1	0.5	0.7	1.4	2.9	5.1	12.8	48.5	154.6
2005	7.3	*	0.2	0.1	0.5	0.7	1.4	2.8	4.8	12.5	46.2	150.2
2004	7.0	*	0.3	0.1	0.6	0.7	1.4	2.9	4.8	12.3	44.5	145.5
2003	6.5	*	0.3	0.1	0.6	0.7	1.4	2.5	4.3	11.3	41.5	137.9
2002	6.2	*	0.2	0.1	0.6	0.7	1.5	2.5	4.0	10.8	38.6	132.6
2001	5.7	*	0.2	0.1	0.6	0.8	1.4	2.6	3.9	10.0	35.8	123.1
2000	5.1	*	0.2	0.1	0.6	0.7	1.3	2.3	3.9	9.1	31.2	113.5
1999	5.1	*	0.4	0.1	0.6	0.8	1.4	2.2	3.7	9.0	31.7	110.7
Non-Hispanic 4/, male												
2006	8.0	*	0.3	*	0.9	1.1	2.2	4.6	7.2	16.6	60.4	185.4
2005	7.6	*	*	0.1	0.9	1.2	2.2	4.2	6.9	16.2	58.4	179.2
2004	7.4	*	0.4	0.1	0.9	1.2	2.3	4.4	6.9	16.7	55.6	177.2
2003	6.8	*	0.5	0.1	0.9	1.1	2.3	3.9	6.0	14.6	52.3	168.3
2002	6.5	*	0.3	0.1	0.9	1.3	2.3	3.9	5.9	14.4	48.4	161.3
2001	6.2	*	0.3	0.2	1.1	1.4	2.3	4.2	5.8	13.1	46.4	159.0
2000	5.5	*	*	0.2	1.0	1.2	2.2	3.6	5.7	11.9	40.2	146.5
1999	5.5	*	0.5	0.2	1.0	1.3	2.3	3.6	5.5	12.4	40.6	142.9
Non-Hispanic 4/, female												
2006	7.4	*	*	*	0.2	0.3	0.6	1.4	3.1	9.5	40.3	140.2
2005	7.0	*	*	*	0.2	0.2	0.6	1.4	2.8	9.3	37.9	136.9
2004	6.7	*	*	*	0.2	0.2	0.6	1.4	2.8	8.7	37.0	131.3
2003	6.2	*	*	*	0.2	0.3	0.6	1.2	2.7	8.5	34.3	124.6
2002	5.9	*	*	*	0.2	0.2	0.7	1.2	2.3	7.7	32.1	120.2
2001	5.3	*	*	*	0.1	0.3	0.5	1.0	2.2	7.4	28.8	108.1
2000	4.7	*	*	*	0.2	0.2	0.5	1.0	2.1	6.8	25.3	100.1
1999	4.6	*	*	*	0.2	0.3	0.5	0.9	2.0	6.3	25.9	97.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
Non-Hispanic white, both sexes												
2006 5/	9.0	*	*	0.1	0.6	0.8	1.5	3.0	5.1	13.3	51.4	164.5
2005 5/	8.5	*	*	0.1	0.6	0.7	1.5	2.8	4.8	13.0	48.5	159.6
2004 5/	8.2	*	0.3	0.1	0.7	0.8	1.5	2.9	4.8	12.8	47.2	155.1
2003 5/	7.5	*	0.3	0.1	0.6	0.8	1.5	2.5	4.3	11.6	43.6	146.8
2002	7.1	*	*	0.1	0.7	0.8	1.5	2.5	4.1	11.1	40.8	140.4
2001	6.6	*	0.2	0.2	0.8	0.9	1.4	2.6	3.9	10.3	37.6	129.9
2000	5.8	*	*	0.1	0.7	0.7	1.4	2.3	3.8	9.2	32.7	120.2
1999	5.7	*	0.4	0.2	0.7	0.8	1.5	2.1	3.5	9.1	33.1	116.6
Non-Hispanic white, male												
2006 5/	9.2	*	*	*	1.0	1.2	2.3	4.6	7.2	17.1	63.3	196.1
2005 5/	8.7	*	*	*	1.0	1.2	2.4	4.2	6.9	16.8	60.9	188.7
2004 5/	8.5	*	*	*	1.1	1.3	2.4	4.5	6.7	17.3	58.6	186.9
2003 5/	7.7	*	0.5	0.2	1.0	1.2	2.4	3.8	6.0	14.8	54.6	178.7
2002	7.3	*	*	*	1.0	1.4	2.3	3.9	5.9	14.8	50.6	169.2
2001	7.0	*	*	0.2	1.3	1.5	2.3	4.1	5.7	13.5	48.2	166.0
2000	6.2	*	*	0.2	1.1	1.3	2.2	3.6	5.6	12.0	41.7	154.1
1999	6.1	*	0.5	0.2	1.1	1.4	2.3	3.4	5.1	12.2	42.0	149.7
Non-Hispanic white, female												
2006 5/	8.8	*	*	*	0.2	0.3	0.6	1.4	3.2	10.1	43.1	149.8
2005 5/	8.3	*	*	*	0.2	0.2	0.6	1.5	2.8	9.8	40.1	146.3
2004 5/	7.9	*	*	*	0.2	0.2	0.5	1.4	2.9	9.0	39.4	140.9
2003 5/	7.3	*	*	*	0.2	0.3	0.6	1.2	2.7	8.9	36.2	132.8
2002	7.0	*	*	*	0.2	0.2	0.7	1.2	2.4	8.1	34.2	128.0
2001	6.2	*	*	*	0.2	0.3	0.5	1.1	2.2	7.7	30.6	114.8
2000	5.5	*	*	*	0.2	0.2	0.5	1.0	2.1	6.9	26.8	106.5
1999	5.4	*	*	*	0.2	0.3	0.6	0.8	1.9	6.4	27.2	103.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
Non-Hispanic black, both sexes												
2006 5/	2.8	*	*	*	*	0.4	1.2	3.1	4.8	9.8	23.9	54.9
2005 5/	2.7	*	*	*	*	0.5	1.0	3.1	4.7	9.0	25.7	53.0
2004 5/	2.7	*	*	*	*	0.6	1.5	2.9	5.6	10.1	20.8	47.5
2003 5/	2.5	*	*	*	*	0.5	1.3	3.2	4.9	9.0	19.3	44.6
2002	2.3	*	*	*	*	0.4	1.3	2.8	4.3	8.2	16.3	49.3
2001	2.3	*	*	*	*	0.5	1.6	3.0	4.4	8.8	18.7	44.0
2000	2.1	*	*	*	*	0.4	1.3	2.7	4.8	8.3	15.2	39.1
1999	2.5	*	*	*	0.4	0.8	1.3	3.5	5.5	9.3	17.2	45.3
Non-Hispanic black, male												
2006 5/	3.6	*	*	*	*	*	1.9	5.4	8.0	14.5	34.5	68.7
2005 5/	3.3	*	*	*	*	*	1.6	5.3	7.7	12.6	35.6	63.8
2004 5/	3.5	*	*	*	*	1.1	2.2	4.8	10.2	14.5	29.9	73.2
2003 5/	3.0	*	*	*	*	*	2.2	5.2	7.2	12.6	24.5	56.5
2002	3.0	*	*	*	*	0.8	2.2	4.7	6.8	11.9	24.9	76.6
2001	3.2	*	*	*	*	*	2.7	5.4	6.9	12.6	29.1	68.1
2000	2.8	*	*	*	*	*	2.1	4.6	8.2	11.4	21.6	61.1
1999	3.4	*	*	*	*	1.4	2.5	5.9	8.7	15.0	24.1	58.2
Non-Hispanic black, female												
2006 5/	2.1	*	*	*	*	*	*	1.2	2.3	6.4	17.8	49.3
2005 5/	2.1	*	*	*	*	*	*	1.2	2.2	6.4	20.0	48.6
2004 5/	1.9	*	*	*	*	*	0.9	1.4	1.9	7.0	15.6	37.3
2003 5/	2.0	*	*	*	*	*	*	1.5	3.0	6.5	16.3	40.0
2002	1.6	*	*	*	*	*	*	1.2	2.2	5.6	11.4	38.9
2001	1.6	*	*	*	*	*	*	*	2.3	6.1	12.8	35.1
2000	1.5	*	*	*	*	*	*	1.0	2.0	6.1	11.6	31.0
1999	1.7	*	*	*	*	*	*	1.4	2.9	5.3	13.3	40.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
All origins 3/, both sexes												
2006	0.2	*	*	0.1	0.5	0.3	0.2	0.2	0.2	0.2	0.3	*
2005	0.3	*	0.1	0.1	0.5	0.3	0.3	0.3	0.2	0.2	0.3	*
2004	0.2	*	*	0.1	0.4	0.2	0.2	0.3	0.1	0.2	0.2	*
2003	0.3	*	*	0.1	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2002	0.3	*	*	0.1	0.5	0.4	0.3	0.2	0.2	0.2	0.3	*
2001	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*
All origins 3/, male												
2006	0.4	*	*	0.2	0.9	0.5	0.3	0.3	0.3	0.4	0.5	*
2005	0.5	*	*	0.2	0.9	0.6	0.4	0.4	0.4	0.4	0.5	*
2004	0.4	*	*	0.2	0.7	0.5	0.4	0.4	0.2	0.4	0.4	*
2003	0.5	*	*	0.2	0.9	0.5	0.5	0.4	0.4	0.4	0.6	*
2002	0.5	*	*	0.2	0.9	0.6	0.4	0.4	0.4	0.4	0.5	*
2001	0.5	*	*	0.2	0.9	0.5	0.5	0.5	0.5	0.3	0.7	*
2000	0.5	*	*	0.3	0.9	0.5	0.6	0.4	0.4	0.4	0.5	*
1999	0.5	*	*	0.3	1.2	0.6	0.5	0.4	0.4	0.4	0.4	*
All origins 3/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
Hispanic, both sexes												
2006	0.1	*	*	*	0.4	*	*	*	*	*	*	*
2005	0.2	*	*	*	0.5	0.3	*	*	*	*	*	*
2004	0.1	*	*	*	0.3	*	*	*	*	*	*	*
2003	0.2	*	*	*	0.5	*	*	*	*	*	*	*
2002	0.2	*	*	*	0.4	*	*	*	*	*	*	*
2001	0.2	*	*	*	0.5	0.3	*	*	*	*	*	*
2000	0.2	*	*	*	0.4	*	*	*	*	*	*	*
1999	0.2	*	*	*	0.5	*	*	*	*	*	*	*
Hispanic, male												
2006	0.3	*	*	*	0.8	*	*	*	*	*	*	*
2005	0.4	*	*	*	0.9	0.5	*	*	*	*	*	*
2004	0.2	*	*	*	0.6	*	*	*	*	*	*	*
2003	0.3	*	*	*	0.8	*	*	*	*	*	*	*
2002	0.3	*	*	*	0.6	*	*	*	*	*	*	*
2001	0.4	*	*	*	0.9	*	*	*	*	*	*	*
2000	0.3	*	*	*	0.7	*	*	*	*	*	*	*
1999	0.3	*	*	*	0.9	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
Non-Hispanic 4/, both sexes												
2006	0.2	*	*	0.1	0.5	0.3	0.2	0.2	0.2	0.2	0.3	*
2005	0.3	*	0.2	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2004	0.2	*	*	0.1	0.4	0.3	0.2	0.3	0.1	0.2	0.2	*
2003	0.3	*	*	0.1	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2002	0.3	*	*	0.1	0.5	0.4	0.3	0.3	0.2	0.2	0.2	*
2001	0.3	*	*	0.2	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000	0.3	*	*	0.2	0.5	0.4	0.4	0.2	0.2	0.2	0.2	*
1999	0.3	*	*	0.2	0.7	0.4	0.3	0.3	0.2	0.2	0.2	*
Non-Hispanic 4/, male												
2006	0.4	*	*	0.2	0.9	0.5	0.3	0.3	0.3	0.4	0.6	*
2005	0.5	*	*	0.2	0.9	0.6	0.5	0.4	0.4	0.4	0.5	*
2004	0.4	*	*	0.2	0.8	0.5	0.4	0.5	0.2	0.4	0.5	*
2003	0.5	*	*	0.2	0.9	0.6	0.5	0.4	0.5	0.4	0.7	*
2002	0.5	*	*	0.2	1.0	0.7	0.5	0.5	0.4	0.4	0.5	*
2001	0.5	*	*	0.3	0.9	0.5	0.6	0.5	0.5	0.3	0.7	*
2000	0.5	*	*	0.3	1.0	0.6	0.6	0.4	0.4	0.4	0.5	*
1999	0.5	*	*	0.3	1.2	0.6	0.5	0.5	0.4	0.4	0.4	*
Non-Hispanic 4/, female												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.1	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
Non-Hispanic white, both sexes												
2006 5/	0.2	*	*	0.1	0.3	0.3	0.2	0.2	0.2	0.2	0.3	*
2005 5/	0.3	*	*	0.1	0.4	0.2	0.3	0.3	0.3	0.3	0.3	*
2004 5/	0.2	*	*	0.1	0.4	0.2	0.3	0.3	0.1	0.2	0.2	*
2003 5/	0.3	*	*	0.1	0.4	0.3	0.3	0.3	0.3	0.2	0.3	*
2002	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.2	0.2	0.2	*
2001	0.3	*	*	0.2	0.5	0.3	0.4	0.3	0.3	0.2	0.3	*
2000	0.3	*	*	0.2	0.5	0.3	0.4	0.2	0.2	0.2	0.2	*
1999	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*
Non-Hispanic white, male												
2006 5/	0.4	*	*	0.2	0.6	0.5	0.3	0.3	0.3	0.4	0.6	*
2005 5/	0.4	*	*	0.2	0.7	0.4	0.4	0.4	0.5	0.5	0.5	*
2004 5/	0.4	*	*	0.2	0.7	0.4	0.4	0.5	0.2	0.4	*	*
2003 5/	0.5	*	*	0.2	0.8	0.5	0.6	0.4	0.5	0.4	0.6	*
2002	0.5	*	*	0.2	0.8	0.6	0.5	0.5	0.4	0.5	0.5	*
2001	0.5	*	*	0.3	0.8	0.6	0.6	0.5	0.5	0.3	0.7	*
2000	0.5	*	*	0.3	0.8	0.5	0.6	0.4	0.4	0.4	0.5	*
1999	0.5	*	*	0.3	1.0	0.6	0.5	0.5	0.4	0.4	*	*
Non-Hispanic white, female												
2006 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
Non-Hispanic black, both sexes												
2006 5/	0.4	*	*	*	1.1	0.6	*	*	*	*	*	*
2005 5/	0.4	*	*	*	0.8	0.7	*	*	*	*	*	*
2004 5/	0.3	*	*	*	0.7	0.6	*	*	*	*	*	*
2003 5/	0.3	*	*	*	0.8	0.7	*	*	*	*	*	*
2002	0.4	*	*	*	1.0	0.8	0.4	*	*	*	*	*
2001	0.3	*	*	*	0.8	*	*	*	*	*	*	*
2000	0.4	*	*	*	0.9	0.6	*	*	*	*	*	*
1999	0.4	*	*	*	1.2	0.4	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.7	*	*	*	2.1	1.1	*	*	*	*	*	*
2005 5/	0.8	*	*	*	1.4	1.5	*	*	*	*	*	*
2004 5/	0.7	*	*	*	1.4	1.2	*	*	*	*	*	*
2003 5/	0.7	*	*	*	1.5	1.3	*	*	*	*	*	*
2002	0.8	*	*	*	1.9	1.6	*	*	*	*	*	*
2001	0.6	*	*	*	1.6	*	*	*	*	*	*	*
2000	0.7	*	*	*	1.6	1.2	*	*	*	*	*	*
1999	0.8	*	*	*	2.2	0.9	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
All origins 3/, both sexes												
2006	1.2	1.2	2.8	0.6	1.5	1.1	1.1	1.1	1.0	1.1	1.5	1.4
2005	1.2	1.6	3.0	0.6	1.5	1.0	1.1	1.2	0.9	1.0	1.4	1.8
2004	1.1	1.5	2.7	0.7	1.4	1.0	1.0	1.1	0.9	1.0	1.2	1.6
2003	1.1	1.4	2.9	0.7	1.4	0.9	1.0	1.0	0.9	1.1	1.3	1.8
2002	1.2	1.6	2.9	0.8	1.5	1.1	1.2	0.9	0.8	0.9	1.3	1.7
2001	1.2	1.7	3.0	0.8	1.5	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000	1.2	2.0	3.2	0.9	1.6	1.1	1.1	0.9	0.9	1.0	1.3	1.5
1999	1.3	1.8	3.2	0.9	1.7	1.1	1.1	1.0	0.9	1.0	1.3	1.6
All origins 3/, male												
2006	1.9	1.3	3.5	0.9	2.5	1.9	1.7	1.8	1.5	1.7	2.4	2.4
2005	1.9	1.3	4.1	0.9	2.7	1.6	1.9	1.9	1.5	1.6	2.3	3.9
2004	1.8	1.8	3.2	0.9	2.4	1.6	1.7	1.8	1.5	1.7	2.2	2.9
2003	1.8	1.7	3.7	0.9	2.5	1.6	1.8	1.6	1.6	1.7	2.0	3.3
2002	1.9	1.9	3.9	1.2	2.8	1.9	2.0	1.5	1.3	1.6	2.1	3.6
2001	1.8	1.5	3.7	1.1	2.7	1.6	1.7	1.6	1.3	1.5	2.0	3.5
2000	2.0	2.2	4.0	1.3	2.9	1.8	1.8	1.6	1.5	1.6	1.8	2.8
1999	2.0	2.0	4.0	1.3	3.0	2.0	1.8	1.7	1.5	1.6	2.3	2.3
All origins 3/, female												
2006	0.5	1.2	2.1	0.3	0.3	0.2	0.4	0.5	0.5	0.7	0.8	0.9
2005	0.5	1.8	2.0	0.3	0.3	0.3	0.4	0.5	0.3	0.6	0.8	0.9
2004	0.5	1.2	2.1	0.4	0.3	0.3	0.3	0.4	0.4	0.5	0.6	1.0
2003	0.5	1.2	2.0	0.3	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1
2002	0.5	1.2	1.9	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.8	0.9
2001	0.5	1.9	2.2	0.5	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999	0.5	1.6	2.4	0.5	0.3	0.2	0.3	0.3	0.4	0.6	0.6	1.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
Hispanic, both sexes												
2006	1.1	*	2.5	0.5	2.0	1.1	0.7	0.6	*	*	*	*
2005	1.2	*	2.9	0.4	2.2	1.0	0.8	0.8	0.9	*	*	*
2004	1.0	*	2.3	0.4	1.4	1.0	1.0	0.7	*	*	*	*
2003	1.1	*	2.5	0.6	1.6	1.2	0.9	0.7	*	*	*	*
2002	1.2	*	2.1	0.4	2.1	1.1	0.9	0.8	*	*	*	*
2001	1.1	*	2.6	0.6	1.5	1.1	1.0	0.7	*	*	*	*
2000	1.2	*	2.5	0.6	2.2	1.0	0.8	0.6	*	*	*	*
1999	1.4	*	2.7	0.7	2.3	1.3	0.9	0.8	*	*	*	*
Hispanic, male												
2006	1.8	*	3.1	0.8	3.6	1.9	1.2	1.0	*	*	*	*
2005	1.9	*	4.0	*	3.8	1.7	1.4	1.5	*	*	*	*
2004	1.7	*	2.7	*	2.5	1.7	1.6	1.4	*	*	*	*
2003	1.9	*	4.0	0.9	2.9	2.0	1.5	1.3	*	*	*	*
2002	2.0	*	3.0	0.6	3.7	2.0	1.6	1.5	*	*	*	*
2001	1.8	*	3.5	0.8	2.7	1.8	1.7	1.2	*	*	*	*
2000	1.9	*	2.9	0.6	3.8	1.7	1.4	*	*	*	*	*
1999	2.3	*	2.9	1.1	4.1	2.2	1.5	1.6	*	*	*	*
Hispanic, female												
2006	0.4	*	1.8	*	*	*	*	*	*	*	*	*
2005	0.4	*	1.8	*	*	*	*	*	*	*	*	*
2004	0.4	*	1.8	*	*	*	*	*	*	*	*	*
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	1.7	*	*	*	*	*	*	*	*	*
2000	0.5	*	2.2	0.6	*	*	*	*	*	*	*	*
1999	0.5	*	2.5	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
Non-Hispanic 4/, both sexes												
2006	1.2	1.1	2.9	0.7	1.3	1.1	1.1	1.2	1.0	1.2	1.5	1.4
2005	1.2	1.5	3.1	0.7	1.4	0.9	1.2	1.2	0.9	1.1	1.4	1.8
2004	1.1	1.4	2.8	0.7	1.4	1.0	1.0	1.1	1.0	1.0	1.2	1.6
2003	1.1	1.5	3.0	0.7	1.3	0.8	1.1	1.0	1.0	1.1	1.3	1.8
2002	1.2	1.4	3.1	0.9	1.4	1.1	1.2	0.9	0.8	0.9	1.3	1.7
2001	1.1	1.6	3.1	0.9	1.5	0.9	1.0	0.9	0.8	0.9	1.3	1.7
2000	1.2	2.1	3.4	1.0	1.5	1.1	1.1	1.0	0.9	1.0	1.3	1.5
1999	1.2	1.7	3.3	0.9	1.5	1.1	1.1	1.0	0.9	1.0	1.3	1.7
Non-Hispanic 4/, male												
2006	1.9	*	3.5	1.0	2.3	1.9	1.8	1.9	1.6	1.7	2.5	2.4
2005	1.9	1.3	4.1	1.0	2.5	1.6	2.0	1.9	1.5	1.6	2.4	3.8
2004	1.8	1.8	3.4	1.0	2.4	1.6	1.7	1.8	1.5	1.7	2.2	2.7
2003	1.8	1.7	3.7	1.0	2.4	1.5	1.8	1.6	1.6	1.8	2.1	3.4
2002	1.9	1.6	4.1	1.3	2.6	1.8	2.0	1.5	1.3	1.6	2.0	3.6
2001	1.8	1.3	3.8	1.2	2.6	1.6	1.7	1.6	1.3	1.5	2.1	3.4
2000	2.0	2.4	4.3	1.4	2.7	1.8	1.9	1.6	1.6	1.6	1.8	2.8
1999	2.0	1.7	4.2	1.4	2.7	1.9	1.8	1.7	1.5	1.6	2.3	2.4
Non-Hispanic 4/, female												
2006	0.6	*	2.2	0.3	0.3	0.2	0.5	0.5	0.5	0.7	0.8	0.9
2005	0.5	1.8	2.0	0.4	0.3	0.3	0.4	0.5	0.3	0.6	0.8	0.9
2004	0.5	*	2.2	0.4	0.3	0.3	0.3	0.4	0.4	0.5	0.6	1.1
2003	0.5	*	2.3	0.3	0.2	0.2	0.3	0.4	0.4	0.5	0.7	1.1
2002	0.5	*	2.1	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.8	0.9
2001	0.5	1.9	2.4	0.5	0.3	0.2	0.4	0.3	0.4	0.4	0.7	1.0
2000	0.5	1.9	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.4	0.6	0.6	1.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
Non-Hispanic white, both sexes												
2006 5/	1.2	1.3	3.1	0.5	1.1	1.0	1.2	1.2	1.0	1.1	1.5	1.4
2005 5/	1.2	1.8	3.4	0.5	1.2	0.9	1.2	1.2	0.9	1.0	1.3	1.8
2004 5/	1.1	1.5	3.2	0.5	1.2	0.9	1.0	1.1	0.9	1.0	1.2	1.5
2003 5/	1.1	1.4	3.3	0.5	1.2	0.8	1.1	1.0	0.9	1.1	1.3	1.7
2002	1.1	1.4	3.3	0.5	1.2	1.0	1.2	0.9	0.8	0.9	1.2	1.6
2001	1.1	1.5	3.5	0.5	1.3	0.8	1.0	0.9	0.8	0.9	1.2	1.7
2000	1.1	1.7	3.7	0.7	1.3	1.0	1.1	0.9	0.9	1.0	1.3	1.4
1999	1.1	1.3	3.6	0.5	1.3	1.0	1.1	1.0	0.9	1.0	1.2	1.6
Non-Hispanic white, male												
2006 5/	1.8	*	3.6	0.7	1.9	1.8	1.8	1.9	1.5	1.8	2.5	2.6
2005 5/	1.8	*	4.6	0.6	2.1	1.5	1.8	1.9	1.4	1.5	2.3	3.9
2004 5/	1.7	1.8	3.8	0.7	2.1	1.5	1.6	1.7	1.4	1.6	2.2	2.6
2003 5/	1.7	*	4.0	0.7	2.1	1.3	1.8	1.5	1.5	1.8	2.1	3.4
2002	1.7	*	4.3	0.7	2.2	1.7	1.9	1.4	1.2	1.5	2.0	3.7
2001	1.6	*	4.3	0.7	2.2	1.4	1.5	1.5	1.2	1.5	2.1	3.5
2000	1.8	1.9	4.7	1.0	2.2	1.7	1.8	1.5	1.6	1.5	1.8	2.7
1999	1.8	*	4.5	0.7	2.2	1.6	1.8	1.7	1.5	1.4	2.2	2.4
Non-Hispanic white, female												
2006 5/	0.6	*	2.5	0.3	0.4	0.2	0.5	0.5	0.5	0.6	0.8	0.9
2005 5/	0.5	2.3	2.3	0.3	0.2	0.3	0.5	0.5	0.3	0.6	0.7	0.8
2004 5/	0.5	*	2.5	0.2	0.2	0.3	0.3	0.4	0.4	0.5	0.5	1.0
2003 5/	0.5	*	2.7	0.3	0.2	0.2	0.3	0.4	0.4	0.5	0.7	0.9
2002	0.5	*	2.2	0.3	0.3	0.3	0.4	0.4	0.4	0.3	0.7	0.7
2001	0.5	1.7	2.6	0.4	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000	0.5	*	2.6	0.4	0.3	0.3	0.4	0.3	0.3	0.5	0.9	0.9
1999	0.5	*	2.6	0.3	0.3	0.3	0.4	0.4	0.4	0.6	0.6	1.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
Non-Hispanic black, both sexes												
2006 5/	1.4	*	2.6	1.3	2.0	1.2	1.0	1.1	1.2	1.4	*	*
2005 5/	1.5	*	2.2	1.4	2.1	1.1	1.3	1.2	1.0	1.5	*	*
2004 5/	1.4	*	1.9	1.6	1.9	1.1	1.1	1.3	1.0	*	*	*
2003 5/	1.3	*	2.1	1.4	1.7	1.1	0.9	1.0	1.2	*	*	*
2002	1.7	*	2.6	2.1	2.1	1.4	1.3	1.3	0.8	*	*	*
2001	1.6	*	2.0	2.1	2.3	1.2	1.3	1.2	1.3	*	*	*
2000	1.7	4.3	2.3	2.1	2.3	1.1	1.4	1.3	*	*	*	*
1999	1.8	3.7	2.5	2.5	2.5	1.6	1.1	0.9	1.2	1.3	*	*
Non-Hispanic black, male												
2006 5/	2.4	*	3.7	1.9	3.7	2.4	1.9	2.1	2.2	*	*	*
2005 5/	2.6	*	2.8	2.3	3.9	2.1	2.4	2.3	2.2	*	*	*
2004 5/	2.4	*	2.8	2.3	3.4	2.0	2.0	2.3	1.8	*	*	*
2003 5/	2.3	*	3.0	2.0	3.1	2.2	1.7	2.0	2.4	*	*	*
2002	2.9	*	3.5	3.3	4.1	2.7	2.5	2.3	*	*	*	*
2001	2.8	*	2.4	3.1	4.4	2.5	2.2	2.1	2.4	*	*	*
2000	2.8	*	2.8	3.1	4.4	2.1	2.6	2.4	*	*	*	*
1999	3.2	*	3.4	3.8	4.6	3.2	2.1	1.9	2.1	*	*	*
Non-Hispanic black, female												
2006 5/	0.4	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.4	*	1.6	*	*	*	*	*	*	*	*	*
2004 5/	0.5	*	*	0.9	*	*	*	*	*	*	*	*
2003 5/	0.4	*	*	0.7	*	*	*	*	*	*	*	*
2002	0.5	*	*	0.8	*	*	*	*	*	*	*	*
2001	0.5	*	*	1.0	*	*	*	*	*	*	*	*
2000	0.6	*	1.8	1.0	*	*	*	*	*	*	*	*
1999	0.6	*	*	1.1	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
All origins 3/, both sexes												
2006	1.0	0.7	1.2	0.4	0.4	0.5	0.7	1.1	1.5	2.1	3.4	4.3
2005	1.1	0.8	1.3	0.5	0.4	0.6	0.8	1.2	1.3	2.1	3.6	5.0
2004	1.1	0.6	1.4	0.6	0.4	0.6	0.7	1.2	1.4	1.9	3.5	5.6
2003	1.2	0.7	1.4	0.5	0.5	0.6	0.9	1.3	1.4	2.3	3.6	5.4
2002	1.1	0.9	1.4	0.6	0.5	0.7	0.8	1.1	1.3	2.1	3.4	4.8
2001	1.2	1.2	1.5	0.6	0.5	0.6	1.0	1.1	1.5	2.2	3.4	5.7
2000	1.2	1.0	1.9	0.6	0.5	0.6	0.9	1.2	1.5	2.2	3.8	6.2
1999	1.2	1.1	2.0	0.6	0.5	0.7	0.9	1.0	1.3	2.3	4.0	5.9
All origins 3/, male												
2006	1.2	*	1.4	0.4	0.5	0.6	0.8	1.4	1.9	2.6	4.5	6.3
2005	1.3	*	1.5	0.6	0.5	0.7	1.0	1.6	1.7	2.6	4.6	7.0
2004	1.3	*	1.5	0.7	0.5	0.8	0.9	1.6	1.8	2.5	4.5	7.2
2003	1.4	*	1.6	0.5	0.6	0.8	1.2	1.6	1.9	2.9	4.4	7.4
2002	1.4	1.0	1.6	0.7	0.6	0.8	1.1	1.5	1.7	2.9	4.5	7.0
2001	1.4	1.5	1.8	0.7	0.7	0.8	1.4	1.5	1.9	2.8	4.1	8.3
2000	1.5	1.0	2.2	0.7	0.6	0.8	1.2	1.6	2.1	2.8	5.0	9.4
1999	1.5	1.1	2.2	0.7	0.7	1.0	1.3	1.4	1.7	2.9	5.1	9.3
All origins 3/, female												
2006	0.8	*	1.0	0.5	0.3	0.3	0.5	0.8	1.0	1.7	2.7	3.4
2005	0.9	*	1.0	0.5	0.3	0.4	0.5	0.8	1.0	1.6	2.9	4.0
2004	0.9	*	1.3	0.5	0.4	0.4	0.5	0.8	1.1	1.4	2.9	4.9
2003	1.0	1.1	1.2	0.5	0.4	0.5	0.6	0.9	1.0	1.8	3.0	4.5
2002	0.8	*	1.2	0.5	0.3	0.5	0.5	0.8	0.9	1.4	2.6	3.9
2001	0.9	1.0	1.1	0.5	0.4	0.4	0.6	0.7	1.1	1.8	2.9	4.6
2000	0.9	*	1.5	0.6	0.4	0.4	0.6	0.7	1.0	1.7	3.0	4.9
1999	0.9	1.1	1.7	0.6	0.4	0.4	0.5	0.6	1.0	1.7	3.2	4.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
Hispanic, both sexes												
2006	0.5	*	0.8	0.3	0.4	0.4	0.4	0.6	1.0	*	*	*
2005	0.5	*	0.8	*	*	0.3	0.4	0.6	*	*	3.4	*
2004	0.4	*	*	*	*	0.3	*	*	*	1.7	3.8	*
2003	0.5	*	0.8	0.3	*	0.3	0.4	0.8	*	*	*	*
2002	0.6	*	1.2	0.4	0.4	*	0.5	0.6	*	1.7	*	*
2001	0.5	*	*	0.3	*	0.3	0.5	*	*	*	*	*
2000	0.5	*	0.7	*	0.3	*	0.5	*	*	*	*	*
1999	0.6	*	1.4	0.4	0.4	*	0.4	*	*	*	*	*
Hispanic, male												
2006	0.7	*	*	*	*	0.6	0.6	0.9	*	*	*	*
2005	0.6	*	*	*	*	*	*	*	*	*	*	*
2004	0.5	*	*	*	*	*	*	*	*	*	*	*
2003	0.6	*	*	*	*	*	*	1.4	*	*	*	*
2002	0.7	*	*	*	*	*	0.7	*	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*	*	*
1999	0.7	*	1.6	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.3	*	*	*	*	*	*	*	*	*	*	*
2005	0.4	*	*	*	*	*	*	*	*	*	*	*
2004	0.4	*	*	*	*	*	*	*	*	*	*	*
2003	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
Non-Hispanic 4/, both sexes												
2006	1.1	0.8	1.3	0.5	0.4	0.5	0.7	1.2	1.5	2.2	3.6	4.5
2005	1.2	0.8	1.4	0.6	0.4	0.6	0.8	1.2	1.4	2.1	3.6	5.1
2004	1.2	0.7	1.6	0.7	0.5	0.7	0.8	1.3	1.5	1.9	3.5	5.6
2003	1.3	0.8	1.6	0.6	0.5	0.7	1.0	1.3	1.5	2.4	3.6	5.4
2002	1.2	1.0	1.5	0.7	0.5	0.8	0.9	1.2	1.3	2.1	3.4	4.8
2001	1.3	1.3	1.7	0.7	0.6	0.7	1.0	1.1	1.5	2.3	3.4	5.7
2000	1.3	1.0	2.2	0.7	0.5	0.7	0.9	1.2	1.5	2.2	3.8	6.3
1999	1.3	1.2	2.1	0.7	0.5	0.8	1.0	1.0	1.4	2.3	4.0	6.0
Non-Hispanic 4/, male												
2006	1.3	*	1.5	0.4	0.5	0.7	0.9	1.5	2.0	2.7	4.6	6.5
2005	1.4	*	1.6	0.7	0.5	0.8	1.1	1.6	1.7	2.7	4.7	7.2
2004	1.4	*	1.7	0.8	0.6	0.9	1.0	1.7	1.9	2.5	4.5	7.3
2003	1.5	*	1.9	0.6	0.6	0.8	1.3	1.7	1.9	3.0	4.5	7.3
2002	1.5	*	1.7	0.8	0.7	1.0	1.2	1.5	1.7	2.9	4.6	7.0
2001	1.6	1.5	2.0	0.8	0.8	0.9	1.5	1.6	2.0	2.8	4.2	8.3
2000	1.6	*	2.6	0.8	0.7	0.9	1.3	1.7	2.1	2.8	5.0	9.3
1999	1.6	*	2.3	0.8	0.7	1.1	1.4	1.4	1.7	3.0	5.1	9.2
Non-Hispanic 4/, female												
2006	0.9	*	1.2	0.5	0.4	0.4	0.6	0.8	1.1	1.8	2.8	3.5
2005	0.9	*	1.1	0.5	0.4	0.5	0.5	0.8	1.0	1.7	2.8	4.1
2004	1.0	*	1.5	0.6	0.4	0.4	0.5	0.8	1.1	1.4	2.9	4.9
2003	1.0	1.3	1.3	0.5	0.4	0.6	0.7	1.0	1.0	1.8	3.1	4.5
2002	0.9	*	1.2	0.6	0.3	0.5	0.5	0.8	0.9	1.4	2.6	3.8
2001	1.0	*	1.3	0.5	0.4	0.5	0.6	0.7	1.1	1.9	2.9	4.6
2000	1.0	*	1.7	0.7	0.4	0.5	0.6	0.7	1.0	1.7	3.1	5.0
1999	1.0	*	1.8	0.6	0.4	0.5	0.6	0.6	1.1	1.8	3.3	4.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
Non-Hispanic white, both sexes												
2006 5/	1.1	*	0.9	0.3	0.4	0.5	0.7	1.1	1.4	2.1	3.1	3.9
2005 5/	1.1	*	1.0	0.5	0.4	0.6	0.8	1.2	1.1	1.9	3.1	4.3
2004 5/	1.1	*	1.3	0.5	0.5	0.6	0.7	1.1	1.3	1.6	3.0	4.9
2003 5/	1.2	*	1.3	0.4	0.5	0.7	1.0	1.2	1.3	2.1	3.1	4.7
2002	1.0	*	1.0	0.4	0.4	0.7	0.8	1.1	1.1	1.8	3.0	3.9
2001	1.1	1.1	1.3	0.5	0.5	0.6	0.9	1.0	1.3	2.0	2.9	4.7
2000	1.1	*	1.5	0.5	0.5	0.6	0.8	1.1	1.3	1.8	3.1	5.1
1999	1.1	0.9	1.5	0.5	0.5	0.8	0.9	0.9	1.2	1.9	3.2	4.7
Non-Hispanic white, male												
2006 5/	1.3	*	1.2	0.3	0.5	0.6	0.8	1.4	1.8	2.4	4.0	6.2
2005 5/	1.3	*	1.4	0.5	0.5	0.7	1.1	1.5	1.4	2.3	4.1	6.1
2004 5/	1.3	*	1.4	0.6	0.6	0.9	0.9	1.5	1.6	2.0	3.9	6.5
2003 5/	1.4	*	1.6	0.4	0.7	0.8	1.3	1.5	1.7	2.6	3.6	6.5
2002	1.3	*	1.1	0.6	0.6	0.9	1.1	1.4	1.4	2.4	4.2	5.7
2001	1.4	*	1.7	0.6	0.7	0.8	1.3	1.3	1.7	2.3	3.5	6.9
2000	1.4	*	1.8	0.6	0.6	0.8	1.1	1.6	1.7	2.3	4.1	8.1
1999	1.4	*	1.7	0.6	0.7	1.1	1.3	1.2	1.5	2.3	4.2	7.8
Non-Hispanic white, female												
2006 5/	0.9	*	0.7	0.4	0.3	0.4	0.6	0.8	1.0	1.7	2.5	2.8
2005 5/	0.9	*	0.7	0.4	0.3	0.4	0.5	0.8	0.9	1.6	2.4	3.5
2004 5/	0.9	*	1.2	0.4	0.4	0.4	0.5	0.8	1.0	1.2	2.4	4.2
2003 5/	1.0	*	1.0	0.4	0.4	0.6	0.6	0.9	0.9	1.6	2.8	3.9
2002	0.8	*	0.8	0.3	0.3	0.5	0.5	0.7	0.8	1.3	2.2	3.2
2001	0.9	*	0.9	0.4	0.3	0.4	0.6	0.6	1.0	1.7	2.5	3.8
2000	0.9	*	1.1	0.5	0.3	0.4	0.5	0.7	0.8	1.5	2.4	3.9
1999	0.8	*	1.3	0.5	0.3	0.4	0.5	0.6	0.8	1.5	2.6	3.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
Non-Hispanic black, both sexes												
2006 5/	1.8	*	3.1	1.1	0.6	0.9	1.1	2.0	2.8	4.2	9.6	12.5
2005 5/	2.0	*	2.8	1.3	0.6	1.0	1.2	2.0	3.4	4.4	9.4	14.5
2004 5/	2.1	*	2.9	1.7	0.6	1.0	1.2	2.5	3.5	5.0	9.7	15.3
2003 5/	2.1	*	3.1	1.5	0.6	0.9	1.2	2.6	2.9	5.4	10.3	14.9
2002	2.1	*	3.7	1.5	0.8	1.2	1.1	2.2	3.3	5.0	8.7	16.0
2001	2.3	*	3.4	1.3	0.7	1.2	1.8	2.4	3.6	6.1	9.9	17.8
2000	2.6	*	5.1	1.6	0.7	1.2	1.7	2.4	4.2	6.1	13.0	20.7
1999	2.6	*	4.7	1.6	0.7	1.0	1.8	2.2	3.4	7.0	13.5	22.0
Non-Hispanic black, male												
2006 5/	2.1	*	3.1	0.9	*	1.2	1.4	2.6	4.0	5.6	13.9	*
2005 5/	2.4	*	2.8	1.4	0.6	1.1	1.8	3.1	5.2	6.4	13.5	21.9
2004 5/	2.7	*	3.0	2.0	0.7	1.3	1.7	3.9	4.5	7.7	13.4	*
2003 5/	2.5	*	3.4	1.5	*	1.1	1.5	3.4	4.4	7.7	16.2	*
2002	2.6	*	4.3	1.6	1.0	1.4	1.7	3.1	4.6	7.8	11.1	26.3
2001	3.0	*	3.7	1.6	0.8	1.4	2.8	3.9	5.1	9.0	13.2	26.6
2000	3.2	*	5.8	1.7	0.9	1.6	2.4	3.5	6.3	9.0	17.9	28.2
1999	3.1	*	5.3	1.8	0.9	1.2	2.7	3.3	4.3	10.4	18.4	29.7
Non-Hispanic black, female												
2006 5/	1.6	*	3.1	1.2	*	*	0.7	1.4	1.7	3.1	7.1	12.0
2005 5/	1.5	*	2.7	1.2	*	0.9	*	1.1	2.0	2.9	7.2	11.5
2004 5/	1.7	*	2.9	1.3	*	*	0.7	1.3	2.6	3.0	7.6	13.7
2003 5/	1.8	*	2.8	1.4	0.7	*	0.9	2.0	1.7	3.8	7.0	13.2
2002	1.6	*	3.0	1.3	*	1.0	*	1.4	2.2	3.1	7.3	12.1
2001	1.8	*	3.1	1.0	*	1.1	1.0	1.1	2.4	4.0	8.1	14.5
2000	2.0	*	4.3	1.4	*	0.9	1.2	1.4	2.5	4.0	10.2	17.9
1999	2.1	*	4.1	1.4	*	0.8	1.1	1.3	2.8	4.7	10.7	19.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
All origins 3/, both sexes												
2006	9.2	*	0.2	0.1	6.9	13.0	17.3	19.0	7.6	2.6	2.6	3.8
2005	8.0	0.5	0.1	0.1	5.9	10.9	15.3	16.4	6.6	2.3	2.4	3.5
2004	7.1	*	*	0.1	5.4	9.1	14.6	14.5	5.4	2.3	2.4	3.4
2003	6.7	0.5	0.3	0.1	4.9	8.6	14.0	13.3	4.9	2.2	2.2	3.5
2002	6.1	0.6	0.2	0.1	4.1	7.8	13.4	11.7	4.1	2.1	2.3	3.9
2001	4.9	*	0.2	0.1	3.4	6.3	11.2	9.1	3.2	1.6	2.2	3.5
2000	4.5	*	0.2	0.1	3.0	6.0	10.3	8.1	2.8	1.5	2.0	4.3
1999	4.4	*	0.2	0.1	2.5	5.9	10.1	7.8	2.8	1.6	2.2	3.4
All origins 3/, male												
2006	12.6	*	*	0.2	10.6	18.9	23.0	24.9	9.9	2.9	2.9	4.6
2005	10.9	*	*	0.1	9.0	15.6	20.2	21.7	8.7	2.7	3.0	4.6
2004	9.6	*	*	0.1	8.2	13.1	19.0	19.1	6.6	2.6	2.7	4.2
2003	9.2	*	0.3	0.1	7.4	12.4	18.8	18.0	6.3	2.7	2.8	4.0
2002	8.5	1.0	*	0.1	6.3	11.5	18.4	15.9	5.4	2.5	2.7	5.1
2001	7.1	*	0.3	0.1	5.3	9.3	15.7	13.0	4.1	2.0	2.7	3.8
2000	6.6	*	0.3	0.1	4.6	9.0	15.1	12.0	3.4	1.8	2.1	6.2
1999	6.5	*	*	0.1	3.8	9.0	14.9	11.7	3.9	2.1	2.4	4.4
All origins 3/, female												
2006	5.9	*	*	0.1	3.0	6.9	11.6	13.4	5.5	2.3	2.4	3.4
2005	5.1	*	*	0.1	2.7	6.0	10.5	11.3	4.7	2.0	2.0	3.0
2004	4.7	*	*	0.1	2.4	4.9	10.3	10.0	4.3	2.0	2.2	3.0
2003	4.3	*	0.3	0.2	2.2	4.7	9.3	8.8	3.7	1.8	1.8	3.3
2002	3.7	*	*	*	1.9	4.0	8.4	7.6	3.0	1.7	2.1	3.4
2001	2.9	*	*	0.1	1.4	3.3	6.7	5.2	2.3	1.2	1.9	3.3
2000	2.5	*	*	*	1.3	2.8	5.6	4.4	2.3	1.3	1.9	3.6
1999	2.3	*	*	*	1.1	2.6	5.3	4.0	1.8	1.3	2.0	3.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
Hispanic, both sexes												
2006	5.2	*	*	*	3.3	7.0	10.6	12.3	7.0	1.5	*	*
2005	4.7	*	*	*	3.4	6.2	9.9	10.9	5.5	2.2	*	*
2004	4.2	*	*	*	2.8	4.9	9.2	9.7	5.4	2.4	*	*
2003	4.5	*	*	*	3.1	5.7	10.3	10.5	4.4	1.8	*	*
2002	4.3	*	*	*	2.9	5.0	10.1	10.1	4.0	2.2	*	*
2001	3.6	*	*	*	2.2	4.9	8.7	7.7	3.5	*	*	*
2000	3.7	*	*	*	2.0	5.3	9.6	8.0	3.1	*	*	*
1999	4.4	*	*	*	2.3	6.1	11.9	9.8	3.5	*	*	*
Hispanic, male												
2006	7.8	*	*	*	4.8	10.3	15.9	18.1	11.7	*	*	*
2005	7.3	*	*	*	5.2	9.3	15.0	16.8	9.3	*	*	*
2004	6.2	*	*	*	4.2	7.6	13.5	14.2	8.4	*	*	*
2003	6.9	*	*	*	4.8	8.6	15.6	16.4	7.4	*	*	*
2002	6.6	*	*	*	4.4	7.8	15.4	15.8	6.4	*	*	*
2001	5.6	*	*	*	3.1	7.4	13.6	12.9	5.8	*	*	*
2000	6.0	*	*	*	3.2	8.4	15.6	13.6	4.4	*	*	*
1999	7.3	*	*	*	3.6	9.8	19.4	16.4	6.6	*	*	*
Hispanic, female												
2006	2.4	*	*	*	1.5	2.9	4.8	6.4	2.8	*	*	*
2005	2.0	*	*	*	1.4	2.3	4.3	5.0	1.9	*	*	*
2004	1.9	*	*	*	1.3	1.6	4.4	5.1	2.6	*	*	*
2003	1.9	*	*	*	1.2	2.2	4.5	4.6	*	*	*	*
2002	1.7	*	*	*	1.1	1.6	4.3	4.4	*	*	*	*
2001	1.5	*	*	*	1.1	1.9	3.4	2.6	*	*	*	*
2000	1.3	*	*	*	*	1.6	3.1	2.5	*	*	*	*
1999	1.4	*	*	*	0.8	1.7	3.9	3.3	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
Non-Hispanic 4/, both sexes												
2006	9.8	*	0.2	0.1	7.7	14.5	18.4	19.7	7.6	2.6	2.6	3.9
2005	8.5	*	*	0.1	6.4	12.0	16.2	16.9	6.7	2.3	2.5	3.5
2004	7.6	*	*	0.1	5.9	10.1	15.4	14.9	5.4	2.3	2.4	3.4
2003	7.0	*	0.3	0.2	5.2	9.2	14.5	13.5	4.9	2.2	2.2	3.5
2002	6.3	0.6	0.2	0.1	4.4	8.4	13.8	11.8	4.1	2.1	2.3	4.1
2001	5.1	*	0.2	0.1	3.6	6.6	11.4	9.1	3.1	1.6	2.3	3.6
2000	4.6	*	0.2	0.1	3.1	6.0	10.3	8.0	2.8	1.5	2.0	4.4
1999	4.3	*	0.2	0.1	2.5	5.8	9.7	7.5	2.7	1.6	2.2	3.5
Non-Hispanic 4/, male												
2006	13.4	*	*	0.2	11.9	21.3	24.2	25.5	9.7	2.9	2.9	4.7
2005	11.5	*	*	0.2	9.8	17.3	21.0	22.1	8.5	2.6	3.1	4.7
2004	10.2	*	*	0.1	9.1	14.5	19.8	19.5	6.4	2.6	2.7	4.2
2003	9.5	*	0.3	0.2	7.9	13.3	19.2	17.9	6.1	2.6	2.8	4.0
2002	8.8	*	*	0.1	6.6	12.3	18.7	15.9	5.2	2.4	2.6	5.3
2001	7.2	*	*	0.1	5.7	9.6	15.9	12.9	3.9	2.0	2.8	3.9
2000	6.6	*	*	0.1	4.9	9.1	14.8	11.7	3.3	1.7	2.1	6.2
1999	6.3	*	*	0.1	3.8	8.8	14.2	11.1	3.6	2.0	2.5	4.4
Non-Hispanic 4/, female												
2006	6.4	*	*	*	3.3	7.8	12.7	14.1	5.8	2.4	2.4	3.5
2005	5.6	*	*	*	2.9	6.8	11.5	12.0	4.9	2.1	2.0	3.0
2004	5.1	*	*	0.1	2.7	5.6	11.1	10.5	4.4	2.0	2.2	3.1
2003	4.6	*	*	0.2	2.4	5.2	9.9	9.2	3.8	1.9	1.8	3.3
2002	4.0	*	*	*	2.0	4.5	8.9	7.9	3.0	1.8	2.1	3.5
2001	3.1	*	*	*	1.5	3.5	7.1	5.4	2.3	1.2	1.9	3.4
2000	2.7	*	*	*	1.4	3.0	5.9	4.4	2.3	1.3	2.0	3.6
1999	2.4	*	*	*	1.2	2.8	5.4	4.0	1.8	1.3	2.0	3.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
Non-Hispanic white, both sexes												
2006 5/	10.5	*	*	0.1	9.5	17.2	19.8	19.7	7.1	2.4	2.6	3.8
2005 5/	9.0	*	*	0.1	7.9	14.0	17.6	16.9	6.1	2.1	2.4	3.6
2004 5/	8.2	*	*	0.1	7.3	11.8	16.8	15.0	5.0	2.2	2.4	3.5
2003 5/	7.4	*	*	0.1	6.4	10.6	15.5	13.4	4.8	2.0	2.3	3.5
2002	6.6	*	*	0.1	5.3	9.6	14.5	11.4	3.7	2.0	2.3	4.1
2001	5.2	*	*	0.1	4.3	7.3	11.7	8.3	2.7	1.5	2.1	3.4
2000	4.6	*	*	0.1	3.7	6.5	10.4	7.2	2.5	1.4	2.1	4.1
1999	4.2	*	*	0.1	2.8	6.0	9.6	6.7	2.4	1.5	2.3	3.3
Non-Hispanic white, male												
2006 5/	14.2	*	*	0.2	14.7	24.9	25.8	24.8	8.2	2.4	2.9	4.8
2005 5/	12.2	*	*	0.2	12.1	20.1	22.7	21.5	7.3	2.1	3.0	4.9
2004 5/	10.9	*	*	0.2	11.2	17.1	21.4	19.3	5.4	2.3	2.5	4.3
2003 5/	10.1	*	*	*	9.8	15.3	20.3	17.5	5.6	2.1	2.9	3.6
2002	9.1	*	*	0.2	8.0	13.9	19.6	15.0	4.3	2.2	2.5	5.4
2001	7.3	*	*	*	6.7	10.7	16.3	11.5	3.1	1.9	2.5	3.8
2000	6.6	*	*	*	5.8	9.8	14.9	10.3	2.7	1.5	2.2	5.8
1999	6.1	*	*	*	4.3	9.2	14.0	9.6	2.9	1.8	2.5	4.6
Non-Hispanic white, female												
2006 5/	7.0	*	*	*	4.0	9.4	13.8	14.7	5.9	2.4	2.5	3.3
2005 5/	6.1	*	*	*	3.5	7.9	12.4	12.4	5.0	2.2	2.1	2.9
2004 5/	5.5	*	*	*	3.2	6.5	12.1	10.8	4.5	2.1	2.3	3.1
2003 5/	4.9	*	*	*	2.9	5.9	10.6	9.4	4.0	1.9	1.9	3.5
2002	4.3	*	*	*	2.5	5.2	9.3	7.8	3.1	1.9	2.2	3.5
2001	3.1	*	*	*	1.7	3.8	7.1	5.2	2.4	1.2	1.9	3.2
2000	2.7	*	*	*	1.5	3.2	5.8	4.2	2.3	1.4	2.0	3.4
1999	2.4	*	*	*	1.3	2.8	5.1	3.8	1.9	1.2	2.1	2.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
Non-Hispanic black, both sexes												
2006 5/	9.2	*	*	*	2.2	8.3	17.9	26.1	14.6	4.7	2.8	5.6
2005 5/	7.8	*	*	*	1.8	7.3	14.6	22.2	13.1	4.5	2.9	*
2004 5/	6.7	*	*	*	1.8	5.5	13.4	19.2	10.0	3.8	3.4	*
2003 5/	6.4	*	*	*	1.4	5.8	13.9	18.3	7.6	4.1	*	*
2002	6.4	*	*	*	1.5	5.8	14.1	18.3	8.6	2.5	2.8	*
2001	6.0	*	*	*	1.6	5.3	13.4	17.0	6.4	2.4	3.7	6.6
2000	5.7	*	*	*	1.4	5.7	13.2	15.7	5.5	2.5	*	8.3
1999	5.8	*	*	*	1.7	6.3	13.2	15.6	6.0	3.0	*	*
Non-Hispanic black, male												
2006 5/	13.3	*	*	*	3.1	13.0	24.9	38.8	25.0	7.8	*	*
2005 5/	11.0	*	*	*	2.4	10.4	19.0	32.7	22.4	8.2	*	*
2004 5/	9.1	*	*	*	2.6	7.8	17.0	27.5	16.5	6.4	*	*
2003 5/	9.1	*	*	*	2.1	8.3	19.2	26.6	12.9	7.5	*	*
2002	9.3	*	*	*	2.2	8.7	19.8	27.1	15.2	4.5	*	*
2001	8.6	*	*	*	2.3	7.6	18.2	27.0	11.0	3.7	6.0	*
2000	8.4	*	*	*	1.8	8.5	19.0	25.3	9.5	4.3	*	*
1999	8.8	*	*	*	2.8	9.3	19.5	25.9	11.0	4.4	*	*
Non-Hispanic black, female												
2006 5/	5.4	*	*	*	1.3	3.8	11.8	15.2	6.2	2.4	*	*
2005 5/	4.9	*	*	*	1.2	4.5	10.8	13.3	5.7	*	*	*
2004 5/	4.4	*	*	*	1.1	3.4	10.1	12.0	4.7	*	*	*
2003 5/	4.1	*	*	*	0.7	3.6	9.3	11.1	3.4	*	*	*
2002	3.8	*	*	*	0.7	3.2	9.1	10.8	3.3	*	*	*
2001	3.6	*	*	*	0.9	3.2	9.2	8.4	2.6	*	*	*
2000	3.2	*	*	*	1.0	3.1	8.0	7.5	2.4	*	*	*
1999	3.1	*	*	*	*	3.5	7.7	6.7	1.9	2.1	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
All origins 3/, both sexes												
2006	5.8	21.5	1.6	0.6	1.6	2.1	2.9	4.2	5.2	9.1	25.9	89.3
2005	6.3	19.5	1.5	0.6	1.7	2.3	3.1	4.7	5.8	10.2	28.6	95.1
2004	5.8	19.2	1.6	0.6	1.5	2.1	2.9	4.2	5.0	8.8	27.0	95.5
2003	5.9	16.9	1.8	0.5	1.6	2.0	3.0	3.9	5.1	9.6	27.5	102.5
2002	6.0	16.8	1.6	0.7	1.6	2.1	3.0	3.7	5.2	9.8	28.8	105.6
2001	6.2	16.6	1.8	0.6	1.6	2.0	3.1	4.2	5.3	10.0	30.5	115.0
2000	6.2	15.0	1.7	0.6	1.7	2.2	3.2	4.0	5.6	9.9	29.8	117.1
1999	6.6	13.8	1.9	0.8	1.7	2.2	3.2	4.1	5.6	11.0	32.9	128.9
All origins 3/, male												
2006	7.0	25.2	1.7	0.8	2.6	3.5	4.8	6.8	7.7	12.7	32.9	99.4
2005	7.4	21.1	1.8	0.9	2.8	3.9	4.9	7.4	8.6	13.8	35.9	107.1
2004	6.8	21.7	2.0	0.9	2.6	3.4	4.6	6.6	7.5	11.8	33.6	109.8
2003	6.9	19.5	2.2	0.8	2.6	3.4	4.8	6.0	7.8	13.1	34.9	120.2
2002	6.9	18.8	2.0	0.9	2.6	3.4	4.8	6.0	7.9	13.4	36.0	118.6
2001	7.2	18.9	2.1	0.9	2.6	3.4	5.0	6.6	7.7	13.2	38.4	134.0
2000	7.2	16.8	2.1	0.9	2.8	3.8	5.4	6.6	8.3	13.0	36.9	135.2
1999	7.6	16.2	2.3	1.1	2.9	3.7	5.3	6.6	8.3	14.8	40.9	152.7
All origins 3/, female												
2006	4.7	17.6	1.5	0.3	0.5	0.7	1.0	1.8	2.8	6.1	21.2	84.5
2005	5.1	17.9	1.2	0.4	0.6	0.7	1.2	2.0	3.1	7.1	23.6	89.6
2004	4.8	16.5	1.1	0.3	0.4	0.7	1.2	1.8	2.7	6.4	22.6	89.1
2003	4.9	14.1	1.3	0.3	0.5	0.7	1.2	1.9	2.6	6.7	22.6	94.7
2002	5.0	14.6	1.2	0.4	0.5	0.6	1.2	1.5	2.7	6.8	24.0	100.0
2001	5.3	14.2	1.5	0.4	0.6	0.6	1.1	1.9	3.1	7.3	25.4	107.0
2000	5.2	13.2	1.4	0.4	0.5	0.6	1.1	1.5	3.0	7.4	25.1	109.8
1999	5.6	11.3	1.5	0.5	0.5	0.7	1.2	1.7	3.1	8.0	27.7	119.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
Hispanic, both sexes												
2006	2.7	12.4	1.4	0.4	1.5	2.0	2.5	3.1	4.4	6.4	14.7	37.2
2005	2.8	11.0	1.2	0.4	1.7	2.4	2.6	3.8	4.1	6.8	15.1	38.1
2004	2.5	9.8	1.2	0.4	1.7	1.9	2.4	3.3	3.5	5.2	14.6	44.2
2003	2.5	8.2	1.6	0.4	1.6	2.0	2.2	3.0	3.9	6.3	16.9	37.3
2002	2.5	8.2	1.3	0.5	1.5	1.9	2.5	2.8	3.4	6.2	15.3	48.2
2001	2.7	9.5	1.4	0.3	1.5	1.8	2.7	3.6	3.8	7.7	16.0	63.8
2000	2.7	7.5	1.4	0.4	1.6	2.2	2.9	2.7	4.0	6.2	16.6	70.3
1999	2.6	8.6	1.9	0.5	1.6	1.6	2.4	2.8	4.2	8.7	15.0	65.9
Hispanic, male												
2006	3.9	13.7	1.4	0.6	2.5	3.2	4.6	5.4	6.6	9.7	18.3	45.1
2005	4.0	11.5	1.5	0.5	2.9	4.1	4.4	6.4	6.4	9.7	18.0	46.1
2004	3.5	11.7	1.6	0.6	2.8	2.9	3.9	5.6	5.1	6.9	17.5	43.4
2003	3.6	9.1	1.9	0.5	2.9	3.4	3.7	4.7	6.4	9.0	24.9	44.6
2002	3.6	10.1	1.8	0.6	2.5	3.3	4.2	4.8	5.6	8.4	21.3	70.3
2001	3.7	9.6	1.8	*	2.6	3.0	4.6	6.1	5.6	10.8	20.6	68.6
2000	3.8	7.9	1.5	*	2.9	3.7	5.2	4.5	6.6	8.7	19.7	88.7
1999	3.6	9.0	2.5	*	2.6	2.7	4.0	4.8	6.2	13.2	15.6	84.1
Hispanic, female												
2006	1.5	11.0	1.5	*	*	0.6	*	*	2.4	3.7	12.2	33.0
2005	1.5	10.5	*	*	*	*	*	1.3	2.0	4.5	13.0	33.9
2004	1.5	7.9	*	*	*	0.6	*	*	2.0	3.8	12.6	44.7
2003	1.3	7.3	1.3	*	*	*	*	1.3	*	4.1	11.3	33.6
2002	1.2	6.1	*	*	*	*	*	*	*	4.4	11.2	37.3
2001	1.6	9.5	*	*	*	*	*	1.2	2.1	5.4	13.0	61.4
2000	1.4	7.2	*	*	*	*	*	*	*	4.3	14.5	61.3
1999	1.6	8.1	*	*	*	*	*	*	2.4	5.2	14.6	57.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
Non-Hispanic 4/, both sexes												
2006	6.4	24.2	1.6	0.6	1.6	2.2	3.0	4.3	5.2	9.3	26.6	91.8
2005	6.8	21.9	1.6	0.7	1.7	2.3	3.1	4.7	5.8	10.4	29.3	97.7
2004	6.3	21.9	1.7	0.7	1.5	2.1	3.0	4.2	5.1	9.1	27.7	97.7
2003	6.4	19.2	1.8	0.6	1.5	2.0	3.1	4.0	5.2	9.8	28.0	105.1
2002	6.5	18.9	1.6	0.7	1.6	2.1	3.1	3.8	5.3	10.0	29.4	107.6
2001	6.7	18.3	1.9	0.7	1.6	2.0	3.1	4.2	5.4	10.1	31.1	116.6
2000	6.7	16.9	1.8	0.7	1.7	2.2	3.2	4.1	5.7	10.1	30.2	118.5
1999	7.1	15.0	1.9	0.8	1.7	2.3	3.3	4.2	5.6	11.1	33.6	130.7
Non-Hispanic 4/, male												
2006	7.5	28.6	1.7	0.8	2.6	3.6	4.8	6.9	7.8	12.8	33.6	102.2
2005	8.0	23.8	1.9	1.0	2.8	3.8	5.0	7.5	8.7	14.0	36.9	110.2
2004	7.4	24.6	2.1	1.0	2.5	3.5	4.7	6.7	7.6	12.1	34.5	113.1
2003	7.5	22.2	2.3	0.8	2.5	3.4	4.9	6.1	7.9	13.4	35.3	123.5
2002	7.5	21.0	2.0	0.9	2.6	3.4	4.9	6.1	8.1	13.7	36.6	120.4
2001	7.7	21.1	2.1	1.0	2.6	3.5	5.1	6.6	7.8	13.3	39.0	136.5
2000	7.7	19.0	2.2	1.0	2.7	3.8	5.3	6.7	8.4	13.2	37.5	136.9
1999	8.1	17.9	2.3	1.1	2.9	3.9	5.4	6.6	8.4	14.8	41.8	154.8
Non-Hispanic 4/, female												
2006	5.2	19.6	1.5	0.3	0.5	0.8	1.2	1.9	2.8	6.3	21.7	87.0
2005	5.7	19.9	1.3	0.4	0.7	0.8	1.3	2.0	3.2	7.3	24.2	92.0
2004	5.3	19.0	1.2	0.4	0.5	0.7	1.3	1.9	2.8	6.6	23.1	90.9
2003	5.4	16.0	1.3	0.3	0.6	0.7	1.3	1.9	2.7	6.8	23.1	97.0
2002	5.6	16.8	1.3	0.4	0.5	0.7	1.2	1.6	2.8	7.0	24.6	102.2
2001	5.8	15.4	1.6	0.4	0.6	0.6	1.2	2.0	3.2	7.4	25.9	108.4
2000	5.7	14.7	1.4	0.4	0.5	0.7	1.1	1.6	3.1	7.5	25.4	111.1
1999	6.2	12.0	1.5	0.5	0.5	0.7	1.2	1.8	3.1	8.1	28.2	121.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
Non-Hispanic white, both sexes												
2006 5/	6.9	18.9	1.5	0.6	1.7	2.3	3.1	4.3	5.2	9.1	27.3	96.0
2005 5/	7.2	18.6	1.6	0.7	1.8	2.5	3.2	4.5	5.6	10.0	29.1	100.2
2004 5/	6.9	18.4	1.6	0.7	1.7	2.3	3.1	4.3	5.0	8.9	27.9	101.3
2003 5/	6.9	15.5	1.7	0.6	1.6	2.2	3.2	3.8	5.0	9.6	28.3	109.1
2002	7.0	14.5	1.5	0.7	1.7	2.3	3.1	3.8	5.2	9.6	29.4	110.9
2001	7.2	14.5	1.7	0.7	1.8	2.1	3.1	4.1	5.2	9.8	31.2	119.5
2000	7.1	13.7	1.7	0.7	1.7	2.3	3.2	3.9	5.4	9.7	30.3	122.0
1999	7.6	12.1	1.7	0.9	1.9	2.4	3.4	3.9	5.2	10.6	33.1	134.3
Non-Hispanic white, male												
2006 5/	8.0	22.0	1.7	0.9	2.9	3.9	4.9	6.7	7.7	12.6	34.4	106.7
2005 5/	8.4	20.1	2.0	1.0	2.9	4.1	5.1	7.1	8.1	13.5	36.3	113.3
2004 5/	7.9	22.0	2.2	0.9	2.8	3.8	4.8	6.7	7.5	11.9	34.6	116.4
2003 5/	7.9	18.0	2.2	0.8	2.7	3.6	5.0	5.9	7.5	13.1	35.3	127.5
2002	7.9	17.0	1.8	0.9	2.9	3.7	5.1	6.0	7.7	13.0	36.3	123.1
2001	8.1	16.8	2.0	1.0	2.9	3.6	5.1	6.3	7.4	13.0	38.6	139.7
2000	8.0	16.0	2.1	1.1	2.9	4.0	5.3	6.3	7.9	12.6	36.8	139.7
1999	8.4	14.2	2.0	1.2	3.1	4.0	5.5	6.2	7.7	13.9	41.0	157.3
Non-Hispanic white, female												
2006 5/	5.8	15.6	1.4	0.3	0.5	0.7	1.2	1.9	2.8	6.0	22.3	91.0
2005 5/	6.2	17.0	1.2	0.4	0.6	0.8	1.2	1.9	3.2	7.0	24.2	94.2
2004 5/	5.9	14.7	0.9	0.4	0.5	0.7	1.4	2.0	2.7	6.4	23.3	94.5
2003 5/	6.0	12.9	1.2	0.3	0.5	0.7	1.3	1.8	2.6	6.7	23.6	101.0
2002	6.1	11.9	1.2	0.4	0.5	0.8	1.2	1.5	2.7	6.7	24.8	105.7
2001	6.4	12.1	1.5	0.4	0.7	0.6	1.1	1.9	3.1	7.1	26.3	111.2
2000	6.3	11.2	1.3	0.4	0.5	0.7	1.1	1.6	3.1	7.3	26.0	114.9
1999	6.8	9.8	1.5	0.5	0.6	0.7	1.3	1.7	2.8	7.9	28.0	125.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
Non-Hispanic black, both sexes												
2006 5/	5.1	46.4	2.2	0.7	1.1	1.9	3.1	5.5	6.8	12.9	24.9	62.6
2005 5/	6.2	38.8	1.7	0.7	1.6	2.1	3.4	7.1	8.8	15.8	38.4	87.4
2004 5/	4.9	38.0	1.9	0.6	1.0	2.0	3.0	4.8	6.6	11.9	31.3	71.4
2003 5/	5.1	35.7	2.1	0.5	1.3	1.8	3.4	5.5	7.5	13.1	27.8	74.1
2002	5.4	38.6	2.5	0.9	1.2	1.7	3.2	4.8	7.4	15.6	32.9	84.4
2001	5.5	35.5	2.6	0.6	1.1	2.0	3.3	5.7	8.0	14.2	34.2	96.5
2000	5.6	31.7	2.3	0.7	1.6	2.3	3.6	5.8	8.4	15.1	34.1	90.9
1999	6.3	29.5	2.6	0.7	1.3	2.5	3.5	6.7	10.2	18.2	43.1	106.7
Non-Hispanic black, male												
2006 5/	6.6	55.5	2.1	0.9	1.7	2.8	5.4	9.6	10.3	18.2	32.0	76.0
2005 5/	7.9	42.6	1.8	0.8	2.4	3.1	5.3	11.7	15.2	21.8	54.4	99.0
2004 5/	6.3	38.7	1.7	1.1	1.5	3.5	4.9	7.8	10.5	16.7	43.4	89.4
2003 5/	6.6	41.6	2.9	0.7	1.8	3.0	5.3	8.7	12.2	18.8	39.8	92.7
2002	6.8	40.0	3.3	1.2	1.8	2.9	4.8	7.6	12.1	22.1	46.6	107.2
2001	7.0	41.8	2.9	0.9	1.6	3.3	5.2	9.2	12.2	19.9	49.5	115.5
2000	7.5	32.8	2.7	1.0	2.6	3.8	6.1	10.3	14.2	20.8	53.7	124.5
1999	8.3	35.0	3.5	1.0	2.4	4.1	5.9	11.2	16.0	27.4	57.8	155.7
Non-Hispanic black, female												
2006 5/	3.7	37.0	2.3	*	*	1.0	1.1	2.1	4.0	9.1	20.8	57.2
2005 5/	4.7	34.7	1.6	*	0.7	1.1	1.8	3.2	3.7	11.6	29.4	82.7
2004 5/	3.7	37.3	2.0	*	*	*	1.3	2.2	3.5	8.4	24.5	64.3
2003 5/	3.7	29.7	*	*	0.8	0.8	1.6	2.8	3.7	9.0	21.0	66.9
2002	4.2	37.2	*	*	*	*	1.8	2.4	3.6	11.0	25.2	75.7
2001	4.2	28.9	2.3	*	*	0.7	1.6	2.7	4.7	10.3	25.6	89.4
2000	3.9	30.7	2.0	*	*	0.9	1.3	1.9	3.7	11.1	23.1	78.5
1999	4.5	23.9	*	*	*	1.1	1.4	2.8	5.5	11.8	34.8	88.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All origins 3/, both sexes												
2006	11.1	...	...	0.5	9.9	12.3	15.1	17.2	14.5	12.6	15.9	15.9
2005	11.0	...	...	0.7	10.0	12.4	14.9	16.5	13.9	12.6	16.9	16.9
2004	11.0	...	...	0.7	10.3	12.7	15.0	16.6	13.8	12.3	16.3	16.4
2003	10.8	...	...	0.6	9.7	12.7	14.9	15.9	13.8	12.7	16.4	16.9
2002	11.0	...	...	0.6	9.9	12.6	15.3	15.7	13.6	13.5	17.7	18.0
2001 6/	10.8	...	...	0.7	9.9	12.8	14.7	15.2	13.1	13.3	17.4	17.5
2000	10.4	...	...	0.7	10.2	12.0	14.5	14.4	12.1	12.5	17.6	19.6
1999	10.5	...	...	0.6	10.1	12.7	14.3	13.9	12.2	13.4	18.1	19.3
All origins 3/, male												
2006	17.8	...	...	0.7	16.2	19.7	23.2	26.2	22.7	22.7	33.3	43.2
2005	17.7	...	...	1.0	16.2	19.9	23.1	25.2	22.2	22.7	35.8	45.0
2004	17.7	...	...	0.9	16.8	20.4	23.0	24.8	22.1	22.6	34.8	45.0
2003	17.6	...	...	0.9	16.0	20.6	23.2	24.4	22.3	23.4	35.1	47.8
2002	17.9	...	...	0.9	16.5	20.5	23.7	24.4	22.2	24.7	38.1	50.7
2001 6/	17.6	...	...	1.0	16.6	21.0	23.1	23.4	21.1	24.6	37.8	51.1
2000	17.1	...	...	1.2	17.1	19.6	22.8	22.4	19.4	22.7	38.6	57.5
1999	17.1	...	...	0.9	16.8	20.7	22.4	21.5	19.8	24.7	38.8	57.1
All origins 3/, female												
2006	4.6	...	...	0.3	3.2	4.7	7.0	8.4	6.8	4.1	4.0	3.1
2005	4.5	...	...	0.3	3.5	4.7	6.8	8.0	6.1	4.0	4.0	4.0
2004	4.6	...	...	0.5	3.6	4.7	7.1	8.6	6.1	3.8	3.9	3.6
2003	4.3	...	...	0.3	3.0	4.6	6.6	7.7	5.9	3.8	4.0	3.3
2002	4.3	...	...	0.3	2.9	4.6	6.9	7.4	5.7	4.1	4.2	3.8
2001 6/	4.1	...	...	0.3	2.9	4.4	6.4	7.2	5.7	3.9	4.0	3.4
2000	4.0	...	...	0.3	3.0	4.3	6.4	6.7	5.4	4.0	4.0	4.2
1999	4.0	...	...	0.3	3.0	4.6	6.4	6.6	5.2	4.1	4.7	4.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Hispanic, both sexes												
2006	4.9	...	...	0.3	7.4	7.1	6.6	6.5	6.0	5.2	7.6	*
2005	5.1	...	...	0.5	7.7	6.8	7.4	6.6	6.0	5.9	8.8	9.0
2004	5.3	...	...	0.5	8.0	7.2	6.8	7.5	7.0	6.8	9.1	*
2003	5.0	...	...	0.6	7.0	6.8	6.8	7.2	6.9	6.5	8.0	10.9
2002	5.0	...	...	0.5	6.7	6.6	7.0	7.1	7.2	7.5	9.2	12.7
2001 6/	5.0	...	...	0.5	6.2	7.1	7.3	7.0	6.2	7.6	8.8	13.8
2000	5.1	...	...	0.4	6.8	6.5	7.5	7.8	6.0	7.8	11.7	*
1999	5.0	...	...	0.5	6.2	6.7	7.5	7.1	6.8	7.0	13.1	15.4
Hispanic, male												
2006	7.9	...	...	*	11.6	10.9	10.6	10.1	10.5	9.1	16.1	*
2005	8.3	...	...	0.7	12.1	10.7	11.8	11.0	10.2	10.7	18.7	*
2004	8.6	...	...	0.6	12.8	11.2	10.8	12.0	11.3	13.6	19.7	*
2003	8.3	...	...	0.8	11.2	10.8	11.0	11.5	13.0	13.1	18.0	*
2002	8.3	...	...	0.6	10.6	10.7	11.2	11.3	13.2	14.7	19.3	35.1
2001 6/	8.3	...	...	0.7	9.5	11.3	12.4	11.4	11.5	15.4	20.1	41.9
2000	8.4	...	...	0.6	10.9	10.6	12.1	12.5	11.1	15.6	26.1	*
1999	8.2	...	...	0.6	9.9	10.7	12.1	11.6	12.2	13.8	28.2	*
Hispanic, female												
2006	1.7	...	...	*	2.6	2.4	2.2	2.8	1.9	*	*	*
2005	1.7	...	...	*	2.7	2.1	2.4	2.2	2.1	*	*	*
2004	1.8	...	...	*	2.5	2.3	2.4	3.1	3.1	*	*	*
2003	1.5	...	...	*	2.2	1.9	2.2	2.9	*	*	*	*
2002	1.6	...	...	*	2.1	1.7	2.4	2.9	*	*	*	*
2001 6/	1.5	...	...	*	2.3	2.2	1.8	2.7	*	*	*	*
2000	1.5	...	...	*	2.0	1.7	2.6	3.2	*	*	*	*
1999	1.6	...	...	*	1.8	2.0	2.6	2.7	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Non-Hispanic 4/, both sexes												
2006	12.2	...	...	0.6	10.4	13.6	16.6	18.3	15.2	13.1	16.4	16.3
2005	12.0	...	...	0.7	10.5	13.8	16.2	17.5	14.5	13.1	17.3	17.2
2004	11.9	...	...	0.7	10.8	13.9	16.3	17.5	14.3	12.7	16.7	16.8
2003	11.7	...	...	0.6	10.2	14.0	16.0	16.7	14.3	13.1	16.9	17.1
2002	11.8	...	...	0.7	10.5	13.9	16.4	16.5	14.0	13.8	18.1	18.1
2001 6/	11.6	...	...	0.7	10.7	13.9	15.7	15.9	13.6	13.6	17.8	17.5
2000	11.1	...	...	0.8	10.8	13.0	15.4	15.0	12.5	12.7	17.9	19.9
1999	11.2	...	...	0.6	10.8	13.8	15.1	14.4	12.5	13.7	18.3	19.3
Non-Hispanic 4/, male												
2006	19.6	...	...	0.8	17.1	22.1	25.5	28.0	23.8	23.7	34.3	44.4
2005	19.3	...	...	1.1	17.0	22.3	25.0	26.8	23.1	23.6	36.6	45.9
2004	19.2	...	...	1.0	17.5	22.7	25.0	26.1	22.8	23.1	35.6	46.2
2003	19.1	...	...	0.9	16.9	22.9	25.1	25.6	22.9	24.1	36.0	48.6
2002	19.4	...	...	1.0	17.6	22.7	25.5	25.6	22.8	25.2	39.0	50.9
2001 6/	19.0	...	...	1.1	18.1	23.1	24.6	24.5	21.7	25.0	38.5	51.3
2000	18.3	...	...	1.3	18.3	21.4	24.1	23.2	19.9	23.1	39.1	58.4
1999	18.4	...	...	1.0	18.2	22.7	23.6	22.2	20.2	25.2	39.1	57.6
Non-Hispanic 4/, female												
2006	5.1	...	...	0.4	3.3	5.2	7.8	9.0	7.3	4.2	4.2	3.2
2005	4.9	...	...	0.4	3.6	5.3	7.5	8.6	6.4	4.1	4.1	4.1
2004	5.0	...	...	0.5	3.7	5.2	7.8	9.2	6.4	3.9	4.0	3.6
2003	4.6	...	...	0.3	3.2	5.2	7.2	8.1	6.3	4.0	4.1	3.3
2002	4.6	...	...	0.3	3.1	5.2	7.4	7.8	5.9	4.3	4.3	3.9
2001 6/	4.5	...	...	0.3	3.0	4.9	7.0	7.6	6.1	4.1	4.2	3.5
2000	4.3	...	...	0.3	3.1	4.8	6.8	7.0	5.7	4.1	4.1	4.3
1999	4.3	...	...	0.2	3.2	5.0	6.8	6.9	5.4	4.3	4.8	4.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Non-Hispanic white, both sexes												
2006 5/	13.9	...	...	0.6	11.2	15.3	19.1	21.0	17.1	14.5	17.8	17.4
2005 5/	13.6	...	...	0.7	11.4	15.3	18.5	20.0	16.2	14.4	18.7	18.3
2004 5/	13.5	...	...	0.7	11.6	15.4	18.7	19.9	15.8	13.8	17.9	17.9
2003 5/	13.3	...	...	0.6	10.9	15.5	18.3	19.1	15.8	14.3	18.2	18.4
2002	13.4	...	...	0.7	11.5	15.4	18.7	18.7	15.6	15.1	19.4	19.2
2001 6/	13.0	...	...	0.7	11.6	15.4	17.7	17.9	15.0	14.7	19.0	18.7
2000	12.5	...	...	0.9	11.6	14.2	17.2	16.7	13.9	13.8	19.0	21.3
1999	12.4	...	...	0.7	11.5	15.0	16.9	16.0	13.7	14.8	19.5	20.8
Non-Hispanic white, male												
2006 5/	22.3	...	...	0.8	18.5	24.4	29.0	31.7	26.3	25.9	37.2	47.6
2005 5/	22.0	...	...	1.1	18.4	24.5	28.3	30.2	25.7	26.0	39.5	49.4
2004 5/	21.6	...	...	0.9	19.0	24.9	28.4	29.3	24.9	24.9	38.0	49.8
2003 5/	21.6	...	...	1.0	18.2	25.0	28.2	29.0	25.1	26.0	38.6	52.5
2002	21.9	...	...	1.1	19.3	24.6	28.7	28.6	25.0	27.5	41.7	54.3
2001 6/	21.4	...	...	1.1	19.6	25.2	27.4	27.4	23.6	26.9	41.1	55.2
2000	20.4	...	...	1.4	19.5	23.1	26.6	25.5	21.8	24.7	41.7	62.7
1999	20.3	...	...	1.1	19.2	24.3	26.0	24.4	21.9	27.0	41.6	62.2
Non-Hispanic white, female												
2006 5/	5.9	...	...	0.4	3.5	6.1	9.2	10.5	8.3	4.5	4.4	3.4
2005 5/	5.6	...	...	0.3	3.9	5.9	8.7	9.9	7.2	4.4	4.3	4.0
2004 5/	5.7	...	...	0.5	4.0	5.6	9.0	10.6	7.1	4.3	4.2	3.6
2003 5/	5.3	...	...	0.2	3.3	5.8	8.3	9.3	7.0	4.3	4.4	3.4
2002	5.3	...	...	0.3	3.4	6.1	8.6	9.0	6.6	4.6	4.5	4.1
2001 6/	5.0	...	...	0.3	3.3	5.5	8.1	8.6	6.8	4.4	4.4	3.5
2000	4.9	...	...	0.4	3.3	5.2	7.8	7.9	6.4	4.6	4.2	4.6
1999	4.9	...	...	0.3	3.4	5.5	7.8	7.8	6.0	4.6	5.0	4.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Non-Hispanic black, both sexes												
2006 5/	5.1	...	...	0.6	6.4	8.7	7.5	6.0	4.9	4.2	4.4	5.8
2005 5/	5.2	...	...	0.7	6.8	9.0	7.6	6.3	4.7	4.5	5.4	*
2004 5/	5.3	...	...	0.8	7.3	9.7	6.7	5.9	5.8	4.5	6.2	*
2003 5/	5.3	...	...	0.7	7.3	9.6	7.4	5.7	4.9	4.6	4.5	*
2002	5.2	...	...	0.5	6.6	9.6	7.7	5.9	4.9	4.9	5.4	*
2001 6/	5.3	...	...	0.6	7.3	9.2	7.5	6.2	5.1	5.5	5.6	*
2000	5.5	...	...	0.6	8.4	9.0	7.7	6.5	4.2	5.3	5.9	*
1999	5.5	...	...	0.5	8.3	9.5	7.8	5.7	5.0	5.8	5.8	*
Non-Hispanic black, male												
2006 5/	9.1	...	...	0.9	10.9	16.1	13.4	10.6	9.1	8.8	11.7	*
2005 5/	8.9	...	...	0.8	11.9	15.5	13.0	10.5	8.0	8.4	12.6	*
2004 5/	9.2	...	...	0.9	12.4	16.8	11.4	9.7	11.2	10.1	15.4	*
2003 5/	9.0	...	...	1.0	12.5	17.1	12.5	9.3	8.5	8.5	11.3	*
2002	9.3	...	...	0.9	11.6	18.0	13.2	10.2	8.5	9.7	13.5	*
2001 6/	9.3	...	...	0.9	13.4	16.6	12.5	9.8	9.3	10.8	13.2	*
2000	9.6	...	...	1.0	14.6	16.1	13.3	11.1	7.5	11.4	12.3	*
1999	9.9	...	...	0.7	14.8	17.5	13.7	10.1	9.2	11.7	14.1	*
Non-Hispanic black, female												
2006 5/	1.4	...	...	*	1.8	1.9	2.2	2.2	1.6	*	*	*
2005 5/	1.8	...	...	*	1.7	2.9	2.8	2.7	2.1	*	*	*
2004 5/	1.8	...	...	*	2.2	3.3	2.6	2.5	1.5	*	*	*
2003 5/	1.8	...	...	*	2.0	2.7	2.9	2.7	2.0	*	*	*
2002	1.6	...	...	*	1.7	2.1	2.9	2.2	2.0	*	*	*
2001 6/	1.7	...	...	*	1.3	2.4	3.0	3.1	1.8	*	*	*
2000	1.7	...	...	*	2.3	2.6	2.7	2.5	1.6	*	*	*
1999	1.6	...	...	*	2.0	2.4	2.6	2.0	1.6	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
All origins 3/, both sexes												
2006	5.6	...	...	0.2	4.7	5.3	6.5	8.0	8.1	8.9	12.0	10.9
2005	5.7	...	...	0.2	4.7	5.7	6.5	8.2	8.1	9.0	12.6	11.2
2004	5.7	...	...	0.1	5.0	5.7	6.5	8.0	8.0	8.8	12.2	11.1
2003	5.8	...	...	0.2	5.0	6.0	6.6	8.0	8.3	9.3	12.4	11.9
2002	5.9	...	...	0.2	5.1	6.0	7.0	7.8	8.4	9.7	13.2	12.0
2001	5.9	...	...	0.2	5.3	6.5	6.7	7.7	8.2	9.6	13.1	12.2
2000	5.9	...	...	0.3	5.8	6.1	7.1	7.6	7.6	9.2	13.0	13.2
1999	5.9	...	...	0.3	6.0	6.5	6.9	7.3	7.8	9.7	13.1	12.6
All origins 3/, male												
2006	10.0	...	...	0.3	8.2	9.2	10.8	13.7	14.3	17.6	27.7	32.7
2005	10.2	...	...	0.3	8.2	9.7	11.0	14.1	14.6	17.6	29.3	33.8
2004	10.0	...	...	0.2	8.8	9.7	10.8	13.5	14.2	17.5	28.3	34.4
2003	10.4	...	...	0.3	8.9	10.4	11.2	13.8	14.9	18.4	28.7	36.6
2002	10.6	...	...	0.3	9.1	10.3	11.9	13.6	15.1	19.5	31.0	37.8
2001	10.6	...	...	0.3	9.4	11.2	11.4	13.3	14.6	19.2	30.7	38.9
2000	10.5	...	...	0.4	10.2	10.6	11.8	13.0	13.5	18.4	30.8	42.9
1999	10.6	...	...	0.4	10.5	11.2	11.5	12.5	14.0	19.5	31.1	41.6
All origins 3/, female												
2006	1.4	...	...	*	0.9	1.3	2.1	2.6	2.3	1.6	1.2	0.7
2005	1.4	...	...	*	0.9	1.5	2.0	2.4	2.1	1.7	1.3	0.8
2004	1.5	...	...	*	1.1	1.5	2.2	2.8	2.2	1.6	1.4	*
2003	1.4	...	...	*	1.0	1.4	2.1	2.5	2.2	1.6	1.5	0.9
2002	1.4	...	...	*	1.0	1.6	2.2	2.2	2.2	1.6	1.4	0.8
2001	1.5	...	...	0.1	1.0	1.6	2.1	2.4	2.4	1.7	1.5	1.0
2000	1.5	...	...	0.1	1.2	1.7	2.4	2.3	2.1	1.7	1.4	1.1
1999	1.5	...	...	0.1	1.2	1.8	2.3	2.3	2.1	1.7	1.4	0.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Hispanic, both sexes												
2006	1.8	...	...	*	3.1	2.6	2.2	2.1	2.2	2.0	4.3	*
2005	1.9	...	...	*	3.1	2.6	2.6	2.3	2.0	2.5	4.5	*
2004	2.1	...	...	*	3.7	2.7	2.4	3.0	2.7	3.5	5.0	*
2003	2.1	...	...	*	3.3	2.8	2.5	3.0	2.7	3.1	4.1	*
2002	2.2	...	...	*	3.3	2.7	2.5	2.7	3.4	4.4	4.7	*
2001	2.2	...	...	*	2.9	3.3	2.5	3.0	3.2	4.0	4.7	*
2000	2.3	...	...	*	3.4	2.8	2.8	4.0	2.7	4.6	5.3	*
1999	2.3	...	...	*	3.5	2.9	3.0	3.2	2.9	3.6	8.1	*
Hispanic, male												
2006	3.3	...	...	*	5.5	4.2	3.9	3.5	4.3	4.4	10.1	*
2005	3.4	...	...	*	5.4	4.3	4.4	4.3	4.1	5.2	11.0	*
2004	3.7	...	...	*	6.5	4.3	4.1	5.3	5.0	7.1	11.4	*
2003	3.8	...	...	*	5.6	4.8	4.3	5.3	5.6	7.0	9.6	*
2002	3.8	...	...	*	5.7	4.6	4.2	4.8	6.6	9.4	11.5	*
2001	3.8	...	...	*	4.8	5.5	4.5	5.3	6.6	8.8	11.6	*
2000	4.1	...	...	*	5.8	4.7	4.7	6.8	5.7	9.7	12.8	*
1999	4.0	...	...	*	5.8	4.8	5.0	5.5	5.9	7.7	19.8	*
Hispanic, female												
2006	0.3	...	...	*	*	0.6	*	*	*	*	*	*
2005	0.3	...	...	*	*	0.6	*	*	*	*	*	*
2004	0.4	...	...	*	0.6	0.7	*	*	*	*	*	*
2003	0.3	...	...	*	*	*	*	*	*	*	*	*
2002	0.4	...	...	*	*	*	*	*	*	*	*	*
2001	0.4	...	...	*	0.7	0.7	*	*	*	*	*	*
2000	0.4	...	...	*	*	*	*	*	*	*	*	*
1999	0.5	...	...	*	0.8	*	0.9	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Non-Hispanic 4/, both sexes												
2006	6.3	...	...	0.2	5.0	6.0	7.2	8.7	8.6	9.4	12.4	11.2
2005	6.4	...	...	0.2	5.0	6.4	7.2	8.8	8.6	9.4	13.1	11.4
2004	6.3	...	...	0.2	5.3	6.4	7.1	8.6	8.4	9.2	12.6	11.4
2003	6.4	...	...	0.2	5.4	6.7	7.2	8.5	8.7	9.7	12.8	12.1
2002	6.5	...	...	0.2	5.5	6.7	7.7	8.3	8.8	10.0	13.6	12.1
2001	6.5	...	...	0.2	5.8	7.1	7.3	8.1	8.6	9.9	13.5	12.4
2000	6.4	...	...	0.3	6.2	6.8	7.6	7.9	7.9	9.5	13.4	13.4
1999	6.4	...	...	0.3	6.5	7.2	7.3	7.7	8.2	10.1	13.3	12.7
Non-Hispanic 4/, male												
2006	11.2	...	...	0.3	8.8	10.5	12.1	14.8	15.2	18.5	28.8	33.8
2005	11.4	...	...	0.4	8.8	11.1	12.2	15.2	15.4	18.5	30.3	34.7
2004	11.1	...	...	0.3	9.3	11.1	11.9	14.4	15.0	18.2	29.3	35.6
2003	11.5	...	...	0.3	9.6	11.7	12.2	14.7	15.6	19.2	29.7	37.5
2002	11.7	...	...	0.4	9.8	11.6	13.1	14.4	15.7	20.1	31.9	38.4
2001	11.6	...	...	0.4	10.4	12.5	12.3	14.0	15.1	19.8	31.5	39.7
2000	11.4	...	...	0.5	11.1	11.7	12.7	13.5	14.0	18.9	31.6	43.8
1999	11.5	...	...	0.4	11.5	12.5	12.3	13.1	14.5	20.1	31.5	42.2
Non-Hispanic 4/, female												
2006	1.6	...	...	*	1.0	1.5	2.4	2.8	2.5	1.7	1.2	0.7
2005	1.6	...	...	*	1.0	1.6	2.2	2.7	2.3	1.8	1.4	0.8
2004	1.7	...	...	*	1.1	1.7	2.5	3.0	2.3	1.6	1.5	*
2003	1.6	...	...	*	1.0	1.6	2.3	2.6	2.3	1.7	1.6	0.9
2002	1.6	...	...	*	1.1	1.9	2.4	2.4	2.3	1.6	1.4	0.8
2001	1.6	...	...	0.1	1.1	1.8	2.4	2.5	2.5	1.8	1.6	1.0
2000	1.6	...	...	0.1	1.3	1.9	2.6	2.4	2.3	1.8	1.5	1.1
1999	1.6	...	...	*	1.3	1.9	2.4	2.4	2.3	1.8	1.5	0.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Non-Hispanic white, both sexes												
2006 5/	7.3	...	...	0.2	5.4	6.7	8.4	10.2	9.8	10.6	13.7	12.0
2005 5/	7.4	...	...	0.3	5.4	7.2	8.3	10.2	9.8	10.6	14.3	12.5
2004 5/	7.3	...	...	0.2	5.8	7.1	8.3	9.9	9.5	10.2	13.7	12.5
2003 5/	7.4	...	...	0.2	5.8	7.4	8.4	9.9	9.9	10.8	14.0	13.2
2002	7.5	...	...	0.2	5.9	7.5	8.8	9.5	9.9	11.2	14.7	13.1
2001	7.4	...	...	0.3	6.2	8.0	8.3	9.3	9.6	10.9	14.5	13.4
2000	7.3	...	...	0.4	6.7	7.4	8.6	8.9	8.7	10.4	14.5	14.5
1999	7.3	...	...	0.3	6.8	7.9	8.3	8.6	9.0	11.0	14.3	13.7
Non-Hispanic white, male												
2006 5/	13.0	...	...	0.3	9.5	11.6	13.9	17.1	17.0	20.6	31.4	36.3
2005 5/	13.2	...	...	0.5	9.4	12.3	13.9	17.3	17.4	20.6	32.9	37.9
2004 5/	12.8	...	...	0.3	10.1	12.1	13.7	16.5	16.7	19.9	31.5	39.0
2003 5/	13.2	...	...	0.4	10.3	12.7	14.1	16.8	17.4	21.2	32.1	41.0
2002	13.4	...	...	0.4	10.5	12.6	14.9	16.3	17.4	22.2	34.3	41.3
2001	13.2	...	...	0.4	11.0	13.8	13.9	15.9	16.7	21.6	33.8	43.0
2000	12.9	...	...	0.6	11.8	12.7	14.3	15.1	15.3	20.4	34.2	47.6
1999	12.9	...	...	0.5	12.0	13.4	13.7	14.6	15.9	21.8	33.8	45.7
Non-Hispanic white, female												
2006 5/	1.9	...	...	*	1.1	1.7	3.0	3.4	2.9	2.0	1.4	0.7
2005 5/	1.8	...	...	*	1.1	2.0	2.7	3.2	2.6	1.9	1.5	0.9
2004 5/	1.9	...	...	*	1.3	2.0	3.0	3.5	2.7	1.9	1.6	*
2003 5/	1.8	...	...	*	1.1	2.0	2.8	3.1	2.7	1.9	1.7	1.0
2002	1.8	...	...	*	1.2	2.3	2.8	2.8	2.7	1.8	1.6	0.9
2001	1.9	...	...	*	1.2	2.1	2.7	2.9	2.9	1.9	1.7	1.1
2000	1.9	...	...	*	1.4	2.1	3.0	2.8	2.6	2.0	1.5	1.2
1999	1.9	...	...	*	1.4	2.3	2.9	2.7	2.6	2.0	1.6	1.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Non-Hispanic black, both sexes												
2006 5/	2.6	...	...	*	3.8	4.8	3.3	2.5	2.5	2.9	3.8	*
2005 5/	2.7	...	...	*	3.9	4.7	3.4	2.8	2.4	3.3	4.0	*
2004 5/	2.7	...	...	*	4.1	5.0	2.8	2.4	3.1	3.6	4.9	*
2003 5/	2.7	...	...	*	4.4	5.4	2.7	2.5	2.8	3.4	4.0	*
2002	2.9	...	...	*	4.2	5.3	3.6	2.9	2.8	3.3	4.3	*
2001	3.0	...	...	*	5.1	5.0	3.5	2.8	2.7	3.8	5.1	*
2000	3.1	...	...	*	5.4	5.2	3.5	3.2	3.0	3.8	3.6	*
1999	3.2	...	...	*	5.8	5.1	3.6	3.0	3.5	4.0	4.8	*
Non-Hispanic black, male												
2006 5/	5.0	...	...	*	6.9	9.2	6.3	4.8	5.1	6.4	10.3	*
2005 5/	5.1	...	...	*	7.2	8.8	6.6	5.4	4.6	6.4	10.4	*
2004 5/	5.1	...	...	*	7.2	9.5	5.3	4.5	6.0	8.6	13.1	*
2003 5/	5.1	...	...	*	8.0	10.5	5.1	4.8	5.3	6.8	9.8	*
2002	5.5	...	...	*	7.7	10.5	6.5	5.5	5.5	7.0	11.4	*
2001	5.6	...	...	*	9.6	9.5	6.4	4.9	5.5	7.9	12.0	*
2000	5.7	...	...	*	9.9	9.6	6.4	5.6	5.7	9.0	8.3	*
1999	6.1	...	...	*	10.6	10.0	6.9	5.6	7.1	8.9	12.2	*
Non-Hispanic black, female												
2006 5/	0.5	...	...	*	0.7	0.8	*	*	*	*	*	*
2005 5/	0.5	...	...	*	*	0.8	*	*	*	*	*	*
2004 5/	0.5	...	...	*	0.9	1.0	*	*	*	*	*	*
2003 5/	0.5	...	...	*	0.7	*	*	*	*	*	*	*
2002	0.5	...	...	*	0.7	*	1.0	*	*	*	*	*
2001	0.6	...	...	*	*	0.9	1.0	1.0	*	*	*	*
2000	0.7	...	...	*	1.0	1.1	0.8	1.1	*	*	*	*
1999	0.6	...	...	*	1.0	0.8	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
All origins 3/, both sexes												
2006	5.5	...	...	0.4	5.2	7.0	8.6	9.1	6.4	3.7	3.9	5.0
2005	5.3	...	...	0.5	5.3	6.8	8.4	8.3	5.7	3.6	4.2	5.7
2004	5.3	...	...	0.6	5.3	7.0	8.5	8.5	5.8	3.5	4.1	5.4
2003	5.0	...	...	0.4	4.6	6.7	8.3	7.8	5.5	3.5	4.0	5.1
2002	5.0	...	...	0.4	4.7	6.6	8.2	7.9	5.2	3.8	4.5	6.0
2001 6/	4.8	...	...	0.5	4.6	6.3	8.0	7.4	4.9	3.7	4.3	5.3
2000	4.5	...	...	0.5	4.4	5.9	7.5	6.8	4.6	3.2	4.6	6.5
1999	4.5	...	...	0.3	4.1	6.2	7.5	6.6	4.4	3.7	5.0	6.7
All origins 3/, male												
2006	7.8	...	...	0.5	7.9	10.5	12.4	12.5	8.4	5.1	5.6	10.5
2005	7.5	...	...	0.6	8.0	10.2	12.1	11.1	7.6	5.1	6.5	11.2
2004	7.6	...	...	0.7	8.0	10.6	12.2	11.3	7.8	5.1	6.5	10.7
2003	7.3	...	...	0.6	7.1	10.2	12.1	10.6	7.3	5.0	6.4	11.1
2002	7.3	...	...	0.6	7.4	10.2	11.8	10.8	7.1	5.2	7.1	12.9
2001 6/	7.1	...	...	0.7	7.2	9.8	11.7	10.2	6.5	5.4	7.1	12.2
2000	6.6	...	...	0.7	6.9	9.0	10.9	9.4	6.0	4.3	7.8	14.6
1999	6.6	...	...	0.5	6.3	9.5	10.9	8.9	5.8	5.3	7.7	15.5
All origins 3/, female												
2006	3.2	...	...	0.3	2.3	3.4	4.9	5.8	4.5	2.4	2.8	2.4
2005	3.1	...	...	0.3	2.6	3.2	4.8	5.5	4.0	2.3	2.7	3.1
2004	3.1	...	...	0.4	2.5	3.2	4.9	5.9	3.9	2.2	2.5	3.0
2003	2.8	...	...	0.2	2.1	3.2	4.5	5.2	3.7	2.2	2.5	2.4
2002	2.9	...	...	0.2	2.0	3.0	4.7	5.2	3.5	2.5	2.8	3.0
2001 6/	2.6	...	...	0.2	1.9	2.8	4.3	4.8	3.4	2.3	2.5	2.4
2000	2.5	...	...	0.2	1.8	2.7	4.0	4.4	3.3	2.3	2.6	3.2
1999	2.5	...	...	0.1	1.8	2.8	4.1	4.3	3.0	2.4	3.3	3.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
Hispanic, both sexes												
2006	3.1	...	...	*	4.3	4.5	4.4	4.4	3.8	3.1	3.3	*
2005	3.2	...	...	0.4	4.6	4.2	4.8	4.3	4.0	3.4	4.3	*
2004	3.2	...	...	0.5	4.3	4.5	4.4	4.5	4.3	3.3	4.1	*
2003	2.9	...	...	0.5	3.8	3.9	4.3	4.2	4.2	3.4	3.9	*
2002	2.9	...	...	0.4	3.4	3.9	4.5	4.4	3.8	3.1	4.5	*
2001 6/	2.8	...	...	0.5	3.3	3.9	4.8	4.0	3.0	3.6	4.1	*
2000	2.8	...	...	0.3	3.4	3.7	4.7	3.8	3.2	3.3	6.3	*
1999	2.7	...	...	*	2.7	3.8	4.5	3.8	3.9	3.5	5.0	*
Hispanic, male												
2006	4.6	...	...	*	6.0	6.7	6.6	6.6	6.2	4.7	*	*
2005	4.9	...	...	0.6	6.7	6.5	7.4	6.7	6.1	5.4	7.7	*
2004	4.9	...	...	0.6	6.4	6.9	6.8	6.6	6.3	6.5	8.2	*
2003	4.5	...	...	0.6	5.6	6.0	6.7	6.2	7.5	6.1	8.4	*
2002	4.4	...	...	*	4.9	6.1	7.0	6.5	6.5	5.4	*	*
2001 6/	4.4	...	...	0.6	4.7	5.8	7.9	6.1	4.9	6.6	*	*
2000	4.3	...	...	*	5.1	5.9	7.3	5.7	5.4	5.9	13.3	*
1999	4.2	...	...	*	4.2	5.9	7.1	6.0	6.4	6.2	*	*
Hispanic, female												
2006	1.4	...	...	*	2.2	1.9	1.9	2.2	1.6	*	*	*
2005	1.4	...	...	*	2.2	1.5	1.9	1.8	2.0	*	*	*
2004	1.4	...	...	*	1.9	1.5	1.9	2.4	2.5	*	*	*
2003	1.2	...	...	*	1.6	1.5	1.7	2.3	*	*	*	*
2002	1.2	...	...	*	1.6	1.3	1.8	2.4	*	*	*	*
2001 6/	1.2	...	...	*	1.6	1.6	1.4	2.0	*	*	*	*
2000	1.1	...	...	*	1.4	1.1	1.9	2.0	*	*	*	*
1999	1.1	...	...	*	1.0	1.4	1.7	1.7	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
Non-Hispanic 4/, both sexes												
2006	5.9	...	...	0.4	5.4	7.6	9.4	9.6	6.6	3.7	4.0	5.1
2005	5.6	...	...	0.5	5.5	7.4	9.0	8.7	5.8	3.6	4.2	5.8
2004	5.7	...	...	0.6	5.5	7.5	9.2	8.9	5.9	3.5	4.1	5.4
2003	5.3	...	...	0.4	4.8	7.3	8.8	8.2	5.5	3.4	4.0	5.1
2002	5.3	...	...	0.4	5.0	7.2	8.7	8.2	5.3	3.8	4.5	5.9
2001 6/	5.1	...	...	0.5	4.8	6.8	8.4	7.7	5.0	3.7	4.3	5.1
2000	4.8	...	...	0.5	4.6	6.2	7.8	7.1	4.6	3.2	4.5	6.5
1999	4.7	...	...	0.4	4.4	6.6	7.8	6.8	4.4	3.7	5.0	6.6
Non-Hispanic 4/, male												
2006	8.4	...	...	0.5	8.3	11.6	13.5	13.2	8.6	5.2	5.5	10.6
2005	8.0	...	...	0.7	8.2	11.2	12.9	11.6	7.7	5.1	6.3	11.3
2004	8.1	...	...	0.7	8.3	11.6	13.1	11.8	7.9	4.9	6.4	10.6
2003	7.7	...	...	0.6	7.4	11.2	12.8	11.0	7.3	4.9	6.3	11.2
2002	7.7	...	...	0.7	7.8	11.1	12.4	11.1	7.0	5.1	7.1	12.5
2001 6/	7.5	...	...	0.7	7.7	10.6	12.2	10.5	6.6	5.3	7.0	11.6
2000	6.9	...	...	0.8	7.2	9.6	11.4	9.7	6.0	4.2	7.5	14.6
1999	6.9	...	...	0.6	6.7	10.2	11.3	9.1	5.7	5.1	7.6	15.4
Non-Hispanic 4/, female												
2006	3.5	...	...	0.3	2.3	3.7	5.4	6.2	4.8	2.5	2.9	2.5
2005	3.4	...	...	0.3	2.6	3.6	5.2	5.9	4.1	2.4	2.7	3.2
2004	3.4	...	...	0.5	2.6	3.5	5.3	6.2	4.0	2.3	2.5	3.0
2003	3.1	...	...	0.2	2.1	3.5	4.9	5.5	3.9	2.2	2.6	2.4
2002	3.1	...	...	0.2	2.0	3.3	5.1	5.4	3.6	2.6	2.9	3.1
2001 6/	2.9	...	...	0.2	1.9	3.1	4.7	5.1	3.5	2.3	2.6	2.5
2000	2.7	...	...	0.2	1.9	2.9	4.3	4.6	3.4	2.4	2.6	3.2
1999	2.7	...	...	0.1	1.9	3.0	4.4	4.5	3.1	2.5	3.3	3.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
Non-Hispanic white, both sexes												
2006 5/	6.6	...	...	0.4	5.8	8.6	10.7	10.8	7.3	3.8	4.1	5.4
2005 5/	6.2	...	...	0.4	6.0	8.1	10.2	9.8	6.4	3.8	4.3	5.7
2004 5/	6.3	...	...	0.5	5.9	8.3	10.4	9.9	6.3	3.6	4.2	5.4
2003 5/	5.9	...	...	0.4	5.1	8.1	9.9	9.2	6.0	3.5	4.2	5.2
2002	6.0	...	...	0.5	5.6	7.9	9.8	9.2	5.7	3.9	4.7	6.1
2001 6/	5.6	...	...	0.4	5.4	7.5	9.4	8.6	5.3	3.8	4.5	5.3
2000	5.2	...	...	0.5	4.9	6.8	8.6	7.8	5.1	3.3	4.5	6.8
1999	5.2	...	...	0.4	4.6	7.1	8.6	7.4	4.6	3.8	5.1	7.0
Non-Hispanic white, male												
2006 5/	9.3	...	...	0.5	9.0	12.8	15.1	14.6	9.2	5.4	5.8	11.2
2005 5/	8.8	...	...	0.6	9.0	12.2	14.4	12.8	8.3	5.4	6.6	11.5
2004 5/	8.9	...	...	0.6	8.9	12.8	14.8	12.8	8.2	5.0	6.5	10.8
2003 5/	8.4	...	...	0.6	7.9	12.3	14.1	12.2	7.7	4.9	6.5	11.6
2002	8.5	...	...	0.7	8.8	12.0	13.8	12.3	7.6	5.3	7.3	13.0
2001 6/	8.2	...	...	0.7	8.6	11.5	13.5	11.5	6.9	5.3	7.3	12.3
2000	7.5	...	...	0.8	7.7	10.3	12.4	10.4	6.5	4.3	7.5	15.2
1999	7.4	...	...	0.6	7.2	10.8	12.3	9.8	6.0	5.3	7.8	16.5
Non-Hispanic white, female												
2006 5/	4.0	...	...	0.3	2.4	4.3	6.3	7.1	5.4	2.5	3.0	2.7
2005 5/	3.7	...	...	0.2	2.8	4.0	6.0	6.7	4.6	2.5	2.8	3.1
2004 5/	3.7	...	...	0.4	2.7	3.6	6.1	7.1	4.5	2.5	2.6	3.0
2003 5/	3.4	...	...	0.2	2.2	3.9	5.6	6.2	4.3	2.4	2.7	2.4
2002	3.5	...	...	0.2	2.2	3.8	5.8	6.1	3.9	2.7	2.9	3.2
2001 6/	3.2	...	...	0.2	2.0	3.4	5.3	5.7	3.9	2.4	2.7	2.4
2000	3.0	...	...	0.2	1.9	3.1	4.8	5.2	3.8	2.5	2.6	3.4
1999	3.0	...	...	*	2.0	3.3	4.9	5.1	3.4	2.6	3.4	3.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
Non-Hispanic black, both sexes												
2006 5/	2.4	...	...	0.5	2.6	3.9	4.2	3.5	2.4	1.3	*	*
2005 5/	2.6	...	...	0.6	2.9	4.3	4.2	3.5	2.3	1.2	*	*
2004 5/	2.6	...	...	0.6	3.2	4.7	3.9	3.5	2.7	*	*	*
2003 5/	2.5	...	...	0.6	2.9	4.2	4.7	3.2	2.1	1.3	*	*
2002	2.4	...	...	0.4	2.4	4.4	4.2	3.0	2.1	1.6	*	*
2001 6/	2.3	...	...	0.4	2.3	4.2	3.9	3.4	2.4	1.7	*	*
2000	2.4	...	...	0.5	3.0	3.9	4.2	3.3	1.3	1.4	2.3	*
1999	2.3	...	...	0.4	2.5	4.4	4.2	2.7	1.5	1.8	*	*
Non-Hispanic black, male												
2006 5/	4.0	...	...	0.8	4.0	6.9	7.0	5.8	4.0	*	*	*
2005 5/	3.9	...	...	0.7	4.7	6.7	6.4	5.1	3.4	*	*	*
2004 5/	4.1	...	...	0.7	5.1	7.3	6.1	5.3	5.2	*	*	*
2003 5/	3.9	...	...	0.9	4.4	6.6	7.4	4.5	3.3	*	*	*
2002	3.8	...	...	0.6	3.8	7.5	6.7	4.7	3.0	*	*	*
2001 6/	3.7	...	...	0.7	3.8	7.1	6.1	4.9	3.7	3.0	*	*
2000	3.9	...	...	0.9	4.7	6.5	6.8	5.4	*	*	*	*
1999	3.8	...	...	*	4.1	7.5	6.8	4.5	2.1	*	*	*
Non-Hispanic black, female												
2006 5/	1.0	...	...	*	1.1	1.0	1.7	1.6	1.2	*	*	*
2005 5/	1.4	...	...	*	1.2	2.1	2.2	2.1	1.4	*	*	*
2004 5/	1.3	...	...	*	1.3	2.3	2.0	1.9	*	*	*	*
2003 5/	1.3	...	...	*	1.3	2.0	2.3	2.1	*	*	*	*
2002	1.1	...	...	*	1.0	1.6	2.0	1.6	1.4	*	*	*
2001 6/	1.1	...	...	*	0.7	1.5	2.0	2.1	*	*	*	*
2000	1.1	...	...	*	1.3	1.5	1.9	1.5	*	*	*	*
1999	1.0	...	...	*	0.9	1.6	2.0	1.3	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
All origins 3/, both sexes												
2006	6.2	8.1	2.2	1.0	13.5	11.7	6.9	5.1	3.2	2.1	2.1	1.9
2005	6.1	7.5	2.3	0.8	13.0	11.8	7.1	4.8	2.8	2.4	2.2	2.1
2004	5.9	8.0	2.4	0.8	12.2	11.2	6.8	4.8	3.0	2.4	2.2	2.1
2003	6.1	8.5	2.4	0.8	13.0	11.3	7.0	4.9	2.8	2.4	2.5	2.2
2002	6.1	7.5	2.7	0.9	12.9	11.2	7.2	4.8	3.2	2.3	2.3	2.1
2001 6/	7.1	8.2	2.7	0.8	13.3	13.1	9.5	6.3	4.0	2.9	2.5	2.4
2000	6.0	9.2	2.3	0.9	12.6	10.4	7.1	4.7	3.0	2.4	2.4	2.4
1999	6.1	8.7	2.5	1.1	12.9	10.5	7.1	4.6	3.0	2.6	2.5	2.4
All origins 3/, male												
2006	10.0	9.4	2.5	1.2	22.8	19.4	10.4	7.6	4.8	3.0	2.7	2.3
2005	9.8	8.2	2.6	1.0	22.0	19.6	10.6	7.6	4.3	3.3	2.6	2.7
2004	9.4	8.0	2.5	1.0	20.4	18.5	10.1	7.3	4.6	3.6	2.6	2.7
2003	9.7	10.0	2.5	1.0	21.8	18.5	10.4	7.4	4.2	3.4	3.3	3.2
2002	9.6	7.9	2.9	0.9	21.5	18.2	10.7	7.1	4.8	3.3	3.1	3.0
2001 6/	11.1	9.5	3.0	0.9	22.2	20.8	13.9	9.5	5.9	4.0	3.1	3.2
2000	9.3	10.4	2.5	1.1	20.9	16.7	10.3	6.9	4.6	3.4	3.2	3.3
1999	9.3	9.6	2.5	1.2	21.1	16.5	10.3	6.8	4.7	3.7	3.4	3.4
All origins 3/, female												
2006	2.5	6.8	2.0	0.7	3.5	3.7	3.4	2.6	1.7	1.3	1.8	1.8
2005	2.5	6.6	2.0	0.7	3.4	3.8	3.6	2.2	1.4	1.6	1.9	1.9
2004	2.5	7.9	2.2	0.6	3.5	3.7	3.4	2.4	1.6	1.4	1.9	1.9
2003	2.6	6.9	2.3	0.5	3.7	3.9	3.6	2.5	1.5	1.5	1.9	1.8
2002	2.7	7.1	2.5	0.8	3.8	4.2	3.8	2.6	1.6	1.4	1.8	1.7
2001 6/	3.3	6.9	2.4	0.7	3.9	5.3	5.1	3.2	2.2	2.0	2.1	2.0
2000	2.8	7.9	2.1	0.7	3.9	4.1	4.0	2.5	1.6	1.6	2.0	2.0
1999	2.9	7.8	2.4	0.9	4.3	4.4	4.0	2.4	1.5	1.7	1.9	2.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Hispanic, both sexes												
2006	8.0	8.3	1.7	1.2	18.2	12.3	8.1	5.8	3.7	2.3	*	*
2005	8.2	6.4	2.3	0.8	18.2	13.9	8.1	6.2	3.0	2.7	3.2	*
2004	7.9	8.1	2.3	0.9	17.4	12.6	7.7	5.9	3.8	2.3	*	*
2003	8.4	6.9	2.5	0.7	18.4	13.2	8.5	6.4	3.6	3.8	3.6	*
2002	8.1	6.2	2.6	0.9	17.6	12.3	8.0	5.9	4.4	2.5	*	*
2001 6/	9.0	6.7	2.3	0.8	18.3	14.0	10.4	7.3	4.4	4.0	*	*
2000	8.3	7.0	1.9	1.1	17.1	12.4	8.7	6.7	4.5	3.3	*	*
1999	8.4	7.0	1.8	1.4	17.7	12.7	8.8	6.6	4.2	3.2	*	*
Hispanic, male												
2006	13.1	10.1	1.8	1.5	31.0	19.7	12.5	9.1	6.7	3.2	*	*
2005	13.6	6.3	2.5	1.0	31.0	22.3	12.7	10.1	4.7	4.4	*	*
2004	13.0	9.1	2.5	1.0	29.6	20.3	11.8	9.5	6.5	4.1	*	*
2003	13.6	7.9	2.1	0.9	30.3	21.2	12.9	10.2	6.0	6.8	*	*
2002	13.2	6.6	3.1	1.0	29.6	19.8	12.1	9.1	7.7	4.8	*	*
2001 6/	14.5	7.9	2.5	1.1	30.5	21.6	15.9	11.3	6.8	6.6	*	*
2000	13.4	6.6	2.0	1.6	28.5	19.9	13.5	10.2	7.1	4.6	*	*
1999	13.8	6.7	1.7	1.8	29.1	20.6	13.9	10.3	7.3	5.7	*	*
Hispanic, female												
2006	2.4	6.5	1.5	0.8	3.8	3.2	3.1	2.4	*	*	*	*
2005	2.5	6.6	2.0	0.6	3.6	3.7	3.1	2.2	*	*	*	*
2004	2.4	7.0	2.1	0.7	3.3	3.4	3.2	2.2	*	*	*	*
2003	2.8	5.9	2.9	*	4.5	3.5	3.6	2.7	*	*	*	*
2002	2.6	5.9	2.1	0.8	3.8	3.4	3.5	2.7	*	*	*	*
2001 6/	3.2	5.5	2.1	*	4.0	5.1	4.4	3.4	2.4	*	*	*
2000	2.8	7.4	1.9	0.6	3.7	3.7	3.6	3.3	2.2	*	*	*
1999	2.8	7.3	1.9	0.9	4.4	3.5	3.3	3.1	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Non-Hispanic 4/, both sexes												
2006	5.9	8.0	2.4	0.9	12.4	11.5	6.7	5.0	3.1	2.1	2.2	1.9
2005	5.7	7.7	2.3	0.9	11.8	11.3	6.9	4.7	2.8	2.3	2.1	2.2
2004	5.5	7.8	2.4	0.8	11.1	10.8	6.6	4.6	3.0	2.4	2.3	2.0
2003	5.7	8.9	2.3	0.8	11.8	10.8	6.7	4.8	2.7	2.2	2.4	2.1
2002	5.7	7.7	2.7	0.9	11.8	10.9	7.0	4.6	3.0	2.3	2.2	2.1
2001 6/	6.7	8.4	2.8	0.8	12.1	12.8	9.2	6.1	3.9	2.8	2.5	2.4
2000	5.6	9.6	2.4	0.9	11.6	10.0	6.9	4.4	2.9	2.3	2.4	2.3
1999	5.7	8.9	2.6	1.0	11.9	10.0	6.8	4.3	2.9	2.5	2.5	2.4
Non-Hispanic 4/, male												
2006	9.3	9.1	2.7	1.1	21.0	19.2	9.9	7.4	4.6	3.0	2.7	2.1
2005	9.1	8.8	2.6	1.0	19.9	18.8	10.1	7.2	4.3	3.2	2.4	2.8
2004	8.7	7.5	2.5	0.9	18.4	17.9	9.7	6.9	4.4	3.5	2.7	2.4
2003	9.0	10.5	2.6	1.0	19.8	17.7	10.0	7.1	4.0	3.1	3.2	3.0
2002	8.9	8.0	2.9	0.9	19.6	17.6	10.4	6.8	4.5	3.2	3.1	2.9
2001 6/	10.4	9.8	3.2	0.9	20.1	20.3	13.4	9.2	5.8	3.8	3.1	3.1
2000	8.6	11.2	2.6	1.0	19.1	15.9	9.8	6.6	4.4	3.2	3.1	3.1
1999	8.6	10.1	2.6	1.1	19.4	15.6	9.7	6.4	4.5	3.5	3.3	3.4
Non-Hispanic 4/, female												
2006	2.5	6.8	2.1	0.7	3.5	3.8	3.5	2.6	1.7	1.3	1.8	1.8
2005	2.5	6.6	2.0	0.7	3.4	3.8	3.7	2.2	1.4	1.6	1.9	1.9
2004	2.5	8.1	2.3	0.6	3.5	3.7	3.5	2.4	1.6	1.5	2.0	1.8
2003	2.5	7.2	2.1	0.6	3.5	4.0	3.6	2.5	1.5	1.5	1.8	1.8
2002	2.7	7.4	2.6	0.8	3.7	4.3	3.8	2.5	1.7	1.5	1.7	1.7
2001 6/	3.2	6.9	2.4	0.7	3.8	5.3	5.1	3.1	2.2	2.0	2.1	2.1
2000	2.7	7.9	2.2	0.7	3.8	4.1	4.0	2.3	1.5	1.6	1.9	2.0
1999	2.9	7.6	2.5	0.9	4.2	4.6	4.0	2.3	1.4	1.7	2.0	2.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Non-Hispanic white, both sexes												
2006 5/	2.7	6.3	1.5	0.6	3.4	3.9	3.6	3.2	2.3	1.6	2.0	1.7
2005 5/	2.7	6.0	1.5	0.5	3.4	4.0	3.7	3.0	2.1	1.8	1.8	1.9
2004 5/	2.7	5.9	1.4	0.5	3.5	4.0	3.8	3.0	2.2	2.0	1.9	1.8
2003 5/	2.7	6.8	1.3	0.4	3.5	3.9	3.8	3.1	2.1	1.7	2.0	2.0
2002	2.8	5.0	1.8	0.5	3.7	3.9	4.1	3.0	2.4	1.9	1.8	1.8
2001 6/	4.0	5.8	1.7	0.6	4.2	6.5	6.6	4.6	3.3	2.3	2.1	2.2
2000	2.8	6.2	1.6	0.6	3.5	4.0	4.0	2.9	2.2	1.8	2.0	2.1
1999	2.9	6.3	1.7	0.7	3.9	4.3	4.1	2.9	2.2	2.1	2.0	2.1
Non-Hispanic white, male												
2006 5/	3.6	6.5	1.6	0.6	4.7	5.5	4.9	4.5	3.2	2.1	2.2	1.8
2005 5/	3.6	6.9	1.6	0.5	4.7	5.4	4.8	4.3	3.1	2.2	1.9	2.0
2004 5/	3.6	6.1	1.6	0.6	4.7	5.5	5.0	4.3	3.1	2.6	2.1	1.9
2003 5/	3.6	8.0	1.3	0.6	5.0	5.2	4.9	4.3	2.8	2.3	2.6	2.9
2002	3.8	5.8	1.9	0.5	5.2	5.3	5.6	3.9	3.3	2.4	2.3	2.7
2001 6/	5.6	6.8	2.0	0.6	5.7	9.3	9.5	6.7	4.7	2.9	2.6	2.8
2000	3.6	8.3	1.8	0.7	4.7	5.2	5.2	3.9	3.1	2.2	2.4	2.6
1999	3.9	8.0	1.7	0.7	5.1	5.7	5.4	4.2	3.2	2.7	2.6	2.6
Non-Hispanic white, female												
2006 5/	1.8	6.2	1.3	0.6	2.0	2.2	2.4	2.0	1.5	1.3	1.8	1.7
2005 5/	1.8	5.0	1.4	0.4	2.0	2.6	2.7	1.6	1.2	1.4	1.8	1.9
2004 5/	1.8	5.8	1.3	0.4	2.3	2.5	2.5	1.8	1.4	1.4	1.8	1.7
2003 5/	1.8	5.5	1.4	0.3	1.9	2.5	2.6	2.0	1.4	1.3	1.6	1.6
2002	1.9	4.1	1.8	0.6	2.2	2.6	2.5	2.1	1.5	1.4	1.5	1.5
2001 6/	2.4	4.7	1.5	0.7	2.7	3.7	3.7	2.6	1.9	1.8	1.8	2.0
2000	1.9	4.1	1.3	0.6	2.3	2.7	2.7	1.9	1.3	1.4	1.7	1.9
1999	2.0	4.4	1.6	0.7	2.6	2.9	2.8	1.7	1.3	1.6	1.6	1.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Non-Hispanic black, both sexes												
2006 5/	23.6	15.9	6.0	2.4	51.2	49.3	24.5	16.2	9.7	6.1	4.5	*
2005 5/	22.8	14.5	5.5	2.3	48.5	47.9	24.8	16.0	8.2	7.2	5.9	*
2004 5/	21.7	15.8	6.2	2.0	45.1	45.9	23.3	15.4	8.8	6.2	6.4	*
2003 5/	22.7	16.8	6.4	2.3	49.1	47.1	24.2	15.8	8.1	6.3	6.6	*
2002	22.5	17.6	6.5	2.0	48.3	47.0	24.3	15.8	8.4	5.8	7.3	*
2001 6/	22.7	19.5	7.1	1.4	48.7	45.3	25.5	16.2	8.9	7.2	6.8	*
2000	22.1	24.1	6.1	1.9	49.3	42.2	24.5	15.0	8.4	7.6	7.1	*
1999	21.7	20.0	6.4	2.3	49.5	40.6	23.7	14.1	8.4	6.4	7.5	*
Non-Hispanic black, male												
2006 5/	42.0	20.3	6.4	3.3	91.6	90.2	41.2	27.7	17.2	11.5	8.3	*
2005 5/	40.9	15.9	6.9	2.9	87.0	89.4	41.9	27.6	14.6	13.0	10.4	*
2004 5/	38.3	13.9	6.3	2.5	80.7	85.1	39.0	26.1	15.2	11.8	10.3	*
2003 5/	40.1	18.6	7.8	3.0	87.3	86.1	41.6	26.9	14.9	11.3	11.3	*
2002	39.4	16.4	7.3	2.6	85.8	85.3	39.8	27.5	15.8	10.5	13.8	*
2001 6/	39.3	21.7	8.4	2.0	88.3	81.7	39.5	27.2	15.0	12.6	10.2	*
2000	38.3	24.9	6.1	2.1	88.1	76.5	39.4	26.3	15.0	13.9	12.0	*
1999	37.2	20.1	6.2	2.7	88.1	71.5	37.9	23.7	15.4	11.9	12.2	*
Non-Hispanic black, female												
2006 5/	6.8	11.3	5.6	1.4	9.8	11.5	9.7	6.4	3.7	2.2	*	*
2005 5/	6.3	13.0	4.1	1.7	9.0	9.7	9.5	6.1	3.1	3.1	3.3	*
2004 5/	6.6	17.7	6.1	1.6	9.1	10.1	9.3	6.2	3.7	2.1	4.2	*
2003 5/	6.8	14.8	4.9	1.5	10.4	11.6	8.8	6.2	2.7	2.8	3.9	*
2002	7.2	18.9	5.7	1.5	10.6	12.3	10.6	5.7	2.5	2.4	3.6	*
2001 6/	7.6	17.1	5.7	0.8	9.2	12.5	13.2	6.9	4.0	3.3	4.9	*
2000	7.4	23.3	6.0	1.6	11.0	11.4	11.3	5.3	3.2	3.2	4.3	*
1999	7.7	20.0	6.7	1.8	11.4	12.8	11.1	5.9	2.7	2.5	4.9	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All origins 3/, both sexes												
2006	4.3	*	0.3	0.6	11.4	9.3	4.6	2.7	1.5	0.8	0.6	0.4
2005	4.2	*	0.2	0.5	10.7	9.4	4.6	2.6	1.3	1.1	0.7	0.5
2004	4.0	*	0.2	0.5	9.9	8.8	4.3	2.6	1.6	1.0	0.8	0.6
2003	4.1	*	0.3	0.5	10.7	8.9	4.4	2.7	1.4	0.9	0.7	0.5
2002	4.1	*	0.3	0.5	10.6	8.7	4.5	2.6	1.5	0.9	0.9	*
2001	4.0	*	0.4	0.4	10.5	8.4	4.4	2.4	1.4	1.0	0.8	*
2000	3.8	*	0.2	0.5	10.1	7.7	4.2	2.5	1.5	1.0	0.7	0.7
1999	3.9	*	0.3	0.5	10.4	7.6	4.1	2.4	1.6	1.0	0.8	0.6
All origins 3/, male												
2006	7.4	*	0.3	0.8	20.0	16.2	7.3	4.3	2.2	1.2	0.8	*
2005	7.2	*	0.3	0.6	18.9	16.4	7.3	4.1	2.1	1.5	0.8	*
2004	6.9	*	*	0.6	17.3	15.3	7.1	4.0	2.5	1.6	0.8	*
2003	7.1	*	0.2	0.6	18.8	15.5	6.9	4.3	2.2	1.4	0.7	*
2002	7.0	*	0.4	0.6	18.5	14.9	7.3	3.9	2.4	1.2	1.3	*
2001	6.8	*	0.3	0.6	18.5	14.3	6.8	3.8	2.2	1.5	1.1	*
2000	6.5	*	*	0.7	17.7	13.2	6.6	3.9	2.3	1.3	0.8	*
1999	6.5	*	0.3	0.7	18.0	12.9	6.3	3.8	2.6	1.5	1.0	*
All origins 3/, female												
2006	1.3	*	*	0.4	2.2	2.2	1.8	1.3	0.8	0.5	0.6	*
2005	1.2	*	*	0.3	2.0	2.2	1.9	1.1	0.6	0.7	0.7	*
2004	1.1	*	*	0.3	2.0	2.0	1.5	1.1	0.8	0.5	0.8	*
2003	1.2	*	0.3	0.3	2.2	2.1	1.8	1.2	0.7	0.5	0.7	*
2002	1.3	*	*	0.4	2.4	2.3	1.9	1.3	0.8	0.6	0.7	*
2001	1.3	*	0.4	0.3	2.1	2.3	2.0	1.0	0.8	0.7	0.6	*
2000	1.3	*	*	0.2	2.2	2.1	1.8	1.2	0.8	0.8	0.6	*
1999	1.3	*	0.3	0.3	2.5	2.2	2.0	1.2	0.6	0.7	0.6	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Hispanic, both sexes												
2006	5.6	*	*	0.8	15.1	9.3	5.2	3.0	1.6	*	*	*
2005	5.7	*	*	0.5	14.5	10.6	5.3	3.3	1.3	1.5	*	*
2004	5.4	*	*	0.5	14.0	9.4	5.0	2.7	2.0	*	*	*
2003	5.8	*	*	0.4	14.9	9.9	5.3	3.6	1.2	*	*	*
2002	5.6	*	*	0.6	14.3	9.3	5.1	3.2	1.6	*	*	*
2001	5.7	*	*	0.4	14.6	9.5	5.6	3.0	1.1	*	*	*
2000	5.5	*	*	0.6	13.7	8.9	4.9	4.0	2.0	*	*	*
1999	5.7	*	*	0.8	14.4	8.7	5.2	3.8	2.1	*	*	*
Hispanic, male												
2006	9.7	*	*	1.1	26.4	15.5	8.2	4.9	2.8	*	*	*
2005	10.0	*	*	0.8	25.5	17.6	8.3	5.5	2.3	*	*	*
2004	9.5	*	*	0.7	24.4	15.9	8.3	4.5	3.5	*	*	*
2003	10.0	*	*	0.6	25.5	16.8	8.4	5.9	2.2	*	*	*
2002	9.7	*	*	0.8	24.9	15.6	8.1	5.0	2.8	*	*	*
2001	9.9	*	*	0.6	24.9	16.1	9.1	4.8	*	*	*	*
2000	9.6	*	*	1.0	23.5	15.0	7.8	6.4	3.2	*	*	*
1999	9.8	*	*	1.2	24.2	14.7	8.6	5.7	3.5	*	*	*
Hispanic, female												
2006	1.2	*	*	*	2.2	1.8	1.7	1.1	*	*	*	*
2005	1.2	*	*	*	1.9	2.1	1.9	1.2	*	*	*	*
2004	1.1	*	*	*	2.0	1.6	1.4	*	*	*	*	*
2003	1.3	*	*	*	2.7	1.7	1.9	1.4	*	*	*	*
2002	1.2	*	*	*	2.1	1.8	1.8	1.4	*	*	*	*
2001	1.3	*	*	*	2.5	1.8	1.9	1.2	*	*	*	*
2000	1.3	*	*	*	2.2	1.9	1.7	1.7	*	*	*	*
1999	1.4	*	*	*	2.9	1.7	1.7	1.9	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Non-Hispanic 4/, both sexes												
2006	4.0	*	0.3	0.5	10.5	9.3	4.4	2.7	1.5	0.9	0.7	0.4
2005	3.9	*	0.2	0.5	9.9	9.1	4.4	2.5	1.3	1.0	0.8	0.5
2004	3.7	*	0.2	0.4	9.0	8.5	4.1	2.5	1.6	1.1	0.8	0.5
2003	3.8	*	0.2	0.5	9.8	8.6	4.2	2.6	1.4	0.8	0.7	0.5
2002	3.8	*	0.3	0.5	9.8	8.5	4.4	2.5	1.5	0.9	0.9	*
2001	3.7	*	0.3	0.4	9.6	8.0	4.2	2.3	1.5	1.0	0.8	*
2000	3.6	*	0.2	0.4	9.3	7.4	4.1	2.4	1.5	1.0	0.7	0.7
1999	3.6	*	0.4	0.5	9.6	7.3	4.0	2.3	1.5	1.1	0.8	0.5
Non-Hispanic 4/, male												
2006	6.9	*	*	0.7	18.5	16.3	7.0	4.2	2.2	1.2	0.8	*
2005	6.7	*	*	0.6	17.4	15.9	7.0	3.9	2.1	1.5	0.8	*
2004	6.4	*	*	0.6	15.8	15.1	6.8	3.9	2.4	1.6	0.8	*
2003	6.5	*	*	0.7	17.2	15.0	6.7	4.1	2.1	1.3	0.7	*
2002	6.5	*	0.3	0.6	17.0	14.7	7.1	3.7	2.3	1.2	1.2	*
2001	6.3	*	0.3	0.6	17.0	13.8	6.5	3.7	2.2	1.4	1.1	*
2000	6.0	*	*	0.6	16.3	12.7	6.4	3.7	2.2	1.3	0.8	*
1999	6.0	*	0.4	0.7	16.7	12.4	6.0	3.6	2.5	1.5	0.9	*
Non-Hispanic 4/, female												
2006	1.3	*	*	0.4	2.2	2.3	1.9	1.3	0.8	0.5	0.6	*
2005	1.2	*	*	0.3	2.0	2.2	1.9	1.1	0.6	0.7	0.7	*
2004	1.2	*	*	0.2	2.0	2.1	1.5	1.2	0.8	0.6	0.8	*
2003	1.2	*	*	0.3	2.0	2.2	1.8	1.2	0.7	0.5	0.7	*
2002	1.3	*	*	0.4	2.4	2.3	1.9	1.3	0.8	0.6	0.7	*
2001	1.2	*	0.3	0.3	2.0	2.3	2.0	1.0	0.8	0.7	0.6	*
2000	1.2	*	*	0.2	2.2	2.2	1.9	1.1	0.8	0.7	0.6	*
1999	1.3	*	0.4	0.3	2.4	2.3	2.0	1.1	0.6	0.7	0.6	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Non-Hispanic white, both sexes												
2006 5/	1.4	*	*	0.3	2.2	2.5	2.2	1.6	1.1	0.6	0.6	*
2005 5/	1.4	*	*	0.2	2.1	2.6	2.2	1.6	1.0	0.8	0.7	*
2004 5/	1.5	*	*	0.2	2.1	2.6	2.2	1.6	1.2	0.9	0.7	*
2003 5/	1.4	*	*	0.2	2.4	2.5	2.1	1.7	1.1	0.7	0.6	0.5
2002	1.5	*	0.2	0.3	2.5	2.6	2.4	1.6	1.2	0.7	0.8	*
2001	1.6	*	0.2	0.4	2.4	2.7	2.4	1.6	1.2	0.8	0.7	*
2000	1.4	*	*	0.3	2.2	2.4	2.3	1.6	1.2	0.8	0.6	0.6
1999	1.5	*	0.2	0.3	2.4	2.6	2.3	1.5	1.1	0.9	0.6	0.6
Non-Hispanic white, male												
2006 5/	2.0	*	*	0.3	3.4	3.8	3.1	2.2	1.5	0.8	0.5	*
2005 5/	2.1	*	*	0.3	3.2	3.8	3.0	2.3	1.5	1.0	0.7	*
2004 5/	2.1	*	*	0.3	3.1	4.0	3.1	2.3	1.7	1.2	0.7	*
2003 5/	2.1	*	*	0.3	3.7	3.7	2.9	2.4	1.6	1.0	0.5	*
2002	2.2	*	*	0.3	3.7	3.7	3.4	2.2	1.7	0.9	1.0	*
2001	2.2	*	*	0.4	3.6	3.9	3.4	2.3	1.6	1.0	0.9	*
2000	2.0	*	*	0.4	3.2	3.4	3.2	2.2	1.6	0.9	0.6	*
1999	2.2	*	*	0.4	3.5	3.9	3.2	2.2	1.7	1.2	0.6	*
Non-Hispanic white, female												
2006 5/	0.8	*	*	0.3	1.1	1.2	1.2	1.1	0.8	0.5	0.7	*
2005 5/	0.8	*	*	*	1.0	1.4	1.4	0.9	0.6	0.7	0.7	*
2004 5/	0.8	*	*	*	1.1	1.2	1.2	0.9	0.7	0.7	0.8	*
2003 5/	0.8	*	*	*	1.0	1.3	1.3	1.0	0.7	0.4	0.6	*
2002	0.9	*	*	0.3	1.2	1.4	1.3	1.1	0.7	0.6	0.7	*
2001	0.9	*	*	0.3	1.2	1.5	1.4	0.8	0.8	0.7	0.6	*
2000	0.9	*	*	0.2	1.2	1.4	1.4	0.9	0.8	0.7	0.6	*
1999	0.9	*	*	0.2	1.2	1.4	1.5	0.8	0.6	0.7	0.6	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Non-Hispanic black, both sexes												
2006 5/	18.6	*	*	1.6	46.4	43.1	17.7	9.6	4.3	2.9	*	*
2005 5/	17.7	*	*	1.4	43.6	41.8	17.4	8.7	3.6	3.1	*	*
2004 5/	16.6	*	*	1.3	40.3	39.4	16.0	8.6	4.7	2.4	*	*
2003 5/	17.3	*	*	1.5	43.3	40.5	16.5	8.9	4.0	2.5	*	*
2002	17.1	*	*	1.2	43.1	39.7	16.7	8.6	4.2	2.4	2.1	*
2001	16.3	*	*	0.9	43.0	36.4	15.5	7.7	3.7	2.6	*	*
2000	16.0	*	*	1.0	42.6	34.5	15.4	8.5	3.6	2.8	*	*
1999	15.9	*	*	1.3	43.6	32.4	14.4	8.1	4.5	2.4	2.4	*
Non-Hispanic black, male												
2006 5/	35.0	*	*	2.3	84.7	81.2	31.8	17.9	8.0	5.6	*	*
2005 5/	33.6	*	*	2.0	80.3	80.3	31.8	15.9	6.9	6.1	*	*
2004 5/	31.4	*	*	1.9	74.1	75.4	29.8	15.5	8.5	5.6	*	*
2003 5/	32.6	*	*	2.1	79.4	77.4	30.1	16.4	7.4	4.8	*	*
2002	31.9	*	*	1.6	78.6	75.6	30.0	15.3	7.7	4.7	*	*
2001	30.6	*	*	1.3	80.6	69.4	26.7	14.0	7.1	5.5	*	*
2000	30.1	*	*	1.6	79.1	65.7	27.4	15.3	6.5	5.5	*	*
1999	29.3	*	*	1.8	80.0	61.1	24.6	14.1	9.1	4.7	*	*
Non-Hispanic black, female												
2006 5/	3.7	*	*	0.8	7.2	7.8	5.2	2.5	1.3	*	*	*
2005 5/	3.2	*	*	0.7	6.1	6.4	4.6	2.5	*	*	*	*
2004 5/	3.1	*	*	*	6.0	6.4	3.8	2.6	1.6	*	*	*
2003 5/	3.4	*	*	0.9	6.7	6.9	4.4	2.6	*	*	*	*
2002	3.7	*	*	0.8	7.5	7.2	5.0	2.8	*	*	*	*
2001	3.2	*	*	*	5.6	6.5	5.5	2.2	*	*	*	*
2000	3.3	*	*	*	6.7	6.4	4.7	2.6	*	*	*	*
1999	3.7	*	*	0.7	7.8	6.6	5.5	3.0	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
All origins 3/, both sexes												
2006	1.9	8.0	2.0	0.4	2.1	2.4	2.4	2.4	1.7	1.3	1.5	1.5
2005	1.9	7.3	2.1	0.4	2.3	2.4	2.5	2.3	1.5	1.3	1.4	1.7
2004	2.0	7.8	2.1	0.4	2.3	2.5	2.5	2.3	1.4	1.4	1.5	1.5
2003	2.0	8.3	2.1	0.3	2.3	2.4	2.6	2.2	1.4	1.5	1.8	1.7
2002	2.0	7.3	2.4	0.4	2.2	2.6	2.7	2.2	1.6	1.4	1.4	1.7
2001 6/	3.1	8.0	2.3	0.4	2.7	4.8	5.1	3.9	2.6	1.9	1.7	2.0
2000	2.1	8.9	2.1	0.4	2.5	2.7	2.9	2.1	1.5	1.4	1.8	1.7
1999	2.2	8.5	2.1	0.5	2.5	2.9	3.0	2.1	1.5	1.5	1.7	1.8
All origins 3/, male												
2006	2.6	9.3	2.2	0.4	2.8	3.2	3.1	3.4	2.6	1.8	1.9	1.8
2005	2.6	7.9	2.3	0.4	3.1	3.3	3.3	3.4	2.2	1.8	1.8	2.1
2004	2.5	7.9	2.3	0.3	3.1	3.2	3.0	3.3	2.1	1.9	1.9	1.7
2003	2.6	9.7	2.3	0.4	3.1	3.0	3.5	3.1	2.1	2.0	2.6	2.5
2002	2.6	7.8	2.6	0.3	3.0	3.2	3.4	3.2	2.4	2.1	1.9	2.3
2001 6/	4.3	9.2	2.7	0.3	3.6	6.5	7.1	5.7	3.8	2.5	2.0	2.7
2000	2.8	10.0	2.3	0.4	3.3	3.5	3.7	3.0	2.3	2.1	2.4	2.5
1999	2.8	9.4	2.1	0.5	3.1	3.6	4.0	3.1	2.2	2.2	2.4	2.6
All origins 3/, female												
2006	1.3	6.6	1.8	0.4	1.3	1.5	1.6	1.4	0.9	0.8	1.2	1.4
2005	1.3	6.6	1.8	0.4	1.5	1.6	1.7	1.1	0.8	0.9	1.2	1.5
2004	1.4	7.7	2.0	0.4	1.5	1.7	1.9	1.3	0.8	0.9	1.2	1.5
2003	1.4	6.9	2.0	0.3	1.6	1.8	1.8	1.3	0.8	1.0	1.2	1.4
2002	1.4	6.8	2.2	0.4	1.4	1.9	1.9	1.3	0.9	0.8	1.1	1.4
2001 6/	2.0	6.7	2.0	0.4	1.8	3.1	3.1	2.1	1.5	1.3	1.5	1.7
2000	1.5	7.7	2.0	0.5	1.7	1.9	2.1	1.3	0.8	0.9	1.4	1.4
1999	1.6	7.6	2.1	0.6	1.8	2.2	2.0	1.2	0.8	1.0	1.3	1.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
Hispanic, both sexes												
2006	2.4	8.2	1.4	0.4	3.2	3.0	2.9	2.8	2.1	1.7	*	*
2005	2.5	6.3	2.1	0.3	3.7	3.3	2.8	2.8	1.7	*	*	*
2004	2.5	7.7	2.1	0.4	3.4	3.2	2.7	3.2	1.8	*	*	*
2003	2.6	6.8	2.2	0.3	3.5	3.3	3.2	2.8	2.3	2.4	3.1	*
2002	2.5	6.0	2.2	0.3	3.3	3.1	2.9	2.7	2.8	1.7	*	*
2001 6/	3.3	6.2	1.9	0.4	3.7	4.5	4.8	4.3	3.4	2.7	*	*
2000	2.7	6.9	1.8	0.5	3.4	3.5	3.9	2.7	2.5	2.0	*	*
1999	2.7	6.7	1.6	0.6	3.3	4.0	3.5	2.9	2.1	2.3	*	*
Hispanic, male												
2006	3.4	10.1	1.5	*	4.6	4.3	4.3	4.2	3.9	*	*	*
2005	3.6	6.1	2.3	*	5.5	4.7	4.4	4.6	2.4	*	*	*
2004	3.5	8.9	2.3	*	5.3	4.4	3.6	5.0	3.0	*	*	*
2003	3.6	7.7	1.8	*	4.8	4.5	4.6	4.3	3.7	4.4	*	*
2002	3.5	6.6	2.6	*	4.7	4.3	4.0	4.1	4.9	*	*	*
2001 6/	4.5	7.2	2.2	*	5.6	5.5	6.9	6.5	5.1	4.6	*	*
2000	3.8	6.6	1.8	0.6	5.0	5.0	5.7	3.8	3.9	*	*	*
1999	4.0	6.2	1.5	0.6	4.9	5.9	5.3	4.5	3.8	4.4	*	*
Hispanic, female												
2006	1.3	6.3	1.3	*	1.6	1.3	1.4	1.3	*	*	*	*
2005	1.3	6.6	1.9	*	1.7	1.6	1.2	1.0	*	*	*	*
2004	1.4	6.5	1.9	*	1.3	1.8	1.7	1.3	*	*	*	*
2003	1.5	5.9	2.6	*	1.9	1.8	1.7	1.4	*	*	*	*
2002	1.4	5.4	1.8	*	1.7	1.6	1.7	1.4	*	*	*	*
2001 6/	1.9	5.2	1.6	*	1.5	3.2	2.6	2.2	*	*	*	*
2000	1.6	7.2	1.7	*	1.5	1.9	1.9	1.6	*	*	*	*
1999	1.4	7.3	1.8	*	1.5	1.8	1.6	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
Non-Hispanic 4/, both sexes												
2006	1.8	7.8	2.2	0.4	1.9	2.2	2.3	2.3	1.7	1.2	1.5	1.5
2005	1.8	7.6	2.1	0.4	2.0	2.2	2.4	2.2	1.5	1.3	1.4	1.7
2004	1.8	7.7	2.1	0.4	2.0	2.3	2.4	2.1	1.4	1.4	1.5	1.5
2003	1.9	8.6	2.1	0.3	2.1	2.2	2.5	2.1	1.3	1.4	1.7	1.6
2002	1.9	7.5	2.4	0.4	2.0	2.4	2.6	2.1	1.5	1.4	1.3	1.7
2001 6/	3.1	8.2	2.4	0.4	2.5	4.7	5.0	3.8	2.5	1.8	1.7	2.1
2000	2.0	9.2	2.2	0.4	2.3	2.6	2.8	2.0	1.4	1.4	1.7	1.6
1999	2.1	8.7	2.2	0.5	2.3	2.7	2.9	2.0	1.4	1.5	1.7	1.8
Non-Hispanic 4/, male												
2006	2.4	9.0	2.4	0.4	2.4	2.9	2.9	3.3	2.5	1.7	1.9	1.7
2005	2.4	8.5	2.3	0.4	2.5	2.9	3.0	3.3	2.2	1.8	1.6	2.2
2004	2.3	7.4	2.3	0.3	2.6	2.9	2.9	3.0	2.0	1.9	1.9	1.7
2003	2.4	10.1	2.4	0.4	2.6	2.7	3.3	3.0	1.9	1.8	2.5	2.3
2002	2.5	7.8	2.5	0.3	2.6	2.9	3.3	3.1	2.2	2.0	1.8	2.3
2001 6/	4.2	9.5	2.8	0.3	3.2	6.5	7.0	5.5	3.6	2.4	2.0	2.7
2000	2.6	10.6	2.4	0.4	2.9	3.2	3.4	2.9	2.2	2.0	2.3	2.4
1999	2.6	10.0	2.3	0.4	2.8	3.2	3.8	2.9	2.0	2.0	2.4	2.6
Non-Hispanic 4/, female												
2006	1.3	6.6	1.9	0.4	1.3	1.5	1.6	1.4	0.9	0.8	1.2	1.4
2005	1.3	6.6	1.8	0.4	1.4	1.6	1.8	1.1	0.8	0.9	1.2	1.4
2004	1.4	8.0	2.0	0.4	1.5	1.7	1.9	1.3	0.8	0.9	1.2	1.4
2003	1.4	7.1	1.8	0.3	1.5	1.8	1.8	1.3	0.8	1.0	1.1	1.3
2002	1.4	7.1	2.3	0.4	1.4	1.9	1.9	1.2	0.9	0.9	1.0	1.4
2001 6/	2.0	6.8	2.0	0.4	1.9	2.9	3.1	2.1	1.4	1.3	1.5	1.8
2000	1.5	7.8	2.0	0.5	1.7	1.9	2.1	1.2	0.7	0.9	1.3	1.3
1999	1.6	7.3	2.2	0.6	1.8	2.3	2.0	1.2	0.8	1.0	1.3	1.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
Non-Hispanic white, both sexes												
2006 5/	1.3	6.3	1.3	0.3	1.2	1.4	1.4	1.6	1.2	1.0	1.4	1.4
2005 5/	1.2	5.9	1.3	0.2	1.3	1.4	1.5	1.4	1.1	0.9	1.1	1.5
2004 5/	1.3	5.9	1.4	0.3	1.4	1.4	1.6	1.4	1.0	1.1	1.2	1.3
2003 5/	1.2	6.8	1.2	0.2	1.1	1.4	1.6	1.4	1.0	1.1	1.4	1.5
2002	1.3	5.0	1.6	0.3	1.2	1.4	1.7	1.4	1.2	1.1	1.0	1.4
2001 6/	2.4	5.6	1.5	0.3	1.7	3.8	4.2	3.1	2.1	1.5	1.4	1.9
2000	1.3	6.0	1.4	0.4	1.3	1.6	1.7	1.4	1.0	1.0	1.4	1.5
1999	1.4	6.2	1.4	0.4	1.5	1.7	1.8	1.4	1.1	1.2	1.4	1.5
Non-Hispanic white, male												
2006 5/	1.6	6.5	1.5	0.2	1.4	1.7	1.8	2.3	1.7	1.3	1.7	1.6
2005 5/	1.5	6.7	1.4	0.2	1.5	1.7	1.8	2.0	1.6	1.2	1.2	1.6
2004 5/	1.5	6.1	1.5	0.2	1.6	1.5	1.9	1.9	1.4	1.5	1.4	*
2003 5/	1.5	8.0	1.2	0.3	1.3	1.5	2.0	1.9	1.3	1.3	2.1	2.2
2002	1.6	5.8	1.6	0.2	1.5	1.6	2.2	1.8	1.6	1.6	1.3	2.0
2001 6/	3.4	6.7	1.8	0.2	2.0	5.4	6.1	4.5	3.1	1.9	1.7	2.3
2000	1.6	8.1	1.7	0.3	1.5	1.8	2.0	1.8	1.5	1.3	1.8	1.9
1999	1.7	7.8	1.5	0.3	1.6	1.8	2.3	2.0	1.5	1.5	2.0	*
Non-Hispanic white, female												
2006 5/	1.0	6.1	1.1	0.3	1.0	1.0	1.1	0.9	0.7	0.7	1.1	1.3
2005 5/	0.9	5.0	1.2	0.3	1.0	1.1	1.2	0.8	0.7	0.7	1.1	1.4
2004 5/	1.0	5.8	1.2	0.3	1.2	1.3	1.3	0.9	0.6	0.8	1.0	1.3
2003 5/	1.0	5.4	1.2	0.2	1.0	1.2	1.3	1.0	0.7	0.9	1.0	1.2
2002	1.0	4.1	1.6	0.3	0.9	1.2	1.2	1.0	0.8	0.8	0.8	1.2
2001 6/	1.5	4.5	1.2	0.4	1.4	2.2	2.3	1.7	1.2	1.1	1.2	1.7
2000	1.0	3.9	1.2	0.4	1.2	1.3	1.4	1.0	0.5	0.7	1.1	1.3
1999	1.1	4.4	1.4	0.4	1.4	1.6	1.3	0.9	0.7	0.9	1.0	1.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
Non-Hispanic black, both sexes												
2006 5/	5.0	15.3	5.2	0.8	4.7	6.3	6.7	6.6	5.5	3.2	3.1	*
2005 5/	5.1	14.1	5.0	0.9	4.9	6.1	7.4	7.3	4.6	4.1	4.4	*
2004 5/	5.1	15.3	5.4	0.8	4.8	6.5	7.2	6.8	4.1	3.7	4.9	*
2003 5/	5.4	15.8	5.6	0.8	5.8	6.6	7.7	6.8	4.2	3.8	4.9	*
2002	5.4	16.7	5.9	0.8	5.2	7.3	7.6	7.2	4.2	3.4	5.2	*
2001 6/	6.4	19.1	6.3	0.6	5.7	9.0	10.1	8.6	5.1	4.5	4.9	*
2000	6.1	22.9	5.7	0.8	6.7	7.8	9.1	6.5	4.9	4.8	5.2	*
1999	5.8	19.7	5.7	1.0	5.9	8.2	9.2	6.0	3.9	4.0	5.2	*
Non-Hispanic black, male												
2006 5/	7.1	19.7	5.6	0.9	6.8	9.0	9.4	9.8	9.3	5.9	*	*
2005 5/	7.4	15.2	6.2	0.9	6.7	9.1	10.1	11.7	7.7	6.8	7.0	*
2004 5/	6.9	13.6	5.6	0.6	6.6	9.6	9.2	10.6	6.6	6.3	8.3	*
2003 5/	7.6	16.8	6.9	0.9	7.9	8.7	11.5	10.5	7.4	6.5	8.7	*
2002	7.6	15.8	6.6	0.9	7.3	9.7	9.9	12.3	8.1	5.9	9.4	*
2001 6/	8.7	21.4	7.7	0.7	7.7	12.3	12.8	13.2	7.9	7.1	*	*
2000	8.2	22.5	5.7	*	9.1	10.8	12.0	11.0	8.5	8.4	8.3	*
1999	7.9	19.7	5.6	0.9	8.1	10.5	13.3	9.6	6.3	7.2	7.2	*
Non-Hispanic black, female												
2006 5/	3.1	10.6	4.9	*	2.6	3.7	4.4	3.9	2.4	*	*	*
2005 5/	3.1	13.0	3.7	0.9	2.9	3.3	4.9	3.6	2.1	2.1	*	*
2004 5/	3.5	17.1	5.3	1.0	3.1	3.7	5.5	3.5	2.1	*	*	*
2003 5/	3.4	14.8	4.3	0.7	3.7	4.8	4.4	3.6	1.5	*	*	*
2002	3.5	17.7	5.2	0.7	3.1	5.1	5.7	2.9	*	*	*	*
2001 6/	4.4	16.8	5.0	*	3.6	5.9	7.7	4.6	2.9	2.7	4.4	*
2000	4.1	23.3	5.8	1.1	4.3	5.0	6.5	2.7	2.0	2.3	3.5	*
1999	4.0	19.6	5.7	1.1	3.6	6.1	5.6	2.9	1.9	*	4.0	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
All origins 3/, both sexes												
2006	0.1	*	*	*	0.2	0.3	0.3	0.2	*	*	*	*
2005	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2004	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
2003	0.1	*	*	*	0.2	0.3	0.3	0.1	*	*	*	*
2002	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2000	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
1999	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
All origins 3/, male												
2006	0.3	*	*	*	0.4	0.7	0.5	0.3	*	*	*	*
2005	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	*	*	*
2004	0.3	*	*	*	0.4	0.5	0.5	0.3	*	*	*	*
2003	0.3	*	*	*	0.4	0.6	0.5	0.2	*	*	*	*
2002	0.3	*	*	*	0.4	0.5	0.5	0.3	*	*	*	*
2001	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	*	*	*
2000	0.3	*	*	*	0.4	0.5	0.4	0.3	*	*	*	*
1999	0.3	*	*	*	0.3	0.7	0.5	0.3	0.2	*	*	*
All origins 3/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
Hispanic, both sexes												
2006	0.2	*	*	*	0.3	0.5	*	*	*	*	*	*
2005	0.2	*	*	*	0.4	0.3	0.3	*	*	*	*	*
2004	0.2	*	*	*	0.4	0.4	*	*	*	*	*	*
2003	0.2	*	*	*	0.4	0.4	*	*	*	*	*	*
2002	0.2	*	*	*	0.3	0.3	*	*	*	*	*	*
2001	0.2	*	*	*	0.3	0.3	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	0.4	*	*	*	*	*	*
Hispanic, male												
2006	0.4	*	*	*	0.6	0.9	*	*	*	*	*	*
2005	0.4	*	*	*	0.8	0.6	0.6	*	*	*	*	*
2004	0.4	*	*	*	0.8	0.7	*	*	*	*	*	*
2003	0.4	*	*	*	0.7	0.8	*	*	*	*	*	*
2002	0.3	*	*	*	0.6	0.5	*	*	*	*	*	*
2001	0.3	*	*	*	0.6	0.6	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	0.7	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
Non-Hispanic 4/, both sexes												
2006	0.1	*	*	*	0.2	0.3	0.3	0.1	*	*	*	*
2005	0.1	*	*	*	0.1	0.3	0.3	0.1	0.1	*	*	*
2004	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2003	0.1	*	*	*	0.2	0.3	0.3	0.1	*	*	*	*
2002	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2000	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
1999	0.1	*	*	*	0.2	0.3	0.3	0.1	*	*	*	*
Non-Hispanic 4/, male												
2006	0.3	*	*	*	0.3	0.6	0.5	0.3	*	*	*	*
2005	0.3	*	*	*	0.3	0.5	0.5	0.3	0.2	*	*	*
2004	0.2	*	*	*	0.3	0.5	0.5	0.3	*	*	*	*
2003	0.3	*	*	*	0.3	0.6	0.5	0.2	*	*	*	*
2002	0.2	*	*	*	0.4	0.5	0.5	0.3	*	*	*	*
2001	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	*	*	*
2000	0.2	*	*	*	0.3	0.6	0.4	0.3	*	*	*	*
1999	0.3	*	*	*	0.3	0.7	0.5	0.3	*	*	*	*
Non-Hispanic 4/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
Non-Hispanic white, both sexes												
2006 5/	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2005 5/	0.1	*	*	*	0.1	0.2	0.2	0.1	0.1	*	*	*
2004 5/	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2003 5/	0.1	*	*	*	0.1	0.2	0.3	0.1	*	*	*	*
2002	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2000	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
1999	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
Non-Hispanic white, male												
2006 5/	0.2	*	*	*	0.2	0.4	0.4	0.2	*	*	*	*
2005 5/	0.2	*	*	*	0.2	0.4	0.4	0.2	0.2	*	*	*
2004 5/	0.2	*	*	*	0.2	0.3	0.4	0.2	*	*	*	*
2003 5/	0.2	*	*	*	0.2	0.3	0.5	0.2	*	*	*	*
2002	0.2	*	*	*	0.2	0.4	0.4	0.2	*	*	*	*
2001	0.2	*	*	*	0.3	0.4	0.4	0.2	*	*	*	*
2000	0.2	*	*	*	0.2	0.4	0.3	0.3	*	*	*	*
1999	0.2	*	*	*	0.2	0.5	0.4	0.2	*	*	*	*
Non-Hispanic white, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
Non-Hispanic black, both sexes												
2006 5/	0.3	*	*	*	0.4	0.8	0.5	*	*	*	*	*
2005 5/	0.3	*	*	*	0.4	0.7	0.6	*	*	*	*	*
2004 5/	0.2	*	*	*	*	0.6	*	*	*	*	*	*
2003 5/	0.3	*	*	*	0.5	1.0	0.4	*	*	*	*	*
2002	0.3	*	*	*	0.6	0.5	0.5	*	*	*	*	*
2001	0.2	*	*	*	0.4	0.4	0.4	*	*	*	*	*
2000	0.3	*	*	*	0.5	0.6	0.6	*	*	*	*	*
1999	0.4	*	*	*	0.6	0.8	0.7	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.6	*	*	*	0.7	1.7	1.1	*	*	*	*	*
2005 5/	0.6	*	*	*	0.8	1.5	1.2	*	*	*	*	*
2004 5/	0.5	*	*	*	*	1.3	*	*	*	*	*	*
2003 5/	0.7	*	*	*	1.0	2.1	0.8	*	*	*	*	*
2002	0.6	*	*	*	1.2	1.1	1.0	*	*	*	*	*
2001	0.5	*	*	*	0.8	0.8	0.8	*	*	*	*	*
2000	0.6	*	*	*	1.0	1.3	1.3	*	*	*	*	*
1999	0.8	*	*	*	1.1	1.7	1.4	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
All origins 3/, both sexes												
2006	1.7	2.2	0.3	0.1	1.4	2.0	2.9	3.1	1.6	0.9	1.0	1.4
2005	1.6	2.5	0.4	0.2	1.2	1.8	2.7	3.1	1.4	0.8	0.8	1.5
2004	1.7	2.6	0.3	0.2	1.2	2.0	3.1	3.1	1.4	0.8	0.9	1.6
2003	1.7	1.9	0.2	0.1	1.4	2.1	3.6	3.1	1.3	0.7	0.7	1.1
2002	1.7	2.5	0.4	0.2	1.2	2.1	3.4	2.8	1.2	0.8	0.7	1.4
2001	1.5	2.1	0.3	0.1	1.0	1.8	3.1	2.5	0.9	0.5	0.9	1.0
2000	1.4	1.7	0.3	0.1	0.9	1.7	2.8	2.2	0.9	0.6	0.6	1.3
1999	1.4	1.4	0.3	0.1	1.1	1.8	2.9	2.2	0.9	0.6	0.8	1.4
All origins 3/, male												
2006	2.2	2.7	0.3	0.2	2.1	2.8	3.5	3.9	1.9	1.1	1.4	2.2
2005	2.0	3.2	0.5	0.2	1.6	2.3	3.2	3.6	1.7	1.1	0.9	1.6
2004	2.2	3.1	0.3	0.2	1.8	2.7	4.0	4.0	1.7	1.1	1.0	1.9
2003	2.3	2.4	*	0.2	1.9	2.9	4.6	4.0	1.7	0.8	1.0	1.5
2002	2.2	2.5	0.4	0.2	1.7	2.9	4.4	3.5	1.6	1.0	0.9	2.2
2001	2.0	2.4	*	0.2	1.5	2.6	4.2	3.4	1.2	0.8	1.1	1.5
2000	1.9	1.4	0.3	0.2	1.4	2.4	3.9	3.1	1.3	0.8	1.0	1.7
1999	2.0	1.7	0.3	0.2	1.7	2.6	4.0	3.0	1.3	0.9	1.1	2.1
All origins 3/, female												
2006	1.2	1.7	0.3	*	0.7	1.2	2.3	2.4	1.2	0.7	0.8	1.0
2005	1.3	1.8	0.3	0.1	0.6	1.3	2.3	2.6	1.1	0.7	0.7	1.4
2004	1.2	2.0	0.3	0.1	0.6	1.3	2.3	2.3	1.1	0.6	0.9	1.6
2003	1.2	1.3	*	*	0.8	1.3	2.6	2.2	1.0	0.7	0.6	0.9
2002	1.2	2.5	0.4	0.1	0.7	1.2	2.4	2.2	0.8	0.6	0.6	1.1
2001	0.9	1.7	0.3	*	0.5	1.1	2.1	1.6	0.6	0.3	0.7	0.8
2000	0.8	2.0	*	*	0.4	1.0	1.8	1.4	0.6	0.3	0.4	1.2
1999	0.9	1.2	0.3	*	0.5	1.1	1.8	1.4	0.6	0.4	0.6	1.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Hispanic, both sexes												
2006	0.8	*	*	*	0.7	0.7	1.4	1.5	1.1	*	*	*
2005	0.8	*	*	*	0.8	0.8	1.2	1.6	*	*	*	*
2004	0.8	*	*	*	0.7	1.0	1.3	1.4	1.1	*	*	*
2003	0.8	*	*	*	0.7	1.2	1.4	1.7	*	*	*	*
2002	0.8	*	*	*	0.7	1.0	1.5	1.3	*	*	*	*
2001	0.8	*	*	*	0.5	0.9	1.7	1.6	*	*	*	*
2000	0.6	*	*	*	0.5	0.9	1.2	0.8	*	*	*	*
1999	0.7	*	*	*	0.8	1.1	1.2	0.9	*	*	*	*
Hispanic, male												
2006	1.1	*	*	*	1.0	1.0	2.0	1.8	1.6	*	*	*
2005	1.0	*	*	*	1.0	1.0	1.6	1.8	*	*	*	*
2004	1.1	*	*	*	1.0	1.3	1.7	1.9	*	*	*	*
2003	1.3	*	*	*	1.0	1.7	2.1	2.6	*	*	*	*
2002	1.2	*	*	*	1.2	1.5	2.1	2.2	*	*	*	*
2001	1.1	*	*	*	0.7	1.4	2.6	2.1	*	*	*	*
2000	0.9	*	*	*	0.8	1.4	1.8	1.3	*	*	*	*
1999	1.1	*	*	*	1.3	1.6	1.9	1.6	*	*	*	*
Hispanic, female												
2006	0.4	*	*	*	*	*	0.7	1.1	*	*	*	*
2005	0.5	*	*	*	0.6	*	0.7	1.3	*	*	*	*
2004	0.5	*	*	*	*	0.6	0.9	*	*	*	*	*
2003	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	0.9	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Non-Hispanic 4/, both sexes												
2006	1.9	2.3	0.3	0.2	1.6	2.4	3.2	3.3	1.6	0.9	1.1	1.3
2005	1.7	2.7	0.5	0.2	1.2	2.1	3.0	3.2	1.4	0.9	0.8	1.5
2004	1.8	2.8	0.3	0.2	1.3	2.2	3.4	3.3	1.4	0.9	0.9	1.6
2003	1.9	2.2	0.2	0.1	1.5	2.3	3.9	3.2	1.3	0.7	0.7	1.1
2002	1.8	2.7	0.4	0.2	1.3	2.3	3.6	3.0	1.2	0.7	0.8	1.4
2001	1.6	2.4	0.3	0.1	1.1	2.0	3.3	2.6	0.9	0.5	0.9	0.9
2000	1.5	1.9	0.3	0.2	1.0	1.8	3.0	2.4	0.9	0.6	0.6	1.4
1999	1.5	1.6	0.3	0.1	1.2	2.0	3.1	2.3	0.9	0.6	0.8	1.4
Non-Hispanic 4/, male												
2006	2.4	2.7	0.4	0.2	2.3	3.2	3.8	4.1	1.9	1.1	1.5	2.2
2005	2.1	3.6	0.6	0.2	1.8	2.7	3.5	3.8	1.7	1.1	0.9	1.7
2004	2.4	3.3	0.3	0.2	1.9	3.1	4.3	4.1	1.7	1.1	1.0	1.7
2003	2.5	3.0	*	0.2	2.2	3.1	4.9	4.2	1.7	0.8	0.9	*
2002	2.3	2.8	0.3	0.2	1.8	3.2	4.7	3.6	1.6	1.0	0.9	2.1
2001	2.1	2.7	*	0.2	1.7	2.8	4.4	3.5	1.2	0.7	1.2	*
2000	2.0	1.6	0.4	0.3	1.5	2.6	4.2	3.2	1.3	0.8	1.0	1.7
1999	2.1	1.9	*	0.2	1.8	2.7	4.2	3.1	1.3	0.9	1.1	2.1
Non-Hispanic 4/, female												
2006	1.4	1.9	0.3	*	0.8	1.5	2.6	2.6	1.3	0.7	0.8	1.0
2005	1.4	1.9	0.4	*	0.6	1.5	2.5	2.7	1.2	0.7	0.7	1.4
2004	1.3	2.3	0.3	*	0.6	1.4	2.6	2.5	1.1	0.7	0.9	1.5
2003	1.3	1.3	*	*	0.8	1.5	2.8	2.3	1.0	0.7	0.6	1.0
2002	1.3	2.6	0.4	0.2	0.7	1.4	2.6	2.3	0.9	0.5	0.7	1.1
2001	1.0	2.0	0.4	*	0.5	1.2	2.2	1.7	0.6	0.4	0.7	0.8
2000	0.9	2.2	*	*	0.5	1.1	1.9	1.5	0.6	0.3	0.4	1.2
1999	0.9	*	*	*	0.5	1.2	2.0	1.5	0.6	0.4	0.6	1.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Non-Hispanic white, both sexes												
2006 5/	1.9	2.0	0.2	0.2	1.8	2.7	3.3	3.3	1.5	0.9	1.0	1.1
2005 5/	1.8	2.2	0.4	0.2	1.3	2.4	3.1	3.3	1.4	0.9	0.7	1.5
2004 5/	1.9	2.3	0.3	0.1	1.5	2.5	3.6	3.4	1.4	0.8	0.9	1.6
2003 5/	2.0	2.0	*	0.1	1.7	2.6	4.1	3.3	1.3	0.7	0.7	1.2
2002	1.8	1.8	0.4	0.2	1.4	2.4	3.7	2.9	1.2	0.7	0.7	1.5
2001	1.6	1.7	0.3	0.1	1.3	2.2	3.3	2.5	0.9	0.5	0.9	1.0
2000	1.4	1.5	0.2	0.1	1.1	1.9	3.0	2.3	0.8	0.5	0.7	1.3
1999	1.4	1.3	*	0.1	1.2	2.0	3.0	2.1	0.9	0.6	0.8	1.4
Non-Hispanic white, male												
2006 5/	2.4	2.4	*	0.2	2.6	3.6	3.8	3.9	1.7	0.9	1.2	1.9
2005 5/	2.1	3.0	0.4	0.2	1.9	2.9	3.5	3.7	1.6	1.0	0.7	1.8
2004 5/	2.4	2.7	*	0.2	2.2	3.4	4.5	4.1	1.6	1.0	0.9	1.9
2003 5/	2.5	2.4	*	0.2	2.4	3.4	5.1	4.0	1.5	0.7	0.9	*
2002	2.3	1.9	*	0.2	2.0	3.4	4.8	3.4	1.5	0.8	0.9	2.2
2001	2.1	2.0	*	0.2	1.8	3.0	4.3	3.3	1.1	0.7	1.2	*
2000	2.0	*	*	0.2	1.6	2.6	4.1	3.0	1.1	0.7	1.0	*
1999	2.0	*	*	0.2	1.8	2.7	4.0	2.6	1.1	0.8	1.1	2.1
Non-Hispanic white, female												
2006 5/	1.5	*	*	*	0.9	1.7	2.9	2.7	1.3	0.8	0.8	0.7
2005 5/	1.5	*	*	*	0.7	1.8	2.6	2.9	1.3	0.7	0.7	1.4
2004 5/	1.4	2.0	*	*	0.7	1.6	2.8	2.7	1.2	0.6	0.9	1.5
2003 5/	1.4	*	*	*	1.0	1.7	3.1	2.5	1.1	0.6	0.6	1.1
2002	1.3	1.8	*	0.2	0.8	1.5	2.7	2.4	0.9	0.6	0.6	1.1
2001	1.1	*	*	*	0.6	1.4	2.4	1.8	0.7	0.4	0.6	0.8
2000	0.9	*	*	*	0.6	1.2	1.9	1.5	0.6	0.3	0.4	1.2
1999	1.0	*	*	*	0.5	1.3	1.9	1.5	0.7	0.4	0.6	1.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Non-Hispanic black, both sexes												
2006 5/	1.9	3.5	0.8	*	0.9	1.6	3.4	4.1	2.4	1.3	*	*
2005 5/	1.8	4.9	0.9	*	0.9	1.7	3.4	3.8	1.7	*	*	*
2004 5/	1.9	4.9	*	0.4	0.9	1.9	3.4	4.0	1.8	1.3	*	*
2003 5/	1.8	3.3	*	*	0.8	1.8	3.9	4.1	2.0	1.3	*	*
2002	2.0	5.7	*	*	0.9	2.3	4.3	4.3	1.5	*	*	*
2001	1.8	5.9	*	*	0.7	1.7	4.0	3.8	1.2	*	*	*
2000	1.9	*	*	0.3	0.9	2.0	4.1	3.7	1.8	*	*	*
1999	2.0	*	*	*	1.3	2.3	4.3	4.2	1.2	*	*	*
Non-Hispanic black, male												
2006 5/	2.8	*	*	*	1.4	2.6	4.9	6.4	3.7	*	*	*
2005 5/	2.5	*	*	*	1.3	2.4	4.4	5.7	2.9	*	*	*
2004 5/	2.7	*	*	*	1.4	2.8	4.8	6.0	3.0	*	*	*
2003 5/	2.8	*	*	*	1.4	3.0	5.4	6.6	3.3	*	*	*
2002	2.9	*	*	*	1.3	3.4	6.2	6.5	2.6	*	*	*
2001	2.8	6.4	*	*	1.3	2.7	6.3	6.3	2.4	*	*	*
2000	2.9	*	*	*	1.6	3.2	6.1	6.0	3.0	*	*	*
1999	3.1	*	*	*	2.0	3.7	6.6	7.3	2.6	*	*	*
Non-Hispanic black, female												
2006 5/	1.1	*	*	*	*	0.8	2.0	2.1	1.3	*	*	*
2005 5/	1.2	*	*	*	*	1.0	2.6	2.2	*	*	*	*
2004 5/	1.1	*	*	*	*	1.0	2.2	2.3	*	*	*	*
2003 5/	1.0	*	*	*	*	*	2.5	2.0	*	*	*	*
2002	1.2	*	*	*	*	1.3	2.7	2.4	*	*	*	*
2001	0.9	*	*	*	*	0.9	1.9	1.5	*	*	*	*
2000	1.0	*	*	*	*	0.9	2.4	1.8	*	*	*	*
1999	0.9	*	*	*	*	1.0	2.4	1.6	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
All origins 3/, both sexes												
2006	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	*	*
2005	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	*	*
2004	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
2003	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	*	*
2002	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
All origins 3/, male												
2006	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
2005	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
2004	0.1	*	*	*	0.2	0.2	0.1	0.1	*	*	*	*
2003	0.1	*	*	*	0.3	0.2	0.1	0.1	*	*	*	*
2002	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.3	0.2	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
1999	0.2	*	*	*	0.6	0.2	0.2	0.1	*	*	*	*
All origins 3/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
Hispanic, both sexes												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	0.1	*	*	*	*	*	*	*	*	*	*	*
2005	0.1	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
Non-Hispanic 4/, both sexes												
2006	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2005	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	*	*
2004	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2003	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	*	*
2002	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
Non-Hispanic 4/, male												
2006	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
2005	0.1	*	*	*	0.3	0.1	*	0.1	*	*	*	*
2004	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
2003	0.1	*	*	*	0.3	0.2	0.1	0.1	*	*	*	*
2002	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.3	0.2	0.2	*	*	*	*	*
2000	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
1999	0.2	*	*	*	0.6	0.2	0.2	0.1	*	*	*	*
Non-Hispanic 4/, female												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
Non-Hispanic white, both sexes												
2006 5/	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*
2005 5/	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	*	*
2004 5/	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2003 5/	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2002	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
Non-Hispanic white, male												
2006 5/	0.1	*	*	*	0.2	*	0.1	*	*	*	*	*
2005 5/	0.1	*	*	*	0.3	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	0.2	*	0.2	*	*	*	*	*
2003 5/	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2002	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.3	0.2	0.2	*	*	*	*	*
2000	0.1	*	*	*	0.2	0.2	0.1	*	*	*	*	*
1999	0.2	*	*	*	0.5	0.2	0.2	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
Non-Hispanic black, both sexes												
2006 5/	0.1	*	*	*	0.4	*	*	*	*	*	*	*
2005 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	0.6	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	0.3	*	*	*	0.7	*	*	*	*	*	*	*
2005 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2003 5/	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	1.1	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)												
All origins 3/, both sexes												
2006	1.6	2.2	0.3	0.1	1.2	1.9	2.8	3.1	1.5	0.8	1.0	1.4
2005	1.5	2.5	0.4	0.1	1.0	1.8	2.7	3.0	1.3	0.8	0.8	1.4
2004	1.6	2.6	0.3	0.1	1.1	1.9	3.0	3.1	1.3	0.8	0.9	1.6
2003	1.7	1.9	0.2	0.1	1.2	2.0	3.5	3.0	1.2	0.7	0.7	1.1
2002	1.6	2.5	0.4	0.2	1.0	2.0	3.3	2.8	1.2	0.7	0.7	1.4
2001	1.4	2.1	0.3	0.1	0.9	1.7	3.0	2.5	0.8	0.5	0.8	1.0
2000	1.3	1.7	0.3	0.1	0.8	1.6	2.8	2.2	0.9	0.5	0.6	1.3
1999	1.3	1.4	0.3	0.1	0.8	1.7	2.7	2.1	0.9	0.5	0.7	1.3
All origins 3/, male												
2006	2.1	2.7	0.3	0.2	1.8	2.6	3.4	3.8	1.8	1.0	1.4	2.2
2005	1.8	3.2	0.5	0.2	1.3	2.2	3.1	3.5	1.6	1.0	0.9	1.5
2004	2.1	3.1	0.3	0.2	1.5	2.6	3.8	3.8	1.6	0.9	0.9	1.8
2003	2.2	2.4	*	0.1	1.7	2.7	4.5	3.9	1.5	0.7	0.8	1.4
2002	2.1	2.5	0.3	0.2	1.4	2.7	4.2	3.4	1.5	0.9	0.8	2.2
2001	1.9	2.4	*	0.2	1.3	2.4	4.0	3.3	1.1	0.7	1.0	*
2000	1.8	1.4	0.3	0.2	1.1	2.2	3.8	3.0	1.2	0.7	0.9	*
1999	1.8	1.7	*	0.1	1.2	2.4	3.8	2.8	1.2	0.7	0.9	1.8
All origins 3/, female												
2006	1.2	1.7	0.3	*	0.7	1.2	2.3	2.4	1.2	0.7	0.8	1.0
2005	1.2	1.8	0.3	*	0.6	1.3	2.2	2.5	1.1	0.6	0.7	1.4
2004	1.2	2.0	0.3	*	0.6	1.2	2.3	2.3	1.1	0.6	0.8	1.5
2003	1.2	1.3	*	*	0.7	1.3	2.5	2.2	1.0	0.7	0.6	0.9
2002	1.1	2.5	0.4	0.1	0.6	1.2	2.3	2.1	0.8	0.5	0.6	1.1
2001	0.9	1.7	0.3	*	0.5	1.1	2.0	1.6	0.6	0.3	0.6	0.8
2000	0.8	2.0	*	*	0.4	0.9	1.7	1.4	0.6	0.3	0.4	1.2
1999	0.8	1.2	0.3	*	0.4	1.1	1.7	1.4	0.6	0.4	0.5	1.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)												
Hispanic, both sexes												
2006	0.7	*	*	*	0.6	0.6	1.3	1.5	1.1	*	*	*
2005	0.7	*	*	*	0.6	0.7	1.1	1.6	*	*	*	*
2004	0.7	*	*	*	0.6	0.9	1.3	1.3	1.0	*	*	*
2003	0.8	*	*	*	0.5	1.1	1.3	1.6	*	*	*	*
2002	0.8	*	*	*	0.5	0.9	1.5	1.3	*	*	*	*
2001	0.7	*	*	*	0.4	0.9	1.6	1.5	*	*	*	*
2000	0.6	*	*	*	0.3	0.9	1.2	0.8	*	*	*	*
1999	0.6	*	*	*	0.5	1.0	1.2	0.9	*	*	*	*
Hispanic, male												
2006	1.0	*	*	*	0.7	0.9	1.8	1.8	*	*	*	*
2005	0.9	*	*	*	0.7	0.9	1.5	1.8	*	*	*	*
2004	1.0	*	*	*	0.8	1.1	1.6	1.7	*	*	*	*
2003	1.1	*	*	*	0.7	1.6	1.9	2.5	*	*	*	*
2002	1.1	*	*	*	0.9	1.3	2.0	2.2	*	*	*	*
2001	1.1	*	*	*	*	1.3	2.5	2.0	*	*	*	*
2000	0.8	*	*	*	*	1.3	1.7	1.3	*	*	*	*
1999	1.0	*	*	*	0.9	1.5	1.8	1.5	*	*	*	*
Hispanic, female												
2006	0.4	*	*	*	*	*	0.7	1.1	*	*	*	*
2005	0.5	*	*	*	*	*	0.7	1.3	*	*	*	*
2004	0.5	*	*	*	*	0.6	0.9	*	*	*	*	*
2003	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	0.9	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)												
Non-Hispanic 4/, both sexes												
2006	1.8	2.3	0.3	0.2	1.4	2.2	3.1	3.3	1.5	0.8	1.1	1.3
2005	1.7	2.7	0.4	0.1	1.0	2.0	2.9	3.2	1.4	0.8	0.8	1.5
2004	1.7	2.8	0.3	0.2	1.1	2.1	3.3	3.2	1.3	0.8	0.9	1.5
2003	1.8	2.2	0.2	0.1	1.3	2.2	3.8	3.2	1.3	0.7	0.7	1.1
2002	1.7	2.7	0.3	0.2	1.1	2.2	3.5	2.9	1.2	0.7	0.7	1.4
2001	1.5	2.4	0.3	0.1	1.0	1.9	3.2	2.5	0.8	0.5	0.8	0.9
2000	1.4	1.9	0.3	0.1	0.9	1.7	2.9	2.3	0.9	0.5	0.6	1.3
1999	1.4	1.6	0.3	0.1	0.8	1.8	2.9	2.2	0.9	0.5	0.7	1.3
Non-Hispanic 4/, male												
2006	2.3	2.7	0.3	0.2	2.0	3.1	3.7	4.1	1.8	1.0	1.4	2.2
2005	2.0	3.6	0.5	0.2	1.5	2.5	3.4	3.7	1.6	1.0	0.8	1.6
2004	2.2	3.3	0.3	0.2	1.7	3.0	4.2	4.0	1.6	0.9	0.9	1.7
2003	2.3	3.0	*	0.1	1.9	3.0	4.8	4.1	1.5	0.7	0.8	*
2002	2.2	2.8	0.3	0.2	1.5	3.0	4.5	3.5	1.5	0.9	0.8	2.1
2001	2.0	2.7	*	0.2	1.4	2.6	4.2	3.4	1.1	0.6	1.0	*
2000	1.9	1.6	0.4	0.2	1.2	2.4	4.1	3.1	1.2	0.7	0.9	*
1999	1.9	1.9	*	0.1	1.2	2.5	4.0	2.9	1.2	0.7	0.9	1.7
Non-Hispanic 4/, female												
2006	1.3	1.9	0.3	*	0.8	1.4	2.6	2.5	1.3	0.7	0.8	1.0
2005	1.3	1.9	0.3	*	0.6	1.5	2.5	2.7	1.1	0.7	0.7	1.4
2004	1.3	2.3	0.3	*	0.6	1.3	2.5	2.4	1.1	0.7	0.9	1.5
2003	1.3	1.3	*	*	0.8	1.4	2.7	2.3	1.0	0.7	0.6	1.0
2002	1.2	2.6	0.4	0.2	0.7	1.3	2.5	2.3	0.9	0.5	0.6	1.1
2001	1.0	2.0	0.4	*	0.5	1.2	2.2	1.6	0.6	0.3	0.7	0.8
2000	0.9	2.2	*	*	0.4	1.0	1.8	1.5	0.6	0.3	0.4	1.2
1999	0.9	*	*	*	0.4	1.2	1.9	1.4	0.6	0.4	0.5	1.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)												
Non-Hispanic white, both sexes												
2006 5/	1.9	2.0	*	0.1	1.6	2.6	3.3	3.3	1.4	0.8	0.9	1.1
2005 5/	1.7	2.2	0.4	0.1	1.2	2.3	3.0	3.2	1.4	0.8	0.7	1.5
2004 5/	1.8	2.3	0.3	0.1	1.3	2.4	3.5	3.3	1.3	0.7	0.9	1.6
2003 5/	1.9	2.0	*	0.1	1.6	2.5	4.0	3.2	1.2	0.6	0.7	1.2
2002	1.7	1.8	0.4	0.2	1.3	2.3	3.6	2.8	1.2	0.6	0.7	1.5
2001	1.5	1.7	0.3	0.1	1.1	2.0	3.2	2.5	0.8	0.4	0.8	0.9
2000	1.4	1.5	0.2	0.1	0.9	1.8	2.9	2.2	0.8	0.5	0.6	1.2
1999	1.3	1.3	*	0.1	0.9	1.9	2.8	2.0	0.9	0.5	0.7	1.3
Non-Hispanic white, male												
2006 5/	2.3	2.4	*	0.2	2.3	3.4	3.7	3.8	1.6	0.9	1.1	1.9
2005 5/	2.0	3.0	0.4	0.2	1.7	2.8	3.4	3.6	1.5	0.9	0.7	1.7
2004 5/	2.3	2.7	*	*	1.9	3.3	4.3	4.0	1.5	0.8	0.8	1.8
2003 5/	2.4	2.4	*	*	2.2	3.3	5.0	3.9	1.3	0.6	0.8	*
2002	2.2	1.9	*	0.2	1.7	3.2	4.6	3.3	1.4	0.7	0.8	2.2
2001	2.0	2.0	*	0.2	1.6	2.7	4.1	3.2	1.0	0.6	1.0	*
2000	1.8	*	*	0.2	1.4	2.4	4.0	2.9	1.0	0.6	0.9	*
1999	1.8	*	*	0.2	1.4	2.5	3.7	2.5	1.0	0.7	0.9	*
Non-Hispanic white, female												
2006 5/	1.4	*	*	*	0.8	1.7	2.8	2.7	1.3	0.7	0.8	0.7
2005 5/	1.4	*	*	*	0.7	1.7	2.6	2.9	1.3	0.7	0.7	1.4
2004 5/	1.4	2.0	*	*	0.7	1.5	2.7	2.6	1.2	0.6	0.9	1.5
2003 5/	1.4	*	*	*	0.9	1.7	3.0	2.5	1.1	0.6	0.6	1.1
2002	1.3	1.8	*	*	0.8	1.5	2.7	2.4	0.9	0.5	0.6	1.1
2001	1.1	*	*	*	0.6	1.3	2.3	1.7	0.7	0.3	0.6	0.8
2000	0.9	*	*	*	0.5	1.1	1.9	1.5	0.6	0.3	0.4	1.2
1999	0.9	*	*	*	0.4	1.3	1.9	1.5	0.7	0.4	0.6	1.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)												
Non-Hispanic black, both sexes												
2006 5/	1.8	3.5	0.8	*	0.5	1.4	3.2	4.0	2.4	1.3	*	*
2005 5/	1.7	4.9	0.8	*	0.6	1.6	3.4	3.8	1.6	*	*	*
2004 5/	1.8	4.9	*	0.3	0.7	1.7	3.4	3.9	1.7	1.2	*	*
2003 5/	1.8	3.3	*	*	0.5	1.7	3.8	4.1	1.9	1.3	*	*
2002	1.9	5.7	*	*	0.6	2.1	4.2	4.3	1.5	*	*	*
2001	1.7	5.9	*	*	0.5	1.7	3.9	3.7	1.1	*	*	*
2000	1.7	*	*	*	0.6	1.7	4.0	3.7	1.6	*	*	*
1999	1.8	*	*	*	0.6	2.0	4.2	4.1	1.2	*	*	*
Non-Hispanic black, male												
2006 5/	2.5	*	*	*	0.7	2.1	4.7	6.3	3.7	*	*	*
2005 5/	2.4	*	*	*	0.8	2.2	4.3	5.7	2.9	*	*	*
2004 5/	2.5	*	*	*	1.0	2.6	4.7	5.8	3.0	*	*	*
2003 5/	2.6	*	*	*	0.9	2.7	5.4	6.5	3.3	*	*	*
2002	2.7	*	*	*	0.8	3.0	6.0	6.5	2.6	*	*	*
2001	2.6	6.4	*	*	0.8	2.5	6.3	6.2	2.1	*	*	*
2000	2.6	*	*	*	1.0	2.7	5.9	5.9	2.7	*	*	*
1999	2.8	*	*	*	0.9	3.2	6.4	7.1	2.5	*	*	*
Non-Hispanic black, female												
2006 5/	1.0	*	*	*	*	0.7	2.0	2.0	1.3	*	*	*
2005 5/	1.1	*	*	*	*	1.0	2.5	2.2	*	*	*	*
2004 5/	1.1	*	*	*	*	0.9	2.2	2.3	*	*	*	*
2003 5/	1.0	*	*	*	*	*	2.4	1.9	*	*	*	*
2002	1.2	*	*	*	*	1.3	2.6	2.4	*	*	*	*
2001	0.9	*	*	*	*	0.9	1.9	1.5	*	*	*	*
2000	0.9	*	*	*	*	0.8	2.3	1.8	*	*	*	*
1999	0.9	*	*	*	*	0.9	2.3	1.6	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
All origins 3/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
Hispanic, both sexes												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
Non-Hispanic 4/, both sexes												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2006	0.0	*	*	*	*	*	*	*	*	*	*	*
2005	0.0	*	*	*	*	*	*	*	*	*	*	*
2004	0.0	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, female												
2006	*	*	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
Non-Hispanic white, both sexes												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2006 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	0.0	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
Non-Hispanic black, both sexes												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2006 5/	*	*	*	*	*	*	*	*	*	*	*	*
2005 5/	*	*	*	*	*	*	*	*	*	*	*	*
2004 5/	*	*	*	*	*	*	*	*	*	*	*	*
2003 5/	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
All origins 3/, both sexes												
2006	0.8	0.6	0.1	*	0.1	0.2	0.3	0.6	1.1	2.5	5.3	8.3
2005	0.9	*	*	*	0.1	0.2	0.3	0.7	1.2	2.6	5.8	9.1
2004	1.0	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.1	10.6
2003	1.0	*	0.2	*	0.1	0.2	0.4	0.7	1.4	2.9	6.2	11.2
2002	1.0	*	*	0.1	0.1	0.2	0.4	0.7	1.3	3.1	6.2	11.3
2001	1.1	0.6	0.2	*	0.1	0.2	0.4	0.7	1.5	3.1	6.9	12.8
2000	1.1	*	0.1	0.1	0.1	0.2	0.3	0.8	1.5	3.5	6.7	13.6
1999	1.0	1.4	0.2	*	0.1	0.2	0.3	0.6	1.3	3.2	6.6	12.7
All origins 3/, male												
2006	0.8	*	*	*	0.1	0.1	0.2	0.7	1.1	2.4	6.0	8.8
2005	0.8	*	*	*	*	0.1	0.3	0.6	1.3	2.9	6.3	10.8
2004	0.9	*	*	*	0.1	0.2	0.3	0.7	1.6	3.3	7.4	11.9
2003	0.9	*	*	*	0.1	0.2	0.4	0.7	1.5	3.2	6.9	12.9
2002	0.9	*	*	0.1	*	0.2	0.4	0.6	1.3	3.1	6.3	12.3
2001	1.0	*	*	*	*	0.2	0.4	0.7	1.6	3.2	7.7	14.5
2000	1.0	*	*	*	*	0.2	0.3	0.8	1.6	4.0	8.2	16.0
1999	1.0	1.6	*	*	0.1	0.2	0.3	0.5	1.4	3.4	7.8	16.5
All origins 3/, female												
2006	0.9	*	*	*	0.1	0.2	0.3	0.6	1.1	2.6	4.8	8.1
2005	1.0	*	*	*	*	0.2	0.4	0.7	1.1	2.3	5.5	8.3
2004	1.0	*	*	*	0.1	0.2	0.4	0.7	1.2	2.8	5.2	9.9
2003	1.0	*	*	*	*	0.2	0.4	0.6	1.3	2.7	5.7	10.5
2002	1.1	*	*	*	*	0.2	0.4	0.7	1.3	3.0	6.2	10.9
2001	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.3	12.1
2000	1.1	*	*	*	*	0.2	0.3	0.8	1.5	3.1	5.7	12.7
1999	1.1	1.2	*	*	*	0.2	0.3	0.6	1.2	3.0	5.9	11.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
Hispanic, both sexes												
2006	0.3	*	*	*	*	*	*	*	*	2.3	3.3	*
2005	0.4	*	*	*	*	*	*	0.6	*	2.3	4.9	*
2004	0.4	*	*	*	*	*	*	0.6	1.1	2.3	4.7	*
2003	0.4	*	*	*	*	*	0.4	*	1.0	3.1	3.7	*
2002	0.4	*	*	*	*	*	*	*	*	2.4	4.8	*
2001	0.4	*	*	*	*	*	*	0.6	*	2.2	5.2	*
2000	0.4	*	*	*	*	*	*	*	*	3.3	5.1	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2006	0.2	*	*	*	*	*	*	*	*	*	*	*
2005	0.3	*	*	*	*	*	*	*	*	*	*	*
2004	0.4	*	*	*	*	*	*	*	*	*	*	*
2003	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2006	0.3	*	*	*	*	*	*	*	*	*	*	*
2005	0.5	*	*	*	*	*	*	*	*	*	5.6	*
2004	0.5	*	*	*	*	*	*	*	*	*	*	*
2003	0.4	*	*	*	*	*	*	*	*	3.1	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
Non-Hispanic 4/, both sexes												
2006	0.9	*	*	*	0.1	0.2	0.3	0.7	1.2	2.5	5.4	8.6
2005	1.0	*	*	*	0.1	0.2	0.4	0.7	1.2	2.6	5.8	9.2
2004	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	3.1	6.1	10.8
2003	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	2.9	6.3	11.4
2002	1.1	*	*	0.1	0.1	0.2	0.4	0.7	1.3	3.1	6.2	11.6
2001	1.2	0.7	*	*	0.1	0.2	0.4	0.7	1.6	3.1	6.9	12.9
2000	1.2	*	*	0.1	0.1	0.2	0.3	0.8	1.6	3.5	6.7	13.7
1999	1.1	1.6	0.2	*	0.1	0.2	0.4	0.6	1.3	3.3	6.8	12.8
Non-Hispanic 4/, male												
2006	0.9	*	*	*	0.1	0.1	0.3	0.7	1.1	2.4	6.2	9.1
2005	0.9	*	*	*	*	0.1	0.3	0.7	1.3	3.0	6.4	11.2
2004	1.0	*	*	*	0.1	0.2	0.3	0.7	1.6	3.4	7.4	12.5
2003	1.0	*	*	*	0.1	0.2	0.4	0.7	1.5	3.2	7.0	13.2
2002	1.0	*	*	0.1	*	0.3	0.4	0.7	1.4	3.1	6.3	12.9
2001	1.1	*	*	*	*	0.2	0.4	0.8	1.7	3.3	7.7	14.5
2000	1.1	*	*	*	*	0.3	0.4	0.8	1.6	4.0	8.2	16.1
1999	1.0	1.8	*	*	*	0.2	0.4	0.6	1.5	3.5	7.9	16.8
Non-Hispanic 4/, female												
2006	1.0	*	*	*	*	0.2	0.3	0.6	1.2	2.6	4.9	8.3
2005	1.0	*	*	*	*	0.2	0.4	0.7	1.1	2.3	5.4	8.4
2004	1.1	*	*	*	0.1	0.2	0.4	0.7	1.2	2.8	5.3	10.1
2003	1.1	*	*	*	*	0.2	0.4	0.6	1.3	2.7	5.8	10.6
2002	1.2	*	*	*	*	0.2	0.4	0.7	1.3	3.1	6.2	11.1
2001	1.2	*	*	*	*	0.2	0.4	0.7	1.5	3.0	6.4	12.2
2000	1.2	*	*	*	*	0.2	0.3	0.8	1.6	3.1	5.7	12.7
1999	1.2	1.4	*	*	*	0.2	0.4	0.6	1.2	3.1	6.1	11.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
Non-Hispanic white, both sexes												
2006 5/	1.0	*	*	*	0.1	0.2	0.3	0.6	1.0	2.4	5.2	8.5
2005 5/	1.0	*	*	*	0.1	0.2	0.3	0.6	1.1	2.3	5.7	9.4
2004 5/	1.1	*	*	*	0.1	0.2	0.3	0.7	1.3	2.9	5.9	10.6
2003 5/	1.1	*	*	*	0.1	0.2	0.4	0.6	1.3	2.8	6.2	11.2
2002	1.1	*	*	*	0.1	0.2	0.3	0.6	1.2	3.0	6.1	11.5
2001	1.2	*	*	*	0.1	0.2	0.3	0.6	1.4	2.8	6.8	12.2
2000	1.2	*	*	*	*	0.2	0.3	0.8	1.4	3.3	6.5	13.2
1999	1.1	1.3	*	*	0.1	0.2	0.3	0.5	1.2	3.1	6.5	12.1
Non-Hispanic white, male												
2006 5/	0.9	*	*	*	*	*	0.3	0.6	1.0	2.3	6.1	9.1
2005 5/	1.0	*	*	*	*	0.2	0.3	0.6	1.2	2.8	6.3	11.6
2004 5/	1.1	*	*	*	*	0.2	0.3	0.7	1.5	3.3	7.4	12.3
2003 5/	1.1	*	*	*	*	0.2	0.4	0.6	1.4	3.1	7.1	13.1
2002	1.0	*	*	*	*	0.3	0.4	0.6	1.3	2.9	6.3	12.4
2001	1.0	*	*	*	*	0.2	0.3	0.6	1.4	2.9	7.8	13.5
2000	1.2	*	*	*	*	0.2	0.4	0.8	1.4	3.7	8.2	15.9
1999	1.1	*	*	*	*	0.2	0.3	0.5	1.3	3.3	7.7	15.8
Non-Hispanic white, female												
2006 5/	1.0	*	*	*	*	0.2	0.3	0.5	1.0	2.5	4.6	8.2
2005 5/	1.0	*	*	*	*	0.2	0.4	0.6	1.1	1.9	5.3	8.3
2004 5/	1.1	*	*	*	*	0.2	0.3	0.6	1.1	2.6	5.0	9.9
2003 5/	1.1	*	*	*	*	*	0.3	0.6	1.1	2.5	5.5	10.4
2002	1.2	*	*	*	*	*	0.3	0.6	1.1	3.0	6.0	11.1
2001	1.3	*	*	*	*	0.2	0.3	0.6	1.4	2.7	6.2	11.6
2000	1.2	*	*	*	*	0.2	0.3	0.7	1.4	3.0	5.4	12.0
1999	1.2	*	*	*	*	*	0.3	0.6	1.1	3.0	5.7	10.7

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
Non-Hispanic black, both sexes												
2006 5/	1.1	*	*	*	*	*	0.6	1.4	2.7	4.1	9.0	10.4
2005 5/	1.1	*	*	*	*	*	0.5	1.6	2.5	5.5	9.0	9.3
2004 5/	1.2	*	*	*	*	*	0.8	1.3	2.6	5.1	9.5	14.1
2003 5/	1.2	*	*	*	*	*	0.6	1.4	2.8	4.6	9.0	15.2
2002	1.2	*	*	*	*	*	0.9	1.7	2.7	4.7	8.5	14.2
2001	1.4	*	*	*	*	0.4	0.9	1.6	3.4	6.4	9.3	23.7
2000	1.4	*	*	*	*	0.4	0.6	1.4	3.6	6.2	10.6	21.3
1999	1.3	*	*	*	*	*	0.6	1.3	2.6	5.2	10.6	20.4
Non-Hispanic black, male												
2006 5/	1.0	*	*	*	*	*	*	1.4	2.7	4.0	9.7	*
2005 5/	1.0	*	*	*	*	*	*	1.5	3.1	5.0	9.5	*
2004 5/	1.0	*	*	*	*	*	*	1.0	3.0	4.7	10.3	*
2003 5/	1.0	*	*	*	*	*	*	1.9	2.9	3.9	7.8	*
2002	1.1	*	*	*	*	*	*	1.4	2.6	5.4	7.3	*
2001	1.5	*	*	*	*	*	1.0	1.9	4.3	7.9	9.0	30.0
2000	1.2	*	*	*	*	*	*	1.3	3.2	7.8	10.2	*
1999	1.2	*	*	*	*	*	*	1.4	2.9	6.0	11.9	23.8
Non-Hispanic black, female												
2006 5/	1.2	*	*	*	*	*	0.8	1.3	2.8	4.2	8.5	10.1
2005 5/	1.3	*	*	*	*	*	0.7	1.6	2.1	5.8	8.7	9.2
2004 5/	1.4	*	*	*	*	*	1.0	1.4	2.4	5.4	9.0	13.7
2003 5/	1.3	*	*	*	*	*	0.8	1.0	2.6	5.0	9.6	15.3
2002	1.4	*	*	*	*	*	1.1	1.9	2.8	4.2	9.1	11.7
2001	1.4	*	*	*	*	*	0.7	1.4	2.7	5.3	9.4	21.4
2000	1.6	*	*	*	*	*	*	1.5	3.9	5.2	10.8	22.2
1999	1.4	*	*	*	*	*	*	1.2	2.4	4.7	9.8	19.1

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/												
All origins 3/, both sexes												
2006	7.4	*	*	*	0.4	1.6	7.4	17.6	19.4	14.7	9.7	4.9
2005	7.3	*	*	*	0.4	1.3	7.5	17.6	19.5	15.1	9.2	4.6
2004	7.2	*	*	*	0.3	1.5	7.6	17.3	18.7	15.7	9.3	4.3
2003	7.1	*	*	*	0.3	1.5	8.1	17.3	18.5	15.1	9.2	4.1
2002	7.0	*	*	*	0.3	1.5	8.1	16.8	18.4	15.5	9.3	4.4
2001	7.1	*	*	*	0.3	1.6	8.3	17.2	18.2	15.5	9.6	5.0
2000	7.0	*	*	*	0.2	1.6	8.5	16.3	18.7	15.8	9.9	5.4
1999	7.0	*	*	*	0.3	1.6	8.5	16.4	18.7	15.9	10.6	5.5
All origins 3/, male												
2006	11.2	*	*	*	0.6	2.4	10.3	26.5	31.1	24.9	17.1	9.5
2005	11.1	*	*	*	0.6	2.0	10.6	26.7	31.6	25.0	17.0	9.3
2004	11.0	*	*	*	0.5	2.3	11.0	26.4	30.2	26.5	17.1	9.0
2003	10.9	*	*	*	0.5	2.3	11.4	27.0	29.8	25.4	16.8	7.8
2002	10.8	*	*	*	0.4	2.3	11.5	26.3	30.0	26.0	16.6	9.7
2001	10.8	*	*	*	0.4	2.3	11.9	26.6	29.7	26.3	17.2	11.6
2000	10.9	*	*	*	0.4	2.5	12.5	25.9	30.0	27.1	18.8	11.9
1999	10.9	*	*	*	0.4	2.3	12.6	26.2	30.5	27.4	19.4	11.8
All origins 3/, female												
2006	3.7	*	*	*	0.1	0.7	4.5	8.9	8.5	6.1	4.7	2.7
2005	3.6	*	*	*	0.1	0.7	4.3	8.8	8.4	6.8	3.9	2.5
2004	3.5	*	*	*	0.2	0.7	4.2	8.6	8.1	6.6	3.9	2.3
2003	3.4	*	*	*	0.1	0.6	4.7	7.9	8.1	6.5	4.1	2.5
2002	3.4	*	*	*	*	0.7	4.7	7.7	7.6	6.8	4.5	2.1
2001	3.4	*	*	*	*	0.9	4.8	8.1	7.5	6.6	4.6	2.3
2000	3.2	*	*	*	*	0.7	4.5	7.0	8.3	6.6	4.1	2.7
1999	3.2	*	*	*	0.1	0.9	4.5	7.0	7.7	6.5	4.8	3.0

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/												
Hispanic, both sexes												
2006	6.3	*	*	*	0.5	2.3	8.0	20.2	26.3	21.7	19.0	9.4
2005	6.2	*	*	*	0.6	2.0	8.4	20.0	25.0	25.9	14.4	*
2004	5.8	*	*	*	0.6	1.9	8.1	20.6	22.5	20.7	17.1	*
2003	6.1	*	*	*	0.6	1.8	8.3	20.2	27.6	23.0	17.8	*
2002	6.2	*	*	*	0.5	2.0	8.7	21.2	26.4	27.0	14.5	*
2001	6.4	*	*	*	0.5	2.1	9.0	22.5	28.3	26.1	20.4	*
2000	6.6	*	*	*	0.5	2.1	10.4	22.3	29.5	28.9	19.0	*
1999	6.4	*	*	*	0.4	2.1	10.2	22.5	28.7	26.2	18.7	14.0
Hispanic, male												
2006	10.2	*	*	*	0.8	3.7	12.3	33.5	46.1	41.4	34.5	*
2005	10.3	*	*	*	1.1	3.2	13.1	33.7	45.0	49.8	29.7	*
2004	9.6	*	*	*	1.0	3.2	13.1	36.0	39.3	37.7	33.3	*
2003	9.9	*	*	*	0.9	2.8	13.2	35.2	50.5	40.4	34.9	*
2002	10.3	*	*	*	0.8	3.4	14.2	36.3	48.6	52.8	26.7	*
2001	10.7	*	*	*	0.7	3.3	14.7	38.7	51.8	52.3	38.0	*
2000	11.1	*	*	*	0.8	3.7	16.8	39.1	54.8	58.7	42.3	*
1999	10.7	*	*	*	0.7	3.4	16.7	39.3	52.8	51.2	36.0	*
Hispanic, female												
2006	2.2	*	*	*	*	*	3.2	6.6	8.0	5.7	8.2	*
2005	1.9	*	*	*	*	0.6	3.1	6.0	6.7	6.6	*	*
2004	1.8	*	*	*	*	*	2.5	5.0	7.3	7.1	5.9	*
2003	1.9	*	*	*	*	0.6	2.9	5.2	6.9	9.1	6.1	*
2002	1.8	*	*	*	*	*	2.7	6.2	6.4	6.5	6.2	*
2001	2.0	*	*	*	*	0.8	2.9	6.7	7.4	5.4	8.5	*
2000	1.7	*	*	*	*	*	3.5	5.8	7.1	5.5	*	*
1999	1.9	*	*	*	*	*	3.3	6.2	7.4	6.5	7.3	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/												
Non-Hispanic 4/, both sexes												
2006	7.5	*	*	*	0.3	1.4	7.3	17.2	18.7	14.1	9.1	4.6
2005	7.4	*	*	*	0.3	1.2	7.3	17.2	19.0	14.2	8.9	4.4
2004	7.4	*	*	*	0.3	1.4	7.5	16.9	18.3	15.2	8.8	4.4
2003	7.2	*	*	*	0.2	1.4	8.0	16.9	17.7	14.5	8.7	4.1
2002	7.1	*	*	*	0.2	1.4	7.9	16.3	17.6	14.6	9.0	4.1
2001	7.1	*	*	*	0.2	1.5	8.2	16.5	17.2	14.7	9.0	5.0
2000	7.0	*	*	*	0.2	1.5	8.2	15.6	17.8	14.9	9.4	5.3
1999	7.0	*	*	*	0.2	1.5	8.2	15.7	17.8	15.2	10.2	5.2
Non-Hispanic 4/, male												
2006	11.3	*	*	*	0.5	2.0	9.8	25.5	29.7	23.5	15.9	8.8
2005	11.2	*	*	*	0.5	1.7	10.1	25.7	30.2	23.0	16.2	8.6
2004	11.2	*	*	*	0.4	2.1	10.6	25.1	29.2	25.6	16.2	8.9
2003	11.0	*	*	*	0.4	2.2	11.1	26.0	27.9	24.2	15.8	7.8
2002	10.7	*	*	*	0.3	2.0	11.0	25.0	28.3	24.0	16.0	9.1
2001	10.8	*	*	*	0.4	2.1	11.4	25.2	27.7	24.4	16.2	11.4
2000	10.7	*	*	*	0.3	2.2	11.8	24.4	27.9	25.0	17.6	11.7
1999	10.8	*	*	*	0.3	2.0	11.9	24.8	28.7	25.8	18.7	11.1
Non-Hispanic 4/, female												
2006	3.9	*	*	*	0.1	0.7	4.8	9.2	8.5	6.2	4.5	2.7
2005	3.8	*	*	*	*	0.7	4.5	9.0	8.5	6.8	3.9	2.5
2004	3.7	*	*	*	0.2	0.8	4.4	8.9	8.1	6.5	3.8	2.3
2003	3.6	*	*	*	*	0.6	4.9	8.2	8.2	6.3	4.0	2.5
2002	3.6	*	*	*	*	0.8	5.0	7.8	7.7	6.8	4.4	1.9
2001	3.6	*	*	*	*	0.9	5.0	8.2	7.5	6.7	4.4	2.3
2000	3.4	*	*	*	*	0.8	4.7	7.1	8.4	6.6	4.2	2.6
1999	3.4	*	*	*	*	1.0	4.6	7.0	7.7	6.5	4.7	2.9

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/												
Non-Hispanic white, both sexes												
2006 5/	8.0	*	*	*	0.3	1.3	7.5	17.8	18.8	14.1	9.4	4.7
2005 5/	7.9	*	*	*	0.3	1.1	7.5	17.5	19.0	14.2	8.9	4.4
2004 5/	7.7	*	*	*	0.3	1.4	7.8	17.2	17.9	14.8	8.9	4.5
2003 5/	7.5	*	*	*	0.2	1.4	8.1	16.9	17.1	14.4	8.8	4.1
2002	7.3	*	*	*	0.2	1.3	8.1	16.1	16.8	14.4	9.0	4.0
2001	7.2	*	*	*	0.2	1.4	8.0	15.9	16.4	14.3	9.0	4.7
2000	7.0	*	*	*	0.2	1.4	7.9	14.9	16.8	14.5	9.3	5.2
1999	6.9	*	*	*	0.2	1.4	7.7	14.8	16.6	14.5	9.8	5.0
Non-Hispanic white, male												
2006 5/	12.0	*	*	*	0.5	2.0	10.2	26.2	29.4	23.1	16.3	8.9
2005 5/	11.8	*	*	*	0.5	1.6	10.5	25.8	30.0	22.5	15.9	8.7
2004 5/	11.7	*	*	*	0.4	2.0	10.9	25.3	28.3	24.4	16.3	9.3
2003 5/	11.4	*	*	*	0.4	2.2	11.3	25.5	26.7	24.0	15.7	7.7
2002	11.1	*	*	*	0.3	1.9	11.1	24.8	26.9	23.0	15.8	8.7
2001	10.8	*	*	*	0.3	2.0	11.0	24.2	26.0	23.2	16.0	10.8
2000	10.8	*	*	*	0.3	2.1	11.4	23.3	26.2	23.9	17.3	11.4
1999	10.7	*	*	*	0.3	2.0	11.3	23.4	26.6	24.1	18.0	10.5
Non-Hispanic white, female												
2006 5/	4.2	*	*	*	*	0.7	4.8	9.5	8.7	6.4	4.6	2.7
2005 5/	4.0	*	*	*	*	0.7	4.5	9.3	8.5	7.0	4.0	2.5
2004 5/	3.9	*	*	*	*	0.8	4.6	9.1	8.0	6.6	3.9	2.4
2003 5/	3.8	*	*	*	*	0.7	4.9	8.3	8.0	6.3	4.2	2.5
2002	3.7	*	*	*	*	0.7	5.0	7.7	7.3	7.0	4.4	2.1
2001	3.6	*	*	*	*	0.9	5.0	7.6	7.4	6.8	4.4	2.2
2000	3.4	*	*	*	*	0.8	4.4	6.5	7.8	6.5	4.1	2.7
1999	3.3	*	*	*	*	0.9	4.2	6.4	7.2	6.4	4.5	2.8

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/												
Non-Hispanic black, both sexes												
2006 5/	5.6	*	*	*	*	1.0	5.4	14.9	20.7	15.8	8.2	*
2005 5/	6.1	*	*	*	*	0.9	5.8	17.3	22.0	16.7	9.9	5.5
2004 5/	6.3	*	*	*	*	1.0	5.8	17.4	23.9	20.3	7.8	*
2003 5/	6.5	*	*	*	*	1.2	7.0	18.7	24.5	17.1	9.1	*
2002	6.6	*	*	*	*	1.4	6.9	18.9	25.4	18.4	11.0	*
2001	7.5	*	*	*	*	1.4	8.9	22.8	26.2	20.7	11.2	*
2000	7.6	*	*	*	*	1.8	9.3	23.1	26.9	21.2	11.6	6.4
1999	8.0	*	*	*	*	1.4	10.1	23.7	29.6	23.4	14.8	7.8
Non-Hispanic black, male												
2006 5/	8.7	*	*	*	*	1.3	7.2	23.4	36.4	30.2	15.6	*
2005 5/	9.4	*	*	*	*	1.2	7.8	27.7	37.2	31.0	21.0	*
2004 5/	10.0	*	*	*	*	1.7	9.0	27.5	42.2	40.0	14.8	*
2003 5/	10.3	*	*	*	*	1.9	10.0	31.6	42.4	30.9	19.3	*
2002	10.3	*	*	*	*	2.1	9.6	30.0	43.7	36.0	21.7	*
2001	11.9	*	*	*	*	1.9	13.5	36.2	46.8	40.5	22.2	*
2000	11.7	*	*	*	*	2.6	13.7	36.6	44.7	40.3	24.4	*
1999	12.5	*	*	*	*	1.8	14.7	38.0	51.6	45.7	28.8	*
Non-Hispanic black, female												
2006 5/	2.9	*	*	*	*	0.7	3.9	7.7	8.1	5.5	3.9	*
2005 5/	3.1	*	*	*	*	*	3.9	8.4	9.7	6.4	3.7	*
2004 5/	2.9	*	*	*	*	*	3.0	8.8	9.3	6.3	3.9	*
2003 5/	3.0	*	*	*	*	*	4.4	7.6	10.1	7.3	3.3	*
2002	3.3	*	*	*	*	*	4.5	9.5	10.8	5.9	5.0	*
2001	3.5	*	*	*	*	0.9	4.8	11.2	9.5	6.7	5.1	*
2000	3.9	*	*	*	*	1.1	5.4	11.5	12.5	7.8	4.3	*
1999	4.0	*	*	*	*	1.0	6.1	11.4	11.9	7.7	7.0	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/												
All origins 3/, both sexes												
2006	12.8	1.1	0.2	0.1	8.6	16.8	23.3	26.7	12.2	5.3	6.0	8.1
2005	11.3	0.9	0.2	0.1	7.4	14.3	21.3	23.6	10.7	4.8	5.4	7.6
2004	10.5	0.7	0.2	0.2	7.0	12.6	20.9	21.7	9.1	4.3	4.8	6.3
2003	9.9	0.6	0.2	0.1	6.4	12.2	20.6	20.0	8.0	4.1	4.2	6.0
2002	9.0	0.7	0.2	0.1	5.5	11.1	19.7	17.9	6.9	3.6	3.8	5.7
2001	7.6	0.5	0.2	0.1	4.5	9.4	17.0	14.8	5.4	3.0	3.5	5.1
2000	7.0	*	*	0.1	4.0	8.8	16.0	13.2	4.9	2.6	3.5	5.7
1999	6.9	0.6	0.2	0.1	3.5	8.9	15.7	12.6	4.9	3.0	3.8	4.8
All origins 3/, male												
2006	16.6	1.1	0.2	0.2	12.7	23.4	29.4	33.3	14.9	5.8	6.9	10.7
2005	14.5	*	0.3	0.1	10.7	19.4	26.6	29.4	13.1	5.3	6.7	9.9
2004	13.4	*	*	0.1	10.2	17.5	26.0	27.0	10.7	4.7	5.3	8.6
2003	12.9	*	0.3	0.1	9.2	16.6	26.5	25.7	9.4	4.6	4.9	7.8
2002	11.8	1.0	*	0.1	7.8	15.5	25.6	23.0	8.1	3.7	4.2	7.3
2001	10.2	*	0.3	0.1	6.6	13.2	22.7	19.7	6.3	3.5	3.8	5.7
2000	9.5	*	*	0.1	5.8	12.5	21.8	18.2	5.4	2.8	3.8	7.5
1999	9.4	*	*	*	5.0	12.7	21.9	17.5	6.0	3.4	3.8	6.2
All origins 3/, female												
2006	9.1	1.0	*	0.1	4.2	10.0	17.1	20.3	9.8	4.9	5.4	6.8
2005	8.2	*	*	0.1	3.8	8.9	16.0	18.1	8.4	4.3	4.5	6.6
2004	7.6	*	*	0.2	3.6	7.7	15.8	16.5	7.7	3.9	4.4	5.2
2003	7.0	*	*	0.1	3.3	7.6	14.8	14.6	6.7	3.7	3.7	5.2
2002	6.3	*	0.3	*	3.0	6.6	13.9	13.1	5.7	3.4	3.5	4.9
2001	5.1	*	*	0.1	2.4	5.6	11.5	10.0	4.6	2.6	3.3	4.9
2000	4.6	*	*	*	2.1	5.1	10.2	8.5	4.4	2.5	3.2	4.9
1999	4.4	*	*	*	2.0	5.0	9.7	8.0	3.8	2.7	3.8	4.3

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/												
Hispanic, both sexes												
2006	6.5	*	*	*	3.6	8.1	13.2	16.3	9.0	3.2	3.5	*
2005	6.1	*	*	*	3.9	7.4	12.5	15.1	7.8	3.7	4.0	*
2004	5.5	*	*	*	3.3	6.3	12.1	13.1	8.3	3.1	3.5	*
2003	5.9	*	*	*	3.5	7.2	13.3	14.9	6.5	3.1	3.7	*
2002	5.5	*	*	*	3.2	6.2	13.4	13.8	5.6	3.0	3.3	*
2001	4.7	*	*	*	2.4	5.9	11.8	11.0	4.8	2.6	*	*
2000	4.8	*	*	*	2.2	6.9	12.7	10.6	4.6	1.9	*	*
1999	5.8	*	*	*	2.7	8.1	15.0	13.4	5.1	2.7	*	*
Hispanic, male												
2006	9.3	*	*	*	5.1	11.5	18.7	23.2	14.5	4.4	*	*
2005	8.9	*	*	*	5.8	10.8	17.9	22.2	13.0	4.9	*	*
2004	7.8	*	*	*	4.6	9.3	17.2	18.3	12.5	4.8	*	*
2003	8.7	*	*	*	5.2	10.5	19.5	22.1	10.5	5.2	*	*
2002	8.2	*	*	*	4.7	9.3	19.9	21.1	8.7	*	*	*
2001	7.0	*	*	*	3.3	8.7	17.8	17.3	7.9	*	*	*
2000	7.4	*	*	*	3.4	10.5	19.6	17.1	6.5	*	*	*
1999	9.2	*	*	*	4.2	12.7	23.8	21.9	9.0	4.6	*	*
Hispanic, female												
2006	3.4	*	*	*	2.0	4.0	6.9	9.2	4.0	*	*	*
2005	3.0	*	*	*	1.9	3.3	6.4	7.9	3.1	2.7	*	*
2004	2.9	*	*	*	1.8	2.8	6.5	7.8	4.6	*	*	*
2003	2.9	*	*	*	1.5	3.3	6.5	7.8	2.8	*	*	*
2002	2.6	*	*	*	1.4	2.5	6.3	6.6	2.9	*	*	*
2001	2.2	*	*	*	1.4	2.8	5.3	4.8	*	*	*	*
2000	2.1	*	*	*	0.9	2.7	5.2	4.2	3.0	*	*	*
1999	2.2	*	*	*	1.0	2.8	5.7	5.2	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/												
Non-Hispanic 4/, both sexes												
2006	13.9	1.1	0.3	0.2	9.6	18.9	25.0	27.8	12.5	5.5	6.2	8.2
2005	12.1	0.9	0.2	0.1	8.0	15.9	22.7	24.4	10.9	4.8	5.5	7.7
2004	11.2	0.9	0.2	0.2	7.7	14.1	22.3	22.5	9.2	4.3	4.8	6.3
2003	10.4	0.7	0.2	0.1	6.9	13.2	21.6	20.4	8.1	4.2	4.2	6.0
2002	9.5	0.8	0.2	0.1	5.9	12.2	20.5	18.2	6.9	3.6	3.8	5.8
2001	8.0	0.7	0.2	0.1	4.9	10.1	17.6	15.0	5.3	3.0	3.5	5.2
2000	7.2	*	*	0.1	4.3	9.1	16.3	13.3	4.8	2.7	3.6	5.7
1999	6.9	*	0.2	0.1	3.6	9.0	15.6	12.4	4.8	3.0	3.9	4.9
Non-Hispanic 4/, male												
2006	17.9	*	*	0.2	14.4	26.6	31.3	34.3	14.8	5.9	7.1	10.8
2005	15.4	*	0.3	0.1	11.7	21.7	28.0	30.0	13.0	5.3	6.8	10.0
2004	14.3	*	*	0.1	11.4	19.5	27.4	27.8	10.5	4.6	5.3	8.6
2003	13.5	*	*	*	10.1	18.0	27.5	25.9	9.2	4.5	4.9	7.8
2002	12.3	*	*	0.1	8.4	17.0	26.3	23.0	8.0	3.7	4.1	7.5
2001	10.6	*	*	0.1	7.3	14.1	23.2	19.7	6.0	3.5	3.9	5.8
2000	9.7	*	*	0.1	6.3	12.8	21.9	17.9	5.2	2.8	3.9	7.5
1999	9.3	*	*	*	5.1	12.7	21.4	16.8	5.6	3.3	4.0	6.3
Non-Hispanic 4/, female												
2006	10.1	*	*	0.1	4.7	11.3	18.8	21.5	10.3	5.1	5.5	7.0
2005	9.0	*	*	0.2	4.2	10.1	17.4	19.1	8.9	4.4	4.6	6.7
2004	8.3	*	*	0.2	3.9	8.7	17.2	17.4	7.9	4.1	4.5	5.3
2003	7.6	*	*	*	3.7	8.4	15.9	15.2	7.0	3.9	3.7	5.3
2002	6.9	*	*	*	3.3	7.5	14.9	13.6	5.9	3.5	3.6	5.1
2001	5.5	*	*	*	2.6	6.1	12.2	10.5	4.8	2.6	3.3	5.0
2000	4.9	*	*	*	2.3	5.5	10.8	8.7	4.5	2.5	3.3	5.0
1999	4.6	*	*	*	2.1	5.4	10.0	8.2	3.9	2.8	3.9	4.4

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/												
Non-Hispanic white, both sexes												
2006 5/	14.9	*	0.2	0.2	11.9	22.5	27.0	27.9	11.8	5.4	6.4	8.4
2005 5/	13.1	0.9	*	0.1	9.9	18.5	24.6	24.5	10.5	4.9	5.7	8.1
2004 5/	12.1	*	*	0.2	9.5	16.5	24.2	22.6	8.8	4.4	5.0	6.4
2003 5/	11.2	*	*	0.1	8.5	15.2	23.0	20.3	7.9	4.2	4.5	6.3
2002	10.0	*	*	0.1	7.1	13.8	21.6	17.8	6.6	3.7	3.9	5.9
2001	8.3	*	*	0.1	5.9	11.3	18.1	14.2	5.1	3.1	3.7	5.2
2000	7.4	*	*	0.1	5.1	9.9	16.4	12.3	4.7	2.7	3.7	5.7
1999	6.9	*	*	0.1	4.1	9.5	15.4	11.4	4.5	3.0	4.1	4.9
Non-Hispanic white, male												
2006 5/	18.9	*	*	0.2	17.9	31.2	33.3	33.2	12.8	5.3	7.3	11.3
2005 5/	16.4	*	*	*	14.5	25.3	30.2	29.1	11.6	5.1	6.9	10.7
2004 5/	15.3	*	*	0.2	14.0	22.8	29.5	27.3	9.2	4.3	5.4	8.7
2003 5/	14.3	*	*	*	12.5	20.8	29.0	25.1	8.4	4.2	5.1	7.8
2002	12.8	*	*	*	10.1	19.2	27.5	21.7	7.0	3.5	4.1	7.8
2001	10.8	*	*	*	8.6	15.8	23.7	17.8	5.1	3.4	3.8	5.7
2000	9.8	*	*	0.2	7.5	13.8	21.9	16.1	4.6	2.7	4.1	7.2
1999	9.2	*	*	*	5.8	13.2	21.0	14.7	4.7	3.2	4.0	6.6
Non-Hispanic white, female												
2006 5/	11.1	*	*	*	5.7	13.7	20.6	22.6	10.9	5.5	5.7	7.1
2005 5/	9.8	*	*	*	5.0	11.7	19.0	20.0	9.4	4.8	4.8	7.0
2004 5/	9.1	*	*	0.2	4.7	10.1	18.8	18.0	8.4	4.5	4.7	5.4
2003 5/	8.2	*	*	*	4.4	9.5	17.0	15.7	7.4	4.1	4.0	5.7
2002	7.4	*	*	*	3.9	8.4	15.7	13.9	6.3	3.8	3.8	5.1
2001	5.8	*	*	*	3.0	6.8	12.5	10.6	5.1	2.8	3.6	4.9
2000	5.1	*	*	*	2.6	6.0	10.8	8.6	4.8	2.7	3.5	5.1
1999	4.8	*	*	*	2.4	5.7	9.9	8.1	4.2	2.9	4.1	4.2

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/												
Non-Hispanic black, both sexes												
2006 5/	12.5	*	*	*	2.6	10.5	23.8	35.9	21.5	7.1	4.4	6.6
2005 5/	10.8	*	*	*	2.2	9.7	20.5	31.5	17.7	5.2	4.3	*
2004 5/	9.7	*	*	*	2.2	8.2	19.4	28.7	14.7	4.7	3.8	*
2003 5/	9.5	*	*	*	1.9	8.3	21.4	27.5	11.7	4.9	2.2	*
2002	9.4	*	*	*	2.1	8.7	21.2	27.3	11.7	2.9	2.6	*
2001	8.7	*	*	*	1.9	7.8	20.4	25.6	8.7	3.2	2.7	6.2
2000	8.5	*	*	*	2.0	8.5	20.6	24.1	7.5	3.0	2.4	7.6
1999	8.7	*	*	*	2.2	9.3	21.2	23.9	8.2	3.4	2.2	*
Non-Hispanic black, male												
2006 5/	17.8	*	*	*	3.5	16.1	32.6	52.7	35.8	12.9	*	*
2005 5/	14.7	*	*	*	2.9	13.2	25.7	45.3	29.6	8.9	6.4	*
2004 5/	13.1	*	*	*	3.1	11.4	24.6	39.9	24.7	8.7	6.6	*
2003 5/	13.0	*	*	*	2.5	10.9	28.4	39.7	19.0	8.4	*	*
2002	13.2	*	*	*	3.0	12.2	28.3	40.1	20.2	5.0	*	*
2001	12.5	*	*	*	2.9	11.0	27.5	40.1	14.8	5.2	*	*
2000	12.3	*	*	*	2.7	12.3	28.7	38.1	12.3	5.1	*	*
1999	12.9	*	*	*	3.4	13.8	30.4	38.3	15.0	5.3	*	*
Non-Hispanic black, female												
2006 5/	7.6	*	*	*	1.8	5.2	16.1	21.4	10.0	2.9	3.9	7.5
2005 5/	7.2	*	*	*	1.4	6.5	15.9	19.7	8.1	2.5	*	*
2004 5/	6.6	*	*	*	1.3	5.2	14.7	19.1	6.7	*	*	*
2003 5/	6.3	*	*	*	1.2	5.9	15.2	17.1	5.9	2.5	*	*
2002	6.0	*	*	*	1.3	5.6	14.9	16.3	4.9	*	*	*
2001	5.3	*	*	*	0.9	5.0	14.1	13.1	3.7	*	*	*
2000	5.0	*	*	*	1.3	5.0	13.4	12.1	3.6	*	*	*
1999	4.9	*	*	*	1.1	5.3	13.1	11.5	2.7	2.1	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/												
All origins 3/, both sexes												
2006	10.3	*	0.3	0.9	16.9	15.3	11.5	11.2	9.8	10.0	12.9	11.5
2005	10.4	*	0.4	0.8	16.2	15.7	11.6	11.2	9.8	10.3	13.7	12.0
2004	10.1	*	0.3	0.7	15.7	15.0	11.3	11.0	9.8	10.2	13.3	11.9
2003	10.4	*	0.3	0.8	16.6	15.5	11.5	11.2	10.1	10.4	13.5	12.5
2002	10.5	*	0.4	0.8	16.7	15.4	12.1	10.8	10.2	10.9	14.4	12.5
2001	10.4	*	0.5	0.8	16.7	15.5	11.7	10.5	10.1	10.9	14.3	12.8
2000	10.2	*	0.3	0.9	16.8	14.5	11.9	10.5	9.4	10.6	13.9	14.2
1999	10.3	*	0.4	1.0	17.6	14.9	11.6	10.2	9.7	11.0	14.2	13.5
All origins 3/, male												
2006	18.1	*	0.5	1.2	29.8	26.6	18.9	18.5	17.0	19.2	29.1	33.6
2005	18.3	*	0.5	1.2	28.7	27.2	19.2	18.9	17.4	19.7	30.8	35.4
2004	17.6	*	0.4	1.1	27.5	26.1	18.7	18.3	17.1	19.7	29.8	35.9
2003	18.3	*	0.3	1.2	29.2	27.1	19.1	18.8	17.7	20.3	30.2	37.8
2002	18.4	*	0.5	1.2	29.3	26.5	20.1	18.2	18.0	21.3	32.9	38.9
2001	18.2	*	0.5	1.2	29.6	26.7	19.2	17.8	17.4	21.1	32.8	40.2
2000	17.8	*	0.4	1.4	29.4	24.9	19.4	17.6	16.3	20.3	32.2	44.7
1999	18.1	*	0.5	1.5	30.6	25.5	18.9	17.0	17.1	21.5	32.8	43.3
All origins 3/, female												
2006	2.8	*	*	0.4	3.2	3.7	4.1	4.0	3.2	2.2	1.8	1.1
2005	2.7	*	0.3	0.4	3.0	3.8	4.0	3.7	2.8	2.5	2.1	1.3
2004	2.7	*	0.3	0.3	3.2	3.7	3.9	4.0	3.1	2.2	2.3	1.0
2003	2.7	*	0.3	0.3	3.3	3.6	4.0	3.8	2.9	2.2	2.2	1.3
2002	2.8	*	0.3	0.5	3.5	4.0	4.2	3.6	3.1	2.3	2.1	1.1
2001	2.8	*	0.4	0.4	3.2	4.0	4.3	3.5	3.3	2.4	2.2	1.3
2000	2.8	*	*	0.4	3.5	4.0	4.4	3.6	3.0	2.5	2.0	1.7
1999	2.9	*	0.4	0.5	3.9	4.1	4.4	3.6	2.9	2.4	2.1	1.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/												
Hispanic, both sexes												
2006	7.8	*	*	0.9	19.1	12.6	7.8	5.4	4.0	2.5	4.7	*
2005	8.1	*	*	0.6	18.8	13.8	8.3	6.1	3.4	4.2	5.3	*
2004	7.9	*	*	0.6	18.5	12.7	7.7	6.2	4.8	4.4	5.4	*
2003	8.3	*	*	0.7	19.2	13.3	8.1	6.8	4.1	4.7	4.7	*
2002	8.1	*	*	0.8	18.5	12.5	7.9	6.0	5.0	5.3	5.5	*
2001	8.3	*	*	0.6	18.4	13.5	8.6	6.3	4.4	5.3	5.6	*
2000	8.2	*	*	0.8	18.0	12.1	8.1	8.1	4.9	5.9	5.5	*
1999	8.5	*	*	1.1	18.8	12.2	8.7	7.4	5.3	4.9	9.2	*
Hispanic, male												
2006	13.7	*	*	1.3	33.6	20.9	12.9	9.1	7.5	5.0	11.1	*
2005	14.2	*	*	0.9	33.0	22.9	13.4	10.4	6.7	8.1	12.0	*
2004	13.9	*	*	0.8	32.4	21.3	12.9	10.6	8.6	8.8	11.8	*
2003	14.6	*	*	1.0	32.8	22.8	13.3	11.5	8.1	10.0	10.7	*
2002	14.2	*	*	1.0	32.1	21.2	12.9	10.1	9.5	10.9	13.1	*
2001	14.6	*	*	0.9	31.4	22.7	14.4	10.7	8.6	11.0	12.5	*
2000	14.2	*	*	1.3	30.8	20.3	13.2	13.5	9.1	11.4	12.8	*
1999	14.6	*	*	1.5	31.6	20.6	14.3	11.8	9.9	10.1	21.9	*
Hispanic, female												
2006	1.5	*	*	*	2.7	2.4	2.1	1.7	*	*	*	*
2005	1.6	*	*	*	2.6	2.8	2.5	1.7	*	*	*	*
2004	1.5	*	*	*	2.6	2.4	2.0	1.7	*	*	*	*
2003	1.7	*	*	*	3.5	2.1	2.4	2.0	*	*	*	*
2002	1.6	*	*	*	2.8	2.3	2.5	1.9	*	*	*	*
2001	1.7	*	*	*	3.3	2.7	2.3	2.0	*	*	*	*
2000	1.8	*	*	*	2.9	2.6	2.5	2.8	*	*	*	*
1999	2.0	*	*	*	3.8	2.4	2.6	3.0	*	*	*	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/												
Non-Hispanic 4/, both sexes												
2006	10.7	*	0.4	0.8	16.3	15.9	12.1	11.8	10.3	10.6	13.4	11.8
2005	10.7	*	0.4	0.9	15.6	16.1	12.1	11.7	10.3	10.8	14.1	12.3
2004	10.4	*	0.4	0.8	15.0	15.5	11.8	11.5	10.2	10.6	13.8	12.1
2003	10.6	*	0.3	0.8	16.0	16.0	12.0	11.6	10.5	10.8	13.9	12.7
2002	10.8	*	0.4	0.9	16.2	15.9	12.7	11.2	10.6	11.2	14.8	12.7
2001	10.6	*	0.5	0.9	16.3	15.8	12.1	10.9	10.4	11.2	14.7	12.9
2000	10.4	*	0.3	0.9	16.4	14.9	12.3	10.7	9.7	10.8	14.3	14.4
1999	10.6	*	0.5	1.0	17.3	15.3	11.9	10.4	10.0	11.4	14.4	13.6
Non-Hispanic 4/, male												
2006	18.8	*	0.5	1.2	28.8	28.0	19.9	19.6	17.8	20.3	30.2	34.7
2005	18.9	*	0.6	1.3	27.6	28.2	20.2	19.8	18.2	20.5	31.8	36.3
2004	18.2	*	0.4	1.2	26.4	27.2	19.7	19.1	17.7	20.5	30.8	37.0
2003	18.8	*	0.4	1.3	28.3	28.0	19.9	19.5	18.4	21.0	31.2	38.7
2002	19.0	*	0.5	1.2	28.4	27.6	21.2	18.9	18.5	21.9	33.8	39.5
2001	18.7	*	0.5	1.2	29.0	27.5	19.9	18.4	17.9	21.7	33.6	40.8
2000	18.3	*	0.4	1.5	28.9	25.7	20.1	17.9	16.7	20.7	33.0	45.8
1999	18.5	*	0.5	1.5	30.3	26.3	19.3	17.4	17.6	22.1	33.2	43.9
Non-Hispanic 4/, female												
2006	2.9	*	*	0.5	3.2	3.9	4.4	4.2	3.4	2.3	1.9	1.2
2005	2.8	*	*	0.5	3.1	4.0	4.2	3.9	3.0	2.6	2.1	1.3
2004	2.9	*	*	0.3	3.3	3.9	4.2	4.3	3.2	2.3	2.3	1.0
2003	2.9	*	*	0.3	3.2	3.9	4.2	4.0	3.1	2.3	2.3	1.3
2002	3.0	*	*	0.5	3.6	4.3	4.4	3.8	3.2	2.4	2.2	1.2
2001	3.0	*	0.4	0.5	3.1	4.3	4.6	3.7	3.5	2.5	2.3	1.4
2000	3.0	*	*	0.4	3.6	4.2	4.6	3.7	3.1	2.6	2.1	1.7
1999	3.0	*	0.4	0.5	3.9	4.4	4.6	3.6	3.0	2.5	2.2	1.5

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/												
Non-Hispanic white, both sexes												
2006 5/	9.1	*	*	0.6	8.2	9.7	11.1	12.2	11.2	11.6	14.6	12.6
2005 5/	9.2	*	0.3	0.7	8.2	10.3	11.1	12.2	11.3	11.8	15.3	13.3
2004 5/	9.1	*	*	0.6	8.5	10.2	11.1	12.0	10.9	11.5	14.8	13.1
2003 5/	9.3	*	*	0.6	8.8	10.4	11.2	12.0	11.3	11.8	14.9	13.8
2002	9.4	*	0.3	0.6	9.2	10.7	11.8	11.5	11.4	12.3	15.8	13.7
2001	9.4	*	0.3	0.8	9.4	11.4	11.4	11.4	11.2	12.1	15.7	14.0
2000	9.1	*	0.2	0.8	9.6	10.5	11.5	10.8	10.2	11.5	15.4	15.5
1999	9.3	*	0.3	0.9	10.1	11.2	11.2	10.5	10.5	12.2	15.3	14.7
Non-Hispanic white, male												
2006 5/	15.6	*	*	0.9	13.9	16.3	17.8	19.9	19.0	21.9	32.6	37.0
2005 5/	15.9	*	*	1.0	13.9	16.9	17.8	20.3	19.6	22.2	34.2	39.2
2004 5/	15.6	*	*	0.9	14.3	16.9	17.8	19.6	18.7	21.8	32.9	40.1
2003 5/	16.0	*	*	0.9	15.2	17.3	18.1	19.9	19.7	22.7	33.5	42.1
2002	16.3	*	*	0.9	15.6	17.4	19.3	19.2	19.7	23.7	35.9	42.5
2001	16.3	*	*	1.2	16.0	18.9	18.4	19.0	19.0	23.1	35.7	44.0
2000	15.7	*	*	1.3	16.2	17.2	18.4	18.0	17.5	21.9	35.5	49.6
1999	15.9	*	*	1.3	17.2	18.6	17.8	17.6	18.1	23.5	35.1	47.5
Non-Hispanic white, female												
2006 5/	2.8	*	*	0.3	2.2	3.1	4.4	4.6	3.8	2.6	2.1	1.1
2005 5/	2.8	*	*	0.3	2.2	3.5	4.3	4.2	3.3	2.7	2.4	1.5
2004 5/	2.9	*	*	0.3	2.5	3.3	4.3	4.6	3.5	2.6	2.5	1.0
2003 5/	2.7	*	*	0.2	2.2	3.4	4.3	4.2	3.4	2.4	2.4	1.4
2002	2.8	*	*	0.4	2.5	3.8	4.3	4.0	3.5	2.6	2.3	1.2
2001	2.9	*	*	0.4	2.5	3.7	4.4	3.8	3.9	2.7	2.4	1.5
2000	2.9	*	*	0.4	2.7	3.7	4.5	3.8	3.4	2.8	2.2	1.8
1999	2.9	*	*	0.5	2.8	3.8	4.5	3.6	3.3	2.7	2.3	1.6

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/												
Non-Hispanic black, both sexes												
2006 5/	22.0	*	1.1	1.8	52.1	49.4	21.6	12.6	6.9	5.9	5.3	6.4
2005 5/	21.1	*	1.1	1.7	48.9	47.9	21.6	11.9	6.3	6.6	5.7	*
2004 5/	19.9	*	1.1	1.6	45.7	45.7	19.4	11.3	8.1	6.4	6.8	*
2003 5/	20.7	*	0.9	1.8	49.3	47.4	19.7	11.7	6.9	6.2	6.0	*
2002	20.7	*	*	1.6	49.2	46.4	21.0	11.9	7.2	5.8	6.9	*
2001	19.8	*	1.0	1.2	49.6	42.2	19.5	10.7	6.9	6.6	7.4	*
2000	19.8	*	*	1.4	49.8	40.9	19.7	12.1	6.8	7.0	5.6	*
1999	20.0	*	1.1	1.6	51.8	38.9	18.8	11.6	8.4	6.8	7.6	*
Non-Hispanic black, male												
2006 5/	41.4	*	*	2.7	95.2	93.3	39.4	23.4	13.4	12.5	14.2	*
2005 5/	40.0	*	*	2.5	90.1	92.0	40.0	21.9	11.8	13.0	14.3	*
2004 5/	37.6	*	*	2.4	83.7	87.5	36.2	20.6	14.9	14.8	16.3	*
2003 5/	39.1	*	*	2.6	90.5	91.1	36.1	21.6	13.1	12.0	13.6	*
2002	38.9	*	*	2.3	89.8	89.0	37.9	21.7	13.7	12.0	17.0	*
2001	37.3	*	*	1.8	93.1	80.4	34.1	19.3	13.6	13.8	17.7	*
2000	37.1	*	*	2.2	92.2	77.6	35.3	21.7	12.8	15.4	12.3	*
1999	37.1	*	*	2.4	95.0	73.8	33.1	20.7	16.8	14.3	18.4	*

Worktable Orig290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/												
Non-Hispanic black, female												
2006 5/	4.2	*	*	0.9	7.9	8.7	5.8	3.3	1.7	*	*	*
2005 5/	3.8	*	*	0.9	6.9	7.3	5.2	3.3	1.8	2.1	*	*
2004 5/	3.8	*	*	0.7	7.1	7.4	4.5	3.4	2.6	*	*	*
2003 5/	4.0	*	*	0.9	7.6	7.6	5.2	3.2	2.0	2.1	*	*
2002	4.3	*	*	0.9	8.4	7.8	6.1	3.5	2.1	*	*	*
2001	4.0	*	*	*	6.3	7.7	6.7	3.3	1.5	*	*	*
2000	4.2	*	*	*	7.9	7.7	5.9	3.9	2.0	*	*	*
1999	4.5	*	*	0.9	9.3	7.6	6.2	3.8	1.6	*	*	*

* Figure does not meet standards of reliability or precision.
0.0 Quantity more than zero but less than 0.05.
... Category not applicable.
1/ Figures for age not stated included in "All ages" but not distributed among age groups.
2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).
3/ Figures for origin not stated are included in "All origins" but not distributed among specified origins.
4/ Includes races other than white and black.
5/ Multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2006, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2006 data-mortality."
6/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."
7/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.